Main manuscript for

Trajectories for the evolution of bacterial CO_2-concentrating mechanisms

Avi I. Flamholz[1,2,3,*,†], Eli Dugan[1,*], Justin Panich[4], John J. Desmarais[1], Luke M. Oltrogge[1], Woodward W. Fischer[3,5], Steven W. Singer[4], David F. Savage[1,6,†]

[1] Department of Molecular and Cell Biology, University of California, Berkeley, California 94720, United States
[2] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125
[3] Resnick Sustainability Institute, California Institute of Technology, Pasadena, CA 91125, USA
[4] Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
[5] Division of Geological & Planetary Sciences, California Institute of Technology, Pasadena, CA 91125
[6] Howard Hughes Medical Institute, University of California, Berkeley, California 94720

* Equal contribution

† Corresponding Authors: Avi I. Flamholz, David F. Savage
Emails: aflamhol@caltech.edu, savage@berkeley.edu

Author contributions:
Avi I. Flamholz: Conceived and designed the study with input from E.J.D. and D.F.S. Conducted preliminary *E. coli* experiments, mathematical modeling, analysis of experimental results, generated figures and wrote the paper with input from all authors.
Eli J. Dugan: Contributed to study design and paper writing. Performed *H. neapolitanus* fitness assays with J.J.D., generated *C. necator* mutants and plasmids for heterologous expression in *E. coli* and *C. necator*, performed *E. coli* growth assays.
Justin Panich: Performed *C. necator* growth experiments.
John J. Desmarais: Performed *H. neapolitanus* fitness assays with E.J.D. Assisted in analysis.
Luke M. Oltrogge: Collaborated with A.I.F. in modeling autotrophic metabolism.
Woodward Fischer: Provided guidance on geological context for this study.
Steven W. Singer: Provided expertise, lab space and materials for *C. necator* cultivation.
David F. Savage: Provided guidance in research design and analysis of experimental results, as well as lab space and materials for most experiments.

Competing interests: D.F.S. is a co-founder of Scribe Therapeutics and a scientific advisory board member of Scribe Therapeutics and Mammoth Biosciences. These companies were not involved in this work in any way.

Classification: Biological Sciences/Microbiology; Physical Sciences/Environmental Sciences

Keywords: physiology, evolution, photosynthesis, Earth history, synthetic biology.

This PDF file includes:

Main Text
Figures 1 to 7

Abstract

Cyanobacteria rely on CO_2 concentrating mechanisms (CCMs) that depend on ≈15 genes to produce two protein complexes: an inorganic carbon (Ci) transporter and a 100+ nm carboxysome compartment that encapsulates rubisco with a carbonic anhydrase (CA) enzyme. Mutations disrupting CCM components prohibit growth in today's atmosphere (0.04% CO_2), indicating that CCMs evolved to cope with declining environmental CO_2. Indeed, geochemical data and models indicate that atmospheric CO_2 has been generally decreasing from high concentrations over the last ≈3.5 billion years. We used a synthetic reconstitution of a bacterial CCM in *E. coli* to study the co-evolution of CCMs with atmospheric CO_2. We constructed strains expressing putative ancestors of modern CCMs — strains lacking one or more CCM components — and evaluated their growth in a variety of CO_2 concentrations. Partial forms expressing CA or Ci uptake genes grew better than controls in intermediate CO_2 levels (≈1%); we observed similar phenotypes in genetic studies of two autotrophic bacteria, *H. neapolitanus* and *C. necator*. To explain how partial CCMs improve growth, we advance a model of co-limitation of autotrophic growth by CO_2 and HCO_3^-, as both are required to produce biomass. Our model and results delineated a viable trajectory for bacterial CCM evolution where decreasing atmospheric CO_2 induces an HCO_3^- deficiency that is alleviated by acquisition of CAs or Ci uptake genes, thereby enabling the emergence of a modern CCM. This work underscores the importance of considering physiology and environmental context when studying the evolution of biological complexity.

Significance

The greenhouse gas content of the ancient atmosphere is estimated using models and measurements of geochemical proxies. Some inferred high ancient CO_2 levels using models of biological CO_2 fixation to interpret the C isotopes found in preserved organic matter. Others argued that elevated CO_2 could reconcile a faint young Sun with evidence for liquid water on Earth. We took a complementary "synthetic biological" approach to understanding the composition of the ancient atmosphere by studying present-day bacteria engineered to resemble ancient autotrophs. By showing that it is simpler to rationalize the emergence of modern bacterial autotrophs if CO_2 was once high, these investigations provided independent evidence for the view that CO_2 concentrations were significantly elevated in the atmosphere of early Earth.

Introduction

Nearly all carbon enters the biosphere through CO_2 fixation in the Calvin-Benson-Bassham (CBB) cycle. Rubisco is the carboxylating enzyme of that pathway and the most abundant enzyme on Earth [1, 2]. Rubisco is often considered inefficient due to relatively slow carboxylation kinetics [3–5] and non-specific oxygenation of its five-carbon substrate ribulose 1,5-bisphosphate, or RuBP [6, 7]. However, rubisco arose more than 2.5 billion years ago, when the Earth's atmosphere contained virtually no O_2 and, many argue, far more CO_2 than today [8, 9]. Over geologic timescales, photosynthetic O_2 production [8] and CO_2-consuming silicate weathering reactions [10] are thought to have caused a gradual increase in atmospheric levels of O_2 ($\approx$20% of 1 bar atmosphere today) and depletion of atmospheric CO_2 to present-day levels of a few hundred parts per million ($\approx$280 ppm pre industrial, $\approx$420 ppm or $\approx$0.04% today). The amount of atmospheric CO_2 during the Archean Eon (4-2.5 Ga) is challenging to estimate accurately from the geological record, but is thought to have been substantially higher than today, perhaps as high as 0.1-1 bar [9, 11, 12]. It is likely, therefore, that contemporary autotrophs grow on drastically lower levels of CO_2 than their ancestors did.

91
Fig. 1: Mechanism and potential routes for the evolution of the bacterial CO$_2$ concentrating mechanism. (A) Today
the bacterial CCM functions through the concerted action of two protein complexes - an inorganic carbon (Ci)
transporter at the cell membrane and carboxysome encapsulation of rubisco with carbonic anhydrase (CA). Ci uptake
leads to a high intracellular HCO$_3^-$ concentration, well above equilibrium with the external environment. Elevated
HCO$_3^-$ is converted to a high carboxysomal CO$_2$ concentration by CA activity located only there, which promotes
carboxylation by rubisco. (B) Mutants lacking genes coding for essential CCM components grow in elevated CO$_2$ but
fail to grow in ambient air, exemplified here by the chemoautotroph *H. neapolitanus*. Strains lacking the
carboxysomal CA (Δ*csosCA*) or an unstructured protein required for carboxysome formation (Δ*csos2*) failed to grow
in ambient air, but grew robustly in 5% CO$_2$ (> 10^8 colony forming units/ml, Figure S1). See Table S4 for description
of mutant strains. (C) We consider the CCM to be composed of three functionalities beyond rubisco itself: a CA
enzyme (magenta), an Ci transporter (dark brown) and carboxysome encapsulation of rubisco (light brown). Since
atmospheric O$_2$ levels were extremely low ($\approx$1 ppm, blue triangle) and CO$_2$ was likely quite high (perhaps 0.1-1 bar,
red triangle) during the Archean epoch, primordial CO$_2$-fixing bacteria would not have needed a CCM. We sought to
discriminate experimentally between the six sequential trajectories (dashed arrows) in which CCM components
could have been acquired.

107
Many photosynthetic organisms evolved CO$_2$ Concentrating Mechanisms (CCMs), which help meet the

challenge of fixing carbon in a low CO$_2$ atmosphere. CCMs concentrate CO$_2$ near rubisco and are found in

all Cyanobacteria, some Proteobacteria, as well as many eukaryotic algae and diverse plants (13). Because

CO$_2$ and O$_2$ addition occur at the same active site in rubisco (6), elevated CO$_2$ has the dual effects of

accelerating carboxylation and suppressing oxygenation of RuBP by competitive inhibition (14–16). As

shown in Figure 1A, bacterial CCMs are encoded by $\approx$15 genes comprising three primary features: (i) an

energy-coupled inorganic carbon (Ci) transporter at the cell membrane and (ii) a cytosolic 100+ nm protein

compartment called the carboxysome found that (iii) co-encapsulates rubisco with a carbonic anhydrase

(CA) enzyme (4, 17). Energized Ci transport produces a high HCO_3^- concentration in the cytosol (≈30 mM,

Fig. 1A), which is converted into a high carboxysomal CO_2 concentration by carbonic anhydrase activity,

localized exclusively to the carboxysome (16, 18, 19).

119
CCM genes are straightforward to identify as mutations disrupting essential CCM components prohibit

growth in ambient air (Fig. 1B) and mutants are typically grown in 1% CO_2 or more (14, 15, 20–22). At first

glance, therefore, the CCM appears to be "irreducibly complex" as all plausible recent ancestors - e.g.

strains lacking individual CCM genes - are not viable in the present-day atmosphere. Irreducible

complexity is incompatible with evolution by natural selection, so we and others supposed that bacterial

CCMs evolved over a protracted interval of Earth history when atmospheric CO_2 concentrations were

much greater than today (13, 23–26). We therefore hypothesized that ancestral forms of the bacterial

CCM (i.e. those lacking some genes and complexes required today) would have improved organismal

fitness in the elevated CO_2 environments that prevailed when they arose.

129
To test the hypothesis, we reconstructed present-day analogues of plausible CCM ancestors (henceforth

"analogues of ancestral CCMs") and tested their growth across a range of CO_2 partial pressures. Our goal

was to identify a stepwise pathway of gene acquisition supporting the evolutionary emergence of a

bacterial CCM by improving growth in ever-decreasing CO_2 concentrations (Fig. 1C). We focused on

trajectories involving sequential acquisition of genetic components because carbonic anhydrases (27), Ci

transporters (22), and homologs of carboxysome shell genes (28, 29) are widespread among bacteria and

could therefore be acquired horizontally.

One approach to constructing contemporary analogues of CCM ancestors is to remove CCM genes from a

native host. If CCM components were acquired sequentially, some single gene knockouts would be

analogous to recent ancestors, e.g. those lacking a complete carboxysome shell (30). We tested this

approach by assaying a whole-genome knockout library of a ɣ-proteobacterial chemoautotroph, *H.*

neapolitanus, in five CO_2 partial pressures (22, 31). As shown in Figure 2 and elaborated below, we found

that many CCM genes contribute substantially to organismal fitness even at CO_2 concentrations tenfold

greater than present-day atmosphere (0.5% CO_2), supporting the view that CCM components play an

important physiological role even in relatively high environmental CO_2 concentrations.

Removing single CCM genes from a native host can only produce analogues of recent ancestors, however.

We recently constructed a functional CCM in an engineered *E. coli* strain called CCMB1 (4). This strain

depends on rubisco carboxylation for growth and the expression of a full complement of CCM genes

enabled growth in ambient air. Here we used CCMB1 to construct analogues of ancestral CCMs, including

several lacking two or more essential components of modern CCMs. We assayed the growth of these

putative ancestors across a range of CO_2 pressures to determine whether any ancestral forms contribute

to organismal fitness — i.e. improve growth relative to a control strain expressing only rubisco — in

various CO_2 pressures. In the following sections we describe the growth of these strains and discuss how

it can inform our entwined understandings of bacterial physiology, CCM evolution and the CO_2 content of

Earth's ancient atmosphere.

Results and Discussion

CCM genes contribute to fitness even in elevated CO$_2$

Using a barcoded genome-wide transposon mutagenesis screen, we previously demonstrated

that a 20-gene cluster in *H. neapolitanus* contains all the genes necessary for a functional CCM

(4, 22). The original screen measured gene fitness across the entire genome via batch

competition assays, comparing the abundance of strains in high CO$_2$ (5%) and ambient air (0.04%)

via high-throughput sequencing (22). This effort demonstrated that at least 12/20 genes are

necessary components of the CCM as disruption of any one produced substantial and

reproducible growth defects in ambient air. To mimic the changes in atmospheric CO$_2$ that likely

occurred over Earth history, we quantified the same library in three additional CO$_2$ pressures to

cover five CO$_2$ levels: ambient ($\approx$0.04% CO$_2$), low (0.5%), moderate (1.5%), high (5%), and very

high (10%). Replicate experiments were strongly correlated (R > 0.85, Figure S2), implying a high

degree of reproducibility. We therefore proceeded to ask whether *H. neapolitanus* CCM genes

contribute to fitness in elevated CO$_2$ conditions.

Fig. 2: The contribution of *H. neapolitanus* CCM genes to organismal fitness depends on the environmental CO$_2$ concentration. *H. neapolitanus* is a chemoautotroph that natively utilizes a CCM in low CO$_2$ environments. (A) Using a barcoded transposon library constructed in this organism (22), we profiled the contributions of CCM genes to

organismal fitness during autotrophic growth across a range of CO_2 levels (*Methods*). Fitness is estimated as the $\log_2$
ratio of barcode counts between the endpoint sample and the pre-culture for each mutant (22, 32). Each point gives
the average fitness of multiple mutants to a single CCM gene in a given CO_2 condition. The library contained 30
mutants disrupting the carboxysomal CA gene (*csosCA*) and an average of ≈40 mutants per CCM gene. A fitness of -
2 implies that gene disruption was, on average, associated with a fourfold decrease in strain abundance after a
defined period of growth. Biological replicates are indicated by shading, and the gray bar gives the interquartile
range of fitness effects for all ≈1700 mutants across all CO_2 levels (-0.15-0.065). "Ci transport" includes 4 DAB genes
in two operons (22), "carboxysome" includes 6 nonenzymatic carboxysome genes, "rubisco" denotes the two
subunits of the carboxysomal rubisco, and "CA" the carboxysomal CA gene. *H. neapolitanus* also expresses a
secondary rubisco, which explains why disruption of the carboxysomal rubisco is not lethal in high CO_2 (Figure S3).
(B) On average, CCM genes contribute less to organismal fitness as environmental CO_2 levels rise. Error bars give the
standard deviation of fitness values across all CCM genes. See Figures S2-3 for analysis of reproducibility and Tables
S1-3 for detailed description of genes.
190
Figure 2 plots the effect of disrupting CCM genes across five CO_2 pressures, with genes grouped by their

known roles in the CCM. The library contained an average of ≈35 mutants per gene in the *H. neapolitanus*

genome (22), so each point in Figure 2A represents the average fitness of 5-50 individual mutants.

Surprisingly, we found that many CCM genes also contributed substantially to organismal fitness in 0.5%

and 1.5% CO_2, indicated by large fitness defects (negative values in Figure 2A), resulting in a negative

average fitness of CCM mutants when the CO_2 pressure was 1.5% or less (Figure 2B). Carboxysome genes,

for example, were critical for growth in 0.5% CO_2, while certain Ci transport genes contributed

substantially to growth in 0.5% and 1.5% CO_2.

Though several *H. neapolitanus* CCM genes contributed to fitness in intermediate CO_2 levels, the CCM

appeared to be dispensable in high CO_2 (5-10%, Fig. 2B). As such, the data presented in Figure 2 indicated

that individual CCM components like the carboxysome, CA, or Ci transporter may improve autotrophic

growth in intermediate CO_2 levels (≈1%) even in the absence of a complete and functional CCM. We

considered testing this hypothesis in *H. neapolitanus* directly by constructing "ancestral-like" CCMs lacking

carboxysome shell genes or Ci transporters. However, genetic manipulation of *H. neapolitanus* is

cumbersome, and the native host's regulatory network could complicate interpretation — a concern

highlighted by three DNA-binding proteins that emerged in our screen as likely regulators of the CCM

(Figure S3 and Table S1). We therefore decided to construct and test analogues of CCM ancestors in a

non-native host, namely *E. coli*.

Evaluating putative ancestral CCMs in a rubisco-dependent *E. coli*

We recently developed an *E. coli* strain, CCMB1, that depends on rubisco carboxylation for growth in

minimal medium. This strain requires elevated CO_2 for rubisco-dependent growth, but expressing the *H.*

neapolitanus CCM from two plasmids enabled growth in ambient air (4). One of these plasmids, pCB',

encodes the carboxysome genes along with the encapsulated rubisco and carbonic anhydrase enzymes

(4). The other plasmid, pCCM', encodes the DAB1 inorganic carbon transporter (4, 22), two rubisco

chaperones (22, 33, 34) and a carboxysome positioning system (35). We used this two-plasmid system to

express analogues of putative CCM ancestors in CCMB1 and assayed these strains' growth in ambient air,

0.5%, 1.5% and 5% CO_2. We compared the growth of strains expressing partial CCMs to that of a reference

CCMB1 strain expressing the *H. neapolitanus* rubisco on the p1A vector but no other CCM components

(4). The reference strain grew robustly in 5% CO_2 but failed to grow in ambient air (Fig. 3A, 'Rubisco

Alone').

Consistent with our previous work (4), expression of the full complement of CCM genes permitted growth

in all CO_2 concentrations tested (Fig. 3A, 'Full CCM'). Recent modeling efforts endeavored to support the

hypothesis that encapsulation of rubisco in a semi-permeable barrier could improve CO_2 fixation by

generating an acidic local pH (26). This proposal does not require Ci transport or CA activity and, if correct,

would therefore support an "encapsulation first" model of CCM evolution. To evaluate the effect of

encapsulating rubisco alone in a protein compartment, we replaced the pCCM' plasmid with a vector

control so that no Ci transporter was expressed and further deactivated the carboxysomal CA by mutating

a crucial cysteine residue (pCB' CsosCA C173S, 'Encap. Rub.' in Fig. 3A). Encapsulating rubisco on its own

did not improve growth over the reference strain in any CO_2 concentration tested. However, when we left

the CA active site intact (`En. Rub.+CA'), growth improved substantially in intermediate CO_2 levels (0.5%

and 1.5%). These data indicated that CA plays a pivotal role in autotrophic growth in low CO_2, but did not

support the hypothesis that encapsulation alone improves CO_2 fixation by rubisco (26).

236
Fig. 3: Encapsulation of rubisco alone does not improve rubisco dependent growth of CCMB1 *E. coli* in any CO₂
level tested. (A) Using our reconstitution of the *H. neapolitanus* CCM in *E. coli* (4) we tested whether rubisco
encapsulation has any effect on growth at four CO_2 partial pressures (Methods). Each diamond represents one
technical replicate of four biological replicates. The CCMB1 strain grows in elevated CO_2 (1.5 and 5%) when rubisco
is expressed ("Rubisco Alone", left). As previously reported, expressing the full complement of CCM genes from the
pCB' and pCCM' plasmids ("Full CCM") enabled growth in all CO_2 levels. By replacing pCCM with a vector control and
making an inactivating mutation to the carboxysomal CA (CsosCA C173S) we were able to express rubisco in a
carboxysome without CA or Ci transport activities ("Encap. Rub."). This strain grew similarly to the reference
"Rubisco Alone" in all conditions. When the CA active site was left intact ("En. Rub.+CA"), growth improved above
the "Rubisco Alone" baseline in 0.5% and 1.5% CO_2. Finally, a negative control strain carrying inactive rubisco ("En.
Rub.⁻", CbbL K194M) failed to grow in all CO_2 conditions. (B) Focusing on the growth in 1.5% CO_2 highlights the
contribution of CA activity to rubisco-dependent growth. See Tables S4-5 for description of strains and plasmids, see
Figures S4-5 for growth curves and analysis of statistical significance.

Carbonic anhydrase and energy-coupled inorganic carbon transport
improve the growth of rubisco-dependent E. coli

Our observation that carboxysomal CA activity improves rubisco-dependent growth on its own (i.e.

without also expressing a Ci transporter) motivated us to test the effects of CA and Ci transport

independently of carboxysome expression. We designed plasmids that express the *H. neapolitanus* Dab2

Ci transporter (22), *E. coli*'s native CA (Can) or both. This was achieved by cloning both Dab2 and Can into

a dual expression vector and making active site mutants (DabA C539A, Can C48A, D50A) to isolate each

activity. These vectors were co-transformed into CCMB1 with a constitutive version of p1A (p1Ac) so that

rubisco expression would not be affected by induction of Dab2 or Can (*Methods*).

260
Fig. 4: Expression of carbonic anhydrase or Ci transport improves rubisco-dependent growth of CCMB1 *E. coli* in
intermediate CO_2 levels even in the absence of other CCM components. As in Figure 3, the reference strain
("Rubisco Alone") grew only in elevated CO_2 (1.5% and 5%). The remaining strains expressed the *E. coli* carbonic
anhydrase (Can) and the DAB2 Ci transporter from a second plasmid. These activities were isolated by means of
active site mutations. The negative control "+DAB⁻ +CA⁻" expressed inactive Can (C48A, D50A) and DAB2 (DabA2
C539A) and grew less robustly than the reference in all conditions. If either active site was left intact ("+DAB+CA⁻"
or "+DAB⁻+CA") we observed a sizable growth improvement in both 0.5 and 1.5% CO_2. This growth improvement
remained when both active sites were left intact ("+DAB+CA"). Panel (B) emphasizes this effect clearly by focusing
on growth in 1.5% CO_2. See Table S4 for strain genotypes, Figures S6-7 for growth curves and analysis of statistical
significance.
271
Expressing active Can and Dab2, whether alone or together, improved growth substantially in 1.5% CO_2

(compare '+DAB+CA⁻', '+DAB⁻+CA', and '+DAB+CA' to 'Rubisco Alone' in Figure 4). This effect was even

more pronounced in 0.5% CO_2, where the reference strain failed to grow (Figures 4A and S7). Similar to

the reference strain, a double-negative control strain expressing inactivated versions of both Dab2 and

Can ('+Dab⁻+CA⁻') grew poorly or not at all in 0.5% and 1.5% CO_2, implying that the observed growth

improvements were due to activity and not a side-effect of heterologous gene expression.

It was not immediately obvious to us how CA and Ci uptake activities improve rubisco-dependent growth

of CCMB1 *E. coli*. We found the effects of Dab2 expression especially perplexing because Ci uptake is

expected to generate intracellular HCO_3^- (16–18, 22, 36) while the rubisco substrate is CO_2 (37). To confirm

that these results are not a side-effect of working in an engineered *E. coli* strain, we pursued genetic

experiments in a natively autotrophic proteobacterium, *C. necator*.

284   *C. necator* depends on CA for autotrophic growth at intermediate pCO2

285   While all photosynthetic Cyanobacteria rely on the CBB cycle and a full complement of CCM genes (38),

286   some chemoautotrophic bacteria depend on the CBB cycle but lack identifiable genes encoding

287   carboxysome components or Ci transporters (39–42). As most characterized bacterial rubiscos are not

288   CO_2-saturated in ambient air and are, in addition, substantially inhibited by atmospheric levels of O_2 (7),

289   we expected that such organisms would require elevated CO_2 for robust growth.

290
Fig. 5: *C. necator* requires carbonic anhydrase or Ci uptake for robust autotrophic growth in 0.5% and 1.5% CO_2. *C.*
necator strains were grown autotrophically in minimal medium at a variety of CO_2 levels and endpoint optical density
was measured after 48 hours (*Methods*). (A) Growth of the *C. necator* double carbonic anhydrase knockout (ΔCA)
was greatly impaired growth in 0.5% and 1.5% CO_2. Panel (B) focuses on 1.5% CO_2. Compared to wild-type *C. necator*
(WT), which grew to a final OD600 of 0.73±0.28 in 1.5% CO_2 (6 biological replicates), growth of ΔCA was greatly
impaired, reaching a final OD of 0.23±0.17 (3 biological replicates). Expression of either the human carbonic
anhydrase II (ΔCA+CA) or the DAB2 Ci transporter from *H. neapolitanus* (ΔCA+DAB) recovered robust growth which
exceeded even the wild type, indicating that the wild type may not express saturating levels of CA. See Figure S8 for
statistical analysis.

300
Cupriavidus necator, formerly known as *Ralstonia eutropha*, is one such bacterium (43, 44). *C. necator* is

a facultative chemolithoautotroph typically found at the interface between oxic and anoxic environments

where H_2 and O_2 coexist. Such "knallgas" environments include soils, sediments and geothermal sites (45)

that are often characterized by elevated CO_2 (45–47). While *C. necator* is an obligate aerobe capable of

chemoautotrophic growth on H_2, CO_2 and O_2 via the CBB cycle, it has no carboxysome genes and no

obvious Ci transporters (41). Consistent with previous reports (48), autotrophic growth of wild-type *C.*

necator was very poor in ambient CO_2 ('*C. necator* WT' in Figure 5A). We generated a double CA knockout

strain, *C. necator* Δcan Δcaa (42, 49) and found that CA removal greatly attenuated autotrophic growth in

0.5% and 1.5% CO_2 ('*C. necator* ΔCA'). Consistent with our experiments in *E. coli*, this growth defect was

complemented by heterologous expression of a human CA ('ΔCA+CA') or a DAB-type Ci uptake system

('ΔCA+DAB'). Moreover, as in *E. coli*, co-expression of Ci uptake with native CAs was not deleterious

('WT+DAB'). Rather, this strain grew to higher densities than wild-type in 1.5% and 5% CO_2 (e.g. Fig. 5B).

A nutritional requirement for HCO_3^- explains the observed
phenotypes

We found that expression of a CA, Ci transporter, or both improved rubisco-dependent growth of *C.*

necator and CCMB1 *E. coli* in intermediate CO_2 concentrations (0.5 and 1.5%, Figs. 4-5). Furthermore, to

our surprise, a CCMB1 strain expressing active Dab2 and Can grew reproducibly, if slowly, in ambient air

('+Dab+CA' in Figure 6). This was remarkable because biological membranes are very permeable to CO_2,

with a permeability coefficient $P_C \approx 0.1$-1 cm/s (16, 50, 51) that is several orders-of-magnitude larger than

those of small charged species like acetate, phosphate and bicarbonate (50–53). Given this high

permeability, we expected co-expression of energized HCO_3^- uptake with a CA to generate a deleterious

"futile cycle" where energy expended pumping HCO_3^- was wasted when CO_2 produced by CA activity

"leaks" back out of the cell. Instead we found that such a cycle is compatible with rubisco-dependent

growth of two bacteria in relatively low CO_2 (1.5% or lower).

A naive explanation of how CA expression might improve growth is that it increases the intracellular CO_2

concentration relative to the reference strain expressing only rubisco. For this hypothesis to explain the

benefit of CA expression, rubisco activity must deplete intracellular CO_2 (C_{in}) significantly below its

extracellular level (C_{out}) in the reference strain. Otherwise CA can have no effect, as carbonic anhydrases

are not coupled to any energy source and so cannot cause C_{in} to exceed C_{out}. However, this naive model

cannot explain the growth benefits associated with expressing Ci uptake systems, which provide HCO_3^-

and not CO_2. Moreover, the following calculation shows that this hypothesis is unreasonable precisely

because CO_2 is so membrane-permeable that rubisco cannot deplete C_{in} much beneath C_{out}.

Fig. 6: Co-expression of carbonic anhydrase and Ci uptake enabled rubisco-dependent growth of CCMB1 *E. coli* **in**
ambient air. Inspecting the ambient CO_2 growth data presented in Figure 4 revealed that coexpression of CA and Ci
transport ('Rub.+DAB+CA') substantially improved rubisco-dependent growth of CCMB1 *E. coli* in ambient CO_2
concentrations. This effect was modest (≈0.1 OD units above the 'Rubisco Alone' control) but reproducible, as
indicated by endpoint data plotted on the inset. Curves are colored to match labels on the inset. See Figure S9 for
statistical analysis.

In a bacterium, rubisco might make up 20% of soluble protein at the very most. This amounts to a mass

concentration of roughly 0.2 x 300 mg/ml ≈ 60 mg/ml rubisco (54). As each rubisco active site is attached

to ≈60 kDa of protein (BNID 105007), the maximum rubisco active site concentration is ≈1 mM. In this

naive model, C_{in} is set by the balance of passive uptake through the membrane, with effective permeability

α = P_C x SA/V ≈ 10^4 s^{-1}, and fixation by rubisco, with an effective rate constant of γ = [rubisco] x k_{cat}/K_M <

10^3 s^{-1}. These values give a steady-state C_{in} of α C_{out} / (α + γ) > 0.9 C_{out}. That is, even an extreme level of

rubisco activity cannot draw C_{in} beneath 90% of C_{out}. In such conditions, rubisco would fix ≈10^{10} CO_2/hour,

supporting a 1-2 hr doubling time (see *Supplementary Information* for full calculation). So, although the

CO_2 concentration is low in aqueous environments equilibrated with present-day atmosphere (≈10 μM),

passive CO_2 uptake can support substantial rubisco flux and, therefore, CO_2 limitation of rubisco

carboxylation cannot explain the observed phenotypes.

We argue that the effects of expressing CA or Ci transport can be explained by the ubiquitous dependence

of growth on HCO_3^- (55–57). It has been clear for at least 80 years that heterotrophs also require inorganic

carbon for growth. Seminal investigations in the 1930s and 1940s advanced the hypothesis that this

dependence is due to a specific requirement for HCO_3^- (58–61), which is the substrate of several

carboxylation reactions involved in lipid, nucleotide and arginine biosynthesis. This explanation is now

supported by detailed experiments in several heterotrophs demonstrating that growth in ambient air can

be supported either by CA activity (55–57) or by providing the products of central-metabolic carboxylation

reactions in the growth media (56, 57). The metabolic networks of chemoautotrophs, Cyanobacteria and

plants imply an equivalent dependence on HCO_3^- for biosynthesis (62–64). We therefore advance a model

of co-limitation of autotrophic growth by both CO_2 and HCO_3^--dependent carboxylation fluxes, where

most of biomass carbon derives from rubisco-catalyzed carboxylation of CO_2 and a small minority derives

from HCO_3^- (Figure 7A). We formalized this notion as an idealized mathematical model fully described in

Supplementary Information.

367
368

Fig. 7: Co-limitation of autotrophic growth by CO₂ and HCO₃⁻-dependent carboxylation reactions explains the growth improvements associated with expressing carbonic anhydrases and Ci transporters. (A) In autotrophs using the CBB cycle, nearly all biomass carbon derives from rubisco-catalyzed CO₂ fixation. However, autotrophs also require HCO₃⁻ for carboxylation reactions in lipid, nucleic acid, and arginine biosynthesis (55–57). We expressed this diagram as a mathematical model, which we applied to understand why CA and Ci uptake improved rubisco-dependent growth. (B) The model exhibited two regimes: one wherein growth was limited by rubisco flux another where it was limited by HCO₃⁻-dependent carboxylation ("bicarboxylation") flux. At low rubisco levels (lighter-colored lines), growth was rubisco-limited: increased rubisco activity produced faster growth, but the growth rate was insensitive to CA activity because slow CO₂ hydration provided sufficient HCO₃⁻ to keep pace with rubisco. At higher rubisco levels (maroon lines), growth was bicarboxylation-limited and increased CA activity was required for increasing rubisco activity to translate into faster growth. Increasing Ci uptake led to similar effects (Figure S11). In panel (C), color indicates the ratio of total Ci leakage ($J_{L,tot} = J_{L,C} + J_{L,H}$) to biomass production flux (J_B) at a fixed rubisco activity across a wide range of CA activities and Ci uptake rates. $J_{L,tot}$ was calculated as the sum of CO₂ and HCO₃⁻ leakage rates ($J_{L,C} + J_{L,H}$) with $J_{L,C} \gg J_{L,H}$ in most conditions due to the much greater membrane permeability of CO₂. So-called "futile cycling," where leakage greatly exceeds biomass production ($J_{L,tot} / J_B \gg 1$), occurs when CA and Ci uptake are co-expressed at extreme levels (redder colors). See *Supplementary Information* for detailed description of the co-limitation model.

When the HCO₃⁻-dependence of biomass production was included in our model, it became possible to

rationalize the growth improvements associated with heterologous expression of a CA or Ci transporter,

as both of these activities can supply HCO₃⁻ needed for biosynthesis (Figure 7A). When the modeled

rubisco activity was low, growth was rubisco-limited: increasing rubisco activity (lighter colored lines in

Figure 7B) or the CO₂ concentration produced faster growth, but the growth rate was insensitive to CA

activity because slow CO₂ hydration (whether spontaneous or CA catalyzed) provided sufficient HCO₃⁻ for

the anabolic carboxylations involved in biomass production (Figures S11 and S12). When rubisco activity

was set to higher values (darker lines in Figure 7B) the model entered a "bicarboxylation-limited" regime

where increased rubisco activity did not affect the growth rate until CA activity was increased to supply

HCO₃⁻ (i.e. bicarbonate) for anabolic carboxylations. In bicarboxylation-limited conditions, additional

HCO_3^- could be supplied by CA, Ci uptake, or both (Figures S11). However, these activities are not

equivalent, as we discuss below.

The co-limitation model also gave insight into the nature of the supposed "futile cycle," which could now

be framed quantitatively by comparing the leakage of inorganic carbon from the cell (flux $J_{L,tot} = J_{L,C} + J_{L,H}$)

to biomass production by rubisco and HCO_3^--dependent carboxylases (J_B, Fig. 7A). As CO_2 is orders of

magnitude more membrane-permeable than HCO_3^- near neutral pH (16), CO_2 leakage ($J_{L,C}$) greatly

exceeded the HCO_3^- leakage flux ($J_{L,H}$) in most conditions. Low Ci leakage is desirable; when $J_{L,tot} \approx 0$ the

flux of Ci pumped into the cell is balanced by fixation of CO_2 and HCO_3^- such that no energy is "wasted"

on Ci uptake. When CA (δ) and Ci uptake (χ) activities were low, the model predicted slow growth and net

diffusive Ci uptake ($J_{L,C}, J_{L,H} < 0$) to balance carboxylation (Figure 7C and S14). Increasing χ could cause $J_{L,tot}$

to change sign from negative (connoting passive uptake) to positive (connoting substantial leakage)

passing through $J_{L,tot} = 0$. If CA was also expressed - i.e. if δ was increased above a baseline value - $J_{L,tot} \approx 0$

could be achieved at lower χ values due to CA-catalyzed dehydration of imported HCO_3^- producing CO_2

used by rubisco. In other words, CA activity lowers the Ci uptake rate and, consequently, energy

expenditure, required to achieve the same rate of biomass production (Figures 7C & S14). When δ and χ

were both high, the model predicted substantial leakage, with $J_{L,tot} / J_B \approx 100$ in extreme cases. Such

extreme levels of futile cycling are likely incompatible with growth. If χ was low, in contrast, δ could be

increased arbitrarily without incurring leakage because CAs do not pump Ci into the cell.

A second, less plausible way in which co-expression might improve growth is if CA and Ci transport are

fast and membrane permeability to CO_2 is far lower than typically assumed. In this case, the combination

of Ci uptake and CA activity might form a CO_2 pump that elevates C_{in} substantially above C_{out} to accelerate

rubisco carboxylation (Figure S13). As HCO_3^- is supplied sufficiently by Ci uptake in this case, the co-

limitation model predicted that increased rubisco fluxes would translate into faster growth. For this effect

to arise, however, the membrane permeability to CO_2 would have to be 100-1000 times lower than the

measured (50) or calculated (51), so CO_2 pumping is unlikely to explain the capacity of a CCMB1 to grow

in ambient air when co-expressing Dab2 and Can (Fig. 6).

Taken together, our experiments and model helped outline a plausible trajectory for the co-evolution of

the bacterial CCM with atmospheric CO_2 levels. Presuming an ancestral autotroph with only a rubisco-

driven CBB cycle and no CCM components, our data and model support a trajectory where CA and Ci

transport are acquired together or serially (in either order) to support growth as atmospheric CO_2 levels

decreased (darker arrows in Fig. 1C). The order of acquisition might depend on the environmental pH,

which strongly affects the extracellular HCO_3^- concentration and, thereby, the expected efficacy of HCO_3^-

uptake (16). The co-limitation model helped us understand the potential advantages of expressing CA and

Ci uptake together: modest co-expression can reduce energy expended on Ci pumping and balance the

supply of CO_2 and HCO_3^- with the cellular demand for rubisco and bicarboxylation products (Figure 7C and

S14). In CCMB1 we found that co-expression of CA and Ci transport supported growth in low CO_2

environments (≈1%, Figure 4) and even permitted modest growth in atmosphere (Figure 6); cells

expressing both activities would have been "primed" for the subsequent acquisition and refinement of

proto-carboxysome structures that co-encapsulate rubisco and CA to enable robust growth at yet lower

CO_2 levels (e.g. ambient air, Figure 3). Notably, carboxysome shell proteins are structurally related to two

ubiquitous protein families (65) and homologous to other metabolic microcompartments (28, 65),

suggesting two plausible routes for the acquisition of carboxysome genes.

Results from our *E. coli* experiments indicated that such an evolutionary trajectory is "fitness positive" in

that each step improves organismal fitness at some environmental CO_2 level (Figs. 3, 4, and 6). Moreover,

the fitness contribution of CA and Ci transport activities was only realized at intermediate $pCO_2 \approx 1\%$ in

both *E. coli* (Fig. 4) and *C. necator* (Fig. 5), supporting the view that CO_2 concentrations declining from high

levels early in Earth history promoted the evolutionary emergence of bacterial CCMs.

Concluding Remarks

There is great and longstanding interest in characterizing the composition of the atmosphere over Earth

history (9, 11, 66). This interest is surely justified as the contents of the atmosphere affected the

temperature, climate, and the chemical conditions in which life arose, evolved, and has been maintained.

In the case of reactive species like O_2, present-day measurements of old sedimentary rocks are quite

informative: measurements of proxies for O_2 reactivity in geological samples demonstrate that the

Archean atmosphere contained very little dioxygen, with inferred O_2 levels of 1 ppm or less (8, 9). Many

posit that the Archean atmosphere also contained very high levels of CO_2. However, since CO_2 is far less

reactive in Earth surface environments than O_2, available geological constraints are meager and this

inference stands on shakier ground.

One reason for presuming high CO_2 levels in the early atmosphere is the paradox of the "Faint Young Sun,"

which would not have been sufficiently luminous to permit liquid water on Earth without a very

substantial greenhouse effect (67) or some other physical mechanism warming the Earth system (68).

Other approaches to estimating ancient CO_2 pressures rely indirectly on geochemical observations and

arguments from the geological record. For example, some studies infer high Archean CO_2 from the

presence or absence of certain carbonate minerals in ancient surface environments (9), while others rely

on models of biological CO_2 fixation to estimate CO_2 concentrations from the carbon isotope ratios of

organic matter in sedimentary rocks (11, 12). These diverse approaches vary in the magnitude of their

estimates, but each gives the impression that Archean CO_2 greatly exceeded present-day levels of roughly

0.04% of a 1 bar atmosphere; indeed, some estimates of Archean CO_2 pressures are as high ≈1 bar (9, 67).

Silicate weathering coupled to the deposition of carbonate-bearing rocks represents a long-term crustal

sink that — combined with subequal amounts of organic carbon burial — would have drawn a large

reservoir of atmospheric CO_2 down to pre-industrial levels (10, 69). Furthermore, modern plants, algae,

and autotrophic bacterial lineages collectively evolved four distinct classes of CCM at least 100 times

indicating that autotrophy has repeatedly and convergently adapted to decreasing environmental CO_2

levels (13).

Carbon isotope methods for inferring historical changes in CO_2 rely on uniformitarian models of biological

carbon fixation to calculate atmospheric CO_2 pressures from isotope ratio measurements of biogenic

organic phases preserved in sediments and rocks (12). Yet autotrophy has undergone considerable

evolution and diversification over the last ≈3 billion years, which complicates inference of ancient CO_2

levels with models based on contemporary autotrophs. For example, rubiscos from different lineages

display variable $^{12}CO_2$ preferences (23, 70) and contemporary CCMs rely on Ci uptake systems that may

fractionate C isotopes to a degree similar to rubisco (71). As demonstrated in the companion paper by

Renee Wang and co-authors, the evolution of CCMs must be considered when attempting to reconstruct

historical CO_2 concentrations from the C isotope record.

Here we attempted to reconstruct the evolution of a bacterial CCM by constructing analogues of plausible

ancestors. This investigation was initially motivated by a basic curiosity about the evolution of

physiological complexity, hoping to resolve the apparent "irreducible complexity" of bacterial CCMs

(Figure 1). Irreducible complexity is incompatible with evolution by natural selection, and so we

hypothesized that ancestral CCMs conferred a growth advantage in the historical environments in which

they arose, which we and others presume were characterized by high CO_2 (5, 24–26).

Results from our experiments with rubisco-dependent *E. coli* and two bacterial autotrophs (*H.*

neapolitanus and *C. necator*) supported this rationalization of CCM evolution by showing (i) that $CO_2 \approx 1\%$

improves the growth of "ancestral" CCMs (Figs. 3-6) and (ii) that all CCM genes are dispensable during

growth in 5-10% CO_2 (Fig. 2-5). This latter result suggested to us that atmospheric CO_2 concentrations

exceeded 1% of a 1 bar atmosphere when rubisco arose, which was likely more than 3 billion years ago

(24, 72). If geological CO_2 sinks later brought atmospheric CO_2 to ≈1%, all organisms, including autotrophs,

would have begun to evolve or acquire carbonic anhydrases and/or Ci transporters to provide the HCO_3^-

required for biosynthesis (Figure 7). An ancestral autotroph expressing both these activities may have had

a growth advantage in relatively lower CO_2 pressures < 1% (Fig. 6A), and would have been "primed" for

the evolution of a CCM as the only missing component, the carboxysome, could have evolved from

oligomeric host proteins (65) or adapted from a different metabolic microcompartment (28).

Alternatively, it is possible that CAs (or Ci transporters) arose prior to rubisco and were already widespread

at the time of rubisco evolution, in which case we might expect $CO_2 \gtrsim 1\%$ when rubisco arose.

Unfortunately, the convergent evolution of CA activity in several protein families (73) makes it very

challenging to constrain the timing of CA evolution with comparative biological and molecular clock

approaches; this issue concerns bacterial Ci transporters as well (17, 22).

It is intrinsically difficult to answer questions about Earth's deep biological history; addressing such

questions will surely require cooperation between scientific disciplines. Here, and in the companion paper

by Renee Wang et. al., we took a "synthetic biological" approach to study the molecular evolution of

bacterial CO_2 fixation by constructing contemporary cells intended to resemble ancient ones. In both

cases, our work highlighted the impact of environmental context (e.g. CO_2 concentrations) and whole cell

physiology (the requirement for HCO_3^-) on the evolution of CO_2 fixation. That is, neither rubisco nor the

CCM should be considered in isolation, but rather in the context of a metabolism that demands both CO_2

and HCO_3^- to produce biomass (55–61, 71). We hope that future research advances the synthetic approach

to studying evolution and fully expect that this approach will enrich our understanding of biological

processes that have shaped the evolution of biogeochemical cycles on Earth.

Methods

Strains, plasmids and genomic modifications

Strains and plasmids used in this study are documented in Supplementary Tables S4 and S5. The rubisco-dependent *E. coli* strain CCMB1 was derived from *E. coli* BW25113 and has the genotype BW25113 $\Delta rpiAB$ Δedd $\Delta cynT$ Δcan, as documented in (4). To construct CA deficient mutants of *C. necator* H16, we first knocked out the hdsR homolog *A0006* as removal of this restriction enzyme increases electroporation efficiency (41). The double CA knockout, *C. necator* H16 $\Delta A0006$ Δcan Δcaa, was constructed by repeated rounds of selection and counter-selection by integrating a construct encoding both kanamycin resistance and *sacB* for counter-selection. Protocols for plasmid construction and genomic modification of *C. necator* are fully described in the *Supplementary Information*.

Genome-wide fitness measurements in *H. neapolitanus*

Competitive fitness assays were performed following (22). The barcoded *H. neapolitanus* transposon library generated for that work was thawed and used to inoculate three 33 ml cultures that were grown overnight at 30 C in DSMZ-68 with 10 μg/ml Kanamycin. These pre-cultures were grown to an OD ≈ 0.07 (600 nm) in 5% CO_2. The library was subsequently back-diluted 1:64 and grown in various CO_2 concentrations (10%, 5%, 1.5%, 0.5% and ambient CO_2) on a platform shaker (New Brunswick Scientific Innova 2000, 250 RPM) in a Percival Intellus Incubator; 20 ml of pre-culture was pelleted by centrifugation (15 min at 4000g) and saved as a T_0 reference. Upon reaching 6.5-7.5 doublings, 50 mls of culture was spun down and gDNA was extracted for barcode PCRs as described in (6); barcodes were previously mapped to the genome via TnSeq (22). PCRs were purified (Zymo Research Clean and Concentrator kit) and pooled for sequencing on an Illumina MiSeq with 150 bp single end reads. We used the software pipeline from (32) to analyze barcode abundance data. Briefly, fitness of individual mutant strains was calculated as the $\log_2$ of the ratio of barcode abundance in the experimental condition over abundance in the T_0. Gene-level fitness values were then calculated considering all transposon insertions expected to

547 disrupt an individual gene. Each CO_2 concentration was tested in biological duplicate except for 5% CO_2,

548 which was assayed in biological quadruplicate.

E. coli growth conditions

E. coli strains were grown at 37 °C. 60 µg/ml kanamycin and 25 µg/ml chloramphenicol were used for

selection during routine cloning and propagation. For strains carrying two selectable markers, antibiotics

were used at half concentration (30 µg/ml kanamycin, 12.5 µg/ml chloramphenicol). All CCMB1 cultures

were grown in 10% CO_2 unless otherwise specified. When rubisco-independent growth was desired,

CCMB1 was propagated in rich LB media. CCMB1 was cultured in a rubisco-dependent manner in M9

minimal media supplemented with trace elements and 0.4% glycerol (v/v) as described in (4). Experiments

presented in Figures 3, 4 and 6 were conducted in 96-well plates in a gas-controlled plate reader (Tecan

Spark). For the data in Figure 3, 100 nM anhydrous tetracycline (aTc) was supplied to induce expression

from p1A, pCB' and pCCM' plasmids. For the data presented in Figures 4 and 6, a constitutive version of

p1A was used (p1Ac) and aTc was omitted from pre-cultures. These strains also carried a dual-expression

plasmid, pFC, for inducible expression of the DAB2 Ci transport operon and the *can* carbonic anhydrase.

100nM aTc was supplied to induce DAB2 and 1 mM IPTG for *can*.

Pre-cultures were inoculated into 5 ml of M9 glycerol with 30 µg/ml kanamycin, 12.5 µg/ml

chloramphenicol and appropriate induction. 1 ml of each culture was transferred to a separate tube,

which was incubated in ambient CO_2 as a negative control; the remaining 4 ml was incubated in 10% CO_2.

CCMB1 strains carrying active site mutants of rubisco (cbbL K194M) were pre-cultured in LB with the same

antibiotic concentrations. Once cultures reached saturation they were centrifuged at 4000 g for 8 min.

Pellets were washed in 10 ml of M9 with no carbon source (M9 NoC) and resuspended in 5 ml M9 NoC.

Optical density was measured in five-fold dilution at 600 nm (Genesys 20 spectrophotometer, Thermo

Scientific) and cultures were normalized to 0.5 OD units. 96-well plates were inoculated by adding 2.5 µl

of OD-normalized preculture to 247.5 µl M9 glycerol supplemented with appropriate antibiotics and

induction. Kanamycin was omitted as the plasmid carrying kanamycin resistance also expresses rubisco,

which is required for growth in glycerol media (4). To minimize evaporation during multi-day cultivations,

150 µl sterile water was added to the reservoirs between the wells and the plate was incubated inside of

a small humidity cassette (Tecan) with 3 ml sterile water added to each reservoir. Plates were incubated

with shaking for at least 4 days in a Tecan Spark plate reader configured to control the CO_2 concentration.

The humidity cassette was replenished after 48 hours. All experiments were performed in biological

quadruplicate and technical triplicate.

C. necator growth conditions

C. necator strains were grown in ambient CO_2 unless otherwise specified, and 200 µg/ml kanamycin was

added to select for plasmid retention. When rubisco-independent heterotrophic growth was desired, *C.*

necator was cultured in LB at 30 °C. For autotrophic growth experiments, strains were pre-cultured in 5

mL of LB media in 10% CO_2 in 25 mL tubes with 20 mm butyl stoppers sealed by aluminum crimping. Pre-

cultures were incubated for two days at 30 °C with 200 rpm shaking, washed three times in *Cupriavidus*

minimal media, and inoculated to an OD of 0.1 (550 nm) in minimal media in a 165 mL flask with a 20 mm

butyl stopper sealed by aluminum crimping. *Cupriavidus* minimal growth media contained 3.24 mM

$MgSO_4$, 0.42 mM $CaCl_2$, 33.5 mM NaH_2PO_4, 32.25 mM Na_2HPO_4, 2.6 mM K_2SO_4, 1 mM NaOH, 1.87 mM

NH_4Cl, and 1 mL/L of a 1000x trace minerals solution containing 480 mg/L $CuSO_4 \cdot 5H_2O$, 2.4 g/L

$ZnSO_4 \cdot 7H_2O$, 2.4 g/L $MnSO_4 \cdot H_2O$, and 15 g/L $FeSO_4 \cdot 7H_2O$ (pH 6.7). The flasks were evacuated and the

headspace was filled with 60% H_2, 10% O_2, and indicated concentrations of CO_2. The balance was

atmospheric air. Cells were grown for 48 hours at 30° C with 200 rpm shaking, and OD_{550} values were

taken using a Molecular Devices SpectraMax M2 spectrophotometer.

Acknowledgements

We dedicate this paper to the memory of Danny Salah Tawfik (z"l), a luminary to the scientific community and a dear friend and mentor to AIF. Danny's many studies of enzyme evolution taught us that evolutionary timescales are accessible in carefully-designed lab-scale experiments and his active cultivation of relationships that transcend generational, professional and cultural divides continues to inspire. We are also indebted to Arren Bar-Even (z"l) for formative conversations guiding this work. Many thanks to Darcy McRose, Elad Noor, and Renee Wang for extensive comments on the manuscript and to Cecilia Blikstad, Julia Borden, Vahe Galstyan, Josh Goldford, Ron Milo, Robert Nichols, Naiya Phillips, Noam Prywes, and Gabe Salmon for helpful input. This research was supported in part by an NSF Graduate Research Fellowship (to A.I.F), NSF Grant No. PHY-1748958, the Gordon and Betty Moore Foundation Grant No. 2919.02, and the Kavli Foundation (to A.I.F), the US Department of Energy (DE-SC00016240 to D.F.S.) and Royal Dutch Shell (Energy and Biosciences Institute project CW163755 to D.F.S. and S.S.).

References

1. S. G. Wildman, Along the trail from Fraction I protein to Rubisco (ribulose bisphosphate
carboxylase-oxygenase). *Photosynth. Res.* **73**, 243–250 (2002).

2. Y. M. Bar-On, R. Milo, The global mass and average rate of rubisco. *Proc. Natl. Acad. Sci. U. S. A.*
116, 4738–4743 (2019).

3. C. Bathellier, G. Tcherkez, G. H. Lorimer, G. D. Farquhar, Rubisco is not really so bad. *Plant Cell*
Environ. **41**, 705–716 (2018).

4. A. I. Flamholz, *et al.*, Functional reconstitution of a bacterial CO2 concentrating mechanism in E.
coli. *Elife* **9** (2020).

5. C. Iñiguez, *et al.*, Evolutionary trends in RuBisCO kinetics and their co-evolution with CO2
concentrating mechanisms. *Plant J.* **101**, 897–918 (2020).

6. G. Bowes, W. L. Ogren, R. H. Hageman, Phosphoglycolate production catalyzed by ribulose
diphosphate carboxylase. *Biochem. Biophys. Res. Commun.* **45**, 716–722 (1971).

7. A. I. Flamholz, *et al.*, Revisiting Trade-offs between Rubisco Kinetic Parameters. *Biochemistry* **58**,
3365–3376 (2019).

8. W. W. Fischer, J. Hemp, J. E. Johnson, Evolution of Oxygenic Photosynthesis. *Annu. Rev. Earth*
Planet. Sci. **44**, 647–683 (2016).

9. D. C. Catling, K. J. Zahnle, The Archean atmosphere. *Sci Adv* **6**, eaax1420 (2020).

10. R. A. Berner, A. C. Lasaga, R. M. Garrels, The carbonate-silicate geochemical cycle and its effect on
atmospheric carbon dioxide over the past 100 million years. *Am. J. Sci.* **283**, 641–683 (1983).

11. K. H. Freeman, J. M. Hayes, Fractionation of carbon isotopes by phytoplankton and estimates of
ancient CO 2 levels. *Global Biogeochem. Cycles* **6**, 185–198 (1992).

12. C. R. Witkowski, J. W. H. Weijers, B. Blais, S. Schouten, J. S. Sinninghe Damsté, Molecular fossils
from phytoplankton reveal secular Pco2 trend over the Phanerozoic. *Sci. Adv.* **4**, eaat4556 (2018).

13. A. Flamholz, P. M. Shih, Cell biology of photosynthesis over geologic time. *Curr. Biol.* **30**, R490–R494
(2020).

14. Y. Marcus, R. Schwarz, D. Friedberg, A. Kaplan, High CO2 Requiring Mutant of Anacystis nidulans R2.
Plant Physiol. **82**, 610–612 (1986).

15. J. Pierce, T. J. Carlson, J. G. Williams, A cyanobacterial mutant requiring the expression of ribulose
bisphosphate carboxylase from a photosynthetic anaerobe. *Proc. Natl. Acad. Sci. U. S. A.* **86**, 5753–
5757 (1989).

16. N. M. Mangan, A. Flamholz, R. D. Hood, R. Milo, D. F. Savage, pH determines the energetic
efficiency of the cyanobacterial CO2 concentrating mechanism. *Proc. Natl. Acad. Sci. U. S. A.* **113**,
E5354–62 (2016).

17. B. D. Rae, B. M. Long, M. R. Badger, G. D. Price, Functions, compositions, and evolution of the two
types of carboxysomes: polyhedral microcompartments that facilitate CO2 fixation in cyanobacteria
and some proteobacteria. *Microbiol. Mol. Biol. Rev.* **77**, 357–379 (2013).

18. G. D. Price, M. R. Badger, Expression of Human Carbonic Anhydrase in the Cyanobacterium
Synechococcus PCC7942 Creates a High CO2-Requiring Phenotype Evidence for a Central Role for
Carboxysomes in the CO2 Concentrating Mechanism. *Plant Physiol.* **91**, 505–513 (1989).

19. L. Whitehead, B. M. Long, G. D. Price, M. R. Badger, Comparing the in Vivo Function of α-
Carboxysomes and β-Carboxysomes in Two Model Cyanobacteria. *Plant Physiol.* **165**, 398–411
(2014).

20. T. Ogawa, T. Kaneda, T. Omata, A Mutant of Synechococcus PCC7942 Incapable of Adapting to Low
CO2 Concentration. *Plant Physiol.* **84**, 711–715 (1987).

21. G. D. Price, M. R. Badger, Isolation and characterization of high CO2-requiring-mutants of the
cyanobacterium Synechococcus PCC7942: two phenotypes that accumulate inorganic carbon but
are apparently unable to generate CO2 within the carboxysome. *Plant Physiol.* **91**, 514–525 (1989).

22. J. J. Desmarais, *et al.*, DABs are inorganic carbon pumps found throughout prokaryotic phyla. *Nat*
Microbiol **4**, 2204–2215 (2019).

23. P. D. Tortell, Evolutionary and ecological perspectives on carbon acquisition in phytoplankton.
Limnol. Oceanogr. **45**, 744–750 (2000).

24. E. G. Nisbet, *et al.*, The age of Rubisco: the evolution of oxygenic photosynthesis. *Geobiology* **5**,
311–335 (2007).

25. J. N. Young, R. E. M. Rickaby, M. V. Kapralov, D. A. Filatov, Adaptive signals in algal Rubisco reveal a
history of ancient atmospheric carbon dioxide. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **367**, 483–492
(2012).

26. B. M. Long, B. Förster, S. B. Pulsford, G. D. Price, M. R. Badger, Rubisco proton production can drive
the elevation of CO2 within condensates and carboxysomes. *Proc. Natl. Acad. Sci. U. S. A.* **118**
(2021).

27. K. S. Smith, C. Jakubzick, T. S. Whittam, J. G. Ferry, Carbonic anhydrase is an ancient enzyme
widespread in prokaryotes. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 15184–15189 (1999).

28. S. D. Axen, O. Erbilgin, C. A. Kerfeld, A taxonomy of bacterial microcompartment loci constructed by
a novel scoring method. *PLoS Comput. Biol.* **10**, e1003898 (2014).

29. M. Krupovic, V. V. Dolja, E. V. Koonin, Origin of viruses: primordial replicators recruiting capsids
from hosts. *Nat. Rev. Microbiol.* **17**, 449–458 (2019).

30. N. Mangan, M. Brenner, Systems analysis of the CO2 concentrating mechanism in cyanobacteria.
Elife, e02043 (2014).

31. J. M. Shively, F. Ball, D. H. Brown, R. E. Saunders, Functional organelles in prokaryotes: polyhedral
inclusions (carboxysomes) of Thiobacillus neapolitanus. *Science* **182**, 584–586 (1973).

32. K. M. Wetmore, *et al.*, Rapid quantification of mutant fitness in diverse bacteria by sequencing
randomly bar-coded transposons. *MBio* **6**, e00306–15 (2015).

33. N. M. Wheatley, C. D. Sundberg, S. D. Gidaniyan, D. Cascio, T. O. Yeates, Structure and identification
of a pterin dehydratase-like protein as a ribulose-bisphosphate carboxylase/oxygenase (RuBisCO)
assembly factor in the α-carboxysome. *J. Biol. Chem.* **289**, 7973–7981 (2014).

34. O. Mueller-Cajar, The Diverse AAA+ Machines that Repair Inhibited Rubisco Active Sites. *Front Mol*
Biosci **4**, 31 (2017).

35. J. S. MacCready, L. Tran, J. L. Basalla, P. Hakim, A. G. Vecchiarelli, The McdAB system positions α-
carboxysomes in proteobacteria. *Mol. Microbiol.* **116**, 277–297 (2021).

36. K. M. Scott, *et al.*, Diversity in CO2-Concentrating Mechanisms among Chemolithoautotrophs from
the Genera Hydrogenovibrio, Thiomicrorhabdus, and Thiomicrospira, Ubiquitous in Sulfidic Habitats
Worldwide. *Appl. Environ. Microbiol.* **85**, 1–19 (2019).

37. T. G. Cooper, D. Filmer, The active species of "CO2" utilized by ribulose diphosphate carboxylase. *J.*
Biol. Chem. **244**, 1081–1083 (1969).

38. C. A. Kerfeld, M. R. Melnicki, Assembly, function and evolution of cyanobacterial carboxysomes.
Curr. Opin. Plant Biol. **31**, 66–75 (2016).

39. A. Pohlmann, *et al.*, Genome sequence of the bioplastic-producing "Knallgas" bacterium Ralstonia
eutropha H16. *Nat. Biotechnol.* **24**, 1257–1262 (2006).

40. M. R. Badger, E. J. Bek, Multiple Rubisco forms in proteobacteria: their functional significance in
relation to CO2 acquisition by the CBB cycle. *J. Exp. Bot.* **59**, 1525–1541 (2008).

41. J. Panich, B. Fong, S. W. Singer, Metabolic Engineering of Cupriavidus necator H16 for Sustainable
Biofuels from CO2. *Trends Biotechnol.* (2021).

42. C. S. Gai, J. Lu, C. J. Brigham, A. C. Bernardi, A. J. Sinskey, Insights into bacterial CO2 metabolism
revealed by the characterization of four carbonic anhydrases in Ralstonia eutropha H16. *AMB*
Express **4**, 2 (2014).

43. H. G. Schlegel, H. Kaltwasser, G. Gottschalk, Ein Submersverfahren zur Kultur
wasserstoffoxydierender Bakterien: Wachstumsphysiologische Untersuchungen. *Arch. Microbiol.*
38, 209–222 (1961).

44. W. Ahrens, H. G. Schlegel, Carbon dioxide requiring mutants of Hydrogenomonas eutropha strain H
16. I. Growth and Co2-fixation. *Arch. Mikrobiol.* **85**, 142–152 (1972).

45. M. Aragno, "Thermophilic, Aerobic, Hydrogen-Oxidizing (Knallgas) Bacteria" in *The Prokaryotes: A*
Handbook on the Biology of Bacteria: Ecophysiology, Isolation, Identification, Applications, A.
Balows, H. G. Trüper, M. Dworkin, W. Harder, K.-H. Schleifer, Eds. (Springer New York, 1992), pp.
3917–3933.

46. D. de Beer, A. Glud, E. Epping, M. Kûhl, A fast-responding CO2 microelectrode for profiling
sediments, microbial mats, and biofilms. *Limnol. Oceanogr.* **42**, 1590–1600 (1997).

47. Z. S. Brecheisen, C. W. Cook, P. R. Heine, J. Ryang, D. D. Richter, Development and deployment of a
field-portable soil O2 and CO2 gas analyzer and sampler. *PLoS One* **14**, e0220176 (2019).

48. N. J. Claassens, *et al.*, Phosphoglycolate salvage in a chemolithoautotroph using the Calvin cycle.
Proc. Natl. Acad. Sci. U. S. A. **117**, 22452–22461 (2020).

49. B. Kusian, D. Sültemeyer, B. Bowien, Carbonic Anhydrase Is Essential for Growth of Ralstonia
eutropha at Ambient CO2 Concentrations. *J. Bacteriol.* **184**, 5018–5026 (2002).

50. J. Gutknecht, M. A. Bisson, F. C. Tosteson, Diffusion of carbon dioxide through lipid bilayer
membranes: effects of carbonic anhydrase, bicarbonate, and unstirred layers. *J. Gen. Physiol.* **69**,
779–794 (1977).

51. C. Hannesschlaeger, A. Horner, P. Pohl, Intrinsic Membrane Permeability to Small Molecules. *Chem.
Rev.* **119**, 5922–5953 (2019).

52. A. C. Chakrabarti, D. W. Deamer, Permeability of lipid bilayers to amino acids and phosphate. *BBA -
Biomembranes* **1111**, 171–177 (1992).

53. T.-X. Xiang, B. D. Anderson, The relationship between permeant size and permeability in lipid
bilayer membranes. *J. Membr. Biol.* **140**, 111–122 (1994).

54. R. Milo, What is the total number of protein molecules per cell volume? A call to rethink some
published values. *Bioessays* **35**, 1050–1055 (2013).

55. C. Merlin, M. Masters, S. McAteer, A. Coulson, Why is carbonic anhydrase essential to Escherichia
coli? *J. Bacteriol.* **185**, 6415–6424 (2003).

56. J. Aguilera, J. P. Van Dijken, J. H. De Winde, J. T. Pronk, Carbonic anhydrase (Nce103p): an essential
biosynthetic enzyme for growth of Saccharomyces cerevisiae at atmospheric carbon dioxide
pressure. *Biochem. J.* **391**, 311–316 (2005).

57. P. Burghout, *et al.*, Carbonic anhydrase is essential for Streptococcus pneumoniae growth in
environmental ambient air. *J. Bacteriol.* **192**, 4054–4062 (2010).

58. G. P. Gladstone, P. Fildes, G. M. Richardson, Carbon Dioxide as an Essential Factor in the Growth of
Bacteria. *Br. J. Exp. Pathol.* **16**, 335 (1935).

59. H. G. Wood, C. H. Werkman, A. Hemingway, A. O. Nier, MECHANISM OF FIXATION OF CARBON
DIOXIDE IN THE KREBS CYCLE. *J. Biol. Chem.* **139**, 483–484 (1941).

60. H. A. Krebs, Carbon Dioxide Assimilation in Heterotrophic Organisms. *Nature* **147**, 560–563 (1941).

61. A. Lwoff, J. Monod, Essai d'analyse du role de l'anhydride carbonique dans la croissance
microbienne. *Ann. Inst. Pasteur* (1947).

62. J. M. Park, T. Y. Kim, S. Y. Lee, Genome-scale reconstruction and in silico analysis of the Ralstonia
eutropha H16 for polyhydroxyalkanoate synthesis, lithoautotrophic growth, and 2-methyl citric acid
production. *BMC Syst. Biol.* **5**, 101 (2011).

63. H. Knoop, *et al.*, Flux Balance Analysis of Cyanobacterial Metabolism: The Metabolic Network of

Synechocystis sp. PCC 6803. *PLoS Comput. Biol.* **9** (2013).

64.   K. M. Hines, V. Chaudhari, K. N. Edgeworth, T. G. Owens, M. R. Hanson, Absence of carbonic
anhydrase in chloroplasts affects C3 plant development but not photosynthesis. *Proc. Natl. Acad.*
Sci. U. S. A. **118** (2021).

65.   M. Krupovic, E. V. Koonin, Cellular origin of the viral capsid-like bacterial microcompartments. *Biol.*
Direct **12**, 25 (2017).

66.   L. V. Berkner, L. C. Marshall, N.A.S. Symposium on the Evolution of the Earth's Atmosphere:
HISTORY OF MAJOR ATMOSPHERIC COMPONENTS. *Proc. Natl. Acad. Sci. U. S. A.* **53**, 1215 (1965).

67.   J. F. Kasting, Faint young Sun redux. *Nature* **464**, 687–689 (2010).

68.   M. T. Rosing, D. K. Bird, N. H. Sleep, C. J. Bjerrum, No climate paradox under the faint early Sun.
Nature **464**, 744–747 (2010).

69.   Z. Liu, W. Dreybrodt, H. Liu, Atmospheric CO2 sink: Silicate weathering or carbonate weathering?
Appl. Geochem. **26**, S292–S294 (2011).

70.   D. B. McNevin, *et al.*, Differences in carbon isotope discrimination of three variants of D-ribulose-
1,5-bisphosphate carboxylase/oxygenase reflect differences in their catalytic mechanisms. *J. Biol.*
Chem. **282**, 36068–36076 (2007).

71.   E. B. Wilkes, A. Pearson, A general model for carbon isotopes in red-lineage phytoplankton:
Interplay between unidirectional processes and fractionation by RubisCO. *Geochim. Cosmochim.*
Acta (2019) https://doi.org/10.1016/j.gca.2019.08.043.

72.   P. M. Shih, *et al.*, Biochemical characterization of predicted Precambrian RuBisCO. *Nat. Commun.* **7**,
10382 (2016).

73.   C. T. Supuran, Carbonic Anhydrases and Metabolism. *Metabolites* **8** (2018).

772