Metabolic influence of core ciliates within the rumen microbiome

Thea O. Andersen[1], Ianina Altshuler[1], Arturo V.P. de Leon[2], Juline Walter[2], Emily McGovern[3], Kate Keogh[3], Cécile Martin[4], Laurence Bernard[4], Diego P. Morgavi[4], Tansol Park[5,6], Zongjun Li[7], Yu Jiang[7], Jeffrey L. Firkins[6], Zhongtang Yu[6], Torgeir R. Hvidsten[2], Sinead M. Waters[3], Milka Popova[4], Magnus Ø. Arntzen[2], Live H. Hagen[2], Phillip B. Pope[1,2*]

[1] Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian University of Life Sciences, Ås, Norway

[2] Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås, Norway

[3] Teagasc, Animal and Bioscience Research Department, Animal and Grassland Research and Innovation Centre, Teagasc, Grange, Dunsany, County Meath, Ireland

[4] Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, F-63122 Saint-Genes-Champanelle, France

[5] Department of Animal Science and Technology, Chung-Ang University, Anseong-si, Gyeonggi-do 17546, Republic of Korea

[6] Department of Animal Sciences, The Ohio State University, Columbus, OH, United States

[7] Center for Ruminant Genetics and Evolution, College of Animal Science and Technology, Northwest A&F University, Yangling, 712100, China

* Correspondence: phil.pope@nmbu.no

ABSTRACT

Protozoa comprise a major fraction of the microbial biomass in the rumen microbiome, of which the entodiniomorphs (order: *Entodiniomorphida*) and holotrichs (order: *Vestibuliferida*) are consistently observed to be dominant across a diverse genetic and geographical range of ruminant hosts. Despite the apparent core role that protozoal species exert, their major biological and metabolic contributions to rumen function remain largely undescribed *in vivo*. Here, we have leveraged (meta)genome-centric metaproteomes from rumen fluid samples originating from both cattle and goats fed diets with varying inclusion levels of lipids and starch, to detail the specific metabolic niches that protozoa occupy in the context of their microbial co-habitants. Initial proteome estimations via total protein counts and label-free quantification highlight that entodiniomorph species *Entodinium* and *Epidinium* as well as the holotrichs *Dasytricha* and *Isotricha* comprise an extensive fraction of the total rumen metaproteome. Proteomic detection of protozoal metabolism such as hydrogenases (*Dasytricha*, *Isotricha, Epidinium, Enoploplastron*), carbohydrate-active enzymes (*Epidinium, Diplodinium, Enoploplastron*, *Polyplastron*), microbial predation (*Entodinium*) and volatile fatty acid production (*Entodinium* and *Epidinium*) was observed at increased levels in high methane-emitting animals. Despite certain protozoal species having well-established reputations for digesting starch, they were

[1]

unexpectedly less detectable in low methane emitting-animals fed high starch diets, which were

instead dominated by propionate/succinate-producing bacterial populations suspected of being

resistant to predation irrespective of host. Finally, we reaffirmed our abovementioned observations

in geographically independent datasets, thus illuminating the substantial metabolic influence that

under-explored eukaryotic populations have in the rumen, with greater implications for both digestion

and methane metabolism.

Keywords: Protozoa, entodiniomorphs, holotrichs, metaproteomics, multi-omics, methane, rumen

BACKGROUND

Ruminants operate in symbiosis with their intrinsic rumen microbiome, which is responsible for the

degradation of forage into nutrients, in the form of volatile fatty acids (VFAs), supplying ~70% of net

energy for the host[1]. The rumen microbiome itself is a complex assemblage of bacterial, fungal,

archaeal, viral, and protozoal microorganisms whose intricate composition and function is connected

to host productivity traits, such as feed efficiency, milk yield, animal health, and greenhouse gas (GHG)

emissions[2-5]. Large collaborative research efforts have been made to identify and characterize the core

rumen microbiome including creating a publicly available catalogue for cultivated and sequenced

genomes[6-8]. In the rumen, bacteria are estimated to constitute 50-90 %, protozoa 10-50 %, fungi 5-

10% and archaea less than 4% of the total microbial biomass[9,10]. Due to the difficulties of axenically

culturing rumen eukaryotic populations and their complex genomic features that are obstinate to

current metagenomic assembly and binning technologies, the reconstruction of the rumen

microbiome has been heavily biased towards bacterial and archaeal members, whereas the fungal and

protozoal contributions of the rumen currently remain poorly characterized. While anaerobic fungi

have a reputable role as fibre degraders in the rumen, only 18 anaerobic gut fungi from herbivores

are currently described, with only 11 genomes available[11-13]. Similarly, to date few rumen protozoal

genomes are sequenced and publicly available, chief among them, the rumen ciliate protozoa

Entodinium caudatum[14]. More recently, single-cell amplified genomes (SAGs) have been recovered

from rumen microbiome samples representative of 5 holotrich and 14 entodiniomorph species

spanning 13 genera including *Isotricha*, *Dasytricha*, *Diplodinium*, *Enoploplastron*, *Metadinium*,

Eremoplastron, *Ostracodinium*, *Polyplastron*, *Ophryoscolex*, *Epidinum* and *Entodinium*[15].

Entodinium and *Epidinium* are entodiniomorphs of the order *Entodiniomorphida* and represent two

of the most dominant genera of rumen protozoa, previously being detected in more than 99% of 592

rumen samples at a mean protozoal relative abundance of ~ 38% and 16% respectively (2015 rumen

census: 32 animal species, 35 countries)[16]. Entodiniomorphs have been previously observed to

stimulate methane production[17], but while some possess hydrogenosomes such as *Epidinium*

caudatum[18], it is believed the numerically dominant *Entodinium caudatum* does not, and instead

encodes mitosomes and iron hydrogenases that may indicate hydrogen production, beneficial for

methanogenic endosymbionts[19]. Holotrich species of the order *Vestibuliferida* such as *Dasytricha* and

Isotricha are generally found at lower abundances compared to entodiniomorphs but are renowned

for their fermentation of soluble carbohydrates and their production of VFAs and hydrogen[20,21]. Here,

we present a genome-centric metaproteomics analysis of the rumen microbiome from two different

host species; Holstein dairy cattle (*Bos taurus*) and alpine goats (*Capra hircus*), that were fed diets of

first-cut grassland hay with a 45:55 forage:concentrate ratio, with concentrates supplemented with

either no additional lipid (CTL), or corn oil and wheat starch (COS)[5,22,23]. Previous microscopy analysis

revealed high counts of entodiniomorphs and to a lesser extent holotrichs, while metadata revealed

that animals fed COS displayed reduced methane emissions, irrespective of host. To describe how

these diets affect digestion and production of methane and VFAs, we sought to investigate changes in

function and composition in the complex rumen microbiome. By using shotgun metagenomic

sequencing we recovered in total 244 prokaryote metagenome-assembled genomes (MAGs) that

together with selected isolate- and SAG-derived eukaryote genomes[14,15,24-29] formed the database for

our genome-centric metaproteomic analysis of rumen protozoal species. Key protozoal metabolisms

were detected related to plant fiber and starch degradation, bacterial predation, and hydrogen and

VFA production. Despite numerous protozoal species having the genetic ability to degrade starch, our

analysis showed contrasting data that suggests rumen protozoa were less metabolically active in the

rumen microbiome of animals fed a starch-rich diet. In such a scenario other starch-degrading and/or

propionate and succinate producing bacterial genera, as *Prevotella* and *Fibrobacter* and members of

the families *Succinivibrionaceae* and *Aminobacteriaceae* appeared to be more prevalent. In concert,

our analysis showed that reduced methane production in both cattle and goats eating feeds

supplemented with COS is likely caused via a redirection of hydrogen to succinate and propionate

production instead of methanogenesis. Finally, by analysing a secondary, geographically independent

dataset, we reaffirmed our primary observations of protozoal dominance and patterns concerning

hydrogen, plant fiber and starch-related metabolism, thus supporting our hypothesis that this

protozoal species plays a core role in rumen microbiome function.

METHODS

Animal trial and sample handling

The experimental procedures were approved by the Auvergne-Rhône-Alpes Ethics Committee for

Experiments on Animals (France; DGRI agreement APAFIS#3277–2015121411432527 v5) and

complied with the European Union Directive 2010/63/EU guidelines. Experiments were performed at

the animal experimental facilities of HerbiPôle site de Theix at the Institut National de la Recherche

pour l'Agriculture, l'Alimentation l'Environnement (INRAE, Saint-Genès-Champanelle, France) from

February to July 2016. Experimental design, animals and diets were as described elsewhere[5,23]. Briefly,

four Holstein cows and four Alpine goats, all lactating, were enrolled in respectively two 4 x 4 Latin

square design trials to study the effects of 4 diets over four 28-d experimental periods (**Supplementary**

Table S1). The original study included four different experimental grassland hay basal diets with

concentrates supplemented with various lipid sources; control diet with no added lipids (CTL), diet

supplemented with corn oil and wheat starch (COS), diet supplemented with marine algae powder

(MAP) and diet supplemented with hydrogenated palm oil (HPO) (**Supplementary Table S2**)[23]. In the

present study, we focused on the CTL and COS diets, which were associated with the most extreme

methane (CH_4) emission profiles in both ruminant species. The CTL diet composed of grass hay *ad*

libitum with concentrates containing no additional lipid, whereas COS contained corn oil (5.0% total

dry matter intake (DMI)) and wheat starch -5.0 % of total DMI (COS) (**Table 1**). Corn oil (Olvea, Saint

Léonard, France) was added to the concentrate, at 5% of total DMI and contained (g/kg of total FA):

16:0 (114), 18:0 (16.4), cis-9 18:1 (297), cis-11 18:1 (6.30), 18:2n-6 (535), 18:3n-3 (7.57), 20:0 (3.48),

22:0 (1.0), 24:0 (1.5), and total FA (1000 g/kg). Detailed diet composition is available in Martin *et al.*[5].

Each experimental period lasted for 28 days. Diets were offered as two equal meals at 0830 and 1600h.

Animals had access to a constant supply of freshwater *ad libitum*. Concentrate and hay refusals were

weighed daily. The amounts of feed offered the following day was adjusted regarding to refusals to

maintain the targeted the dietary 45 % dry matter (DM) forage and 55 % DM concentrate ratio. We

acknowledge that by only including one sampling point for rumen fluid we are reducing the power

that the original experimental Latin square design gives regarding minimizing individual error between

the animals.

Rumen fluid was collected through stomach-tubing before the morning feeding on day 27 of each

experimental period. The stomach tube consisted of a flexible 23 mm external diameter PVC hose

fitted to a 10 cm-strainer at the head of the probe for cattle, and a flexible 15 mm PVC hose with a 12

cm-strainer for goats. The first 200 ml of rumen fluid was discarded from to minimize contamination

from saliva. Samples were filtered through a polyester monofilament fabric (280 μm pore size),

dispatched in 2-ml screw-cap tubes, centrifuged at 15000 x g for 10 mins and the pellet snap-frozen

in liquid nitrogen. Samples were stored at -80°C until DNA extraction using the Yu and Morrison bead-

beating procedure[30]. In total, 32 rumen fluid samples (four cows and four goats fed four diets included

in the original study[23]) were sent to the Norwegian University of Life Sciences (NMBU) for

metagenomic and metaproteomic analysis. Respiration chambers were used to measure methane

emissions over a 5-day period, while VFA and NH_3 concentrations were determined by gas

chromatography using a flame ionization detector[31]. Protozoa were counted by microscopy and

categorized as either small entodiniomorphs (<100 μm), large entodiniomorphs (>100 μm) or as

holotrichs *Dasytricha* and *Isotricha*[9]. Further specifics about diets and measurements can be found in

Martin *et al.*[5] and VFA and methane measurements are summarized in **Table 1**.

Metagenomic sequencing and analysis

Metagenomic shotgun sequencing was performed at the Norwegian Sequencing Centre on two lanes

of the HiSeq 3/4000 (Illumina) generating 150 bp paired-end reads in both lanes. Sequencing libraries

were prepared using the TruSeq DNA PCR-Free High Throughput Library Prep Kit (Illumina) prior to

sequencing. All 32 samples (four cows and four goats fed four diets included in the original study[23])

were run on both lanes to prevent potential lane-to-lane sequencing bias. FASTQ files were quality

filtered and Illumina adapters removed using Trimmomatic[32] (v. 0.36) with parameters -phred33 for

base quality encoding, leading and trailing base threshold set to 20. Sequences with an average quality

score below 15 in a 4-base sliding window were trimmed and the minimum length of reads was set to

36 bp. MEGAHIT[33] (v.1.2.9) was used to co-assemble reads originating from samples collected from

cattle and goats separately, with options –kmin-1pass, --k-list 27,37,47,57,67,77,87, --min-contig-len

1000 in accordance with[7]. Bowtie2[34] (v. 2.3.4.1) was used to map reads back to the assemblies and

SAMtools[35] (v. 1.3.1) was used to convert SAM files to BAM format and index sorted BAM files.

The two co-assemblies (one from the samples originating from cattle and the other originating from

the samples of goats) were binned using Maxbin2[36], MetaBAT2[37] and CONCOCT[38]. MetaBAT2 (v.

2.12.1) was run using parameters –minContig 2000 and –numThreads 4, Maxbin2 (v. 2.2.7) ran with

default parameters and -thread 4, min_contig_length 2000, and CONCOCT (v. 1.1.0) ran with default

parameters and –length_threshold 2000. Further, bins were filtered, dereplicated and aggregated

using DASTool[39](v. 1.1.2) with the parameters –write_bins 1, --threads 2 and BLAST[40] as search engine.

This resulted in a total of 244 dereplicated MAGs across the two host species (104 originating from

cattle and 140 from goats). CheckM[43](v. 1.1.3) lineage workflow was used to calculate completeness

and contamination of each MAG, with parameters –threads 8, --extension fa, and CoverM (v. 0.5.0)

(https://github.com/wwood/CoverM) was used to calculate relative abundance of each MAG, while

GTDB-tk[44,45] (v. 1.3.0) was used for taxonomic annotation. Approximately 90% (219 of 244) of the

recovered MAGs were considered high or medium quality MAGs according to MIMAGs threshold for

completeness and contamination for genome reporting standards[46] (**Supplementary Table S3**). Gene

calling and functional annotation of the final MAGs were performed using the DRAM[47] pipeline with

the databases dbCAN[48], Pfam[49], Uniref90[50], Merops[51], VOGdb and KOfam[52]. The translated amino acid

sequences from the publicly available drafted *En. caudatum* macronucleus genome were annotated

with the KEGG metabolic pathway database using BlastKOALA[53] by Park *et al.*[14]. Proteins with resulting

KEGG Orthology (KO) numbers were functionally assigned to metabolic pathways using KEGG Mapper

Reconstruct Tool[54].

The resulting protein sequences for each MAG, as well as those from the host genomes of goat (*Capra*

hircus, NCBI ID: 10731) and cattle (*Bos taurus*, NCBI ID: 82) were compiled into two databases, from

now on referred to as sample specific RUmen DataBase for Goat (RUDB-G) and sample specific RUmen

DataBase for Cattle (RUDB-C). In addition, both databases were supplemented with the genome of

the protozoa *Entodinium caudatum*[14], as well as 18 SAGs from Li et al.[15] including *Dasytricha*

ruminantium SAG3, *Diplodinium dentatum* SAGT1, *Diplodinium flabellum* SAG1, *Enoploplastron*

triloricatum SAGT1, *Entodinium bursa* SAG3, *Entodinium longinucleatum* SAG4, *Epidinium cattanei*

SAG3, *Epidinium caudatum* SAG1, *Eremoplastron rostratum* SAG2, *Isotricha intestinalis* SAGT2,

Isotricha prostoma SAG3, *Isotricha* sp YL-2021a SAG1, *Isotricha* sp YL-2021b SAG3, *Metadinium*

minomm SAG1, *Ophryoscolex caudatus* SAGT3, *Ostracodinium dentatum* SAG1, *Ostracodinium gracile*

SAG1 and *Polyplastron multivesiculatum* SAGT3. The genome of *Fibrobacter succinogenes* S85 (NCBI

ID: 932) was also added to both rumen databases since it is well recognized as a primary cellulolytic

bacterium in the rumen microbiome and has previously been observed as an active microorganism in

similar studies yet was not thoroughly binned as a unique MAG in this study. Nine available fungal

genomes downloaded from Joint Genome Institute (JGI) Mycocosm[55] were added to RUDB-C

(*Anaeromyces* sp. S4[25], *Caecomyces churrovis*[28], *Neocallimastix californiae*[25], *Neocallimastix lanati*[29],

Piromyces finnis[25], *Piromyces* sp. E2[25], *Piromyces* UH3-1[24,25,27], *Piromyces* eukMAG[56], *Orpinomyces*

sp.[26]). In total the complete databases consisted of 1 291 083 and 1 190 938 protein entries for RUDB-

G and RUDB-C, respectively.

Metaproteomic data generation

To 300 μL of rumen fluid sample (in total 32 rumen fluid samples originating from four cows and four

goats fed four diets included in the original study[23]) 150 μL lysis buffer (30 mM DTT, 150 mM Tris-HCl

(pH=8), 0.3% Triton X-100, 12% SDS) was added together with 4 mm glass beads (≤ 160 μm), followed

by brief vortexing and resting on ice for 30 mins. Lysis was performed using FastPrep-24 Classic Grinder

(MP Biomedical, Ohio, USA) for 3 × 60 seconds at 4.0 meter/ second[57]. Samples were centrifuged at

16 000 × g for 15 minutes at 4°C and lysate was carefully removed. Protein concentration was

measured using the Bio-Rad DC™ Protein Assay (Bio-Rad, California USA) with bovine serum albumin

as standard. Absorbance of sample lysates was measured at A750 on BioTek Synergy H4 Hybrid

Microplate Reader (Thermo Fisher Scientific Inc., Massachusetts, USA). 40-50 µg of protein was

prepared in SDS-buffer, heated in water bath for 5 minutes at 99°C and analysed by SDS-PAGE using

Any-kD Mini-PROTEAN TGX Stain-Free gels (Bio-Rad, California, USA) in a 2-minute run for sample

clean-up purposes, before staining with Coomassie Blue R-250. The visible bands were carefully

excised from the gel and divided as 1×1 mm pieces before reduction, alkylation, and digestion with

trypsin. Peptides were concentrated and eluted using C18 ZipTips (Merck Millipore, Darmstadt,

Germany) according to manufacturer's instructions, dried, and analysed by nano-LC-MS/MS using a

Q-Exactive hybrid quadrapole Orbitrap MS (Thermo Fisher Scientific Inc., Massachusetts, USA) as

previously described[58]

Metaproteomic data analysis

Mass spectrometry (MS) raw data were analysed with FragPipe version 19 and searched against the

sample-specific protein sequence database (1 291 083 and 1 190 938 protein entries for RUDB-G and

RUDB-C, respectively) with MSFragger[59] (**Supplementary Table S4**). The database was supplemented

with contaminant protein entries, such as human keratin, trypsin, and bovine serum albumin, in

addition to reversed sequences of all protein entries for estimation of false discovery rates (FDR)[60].

Oxidation of methionine and protein N-terminal acetylation were used as variable modifications, while

carbomidomethylation of cysteine residues were used as fixed modification. Trypsin was chosen as

digestive enzyme, maximum missed cleavages allowed was one and matching tolerance levels for both

MS and MS/MS were 20 ppm. The results were filtered to 1% FDR and quantification was done using

IonQuant[61] including normalization between samples and the feature 'match between runs' to

maximize protein identifications. Perseus[62] version 1.6.2.3 was used for further analysis. A protein

group was considered valid if it was quantified in at least 50% of the replicates in at least one

condition. Protein groups identified as potential contaminants were removed. Calculations of

individual genome/SAG/MAG contributions were calculated using the workflow outlined by Kleiner et

al.[63], which sums protein abundances (label free quantification (LFQ) values) for all proteins assigned

to each genome/SAG/MAG, and differential abundance between diets were detected by an unpaired

two-sided Student's t-test (p-value <0.05).

After filtration, we resolved 3657 unique protein groups across the 16 samples from cattle and 3818

unique protein groups across 15 samples from goats (**Supplementary Table S5** and **Supplementary**

Table S6). Dot plots for **Figure 1** were made with ggplot2[64] in R (v. 4.2.0)[65]. To determine which

expressed metabolic pathways *En. caudatum* were significantly enriched for in each diet/animal, we

used the hypeR() function from the hyperR package[66] in R which employs the hypergeometric test for

significance. The 'geneset' for hyperR was generated by using the keggGet() functionof KEGGREST R

package to retrieve entries from the KEGG database and determine which pathways the *En. caudatum*

KOs belong to. The geneset was then manually curated to only include metabolic pathways of interest

(i.e., we remove pathways such as "Huntington disease"). For the 'background' setting in hyperR(), to

be conservative, we used the total number of unique KOs (7592) in the *En. caudatum* genome that

could possibly be expressed. Significantly enriched KOs with *p*-value < 0.05 were manually filtered.

Both *p*-values and FDR-adjusted *p*-values are available in **Supplementary Table S7**, with FDR- adjusted

p-values < 0.05 highlighted in bold.

Animal trial, sample handling and metagenomic data generation for independent validating dataset

Samples were also analysed from previously performed feeding experiments with Holstein Friesian

bulls[67]. In brief, these bulls were subjected to either *ad libitum* or restricted feeding regime in a

compensatory growth model detailed in Keogh *et al.*, 2015[67]. Both feeding groups received the same

ratio of concentrate and grass silage, respectively 70% and 30%, of which the concentrate was mainly

composed of starch-rich rolled barley (72.5%) and soya (22.5%). Rumen samples were collected at

slaughter and stored at -80C prior to metagenomics and metaproteomic analysis in this study.

Sample preparation, cell lysis and extraction of DNA was carried out as previously described by

McCabe *et al.*[68] Quality check of fastq files and removal of low-quality reads was performed using fastp

(V.0.19.5). Sequence reads were mapped against the bovine genome (ARS-UCD1.3) using minimap2

(V.2.16), and host sequences were removed. Reads were co-assembled using Megahit (V1.2.6) with

"−meta-large" pre-set option as the metagenome was complex. Metagenomic binning was applied to

the co-assembly using MetaBAT2 using standard parameters (V.2.12.1). MAGs were then dereplicated

using dRep (V.1.4.3), and the resulting MAGs were taxonomically assigned using Bin Annotation Tool

(BAT), available on (https://github.com/dutilh/CAT). This tool internally uses prodigal (V.2.6.3) for

gene prediction and DIAMOND (V.0.9.14) for the alignment against the non-redundant (nr) protein

database (as of Feb 2020).

Sample preparation for metaproteomics was done by lysing cells with bead beating with two glass
bead sizes (≤106 μm and 0.5 mm), in 100 mM Tris, pH8, 5% SDS and 10 mM DTT. A FastPrep 24
instrument was operated for 3 × 45 seconds at a speed of 6.5 m/s. The samples were centrifuged for
15 minutes at 20.000 × g and the protein extracts were cleaned by Wessel-Flügge precipitation[69];
pellets were dissolved in 5% SDS, 100 mM Tris-Cl, pH8, 10 mM DTT and kept at -20 $^{\circ}$C until further
processing. Protein digestion was performed using suspension trapping (STrap)[70], dried in a SpeedVac
(Eppendorf Concentrator Plus) and re-dissolved in 0.05 % trifluoroacetic acid, 2% acetonitrile for
peptide concentration estimation using a Nanodrop One instrument, and subsequent MS/MS-
analysis. The samples were analyzed using an Ultimate3000 RSLCnano UHPLC coupled to a QExactive
hybrid quadrupole-orbitrap mass spectrometer (Thermo Fisher, Bremen, Germany) as described
previously[58]. MS raw data were analysed with FragPipe and searched against the Holstein Friesian
bulls sample-specific protein sequence database (2 533 362 protein entries) using the same workflow
described above.
285
RESULTS AND DISCUSSION
Protozoal populations have large proteomes in the rumen microbiome
Because of their large size protozoal species can comprise a significant fraction of the microbial
biomass in the rumen[9]. While the total count and diversity of protozoal species are lesser than their
bacterial counterparts in the rumen, their genome size and total number of genes encoded within are
considerably larger. In addition, due to alternative splicing and post-translational modifications, the
protein representation, or proteome, of protozoal populations will be larger than the number of genes
in their protozoal genome[71]. Thus, the amount of protein in a protozoa species can be expected to far
exceed the amount of protein in a bacterial species that can be identified and quantified in proteomic
studies. In this context, our metaproteomic data showed an extensive fraction of detectable proteins
affiliated to protozoal species, and other closely related species, in both dairy cattle and goats in
proportion to the combined bacterial species that were represented in our genome databases (**Figure**
1a). Protein detection intensity of proteins assigned to protozoa were also proportionally greater than
that of the bacterial fraction of the rumen microbiome in animals fed the control (CTL) diet (**Figure 1b**)
further supporting the significant levels of protozoal detection in our samples. Proteomes from
entodiniomorph species were detected at high levels in both animal hosts, with *Ep. cattanei* and *En.*
caudatum dominating the rumen metaproteome of cattle and goats respectively (**Figure 2**). In cattle
we also detected entodinomorphs *D. dentatum*, *En. bursa* and *Ep. caudatum* (**Figure 2a**,
Supplementary Table S6) at high levels whereas in goats *Entodinium* species *En. caudatum*, *En.*
longinucleatum and *En. bursa* where among the highest detected entodinomorphs (**Figure 2c**). More

than three times as many proteins from *En. caudatum* were detected in goats than cattle, however
this was somewhat expected given the 7x higher counts of entodiniomorph concentration (cells/mL)
previously observed in the goat samples compared to cattle (**Table 1**)[5]. In agreement with microscopic
cell counts of holotrichs, we detected proteins from both *Dasytricha* spp. and *Isotricha* spp (**Table 1**)
in both cattle and goats (**Figure 2a and 2c**).

Entodiniomorphs and holotrichs have influential roles in rumen carbohydrate metabolism
Rumen protozoa have long been characterized as degraders of plant fibers as well as non-structural
storage polysaccharides such as starch[72]. Closer examination of the proteomes from both highly
detected *Epidinium species Ep. cattanei* and *Ep. caudatum* in cattle highlighted a plethora of detected
carbohydrate active enzymes (CAZymes) targeting lignocellulosic polysaccharides such as cellulose
(GH5, GH9), xylan (GH10, GH11, GH30, CE1), mannan (GH26) and fiber-derived oligosaccharides (GH2,
GH3, GH94) (**Figure 2a**, **Supplementary Table S6**). In goats similar CAZy families were detected but
from different entodinomorphs species, such as *E. triloricatum* and *P. multivesiculatum* (**Figure 2c**,
Supplementary Table S6). Our *in vivo* CAZyme observations were in strong agreement to long standing
in vitro data, which have shown entodinomorphs such as *Epidinium*, *Polyplastron* and *Enoploplastron*
have greater endoglucanase and xylanase activity, while *Entodinium* spp. have only weak activity[72].
Similarly, it has previously been shown *in vitro* that holotrichs such as *Dasytricha* spp. have glucosidase
and cellobiosidase activity but negligible fibrolytic activity[73], which our *in vivo* data also supported
(**Figure 2a and 2c**). The composition of protozoal species metabolising starch (both degrading and
synthesizing) also varied between the two animal hosts with goats dominated by *Entodinium* spp. and
D. ruminantium while protozoal starch metabolism in cattle was *Epidinium* spp., *D. ruminantium and*
Isotricha spp (**Figure 2**, **Supplementary Table S6**). While bacterial CAZymes targeting lignocellulose
were also detected (e.g. GH48 cellulases from *Ruminococcaceae* populations in cattle), CAZymes
targeting lignocellulose and starch were detected at higher frequency and protein quantification levels
in protozoal species suggesting they have a central role in ruminal fibre digestion.

Metabolism of En. caudatum shows predatory activity and metabolism of VFAs
En. caudatum is renowned for its predatory activity and is acknowledged as the most abundant
protozoa in the rumen, whereby it has been estimated that 0.1% of rumen prokaryotes are digested
by the rumen protozoal population every minute[74]. The dominance of *En. caudatum* in our data
presented an opportunity to better understand the metabolic influence of this universal species. While
previous efforts have investigated the genome and transcriptome of *En. caudatum* grown in
monoculture[14,75], our metaproteomic analysis sought to reveal *in vivo* metabolism and functions of *En.*

caudatum within the rumen microbiome. In accordance with Wang *et al.*[75], our metaproteomic

analysis revealed expressed proteins attributed to metabolic pathways such as carbon metabolism,

glycolysis/gluconeogenesis, starch and sucrose (and glycogen) metabolism, pyruvate metabolism,

oxidative phosphorylation and production of alcohol (**Supplementary Table S6**). Wang *et al.* found

that as for most rumen microbes, *En. caudatum* uses carbohydrates such as starch as its primary

substrate, as well as cellulose and hemicellulose to a certain degree[75], and their transcript analyses

showed that *En. caudatum* had high levels of expression of amylases and low-level expression of

hemicellulases, cellulases and pectinases. Similarly, our metaproteomic analysis reveals expression of

amylases by *En. caudatum* that are predicted to enable *En. caudatum* to engulf and degrade starch

granules to simpler sugars and to produce glycogen, its most important storage carbohydrate[76].

However, no detection of *En. caudatum* CAZymes related to hemicellulose or pectin were observed in

any of our metaproteomes, suggesting that it is not engaging in the deconstruction of these

carbohydrates at the time our samples were collected for analysis (before feeding). It should be noted

that ruminal fermentation activity as well as production of VFAs and methane will be at its highest

after feeding, as a result of an increased availability of fermentable substrate[77]. While sampling time

can influence the recovered microbial composition and hence function, any differences in metabolic

parameters or species abundance in this study is comparable across both diets given the consistent

sampling times.

While monoculture cultures of *En. caudatum* have not been established to verify the VFAs it can

produce, Wang *et al.* found transcripts of enzymes involved in fermentative formation of acetate and

butyrate[75]. Similarly, we detected proteins inferred in metabolism of acetate, butyrate, and alcohol in

En. caudatum. Goats fed the CTL diet had a significantly higher proportion of proteins from *En.*

caudatum and concurrently had increased relative levels of acetate and butyrate compared to animals

fed the corn oil and wheat starch diet (COS), which had fewer *En. caudatum* proteins and lower

acetate/butyrate levels (**Figure 2** and **Table 1**). As *En. caudatum* populations were seemingly most

abundant in goats fed the CTL diet, we used these metaproteomes to reconstruct metabolic features

(**Figure 3**). Of the 538 *En. caudatum* proteins identified in CTL-fed goats, 244 had unique KO numbers

assigned, from which KEGG Mapper reconstructions[54] enabled functional assignment of 217 proteins

to metabolic pathways. Our metabolic reconstructions showed expressed proteins involved in

endocytosis, phagosome and lysosome processes for predatory activity, engulfment, and digestion of

bacteria (**Figure 3**, **Supplementary Table S6**). For the rumen samples used in this study Martin *et al.*

previously observed higher NH_3 concentrations in goats compared to cattle and hypothesised that it

might have resulted from increased bacterial protein breakdown and feed protein degradability due

to higher density of entodiniomorphs known for their predatory activity[5]. In support of these

observations, we performed metaproteomic pathway enrichment analysis of *En. caudatum* (**Figure 3,**

Supplementary Table S7) proteins detected in goats, which revealed significantly enriched

endocytosis metabolism. Other biological processes such as glycolysis and starch and sucrose

metabolism were significantly enriched pathways in *En. caudatum* (**Supplementary Table S7**).

Although suspected of having metabolic interactions with methanogenic archaea, several protozoal

populations such as *Entodinium* are hypothesized as having associations with certain members of the

Gram-negative *Gammaproteobacteria*, which multiple studies have speculated are resistant to

protozoal engulfment[78-80]. In contrast, Gutierrez and Davis previously demonstrated that *Entodinium*-

species engulf Gram positive starch-degraders[80]. In the context of our data, we speculate that CTL fed

animals provided *En. caudatum*-like populations optimal conditions for predation, whereas increased

starch levels in the COS diets facilitated increases in Gram-negative species that are possibly resistant

to protozoal engulfment and/or reduced pH levels, leading to sub-optimal conditions for *Entodinium*

and a decrease in its levels (**Figure 2**).

Protozoa were less active in diets supplemented with starch regardless of their starch metabolising

reputations.

The changes in VFA and methane levels in animals fed the high starch COS diet, previously measured

by Martin *et al.*[5], suggested significant alterations in composition and thus functions of the rumen

microbiome irrespective of host species. In particular, a decrease in proportions of acetate and

butyrate, decrease in the acetate:propionate ratio and an increase in relative propionate levels were

observed in animals fed the COS diet, compared to the CTL diet (**Table 1**). Diets that are high in starch

content or with low forage:concentrate ratios have previously been shown to result in higher

production of propionate and succinate, as they are easily fermented in the rumen and accordingly

have high passage rates[81,82]. We therefore leveraged our genome-centric metaproteomic data from

both cattle (**Figure 2a-b)** and goats (**Figure 2c-d**) fed either the COS or CTL diet to gain an overview of

protein expression from individual populations. We specifically focused on pathways involved in the

degradation of starch (CTL: corn starch, COS: corn + wheat starch) to pyruvate through glycolysis and

finally formation of acetate, butyrate, and propionate (via succinate). Irrespective of host, and despite

their starch-metabolising reputation[75,76], *Entodinium*, *Epidinium*, *Isotricha* and *Dasytricha* spp had

lower abundance and less proteins involved in starch degradation in animals fed the COS diet

compared to those fed the CTL diet (**Figure 2a** and **2c**). Further, we observed opposing patterns for

proteins identified as *Entodinium* and *Epidinium*-spp. involved in glycolysis, and production of

[12]

pyruvate, acetate, and butyrate, which were detected in higher levels in both cattle and goats fed the CTL diet compared to the COS diet.

While several putative protozoal amylases were detected across all animals and diets, their quantification levels (i.e., protein abundances) did not increase as expected when higher levels of starch were available (**Figure 2a** and **2c**). We therefore hypothesized that the observed shift in VFA profiles in response to increased starch was additionally influenced by the bacterial fraction of the rumen microbiome. In contrast to lower levels of protozoa in the animals fed the COS diet, we observed an increase in bacterial proteome detection irrespective of host (**Figure 4**), including suspected starch-degrading and succinate- and propionate-producing bacterial species (**Figure 2b** and **2d**). For example, starch fermentation pathways from population genomes affiliated with the *Succinivibrionaceae* family, *Prevotella* species and, additionally for goats, members of the *Aminobacteriaceae* families, were detected at higher proteomic levels in the animals fed the COS diet compared to those fed the CTL diet (**Figure 2b** and **2d**).

Protozoa are less active in animals that produce lower methane yield

For the animals sampled in this study, Martin *et al.* demonstrated a ~25-30% reduction in methane emissions in both cattle and goats fed the COS diet compared to the control (**Table 1**)[5]. Previous comparisons between defaunated and faunated animals have shown decrease in methane production in protozoa-free ruminants, suggesting symbiotic interactions between methanogenic archaea and protozoal species[78]. Methanogen's epi- and endo-symbiotic relationships with protozoa have also been suggested to contribute to 9-37% of rumen methanogenesis[78,83-85]. Moreover, studying microcosms with the presence and absence of protozoal species Solomon *et al.* reported higher levels of acetate and butyrate in microcosms with protozoa present in addition to increased methane emissions[78], which supports the main findings of animals fed the CTL diet in this study (**Table 1**). Several of the protozoal species detected in our metaproteomic data are known to have hydrogenosomes and closer examination showed that [FeFe]-hydrogenases from both holotrich and entodinomorph genera, such as *Dasytricha, Isotricha, Diplodinium, Epidinium, Entodinium* and *Enoploplastron*, were detected at higher quantification levels in animals fed the CTL compared to those fed COS (**Figure 2a and 2c**, **Supplementary Table S6**). We also observed partial evidence in our proteomic data that the hydrogenosome-lacking *En. caudatum* makes contributions to ruminal hydrogen production, supporting previous reports associating its abundance with higher methane levels[78,85,86]. Only one of its eight [FeFe]-hydrogenases were detected in goats (absent in cattle), though it showed no changes in protein abundance in either the high (CTL) or low (COS) methane

[13]

yielding animals. Given our observations, we speculate that predicted protozoal-methanogen

relationships that lead to increased methane levels in this study are centred on hydrogen transfer.

The addition of lipids in diets for ruminants have been largely studied as a methane mitigation

strategy[87,88]. However, it is widely believed that lipids are not fermented in the rumen and rather

modified/hydrogenated and hence do not contribute to hydrogen production or methane production

in the rumen. While biohydrogenation of unsaturated fatty acids such as corn oil can serve as a

hydrogen sink, it has been observed that very small amounts of metabolic hydrogen (1-2%) are used

for biohydrogenation[87,88]. Several studies have also shown that the CH_4 mitigation effects of lipids are

dependent on both dose and fatty acid composition[89,90]. For example, medium chain fatty acids

(MCFAs), such as lauric acid, have been proven to be a more effective inhibitor of protozoa than long

chain polyunsaturated fatty acids (PUFAs), such as corn oil, which mainly consists of linoleic acid[88-91].

In a study conducted by Zhang *et al.*[90], goats fed corn oil as a supplement decreased ruminal hydrogen

concentrations and total methane emissions. Nevertheless, there was seemingly no effect on rumen

protozoal populations, which suggests that the corn oil dosage used in the Zhang study did not act as

an anti-protozoal agent[90]. In addition, high levels of PUFAs in the diet will also likely impact the

metabolism of various bacterial and archaeal species[88-90], particularly keystone fibre degrading Gram-

positive bacterial species that produce various levels of VFAs and hydrogen[92,93]. Collectively, these

previous findings were in agreement with the decreased CH_4 production in cattle and goats in this

study fed the COS diet, which was observed to additionally impact other bacterial metabolism and

ruminal fermentation parameters, such as increased propionate and decreased butyrate and acetate

levels (**Table 1**)[5]. Thus, while we cannot unequivocally rule out that diets supplemented with lipids at

higher levels could be directly affecting the abundance of protozoa, we believe it is also possible that

corn oil is influencing other fibre-degrading and hydrogen-producing species in the rumen, hence

effecting ruminal fermentation and methane emissions[88,94]. Nonetheless, validation of these results

under different conditions and extended sampling is necessary to verify our findings.

Increases in dietary starch for ruminants is known to stimulate the propionate and succinate pathways

of starch-degrading bacteria, which due to their net incorporation of metabolic hydrogen [H]

represent a [H] sink in rumen fermentation besides hydrogenotrophic methanogenesis[83,95]. Diets rich

in starch are more fermentable in the rumen, which can decrease the ruminal pH to levels that can

inhibit methanogenic archaea and fibre-degrading bacterial species[96,97]. Yet, lowered pH levels in the

rumen can also lead to clinical (or sub-clinical in most production scenarios) ruminal acidosis[98,99].

Hence, high concentrate diets, which increase production of propionate at the expense of methane,

does not necessarily opt for a viable methane mitigation strategy in the long term. Our results suggests

that decreased methanogenesis in COS-fed animals is likely due to a decrease in available hydrogen

and/or decrease in pH levels, which we predict is caused by the lower levels of hydrogen-producing

protozoa and bacteria as well as the metabolism of dominant bacterial populations that likely do not

produce exogenous hydrogen due to their own [H]-utilizing succinate and propionate metabolism.

This prediction was partially supported by the detection of several putative hydrogenases believed to

use molecular hydrogen in putative fumarate-reducing *Aminobacteriaceae* populations that were

dominant in the COS-fed goat rumen. In particular, two [FeFe]-hydrogenases were detected in COS-

fed animals from RUDB_G023 (ORF_4456741_13) and RUDBG016 (ORF_ 1143158_19) that were

annotated as NADP-reducing hydrogenase subunits (EC:1.12.1.3) (**Figure 2d**, **Supplementary Table**

S6).

Protozoal dominance is validated in geographically independent datasets

To further test our hypothesis that protozoal species play a central role in the rumen ecosystem, we

explored additional metagenome-centric metaproteomic datasets originating from an independent

feeding experiment performed in Ireland on 60 Holstein Friesian bulls[67]. In brief, these bulls were

subjected to the same ratio of concentrate and grass silage at either an *ad libitum* or restricted feeding

regime in a compensatory growth model detailed in Keogh et al, 2015[67]. We applied the same strategy

as for the described Holstein dairy cattle and alpine goats to resolve the metaproteomic dataset for a

subset of 15 animals (7 restricted and 8 *ad libitum*) against 781 reconstructed sample-specific MAGs

(RUDB-HF), which were supplemented with the genome of *En. caudatum,* 18 protozoal SAGs[15] as well

as genomes of available anaerobic fungi. This collection of microbial prokaryote and eukaryote

genomes was then used as a sequence database for the generated protein spectra. Consistent with

our previous observation in dairy cattle, a substantial proportion of the detected proteins were

affiliated to *Epidinium* (*Ep. cattanei* and *Ep. caudatum*), *Entodinium* (*En. caudatum* and *En. bursa*) and

Isotricha spp. providing further support that protozoa are an important and metabolically active

contributor to the rumen microbiome (**Figure 5, Supplementary Table S8**). The protein quantification

(measured as sum of LFQ intensities affiliated to each MAG/genome, averaged for each diet) was also

higher in the rumen sample from bulls on the restricted diet, which likely had less starch available

compared to the *ad libitum* group and a higher retention time (**Figure 5a, Supplementary Table S8**).

A previously published 16S rRNA amplicon investigation of the phylogenetic differences between the

rumen microbiomes of these two diet groups highlighted an increase in *Succinivibrionaceae* in the

starch-rich *ad libitum* diet[68]. Our metaproteomic analysis confirmed a significantly higher (unpaired t-

test, *p*-value < 0.05) proteomic detection of several *Succinivibrionaceae*-MAGs under the *ad libitum*

group (**Supplementary Table S8**), accompanied with a reduced acetate:propionate ratio in the rumen,

which is often associated with increased feed efficiency and reduced production of methane[68]. These

observations largely mirror the dominance of *Succinivibrionaceae*-MAGs in the dairy cattle and goats

fed the COS diet, further strengthening our hypothesis that *Entodinium* and *Epidinium* spp do not

metabolically respond to increases in available starch in the host animals' diets and have other roles

than being a primary starch degrader.

To assess whether similar patterns of diet-induced microbiome shifts were consistent across the two

geographically distinct datasets used in this study, we compared ratios of protein abundance level for

commonly observed taxonomic families. Spearman's rank correlation coefficients were calculated for

the comparisons, where two tests were performed: correlation between common taxonomic families

(rarely observed families with < 2 MAGs with detected proteins across both conditions were excluded)

in the rumen of beef (RUDB_HF) and dairy (RUDB_C) cattle as well as beef cattle (RUDB_HF) and alpine

goats (RUDB_G), respectively. Changes in protein abundance levels from a total of 16 taxonomic

families were found in common between the rumen microbiomes of beef and dairy cattle (significant

correlation: *p*-value < 0.05 and correlation coefficient rho of 0.68), further strengthening our

hypothesis that similar structural rumen microbiome shifts occur across independent datasets of

cattle subjected to diets of high and low starch (**Figure S1, Supplementary Table S9**). However,

unsurprisingly no significant correlations were observed between the rumen microbiomes of goats

(RUDB_G) and beef cattle (RUDB_HF) (*p*-value > 0.05 and correlation coefficient rho of 0.19,

Supplementary Table S9, Supplementary Figure S1), despite similar trends in protein abundance

levels across taxa being observed (**Figure 2, Figure 5,** and **Supplementary Figure S1)**.

In conclusion, by using a (meta)genome-centric metaproteomics approach we primarily investigated

the role of the rumen protozoa in the rumen microbiome of beef and dairy cattle as well as dairy goats

that were subjected to varying dietary conditions. We showed that the proteomes of core

entodinomorph and holotrich genera such as *Entodinium*, *Epidinium*, *Dasytricha* and *Isotricha*

constitute a substantial fraction of the recovered rumen microbial proteome, which supports previous

16S/18S rRNA gene-based rumen census data that have highlighted their global dominance across a

plethora of ruminant species. In both animal hosts, protozoal CAZymes targeting lignocellulose were

detected though more frequently and at higher quantification levels in animals fed the CTL diet

compared to the COS supplemented diet, presumably due to lower cell density in the later (**Table 1**).

Proteins identified as *En. caudatum*, *Epidinium, Isotricha* and *Dasytricha* spp. were surprisingly

detected at lower levels in animals that were fed increased levels of wheat starch, despite their

[16]

reputable starch-metabolising capabilities (**Figures 2-5**). We hypothesize that protozoa were being out
competed by Gram-negative bacterial species (e.g. *Succinivibrionaceae* in cattle and/or
Aminobacteriaceae in goats) in the COS-fed animals, which were possibly resistant to protozoal
engulfment[78-80] and/or lower pH levels, creating sub-optimal conditions for starch-degrading
protozoal populations. We also observed lower detection of protozoal [FeFe]-hydrogenases in COS-
fed animals that measured a ~25-30% lower CH_4 yield at the time of sampling in this study (prior to
feeding), further suggesting the specific conditions that enable succinate- and propionate-producing
populations to flourish subsequently exert an impact on hydrogen and methane metabolisms in the
rumen microbiome. For animals that were additionally fed corn oil, it is acknowledged that diets
supplemented with lipids can have effects on the decreased abundance of protozoa, however our
study did not provide evidence of a direct effect. While much work is still needed to confirm our
abovementioned hypotheses, our integrated metaproteomics approaches have demonstrated the
future importance of including eukaryote populations for accurate and meaningful analyses of the
rumen microbiome and its impact on GHG mitigation strategies and host productivity traits.

Acknowledgements
PBP and TOA are grateful for support from The Research Council of Norway (FRIPRO program, PBP:
250479), as well as the European Research Commission Starting Grant Fellowship (awarded to PBP:
336355 - MicroDE), and the Novo Nordisk Foundation (awarded to PBP: 0054575 - SuPAcow). LHH was
supported by The Research Council of Norway (FRIPRO program, LHH: 302639 – SeaCow), while MØA
was supported by the Novo Nordisk Foundation, project No. NNF20OC006131. The experimental trial
was financed by APIS-GENE (Paris, France) as part of the NutriLip project. Gas emission measurements
were funded by UMR 1213 Herbivores (INRAE, Saint-Genès-Champanelle, France). The sequencing
service was provided by the Norwegian Sequencing Centre (www.sequencing.uio.no), a national
technology platform hosted by the University of Oslo and supported by the "Functional Genomics"
and "Infrastructure" programs of the Research Council of Norway and the Southeastern Regional
Health Authorities. The authors acknowledge the Orion High Performance Computing Center at the
Norwegian University of Life Sciences and Sigma2 - the National Infrastructure for High Performance
Computing and Data Storage in Norway for providing computational resources that have contributed
to meta-omics computations reported in this paper. Mass spectrometry-based proteomic analyses
were performed by The MS and Proteomics Core Facility, Norwegian University of Life Sciences
(NMBU). This facility is a member of the National Network of Advanced Proteomics Infrastructure
(NAPI), which is funded by the Research Council of Norway INFRASTRUKTUR-program (project
number: 295910).

The authors declare no conflicts of interest.

Data Availability

Raw shotgun metagenomic data has been deposited in the National Center for Biotechnology

Sequence Read Archive (NCBI-SRA) under accessions numbers SRR19524239 to SRR19524270 with

links to BioProject accession number PRJNA844951. All annotated prokaryote MAGs are available

publicly at DOI: 10.6084/m9.figshare.20066972.v1. The mass spectrometry proteomics data have

been deposited to the ProteomeXchange Consortium via the PRIDE[100] partner repository with the

dataset identifiers PXD040467, PXD040454 and PXD040349. NCBI accession numbers for the protozal

SAGs used in this study are as follows: *Dasytricha ruminantium* SAG3 (JAJJKK000000000), *Diplodinium*

dentatum SAGT1 (JAJJLX000000000), *Diplodinium flabellum* SAG1 (JAJJLB000000000), *Enoploplastron*

triloricatum SAGT1 (JAJNAB000000000), *Entodinium bursa* SAG3 (JAJJKR000000000), *Epidinium*

cattanei SAG3 (JAJJKG000000000), *Epidinium caudatum* SAG1 (JAJMZR000000000), *Eremoplastron*

rostratum SAG2 (JAJMZO000000000), *Isotricha intestinalis* SAGT2 (JAJJLT000000000), *Isotricha*

prostoma SAG3 (JAJMZS000000000), *Isotricha* sp. YL-2021a SAG1 (JAJJLH000000000), *Isotricha* sp. YL-

2021b SAG3 (JAJJKW000000000), *Metadinium minomm* SAG1 (JAJMZT000000000), *Ophryoscolex*

caudatus SAGT3 (JAJJLZ000000000), *Ostracodinium dentatum* SAG1 (JAJJKF000000000),

Ostracodinium gracile SAG1 (JAJJKZ000000000), *Polyplastron multivesiculatum* SAGT3

(JAJJLS000000000).

Data access for reviewers:

SRA:
https://dataview.ncbi.nlm.nih.gov/object/PRJNA844951?reviewer=7q1stta7ga42mq5tq27ifj7n85
601
602   Figshare   - https://figshare.com/s/cec40f220249f3ef848c
603   - https://figshare.com/s/ad4eb7e1b7db17f0535b
604
PRIDE - Reviewer must download PRIDE Inspector Tool
https://www.ebi.ac.uk/pride/markdownpage/prideinspector
•   MS Rawfiles for samples from CATTLE:
Username: reviewer_pxd040467@ebi.ac.uk
Password: KPtwrE0J
•   MS Rawfiles for samples from GOAT:
Username: reviewer_pxd040454@ebi.ac.uk
Password: FjGXlkSf
•   MS Rawfiles for samples from Holstein Friesian BULLS:
Username: reviewer_pxd040349@ebi.ac.uk
Password: I1rcyaFi
616

REFERENCES

Bergman, E. N. Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol Rev* **70**, 567-590 (1990).

Wallace, R. J. *et al.* The rumen microbial metagenome associated with high methane production in cattle. *BMC Genomics* **16**, 839 (2015).

Jami, E., White, B. A. & Mizrahi, I. Potential role of the bovine rumen microbiome in modulating milk composition and feed efficiency. *PLoS One* **9**, e85423 (2014).

McCann, J. C. *et al.* Induction of Subacute Ruminal Acidosis Affects the Ruminal Microbiome and Epithelium. *Front Microbiol* **7**, 701 (2016).

Martin, C. *et al.* Diets supplemented with corn oil and wheat starch, marine algae, or hydrogenated palm oil modulate methane emissions similarly in dairy goats and cows, but not feeding behavior. *Anim Feed Sci Technol* **272**, 114783 (2021).

Seshadri, R. *et al.* Cultivation and sequencing of rumen microbiome members from the Hungate1000 Collection. *Nat Biotechnol* **36**, 359-367 (2018).

Stewart, R. D. *et al.* Compendium of 4,941 rumen metagenome-assembled genomes for rumen microbiome biology and enzyme discovery. *Nat Biotechnol* **37**, 953-961 (2019).

Xie, F. *et al.* An integrated gene catalog and over 10,000 metagenome-assembled genomes from the gastrointestinal microbiome of ruminants. *Microbiome* **9**, 137 (2021).

Williams, A. G. & Coleman, G. S. in *The rumen microbial ecosystem*   73-139 (Springer, 1997).

Lin, C., Raskin, L. & Stahl, D. A. Microbial community structure in gastrointestinal tracts of domestic animals: comparative analyses using rRNA-targeted oligonucleotide probes. *FEMS Microbiol Ecol* **22**, 281-294 (1997).

Hagen, L. H. *et al.* Proteome specialization of anaerobic fungi during ruminal degradation of recalcitrant plant fiber. *ISME J* **15**, 421-434 (2021).

Saye, L. M. G. *et al.* The Anaerobic Fungi: Challenges and Opportunities for Industrial Lignocellulosic Biofuel Production. *Microorganisms* **9**, 694 (2021).

Stabel, M. *et al.* Aestipascuomyces dupliciliberans gen. nov, sp. nov., the First Cultured Representative of the Uncultured SK4 Clade from Aoudad Sheep and Alpaca. *Microorganisms* **8**, 1734 (2020).

Park, T., Wijeratne, S., Meulia, T., Firkins, J. L. & Yu, Z. The macronuclear genome of anaerobic ciliate *Entodinium caudatum* reveals its biological features adapted to the distinct rumen environment. *Genomics* **113**, 1416-1427 (2021).

Li, Z. *et al.* Genomic insights into the phylogeny and biomass-degrading enzymes of rumen ciliates. *ISME J* **16**, 2775–2787 (2022).

Henderson, G. *et al.* Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci Rep* **5**, 14567 (2015).

Ranilla, M. J., Jouany, J. P. & Morgavi, D. P. Methane production and substrate degradation by rumen microbial communities containing single protozoal species in vitro. *Lett Appl Microbiol* **45**, 675-680 (2007).

Yarlett, N., Coleman, G. S., Williams, A. G. & Lloyd, D. Hydrogenosomes in known species of rumen entodiniomorphid protozoa. *FEMS Microbiol Lett* **21**, 15-19 (1984).

Park, T., Meulia, T., Firkins, J. L. & Yu, Z. Inhibition of the Rumen Ciliate Entodinium caudatum by Antibiotics. *Front Microbiol* **8** (2017).

Belanche, A., de la Fuente, G. & Newbold, C. J. Study of methanogen communities associated with different rumen protozoal populations. *FEMS Microbiol Ecol* **90**, 663-677 (2014).

Hillman, D., Lloyd, K., Yarlett, N. & Williams, A. G. Hydrogen Production by Rumen Holotrich Protozoa: Effects of Oxygen and Implications for Metabolic Control by In Situ Conditions. *J Protozool* **36**, 205-213 (1989).

22 Fougere, H. & Bernard, L. Effect of diets supplemented with starch and corn oil, marine
algae, or hydrogenated palm oil on mammary lipogenic gene expression in cows and goats:
A comparative study. *J Dairy Sci* **102**, 768-779 (2019).
23 Fougere, H., Delavaud, C. & Bernard, L. Diets supplemented with starch and corn oil, marine
algae, or hydrogenated palm oil differentially modulate milk fat secretion and composition
in cows and goats: A comparative study. *J Dairy Sci* **101**, 8429-8445 (2018).
24 Solomon, K. V. *et al.* Early-branching gut fungi possess a large, comprehensive array of
biomass-degrading enzymes. *Science* **351**, 1192-1195 (2016).
25 Haitjema, C. H. *et al.* A parts list for fungal cellulosomes revealed by comparative genomics.
Nat Microbiol **2**, 17087 (2017).
26 Youssef, N. H. *et al.* The genome of the anaerobic fungus Orpinomyces sp. strain C1A reveals
the unique evolutionary history of a remarkable plant biomass degrader. *Appl Environ*
Microbiol **79**, 4620-4634 (2013).
27 Hooker, C. A. *et al.* Hydrolysis of untreated lignocellulosic feedstock is independent of S-
lignin composition in newly classified anaerobic fungal isolate, Piromyces sp. UH3-1.
Biotechnol Biofuels **11**, 293 (2018).
28 Brown, J. L. *et al.* Cocultivation of the anaerobic fungus Caecomyces churrovis with
Methanobacterium bryantii enhances transcription of carbohydrate binding modules,
dockerins, and pyruvate formate lyases on specific substrates. *Biotechnol Biofuels* **14**, 234
(2021).
29 Wilken, S. E. *et al.* Experimentally Validated Reconstruction and Analysis of a Genome-Scale
Metabolic Model of an Anaerobic Neocallimastigomycota Fungus. *mSystems* **6**, e00002-
00021 (2021).
30 Yu, Z. & Morrison, M. Improved extraction of PCR-quality community DNA from digesta and
fecal samples. *Biotechniques* **36**, 808-812 (2004).
31 Morgavi, D. P., Jouany, J.-P. & Martin, C. Changes in methane emission and rumen
fermentation parameters induced by refaunation in sheep. *Aust J Exp Agric* **48**, 69-72 (2008).
32 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114-2120 (2014).
33 Li, D., Liu, C. M., Luo, R., Sadakane, K. & Lam, T. W. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics **31**, 1674-1676 (2015).
34 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
357-359 (2012).
35 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079 (2009).
36 Wu, Y. W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning algorithm to
recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605-607 (2016).
37 Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome
reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).
38 Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat Methods*
11, 1144-1146 (2014).
39 Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a dereplication,
aggregation and scoring strategy. *Nat Microbiol* **3**, 836-843 (2018).
40 Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res* **36**, W5-9 (2008).
41 Hyatt, D., LoCascio, P. F., Hauser, L. J. & Uberbacher, E. C. Gene and translation initiation site
prediction in metagenomic sequences. *Bioinformatics* **28**, 2223-2230 (2012).
42 Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood and
Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics **11**, 538 (2010).

43   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043-1055 (2015).

44   Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat Biotechnol* **36**, 996-1004 (2018).

45   Parks, D. H. *et al.* A complete domain-to-species taxonomy for Bacteria and Archaea. *Nat*
Biotechnol **38**, 1079-1086 (2020).

46   Bowers, R. M. *et al.* Minimum information about a single amplified genome (MISAG) and a
metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol* **35**, 725-
731 (2017).

47   Shaffer, M. *et al.* DRAM for distilling microbial metabolism to automate the curation of
microbiome function. *Nucleic Acids Res* **48**, 8883-8900 (2020).

48   Zhang, H. *et al.* dbCAN2: a meta server for automated carbohydrate-active enzyme
annotation. *Nucleic Acids Res* **46**, W95-W101 (2018).

49   Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**, D412-D419
(2021).

50   Suzek, B. E. *et al.* UniRef clusters: a comprehensive and scalable alternative for improving
sequence similarity searches. *Bioinformatics* **31**, 926-932 (2015).

51   Rawlings, N. D. *et al.* The MEROPS database of proteolytic enzymes, their substrates and
inhibitors in 2017 and a comparison with peptidases in the PANTHER database. *Nucleic Acids*
Res **46**, D624-D632 (2018).

52   Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and
adaptive score threshold. *Bioinformatics* **36**, 2251-2252 (2020).

53   Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG Tools for
Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol* **428**, 726-
731 (2016).

54   Kanehisa, M., Sato, Y. & Kawashima, M. KEGG mapping tools for uncovering hidden features
in biological data. *Protein Sci* **31**, 47-53 (2022).

55   Nordberg, H. *et al.* The genome portal of the Department of Energy Joint Genome Institute:
2014 updates. *Nucleic Acids Res* **42**, D26-31 (2014).

56   Peng, X. *et al.* Genomic and functional analyses of fungal and bacterial consortia that enable
lignocellulose breakdown in goat gut microbiomes. *Nat Microbiol* **6**, 499-511 (2021).

57   Mićić, M., Whyte, J. D. & Karsten, V. in *Sample Preparation Techniques for Soil, Plant, and*
Animal Samples (ed Miodrag Micic) 99-116 (Springer New York, 2016).

58   Michalak, L. *et al.* Microbiota-directed fibre activates both targeted and secondary metabolic
shifts in the distal gut. *Nat Commun* **11**, 5773 (2020).

59   Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I.
MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry–based
proteomics. *Nat. Methods* **14**, 513-520 (2017).

60   da Veiga Leprevost, F. *et al.* Philosopher: a versatile toolkit for shotgun proteomics data
analysis. *Nat Methods* **17**, 869-870 (2020).

61   Yu, F., Haynes, S. E. & Nesvizhskii, A. I. IonQuant Enables Accurate and Sensitive Label-Free
Quantification With FDR-Controlled Match-Between-Runs. *Mol Cell Proteomics* **20**, 100077
(2021).

62   Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of
(prote)omics data. *Nat Methods* **13**, 731-740 (2016).

63   Kleiner, M. *et al.* Assessing species biomass contributions in microbial communities via
metaproteomics. *Nat Commun* **8**, 1558 (2017).

64   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (2016).

65   R Core Team. R: A language and environment for statistical computing. (2013).

66 Federico, A. & Monti, S. hypeR: an R package for geneset enrichment workflows.
Bioinformatics **36**, 1307-1308 (2020).
67 Keogh, K., Waters, S. M., Kelly, A. K. & Kenny, D. A. Feed restriction and subsequent
realimentation in Holstein Friesian bulls: I. Effect on animal performance; muscle, fat, and
linear body measurements; and slaughter characteristics. *J Anim Sci* **93**, 3578-3589 (2015).
68 McCabe, M. S. *et al.* Illumina MiSeq Phylogenetic Amplicon Sequencing Shows a Large
Reduction of an Uncharacterised Succinivibrionaceae and an Increase of the
Methanobrevibacter gottschalkii Clade in Feed Restricted Cattle. *PLOS ONE* **10**, e0133234
(2015).
69 Wessel, D. & Flugge, U. I. A method for the quantitative recovery of protein in dilute solution
in the presence of detergents and lipids. *Anal Biochem* **138**, 141-143 (1984).
70 Zougman, A., Selby, P. J. & Banks, R. E. Suspension trapping (STrap) sample preparation
method for bottom-up proteomics analysis. *Proteomics* **14**, 1006-1000 (2014).
71 Harper, J. W. & Bennett, E. J. Proteome complexity and the forces that drive proteome
imbalance. *Nature* **537**, 328-338 (2016).
72 Williams, A. G. & Coleman, G. S. *The Rumen Protozoa*. (1992).
73 Takenaka, A., Tajima, K., Mitsumori, M. & Kajikawa, H. Fiber Digestion by Rumen Ciliate
Protozoa. *Microbes Environ* **19**, 203-210 (2004).
74 Coleman, G. S. & Sandford, D. C. The engulfment and digestion of mixed rumen bacteria and
individual bacterial species by single and mixed species of rumen ciliate protozoa grown *in*
vivo. *J Agric Sci* **92**, 729-742 (1979).
75 Wang, L. *et al.* The transcriptome of the rumen ciliate *Entodinium caudatum* reveals some of
its metabolic features. *BMC Genomics* **20**, 1008 (2019).
76 Belzecki, G., McEwan, N. R., Kowalik, B., Michalowski, T. & Miltko, R. Effect of Entodinium
caudatum on starch intake and glycogen formation by Eudiplodinium maggii in the rumen
and reticulum. *Eur J Protistol* **57**, 38-49 (2017).
77 Allen, M. S. Relationship between fermentation acid production in the rumen and the
requirement for physically effective fiber. *J Dairy Sci* **80**, 1447-1462 (1997).
78 Solomon, R. *et al.* Protozoa populations are ecosystem engineers that shape prokaryotic
community structure and function of the rumen microbial ecosystem. *ISME J* **16**, 1187-1197
(2022).
79 Park, T. & Yu, Z. Do Ruminal Ciliates Select Their Preys and Prokaryotic Symbionts? *Front*
Microbiol **9**, 1710 (2018).
80 Gutierrez, J. & Davis, R. E. Bacterial Ingestion by the Rumen Ciliates Entodinium and
Diplodinium. *J Protozool* **6**, 222-226 (1959).
81 Bauman, D. E., Davis, C. L. & Bucholtz, H. F. Propionate production in the rumen of cows fed
either a control or high-grain, low-fiber diet. *J Dairy Sci* **54**, 1282-1287 (1971).
82 Jiao, H. P. *et al.* Effect of concentrate feed level on methane emissions from grazing dairy
cows. *J Dairy Sci* **97**, 7043-7053 (2014).
83 Martin, C., Morgavi, D. P. & Doreau, M. Methane mitigation in ruminants: from microbe to
the farm scale. *Animal* **4**, 351-365 (2010).
84 Finlay, B. J. *et al.* Some rumen ciliates have endosymbiotic methanogens. *FEMS Microbiol*
Lett **117**, 157-161 (1994).
85 Newbold, C. J., Lassalas, B. & Jouany, J. P. The importance of methanogens associated with
ciliate protozoa in ruminal methane production in vitro. *Lett Appl Microbiol* **21**, 230-234
(1995).
86 Newbold, C. J., de la Fuente, G., Belanche, A., Ramos-Morales, E. & McEwan, N. R. The Role
of Ciliate Protozoa in the Rumen. *Front Microbiol* **6**, 1313 (2015).
87 Hristov, A. N. *et al.* Special topics--Mitigation of methane and nitrous oxide emissions from
animal operations: I. A review of enteric methane mitigation options. *J Anim Sci* **91**, 5045-
5069 (2013).

88   Jenkins, T. C., Wallace, R. J., Moate, P. J. & Mosley, E. E. Board-invited review: Recent
advances in biohydrogenation of unsaturated fatty acids within the rumen microbial
ecosystem. *J Anim Sci* **86**, 397-412 (2008).
89   Martin, C., Morgavi, D. P. & Doreau, M. Methane mitigation in ruminants: from microbe to
the farm scale. *Animal* **4**, 351-365 (2010).
90   Zhang, X. M. *et al.* Corn oil supplementation enhances hydrogen use for biohydrogenation,
inhibits methanogenesis, and alters fermentation pathways and the microbial community in
the rumen of goats. *J Anim Sci* **97**, 4999-5008 (2019).
91   Guyader, J. *et al.* Influence of rumen protozoa on methane emission in ruminants: a meta-
analysis approach. *Animal* **8**, 1816-1825 (2014).
92   Niu, P. *et al.* A Basic Model to Predict Enteric Methane Emission from Dairy Cows and Its
Application to Update Operational Models for the National Inventory in Norway. *Animals*
(Basel) **11**, 1891 (2021).
93   McAllister, T. A., Cheng, K.-J., Okine, E. K. & Mathison, G. W. Dietary, environmental and
microbiological aspects of methane production in ruminants. *Can J Anim Sci* **76**, 231-243
(1996).
94   Gruninger, R. J. *et al.* Application of 3-nitrooxypropanol and canola oil to mitigate enteric
methane emissions of beef cattle results in distinctly different effects on the rumen
microbial community. *Anim Microbiome* **4**, 35 (2022).
95   Popova, M. *et al.* Effect of fibre- and starch-rich finishing diets on methanogenic Archaea
diversity and activity in the rumen of feedlot bulls. *Anim Feed Sci Technol* **166-167**, 113-121
(2011).
96   Van Kessel, J. A. S. & Russell, J. B. The effect of pH on ruminal methanogenesis. *FEMS*
Microbiol Ecol **20**, 205-210 (1996).
97   Russell, J. B., Muck, R. E. & Weimer, P. J. Quantitative analysis of cellulose degradation and
growth of cellulolytic bacteria in the rumen. *FEMS Microbiol Ecol* **67**, 183-197 (2009).
98   Russell, J. B. The importance of pH in the regulation of ruminal acetate to propionate ratio
and methane production in vitro. *J Dairy Sci* **81**, 3222-3230 (1998).
99   Kleen, J. L., Hooijer, G. A., Rehage, J. & Noordhuizen, J. P. Subacute ruminal acidosis (SARA):
a review. *J Vet Med A Physiol Pathol Clin Med* **50**, 406-414 (2003).
100 Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019:
improving support for quantification data. *Nucleic Acids Res* **47**, D442-D450 (2019).

Table 1. Summary of the animal host, dietary conditions and accompanying metadata that are linked to
the rumen samples used to explore protozoal functions. Effects of control diet (CTL) and diet
supplemented with corn oil and wheat starch (COS) on average VFA concentration in the percentage of
total VFA concentration (mmol/l) and average methane yield (CH_4 g/kg DMI) in addition to average total
protozoa cell counts and average cell counts for small entodiniomorph protozoa for cattle (n=4) and goats
(n=4) (**Supplementary Table S2**)[5]. Total protozoal cell counts, and cell counts of entodiniomorphs and
holotrichs were based on ciliate protozoa morphology in microscopy[5]. Measurements on VFA
concentrations and methane yield were determined by gas chromatography and respiration chambers,
respectively.

Animal	Cattle		Goat	
Diet	CTL	COS	CTL	COS
VFA sum mmol/l	61.8	68.3	33.7	25.3
Acetate (% sum)	72.0	69.1	65.7	64.2
Propionate (% sum)	14.2	21.3	15.6	19.1
Butyrate (% sum)	10.8	6.4	13.7	9.9
CH_4 g/kg dry matter intake	19.5	14.5	19.5	13.5
Standard deviation	2.2	2.9	4.0	4.1
Total protozoa (10^3 cells/ml)	92	100	1382	898
Small entodiniomorphs (<100µm) (10^3 cells/ml)	51	97	1346	898
Large entodiniomorphs (>100µm) (10^3 cells/ml)	25.1	1.7	4.9	0.2
Holotrichs				
Isotricha spp	4.1	1.0	4.6	0.2
Dasytricha spp	11.8	0.1	26.7	0.2

Figure legends

Figure 1. Quantities of identified protozoal proteins in the rumen microbiome vary depending on

host animal and dietary conditions. Dot plots for total proteins identified (**a**), and average recovered

metaproteomic expression (**b:** presented as proportion of summed LFQ intensities) belonging to

protozoal, bacterial, archaeal, or fungal species across the control diet (CTL) and diets supplemented

with corn oil and wheat starch (COS) for dairy cattle (n=4) and goats (n=4). Detected protein

abundances for protozoal and bacterial populations can be found in **Supplementary Table S6**.

Figure 2. Detected proteins mapped to protozoal genomes/SAGs and bacterial metagenome-

assembled genomes (MAGs) in the rumen microbiome of dairy cattle (n=4) and goats (n=4) fed

either a control diet (CTL) or one supplemented with corn oil and wheat starch (COS). The figure

displays metabolically active populations (as genomes, SAGs or MAGs), with selected expressed

proteins (presented as Enzyme Commission (EC) number with short enzyme descriptions) active in

starch degradation, glycolysis and production of pyruvate, butyrate, acetate, and succinate in cattle
(**a-b**) and goats (**c-d**) fed CTL or COS diets. Panels **a** and **c** depict protozoal proteomes that were
detected in cattle and goats respectively are presented separately to bacteria (panels **b** and **d**) as the
scale of their protein quantification values were ~10x larger. Protein quantification values (y-axis)
were calculated by considering both the number of proteins detected per MAG/SAG/genome and
their LFQ intensity: we averaged LFQ intensities for each detected protein across biological replicates
for each dietary condition (CTL: green or COS: orange), which were subsequently summed for all
detected proteins per MAG/SAG/genome. Heatmaps show selected MAG/SAG/genome with
metabolically active proteins, presented as EC numbers, recovered from cattle (RUDB-C) and goats
(RUDB-G) fed either CTL or COS diets. Where average protein abundance level for a given
MAG/SAG/genome was significantly different (determined by paired Wilcoxon test, *p*-value < 0.05)
between the COS and CTL diet, the given MAG/SAG/genome has been marked with a red star. MAGs
are presented with their MAG ID and taxonomic annotation from GTDB-tk. Genome annotations and
LFQ intensities used to create heatmaps can be found in **Supplementary Table S6**.

Figure 3. Reconstructed phagolysosome formation and starch metabolism of *Entodinium caudatum*
within the rumen microbiome of goats fed the control diet (CTL) based on metaproteomic analysis.
KO identifiers for identified proteins were analysed via KEGG mapper to reconstruct expressed key
features in the metabolism of *En. caudatum*. Dashed arrows represent proteins or pathways that
were not detected in our metaproteomes but are key steps in their respective pathways. Detailed
information connecting KO identifiers to their respective gene ID, LFQ and animal/diet can be found
in **Supplementary Table S6**.

Figure 4. Volcano plots indicating different rumen microbiome proteins from dairy cattle (a) and
goats (b) fed either the control or COS diets and which displayed both large magnitude of fold-
changes in LFQ intensities (x axis) and high statistical significance (-log10 of nominal *p*-values using
an unpaired t-test, y axis). Dashed horizontal line denotes a *p*-value <0.05 cut-off. For both animal
hosts we observed an increase in protein detection for bacterial populations (greenblue) in animals
fed high-starch diets compared to protozoal populations (purple), which were detected at higher LFQ
intensities in animals fed the control diet.

Figure 5. The proteomes of rumen microbiome populations from Holstein-Friesian beef cattle are
affected by high starch diets. A total of 60 beef cattle were subjected to two dietary contrasting
condition: 30 animals with *ad libitum* feeding and 30 subjected to 125 days of feed restriction. Dietary

components in both treatments consisted of 70% concentrate, and 30% grass silage, with the

concentrate containing rolled barley 72.5%, soya 22.5%, molasses 3% and calf mineral 2%. Rolled

barley is high in energy and starch content (~50%). Metaproteomes for a subset of 15 animals (7

restricted and 8 *ad libitum*) were analysed against a database that combined 19 protozoal

SAGs/genome (**a**) and 781 MAGs and isolate genomes (**b**), including fungal representatives. Protozoal

proteomes are presented separately as the scale of their protein quantification values were ~10x

larger. Protein quantification values were calculated by considering both the number of proteins

detected per MAG/SAG/genome and their abundance; we averaged LFQ intensities for each detected

protein across biological replicates for each dietary condition (*ad libitum*: green or restricted: orange),

which were subsequently summed for all detected proteins per MAG/SAG/genome. Similar to our

observations in Holstein dairy heifers and Alpine goats (**Figure 2)**, the proteomes of protozoal species

were a major fraction of the total rumen metaproteome, however they were substantially reduced in

dietary conditions where starch content was higher. Further supporting our results in dairy cattle, we

observed in high-starch conditions an increase in protein detection for bacterial populations affiliated

to the *Succinivibrionaceae*. **c.** Volcano plot indicating different rumen microbiome proteins that

displayed both large magnitude of fold-changes in LFQ intensities (x axis) and high statistical

significance (-log10 of nominal *p*-values using an unpaired t-test, y axis). Dashed horizontal line

denotes a *p*-value<0.05 cut-off. Volcano plots showed an increase in protein detection for bacterial

populations (greenblue) in animals fed *ad libitum* compared to protozoal populations (purple), which

were detected at higher LFQ intensities in animals under restricted conditions. LFQ intensities used to

create panel **a** and **b** can be found in **Supplementary Table S8**.

Supplementary Figure S1. Protein abundance level ratios calculated for commonly observed families

between animals fed high/low starch diets. Log 2-Fold change ratios between average LFQ intensities

for dairy cattle (**a**), goats (**b**) or beef cattle from an independent dataset (**c**) fed high or low starch diets.

Positive ratios show higher average protein abundance level for a given taxonomic family in animals

that were fed high starch diets (COS or *ad libitum*). Negative ratios show higher average protein

abundance levels for a given taxonomic family in animals that were fed the low starch diets (CTL or

Restricted). Rarely observed families with < 2 MAGs with detected proteins across both conditions

were excluded from the analysis. All ratios, and subsequent correlation tests are available in

Supplementary Table S9.

[26]

Adoral ciliary zone

Adoral lip

oesophagus

Macronucleus

Micronucleus

Clathrin-coated pit
K04646
Endocytosis

Clathrin-coated vesicle
K18584
K03283

Phagocytosis

K12478
K18464
K18466
K18468

K10364
K05756
early endosome

K07897
K12196
K19476
late endosome

K07897

early phagosome

K02145
K02146
K02149
K02150
mature phagosome

lysosome
K02146 K12391
K04646 K21393

K02145
K05146
K02149
K02150
phagolysosome

Starch

K00688 K00845
K00703 K01176
K00705 K01835
Glycogen

Glucose

K00873 K01834
K00895 K01835
K01006

Pyruvate
K01006 K00024
K00873 K00027

Acetoacetyl-CoA ← ⟵ Acetyl-CoA
K01692
Butanoyl-CoA
K00002

Butyrate   **Acetate**   **Ethanol**

Caudal spine

a Protozoal Single-Cell Amplified Genomes (SAGs)

y-axis: Protozoal protein quantification (summed LFQ intensity per SAG)

Legend:
- *Ad libitum* (high starch) (green)
- Restricted (low starch) (orange)

x-axis categories: *I. intestinalis*, *I. prostoma*, *Isotricha sp YL 2021a*, *Isotricha sp YL 2021b*, *D. ruminantium*, *En. bursa*, *En. caudatum*, *En. longinucleatum*, *D. dentatum*, *D. flabellum*, *E. triloricatum*, *M. minomm*, *E. rostratum*, *O. gracile*, *P. multivesiculatum*, *O. dentatum*, *Ep. cattanei*, *Ep. caudatum*, *O. caudatus*

b Bacterial, archaeal and fungal protein quantification (summed LFQ intensity per MAG)

y-axis: Fungal genomes and prokaryotic Metagenome-Assembled Genomes (MAGs)

Legend:
- *Ad libitum* (high starch) (green)
- Restricted (low starch) (orange)

Succinivibrionaceae-affiliated MAGs

c *Ad libitum* vs Restricted diet

Detected proteins:
- Bacteria/Archaea/Fungi (teal)
- Protozoa (purple)

y-axis: -Log$_{10}$(p-value)

x-axis: Log2(fold change *ad libitum*/restricted)