Increased levels of circulating neurotoxic metabolites in patients with mild Covid19

Estibaliz Santiago-Mujika[1], Kevin Heinrich[2], Sonia George[1], Colt D Capan[1], Cameron Forton[1], Zachary Madaj[3] Amanda R Burmeister[1], Matthew Sims [4,5], J Andrew Pospisilik[1], Patrik Brundin[1], Stewart F Graham[6,7], and Lena Brundin[1, 8][7]

[1] Parkinson's Disease Center, Department of Neurodegenerative Science, Van Andel Institute, Grand Rapids, MI 49503, USA.
[2] Quire Inc., Memphis, TN 38103, USA
[3] Bioinformatics & Biostatistics core, Van Andel Institute, Grand Rapids, MI 49503, USA
[4] Department of Internal Medicine, Oakland University William Beaumont School of Medicine, Rochester, Michigan, USA.
[5] Department of Foundational Medical Studies, Oakland University William Beaumont School of Medicine, Rochester, Michigan, USA.
[6] Metabolomics Department, Beaumont Research Institute, Beaumont Health, Royal Oak, Michigan 48073, USA.
[7] Oakland University-William Beaumont School of Medicine, Rochester, MI 48309, USA.
[8] Department of Psychiatry, Michigan State University College of Human Medicine, Grand Rapids, MI 49503, USA.
Corresponding author, Lena.Brundin@vai.org, Van Andel Institute, 333 Bostwick Ave NE, Grand Rapids, MI 49503, USA.

Abstract

SARS-CoV-2 corona virus causes a multi-faceted and poorly defined clinical and pathological phenotype involving hyperinflammation, cytokine release. And long-term cognitive deficits, with an undefined neuropathological mechanism. Inflammation increases the activity of the kynurenine pathway, which is linked to neurodegenerative and psychiatric disorders. We sought to determine whether the kynurenine pathway is impacted in patients with mild COVID-19, leading to elevated neurotoxic metabolites in blood, and whether such changes are associated with pro-inflammatory cytokines. Serum samples were taken from 150 patients and analyzed by ELISA and ultra-high performance liquid chromatography (UHPLC). The data were analyzed using multiple linear regression models adjusted for age and sex. We found increased levels of kynurenine, quinolinic acid and 3-hydroxykynurenine in serum from patients with mild COVID-19, together with increased levels of IL-6, ICAM-1, VCAM-1 and neopterin. The levels of neurotoxic metabolites were significantly associated with key inflammatory cytokines including IL-6 and TNFα. The COVID-19 risk-factor hypertension was associated with the highest levels of neurotoxic metabolites in plasma. These neuroactive metabolites could be part of the pathological mechanisms underlying cognitive impairment during and post-COVID and should be explored as potential biomarkers for long-COVID symptoms.

Keywords: Mild COVID-19, Tryptophan metabolism, 3-HK, quinolinic acid

Introduction

COVID-19, caused by SARS-CoV-2, can lead to systemic disease, including pneumonia, acute respiratory distress syndrome, and impaired consciousness (1). COVID-19 also leads to an activation of innate and adaptive immune responses which result in a substantial inflammatory response (2). In a certain percentage of patients, symptoms persist over time with the potential of symptoms lasting months and up to years after initial illness. In these patients, there is a predominance for those effected by the illness to present neuropsychiatric symptoms (3). Recent studies have shown an increased risk for acute and long-term sequalae after COVID-19 in both vaccinated and unvaccinated patients (4). However, there is currently a lack of understanding for the factors that drive the neuropsychiatric symptoms present during acute and long-term COVID-19. Thus, there is an urgent need to identify biomarkers that can indicate the disease progression during long-COVID, as well as contribute to the understanding of its underlying pathology.

The kynurenine pathway is the major route for tryptophan (TRP) metabolism, and it contributes to several fundamental biological processes, all converging in energy metabolism. Infections and inflammatory conditions can alter the activity of the enzymes in this pathway (5). Several neuroactive metabolites of the kynurenine pathway that bind neuronal receptors have an underlying role in both neurological and psychiatric symptoms (5). As such, quinolinic acid (QUIN) is an agonist of the glutamatergic N-methyl-D-aspartate (NMDA) receptor and kynurenic acid (KYNA) is an antagonist of the same receptor (6). In high concentrations, QUIN induces excitotoxic neuronal death by allowing excessive amounts of calcium to enter the cell (6). While KYNA blocks the cholinergic α7

nicotinic receptor, it also antagonizes the glycine site of the NMDA-receptor, thus

preventing calcium influx (7, 8).

Another kynurenine pathway metabolite, 3-hydroxykynurenine (3-HK) is also neurotoxic

and pro-inflammatory (9). 3-HK promotes reactive oxygen species (ROS) generation

through several oxidative conversions (10), and 3-HK can accelerate endothelial cell

apoptosis (11). The neurotoxic effects of QUIN and 3-HK are additive (5). Entry of QUIN

and KYNA into the central nervous system (CNS) is likely partially prohibited by an intact

blood brain barrier (BBB). KYN (the metabolite produced from TRP metabolism) and 3-

HK, can freely pass the BBB (12). Within the brain, KYN is metabolized to KYNA by

astrocytes (5, 13), or to 3-HK and then further to QUIN by microglia and macrophages (5,

14). SARS-CoV-2 infection is known to affect the KYN levels by inducing the production

of the pro-inflammatory cytokine, interferon-gamma (IFN-ɣ) that stimulates the rate-

limiting enzyme in the kynurenine pathway, indoleamine-2,3-dioxygenase (IDO), thus

affecting KYN levels (15).

As previously mentioned, SARS-CoV-2 can lead to a substantial inflammatory response,

which affects the levels of different metabolites from the kynurenine pathway. However,

there are also other proteins involved in the inflammatory response, such as the

intercellular cell adhesion molecule-1 (ICAM-1) and the vascular cell adhesion protein-1

(VCAM-1), in which we were interested. Both ICAM-1 and VCAM-1 are cell surface

glycoproteins that govern immune cell migration to sites of inflammation and T-cell-

mediated immunity in tissues (16-18). They are expressed in vascular endothelial cells

and in response to inflammation, their expression is induced in epithelial and immune

cells (19, 20). ICAM-1 and VCAM-1 play a central role in leukocyte trafficking, lymphocyte

activation and several immune responses, and the upregulation of ICAM-1 is a signature

event during inflammation (21). A recent study showed that monocyte-derived

macrophages drive the inflammatory response to SARS-CoV-2 and long-term changes

in inflammatory response of monocytes can be detected in convalescent SARS-CoV-2

patients following mild infection (22).Therefore, it is of interest to determine whether the

levels of ICAM-1 and VCAM-1 are affected in patients with mild COVID-19, and whether

they are linked to kynurenine pathway activation.

In the current study, we analyzed serum samples from 150 patients, 44 of whom tested

positive for COVID-19 but exhibited mild disease and were non-hospitalized. We sought

to determine whether the production of neuroactive and neurotoxic metabolites along the

kynurenine pathway, as well as several proteins involved in inflammation, is altered in

patients with mild COVID-19, as this could be associated to the underlying

neuropsychiatry symptoms. Kynurenine pathway metabolites are correlated with the

severity and predicted negative outcomes of symptoms in COVID-19 patients; and could

potentially serve as biomarkers or predictors of neuropsychiatric long-covid symptoms.

Results

Demographics of study participants

Serum samples were taken from 150 individuals, 44 were diagnosed with mild COVID-19 (defined for this purpose as positive, but not requiring hospitalization or treatment) and 106 were controls who tested negative for COVID-19. Demographics of study participants are shown in **Table 1**. The average age of SARS-CoV-2 positive individuals was 44.2 $\pm$ 13.1 years and controls were 45.6 $\pm$ 13.4 years. 47 females (44.3%) were included as controls and 25 females (56.8%) were enrolled with mild COVID-19. Of all participants, 8 controls (7.5%) and 7 COVID +ve (15.9%) were Asian, 5 controls (4.7%) and 0 COVID +ve (0%) were Black/African, 84 controls (79.2%) and 35 COVID +ve (79.5%) were White/Caucasian, 4 controls (3.8%) and 2 COVID +ve (4.5%) were Other, and 5 control (4.7%) and 0 COVID +ve (0%) opted to not answer.

Higher levels of IL-6 present in patients with mild COVID-19

Individuals with mild COVID-19 demonstrated significantly higher levels of IL-6 when compared to negative controls (ANOVA F: 5.260, p= 0.0028**) (**Figure 1a**). TNF-α levels were not altered when both groups were compared (ANOVA F: 2.347, p= 0.075 ns) (**Figure 1b**). No significant differences were observed for the other inflammatory markers including IL-13 (F: 1.847, p= 0.141 ns), IL-8 (F: 1.223, p= 0.304 ns), IFN-ɣ (F: 2.054, p= 0.109 ns, data not shown). All data was corrected for age and sex.

Higher levels of several proteins related to inflammation present in patients with mild COVID-19

The level of ICAM-1 was increased in patients who tested positive for COVID-19 when compared to the ones that did not (data corrected for sex and age, ANOVA test F: 5.823, $p<0.001$ ***) (**Figure 2a**). Similarly, patients positive for COVID-19 also showed increased levels of VCAM-1 when compared to the PCR-negative patients (data adjusted for sex and age, ANOVA F: 3.307, $p= 0.022$ *). The inflammatory related protein, neopterin was also increased in COVID-19 positive patients (data adjusted for age and sex, ANOVA test F: 3.309, $p= 0.022$ *) (**Figure 2b and 2c**)

Metabolites of the kynurenine pathway are increased in patients with mild COVID-19

Previous research demonstrated alterations in kynurenine pathway activity following infection (23-25).   Therefore, we investigated alterations in kynurenine pathway metabolite levels. Kynurenine (KYN), the first metabolite of the pathway, was significantly increased in patients with mild COVID-19 when compared to patients who tested negative for SARS-CoV-2 (ANOVA test F: 11.195, $p= 0<0.001$ ***, **Figure 3**). Additionally, 3-HK and QUIN, were increased in patients who tested positive for COVID-19 when compared to patients who did not (ANOVA test F: 3.990, $p= 0.009$ **; F: 8.492, $p<0.001$ ***, respectively, **Figure 3**). Further, picolinic acid (PIC) levels in COVID-19 positive patients were significantly decreased when compared to COVID negative patients (ANOVA test F: 4.399, $p= 0.005$ **). Finally, anthranilic acid (AA) was also significantly increased in patients with mild COVID-19 when compared to the patients who tested negative for the

virus (ANOVA test F: 4.024, p= 0.009 **, **Figure 2d**). All data was corrected for age and

sex.

The ratio of several metabolites was used to further investigate the induction of the

kynurenine pathway. The KYN/TRP ratio was quantified to determine the induction of the

kynurenine pathway. When both groups were compared, COVID-19 patients showed

significantly increased levels of the induction of the kynurenine pathway (ANOVA test F:

6.377, p< 0.001 ***, **Figure 3**). The QUIN/TRP ratio, which has been previously identified

as a biomarker for neurological diseases (26), was also analyzed. COVID-19 positive

cases showed a statistically significant increase in this ratio when compared to the non-

COVID-19 patients (ANOVA test F: 5.837, p<0.001 ***, **Figure 3**). Another neurotoxic

ratio that was measured was QUIN/KYNA. In this study, COVID-19 positive patients

presented significantly higher ratios when compared to the control group (ANOVA test F:

2.847, p= 0.040 *, **Figure 3**). All data was corrected for age and sex.

Correlation between neurotoxic metabolites of the kynurenine pathway and pro-

inflammatory cytokines

Previous research has shown the relationship between inflammation and the kynurenine

pathway activity. Therefore, we subsequently investigated whether the levels of

kynurenine metabolites and inflammatory cytokines were correlated. As shown in **Figure**

4, there was a positive correlation between the levels of TNF-α and KYN (Pearson

correlation: 0.453; p=0.002**), 3-HK (Pearson correlation: 0.527; p<0.001***) and QUIN

(Pearson correlation: 0.482; p<0.001***). Furthermore, a positive correlation between IL-

6 and both 3-HK (Pearson correlation: 0.328; p=0.03*) and QUIN (data adjusted for age

and sex, Pearson correlation: 0.418; p=0.005**) was also found. All data was corrected

for age and sex.

Increased levels of neurotoxic metabolites in COVID-19 patients with hypertension

We then analyzed patients who presented with hypertension, a well-known risk factor for

severe COVID-19, and compared them with the patients that were not hypertensive in

both COVID-19 positive and negative patients. Regression analysis adjusted for age and

sex revealed that COVID-19 hypertensive patients presented evidence of higher levels

when compared to non-hypertensive COVID-19 patients in the following proteins: IL-2

(fold change 2.66, p=0.003**), IL-6 (fold change 1.45, p-value=0.014*), TNF-α (fold

change 1.38, p=0.012*), 3-HK (fold change 1.26, p=0.08), and the QUIN/TRP ratio (fold

change 1.14, p = 0.089) (**Figure 5**).

Discussion

In the current study, we investigated the inflammatory and kynurenine pathway metabolite signatures between mild COVID-19 cases and controls. We found increases in several kynurenine pathway metabolites, such as KYN, QUIN, 3-HK and PIC, together with higher levels of IL-6. Additionally, the increased levels in QUIN/TRP, QUIN/KYN, QUIN/KYNA and KYN/TRP, further support a recent published study by Cihan and colleagues (23). Pro-inflammatory pathway proteins, along with metabolites from the kynurenine pathway were significantly increased in patients that presented with both COVID-19 and hypertension when compared to COVID-19 patients that were not hypertensive. Those proteins included IL-2, IL-6, TNFα, ICAM-1, VCAM-1, 3-HK, QUIN, QUIN:TRP and KYN:TRP.

Kynurenine pathway metabolites are correlated with severity and predicted negative outcomes of symptoms in COVID-19 patients; therefore, it is important to understand the role of kynurenine metabolites in mild COVID-19 patients and long-haulers, in particular those with neuropsychiatric symptoms (15). In this study, we found the levels of ICAM-1 and VCAM-1 to be significantly increased in patients with mild SARS-CoV-2, in particular those with hypertension. Increased levels of ICAM-1 and VCAM-1 in mild and severe cases of COVID-19 infection has already been reported in a small study (27), where it was observed that the severity of COVID-19 disease was associated with increased levels of ICAM-1 and VCAM-1. The main caveat with that study was the small number of patients analyzed. Our new findings support the prior observations and link increases in endothelial cell adhesion molecules to kynurenine pathway metabolites, such as IDO.

IDO is expressed in endothelial cells of vessel walls, and under pathological states, a

decrease in IDO leads to an increase in VCAM-1 (28). Therefore, understanding how the

kynurenine pathway is involved in inflammatory diseases and how it can alter the levels

of ICAM-1 and VCAM-1 will be important to understand, especially in the context of

COVID-19.

Lionetto et al., measured the levels of KYN and TRP in the serum of healthy patients,

SARS-CoV-2 negative patients, and SARS-CoV-2 positive patients (29). In SARS-CoV-2

positive patients, the KYN/TRP ratio was higher when compared to the negative and the

healthy controls (29). We also found an increase in the KYN/TRP ratio in SARS-CoV-2

positive patients when compared to those who tested negative. The KYN/TRP ratio is

usually used as an indirect measure of the activity of the IDO enzyme (30); which

catalyzes tryptophan. Previous research has demonstrated that IDO is regulated by IFN-

ɣ (31, 32). We observed no differences in the levels of serum IFN-ɣ in mild COVID-19

samples. However, IDO activation and gene expression has also been shown to be

altered by noncanonical pathways in addition to IFN-ɣ (33). This potentially explains the

increased kynurenine pathway activation that we observed in our cohort. The activation

of the kynurenine pathway may cause long-lasting inflammation rather than the acute

inflammation observed early in COVID-19. Therefore, it will be of interest to follow the

kynurenine metabolites as markers in patients with symptoms of long-COVID.

Together with the KYN/TRP ratio, the QUIN/TRP ratio was increased in patients with

COVID-19 when compared to controls. In a previous study, Drewes and colleagues

suggest that cerebrospinal fluid (CSF) QUIN/TRP ratio could be an early, predictive

marker of CNS disease (26). In simian-immunodeficient virus (SIV)-infected macaques,

the ratio of QUIN/TRP was significantly increased in cases that led to severe encephalitis

(26). Encephalitis is one of the many potential consequences of COVID-19 patients (34);

and the QUIN/TRP ratio could be used to determine the outcome of patients. Our data

show a significant increase in this ratio in COVID-19 patients, when compared to those

that tested negative for the virus. We also found that the ratio between QUIN/TRP was

also increased in hypertensive COVID-19 patients, suggesting that patients with

hypertension might be prone to a worse outcome.

Cihan and colleagues analyzed kynurenine pathway metabolites and inflammatory

cytokines and found a positive correlation between IL-6 and various metabolites from the

kynurenine pathway (23). Our study supports a positive correlation between IL-6 and

QUIN as well as between IL-6 and 3-HK. Since Cihan's study analyzed severe and ICU

cases of COVID-19 patients, it is possible that the lack of correlation between IL-6 and

other kynurenine metabolites in our study is because only mild COVID-19 cases were

analyzed. The correlation of IL-6 and kynurenine metabolites could also potentially be

used as a biomarker of disease severity.

The correlation between IL-6, IFN-ɣ, and TNF-α and kynurenine metabolites supports the

link between inflammation, SARS-CoV-2 and the kynurenine pathway. We found a

positive correlation between TNF-α and three metabolites: 3-HK, KYN, and QUIN. TNF-

α affects the kynurenine pathway in patients with schizophrenia, as a positive correlation

between the levels of TNF-α and KYN has been reported (35). TNF-α may accelerate the

formation of KYN from the catabolism of TRP (35). We did not observe differences in

TNF-α levels between negative and positive SARS-CoV-2 patients, however, we did find

a correlation between TNF-α and KYN levels, similar to what has been reported for

schizophrenia (35). In this study, QUIN, 3-HK and KYN were significantly increased in

patients with mild COVID-19. Both QUIN and 3-HK are neurotoxic metabolites and were

positively correlated with TNF-α and IL-6 in the case of QUIN, and TNF-α in the case of

3-HK, it is possible that the inflammation observed in COVID-19 patients further

contributes to the neuronal damage caused by the neurotoxic metabolites.

To summarize, we found an increase in neurotoxic metabolites of the kynurenine pathway

in patients with mild COVID-19. Furthermore, the neurotoxic metabolites were correlated

with inflammatory markers and vascular injury markers, such as TNF-α, IL-6, VCAM-1

and ICAM-1. We hypothesize that the activation of the neurotoxic branch of the

kynurenine pathway might contribute to neurological, cognitive, and psychiatric

symptoms experienced in COVID-19 and its aftermath. We suggest that these

metabolites should be studied further for their potential as biomarkers of long COVID and

as potential contributors to the disease mechanisms underlying long COVID.

Materials and Methods

Blood samples

Blood was drawn by venipuncture of the right or left antecubital vein. Blood was allowed to clot at room temperature prior to centrifugation. Serum was aliquoted into cryovials and immediately transferred to -80°C for storage until biological assays.

Detection of tryptophan, serotonin, and kynurenine metabolites

Serum samples were mixed with extraction solution, briefly vortexed and centrifuged. The supernatant, which contains the metabolites of interest was removed and dried under reduced pressure conditions for ninety minutes in a GeneVac EZ-2 Plus speedvac (SP Scientific, Warminster,PA). Dried down extracts were then resuspended in 0.1% formic acid in Milli-Q water, once resuspended samples were centrifuged through a COSTAR Spin-X 0.22-um filter tube and transferred to an amber vial containing a glass insert.

Quantification of kynurenine metabolites

Kynurenine pathway metabolites (KYN, KYNA, 3-HK, QUIN, PIC, NTA, NIC, AA), TRP and serotonin were quantified using reverse phase ultra-high-performance liquid chromatography (UHPLC) coupled to a triple quadrupole mass spectrometer (1290 Infinity II LC System, 6470 Triple quadrupole, Agilent Technologies, Santa Clara, CA). Five-microliters of samples were injected onto a Vanguard HSS T3 Pre-column that was connected to an Acquity HSS T3 analytical column. Elution conditions used a combination of Solvent A (0.1% formic acid in LC/MS grade Water) and Solvent B (0.1% formic acid

in 90% LC/MS grade acetonitrile) at a flow rate of 0.4mL/min. Agilent Masshunter

Quantitative Analysis Software (v9.0, Agilent) was used to analyze and export data.

Intra-assay coefficients of variability (CV) for plasma analytes: TRP 4.1%, KYN 2.5%,

KYNA 1.6%, 3-HK 5.0%, QUIN 2.8%, PIC 3.3%, NTA 1.6%, NIC 5.4%, AA 8.4% and 5-

HT 2.2%.

Inter-assay CVs: TRP 4.8%, KYN 1.8%, KYNA 1.5%, 3-HK 4.9%, QUIN 1.8%, PIC 1.9%,

NTA 1.8%, NIC 6.5%, AA 7.2% and 5-HT 4.9%.

Lower limits of detection (LLOD) were found to be as follows: TRP 36.6 nM, KYN 2.2 nM,

KYNA 0.16 nM, 3-HK 0.29 nM, QUIN 4.15 nM, PIC 0.63 nM, NTA 0.98 nM, NIC 0.07 nM,

AA 0.98 nM and 5-HT 0.73 nM.

Quantification of cytokines and alpha-synuclein

Three Meso Scale Discovery (MSD) multiplex kits (Meso Scale Diagnostics LLC,

Rockville, MD) were read using a MESO QuickPlex SQ 120 plate reader. Samples were

run in duplicate according to the manufacturer's instructions and sample concentrations

were generated through MSD Discovery Workbench 4.0 software. Samples below the

average LLOD were denoted as the average LLOD across plates. For α-synuclein, the

U-PLEX human α-synuclein kit was used. Samples were diluted 1:8 with sample diluent

and generated an inter-plate CV of 3%, an average intra-assay CV of 5.2%, and the LLOD

of 0.876 pg/mL.

Using the MSD V-PLEX human vascular injury II kit, we quantified C-reactive protein

(CRP), serum amyloid A (SAA), ICAM-1, and VCAM-1. Samples were diluted 1:1000 with

manufacturer's diluent reagent. The inter-plate CV was 10.3% (CRP: 8.3%, SAA: 9.7%,

ICAM-1: 15.2%, VCAM-1: 8.1 %), average intra-assay CV 5.6% (CRP 4.2%, SAA 6.5%,

ICAM-1 6.6%, and VCAM-1 5.2%), the LLOD were 1.077pg/mL CRP, 12.839 pg/mL SAA,

1.037 pg/mL ICAM-1, and 6.618 pg/mL VCAM-1.

The MSD V-PLEX human proinflammatory I kit was used to determine the level of IFN-ɣ,

interleukin (IL)-10, IL-12p70, IL-13, IL-1β, IL-2, IL-4, IL-6, IL-8, and tumor necrosis factor

alpha (TNF-α). Samples were diluted 1:1 with sample diluent. Inter-plate CV were

calculated (**see Supplementary Table 1**), and the LLOD was IFN-ɣ: 0.087 pg/mL, IL-10:

0.025 pg/mL, IL-12p70: 0.0385 pg/mL, IL-13: 0.209 pg/mL, IL-1β: 0.032 pg/mL, IL-2:

0.024 pg/mL, IL-4: 0.004 pg/mL, IL-6: 0.045 pg/mL, IL-8: 0.017 pg/mL, and TNF-α: 0.049

pg/mL.

Quantification of Neopterin and S100B

Neopterin ELISA kits were purchased from IBL America (Immuno-Biological Laboratories

Inc., Minneapolis, MN). Undiluted serum samples were used following the manufacturer's

protocol. S100B ELISA kits were purchased from Millipore (EMD Millipore, St. Louis, MO)

and samples were diluted 1:1. Plates were read using a Tecan Infinite M200 Pro plate

reader (Tecan Group Ltd, Männedorf, Switzerland). Sample concentrations were

generated using a 4 Parameter Logistic Curve Calculator (AAT Bioquest, "Quest Graph

Four Parameter Logistic (4PL) Curve Calculator" 15 Jul. 2021,

https://www.aatbio.com/tools/four-parameter-logistic-4pl-curve-regression-online-

calculator). The average inter-plate CV for Neopterin was 12.1%, the average intra-assay

CV was 2.2%, and the LLOD given by the manufacturer is 0.177 ng/mL. For S100B the

average inter-plate CV was 5.0%, the average intra-assay CV was 1.9%, and the LLOD

was 2.7 pg/mL as specified by the manufacturer.

Statistical analysis

Statistical analysis was done using SPSS Statistics (version 28.0.1.0). Linear regression

models were adjusted for age and sex in all instances. The ANOVA test was used to

compare both groups and statistical significance was considered when $p<0.05$.

Correlation analyses were also done with SPSS Statistics and were also adjusted for age

and sex in all instances. The graphs were generated using GraphPad Prism version 9.0

(GraphPad Software, La Jolla, CA).

Study approval

This study utilized a subset of samples from the Beaumont Health Large-Scale

Automated Serologic Testing for COVID-19 study (36) and was approved by the

Institutional Review Board (IRB) at the Beaumont Research Institute Detroit, Michigan,

USA (2021-110). The final cohort consisted of 150 individuals (44 Mild COVID-19 cases

and 106 controls) randomly selected from the registry. Individuals testing positive for

SARS-CoV-2 in a qPCR test at the time of sampling were assigned "Covid19 positive"

whereas individuals testing negative in a qPCR test for C SARS-CoV-2 at the time of

blood sample were assigned "Covid19 negative" for the purpose of this study.

Author Contributions

Conceptualization, LB SFG, PB; data curation, ESM, ARB, CDC, CF, KH; formal analysis, ESM, LB, SFG, PB, regression analysis with R, ZM; methodology, LB; visualization, ESM, ARB; writing—original draft, ESM, ARB; writing—review and editing, SG, MXH, JAP, PB, SFG, LB.

Acknowledgements

We would like to thank the Farmer Family Foundation for their generous contribution, which allowed for this research to be conducted. Additionally, we would like to thank the Van Andel Institute Metabolomics and Bioenergetics Core for providing technical assistance.   This project was also supported with Beaumont institutional funds as well as through the Beaumont Health Foundation with philanthropic gifts provided by the following: Sidney & Madeline Forbes, Nathan & Catherine Forbes, Edward C. Levy & the Linda Dresner Foundation, Stephen & Bobbi Polk, Warren & Carol Ann Rose & Family, Elizabeth Rose, Mickey Shapiro & Family, S. Evan & Gwen Weiner and The Hearst Foundations.

List of abbreviations

List of abbreviations

3-HK: 3-HydroxyKynurenine

AA: Anthranilic Acid

BBB: Blood Brain Barrier

COVID-19: Coronavirus Disease (2019)

CI: Confidence Interval

CSF: Cerebrospinal Fluid

Glu: Glutamate

ICAM-1: Intercellular Adhesion Molecule 1

ICU: Intensive Care Unit

IDO: Indoleamine 2,3-Dioxygenase

IFN-γ: Interferon Gamma

IL: Interleukins

KYN: Kynurenine

KYNA: Kynurenic Acid

PA: Picolinic Acid

PCR: Polymerase Chain Reaction

ROS: Reactive Oxygen Species

S100B: S100- Calcium-Binding Protein B

SAA: Serum Amyloid A

SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus 2

SIV: Simian Immunodeficient Virus

TNF-α: Tumor Necrosis Alpha

431 TRP: Tryptophan

432 QUIN: Quinolinic acid

434 **Data availability**

435 Data generated is available upon request.

437 **Competing interests**

439 P.B. has received support as a consultant from Calico Life Sciences, CureSen,

440 Enterin Inc, Idorsia Pharmaceuticals, Lundbeck A/S, AbbVie, Fujifilm-Cellular Dynamics

441 International, and Axial Therapeutics. He has received commercial support for research

442 from Lundbeck A/S and F. Hoffman-La Roche. He has ownership interests in Acousort

443 AB, Axial Therapeutics, Enterin Inc and RYNE Biotechnology. During the time that this

444 paper was written he became an employee of F. Hoffman-La Roche, although none of

445 the data were generated by this company.

References

1. Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, et al. Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China. JAMA Neurol. 2020;77(6):683-90. doi: 10.1001/jamaneurol.2020.1127. PubMed PMID: 32275288; PubMed Central PMCID: PMC7149362.

2. Anka AU, Tahir MI, Abubakar SD, Alsabbagh M, Zian Z, Hamedifar H, et al. Coronavirus disease 2019 (COVID-19): An overview of the immunopathology, serological diagnosis and management. Scand J Immunol. 2021;93(4):e12998. Epub 20201203. doi: 10.1111/sji.12998. PubMed PMID: 33190302; PubMed Central PMCID: PMC7744910.

3. Roy D, Ghosh R, Dubey S, Dubey MJ, Benito-León J, Kanti Ray B. Neurological and Neuropsychiatric Impacts of COVID-19 Pandemic. Can J Neurol Sci. 2021;48(1):9-24. Epub 20200805. doi: 10.1017/cjn.2020.173. PubMed PMID: 32753076; PubMed Central PMCID: PMC7533477.

4. Al-Aly Z, Bowe B, Xie Y. Long COVID after breakthrough SARS-CoV-2 infection. Nat Med. 2022. Epub 20220525. doi: 10.1038/s41591-022-01840-0. PubMed PMID: 35614233.

5. Schwarcz R, Bruno JP, Muchowski PJ, Wu HQ. Kynurenines in the mammalian brain: when physiology meets pathology. Nat Rev Neurosci. 2012;13(7):465-77. doi: 10.1038/nrn3257. PubMed PMID: 22678511; PubMed Central PMCID: PMC3681811.

6. Lugo-Huitrón R, Blanco-Ayala T, Ugalde-Muñiz P, Carrillo-Mora P, Pedraza-Chaverrí J, Silva-Adaya D, et al. On the antioxidant properties of kynurenic acid: free radical scavenging activity and inhibition of oxidative stress. Neurotoxicol Teratol. 2011;33(5):538-47. Epub 20110713. doi: 10.1016/j.ntt.2011.07.002. PubMed PMID: 21763768.

7. Hilmas C, Pereira EF, Alkondon M, Rassoulpour A, Schwarcz R, Albuquerque EX. The brain metabolite kynurenic acid inhibits alpha7 nicotinic receptor activity and increases non-alpha7 nicotinic receptor expression: physiopathological implications. J Neurosci. 2001;21(19):7463-73. doi: 10.1523/jneurosci.21-19-07463.2001. PubMed PMID: 11567036; PubMed Central PMCID: PMC6762893.

8. Stone TW. Neuropharmacology of quinolinic and kynurenic acids. Pharmacol Rev. 1993;45(3):309-79. PubMed PMID: 8248282.

9. Wang Q, Liu D, Song P, Zou MH. Tryptophan-kynurenine pathway is dysregulated in inflammation, and immune activation. Front Biosci (Landmark Ed). 2015;20(7):1116-43. Epub 20150601. doi: 10.2741/4363. PubMed PMID: 25961549; PubMed Central PMCID: PMC4911177.

10. Ishii T, Iwahashi H, Sugata R, Kido R. Formation of hydroxanthommatin-derived radical in the oxidation of 3-hydroxykynurenine. Arch Biochem Biophys. 1992;294(2):616-22. doi: 10.1016/0003-9861(92)90733-d. PubMed PMID: 1314547.

11. Wang Q, Zhang M, Ding Y, Wang Q, Zhang W, Song P, et al. Activation of NAD(P)H oxidase by tryptophan-derived 3-hydroxykynurenine accelerates endothelial apoptosis and dysfunction in vivo. Circ Res. 2014;114(3):480-92. Epub 20131126. doi: 10.1161/circresaha.114.302113. PubMed PMID: 24281189; PubMed Central PMCID: PMC4104160.

12. Skorobogatov K, De Picker L, Verkerk R, Coppens V, Leboyer M, Müller N, et al. Brain Versus Blood: A Systematic Review on the Concordance Between Peripheral and Central Kynurenine Pathway Measures in Psychiatric Disorders. Front Immunol. 2021;12:716980. Epub

20210923. doi: 10.3389/fimmu.2021.716980. PubMed PMID: 34630391; PubMed Central
PMCID: PMC8495160.
13.   Guillemin GJ, Kerr SJ, Smythe GA, Smith DG, Kapoor V, Armati PJ, et al. Kynurenine
pathway metabolism in human astrocytes: a paradox for neuronal protection. J Neurochem.
2001;78(4):842-53. doi: 10.1046/j.1471-4159.2001.00498.x. PubMed PMID: 11520905.
14.   Agudelo LZ, Femenía T, Orhan F, Porsmyr-Palmertz M, Goiny M, Martinez-Redondo V,
et al. Skeletal muscle PGC-1α1 modulates kynurenine metabolism and mediates resilience to
stress-induced depression. Cell. 2014;159(1):33-45. doi: 10.1016/j.cell.2014.07.051. PubMed
PMID: 25259918.
15.   Oxenkrug G, Summergrad P. Peripheral kynurenines as biomarkers and targets for
prevention and treatment of psychiatric conditions associated with SARS-CoV-2 infection.
Personalized   Medicine   in   Psychiatry.   2021;29-30:100088.   doi:
https://doi.org/10.1016/j.pmip.2021.100088.
16.   Lembas A, Zawartko K, Sapuła M, Mikuła T, Kozłowska J, Wiercińska-Drapało A.
VCAM-1 as a Biomarker of Endothelial Function among HIV-Infected Patients Receiving and
Not Receiving Antiretroviral Therapy. Viruses. 2022;14(3). Epub 20220311. doi:
10.3390/v14030578. PubMed PMID: 35336985; PubMed Central PMCID: PMC8955345.
17.   Chen YH, Lightman S, Eskandarpour M, Calder VL. Adhesion Molecule Targeted Therapy
for Non-Infectious Uveitis. Int J Mol Sci. 2022;23(1). Epub 20220103. doi:
10.3390/ijms23010503. PubMed PMID: 35008929; PubMed Central PMCID: PMC8745221.
18.   Kang L, Kim M, Lee YM. Expression of ICAM-1 in Blood Vascular Endothelium and
Tissues in Human Premalignant Lesion and Gastric/Hepatocellular Carcinomas. Korean J
Gastroenterol. 2022;79(4):170-6. doi: 10.4166/kjg.2022.008. PubMed PMID: 35473775.
19.   Bui TM, Wiesolek HL, Sumagin R. ICAM-1: A master regulator of cellular responses in
inflammation, injury resolution, and tumorigenesis. J Leukoc Biol. 2020;108(3):787-99. Epub
20200317. doi: 10.1002/jlb.2mr0220-549r. PubMed PMID: 32182390; PubMed Central PMCID:
PMC7977775.
20.   Hyun YM, Lefort CT, Kim M. Leukocyte integrins and their ligand interactions. Immunol
Res. 2009;45(2-3):195-208. Epub 20090129. doi: 10.1007/s12026-009-8101-1. PubMed PMID:
19184539; PubMed Central PMCID: PMC2990789.
21.   Ramos TN, Bullard DC, Barnum SR. ICAM-1: isoforms and phenotypes. J Immunol.
2014;192(10):4469-74. doi: 10.4049/jimmunol.1400135. PubMed PMID: 24795464; PubMed
Central PMCID: PMC4015451.
22.   Bohnacker S, Hartung F, Henkel F, Quaranta A, Kolmert J, Priller A, et al. Correction to:
Mild COVID-19 imprints a long-term inflammatory eicosanoid- and chemokine memory in
monocyte-derived macrophages. Mucosal Immunol. 2022:1. Epub 20220513. doi:
10.1038/s41385-022-00526-7. PubMed PMID: 35562559; PubMed Central PMCID:
PMC9098895.
23.   Cihan M, Doğan Ö, Ceran Serdar C, Altunçekiç Yıldırım A, Kurt C, Serdar MA.
Kynurenine pathway in Coronavirus disease (COVID-19): Potential role in prognosis. J Clin Lab
Anal. 2022;36(3):e24257. Epub 20220129. doi: 10.1002/jcla.24257. PubMed PMID: 35092710;
PubMed Central PMCID: PMC8906035.
24.   Holtze M, Asp L, Schwieler L, Engberg G, Karlsson H. Induction of the kynurenine
pathway by neurotropic influenza A virus infection. J Neurosci Res. 2008;86(16):3674-83. doi:
10.1002/jnr.21799. PubMed PMID: 18655201.

25.   Wickström R, Fowler Å, Goiny M, Millischer V, Ygberg S, Schwieler L. The Kynurenine
Pathway is Differentially Activated in Children with Lyme Disease and Tick-Borne Encephalitis.
Microorganisms. 2021;9(2). Epub 20210204. doi: 10.3390/microorganisms9020322. PubMed
PMID: 33557172; PubMed Central PMCID: PMC7913947.
26.   Drewes JL, Meulendyke KA, Liao Z, Witwer KW, Gama L, Ubaida-Mohien C, et al.
Quinolinic acid/tryptophan ratios predict neurological disease in SIV-infected macaques and
remain elevated in the brain under cART. J Neurovirol. 2015;21(4):449-63. Epub 20150317. doi:
10.1007/s13365-015-0334-2. PubMed PMID: 25776527; PubMed Central PMCID:
PMC4512924.
27.   Tong M, Jiang Y, Xia D, Xiong Y, Zheng Q, Chen F, et al. Elevated Expression of Serum
Endothelial Cell Adhesion Molecules in COVID-19 Patients. J Infect Dis. 2020;222(6):894-8. doi:
10.1093/infdis/jiaa349. PubMed PMID: 32582936; PubMed Central PMCID: PMC7337874.
28.   Kiluk M, Lewkowicz J, Pawlak D, Tankiewicz-Kwedlo A. Crosstalk between Tryptophan
Metabolism via Kynurenine Pathway and Carbohydrate Metabolism in the Context of Cardio-
Metabolic Risk-Review. J Clin Med. 2021;10(11). Epub 20210604. doi: 10.3390/jcm10112484.
PubMed PMID: 34199713; PubMed Central PMCID: PMC8199979.
29.   Lionetto L, Ulivieri M, Capi M, De Bernardini D, Fazio F, Petrucca A, et al. Increased
kynurenine-to-tryptophan ratio in the serum of patients infected with SARS-CoV2: An
observational cohort study. Biochim Biophys Acta Mol Basis Dis. 2021;1867(3):166042. Epub
20201216. doi: 10.1016/j.bbadis.2020.166042. PubMed PMID: 33338598; PubMed Central
PMCID: PMC7834629.
30.   Badawy AA-B, Guillemin G. The Plasma [Kynurenine]/[Tryptophan] Ratio and
Indoleamine 2,3-Dioxygenase: Time for Appraisal. International Journal of Tryptophan Research.
2019;12:1178646919868978. doi: 10.1177/1178646919868978. PubMed PMID: 31488951.
31.   Badawy AA. Kynurenine Pathway of Tryptophan Metabolism: Regulatory and Functional
Aspects. Int J Tryptophan Res. 2017;10:1178646917691938. Epub 20170315. doi:
10.1177/1178646917691938. PubMed PMID: 28469468; PubMed Central PMCID:
PMC5398323.
32.   Sarkar SA, Wong R, Hackl SI, Moua O, Gill RG, Wiseman A, et al. Induction of
indoleamine 2,3-dioxygenase by interferon-gamma in human islets. Diabetes. 2007;56(1):72-9.
doi: 10.2337/db06-0617. PubMed PMID: 17192467.
33.   Liang Y, Yu Z, Song Y, Wang T, Xiao B. Indoleamine 2,3-Dioxygenase Activation by
Interferon Gamma in Vascular Endothelial Rat Cells Requires Noncanonical NF-κB Signaling.
Transplant Proc. 2019;51(6):2141-5. Epub 20190712. doi: 10.1016/j.transproceed.2019.03.043.
PubMed PMID: 31307771.
34.   Ellul MA, Benjamin L, Singh B, Lant S, Michael BD, Easton A, et al. Neurological
associations of COVID-19. Lancet Neurol. 2020;19(9):767-83. Epub 20200702. doi:
10.1016/s1474-4422(20)30221-0. PubMed PMID: 32622375; PubMed Central PMCID:
PMC7332267.
35.   Okamoto N, Natsuyama T, Igata R, Konishi Y, Tesen H, Ikenouchi A, et al. Associations
Between the Kynurenine Pathway, Proinflammatory Cytokines, and Brain-Derived Neurotrophic
Factor in Hospitalized Patients With Chronic Schizophrenia: A Preliminary Study. Front
Psychiatry. 2021;12:696059. Epub 20210728. doi: 10.3389/fpsyt.2021.696059. PubMed PMID:
34393855; PubMed Central PMCID: PMC8357143.
36.   Sims MD, Maine GN, Childers KL, Podolsky RH, Voss DR, Berkiw-Scenna N, et al.
Coronavirus Disease 2019 (COVID-19) Seropositivity and Asymptomatic Rates in Healthcare

Workers Are Associated with Job Function and Masking. Clin Infect Dis. 2021;73(Suppl 2):S154-
s62. doi: 10.1093/cid/ciaa1684. PubMed PMID: 33150375; PubMed Central PMCID:
PMC7665441.

616   **Table 1. Demographics of the patients included in the current study**

Mild SARS-CoV-2 Demographics		
	SARS-CoV-2 Negative (n = 106)	SARS-CoV-2 Positive (n = 44)
Age (Mean $\pm$ SD)	45.65 $\pm$ 13.39	44.18 $\pm$ 13.12
Sex F (%)	47 (44.3%)	25 (56.8%)
Race n (%)		
Asian	8 (7.5%)	7 (15.9%)
Black/African	5 (4.7%)	0 (0%)
White/Caucasian	84 (79.2%)	35 (79.5%)
Other	4 (3.8%)	2 (4.5%)
No Answer	5 (4.7%)	0 (0%)

617
618   SD: Standard Deviation
619   F: female

 Figures:

Figure 1. Significantly higher levels IL-6 in patients with mild COVID-19 when compared to controls. a) Significantly higher levels of IL-6 were found in patients with COVID-19 when compared to the negative controls (data adjusted for age and sex, ANOVA test F: 5.260, p=0.002 **). **b)** No differences were found in the level of TNF-α between the two groups (data adjusted for age and sex, ANOVA test F: 2.347, p= 0.075 ns). Graphs are represented by median with 95% of confidence interval (CI).

673
Figure 2. Significantly higher levels of serum ICAM-1, VCAM-1, and neopterin in patients with COVID-19. When the levels of all three proteins were analyzed, patients with COVID-19 presented significantly higher levels of **a)** ICAM-1 (data adjusted for age and sex, ANOVA test F: 5.823, p<0.001 ***), **b)** VCAM-1 (data adjusted for age and sex, ANOVA test F: 3.307, p= 0.022 *), and **c)** neopterin (data adjusted for age and sex, ANOVA test F:3.309, p= 0.022*). Graphs are represented by median with 95% of confidence interval (CI).

681
Figure 3. The kynurenine pathway is altered in patients with mild COVID-19, who present increased levels of neurotoxic metabolites. Significantly increased levels of kynurenine (data adjusted for age and sex, ANOVA test F: 11.195, p<0.001 ***), 3-hydroxykynurenine (data adjusted for age and sex, ANOVA test F: 3.390, p=0.009 **), anthranilic acid (data adjusted for age and sex, ANOVA test F: 4.024, p=0.009 **), and quinolinic acid (data adjusted for age and sex, ANOVA test F: 8.492, p<0.001 ***) were found in patients with mild COVID-19 when compared to controls. When the ratio of the metabolites was analyzed, significantly increased levels of KYN/TRP (data adjusted for age and sex, ANOVA test F: 6.377, p<0.001 ***) and QUIN/TRP (data adjusted for age and sex, ANOVA test F: 5.837, p<0.001 ***), as well as QUIN/KYNA (data adjusted for age and sex, ANOVA test F: 2.847, p= 0.040 *) were found in patients with COVID-19. Graphs show the median with 95% of CI. Abbreviations: IDO, Indoleamine 2,3-dioxygenase; TDO, Tryptophan 2,3-dioxygenase; KATs, Kynurenine aminotransferase; KYNA, kynurenic acid; KMO, Kynurenine 3-monooxygenase; AA, anthranilic acid; 3-HK, 3-hydroxykynurenine; KYNU, Kynureninase; 3-HAA, 3-hydroxyanthranilic acid; 3HAO, 3-hydroxyanthranilate oxidase; ACMSD, Aminocarboxymuconate-semialdehyde decarboxylase, and NAD, Nicotinamide adenine dinucleotide.

Figure 4. There is correlation between inflammatory cytokines and metabolites of the kynurenine pathway in patients with mild COVID-19. Patients with mild COVID-19 present a positive correlation between TNF-α and kynurenine (data adjusted for age and sex, Pearson R: 0.4512; p=0.0021 **), TNF-α and 3-hydroxykynurenine (data adjusted for age and sex, Pearson R: 0.4567; p=0.0018 **), TNF-α and quinolinic acid (data adjusted for age and sex, Pearson R: 0.4812; p=0.0009 ***), IL-6 and 3-hydroxykynurenine (data adjusted for age and sex, Pearson R: 0.3252, p=0.0312 *) and between IL-6 and quinolinic acid (data adjusted for age and sex, Pearson R: 0.4161, p=0.0050 **).

725
726
Figure 5. Increased metabolites and ratio in COVID-19 positive hypertensive
patients when compared to non-hypertensive COVID-19 patients. Higher levels of IL-
2 (fold change 2.66, p=0.003), IL-6 (fold change 1.45, p-value=0.014), and TNF-α (fold
change 1.38, p=0.012) in COVID-19 hypertensive patients when compared to COVID-19
non-hypertensive patients. Trend towards increased levels in 3-HK (fold change 1.26,
p=0.08), and the QUIN/TRP ratio (fold change 1.14, p = 0.089). Graphs are represented
by median with 95% of confidence interval (CI).