Over the hills and far away: linking landscape factors with cavity excavation on living forest trees by the Black Woodpecker (*Dryocopus martius,* L. 1758)

Cedric Cabrera[1], Jean-Matthieu Monnet[1], Jean-Jacques Boutteaux[2], Baptiste Doutau[3], Pascal Denis[4], Yoan Paillet[1,*]

1. Univ. Grenoble Alpes, INRAE, Lessem, 2 rue de la Papeterie, 38400 Saint-Martin d'Hères, France

2. ONF, Unité Territoriale d'Auberive & Groupe Cigogne noire, 6 rue de l'Abbatiale, 52160 Auberive, France

3. LPO Auvergne Rhone-Alpes, Délégation territoriale Haute-Savoie, 46 route de la fruitière, 74650 Chavanod, France

4. ONF, Réseau Avifaune, Direction Forêt et Risques Naturels, 68 rue Maurice Burrus, 68160 Ste Croix aux Mines, France.

*corresponding author: yoan.paillet@inrae.fr

Abstract

The Black Woodpecker (*Dryocopus martius*, L. 1758) is the largest primary cavity excavator in

Western Europe. Its cavities represent an essential microhabitat for many other forest species and the

landscape factors linked with cavity excavation by the Black Woodpecker are needed to support the

conservation of this species. Such relationships should thus be quantified at the scale of the home

range in Western Europe.

We used cavity maps by foresters and naturalists to build a large (2689 cavity bearing trees) database

distributed over 11 sites in France. Based on this and on a set of background points, we analysed the

effects of stand composition and landscape features (forest cover, landscape connectivity and

fragmentation) at three different scales around each plot corresponding to a forest management unit

(10ha), the core (100ha) and extended (250ha) home range scales.

We showed that landscape shape index and forest composition (especially the presence of mixed

forests) had significant effects but that the magnitude varied across the three scales. We notably

observed the strongest effects at the core home range scale (100ha), indicating that Black

Woodpecker requirements for cavity excavation are more pronounced at this scale. The Black

Woodpecker tend to avoid conifer-dominated stands to excavate cavities, but benefits from mixed

forests, that couple favourable foraging and cavity excavation sites. The bird also prefers continuous

forest landscapes with low edge densities. We also showed that effects of landscape were stronger in

wetter climates and higher slopes, indicating context-dependence.

Forest planning rarely integrates the landscape patterns in western temperate Europe, a better

understanding of the features linked with cavity excavation by the Black Woodpecker may hence help

to better integrate their conservation in forest management planning. Our results also show the

importance to maintain mixed broadleaf-conifer forests as well as continuous and well-connected

forest landscapes to favour features that benefit secondary cavity nesters at the large scale.

Keywords: Black Woodpecker, connectivity, landscape features, forest management, Tree-related

microhabitat, cavity

Introduction

The conservation of forest biodiversity relies on an ensemble of methods including setting aside forest reserves and small conservation patches and preserving, within managed forests, elements that favor the biodiversity of forest dwelling species (Bollmann and Braunisch, 2013; Kraus and Krumm, 2013). Tree-related microhabitats (hereafter microhabitats, sensu Larrieu *et al.*, 2018), defined as tree-born singularities, are structural elements that support various species and communities. Among the different microhabitats, Woodpecker cavities are, so far, the most documented regarding the biodiversity they harbor (Cockle *et al.*, 2011; Larrieu *et al.*, 2018). Black Woodpecker (*Dryocopus martius* L. 1758) cavities are the largest excavated by a bird in Europe, and, as such, they have a crucial role for many secondary cavity nesters (e.g. Johnsson, 1993). The Black Woodpecker excavates for nesting, but also for roosting, and both cavity types may be used by other nesters. The Black Woodpecker is qualified as an ecosystem engineer, because it significantly modifies its environment and thus allows species to live and reproduce (Johnsson, 1993; Kosiński *et al.*, 2010; Pirovano and Zecca, 2014): e.g. the Boreal Owl (*Aegolius funereus*, L. 1758), the Stock Dove (*Columba oenas*, L. 1758), the Greater Noctule bat (*Nyctalus lasiopterus*, S. 1780), the Pine Marten (*Martes martes*, L. 1758), and numerous saproxylic beetles and fungi. This effect is accentuated by the fact that this bird has a large home range (up to 500ha in Sweden and Norway, Tjernberg *et al.*, 1993; Rolstad *et al.*, 1998; 316ha in the Italian Alps, Bocca *et al.*, 2007) and also needs forests with old-growth conditions (particularly old and large trees) to accomplish its life cycle (Khanaposhtani *et al.*, 2012; Olano, 2015; but see Basile *et al.*, 2020). Understanding the factors linked with cavity excavation by the Black Woodpecker is essential to better integrate them in forest management at different scales. However, such knowledge either relies on expertise (e.g. Cuisin, 1967) or has seldom been quantified for western Europe using systematic approaches on large datasets (see Tjernberg *et al.*, 1993; Gil-Tena *et al.*, 2013; Brambilla and Saporetti, 2014).

At the tree scale, several studies have shown that the Black Woodpecker excavates in less crowded environments (i.e. with less undergrowth), on trees devoid of low branches and trees with a diameter at breast height larger than the others in the vicinity (Puverel *et al.*, 2019; Basile *et al.*, 2020). Some papers also mention fungi infection as a strong determinant of Black Woodpecker excavation (Jackson and Jackson, 2004; Zahner *et al.*, 2012). At larger scales, several stand and landscape factors are also linked with cavity excavation, notably forest fragmentation and landscape heterogeneity (e.g. Gil-

Tena *et al.*, 2013; Brambilla and Saporetti, 2014; Saporetti *et al.*, 2016). As a forest specialist, the

Black Woodpecker depends on forest habitats to reproduce (Tjernberg *et al.*, 1993; Mikusinski, 1995;

Angelstam *et al.*, 2002), but also on more open forest habitats like clearings to feed (Dorresteijn *et al.*,

2013; Brambilla and Saporetti, 2014; Pirovano and Zecca, 2014). Cavity excavation may first be

influenced by the forest cover, that would benefit both nesting and feeding of the species (Dorresteijn

et al., 2013; Pirovano and Zecca, 2014). This effect of forest cover could also be context-dependent

and stronger in heterogeneous (mixed mature stands combined with open forest habitats) and

continuous forest landscapes (see e.g. Karimi *et al.*, 2018 for an example in Iran). Then, landscape

connectivity would positively influence the Black Woodpecker presence and cavity excavation, with

higher probabilities in places where favorable patches are connected (Tobalske and Tobalske, 1999;

Gil-Tena *et al.*, 2013).

In this context, we aimed to quantify the effects of forest composition and landscape configuration on

cavity excavation by the Black Woodpecker to better integrate its cavities in forest management and

planning. We used a large database of Black Woodpecker cavities maps provided by local managers

and naturalists on nine different sites in France covering a wide range of ecological conditions (forest

types, elevation, climate). We worked at three different scales to provide management and biologically

relevant answers: (i) 10ha (0.1km2) scale that corresponds to a typical forest management unit in

France; (ii) 100ha (1km2) scale that corresponds to the mean core home range of the Black

Woodpecker where it defends its territory and breeds (Bocca *et al.*, 2007); (iii) 250ha (2.5km2) scale

that corresponds to the mean home range over the year (Olano, 2015). For each scale, we extracted

the same landscape variables to assess which ones had most influence on cavity excavation, and

understand whether the landscape factors were the same across all scales. We hypothesized that:

(i)   The presence of both broadleaf stands for nesting and conifer stands for feeding would

96   positively influence the presence of cavities;

97   (ii)   The presence of large, continuous, forest areas would positively influence the presence of

98   cavities. This would consist in zones with high forest cover and high values of connectivity

99   between patches;

100   (iii)   The landscape heterogeneity, measured as the size and distribution of the patches, would

101   negatively influence the presence of cavities.

Materials and methods

<u>Study sites and dataset</u>

We worked on nine French forest sites where Black Woodpecker cavities have been mapped by forest

managers and naturalists (Figure 1).

Figure 1: Map of the study sites.

The dataset has been compiled from different sources (National Forest Service, non-governmental

organizations, local naturalists, National and Regional Parks) and originates from different field

campaigns and methods: trees identified during tree-marking operations in forest management or

active prospections dedicated to these cavities (see Table 1 for specifications on methods and data

providers). Maps cannot be assumed to be fully exhaustive and some cavities may have been missed

by the inventories, but regarding the quantity of cavities mapped, and the methods used to analyze

this data (see below), we remain confident that our results highlight unbiased trends on the link

between cavities and landscape features. Most cavities were observed from the ground, without

relying on pole-cameras or other devices except binoculars (see discussion for the potential bias

linked to this method of observation). Since Black Woodpecker cavities cannot be confounded in the

field (Larrieu *et al.*, 2018) and because many professional birders and naturalists were involved in the

process, we assume that these inventories were reliable. It is important to note that we did not

considered only breeding cavities, but all cavities excavated by the bird. We are conscious that this is

a strong limitation in the interpretations we could draw on the bird biology, however, since the

presence of such cavities (whatever their use by the Black Woodpecker) in the landscape are crucial

for many species, we still think that studying landscape features around them is valuable in terms of

management.

We harmonized the different datasets and selected data to control for dependence and heterogeneity.

We first excluded observations made before 2010, since we considered that landscape (forest)

variables could have changed significantly since then because of e.g. tree mortality and anthropogenic

or natural disturbances (e.g. harvesting, tree dieback, storms). Cavities were rarely observed on dead

trees and snags since most managers target living trees for forest management purposes (in total 18

cavities were reported on dead trees on the initial dataset). Although they constitute quality breeding

and foraging sites for the birds, dead trees and snags are relatively ephemeral habitats that can be

more easily missed out by the inventories. We excluded the few cavities on dead trees since the

conditions of excavation may rely more on tree vitality than landscape in this case and it is also difficult

to assess whether a cavity on a dead tree was excavated before or after the tree died (Puverel *et al.*,

2019).

Based on this first selection, we kept 2689 Black Woodpecker cavities observations with spatial

coordinates (Table 1). As indicated by the data providers, all locations were measured with

commercial-grade GPS receivers or directly plotted on maps. Precision is thus comprised between

10 m and 100 m, which is consistent with the minimum scale of analysis (10ha). Note that we decided

to keep the "Ecouges" site although it contained only one cavity (removal from the dataset did not

affected the results – not shown).

Site	Cavity number	Elevation (m)		Mean annual temperature (°C)		Mean annual precipitation (mm)		Slope (deg)		Methods of cavity inventory	Data provider
		Min/Max	Mean	Min/Max	Mean			Min/Max	Mean		
Aigoual	1819	796/1398	1218	5.6/10.2	7.3	774/979	900.9	1.1/51.4	16.4		
Causses	251	821/1400	1122	5.9/10.2	7.8	769/966	864.7	1.2/48.1	22.5	Active and systematic search by foresters and National Park staff	National Forest Service (ONF) and Cévennes National Park
Mont Lozère	14	941/1187	1107	6.8/8.7	7.5	787/898	857.8	10/33.3	17.2		
Vallées Cévenoles	10	919/1248	1156	7.1/9.3	7.8	790/881	850.3	7.7/25.6	17.3		
Auberive	194	360/504	440	8.7/9.6	9.1	802/911	853	0/31.2	7.8	Inventory during tree selection operations	National Forest Service (ONF)
Ecouges	1	1218	1218	5.7	5.7	1187	1187	16.4	16.4	Active and systematic searching by natural areas managers	Conseil Départemental de l'Isère (Natural area manager)
Loches	111	91/141	124	11/11.3	11.1	690/716	703.8	0/9.7	1.9	Active searching by a naturalist	M. Dubois (naturalist)
Semnoz	37	1017/1500	1353	4.6/7.5	5.6	1154/1430	1335	4.3/28	12.2	Active and systematic searching by a NGO	Ligue pour la Protection des Oiseaux (LPO, B. Doutau)
Volcans	252	685/1154	935	6.5/8.7	7.6	765/912	841	0.9/40.7	10.3	Active and systematic searching by a regional park	Parc Naturel Régional des Volcans d'Auvergne

Table 1: Distribution and characteristics of Black Woodpecker cavity locations selected for the study

The dataset is presence-only data and we chose to generate background points to process statistical

analyses (Renner *et al.*, 2015). Random generated background points is a powerful method to deal

with presence-only data, notably because background points are less time-consuming than true

absence data to gather, even if they are sometimes less precise (e.g. Iturbide *et al.*, 2015). We

preferred this method over other presence only analysis methods (see below) because it allowed a

better control of the location and distribution of absence data.

We selected 5km radius areas around the presence locations and kept only forested areas around

(from the satellite images derived layer OCSOL 2018, https://www.theia-land.fr/ceslist/ces-occupation-

des-sols/, Inglada *et al.*, 2017). Then, we excluded an area of 250m radius around each presence

location to limit the spatial auto-correlation between presence and background point locations.

Although this distance does not insure independence of the observations (250m radius corresponds to

approximately 20ha), it represents a compromise between spatial autocorrelation and power of the

analyses.

We then compiled 2689 presence data with approximately ten times more (22326) background points

drawn at random in the envelopes described above, for a total of 25015 locations. To ensure spatial

independence between the observations, we used spatial thinning to conserve presence data that

were distant of at least 250m (i.e. we randomly selected locations that were at least 250m apart), and

selected ten-times more background point locations within each of the nine sites (R package spThin,

Aiello-Lammens *et al.*, 2015). The training dataset finally comprises 7348 locations (668 presence and

6680 background points). We also generated 10 datasets with different random spatial thinnings for

model validation (see below).

We then delineated the three buffer zones corresponding to our three study scales (radius 179m

(10ha); radius 565m (100ha) and radius 879m (250ha) around each location) and extracted forest,

landscape and biogeoclimatic variables (Table 2).

Group	Variable / Indices	Unit	Source (resolution)
Topographic and climatic	Elevation	Meters	BD ALTI v.2.0: digital terrain model (25 m)
	Slope	Percent	BD ALTI v.2.0: digital terrain model (25 m)
	Cosinus(Aspect)		BD ALTI v.2.0: digital terrain model (25 m)
	Sinus(Aspect)		BD ALTI v.2.0: digital terrain model (25 m)
	Mean annual temperature	Celsius	Worldclim v2 (1 km)
	Mean annual precipitations	Millimeters	Worldclim v2 (1 km)
Forest composition	Proportion of broadleaves	Percent	BD Forêt v2 (0.5 ha)
	Proportion of conifers	Percent	BD Forêt v2 (0.5 ha)
	Proportion of mixed forests	Percent	BD Forêt v2 (0.5 ha)
Forest diversity	Shannon diversity index for forest types		BD Forêt v2 (0.5 ha)
	Simpson diversity index for forest types		BD Forêt v2 (0.5 ha)
Forest cover	Proportion of forest cover	Percent	BD Forêt v2 (0.5 ha)
Landscape connectivity (forest patches)	Contagion index	Percent	BD Forêt v2 (0.5 ha)
	Cohesion index	Percent	BD Forêt v2 (0.5 ha)
	Aggregation index	Percent	BD Forêt v2 (0.5 ha)
Landscape heterogeneity	Landscape Shannon diversity index		Corine Land Cover (25 ha)
	Landscape Simpson diversity index		Corine Land Cover (25 ha)
Landscape fragmentation	Landscape shape index		OSO (10 m)
	Edge density	Meters / hectares	OSO (10 m)

174 Table 2: Explanatory variables and sources used to characterize the environment of Black
175 Woodpecker cavities
176

<u>Forest variables</u>

We used QGIS (version 3.16.6-Hannover) and R v 4.0.3 (R Core Team, 2020) to extract forest

characteristics in the different buffers around presence and background points locations. We extracted

forest composition summarized as the proportion of broadleaves, conifers and mixed forests types

from the BD Forêt version 2.0 (National Forest Inventory, https://inventaire -

forestier.ign.fr/IMG/pdf/dc_bdfor et_2-0.pdf, Table 2).

We then calculated Shannon and Simpson diversity indices based on the forest types issued from the

BD Forêt v2 (27 forest types in total). Both indices account for the number of types and their

abundance within each buffer, they vary from 0 (only 1 type within the buffer) to plus infinite (types

equitably distributed within each buffer). The Shannon index accounts more for rare elements than the

Simpson does, Table 2).

We finally calculated the proportion of forest within each buffer.

<u>Landscape variables</u>

Using the R package landscapemetrics (Hesselbarth *et al.*, 2019), we calculated different landscape

variables that account for the different approaches in landscape ecology: the patch characterizing a

homogeneous unit, the class characterizing all patches within a given typology and the landscape

characterizing all patches and classes within a given study window (Sertel *et al.*, 2018). We first

calculated landscape connectivity indices using the BD Forêt v2 taking into account forest cover only

(see McGarigal, 2012):

(i)   the contagion index measures the probability that two patches belong to the same class.

The index varies from 0 (a lot of dispersed patches) to 100% (large contiguous patches);

(ii)   the patch cohesion index measures the connectivity of the patch type studied. The index

also varies from 0 (when the studied class is fragmented) to 100% (when the studied

class is aggregated);

(iii) the aggregation index is calculated using a contiguity matrix and measures, for each

combination of patches, the frequency at which patches appear contiguous within a given

window. The index varies from 0 (when the landscape is totally fragmented) to 100%

(when the landscape is composed of fully contiguous patches, He *et al.*, 2000).

Second, we calculated indices characterizing landscape heterogeneity: Shannon and Simpson indices

measure the landscape diversity accounting for the number of patch classes and their abundance.

Both indices were calculated from Corine Land Cover 2018 (https://land.copernicus.eu/pan-

european/corine-land-cover/clc2018, Büttner *et al.*, 2021).

Third, we calculated several landscape fragmentation indices based on the forest cover of BD Forêt

V2:

(i)   the landscape shape index measures the shape of the study class compared to a maximal

level of aggregation within this class. The index varies from 1 (when the landscape is

composed of only one patch) to the infinite;

(ii)   the edge index measures edge length per surface unit (meters per hectare in our case).

The index varies from 0 (no edge) to infinite.

<u>Biogeoclimatic variables</u>

We extracted biogeoclimatic variables to contextualize the relationships between stand and landscape

variables and the occurrence of Black Woodpecker cavities. For each location, we extracted the

elevation, slope, aspect (cosinus and sinus transformed) based on the digital terrain model (BDAlti

V2.0) , mean annual temperature and precipitations issued of Worldclim 2 (Fick and Hijmans, 2017,

Table 2). For these variables only, we limited extraction to the location of cavity presences and

background points, and not to buffers, since we used them to contextualize the relationships between

landscape variables and presence of cavities.

<u>Statistical analyses</u>

We used R v4.0.3 (R Core Team, 2020) for all analyses. Following Zuur *et al.* (2010), we explored the

data for variance homogeneity and unaccounted sources of variation, outliers, collinearity and

interactions. We used linear mixed model to analyze the occurrence of Black Woodpecker cavities as

a function of the different explanatory variables listed above (Table 2). We fitted models using the

function glmmTMB from the glmmTMB package (Magnusson *et al.*, 2017) with binomial error

distribution for occurrence data (presences and background points), logit link and site as random

effect to account for the fact that observations from the same site may be more similar than

observations issued from two different sites. Before processing for model selection, we used the

Variance Inflation factor (VIF, package car, Fox and Weisberg, 2019) to detect multicollinearity

amongst explanatory variables and excluded variables with a VIF higher than 5 points. We then

processed a forward model selection based on Akaike Information Criterion (AIC, function stepAIC in

the package MuMIn, Barton, 2020), adding variables corresponding to our hypotheses once at the

time and limiting the interactions to first order. We standardized all variables to be able to compare the

magnitudes of the effects. For each scale (10, 100, and 250ha), we added variables and interactions

until the AIC did not changed by more than two points. We then validated the best model by fitting it on

the 10 thinned datasets and comparing the estimates for each scale. The model was judged robust if

the standard deviations of the mean estimates across the 10 datasets was not higher than the

standard errors of the estimates of the training model.

Results

<u>Model selection on the training dataset</u>

At the 10ha scale, contagion index and proportion of broadleaves showed a VIF > 5 and were

excluded. The best model comprised slope, mean annual precipitation, proportion of conifers,

proportion of mixed forests, proportion of forest cover, landscape shape index and interactions

between slope and mean annual precipitations and landscape shape index and mean annual

precipitations (Table 3). All the effects were significant (p<0.05) or marginally significant (p<0.1). In

terms of magnitude, the proportion of forest cover had the highest positive effect on the occurrence of

Black Woodpecker cavities (standardized estimate 1.56) followed by mean annual precipitation

(positive effect, standardized estimate 1.49, Figure 2). Proportion of mixed forests and slope had a

lesser positive effect (standardized estimates resp. 0.15 and 0.13, Figure 2). Conversely, proportion of

conifers and landscape shape index had negative effects on the Black Woodpecker cavities

occurrence (standardized estimates resp. -0.18 and -0.17, Figure 2). Finally, the estimate of the

interaction between slope and mean annual precipitation was negative (standardized estimate -0.12),

while the estimate for the interaction between annual mean precipitation and landscape shape index

was marginally positive (standardized estimate 0.08).

Figure 2: Effects of landscape shape index, proportion of mixed forest and mean precipitations on the

occurrence of Black Woodpecker cavities at three scales (n = 7348 locations). Plain line represents

the mean estimation derived from a generalized linear mixed model with binomial error distribution and

site as a random effect. Ribbons are 95% confidence intervals. All other variables were hold constant

(mean) for the representation.

At the 100ha scale, contagion index showed a VIF > 5 and was excluded. The best model comprised

slope, mean annual precipitations, proportion of broadleaves and mixed forests, edge density index,

landscape shape index and Shannon landscape diversity index. In addition, it comprised the

interactions between slope and mean annual precipitations, edge density index and landscape shape

index, mean annual precipitations and landscape scale index, slope and proportion of broadleaves,

proportion of broadleaves and mean annual precipitations, proportion of conifers and mean annual

precipitations and Shannon landscape diversity index and slope (Table 3). All the effects were

significant (p<0.05). In terms of magnitude, mean annual precipitations had the higher positive effect

on Black Woodpecker cavities occurrence (standardized estimate 1.49, Figure 2) followed by the edge

density index (negative effect, standardized estimate -0.66). Other positive effects comprised the

proportion of mixed forests (0.38, Figure 2), the slope (0.33), the Shannon landscape diversity index

(0.15) and the proportion of broadleaves (0.14), as well as interactions between mean annual

precipitations and landscape shape index (0.22). Landscape shape index (-0.44, Figure 2) and

interactions between slope and mean annual precipitations (-0.17), edge density and landscape shape

index (-0.20), proportion of broadleaves and slope (-0.21) and mean annual precipitations (-0.18), the

proportion of mixed forests and mean annual precipitation (-0.11) and the Shannon index for forest

types and slope (-0.12) all had negative effects.

At the 250ha scale, no variable was excluded based on VIF. The best model comprised the slope, the

mean annual precipitations, the proportion of mixed forests, the edge density index, the proportions or

broadleaves and mixed forests, the proportion of forest cover, the landscape shape index, the

Shannon landscape diversity index, and the interactions between slope and mean annual

precipitations, edge density and landscape shape index, mean annual precipitation and landscape

shape index and proportion of broadleaves and slope. All the effects were significant (p>0.05). In

terms of magnitude, mean annual temperatures had the strongest positive effect (standardized

estimate 1.34, Figure 2) followed by the positive effect of the proportion of forest cover (positive effect,

standardized estimate 0.94). Other positive effects comprised the Shannon landscape diversity index

(0.33), the slope (0.29), the proportion of mixed forests (0.14, Figure 2) as well as the interactions

between mean annual precipitation and landscape shape index (0.14). Landscape shape index (-0.39,

Figure 2), the edge density (-0.36), the proportion of broadleaves (-0.11) as well as the interactions

between slope and mean annual precipitations (-0.16), the edge density index and the landscape

shape index (-0.31) and the proportion of broadleaves and slope (-0.17) all had negative effects.

Variable	10ha scale				100ha scale				250ha scale			
	Estimate	SE	p		Estimate	SE	p		Estimate	SE	p	
Intercept	-4.346	0.937	<0.001	***	-4.075	0.921	<0.001	***	-4.035	0.905	<0.001	***
Slope	0.134	0.067	0.047	*	0.334	0.067	<0.001	***	0.286	0.067	<0.001	***
Mean annual precipitation	1.488	0.083	<0.001	***	1.491	0.089	<0.001	***	1.335	0.091	<0.001	***
%broadleaves					0.141	0.056	0.012	*	-0.110	0.054	0.043	*
%conifers	-0.182	0.052	0.000	***								
%mixed forests	0.152	0.038	<0.001	***	0.384	0.047	<0.001	***	0.138	0.044	0.002	**
Edge density					-0.661	0.076	<0.001	***	-0.364	0.082	<0.001	***
% forest cover	1.563	0.127	<0.001	***					0.938	0.099	<0.001	***
Landscape shape index	-0.170	0.068	0.012	*	-0.447	0.092	<0.001	***	-0.386	0.092	<0.001	***
Shannon landscape diversity					0.158	0.056	0.005	**	0.332	0.061	<0.001	***
Slope x Precipitation	-0.120	0.045	0.008	**	-0.171	0.049	<0.001	***	-0.163	0.047	<0.001	***
Edge density x Landscape shape index					-0.197	0.078	0.012	*	-0.305	0.077	<0.001	***
Landscape shape index x Precipitation	0.083	0.050	0.095	(*)	0.223	0.058	<0.001	***	0.144	0.060	0.016	*
%broadleaves x Slope					-0.207	0.050	<0.001	***	-0.166	0.049	<0.001	***
%broadleaves x Precipitation					-0.178	0.050	<0.001	***				
%mixed forests x precipitation					-0.105	0.024	<0.001	***				
Shannon landscape diversity x Slope					-0.121	0.054	0.024	*				

Table 3. Standardized estimates of the best models predicting the occurrence of Black Woodpecker

cavities at three study scales (10ha, 100ha and 250ha). Grey cells correspond to variables not

selected in the models. SE: Standard Error. *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$, (*) $p < 0.1$.

<u>Model validation</u>

We fitted the selected model for each scale on the 10 different thinned validation datasets. Globally

the standardized estimates were consistent for the different datasets with standard deviations of the

values below – or the same order of magnitude than – the standard errors of the best models

(Table 4).

Variable	10ha scale	100ha scale	250ha scale

Intercept	0.0731	0.0776	0.0725
Slope	0.0210	0.0279	0.0272
Mean annual precipitations	0.0269	0.0348	0.0242
%broadleaves		0.0161	0.0257
%conifers	0.0217		
%mixed forests	0.0107	0.0231	0.0214
Edge density		0.0188	0.0359
% forest cover	0.0420		0.0218
Landscape shape index	0.0251	0.0193	0.0208
Shannon landscape diversity		0.0143	0.0193
Slope x Precipitation	0.0198	0.0191	0.0141
Edge density x Landscape shape index		0.0269	0.0260
Landscape shape index x Precipitation	0.0219	0.0270	0.0313
%broadleaves x Slope		0.0233	0.0207
%broadleaves x Precipitation		0.0188	
%mixed forests x precipitation		0.0136	
Shannon landscape diversity x Slope		0.0280	

Tableau 4. Mean standard errors of estimates for the ten validation datasets.

Discussion

<u>Heterogeneous and diversified forest types favour cavity excavation by the Black Woodpecker</u>

We partially validated the hypothesis that mixed forests favoured the excavation of cavity by the Black

Woodpecker. Indeed, landscapes composed of broadleaves and conifers positively influenced the

presence of cavities: the probability of cavity occurrence increased from 14 to 47% when the

proportion of mixed forests in the landscape increased from 50 to 75% at the 100ha scale, and from

25 to 68% at the 250ha scale (Figure 2). Mixed broadleaf and coniferous forests favour the presence

of the Black Woodpecker since broadleaves are preferentially selected for cavity excavation

(Fernandez and Azkona, 1996; Kosiński and Kempa, 2007; Karimi *et al.*, 2018; Basile *et al.*, 2020)

while conifers provide food resources, in particular wood and carpenter ants (Mikusiński, 1997) and

saproxylic larvae (Rolstad *et al.*, 1998; Brambilla and Saporetti, 2014). Indeed, in central and western

Europe, the Black Woodpecker seems to excavate cavities preferentially in beech (Fagus sylvatica L.),

while in Scandinavia, the bird excavates more on conifers like Scots Pine (Pinus sylvestris L.,

Angelstam and Mikusiński, 1994; Pirovano and Zecca, 2014). Tree species with smooth bark, like

beech, would be preferred to prevent predation by the Pine Marten since climbing on such trees

species seems more difficult (Kosiński and Kempa, 2007; Olano, 2015; Zahner *et al.*, 2017; Puverel *et*

al., 2019). In the Alps, beech stands would also be favoured because they present larger mean

diameter due to a less intensive harvesting and lower densities compared to other forests (De Rosa *et*

al., 2016). The Black Woodpecker may also avoid excavating cavities in coniferous stands with low

branches, that favour predation (Bocca *et al.*, 2007). This is confirmed by our results on forest

composition at different scales. At the 10ha scale, although the variable proportion of broadleaves has

not been selected, the proportion of conifers had a negative effect on the presence of cavities. At the

100ha scale, that corresponds to the core zone of the home range of the species (nesting, breeding,

and active defence of the territory), the effect of broadleaves proportion became positive. Conversely,

at the larger 250ha scale, this effect was negative. Therefore, the Black Woodpecker avoids conifers

at the local scale and selects broadleaves in the core zone of its home range, while the presence of

conifers at the larger scale may be positive as a feeding site. However, since we did not observe

significant effects of forest diversity of forest types on the cavity presence at the two larger scales

(Shannon and Simpson indices of forest types), the presence of cavities seems primarily influenced by

the abundance of broadleaves.

<u>Black Woodpecker cavities occur more in unfragmented forest landscapes</u>

The quantity of edges (between forest and non-forest areas) negatively affected the presence of Black

Woodpecker cavities, which validated our second hypothesis. Indeed, the occurrence of cavities

decreased significantly with increasing edge abundance, and was almost null when the proportion of

edge reached 100m per hectare (Table 3). We observed the larger impact of edges at the 100ha

scale, which shows that the Black Woodpecker needs large continuous forest areas to excavate in its

core home range. The positive effect of the contagion index at the three scales confirmed this

statement: the occurrence probability doubled when the index increased from 80 to 100 (Table 3).

While the Black Woodpecker may cover large distances for foraging, it seems that large unfragmented

forest areas are necessary for cavity excavation. However, this result does not seem to be

generalizable over the whole distribution of the species. Indeed, in Sweden, Tjernberg *et al.* (1993)

showed that the Black Woodpecker also occurs in fragmented forest landscapes, as long as it can fly

from one feeding ground to another. Thus, it can live in landscapes with as little as 26% of forested

areas (Dorresteijn *et al.*, 2013) which confirms the contextualised response of the species to
fragmentation.

We did not validate our third hypothesis on landscape heterogeneity. Indeed, we showed that
increasing heterogeneity favoured the occurrence of the Black Woodpecker cavities, although the
magnitude of this effect was low. The positive effect of heterogeneity may be linked to the fact that the
species also need some open, non-forested habitats, eventually with patches of conifer, to feed, and
that these zones may be located in the home range (translated in our study by an increasing effect at
the 250ha scale).

<u>The probability of cavity excavation is context dependant</u>

Slope and mean annual precipitation were often selected as variables characterizing the general
context as well as in the interactions of the model. While occurrence of cavities generally increased
with slope and precipitations across all scales, these variables also had a negative effect on the
interactions with other variables, notably composition (Table 3). Although elevation was not selected in
our models, it is clear that higher slopes and precipitations are characteristics of mountainous
landscapes, where the effects of forest composition are less pronounced than in lowlands, but where
cavities are also less abundant. These effects probably reflect other factors not included in these
analyses like tree diameter, a minimum of which is necessary for the Black Woodpecker to excavate a
suitable cavity (e.g. Basile *et al.*, 2020).

<u>Limitations and research perspectives</u>

The dataset we analysed was, by essence, heterogeneous since it originated from management and
naturalist prospections without a real systematic approach. Ideally, it would have been worth verifying
at least a sample of the data in the field, for example using a pole camera since, from the ground, full
cavities may be confounded with cavity initiations (Cuisin, 1967). In terms of model, this generates
false positives that may bias the results and that could be taken into account by adding a detection
probability term (aka. N-mixture models). However, given the total number of observations, verifying
all the cavities would have been impossible, and we also assume that the bias remains marginal and

not systematic, since the birds very probably initiate cavities in conditions similar as for full cavities

excavation. Some observers may have also indirectly controlled the effective presence of cavities by

observations of nesting animals or insects, but such validation was not systematically reported. In

addition, model validation did not show strong variations when changing the dataset, we are thus

confident that our results are robust, but would be more precise with full cavity inventories. Such

modelling approach would also be more informative with nesting cavities only, but at such a scale,

would require huge fieldwork to have a dataset allowing the analysis. We think our approach is a good

compromise regarding the value of the compiled dataset, and remains informative regarding the role

of cavities for other species than the Woodpecker itself.

We voluntarily excluded dead trees from the dataset, in order to limit the noise due other uncontrolled

factors like tree dieback. We were conscious that this may have biased the results since dead trees

and snags are an important part of Woodpecker foraging and breeding niche (Rolstad *et al.*, 1998;

Angelstam *et al.*, 2002). However, living trees are under the direct control of forest managers since

they have a production as well as a conservation role. As such, it is likely (but difficult to verify) that

living trees were more systematically observed than dead trees in the inventory process. For this

methodological reason at least, it seems more reasonable to exclude dead trees and snags being

conscious that this may bias our results.

More generally, we could not analyse the effects of forest structure and precise composition at

different scales around cavities observations, notably the degree of stand maturity and the quantity of

deadwood at the large scale. Such data would have been interesting to specify the needs of the

species, notably in terms of food resources linked to deadwood. These variables could be derived

from airborne LiDAR interpretation (Zellweger *et al.*, 2013; Fuhr *et al.*, 2022), but such data was not

available across all the sites we studied, and would have required additional field data to calibrate

LiDAR models. In a near future, the ongoing French nation-wide LiDAR acquisition program could help

completing the present study, using the same processes of data selection, as well as specifying the

requirements of the Black Woodpecker in terms of forest structure. Also, composition could be

assessed using multispectral data such as Sentinel 2 (e.g. Grabska *et al.*, 2019). More precise

mapping of forest and landscape attributes could also be of use in the study of cavity excavation by

the Black Woodpecker. For example, mapping issued of the BD Forêts v2 has a resolution of 0.5ha,

which may cause noise for the results at the 10ha scale, especially for landscape indices (such as

fragmentation, contagion, etc). However, it was the most precise data systematically available for the

whole dataset, and our results are reproducible since they gather field data and public data sources.

Finally, generalisation of our approach in other locations of the distribution range of the species would

be interesting, since it seems that its ecological preferences may vary with the biogeographical

context. In turn, all this would help the preservation of this keystone species for the forest ecosystem,

and the cavities excavated that support numerous secondary species from other taxa.

<u>Conclusions and implications for forest management.</u>

We studied the forest composition and landscapes variables that influenced cavity excavation by the

Black Woodpecker. From a strict forest management point of view, landscape variables are rarely

under the control of forest management. However, some results may be transferred to management

planning at the forest or a larger scale. At the 10ha (management unit) scale, we showed that forest

composition (mixed forests) was the most influential variable, while at larger 100ha (core home range)

and 250ha (home range) scales, fragmentation and connectivity had stronger effects. Therefore,

maintaining and favouring mixed stands locally would be beneficial to the Black Woodpecker and all

the species that depend on its cavities for living. Mixed forests have proved to be beneficial both for

biodiversity conservation and a large set of other services (e.g. Liang *et al.*, 2016). Black Woodpecker

cavities seem to benefit from either a mix at the stand scale or the juxtaposition of broadleaved and

coniferous stands, providing that broadleaved stands are large enough. Such action should then be

integrated at the management planning level. At a larger scale, large continuous forested areas reveal

the forest character of the Black Woodpecker, but also the need to maintain connectivity at the

landscape scale to favour forest species. Cavity excavation by the Black Woodpecker is also sensitive

to edges, as are many other forest bird species, the conservation of which is then conditioned by the

conservation of large unfragmented areas (Robinson *et al.*, 1995; Villard *et al.*, 1999). These

recommendations may ultimately be applied at the larger scale, e.g. for a regional or national parks,

that have an objective of biodiversity conservation.

Acknowledgements

We are indebted to all forest managers, naturalists and other Woodpecker lovers who contributed to

cavity mapping. Without their huge work and dedication, this study would not have been possible.

B. Algoët, H. Caroff, D. Joud, A. Savine, A. Porte, L. Belenger and B. Guérin especially facilitated the

data exchange and interpretation. CC did all the analyses during the Master's degree training period.

Literature cited

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B., Anderson, R.P., 2015. spThin: An R
package for spatial thinning of species occurrence records for use in ecological niche models.
Ecography 38, 541-545.

Angelstam, P., Breuss, M., Mikusiński, G., Stenstrom, M., Stighall, K., Thorell, D., 2002. Effects of
forest structure on the presence of woodpeckers with different specialisation in a landscape history
gradient in NE Poland. Avian Landscape Ecology: Pure and Applied Issues in the Large-Scale
Ecology of Birds, 25-38.

Angelstam, P., Mikusiński, G., 1994. Woodpecker assemblages in natural and managed boreal and
hemiboreal forest — a review. Ann Zool Fenn 31, 157-172.

Barton, K., 2020. MuMIn: multi-model inference. in R package version 1.43.13. In.

Basile, M., Asbeck, T., Pacioni, C., Mikusiński, G., Storch, I., 2020. Woodpecker cavity establishment
in managed forests: Relative rather than absolute tree size matters. Wildl Biol 2020.

Bocca, M., Carisio, L., Rolando, A., 2007. Habitat use, home ranges and census techniques in the
Black Woodpecker *Dryocopus martius* in the Alps. Ardea 95, 17-29.

Bollmann, K., Braunisch, V., 2013. To integrate or to segregate: balancing commodity production and
biodiversity conservation in European forests. In: Kraus, D., Krumm, F. (Eds.), Integrative approaches
as an opportunity for the conservation of forest biodiversity. European Forest Institute, Freiburg,
Germany, pp. 18-31.

Brambilla, M., Saporetti, F., 2014. Modelling distribution of habitats required for different uses by the
same species: Implications for conservation at the regional scale. Biol Conserv 174, 39-46.

Büttner, G., Kosztra, B., Maucha, G., Pataki, R., Kleeschulte, S., Hazeu, G., Vittek, M., Schröder, C.,
Littkopf, A., 2021. CORINE Land Cover User manual. In: Copernicus Land Monitoring Service, E.E.A.
(Ed.). Copernicus Land Monitoring Service, European Environment Agency.

Cockle, K.L., Martin, K., Wesołowski, T., 2011. Woodpeckers, decay, and the future of cavity-nesting
vertebrate communities worldwide. Frontiers in Ecology and the Environment 9, 377-382.

Cuisin, M., 1967. Essai d'une monographie du Pic noir (*Dryocopus martius* (L.)). L'Oiseau et la Revue
française d'Ornithologie 37, 164-224.

De Rosa, D., Andriuzzi, W.S., Di Febbraro, M., 2016. Breeding habitat selection of the black
woodpecker dryocopus martius L. in mediterranean forests. Avocetta 40, 63-69.

Dorresteijn, I., Hartel, T., Hanspach, J., von Wehrden, H., Fischer, J., 2013. The conservation value of

traditional rural landscapes: the case of woodpeckers in Transylvania, Romania. PLoS ONE 8.

Fernandez, C., Azkona, P., 1996. Influence of forest structure on the density and distribution of the

white-backed woodpecker *Dendrocopos leucotos* and black woodpecker *Dryocopus martius* in quinto

real (spanish western pyrenees). Bird Study 43, 305-313.

Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global

land areas. International Journal of Climatology 37, 4302-4315.

Fox, J., Weisberg, S., 2019. An R Companion to Applied Regression, Third Edition. Thousand Oaks,

CA: Sage.

Fuhr, M., Lalechère, E., Monnet, J.M., Bergès, L., 2022. Detecting overmature forests with airborne

laser scanning (ALS). Remote Sens. Ecol. Conserv.

Gil-Tena, A., Brotons, L., Fortin, M.J., Burel, F., Saura, S., 2013. Assessing the role of landscape

connectivity in recent woodpecker range expansion in Mediterranean Europe: Forest management

implications. European Journal of Forest Research 132, 181-194.

Grabska, E., Hostert, P., Pflugmacher, D., Ostapowicz, K., 2019. Forest stand species mapping using

the Sentinel-2 time series. Remote Sens. 11, 1197.

He, H.S., DeZonia, B.E., Mladenoff, D.J., 2000. An aggregation index (AI) to quantify spatial patterns

of landscapes. Landsc Ecol 15, 591-601.

Hesselbarth, M.H.K., Sciaini, M., With, K.A., Wiegand, K., Nowosad, J., 2019. landscapemetrics : an

open‑source R tool to calculate landscape metrics. Ecography 42, 1648-1657.

Inglada, J., Vincent, A., Arias, M., Tardy, B., Morin, D., Rodes, I., 2017. Operational high resolution

land cover map production at the country scale using satellite image time series. Remote Sens. 9, 95.

Iturbide, M., Bedia, J., Herrera, S., del Hierro, O., Pinto, M., Gutiérrez, J.M., 2015. A framework for

species distribution modelling with improved pseudo-absence generation. Ecol Model 312, 166-174.

Jackson, J.A., Jackson, B.J.S., 2004. Ecological Relationships Between Fungi and Woodpecker

Cavity Sites. The Condor 106, 37-37.

Johnsson, K., 1993. The Black Woodpecker *Dryocopus martius* as a keystone species in forest.

Karimi, S., Moradi, H.V., Rezaei, H.R., Brambilla, M., Ghadimi, M., 2018. Fine-scale habitat use by

Black Woodpecker *Dryocopus martius* : a year-round study in the Hyrcanian forest, Iran. North-

Western Journal of Zoology 14.

Khanaposhtani, M.G., Najafabadi, M.S., Kaboli, M., Farashi, A., Spiering, D., 2012. Habitat

requirements of the black woodpecker, *Dryocopus martius*, in hyrcanian forests, Iran. Zool Middle East

55, 19-25.

Kosiński, Z., Bilińska, E., Dereziński, J., Jeleń, J., 2010. The Black Woodpecker *Dryocopus martius*

and the European Beech *Fagus sylvatica* as keystone species for the Stock Dove *Columba oenas* in

western Poland. Ornis Polonica 51, 1-13.

Kosiński, Z., Kempa, M., 2007. Density, distribution and nest-sites of woodpeckers picidae, in a

managed forest of western Poland. Pol J Ecol 55, 519-533.

Kraus, D., Krumm, F., 2013. Integrative approaches as an opportunity for the conservation of forest

biodiversity.

Larrieu, L., Paillet, Y., Winter, S., Bütler, R., Kraus, D., Krumm, F., Lachat, T., Michel, A.K., Regnery, B., Vandekerkhove, K., 2018. Tree related microhabitats in temperate and Mediterranean European forests: a hierarchical typology for inventory standardization. Ecol Indic 84, 194-207.

Liang, J., Crowther, T.W., Picard, N., Wiser, S., Zhou, M., Alberti, G., Schulze, E.D., McGuire, A.D., Bozzato, F., Pretzsch, H., De-Miguel, S., Paquette, A., Hérault, B., Scherer-Lorenzen, M., Barrett, C.B., Glick, H.B., Hengeveld, G.M., Nabuurs, G.J., Pfautsch, S., Viana, H., Vibrans, A.C., Ammer, C., Schall, P., Verbyla, D., Tchebakova, N., Fischer, M., Watson, J.V., Chen, H.Y.H., Lei, X., Schelhaas, M.J., Lu, H., Gianelle, D., Parfenova, E.I., Salas, C., Lee, E., Lee, B., Kim, H.S., Bruelheide, H., Coomes, D.A., Piotto, D., Sunderland, T., Schmid, B., Gourlet-Fleury, S., Sonké, B., Tavani, R., Zhu, J., Brandl, S., Vayreda, J., Kitahara, F., Searle, E.B., Neldner, V.J., Ngugi, M.R., Baraloto, C., Frizzera, L., Bałazy, R., Oleksyn, J., Zawiła-Niedźwiecki, T., Bouriaud, O., Bussotti, F., Finér, L., Jaroszewicz, B., Jucker, T., Valladares, F., Jagodzinski, A.M., Peri, P.L., Gonmadje, C., Marthy, W., O'Brien, T., Martin, E.H., Marshall, A.R., Rovero, F., Bitariho, R., Niklaus, P.A., Alvarez-Loayza, P., Chamuya, N., Valencia, R., Mortier, F., Wortel, V., Engone-Obiang, N.L., Ferreira, L.V., Odeke, D.E., Vasquez, R.M., Lewis, S.L., Reich, P.B., 2016. Positive biodiversity-productivity relationship predominant in global forests. Science 354.

Magnusson, A., Skaug, H.J., Nielsen, A., Berg, C.W., Kristensen, K., Maechler, M., van Bentham, K.J., Bolker, B.M., Brooks, M.E., 2017. glmmTMB: Generalized Linear Mixed Models using Template Model Builder. In.

McGarigal, K., 2012. FRAGSTATS v4: Spatial Pattern Analysis Program for Categorical and Continuous Maps. In, University of Massachusetts, Amherst, MA, p. 77.

Mikusinski, G., 1995. Population trends in black woodpecker in relation to changes and characteristics of European forests. Ecography 18, 363-369.

Mikusiński, G., 1997. Winter foraging of the Black Woodpecker *Dryocopus martius* in managed forest in south-central Sweden. Ornis Fenn 74, 161-166.

Olano, M., 2015. Black woodpecker *Dryocopus martius* (L., 1758) distribution, abundance, habitat use and breeding performance in a recently colonized region in SW Europe. Munibe, Cienc. Nat. 63, 049-071.

Pirovano, A.R., Zecca, G., 2014. Black woodpecker (*Dryocopus martius*) habitat selection in the italian alps: Implications for conservation in Natura 2000 network. Bird Conserv Int 24, 299-315.

Puverel, C., Abourachid, A., Böhmer, C., Leban, J.M., Svoboda, M., Paillet, Y., 2019. This is my spot: What are the characteristics of the trees excavated by the Black Woodpecker? A case study in two managed French forests. For Ecol Manag 453.

R Core Team, 2020. R: A language and environment for statistical computing. In. R Foundation for Statistical Computing, Vienna, Austria.

Renner, I.W., Elith, J., Baddeley, A., Fithian, W., Hastie, T., Phillips, S.J., Popovic, G., Warton, D.I., 2015. Point process models for presence-only analysis. Meth Ecol Evol 6, 366-379.

Robinson, S.K., Thompson, F.R., Donovan, T.M., Whitehead, D.R., Faaborg, J., 1995. Regional forest fragmentation and the nesting success of migratory birds. Science 267, 1987-1990.

Rolstad, J., Majewski, P., Rolstad, E., 1998. Black woodpecker use of habitats and feeding substrates
in a managed Scandinavian forest. The Journal of Wildlife Management 62, 11-23.

Saporetti, F., Colaone, S., Guenzani, W., Zarbo, T., 2016. Nest-site characteristics and breeding
biology of the Black Woodpecker in north-western Lombardy, Italy. Rivista Italiana di Ornitologia 86,
39-48.

Sertel, E., Topaloğlu, R., Şallı, B., Yay Algan, I., Aksu, G., 2018. Comparison of landscape metrics for
three different level land cover/land use maps. IJGI 7, 408.

Tjernberg, M., Johnsson, K., Nilsson, S.G., 1993. Density variation and breeding success of the Black
Woodpecker Dryocupus martius in relation to forest fragmentation. Ornis Fenn 70, 155-162.

Tobalske, C., Tobalske, B.W., 1999. Using atlas data to model the distribution of woodpecker species
in the Jura, France. The Condor 101, 472-483.

Villard, M.A., Kurtis Trzcinski, M., Merriam, G., 1999. Fragmentation effects on forest birds: Relative
influence of woodland cover and configuration on landscape occupancy. Conserv Biol 13, 774-783.

Zahner, V., Bauer, R., Kaphegyi, T.A.M., 2017. Are Black Woodpecker (*Dryocopus martius*) tree
cavities in temperate Beech (*Fagus sylvatica*) forests an answer to depredation risk? Journal of
Ornithology 158, 1073-1079.

Zahner, V., Sikora, L., Pasinelli, G., 2012. Heart rot as a key factor for cavity tree selection in the Black
woodpecker. For Ecol Manag 271, 98-103.

Zellweger, F., Braunisch, V., Baltensweiler, A., Bollmann, K., 2013. Remotely sensed forest structural
complexity predicts multi species occurrence at the landscape scale. For Ecol Manag 307, 303-312.

Zuur, A.F., Ieno, E.N., Elphick, C.S., 2010. A protocol for data exploration to avoid common statistical
problems. Meth Ecol Evol 1, 3-14.

582