Using a Whole Genome Co-expression Network to Inform the Functional Characterisation of Predicted Genomic Elements from *Mycobacterium tuberculosis* Transcriptomic Data

Authors

Jennifer Stiens, Yen Yi Tan, Rosanna Joyce, Kristine B. Arnvig, Sharon L. Kendall, Irene Nobeli

ABSTRACT

A whole genome co-expression network was created using *Mycobacterium tuberculosis* transcriptomic data from publicly available RNA-sequencing experiments covering a wide variety of experimental conditions. The network includes expressed regions with no formal annotation, including putative short RNAs and untranslated regions of expressed transcripts, along with the protein-coding genes. These unannotated expressed transcripts were among the best-connected members of the module sub-networks, making up more than half of the 'hub' elements in modules that include protein-coding genes known to be part of regulatory systems involved in stress response and host adaptation. This dataset provides a valuable resource for investigating the role of non-coding RNA, and conserved hypothetical proteins, in transcriptomic remodelling. Based on their connections to genes with known functional groupings and correlations with replicated host conditions, predicted expressed transcripts can be screened as suitable candidates for further experimental validation.

Abbreviations

CDS, coding sequence

ME, module eigengene

MM, module membership

Mtb, *Mycobacterium tuberculosis*

MTBC, *Mycobacterium tuberculosis* complex

ncRNA, non-coding RNA

ORF, open reading frame

RNA-seq, RNA sequencing

RNAP, RNA polymerase

42   sORF, short open reading frame

43   sRNA, short non-coding RNA

44   TSS, transcription start site

45   UTR, untranslated region

46   WGCNA, weighted gene co-expression analysis

INTRODUCTION

Tuberculosis continues to be a leading cause of death worldwide, causing over 1.5 million deaths, and infecting over 10 million people in 2020 (World Health Organization, 2021). The human-adapted pathogen causing tuberculosis, *Mycobacterium tuberculosis* (Mtb), has a complex lifestyle that requires rapid adaptation to host defences and immune pressure, including nutritional immunity, hypoxia and lipid-rich environments. In order to eradicate the disease, it is crucial to understand how the pathogen survives attacks from host immune cells and persists in an extended latent state inside the host. To adapt to these environmental challenges, bacterial cells must make complex transcriptomic adjustments, and these are thought to be complemented and fine-tuned by post-transcriptional regulation.

The mycobacterial genome produces a range of conditionally expressed transcripts, many of which are poorly annotated and understood. In this paper, 'non-coding' RNA (ncRNA) refers to non-ribosomal RNA transcripts not known to be translated into peptides, such as short RNAs (sRNAs) acting on either distant or antisense mRNA targets and the expressed untranslated regions (UTRs) flanking coding genes (which may also contain short open reading frames (sORFs), upstream from coding regions). Non-coding RNA can alter the abundance of gene products by controlling mRNA stability and processing, access to ribosome binding sites and the translation of overlapping open reading frames (ORFs). Discovering the contribution of the non-coding genome to specific adaptation-response pathways may improve our ability to design therapeutics and prevent the evolution of persistent phenotypes.

Uncovering the role of non-coding RNA in adaptation and transcriptomic remodelling

The proportion of non-ribosomal, ncRNA in the Mtb transcriptome has been shown to increase in stationary and hypoxic conditions, indicating a potential role in adjusting to environmental cues (Aguilar-Ayala et al., 2017; Arnvig et al., 2011; Gerrick et al., 2018; Ignatov et al., 2015). Several mycobacterial ncRNA transcripts (particularly, sRNA) have been extensively studied and found to be associated with regulatory systems controlling adaptation to stress conditions or growth phase, linked to virulence pathways and to access to lipid media (Arnvig et al., 2011; Gerrick et al., 2018; Girardin & McDonough, 2020; Mai et al., 2019; Moores et al., 2017; Solans et al., 2014). Non-coding regulation in Mtb appears to function quite differently compared to model organisms, eschewing the use of any known chaperone proteins for RNA-RNA interactions and with few sRNA homologs found outside the phyla (Gerrick et al., 2018; Mai et al., 2019; Schwenk & Arnvig, 2018). The discovery and characterisation of ncRNA in Mtb, especially sRNAs, has progressed using both molecular biology methods and high-throughput sequence-based approaches (reviewed in Schwenk & Arnvig, 2018) but characterising the gene interactions of a particular sRNA is an experimentally-expensive process, and the number of fully-characterised ncRNAs remains limited. Annotation of identified transcripts remains incomplete, as well, with only 30 listed in the Mtb H37Rv reference sequence (GenBank AL123456.3). Efforts to compile a comprehensive list of annotated ncRNAs for Mtb faces challenges of non-standardised nomenclature, different standards of experimental validation, incomplete reference annotations (especially for the animal-adapted species of the Mycobacterium tuberculosis complex (MTBC)) and the variable expression of non-coding transcripts in response to different experimental conditions (Stiens et al., 2022).

Prediction of ncRNA from RNA-sequencing (RNA-seq) data in the compact Mtb genome is challenging. Paradoxically, more sensitive, high-depth sequencing can make it more difficult to identify the small, low-abundance, functional transcripts above stochastic gene expression and technical noise. Parameters of detection must therefore be carefully considered for each dataset to account for variation in expression levels. Though RNA-seq-based ncRNA prediction algorithms are often assumed to overpredict putative ncRNAs, especially at the 5' and 3' ends of coding genes, there are biological and technical reasons for detecting abundant signal in the unannotated regions of the genome. Ribosome profiling (Ribo-seq) methods that sequence the ribosome-protected fragments of mRNA have identified actively translated RNA in the 5' UTRs of annotated protein-coding mRNA transcripts (Canestrari et al., 2020; D'Halluin et al., 2022; Sawyer et al., 2021; Shell et al., 2015; C. Smith et al., 2022). These unannotated sORFs may represent functional peptides or function to regulate the translation of the downstream transcript; however, it is impossible to tell the difference between a putative ncRNA and a sORF from RNA-seq signal alone. Additionally, post-transcriptional processing may be the norm for prokaryotes at both the 5' and 3' ends of coding transcripts, with 3' ends in mycobacteria often lacking clear signal termination (Dar & Sorek, 2018; D'Halluin et al., 2022; Wang et al., 2019). Finally, polycistronic transcripts often include non-coding sequence between the genes of an operon, and this may contain functional elements and/or processing sites (Martini et al., 2019).

The location of a transcription start site (TSS) in the 5' end of a predicted transcript supports the biological relevance of a predicted ncRNA. However, the available lists of Mtb TSS sites (Cortes et al., 2013; Shell et al., 2015) have been mapped only in starvation and exponential growth and may not include TSSs that are expressed under different experimental conditions. New TSS maps, published subsequent to this analysis may

increase the number of predicted transcripts with a TSS (D'Halluin et al., 2022).

Furthermore, functional ncRNA elements generated from the 3' UTRs of coding genes

through RNase processing would presumably lack a TSS. 3' UTRs that are functionally

independent from their cognate coding sequence (CDS) have been identified in other

bacteria (Desgranges et al., 2021; Menendez-Gil et al., 2020; Ponath et al., 2022).

Therefore, it is important to consider predicted UTRs as separate annotated elements

from protein-coding transcripts when quantifying differential expression.

To include a complete picture of the interaction of the non-coding genome with coding

genes involved in adaptation pathways, we have generated a novel set of ncRNA sequence-

based predictions (sRNAs and UTRs) from the same datasets using our in-house software

package, *baerhunter* (Ozuna et al., 2019). Some of these predicted transcripts overlap with

predictions from previous studies, but many represent novel predictions. The expression

of these transcripts is quantified along with the protein-coding genes and used in network

analysis to provide a more complete picture of the functional groupings involved in

adaptation to environmental changes. Including a variety of culture conditions that

replicate aspects of the host environment improves the chances that the expression of any

ncRNA that is restricted to one or more conditions is included in the network (Ami et al.,

2020).

Using WGCNA to implicate functional associations of non-coding RNA
146
147
148 Weighted gene co-expression network analysis (WGCNA) (B. Zhang & Horvath, 2005) has

149 been widely used to identify functional groups of genes, called 'modules', through the

150 application of hierarchical clustering to differential expression levels of RNA transcripts

151 in microarray or RNA-seq experiments. Recent studies have focussed entirely on the

152 protein-coding portion of the transcriptome, using WGCNA with RNA-seq to cluster the

differentially expressed genes of *Mycobacterium marinum* in response to resuscitation after hypoxia (Jiang et al., 2020) and *Mycobacterium aurum* infected macrophages (Lu et al., 2021). Mtb microarray data have been used to cluster protein-coding genes that show differential expression among species-specific strains (Puniya et al., 2013) and in response to two different hypoxic models to identify potential transcription factors (Jiang et al., 2016). Another recent network analysis, using a matrix deconvolution method followed by module clustering, uses a large number of RNA-seq samples including deletion mutants, infection models and antibiotic-treated samples as well as restricted media and culture conditions (Yoo, et al., 2022). They identify 80 modules of protein-coding genes that each approximate an isolated source of variance, together estimated to account for 61% of the total variance seen in in the dataset. This proportion is reportedly lower than results from similar analyses in other organisms, potentially due to the bias in the types of conditions available in the database and/or the complex nature of regulation in Mtb (Yoo, et al., 2022). However, the contribution of regulatory ncRNA elements may be a considerable unexplored source of variance in this complex system. Here we use an alternative, complementary approach by including ncRNA, as well as annotated protein-coding genes, in the modules.

In this study, WGCNA was applied to multiple Mtb H37Rv datasets covering 15 different culture conditions replicating various growth conditions, nutrient sources and stressors encountered in the host environment. We present a global view of the non-coding genome across an extensive WGCNA network and interrogate selected modules to identify functional groupings between protein-coding and non-coding transcripts, as well as between well-characterised genes and those with little functional annotation. The correlation of the modules with the various conditions can identify participants in large-

178 scale transcriptomic remodelling programs in response to changes in environmental

179 conditions.

182 MATERIALS AND METHODS

184 The overall workflow for this analysis is presented in Figure 1. All scripts for *baerhunter*,

185 WGCNA and subsequent analysis are available at:

186 https://github.com/jenjane118/mtb_modules.

188 Figure 1. Analysis workflow

189
190

Data Acquisition and Mapping

Datasets were downloaded from SRA (https://www.ncbi.nlm.nih.gov/sra/docs/) and Array Express (https://www.ebi.ac.uk/arrayexpress/) using the accession numbers listed in Table 1. To minimise batch effects and ensure compatibility with RNA prediction software, we limited analysis to datasets with similar library strategies. Samples were included based on inspection to confirm that 1) samples were from monocultures of wild-type Mtb H37Rv strain and 2) sequencing was using a paired-end, stranded protocol. Reads from samples that passed quality control thresholds were trimmed using *Trimmomatic* (Bolger et al, 2014) to remove adapters and low-quality bases from the 5' and 3' ends of the sequences. Trimmed reads were mapped to the H37Rv reference genome (GenBank AL123456.3) using *BWA-mem* in paired-end mode (Li, Heng, 2013). All samples had >70% percent reads mapped with an overall mean of ~ 27.75M mapped reads and a range of 3.97M to 60.68M mapped reads per sample (Supp Table 1, 'Samples' tab).

Table 1. Datasets used in analysis. Accession numbers from SRA and Array Express.

Dataset	Num of samples	Instrument	Library Layout	Library Strand	Library Strategy	Avg Spot Length	Ribo depleted
PRJEB65014_3 E-MTAB-6011	3	Illumina MiSeq	paired end	reversely stranded	cDNA	150	Y
PRJNA278760 GSE67035	22	Illumina HiSeq 2000	paired end	reversely stranded	cDNA	50	Y
PRJNA327080 GSE83814	15	Illumina HiSeq 2000	paired end	reversely stranded	cDNA	180	Y
PRJNA390669 GSE100097	12	Illumina NextSeq 500	paired end	reversely stranded	cDNA	287	N

Non-coding RNA prediction

Each dataset was run through the R-package, *baerhunter* (Ozuna et al., 2019), using the '*feature_file_editor*' function optimised to the most appropriate parameters for the sequencing depth (https://github.com/jenjane118/mtb_modules). '*Count_features*' and '*tpm_norm_flagging*' functions were used for transcript quantification and to identify low

expression hits (less than or equal to 10 transcripts per million) in each dataset, which

were subsequently eliminated. When viewed on a genome browser, coverage at the 3' ends

of putative sRNA and UTRs often appears to decrease gradually, with the actual end of

the transcript appearing indistinct, compared to the 5' end. Prokaryotic ncRNA

transcripts may not demonstrate a clear fall-off of expression signal in RNA-seq, as

pervasive transcription is regulated by the changing levels of Rho protein observed in

different conditions (Bidnenko & Bidnenko, 2018; Wade & Grainger, 2014). These very

long predictions can mask predicted transcripts in the same region from other samples,

obscuring potentially interesting shorter transcripts expressed in different conditions. For

this reason, transcripts longer than 1000 nucleotides were eliminated before combining

the predictions between datasets. The predicted annotations for each dataset were

combined into a single annotation file, adding the union of the predicted boundaries to

the reference genome for H37Rv (AL123456.3). Predictions that overlapped with

annotated ncRNAs and UTR predictions that overlapped sRNA predictions from a

different dataset were eliminated. Transcript quantification was repeated on each dataset

using the resulting combined annotation file and the count data from each dataset was

merged into a single counts matrix.

DESeq2 v1.30.1 (Love et al., 2014) was used on the complete counts matrix including the

filtered *baerhunter* predictions to calculate size factors, estimate dispersion and

normalise the data with the regularised log transformation function (Supp figures, S1 and

S2). The normalised data was checked for potential batch effects using PCA plots and

hierarchical dendrograms. *Limma* v3.46.0 (Ritchie et al., 2015) *'removeBatchEffect'* was

applied with a single batch argument to remove batch effects associated with the first

component (batching the data according to dataset due to technical differences) while

preserving differences between samples. The final hierarchical dendrogram, post-batch

correction, indicates successful application as samples cluster by similar experimental

conditions, rather than by dataset alone (Figure 2 compared to Supp figure S3). Samples

from experiment PRJEB65014 continue to group together, but as they represent single

replicates in unique conditions, it is difficult to estimate the influence of confounding

batch effects for these samples.

Figure 2. Hierarchical dendrogram of *rlog* transformed and *limma* batch corrected
expression data by sample. The sample labels are coloured by dataset, demonstrating that
they are clustering by condition, rather than experiment.
252

Creation of the WGCNA network

The normalised and batch-corrected expression matrix was used to create a signed co-

expression network using the R package, *WGCNA* v1.69 (Langfelder & Horvath, 2008),

with the following parameters: corType = "bicor", networkType = "signed", power = 14,

TOMType = "signed", minModuleSize = 20, reassignThreshold = 0, mergeCutHeight =

0.15, deepSplit = 2. In this type of network, the 'nodes' are the genes, and the 'edges', or

links, are created when gene expression patterns correlate. In contrast to unweighted

binary networks where links are assigned 0 or 1 to indicate whether or not the genes are

linked, in a weighted network, the links are given a numeric weight based on how closely

correlated the expression is. *WGCNA* first calculates the signed co-expression similarity

for each gene pair. The absolute value of this correlation is raised to a power (determined

by the user, based on a scale-free topology model that mimics biological systems (Supp

figure S4) in order to weight the strong connections more highly than the weaker

connections. The resulting similarity matrix is used to cluster groups of genes with strong

connections to each other in a non-supervised manner (i.e., it doesn't use any previous

information about gene groups or connected regulons). A cluster dendrogram is created

(Supp figure, S8) and closely connected branches of the dendrogram are merged into

modules based on a cut-off value (also a parameter controlled by the user). The modules

are defined by a 'module eigengene' (ME), which explains most of the variance in the

expression values in the module. The connectivity of the MEs define the shape of the

overall network (Supp figure, S9). The modules can then be tested for potential

correlations with experimental conditions without incurring the same punitive penalties

for multiple testing as individual gene correlations would (Supp figure, S10). In signed

networks, correlation of the module with a condition can be in either the positive or

negative direction, as modules include transcripts that are similar in both the degree and

direction of correlation, allowing for a more fine-grained analysis than with unsigned

networks.

To test correlations of modules with experimental conditions, the individual RNA-seq

samples were assigned to a condition based on the experimental description in the project

metadata. Some of these conditions were shared among the different projects, so when

appropriate, samples from different datasets were assigned the same condition, resulting in 15 tested conditions. For example, late-stage reaeration samples were tested along with exponential growth samples, and samples that tested hypoxia and cholesterol utilisation together were included in multiple conditions. Models of hypoxia differed between the RNA-seq projects, and these samples were assigned to different conditions: 'hypoxia' versus 'extended hypoxia' (Supp Table 1, 'Condition summary' tab). All correlations were made using robust biweight midcorrelation tests and all p-values were corrected for multiple testing with the BH-fdr method (Benjamini & Hochberg, 1995). Significance was evaluated as an adjusted p-value (p_{adj}) of < 0.05.

Module Enrichment

Modules were interrogated for enrichment for Gene Onotology (GO) terms (Ashburner et al., 2000; The Gene Ontology Consortium, 2021), Clusters of Orthologous Groups (COG) (Galperin et al., 2021), KEGG pathway genes (Kanehisa et al., 2022), functional categories and literature searches for known regulons. GO terms, COG term and KEGG pathway enrichment were accessed programmatically using the David web service (Huang et al., 2009b, 2009a; Jiao et al., 2012) to query the list of protein-coding genes from each module for enrichment. Enrichment was determined using a modified one-sided Fisher's Exact Test ('EASE' score) with fdr correction for multiple testing, with $p_{adj} < 0.01$ considered significantly enriched for a particular term or pathway, and $p_{adj} < 0.05$ for COG term. Enrichment for the 11 functional categories from Mycobrowser annotation (Kapopoulou et al., 2011) was determined using a one-sided Fisher's Exact Test with fdr correction for multiple testing. Modules were enriched for a particular functional category if $p_{adj} < 0.01$. Lists of genes associated with known regulons were mined from literature and enrichment was tested using the same one-sided Fisher's Exact Test as above with a $p_{adj} < 0.01$ cut-off for enrichment.

313 Non-coding RNA prediction, network analysis and subsequent data manipulation was

314 performed with R (v4.0.5, 2021-03-31). All plots were made in R with the following

315 packages: *WGCNA* (v1.69), *dendextend* (v1.15.2), *ggplot2* (v3.3.5). Scripts and expression

316 data are available at https://github.com/jenjane118/mtb_modules .

RESULTS AND DISCUSSION

Mtb expresses an extensive range of ncRNA transcripts over a wide variety of experimental conditions

Mycobacterium tuberculosis RNA-seq datasets were selected from publicly available data to find experiments using the wild-type H37Rv strain and representing a range of growth conditions the pathogen may encounter in a host environment. Four datasets passing our quality standards were subjected to our analysis pipeline (see Material and Methods) and included 52 samples under 15 different experimental conditions (Supp Table 1, 'Samples' tab). The R package, *baerhunter* (Ozuna et al., 2019), was used to predict ncRNA in intergenic regions, antisense RNA (opposite a protein-coding gene) and UTRs at both the 5' and 3' ends of genes by searching the mapped RNA-seq data for expression peaks outside of the annotated regions in the reference sequence for H37Rv. Non-coding RNA predictions from each dataset were filtered for low expression and combined to create a single set of non-overlapping annotations that encompassed all predictions made from any sample under any experimental condition. In total, 1283 putative sRNAs were predicted (including both truly intergenic transcripts as well as those antisense to a protein-coding gene, or annotated RNA) and 1715 UTRs which includes all transcribed regions outside of annotated protein-coding sequences at both 5' and 3' ends, as well as the non-coding regions between adjacent genes in operons. All putative ncRNA transcripts (sRNAs and UTRs) were searched for a TSS near the start of the predicted 5' boundary using previously published annotations (Cortes et al., 2013; Shell et al., 2015). Annotated TSSs were found within 20 nucleotides of the 5' end in 43% of the predicted sRNA transcripts. Predicted 5' UTRs had a TSS within 10 nucleotides of the start in 42% of cases, compared with 3% of the predicted 3' UTRs. Where the UTR covered the entire sequence between two protein-coding regions (labelled as 'between' UTRs), 9% had a TSS in the first 10

nucleotides of the sequence (Table 2 and Supp Table 2 'putative_sRNAs', 'putative_UTRs' tabs).

Table 2. Tally of predicted expressed elements in the *baerhunter*-generated combined annotation file. 4015 protein-coding genes were included in the annotation. *TSS predictions from (Cortes et al., 2013; Shell et al., 2015).

Predicted element	Number predicted	With predicted TSS* (exponential and starvation)
Total sRNA	1283	553
sRNA 'intergenic'	91	23
sRNA 'antisense'	1192	530
Total UTRs	1715	273
5' UTRs	475	200
3' UTRs	602	16
'Between' UTRs	638	57

The predicted sRNAs were further annotated using the accepted nomenclature (Lamichhane et al., 2013) which identifies the putative ncRNA relative to annotated gene loci and differently signifies truly intergenic sRNAs and those that overlap any part of a protein-coding region on the opposite strand. Most of the putative sRNAs are antisense to the protein-coding region of one or more genes, but 91 putative sRNAs have predicted boundaries that do not overlap an annotated transcript on either strand (or overlap an annotated transcript on the opposite strand by fewer than 10 nucleotides). This number is most probably an underestimate of the truly 'intergenic' sRNAs in the genome, as many of the sRNA predictions appear over-estimated at the 3' end, effectively classifying them as an antisense RNA even though the 5' half of the transcript does not overlap any genes on the opposite strand. Isoforms of annotated sRNAs can be subject to post-transcriptional processing to create an active transcript (Moores et al., 2017) and post-transcriptional

processing of 3' ends *in vivo* is more likely the norm for most prokaryotic transcripts (Wang et al., 2019). However, for our purposes, any RNA-seq transcripts that extend to overlap a protein-coding gene on the other strand in any dataset will be labelled as antisense RNA.

The generated combined annotation file was used to quantify the expression of all 7043 expressed elements, including every annotated CDS, annotated ncRNA and predicted ncRNA, in each sample. Raw counts of expression varied greatly among the datasets due to different sequencing depth, as well as between some samples within datasets (as would be expected with different environmental conditions), and only three protein-coding genes showed no expression in any sample. The raw expression counts were transformed using DESeq2's rlog function (Love et al., 2014), and plots of the dispersion of count data show that the median expression level between samples and between datasets has been normalised (Supp figures S1, S2). The distribution of the normalised expression levels of protein-coding regions alone shows consistent median expression levels across the entire dataset, however distribution of the normalised data restricted to putative sRNAs shows more variability, with certain conditions showing increased or decreased expression of these transcripts (Supp figures S5-S7). This is not unexpected, given that several studies have identified pervasive transcription in hypoxic infection models, stationary phase and dormancy. This is accompanied by a concomitant increase in non-rRNA expression (especially antisense RNA transcripts) and in the number of predicted TSSs in Mtb and *M. smegmatis* (a fast-growing, non-pathogenic strain) (Arnvig et al., 2011; Ignatov et al., 2015; Martini et al., 2019).

Module networks represent groups of co-expressed genes and predicted non-coding RNA

Creation of the WGCNA network

A weighted co-expression network was created from the normalised RNA-seq expression data using *WGCNA* (Langfelder & Horvath, 2008) (see Materials and Methods). This program segregates genetic elements with similar patterns of expression over a range of samples into modules. The modules represent sub-networks of connected genes, and functional relationships can be explored among the members of the individual modules. The 'hub' genes represent the most highly connected gene elements within a module and have highest module membership values. Module membership is measured by correlation of the expression of the individual genes with the module eigengene (ME), the vector that best represents the variation in the module.

The signed co-expression network presented in this paper consists of 56 different modules, assigning 97.6% of the expressed elements (CDS, putative UTRs and putative sRNAs) into 55 modules, with 168 unassigned elements clustered in the 'grey' module (Supp Table 2, 'Module_Overview' tab). Module size ranged from 1086 to 25 expressed elements. The modules (using the ME) were tested for correlations with the various conditions used in the RNA-seq experiments (see Materials and Methods). The RNA-seq data was categorised into 15 different experimental conditions in total with varying numbers of replicates (Supp Table S1, 'Condition Summary' tab), therefore, a statistically significant correlation of modules with every condition was not expected. However, some modules do show significant correlations with conditions such as iron restriction, cholesterol media, hypoxia and growth phase and this can be informative when considering the association of the gene groups with biological processes (Figure 3).

Well-established regulons cluster together in single modules

In many cases, the gene membership of the modules includes well-established regulons

or groups of functionally related genes, establishing the biological relevance of the module

sub-networks and proof of concept for the application of WGCNA on such a heterogenous

dataset. For example, the DosR regulon is a well-studied regulon associated with hypoxia

and stress responses (Du et al., 2016; Rustad et al., 2008; Voskuil et al., 2004). 40 of the

48 DosR-regulated genes are found in a single module, '*greenyellow*', which is negatively

correlated with reaerated culture and exponential growth (Figure 3) and enriched for the

GO term, 'response to hypoxia'. Unsurprisingly, this represents statistically significant

enrichment of DosR-regulated genes in the module (one-sided Fisher's exact test, p_{adj}=

6.6e-50). The '*greenyellow*' module is also enriched for genes from the PhoP regulon (one-

sided Fisher's exact test, p_{adj}=0.021) which is associated with hypoxic response and

coordination with the DosR regulon (Gonzalo-Asensio et al., 2008; Singh et al., 2020). The

KstR regulon includes 74 genes under control of the TetR-type transcriptional repressor,

KstR, known to be involved in lipid catabolism and upregulated during infection (Kendall

et al., 2007, 2010). The '*turquoise*' module is significantly enriched for known KstR-

regulated genes (one-sided Fisher's exact test, p_{adj} = 0.0026) with 35 of 74 KstR-regulated

genes clustering together in the module. This module showed significant positive

correlation with hypoxia, extended hypoxia and stationary growth phase, and a negative

correlation with exponential growth (Figure 3).

Other examples include genes involved in mycobactin synthesis which are nearly all found

in the '*steelblue*' module (positively correlated with the low iron condition), and the genes

of the DIM locus which are significantly enriched (one-sided Fisher's Exact test,

p_{adj}=4.95e-5) in the '*paleturquoise*' module (positively correlated for exponential growth

and reaerated culture and negatively correlated to slow growth conditions) (Figure 3). As

these examples show, known associated genes are co-located in modules which represent

a functional group of genes that have co-regulated expression under various experimental

conditions. The modules can be further explored to identify novel associations.

Figure 3. Heat map of correlation of module eigengene (ME) of each module with selected
experimental conditions. Correlation was calculated using biweight midcorrelation (bicor)
and p-values were adjusted for multiple testing (BH-fdr). Positive correlation is red,
negative correlation is blue. Non-significant correlations in grey ($p_{adj} < 0.05$).

Predicted non-coding RNAs are enriched in certain modules

Putative sRNAs and predicted UTRs were distributed throughout all modules in the

network (Figure 4, Supp Table 2, 'Module_Overview' tab). The number of predicted

elements were enriched in certain modules: the two largest modules, *'turquoise'* and *'blue'*,

are significantly enriched for predicted sRNAs, and eight modules are statistically

enriched for predicted UTRs (one-sided Fisher's exact test, p_{adj} < 0.01, Supp Table 2,

'Module_Overview' tab). A roughly linear relationship between the number of CDS and

the number of UTRs, is to be expected, given that UTRs are defined by the *baerhunter*

algorithm by their position at the start or end of protein-coding genes (Ozuna et al., 2019).

However, if the UTRs are positioned in an operon, there will be a smaller increase in the

relative number of UTRs with an increasing number of protein-coding genes, as UTRs

between two protein-coding genes are predicted as a single UTR. As a result, the two

modules with the highest number of predicted operons (from OperonDB, Chetal & Janga,

2015), *'turquoise'* and '*brown*', have a lower relative proportion of UTRs (Figure 5).

Within the module sub-networks, the tight co-expression of protein-coding genes and

ncRNA is reflected by the number of ncRNA found among the most connected elements

in the module. The 'hub' elements are those with the best correlation to the ME and

therefore the most tightly connected elements in the individual module networks. In 12

modules, ncRNA (both predicted and annotated) make up more than half of the elements

with module membership values (MM) > = 0.80 (our threshold for identifying hub

elements) (Supp Table 2, 'Hub_info' tab). This implicates ncRNA as important members

of the regulatory pathways implemented to adapt to conditions such as hypoxia,

cholesterol media and low iron. The 30 annotated ncRNAs in the Mtb reference genome

(AL123456.3) are spread over 15 modules, with 10 of them hubs of the modules, and one

unassigned ('grey' module) (Supp Table 2, 'Annotated ncRNA' tab). For example,

Ms1/MTS2823, observed to be the most abundantly expressed ncRNA in expression

studies over various stress conditions (Arnvig et al., 2011; Arnvig & Young, 2012; Ignatov

et al., 2015; Šiková et al., 2019), is a hub element in a module that is positively correlated

with hypoxia and negatively correlated with exponential and reaerated culture conditions

(*'lightsteelblue1',* Figure 3). Mcr7/ncRv2395A, found to be part of the PhoP regulon

(Solans et al., 2014), is a hub in the *'magenta'* module enriched for elements in the KEGG

pathway for valine, leucine and isoleucine degradation and correlated positively with the

low iron condition (Figure 3). F6/ncRv10243/SfdS, a sRNA upregulated in starvation and

mouse infection models, is thought to be involved in regulating lipid metabolism and long-

term persistence (Houghton et al., 2021). This ncRNA is a hub in a module found to be

enriched for the GO terms 'lipid metabolism' and 'biosynthesis of fatty acids' (*'lightcyan'*)

and found to be correlated positively with low iron and negatively with extended hypoxia

conditions (Figure 3).

Figure 4. Relative proportion of annotated CDS, predicted UTRs and predicted sRNAs in
each module, ordered by module size.
496
497

499
500
501
502 Figure 5. Plot of number of UTRs against number of CDS in each module. Grey shading
503 indicates confidence interval of 0.95.
504

505
506

UTR and adjacent ORF expression differ in nearly 50% of cases

We were interested to see how many of the predicted UTRs were assigned the same

module as the adjacent ORF—indicating whether the ORF and its adjacent UTR were co-

regulated. Intuitively, the UTR of a protein-coding gene would be expected to be expressed

as a single transcript along with the ORF and show similar expression patterns. However,

both 5' and 3' UTRs can act independently of the attached ORF and RNA abundance in

RNA-seq experiments reflects both transcription activity and transcript stability. For

example, some 5' UTRs are known to contain regulatory elements, such as riboswitches,

that alter the transcription of the downstream ORF (Dar et al., 2016; Kipkorir et al., 2021;

Schwenk & Arnvig, 2018; Warner et al., 2007), whereas sRNAs cleaved from 3' UTRs have

been shown to regulate the stability of the remaining transcript--with different half-lives

as a result (Chao et al., 2012; Dar & Sorek, 2018; Menendez-Gil & Toledo-Arana, 2021).

Of the *baerhunter*-predicted UTRs labelled 5' and 3', the UTRs co-segregated with the

ORF they were closest to approximately half the time (Table 3). We would expect

correctly-identified 5' UTRs to utilise a TSS (whether or not there is a known predicted

TSS), whereas it appears functional 3' UTRs are more likely to be cleaved from the longer

mRNA transcript (Dar & Sorek, 2018; Menendez-Gil & Toledo-Arana, 2021; Ponath et al.,

2022). Our data confirms this: transcripts classified as 5' UTRs are much more likely to

have a predicted TSS in the first 10 nucleotides than transcripts classified as 3' UTRs

(42% vs 2.7%). Approximately 11% of the UTRs predicted to be between ORFs (labelled,

'Between' UTRs) have predicted TSS (Table 3). The presence of a TSS in the first 10

nucleotides of the predicted UTR appeared to have little bearing on whether or not the

UTR and its adjacent ORF are assigned to the same module, with 56% of 5' and 44% of 3'

UTRs with a predicted TSS co-assigned with their adjacent ORF partner. A similar

proportion of the 'Between' UTRs (38%) do not segregate with either the ORF upstream

or downstream, indicating their expression is, to some degree, independent of either

adjacent ORF. All UTRs that are in modules independent of their adjacent ORF(s) are

found in Supplementary Table 2, 'independent_UTRs' tab.

Table 3. UTRs and module assignment of adjacent ORFs. DS=downstream,
US=upstream. TSS indicates presence of annotated TSS in first 10 nucleotides of
predicted UTR (Cortes et al., 2013; Shell et al., 2015).
540

	Total (excluding grey)	Number with TSS	Number in same module as adjacent ORF	Proportion of UTRs in same module as ORF
5' UTR	462	196	227 DS	48%
3' UTR	592	16	296 US	49%
BTWN UTR	622	55	117 DS	19%
			140 US	23%
			126 both	20%
			239 none	38%

541
542

Antisense RNAs are hubs in modules independent of cognate ORF

It has been observed that the overall abundance of antisense RNA and other non-

ribosomal RNA increases upon exposure to stress such as hypoxia and nutrient restriction

(Arnvig et al., 2011; Ignatov et al., 2015), and in our network, ncRNA, including antisense

RNAs, were found to be well-connected hubs in module sub-networks associated with

known gene regulons, such as DosR and KstR. This supports the view that antisense RNA

may be part of specific regulatory networks, especially those that are involved in

adaptation to environmental conditions, rather than products of indiscriminate pervasive

transcription (Arnvig et al., 2011; Lloréns-Rico et al., 2016). Not unexpectedly, very few

(7%) of the predicted antisense transcripts were assigned to the same module as the

protein-coding region overlapping on the opposite strand (choosing the most downstream

locus in the event of multiple overlapping ORFs), signifying distinct patterns of expression

for transcripts on opposite strands, possibly due to independent or bi-directional

promoters and/or overlapping transcription termination sites. Bi-directional promoters

have been identified in multiple prokaryotic genomes, and competition for RNA

polymerase (RNAP) binding among divergently transcribed sense/antisense pairs may

function as a mechanism for regulation of gene expression (Ju et al., 2019; Warman et al.,

2021). Long 3' UTRs that overlap with converging protein-coding genes on the opposite

strand (or with the 3' UTR) can create an 'excludon' regulatory arrangement, where

transcription of the two opposite mRNAs is simultaneously regulated by RNase targeting,

or mutually exclusive due to RNAP collision (Sáenz-Lahoya S. et al., 2019; Toledo-Arana

& Lasa, 2020). Examining the module groupings of the antisense RNAs and their base-

pairing target on the other strand may provide insight on which genes are regulated by

antisense transcription.

Focus on Selected Module Networks

The large-scale transcription analysis presented here is useful for the more global

analysis of the overall trends related to ncRNA and transcription, but there is a great deal

of information to be gleaned by more fine-grained inspection of individual module

groupings. To discover novel associations in such a large and complex dataset, we have

selected a few modules for closer examination, focussing on those that contain gene groups

or regulons related to the tested conditions. Many of the modules that contain interesting

correlations or gene regulon enrichments also include an abundance of putative sRNAs

and UTRs. Using the 'guilt by association' principle, we can hypothesise that the well-

connected ncRNAs found among the module hub elements have a role in transcriptional

'remodelling' in response to changes in environmental conditions such as growth on

cholesterol-containing media, restricted iron or hypoxia.

One condition that causes a major shift in the transcriptome is the adaptation of Mtb to

a cholesterol or lipid-rich environment, a process that involves a multitude of gene

pathways to facilitate the pathogen's survival and persistence in the infected macrophage

(Del Portillo et al., 2019; Pandey Amit K. & Sassetti Christopher M., 2008; Pawełczyk et

al., 2021). In fact, a recent study, published after this analysis, observes differential

expression of over 500 protein-coding genes with a switch from glycerol to cholesterol as

the carbon source (Pawełczyk et al., 2021). Our network includes transcriptomes from

several samples that utilise cholesterol and fatty-acid containing media over a range of

growth conditions including hypoxia (SRA project: PRJNA390669) (Aguilar-Ayala et al.,

2017) and although several modules have a significant correlation with the cholesterol

media trait, other modules with clusters of genes related to cholesterol catabolism are

correlated to hypoxia or extended hypoxia conditions. All of these modules are found to

contain a large number of predicted non-coding elements, confirming studies that show

increased ncRNA expression levels in response to lipid conditions and cholesterol,

especially when combined with hypoxia; conditions meant to most resemble those

encountered in host infection models (Aguilar-Ayala et al., 2017; Del Portillo et al., 2019;

Soto-Ramirez et al., 2017).

Several modules correlating with the low iron condition show enrichment of genes

associated with siderophore synthesis, transport and regulation, along with redox sensors

and genes known to be upregulated in response to cholesterol media. Restricting iron

availability to growing cells is meant to replicate a host response to infection and will

stimulate a cascade of pathways to enable the pathogen to survive in a slow-growing, or

latent state. The co-expression of genes involved in metal ion homeostasis and genes

known to be involved in adaptation to cholesterol and lipids is supported by observations

in a recent study that the presence of cholesterol causes changes in metal ion metabolism

(Pawełczyk et al., 2021) and closer inspection may uncover gene interactions related to

the metabolic changes made in anticipation of re-entry from hypoxic environments when

bacteria are particularly vulnerable to oxidative stress (Eoh et al., 2017; Gerrick et al.,

2018).

The data have been organised into an easily-accessible spreadsheet for researchers to

query particular genes or modules of interest and find associated protein-coding genes or

ncRNA (Supp Table 2). We anticipate this to be a useful resource to find ncRNA

candidates for further study, to identify associations of genes with unknown functions,

and to suggest roles for 'moonlighting' proteins that may be associated with unexpected

gene groupings.

The largest module includes the kstR regulon and is enriched for predicted sRNAs

The *'turquoise' module* contains more than 1,000 expressed elements, with over 50% of the hubs being predicted sRNAs. It contains 461 protein-coding genes, including 34 of the 71 KstR regulon genes and 52 transcription factors (Rustad et al., 2014). 26 of the 32 *kstR* regulon genes found to be differentially expressed in Mtb grown with cholesterol versus glycerol as the main carbon source (Pawełczyk et al., 2021) are found in the '*turquoise*' module, with 15 of them hubs. The hubs also include 10 transcription factors and DNA binding proteins, including IdeR, FurA, KstR, KstR2 and SigB, anti-sigma factor ResA, two annotated sRNAs (mcr11/ncRv11264c, and mpr6/ncRv1222) and many predicted ncRNA elements including 131 predicted sRNAs and 26 UTRs (Supp Table 2, 'CDS hubs, 'srna_hubs' tabs). The module has 46 complete predicted operons from OperomeDB (Chetal & Janga, 2015), and the highest number of consecutive ORFs in the genome of all the modules.

The size of the *'turquoise'* module, and the fact that it has resisted splintering into smaller modules during the tree-cutting process, indicates that it includes many highly connected gene operons involved in multiple interconnected stress response pathways. The module shows enrichment for the GO terms 'regulation of transcription' and 'cholesterol catabolic process', as well as for the KEGG pathway for steroid degradation (Supp Table 2, 'Module Overview' tab). Despite the inclusion of genes linked specifically to cholesterol metabolism, a significant correlation of the *'turquoise'* module with the cholesterol-containing media condition was not established; rather, the module shows positive correlations with hypoxia (bicor = 0.41, p_{adj} = 0.001), extended hypoxia (bicor = 0.035, p_{adj} = 0.002) and stationary (bicor = 0.34, p_{adj} = 0.03) conditions, and a negative correlation with exponential growth (bicor = -0.42, p_{adj} = 0.009) (Figure 3). Transcriptomic changes in response to lipid degradation include many genes related to redox maintenance which are

found in the module, including redox-sensing *whiB3* and *whiB4* (Larsson et al., 2012;

Mehta & Singh, 2019).

Among the module hubs, are annotated sRNAs such as mcr11/ncRv11264c, which has

been associated with dormancy and hypoxic conditions and shown to regulate the

expression of genes related to the metabolic remodelling associated with persistence and

slow growth states in Mtb (Girardin & McDonough, 2020). Other annotated ncRNA in

'turquoise' include: mpr6 (ncRv1222), G2 (ncRv11689c), mcr16 (ncRv2243c), C8/4.5S RNA

(ncRv13722Ac), and another experimentally-verified ncRNA, mrsI (ncRv11846) that was

predicted as a somewhat longer transcript in this study (and in a previous study, (Arnvig

et al., 2011)) which extends antisense to the gene Rv1847

(putative_sRNA:m2096739_2097122 / ncRv1847c). MrsI has been found to be upregulated

in several growth states and stress conditions and is implicated in anticipatory regulation

of iron acquisition (Gerrick et al., 2018). Most of the predicted sRNAs in the *'turquoise'*

hubs are classified as antisense transcripts, with 82 having predicted TSSs within 10 nt

of the start. In addition, 7 strictly 'intergenic' predicted sRNAs are among the hubs. Four

of these have predicted TSS within 20 nucleotides of the start. (Supp Table 2,

'intergenic_putative_sRNAs' tab).

Detoxification-linked proteins cluster in the module best correlated with cholesterol

media condition

The *'black'* module showed positive correlation with the cholesterol media condition

(bicor=0.54, p_{adj}=0.002) and negative correlation with low iron (bicor = -0.48, p_{adj} = 0.001)

(Figure 3). Many protein-coding genes involved in detoxification pathways are hubs in the

module, including several encoding transmembrane proteins such as the *mmpL5-mmpS5*

efflux pump operon (Rv0676c-Rv0677c), as well as the next gene downstream, Rv0678,

which was identified as part of a 'core lipid response' in differential expression analysis in lipid-rich media (Aguilar-Ayala, et al., 2017). The 5' UTR for Rv0677c and 3' UTRs for Rv0676c and Rv0677c are also hubs. This operon is involved in siderophore transport and expressed in cholesterol and lipid-rich environments (Aguilar-Ayala, et al., 2017; Pawełczyk et al., 2021). Other detoxification-linked genes in the module, such as the ABC-family transporter efflux system, Rv1216c-1219c and the operon including PPE53 (Rv3159c), Rv3160c and Rv3161c, have also been implicated in transcriptomic remodelling in response to cholesterol (Aguilar-Ayala et al., 2017; Pawełczyk et al., 2021).

Among the hubs are three predicted antisense RNAs. One antisense RNA, ncRv1358c (putative_sRNA:m1530046_1530745) has a TSS near its start and is found antisense to Rv1359. Rv1359 and the upstream gene, Rv1358, on the opposite strand are very similar to each other (43.7% identity in 197 aa overlap) and to another gene elsewhere in the genome, Rv0891c (48.5% identity in 204 aa overlap) (Kapopoulou et al., 2011). All three genes are possible LuxR family transcriptional regulators which are thought to be involved in quorum-sensing adaptations and contain a probable ATP/GTP binding site motif (Chen & Xie, 2011; Modlin et al., 2021). Expression of this antisense sRNA appears to suppress the expression of the transcript on the opposite strand to varying degrees in all conditions (Figure 6). Expression of a shorter transcript appears to begin inside the Rv1359 ORF, where the transcript is not overlapped by the antisense transcript, possibly utilising an internal TSS at 1530774.

Figure 6. Expression of antisense transcript putative_sRNA:m1530046_1530745 (magenta bar) seems to suppress the expression of most of Rv1359 and Rv1358. An internal TSS exists inside the Rv1359 CDS at 1530774 near where expression begins. Sample SRR5689230 from PRJNA27860, exponential growth on cholesterol and fatty acid media. Strand coverage using the 'second' read of each pair mapping to the transcript strand, visualised using Artemis genome browser (Carver et al., 2012).

Two adjacent predictions, the 3' UTR for Rv1772 (putative_UTR:p2006948_2007063) followed by ncRv1773/ putative_sRNA:p2007213_2007377, are hubs in the *'black'* module. Together, they extend to overlap the antisense strand of a large portion of Rv1773c, a probable transcriptional regulator in the IclR-family, found in a different module *('navajowhite2')*. The 3' UTR for Rv1772 was previously identified as an abundant antisense transcript during exponential growth (Arnvig et al., 2011). The start of the predicted sRNA transcript has no known TSS and could instead be an extension of the predicted 3' UTR (Supp figure S11). (When combining predicted annotations from different datasets, long predicted UTRs that overlapped shorter sRNA predictions were discarded, see Methods). In *E.coli*, the IclR-family transcriptional regulators demonstrate both activating and repressing activities on targets such as multidrug efflux pumps and the *aceBAK* operon which regulates the glyoxylate shunt (Zhou et al., 2012). *Icl2a* (Rv1915) is one of the Mtb isoforms of the isocitrate/methylocitrate lyase gene, *aceA*, and may be regulated by Rv1773c, as seen in *E.coli*. Icl2a, Rv1772, its predicted UTR and the antisense RNA (ncRv1773) are all hubs in the *'black'* module. *Icl2a* has been observed to be upregulated with cholesterol as the sole carbon source and likely has a second function as part of the methylcitrate cycle to convert the fatty acid metabolites propionate and

propionyl CoA to less toxic compounds (Bhusal et al., 2017; Pawełczyk et al., 2021).

Another predicted antisense RNA in the *'black'* hubs, ncRv0027c/

putative_sRNA:m31259_31967, has a TSS near its start (31967) and is antisense both to

Rv0027 and Rv0028, conserved hypothetical proteins with no known function found in

different modules.

The module including transcriptional regulator whiB1 and genes of kstR2 regulon, links

metal ion balance with cholesterol utilisation

The *'lightcyan'* module is significantly enriched for genes under control of another TetR-

type repressor, KstR2, (one-sided Fisher's exact test, p_{adj} = 4.27e-06) with 7 of the 15

known regulon genes found in the module. KstR2-regulated genes are known to be

involved in cholesterol utilisation (Kendall et al., 2010) and the protein-coding genes of

this module were enriched for the COG term, 'lipid metabolism', and KEGG pathways,

'Biosynthesis of unsaturated fatty acids' and 'Fatty acid metabolism'. However this

module did not significantly correlate with the cholesterol media condition. Instead, the

ME was positively-correlated with the low iron condition (bicor = 0.59, p_{adj} = 9e-5) and

negatively-correlated with the extended hypoxia condition (bicor = 0.41, p_{adj} = 0.008)

(Figure 3). The correlation of this ME with the low iron condition implies that there are

expressed elements within the module that are involved in iron homeostasis, possibly in

tandem with adaptation to cholesterol. Intriguingly, one of the hub genes of this module

encodes the redox-sensing transcriptional regulator, WhiB1. This transcription factor is

known to be stimulated by a variety of stress conditions and *in vivo,* and binds an iron-

sulfur cluster (Larsson et al., 2012; L. J. Smith et al., 2010).

The hubs of the *'lightcyan'* module include several predicted sRNAs, and the annotated

sRNA, F6. F6/ncRv10243/SfdS is a sigF-dependent ncRNA which has been shown to be

induced in nutrient starvation, oxidative stress, acid stress (Arnvig & Young, 2009;

Houghton et al., 2021) and the fatty acid hypoxia model (Del Portillo et al., 2019). In

addition to being expressed from its own promoter, F6/SfdS has been proposed to be co-

transcribed with the upstream gene *fadA2* (Rv0243), a probable acetyl-CoA

acyltransferase; however, *fadA2* is clustered in a different module from SfdS, one

associated with iron acquisition ('*violet*', see below). One of the predicted sRNAs in the

module hubs is antisense transcript ncRv2489/putative_srna:p2801108_2801678 with a

TSS at 2801108. This overlaps the 3' end of PE-PGRS43 (Rv2490c) (Figure 7). There is a

short reading frame (30 nucleotides, 10 amino acids) initiating from a Methionine at this

TSS that suggests a possible dual-function sRNA or sORF with independent function. The

TSS for the predicted sRNA overlaps the 5' end of Rv2489c, a short, hypothetical 'alanine-

rich protein'. The TSSs for these convergently overlapping transcripts are 42 nts apart

(Rv2489c appears to be a leaderless transcript based on dRNA-seq and position of TSS)

and may involve RNAP collision if both are transcribed simultaneously. Therefore,

transcription of the predicted sRNA could impact either Rv2489c and/or PE-PGRS43

expression through two different mechanisms. Other hub sRNAs in '*lightcyan*' include

ncRv1450/putative_sRNA:p1630466_1631246, which has a TSS at 1630466 and is

antisense to the 3' end of PE-PGRS27 (Rv1450c) and putative_sRNA:p3936733_3936893

/ ncRv3509 which includes a predicted TSS at 3936720 which overlaps the 3' end (and

predicted 3' UTR) of Rv3509c (*ilvX*), a probable acetolactate synthase (found in the '*violet*'

module).

Figure 7. Antisense sRNA, ncRv2489/putative_srna:p2801108_2801678, (magenta bar) overlaps two
transcripts and may encode a short peptide. TSS for sRNA indicated in red and corresponding amino acid
highlighted in pink. Sample SRR5689230 from PRJNA390669, exponential growth on cholesterol and fatty
acid media. Strand coverage using the 'second' read of each pair mapping to the transcript strand, visualised
using Artemis genome browser (Carver et al., 2012).

The module best correlated with the low iron condition includes genes related to metal

ion and fatty acid homeostasis

Another module that is positively correlated to the low iron condition is the *'violet'* module

(bicor=0.61, p_{adj}=6e-05, Figure 3). This module contains most of the ESX-3 genes (Rv0282-

Rv0292) related to siderophore-mediated iron (and zinc) uptake in Mtb (Serafini et al.,

2013; L. Zhang et al., 2020), with two of these representing hubs in the module. The gene

preceding the ESX-3 genes, Rv0281, a possible S-adenosylmethionine-dependent

methyltransferase involved in lipid metabolism (though its position in the genome would

suggest regulation could be linked to ESX-3 (Lunge et al., 2020)), is in the module, as well

as an ESX-5 gene, Rv1797 (*eccE5*). The module also contains another Zur-regulated gene,

Rv0106, which is a potential zinc-ion transporter (Zondervan et al., 2018). Among the

hubs of the module are several genes related to lipid metabolism and fatty acid synthesis,

including: probable triglyceride transporter, Rv1410; the operon consisting of Rv0241c

(*htdX*), Rv0242c (*fabG4*), and Rv0243 (*fadA2*) (Dutta, 2018); and a gene involved in the

pentose phosphate pathway, *zwf2* (Rv1447c).

There are some well-connected ncRNAs in the *'violet'* module, including a predicted

antisense RNA to Rv0281, 'ncRv0281c'. This putative sRNA has a predicted TSS at the 5'

end and is transcribed divergently from Rv0282 *(eccA3)*. This is one of the rarer cases

where the antisense transcript and cognate protein-coding gene (Rv0281) are clustered in

the same module. The prevailing direction of transcription at this locus may be a result of

competition for RNAP binding at a bi-directional promoter in the predicted 5' UTR of

Rv0282 which also clusters in the module. Another predicted sRNA in the module,

ncRv3508/putative_sRNA:m3932046_3932369 has a predicted TSS at 3932369 and

transcribed opposite to a central region of Rv3508c, PE_PGRS54, a gene in the

'darkolivegreen' module which is enriched for PE/PPE genes (p_{adj} = 4.12e-09).

There are several UTRs in the module hubs, including a 3' UTR for the gene Rv1133c,

metE; the gene is found in another module, *'grey60'*. This might be an example of a sRNA

differentially transcribed or cleaved from the 3' UTR of a protein-coding gene. This UTR

was also identified as abundantly expressed in exponential culture (Arnvig et al., 2011).

There is a 3' UTR for Rv0292 *(eccE3,* also a hub in the *'violet'* module) that is antisense to

a large part of the 3' end of Rv0293c which has a converging orientation to Rv0292 (Supp

figure S12). Rv0293c is found in a different module *('turquoise')* and has a 3' UTR in the

'lightsteelblue1' module. The overlapping 3' ends of the genes could function to regulate

transcription, possibly to facilitate bi-directional termination brought about by RNAP

collision.

CONCLUSION

This paper presents a large-scale network analysis of over 7000 transcripts expressed by Mtb under a variety of conditions. The modules group together clusters of co-expressed protein-coding genes, as well as ncRNA transcripts predicted from RNA-Seq signals. The ncRNAs are unevenly distributed among modules; modules with the highest proportion of sRNAs correlated negatively to exponential growth and correlated positively to hypoxia and the extended hypoxia model *('turquoise', 'blue', 'skyblue')* (Figures 3 and 4), supporting the observation that high levels of ncRNA are associated with Mtb's response to hypoxic stress (Arnvig et al., 2011; Ignatov et al., 2015; Martini et al., 2019). The prevalence of antisense RNA in the hubs of these and other modules, and the fact that the complementary ORF is usually excluded, implicates antisense transcription as part of a regulation strategy through mechanisms of divergent transcription or in order to regulate mRNA stability (Vargas-Blanco & Shell, 2020; Warman et al., 2021); strategies that may differ among the members of the MTBC (Dinan, Adam M. et al., 2014). 3' UTR transcripts in modules distinct from their upstream ORF implies independent function from the ORF. sRNAs generated from 3' UTRs have been reported in other prokaryotes and evidence points to widespread mRNA processing that could release independent transcripts at the 3' end (Dar & Sorek, 2018; Desgranges et al., 2021; Updegrove et al., 2019; Wang et al., 2019). In compact bacterial genomes, 3' UTRs are also found to overlap other 3' UTRs in a converging transcription pattern which may provide a mechanism for regulating the expression or stability of either transcript.

The gene modules presented here are somewhat 'blunt-force instruments' applied to transcripts that are part of overlapping, coordinated responses to various environmental cues, but restricted to a single module grouping. Recent work exploring differentially expressed genes in response to various environmental conditions have revealed highly

integrated adaptation responses. In other words, a single environmental change, e.g. hypoxia or growth on fatty acids or cholesterol, stimulates transcriptomic remodelling across diverse cellular functions, perhaps acting as cues to stimulate anticipatory pathways and ready the pathogen for the next challenge (Aguilar-Ayala et al., 2017; Eoh et al., 2017; Gerrick et al., 2018). Confounders such as dual-function, 'moonlighting', proteins may weaken the correlation of a module with a specific condition and may create noise in otherwise well-connected modules. However, focussing on the best connected transcripts in various modules can uncover the unexpected connections between genes of diverse pathways.

Other methods of network analysis, such as those using deconvolution methods, allow genes to be members of more than one module and are considered less 'noisy' than clustering methods, such as WGCNA. However, these methods require extremely large numbers of samples to perform well, may be subject to batch effect issues between experimental datasets and characterise a limited proportion of the protein-coding transcripts expressed by Mtb (Saelens et al., 2018; Yoo, et al., 2022). Predicting ncRNA from different datasets involves a significant degree of quality control, parameter adjustment and manual curation, limiting the number of datasets that could be included in our analysis. Including more data would most likely strengthen the correlations with certain conditions and improve the overall specificity of the modules. However, the work presented here confirms that ncRNA are important players in adaptation responses, and their associations with the protein-coding genes in their assigned modules provides context for their activity.

The few modules discussed in depth in this paper represent a very limited snapshot of this extensive co-expression network. Modules of interest can be identified by correlations

to experimental conditions, associated GO terms, functional categories, or gene group

enrichment. The supplementary tables provide information about the module association,

membership values, TSSs and for UTRs, the module membership of the adjacent ORFs

for each predicted ncRNA. This analysis can add context to the circumstances of

expression of previously identified ncRNAs and conserved hypothetical proteins by

associating their expression with functionally-characterised protein-coding genes in the

same module, as well as identifying novel ncRNA candidates for further investigation

such as structural analysis, target prediction and ultimately, experimental validation.

References

Aguilar-Ayala, D. A., Tilleman, L., Van Nieuwerburgh, F., Deforce, D., Palomino, J. C.,

Vandamme, P., Gonzalez-Y-Merchand, J. A., & Martin, A. (2017a). The

transcriptome of Mycobacterium tuberculosis in a lipid-rich dormancy model through

RNAseq analysis. *Scientific Reports*, *7*(1), 17665. https://doi.org/10.1038/s41598-

017-17751-x

Aguilar-Ayala, D. A., Tilleman, L., Van Nieuwerburgh, F., Deforce, D., Palomino, J. C.,

Vandamme, P., Gonzalez-Y-Merchand, J. A., & Martin, A. (2017b). The

transcriptome of Mycobacterium tuberculosis in a lipid-rich dormancy model through

RNAseq analysis. *Scientific Reports*, *7*(1), 17665–17665. PubMed.

https://doi.org/10.1038/s41598-017-17751-x

Ami, V. K. G., Balasubramanian, R., & Hegde, S. R. (2020). Genome-wide identification of

the context- dependent sRNA expression in Mycobacterium tuberculosis. *BMC*

Genomics, *21*(167), 1–12.

Arnvig, K. B., Comas, I., Thomson, N. R., Houghton, J., Boshoff, H. I., Croucher, N. J.,

Rose, G., Perkins, T. T., Parkhill, J., Dougan, G., & Young, D. B. (2011). Sequence-

Based Analysis Uncovers an Abundance of Non-Coding RNA in the Total

Transcriptome of Mycobacterium tuberculosis. *PLOS Pathogens*, *7*(11), e1002342.

https://doi.org/10.1371/journal.ppat.1002342

Arnvig, K. B., & Young, D. B. (2009). Identification of small RNAs in Mycobacterium

tuberculosis. *Molecular Microbiology*, *73*(3), 397–408.

https://doi.org/10.1111/j.1365-2958.2009.06777.x

Arnvig, K., & Young, D. (2012). Non-coding RNA and its potential role in Mycobacterium

tuberculosis pathogenesis. *RNA Biology*, *9*(4), 427–436.

https://doi.org/10.4161/rna.20105

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P.,

903    Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L.,

904    Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G.

905    M., & Sherlock, G. (2000). Gene Ontology: Tool for the unification of biology.

906    *Nature Genetics*, *25*(1), 25–29. https://doi.org/10.1038/75556

907 Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

908    Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society:*

909    *Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-

910    6161.1995.tb02031.x

911 Bhusal, R. P., Bashiri, G., Kwai, B. X. C., Sperry, J., & Leung, I. K. H. (2017). Targeting

912    isocitrate lyase for the treatment of latent tuberculosis. *Drug Discovery Today*, *22*(7),

913    1008–1016. https://doi.org/10.1016/j.drudis.2017.04.012

914 Bidnenko, E., & Bidnenko, V. (2018). Transcription termination factor Rho and microbial

915    phenotypic heterogeneity. *Current Genetics*, *64*(3), 541–546.

916    https://doi.org/10.1007/s00294-017-0775-7

917 Canestrari, J. G., Lasek-Nesselquist, E., Upadhyay, A., Rofaeil, M., Champion, M. M., Wade,

918    J. T., Derbyshire, K. M., & Gray, T. A. (2020). Polycysteine-encoding leaderless

919    short ORFs function as cysteine-responsive attenuators of operonic gene expression in

920    mycobacteria. *Molecular Microbiology*, *114*(1), 93–108.

921    https://doi.org/10.1111/mmi.14498

922 Chao, Y., Papenfort, K., Reinhardt, R., Sharma, C. M., & Vogel, J. (2012). An atlas of Hfq-

923    bound transcripts reveals 3' UTRs as a genomic reservoir of regulatory small RNAs.

924    *The EMBO Journal*, *31*(20), 4005–4019. https://doi.org/10.1038/emboj.2012.229

925 Chen, J., & Xie, J. (2011). Role and regulation of bacterial LuxR-like regulators. *Journal of*

926    *Cellular Biochemistry*, *112*(10), 2694–2702. https://doi.org/10.1002/jcb.23219

Chetal, K., & Janga, S. C. (2015). OperomeDB: A Database of Condition-Specific

Transcription Units in Prokaryotic Genomes. *BioMed Research International*, *2015*,

318217–318217. PubMed. https://doi.org/10.1155/2015/318217

Cortes, T., Schubert, O. T., Rose, G., Arnvig, K. B., Comas, I., Aebersold, R., & Young, D.

B. (2013). Genome-wide mapping of transcriptional start sites defines an extensive

leaderless transcriptome in Mycobacterium tuberculosis. *Cell Reports*, *5*(4), 1121–

1131. https://doi.org/10.1016/j.celrep.2013.10.031

Dar, D., Shamir, M., Mellin, J. R., Koutero, M., Stern-Ginossar, N., Cossart, P., & Sorek, R.

(2016). Term-seq reveals abundant ribo-regulation of antibiotics resistance in

bacteria. *Science*, *352*(6282), aad9822. https://doi.org/10.1126/science.aad9822

Dar, D., & Sorek, R. (2018). Bacterial noncoding RNAs excised from within protein-coding

transcripts. *MBio*, *9*(5). https://doi.org/10.1128/mBio.01730-18

Del Portillo, P., García-Morales, L., Menéndez, M. C., Anzola, J. M., Rodríguez, J. G.,

Helguera-Repetto, A. C., Ares, M. A., Prados-Rosales, R., Gonzalez-y-Merchand, J.

A., & García, M. J. (2019). Hypoxia Is Not a Main Stress When Mycobacterium

tuberculosis Is in a Dormancy-Like Long-Chain Fatty Acid Environment. *Frontiers in*

Cellular and Infection Microbiology, *8*, 449–449.

Desgranges, E., Barrientos, L., & Caldelari, I. (2021). The 3'UTR-derived sRNA RsaG

coordinates redox homeostasis and metabolism adaptation in response to glucose-6-

phosphate uptake in Staphylococcus aureus. *Molecular Microbiology*.

https://doi.org/10.1111/MMI.14845

D'Halluin, A., Polgar, P., Kipkorir, T., Patel, Z., Cortes, T., & Arnvig, K. B. (2022). Term-

seq reveals an abundance of conditional, Rho-dependent termination in

Mycobacterium tuberculosis. *BioRxiv*, 2022.06.01.494293.

https://doi.org/10.1101/2022.06.01.494293

Dinan, Adam M., Tong, Pin, Lohan, Amanda J., Conlon, Kevin M., Miranda-CasoLuengo

Aleksandra A., Malone, Kerri M., Gordon, Stephen V., & Loftus, Brendan J. (2014).

Relaxed Selection Drives a Noisy Noncoding Transcriptome in Members of the

Mycobacterium tuberculosis Complex. *MBio*, *5*(4), e01169-14.

https://doi.org/10.1128/mBio.01169-14

Du, P., Sohaskey, C. D., & Shi, L. (2016). Transcriptional and physiological changes during

Mycobacterium tuberculosis reactivation from non-replicating persistence. *Frontiers*

in Microbiology, *7*(AUG). https://doi.org/10.3389/fmicb.2016.01346

Dutta, D. (2018). Advance in Research on Mycobacterium tuberculosis FabG4 and Its

Inhibitor. *Frontiers in Microbiology*, *9*.

https://www.frontiersin.org/article/10.3389/fmicb.2018.01184

Eoh, H., Wang, Z., Layre, E., Rath, P., Morris, R., Branch Moody, D., & Rhee, K. Y. (2017).

Metabolic anticipation in Mycobacterium tuberculosis. *Nature Microbiology*, *2*(8),

17084. https://doi.org/10.1038/nmicrobiol.2017.84

Galperin, M. Y., Wolf, Y. I., Makarova, K. S., Vera Alvarez, R., Landsman, D., & Koonin, E.

V. (2021). COG database update: Focus on microbial diversity, model organisms, and

widespread pathogens. *Nucleic Acids Research*, *49*(D1), D274–D281.

https://doi.org/10.1093/nar/gkaa1018

Gerrick, E. R., Barbier, T., Chase, M. R., Xu, R., François, J., Lin, V. H., Szucs, M. J., Rock,

J. M., Ahmad, R., Tjaden, B., Livny, J., & Fortune, S. M. (2018). Small RNA

profiling in mycobacterium tuberculosis identifies mrsi as necessary for an

anticipatory iron sparing response. *Proceedings of the National Academy of Sciences*

of the United States of America, *115*(25), 6464–6469.

https://doi.org/10.1073/pnas.1718003115

Girardin, R. C., & McDonough, K. A. (2020). Small RNA Mcr11 requires the transcription

factor AbmR for stable expression and regulates genes involved in the central

metabolism of Mycobacterium tuberculosis. *Molecular Microbiology*, *113*(2), 504–

520. https://doi.org/10.1111/mmi.14436

Gonzalo-Asensio, J., Mostowy, S., Harders-Westerveen, J., Huygen, K., Hernández-Pando,

R., Thole, J., Behr, M., Gicquel, B., & Martín, C. (2008). PhoP: a missing piece in the

intricate puzzle of Mycobacterium tuberculosis virulence. *PloS One*, *3*(10), e3496–

e3496. PubMed. https://doi.org/10.1371/journal.pone.0003496

Houghton, Joanna, Rodgers, Angela, Rose, Graham, D'Halluin, Alexandre, Kipkorir, Terry,

Barker, Declan, Waddell, Simon J., Arnvig, Kristine B., & Oglesby, Amanda G.

(2021). The Mycobacterium tuberculosis sRNA F6 Modifies Expression of Essential

Chaperonins, GroEL2 and GroES. *Microbiology Spectrum*, *9*(2), e01095-21.

https://doi.org/10.1128/Spectrum.01095-21

Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2009a). Bioinformatics enrichment tools:

Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids*

Research, *37*(1), 1–13. https://doi.org/10.1093/nar/gkn923

Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2009b). Systematic and integrative

analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*,

4(1), 44–57. https://doi.org/10.1038/nprot.2008.211

Ignatov, D. V., Salina, E. G., Fursov, M. V., Skvortsov, T. A., Azhikina, T. L., &

Kaprelyants, A. S. (2015). Dormant non-culturable Mycobacterium tuberculosis

retains stable low-abundant mRNA. *BMC Genomics*, *16*(1), 954.

https://doi.org/10.1186/s12864-015-2197-6

Jiang, J., Lin, C., Zhang, J., Wang, Y., Shen, L., Yang, K., Xiao, W., Li, Y., Zhang, L., &

Liu, J. (2020). Transcriptome Changes of Mycobacterium marinum in the Process of

Resuscitation From Hypoxia-Induced Dormancy. *Frontiers in Genetics*, *10*(February), 1–13. https://doi.org/10.3389/fgene.2019.01359

Jiang, J., Sun, X., Wu, W., Li, L., Wu, H., Zhang, L., Yu, G., & Li, Y. (2016). Construction and application of a co-expression network in Mycobacterium tuberculosis. *Scientific Reports*, *6*(March 2015), 1–18. https://doi.org/10.1038/srep28422

Jiao, X., Sherman, B. T., Huang, D. W., Stephens, R., Baseler, M. W., Lane, H. C., & Lempicki, R. A. (2012). DAVID-WS: a stateful web service to facilitate gene/protein list analysis. *Bioinformatics*, *28*(13), 1805–1806. https://doi.org/10.1093/bioinformatics/bts251

Ju, X., Li, D., & Liu, S. (2019). Full-length RNA profiling reveals pervasive bidirectional transcription terminators in bacteria. *Nature Microbiology*, *4*(11), 1907–1918. https://doi.org/10.1038/s41564-019-0500-z

Kanehisa, M., Sato, Y., & Kawashima, M. (2022). KEGG mapping tools for uncovering hidden features in biological data. *Protein Science*, *31*(1), 47–53. https://doi.org/10.1002/pro.4172

Kapopoulou, A., Lew, J. M., & Cole, S. T. (2011). The MycoBrowser portal: A comprehensive and manually annotated resource for mycobacterial genomes. *Tuberculosis*, *91*(1), 8–13. https://doi.org/10.1016/j.tube.2010.09.006

Kendall, S. L., Burgess, P., Balhana, R., Withers, M., Ten Bokum, A., Lott, J. S., Gao, C., Uhia-Castro, I., & Stoker, N. G. (2010). Cholesterol utilization in mycobacteria is controlled by two TetR-type transcriptional regulators: KstR and kstR2. *Microbiology*, *156*(5), 1362–1371. https://doi.org/10.1099/mic.0.034538-0

Kendall, S. L., Withers, M., Soffair, C. N., Moreland, N. J., Gurcha, S., Sidders, B., Frita, R., Ten Bokum, A., Besra, G. S., Lott, J. S., & Stoker, N. G. (2007). A highly conserved transcriptional repressor controls a large regulon involved in lipid degradation in

Mycobacterium smegmatis and Mycobacterium tuberculosis. *Molecular*

Microbiology, *65*(3), 684–699. https://doi.org/10.1111/j.1365-2958.2007.05827.x

Kipkorir, Terry, Mashabela, Gabriel T., de Wet, Timothy J., Koch, Anastasia, Dawes

Stephanie S., Wiesner, Lubbe, Mizrahi, Valerie, Warner, Digby F., & Henkin, Tina

M. (2021). De Novo Cobalamin Biosynthesis, Transport, and Assimilation and

Cobalamin-Mediated Regulation of Methionine Biosynthesis in Mycobacterium

smegmatis. *Journal of Bacteriology*, *203*(7), e00620-20.

https://doi.org/10.1128/JB.00620-20

Lamichhane, G., Arnvig, K. B., & McDonough, K. A. (2013). Definition and annotation of

(myco)bacterial non-coding RNA. *Tuberculosis*, *93*(1), 26–29.

https://doi.org/10.1016/j.tube.2012.11.010

Langfelder, P., & Horvath, S. (2008). WGCNA: An R package for weighted correlation

network analysis. *BMC Bioinformatics*, *9*. https://doi.org/10.1186/1471-2105-9-559

Larsson, C., Luna, B., Ammerman, N. C., Maiga, M., Agarwal, N., & Bishai, W. R. (2012).

Gene Expression of Mycobacterium tuberculosis Putative Transcription Factors

whiB1-7 in Redox Environments. *PLOS ONE*, *7*(7), e37516.

https://doi.org/10.1371/journal.pone.0037516

Li, Heng. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-*

MEM. https://doi.org/10.48550/arXiv.1303.3997

Lloréns-Rico, V., Cano, J., Kamminga, T., Gil, R., Latorre, A., Chen, W.-H., Bork, P., Glass,

J. I., Serrano, L., & Lluch-Senar, M. (2016). Bacterial antisense RNAs are mainly the

product of transcriptional noise. *Science Advances*, *2*(3), e1501363.

https://doi.org/10.1126/sciadv.1501363

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21.

https://doi.org/10.1186/s13059-014-0550-8

Lu, L., Wei, R., Bhakta, S., Waddell, S. J., & Boix, E. (2021). Weighted gene co-expression

network analysis to identify key modules and hub genes associated with

paucigranulocytic asthma. *Antibiotics*, *10*(97).

https://doi.org/10.3390/antibiotics10020097

Lunge, A., Gupta, R., Choudhary, E., & Agarwal, N. (2020). The unfoldase ClpC1 of

Mycobacterium tuberculosis regulates the expression of a distinct subset of proteins

having intrinsically disordered termini. *Journal of Biological Chemistry*, *295*(28),

9455–9473. https://doi.org/10.1074/jbc.RA120.013456

Mai, J., Rao, C., Watt, J., Sun, X., Lin, C., Zhang, L., & Liu, J. (2019). Mycobacterium

tuberculosis 6C sRNA binds multiple mRNA targets via C-rich loops independent of

RNA chaperones. *Nucleic Acids Research*, *47*(8), 4292–4307.

https://doi.org/10.1093/nar/gkz149

Martini, M. C., Zhou, Y., Sun, H., & Shell, S. S. (2019). Defining the Transcriptional and

Post-transcriptional Landscapes of Mycobacterium smegmatis in Aerobic Growth and

Hypoxia. In *Frontiers in Microbiology* (Vol. 10).

https://www.frontiersin.org/article/10.3389/fmicb.2019.00591

Mehta, M., & Singh, A. (2019). Mycobacterium tuberculosis WhiB3 maintains redox

homeostasis and survival in response to reactive oxygen and nitrogen species. *Free*

Radical Biology and Medicine, *131*, 50–58.

https://doi.org/10.1016/j.freeradbiomed.2018.11.032

Menendez-Gil, P., Caballero, C., Catalan-Moreno, A., Irurzun, N., Barrio-Hernandez, I.,

Caldelari, I., & Toledo-Arana, A. (2020). Differential evolution in 3'UTRs leads to

specific gene expression in Staphylococcus. *Nucleic Acids Research*, *48*.

https://doi.org/10.1093/nar/gkaa047

Menendez-Gil, P., & Toledo-Arana, A. (2021). Bacterial 3′UTRs: A Useful Resource in Post-

transcriptional Regulation. *Frontiers in Molecular Biosciences*, *7*.

https://www.frontiersin.org/article/10.3389/fmolb.2020.617633

Modlin, S. J., Afif, E., Deepika, G., Zlotnicki, A. M., Dillon, N. A., Dhillon, N., Kuo, N.,

Robinhold, C., Chan, C. K., Baughn, A. D., & Valafar, F. (2021). Structure-Aware

Mycobacterium tuberculosis Functional Annotation Uncloaks Resistance, Metabolic,

and Virulence Genes. *MSystems*, *0*(0), e00673-21.

https://doi.org/10.1128/mSystems.00673-21

Moores, A., Riesco, A. B., Schwenk, S., & Arnvig, K. B. (2017). Expression, maturation and

turnover of DrrS, an unusually stable, DosR regulated small RNA in Mycobacterium

tuberculosis. *PLOS ONE*, *12*(3), e0174079.

https://doi.org/10.1371/journal.pone.0174079

Ozuna, A., Liberto, D., Joyce, R. M., Arnvig, K. B., & Nobeli, I. (2019). baerhunter: An R

package for the discovery and analysis of expressed non-coding regions in bacterial

RNA-seq data. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btz643

Pandey Amit K. & Sassetti Christopher M. (2008). Mycobacterial persistence requires the

utilization of host cholesterol. *Proceedings of the National Academy of Sciences*,

105(11), 4376–4380. https://doi.org/10.1073/pnas.0711159105

Pawełczyk, J., Brzostek, A., Minias, A., Płociński, P., Rumijowska-Galewicz, A., Strapagiel,

D., Zakrzewska-Czerwińska, J., & Dziadek, J. (2021). Cholesterol-dependent

transcriptome remodeling reveals new insight into the contribution of cholesterol to

Mycobacterium tuberculosis pathogenesis. *Scientific Reports*, *11*(1), 12396.

https://doi.org/10.1038/s41598-021-91812-0

Ponath, F., Hör, J., & Vogel, J. (2022). An overview of gene regulation in bacteria by small

RNAs derived from mRNA 3' ends. *FEMS Microbiology Reviews*, fuac017.

https://doi.org/10.1093/femsre/fuac017

Puniya, B. L., Kulshreshtha, D., Verma, S. P., Kumar, S., & Ramachandran, S. (2013).

Integrated gene co-expression network analysis in the growth phase of

Mycobacterium tuberculosis reveals new potential drug targets. *Molecular*

BioSystems, *9*(11), 2798–2815. https://doi.org/10.1039/c3mb70278b

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015).

Limma powers differential expression analyses for RNA-sequencing and microarray

studies. *Nucleic Acids Research*, *43*(7), e47–e47. https://doi.org/10.1093/nar/gkv007

Rustad, T. R., Harrell, M. I., Liao, R., & Sherman, D. R. (2008). The enduring hypoxic

response of Mycobacterium tuberculosis. *PLoS ONE*, *3*(1), 1–8.

https://doi.org/10.1371/journal.pone.0001502

Rustad, T. R., Minch, K. J., Ma, S., Winkler, J. K., Hobbs, S., Hickey, M., Brabant, W.,

Turkarslan, S., Price, N. D., Baliga, N. S., & Sherman, D. R. (2014). Mapping and

manipulating the Mycobacterium tuberculosis transcriptome using a transcription

factor overexpression-derived regulatory network. *Genome Biology*, *15*(11), 502.

https://doi.org/10.1186/s13059-014-0502-3

Saelens, W., Cannoodt, R., & Saeys, Y. (2018). A comprehensive evaluation of module

detection methods for gene expression data. *Nature Communications*, *9*(1), 1090.

https://doi.org/10.1038/s41467-018-03424-4

Sáenz-Lahoya S., Bitarte N., García B., Burgui S., Vergara-Irigaray M., Valle J., Solano C.,

Toledo-Arana A., & Lasa I. (2019). Noncontiguous operon is a genetic organization

for coordinating bacterial gene expression. *Proceedings of the National Academy of*

Sciences, *116*(5), 1733–1738. https://doi.org/10.1073/pnas.1812746116

Sawyer, E. B., Phelan, J. E., Clark, T. G., & Cortes, T. (2021). A snapshot of translation in

Mycobacterium tuberculosis during exponential growth and nutrient starvation

revealed by ribosome profiling. *Cell Reports*, *34*(5).

https://doi.org/10.1016/j.celrep.2021.108695

Schwenk, S., & Arnvig, K. B. (2018). Regulatory RNA in Mycobacterium tuberculosis, back

to basics. *Pathogens and Disease*, *76*(4). https://doi.org/10.1093/femspd/fty035

Serafini, A., Pisu, D., Palù, G., Rodriguez, G. M., & Manganelli, R. (2013). The ESX-3

Secretion System Is Necessary for Iron and Zinc Homeostasis in Mycobacterium

tuberculosis. *PLoS ONE*, *8*(10), 1–15. https://doi.org/10.1371/journal.pone.0078351

Shell, S. S., Wang, J., Lapierre, P., Mir, M., Chase, M. R., Pyle, M. M., Gawande, R.,

Ahmad, R., Sarracino, D. A., Ioerger, T. R., Fortune, S. M., Derbyshire, K. M., Wade,

J. T., & Gray, T. A. (2015). Leaderless Transcripts and Small Proteins Are Common

Features of the Mycobacterial Translational Landscape. *PLOS Genetics*, *11*(11),

e1005641. https://doi.org/10.1371/journal.pgen.1005641

Šiková, M., Janoušková, M., Ramaniuk, O., Páleníková, P., Pospíšil, J., Bartl, P., Suder, A.,

Pajer, P., Kubičková, P., Pavliš, O., Hradilová, M., Vítovská, D., Šanderová, H.,

Převorovský, M., Hnilicová, J., & Krásný, L. (2019). Ms1 RNA increases the amount

of RNA polymerase in Mycobacterium smegmatis. *Molecular Microbiology*, *111*(2),

354–372. https://doi.org/10.1111/mmi.14159

Singh Prabhat Ranjan, Vijjamarri Anil Kumar, Sarkar Dibyendu, & Federle Michael J.

(2020). Metabolic Switching of Mycobacterium tuberculosis during Hypoxia Is

Controlled by the Virulence Regulator PhoP. *Journal of Bacteriology*, *202*(7),

e00705-19. https://doi.org/10.1128/JB.00705-19

Smith, C., Canestrari, J. G., Wang, A. J., Champion, M. M., Derbyshire, K. M., Gray, T. A.,

& Wade, J. T. (2022). Pervasive translation in Mycobacterium tuberculosis. *ELife*, *11*,

e73980. https://doi.org/10.7554/eLife.73980

Smith, L. J., Stapleton, M. R., Fullstone, G. J. M., Crack, J. C., Thomson, A. J., Le Brun, N.

E., Hunt, D. M., Harvey, E., Adinolfi, S., Buxton, R. S., & Green, J. (2010).

Mycobacterium tuberculosis WhiB1 is an essential DNA-binding protein with a nitric

oxide-sensitive iron–sulfur cluster. *Biochemical Journal*, *432*(3), 417–427.

https://doi.org/10.1042/BJ20101440

Solans, L., Gonzalo-Asensio, J., Sala, C., Benjak, A., Uplekar, S., Rougemont, J., Guilhot,

C., Malaga, W., Martín, C., & Cole, S. T. (2014). The PhoP-Dependent ncRNA Mcr7

Modulates the TAT Secretion System in Mycobacterium tuberculosis. *PLOS*

Pathogens, *10*(5), e1004183. https://doi.org/10.1371/journal.ppat.1004183

Soto-Ramirez, M. D., Aguilar-Ayala, D. A., Garcia-Morales, L., Rodriguez-Peredo, S. M.,

Badillo-Lopez, C., Rios-Muñiz, D. E., Meza-Segura, M. A., Rivera-Morales, G. Y.,

Leon-Solis, L., Cerna-Cortes, J. F., Rivera-Gutierrez, S., Helguera-Repetto, A. C., &

Gonzalez-y-Merchand, J. A. (2017). Cholesterol plays a larger role during

Mycobacterium tuberculosis in vitro dormancy and reactivation than previously

suspected. *Tuberculosis*, *103*(November 2020), 1–9.

https://doi.org/10.1016/j.tube.2016.12.004

Stiens, J., Arnvig, K. B., Kendall, S. L., & Nobeli, I. (2022). Challenges in defining the

functional, non-coding, expressed genome of members of the Mycobacterium

tuberculosis complex. *Molecular Microbiology*, *117*(1), 20–31.

https://doi.org/10.1111/mmi.14862

The Gene Ontology Consortium. (2021). The Gene Ontology resource: Enriching a GOld

mine. *Nucleic Acids Research*, *49*(D1), D325–D334.

https://doi.org/10.1093/nar/gkaa1113

Toledo-Arana, A., & Lasa, I. (2020). Advances in bacterial transcriptome understanding:

From overlapping transcription to the excludon concept. *Molecular Microbiology*,

113(3), 593–602. https://doi.org/10.1111/mmi.14456

Updegrove, T. B., Kouse, A. B., Bandyra, K. J., & Storz, G. (2019). Stem-loops direct precise

processing of 3′ UTR-derived small RNA MicL. *Nucleic Acids Research*, *47*(3),

1482–1492. https://doi.org/10.1093/nar/gky1175

Vargas-Blanco, D. A., & Shell, S. S. (2020). Regulation of mRNA Stability During Bacterial

Stress Responses. *Frontiers in Microbiology*, *11*(September).

https://doi.org/10.3389/fmicb.2020.02111

Voskuil, M. I., Visconti, K. C., & Schoolnik, G. K. (2004). Mycobacterium tuberculosis gene

expression during adaptation to stationary phase and low-oxygen dormancy.

Tuberculosis, *84*(3–4), 218–227. https://doi.org/10.1016/j.tube.2004.02.003

Wade, J. T., & Grainger, D. C. (2014). Pervasive transcription: Illuminating the dark matter

of bacterial transcriptomes. *Nature Reviews Microbiology*, *12*(9), 647–653.

https://doi.org/10.1038/nrmicro3316

Wang, X., Monford Paul Abishek, N., Jeon, H. J., Lee, Y., He, J., Adhya, S., & Lim, H. M.

(2019). Processing generates 3′ ends of RNA masking transcription termination

events in prokaryotes. *Proceedings of the National Academy of Sciences of the United*

States of America, *116*(10), 4440–4445. https://doi.org/10.1073/pnas.1813181116

Warman, E. A., Forrest, D., Guest, T., Haycocks, J. J. R. J., Wade, J. T., & Grainger, D. C.

(2021). Widespread divergent transcription from bacterial and archaeal promoters is a

consequence of DNA-sequence symmetry. *Nature Microbiology*, *6*(6), 746–756.

https://doi.org/10.1038/s41564-021-00898-9

Warner, D. F., Savvi, S., Mizrahi, V., & Dawes, S. S. (2007). A Riboswitch Regulates

Expression of the Coenzyme B12-Independent Methionine Synthase in

Mycobacterium tuberculosis: Implications for Differential Methionine Synthase

Function in Strains H37Rv and CDC1551. *Journal of Bacteriology*, *189*(9), 3655 LP

– 3659. https://doi.org/10.1128/JB.00040-07

World Health Organization. (2021, October 14). *Tuberculosis Fact Sheet*. Tuberculosis.

https://www.who.int/news-room/fact-sheets/detail/tuberculosis

Yoo, Reo, Rychel, Kevin, Poudel, Saugat, Al-bulushi, Tahani, Yuan Yuan, Chauhan,

Siddharth, Lamoureux, Cameron, Palsson, Bernhard O., Sastry, Anand, & Tringe,

Susannah Green. (2022). Machine Learning of All Mycobacterium tuberculosis

H37Rv RNA-seq Data Reveals a Structured Interplay between Metabolism, Stress

Response, and Infection. *MSphere*, *0*(0), e00033-22.

https://doi.org/10.1128/msphere.00033-22

Zhang, B., & Horvath, S. (2005). A General Framework for Weighted Gene Co-Expression

Network Analysis. *Statistical Applications in Genetics and Molecular Biology*, *4*(1).

https://doi.org/10.2202/1544-6115.1128

Zhang, L., Hendrickson, R. C., Meikle, V., Lefkowitz, E. J., Ioerger, T. R., & Niederweis, M.

(2020). Comprehensive analysis of iron utilization by Mycobacterium tuberculosis.

PLOS Pathogens, *16*(2), e1008337. https://doi.org/10.1371/journal.ppat.1008337

Zhou, Y., Huang, H., Zhou, P., & Xie, J. (2012). Molecular mechanisms underlying the

function diversity of transcriptional factor IclR family. *Cellular Signalling*, *24*(6),

1270–1275. https://doi.org/10.1016/j.cellsig.2012.02.008

1218   Zondervan, N. A., Van Dam, J. C. J., Schaap, P. J., Martins dos Santos, V. A. P., & Suarez-

1219   Diez, M. (2018). Regulation of Three Virulence Strategies of Mycobacterium

1220   tuberculosis: A Success Story. *International Journal of Molecular Sciences*, *19*(2).

1221   https://doi.org/10.3390/ijms19020347

1222