Patterns of hydrolysis initiation in P-loop fold nucleoside triphosphatases -

united in diversity

Maria I. Kozlova[1], Daria N. Shalaeva[1], Daria V. Dibrova[1], Armen Y Mulkidjanian[1,2*]

[1]School of Physics and [2]Center of Cellular Nanoanalytics, University of Osnabrueck,

49069 Osnabrueck, Germany

*For correspondence: Armen Y. Mulkidjanian, School of Physics, University of

Osnabrueck, D-49069, Osnabrueck, Germany. Tel. +49-541-969-2698; E-mail:

amulkid@uni-osnabrueck.de

E-mail addresses:

Maria I Kozlova, makozlova@uni-osnabrueck.de, ORCID: 0000-0003-0699-5064

Daria N. Shalaeva, dshalaeva@ uni-osnabrueck.de, ORCID: 0000-0003-0582-2612

Daria V. Dibrova, udavdasha@gmail.com, ORCID: 0000-0002-6954-6437

Armen Y. Mulkidjanian, amulkid@ uni-osnabrueck.de, ORCID: 0000-0001-5844-3064

Abstract

In ubiquitous P-loop fold nucleoside triphosphatases (also known as Walker NTPases), hydrolysis of ATP or GTP is triggered by interaction with an activating partner (usually another protein domain), which is accompanied by insertion of stimulatory moieties (usually arginine or lysine residues) into the catalytic sites. After inspecting over 3600 Mg-NTP-containing structures of P-loop NTPases, we identified those with stimulator(s) inserted into catalytic sites and analysed the patterns of stimulatory interactions. In most cases, at least one stimulator twists gamma-phosphate counter-clockwise by linking the oxygen atoms of alpha- and gamma-phosphates; the twisted gamma-phosphate is stabilized by a hydrogen bond with the backbone amino group of the fourth residue of the Walker A motif. In the remaining cases, the stimulators only interact with gamma-phosphate. The all-pervasive mechanistic interactions of diverse stimulators with the gamma phosphate group suggests its twisting/turning as the trigger for NTP hydrolysis.

1. Introduction

Hydrolysis of nucleoside triphosphates (NTPs), such as ATP or GTP, by so-called P-loop fold nucleoside triphosphatases (also known as Walker ATPases) is one of the key enzymatic reactions. P-loop NTPases are coded by up to 20% gene products in a typical cell. P-loop NTPase domains drive the activity of rotary ATP synthases, DNA and RNA helicases, kinesins and myosins, ABC-transporters, as well as most GTPases, including ubiquitous translation factors, α-subunits of signaling heterotrimeric G-proteins and oncogenic Ras-like GTPases [1-15]. In the ECOD database [16], the topology-level entry "P-loop_NTPase" contains 193 protein families. In the Pfam database [17], the P-loop NTPase clan CL0023 contains 217 families. The main classes of P-loop NTPases, which are listed in Fig. S1 of Supplementary Figures and characterized in the accompanying article [18], were already present in the Last Universal Cellular Ancestor (LUCA) [7, 8, 13, 15, 19-23].

In P-loop NTPases, the eponymous P-loop, together with the first two residues of the following α-helix, usually follows the sequence pattern [G/A]xxxxGK[S/T], known as the Walker A motif [1]. This motif is stabilizing the triphosphate chain of NTP and the cofactor Mg^{2+} ion, see [1, 2, 24] and Fig. 1A-D. The Walker B motif hhhh[D/E], where "h" denotes a hydrophobic residue, is the other common motif of P-loop NTPases that is found on the C-terminal tip of that β-strand, which is opposite the Walker A α-helix, see Fig. 1A,B and [1, 2, 24]. The conserved Asp/Glu residue of the Walker B motif links it with the last [S/T] residue of the Walker A motif by a hydrogen bond (H-bond), see Fig. 1B-D.

Figure 1. P-loop fold NTPases.
A, conserved motifs in P-loop NTPases; **B**, naming of atoms according to IUPAC recommendations for nucleoside triphosphates [25] and typical Mg^{2+} coordination; **C, D,** crystal structures of typical catalytic sites of P-loop NTPases. **C,** Superfamily 1 helicase Pif1 with a transition state analog ADP-AlF$_4^-$ bound (PDB ID 5FHD, [26]). **D,** K$^+$-dependent GTPase MnmE with a transition state analog GDP-AlF$_4^-$ bound (PDB ID 2GJ8 [27]). Color code: Polypeptide chains are shown as gray cartoons; nucleotides, their analogs, and important amino acid residues are shown as sticks, water molecules – as red spheres, Mg^{2+} ions are shown as lime spheres, K$^+$ - as a purple sphere; the conserved Lys residue of Walker A motif is shown in green, and the amino acid residue three residues before it is highlighted in magenta; the conserved Asp residue of Walker B motif is shown in dark red, and the Arg fingers are shown in blue. In those amino acid residues that are shown as sticks, the oxygen atoms are colored red, and the nitrogen atoms are colored blue. In the AlF$_4^-$ moiety, the Al atom is colored gray, and the fluoride atoms are colored light blue. Switch I/K-loop motif is shown in lilac, the conserved Thr^{SwI} is highlighted in yellow.

An uncontrolled NTP hydrolysis would be perilous for cell survival. Therefore, a specific

feature of most P-loop ATPases is their activation before each turnover. Usually, an ATP- or

GTP-bound P-loop domain interacts with its cognate activating partner, which could be a

domain of the same protein, a separate protein and/or a DNA/RNA molecule. Upon this

interaction, specific stimulatory moieties, usually Arg or Lys residues ("Arg/Lys finger(s)",

Fig. 1C), are inserted into the catalytic site and promote the cleavage of γ-phosphate [4, 12, 27-

36]. As a rule, large-scale conformational changes accompany both the binding of NTP

("closing" of the catalytic pocket) and the NTP hydrolysis ("opening" of the catalytic pocket);

these structural changes are coupled to useful mechanical work and drive most biomolecular

motors, see, e.g. [4, 37-39]. In this paper, however, we focus only on the interaction of

stimulators with the NTP molecules within catalytic pockets.

Important hints for clarifying the events inside these pockets are provided by their structures

with bound transition state (TS) analogs, such as $NDP:AlF_4^-$ or $NDP:MgF_3^-$ or $NDP-VO_4^-$

complexes [4, 28, 29, 35, 40-44]. It is anticipated that these analogues promote the formation

of a full-fledged active site in a pre-catalytic state. In such complexes, a water molecule is often

located apically to the analog of γ-phosphate, which corroborates the earlier suggestions that

γ-phosphate cleavage is triggered by the apical nucleophilic attack of a polarized water

molecule (W_{cat}), after its deprotonation to OH^-_{cat}, on γ-phosphate, see Fig. 1C, D; the

accompanying article [18] and [12, 35, 44-49]. Also, TS analogs were shown to promote the

interaction of the activator with the NTPase domain and therefore often disclose the interaction

of the stimulator with the triphosphate chain, as shown in Fig. 1C, D, scrutinized in the

accompanying article [18], and discussed in [28, 29, 35, 36, 40-42, 44, 50-59].

There is no consensus on how do stimulators initiate the NTP hydrolysis. Warshel and

colleagues suggested that stimulators, owing to their positive charge, compensate the negative

charges of oxygen atoms of triphosphate and make the γ-phosphorus atom (P^G) more prone to

the nucleophilic attack by a water molecule [60-62]. It was also suggested that the pushing of

bound water molecules out of the binding pocket into the bulk solution by the stimulatory

arginine finger provides an entropic gain [63]. Jin and colleagues proposed reorientation of the

W_{cat} molecule into the attack position by the stimulator [44, 64, 65]. On a later reaction step,

stimulators were implied in compensating for the negative charge that develops at β-phosphate

upon the breakaway of γ-phosphate [49, 66]. Recently, Gerwert and colleagues proposed, for

small GTPases, that the stimulatory Arg finger rotates the α-phosphate group towards an

eclipsed conformation with respect to β- and γ-phosphates, which would destabilize the

triphosphate chain [67-69].

Elsewhere [70], we have focused on P-loop NTPases that are stimulated not by Arg/Lys

residues but by K^+ and Na^+ ions, see [27, 33, 71-74]. We performed molecular dynamics (MD)

simulations of the Mg^{2+}-GTP-containing dimer of K^+-dependent tRNA-modifying enzyme

MnmE (see Fig. 1D) and compared their results with MD simulations of water-dissolved Mg^{2+}-

GTP and Mg^{2+}-ATP complexes in the presence of monovalent cations. In the latter case, one

of the monovalent cations bound between the O^{2A} and O^{3G} atoms of α- and γ-phosphates, in

the so-called AG site, which was accompanied by rotation of α-phosphate - as the least

constrained phosphate group - yielding a fully eclipsed configuration of the triphosphate chain

[70]. These data agreed with the results of Gerwert and colleagues who used a Mg^{2+}-

metyltriphosphate complex – as a simple mimic of Mg^{2+}-GTP – in their modeling the

interaction with an Arg finger [67].

However, MD simulations of the GTP-containing MnmE dimer showed that α-phosphate was

fully immobilized by bonds between the GTP molecule and the P-loop; these interactions

stabilize the triphosphate chain in an extended, supposedly catalytically prone conformation

see Fig. 1D and [70]. In this case, only the terminal γ-phosphate retained some mobility. The

K^+ ion, by binding between the O^{2A} and O^{3G} atoms of the protein-bound GTP molecule, caused

twisting of the γ-phosphate group by approx. 30°; the rotation was counterclockwise when

viewed from the side of γ-phosphate, see Fig. S2 and [70]. It is important that the K^+ ion can

hardly link these two oxygen atoms without twisting γ-phosphate. The twisted γ-phosphate was

stabilized by a new H-bond between the O^{2G} atom of γ-phosphate and the backbone amide of

Asn226, the fourth residue of the Walker A motif, see Fig. 1A, 1D, S2 and [70]. Both the

twisting of γ-phosphate and its stabilization by an additional H-bond could potentially promote

hydrolysis [70]. Still, these features were not discussed in relation to the catalytic mechanisms

of any P-loop NTPase before. Therefore, we were interested to find out how widespread are

the features we identified in K^+-stimulated GTPase MnmE.

Here, we performed the first, to our best knowledge, comparative structure analysis of

stimulation patterns in all relevant structures of P-loop NTPases. We screened 1484 available

structures of P-loop NTPases with Mg-NTPs or their analogs bound in their catalytic sites (in

total 3666 individual catalytic sites were analyzed). In this search, we relied on a key

observation that diverse P-loop NTPases, despite playing dramatically different roles in the

cell, catalyze essentially the same or similar chemical reactions [9]. It can be expected that

catalysis of congruent chemical reactions in related enzymes is accomplished by homologous

structural elements.

We have found two major types of interaction between stimulatory moieties and triphosphate

chains. In most cases, as in the MnmE GTPase, at least one stimulator links O^{2A} and O^{3G}, which

enforces a counter-clockwise rotation of γ-phosphate. Otherwise, the stimulators interact only

with the γ-phosphate group. In this case, the direction of twisting of γ-phosphate should be

clarified on a case-to-case basis. Also, we noticed that not all stimulators carry a full positive

charge; in diverse classes of P-loop NTPases we found asparagine, serine, and glycine residues

forming stimulatory bonds with oxygen atoms of the triphosphate chain. In general, the only

feature that seems to be common to all the studied P-loop NTPases is the mechanistic

interaction of diverse stimulators with the γ-phosphate group.

This common feature allowed us to suggest that stimulators may trigger NTP hydrolysis by

displacing the oxygen atoms of γ-phosphate, in particular, the O^{1G} atom that coordinates the

Mg^{2+} ion. In the accompanying article [18] we further consider the functional consequences of

the γ-phosphate movement and suggest a common catalytic mechanism for P-loop fold

NTPases.

150 2. **Results**

151 **2.1. Generic numbering of key amino acid residues for P-loop NTPases**

Historically, different families of P-loop NTPases were studied by different research

communities, each developing its own terminology. Therefore, no generic amino acid

numbering for P-loop NTPases exists. For convenience, and by analogy with the Ballesteros-

Weinstein numbering scheme for the G-protein coupled receptors [75], we introduce here a

generic residue numbering for the conserved regions of P-loop NTPases. Without this generic

nomenclature, our comparison of stimulatory mechanisms in diverse classes of P-loop

NTPases, as performed here and in the accompanying paper of Kozlova et al. [18], would be

hardly possible.

First, we chose highly conserved residues as benchmark references. In the Walker A motif, this

is the conserved Lys residue (K^{WA}) that forms H-bonds with the O^{1B} oxygen atom of β-

phosphate and O^{2G} atom of γ-phosphate, see Fig. 1. In the Walker B motif, we have chosen the

conserved [Asp/Glu]^{WB} residue ([D/E]^{WB}) that makes an H-bond with the last Ser/Thr residue

of the Walker A motif (see Fig. 1) and is involved in the coordination of the Mg^{2+} ion.

In addition to universal Walker A and Walker B motifs, we also invoked the Switch I motif

that is located between the Walker A and Walker B motifs in NTPases of TRAFAC

(TRAnslation FACtors) class, see Fig. S1 and [7, 76]. The Switch I has only a single strictly

conserved [Thr/Ser]^{SwI} residue ([T/S] ^{SwI}, colored yellow in Fig. 1D) that can be used as a

reference. In NTPases of the TRAFAC class , the side chain of this [T/S]^{SwI} residue coordinates

the Mg^{2+} ion, its backbone amino group (hereafter **HN**) forms a H-bond with γ-phosphate, and

its backbone carbonyl group (hereafter **CO**) stabilizes W_{cat}, see Fig. 1D.

In the following, we number the amino acids of the Walker A, Switch I, and Walker B motifs

relatively to the reference residues, as shown in Fig. 1. In this case, the Asp226 residue of

MnmE GTPase, the fourth residue of Walker A motif, which binds the K^+ ion by its side chain

and makes a H-bond with O^{2G} via its backbone **HN** group (see Fig. 1D, S2 and [70]), is denoted

as Asp226^{K-3} or Asp^{K-3}. Its backbone **HN** group proper is then labeled as **HN**$^{K-3}$.

To distinguish the interaction of the P-loop domain with its cognate activating partner from the

consequent insertion of, say, an Arg finger into the catalytic site, we consistently refer to those

elements that are poked into catalytic sites to stimulate hydrolysis as 'stimulatory moieties' or

'stimulators'; see, for instance, the dark blue Arg fingers in Fig. 1C or the K^+ ion in Fig. 1D.

We follow Wittinghofer and colleagues who wrote about "GAP stimulated GTPase activity"

in one of their pioneering works [77]. Furthermore, the original Latin meaning of "stimulus" –

"a sharp stick used to poke cattle to get them to keep moving" (quoted from

https://www.dictionary.com/browse/stimulus) - nicely describes the function of Lys and Arg

fingers in P-loop NTPases that routinely empower movements of cellular structures.

2.2. Global structural analysis of stimulatory patterns in the whole set of P-loop NTPases with bound Mg-NTP complexes or their analogs

The ever-rising four-digit numbers of P-loop NTPase structures with ATP, GTP, and their analogs bound, as deposited in the PDB, demanded computational approaches. The search in the Protein Data Bank (PDB) at www.rcsb.org [78, 79] for proteins assigned to the entry IPR027417 "P-loop containing nucleoside triphosphate hydrolase" of the Interpro database [80] yielded as many as 1484 structure entries with 3666 catalytic sites with NTP or NTP-mimicking molecules bound (as of 11.09.2019; many of the structures contained several catalytic sites). The criteria for selection of full-fledged catalytic sites from this set and the routine of their subsequent structural analysis are described and depicted in the Methods section. After filtering, we obtained 3136 structures of catalytic sites containing complexes of Mg^{2+} ions with NTPs or NTP-like molecules, these structures were subjected to further analysis. The relevant data for all these catalytic sites are presented in Supplementary Tables 1 and 2 (hereafter Tables S1 and S2, data as of 11.09.2019).

Based on the type of the molecule, the complexes could be sorted into four groups: 1043 sites contained native ATP/GTP molecules; 1612 sites contained bound non-hydrolyzable NTP analogs such as Adenosine 5′-[β,γ-imido]triphosphate, (AMP-PNP), Guanosine 5′-[β,γ-imido]triphosphate (GMP-PNP), Adenosine 5′-[$β,γ$-methylene]triphosphate (AMP-PCP), Guanosine 5′-[$β,γ$-methylene]triphosphate (GMP-PCP), Adenosine 5′-[γ-thio]triphosphate (ATP-γ-S), and Guanosine 5′-[γ-thio]triphosphate (GTP-γ-S); 234 sites contained NDP:fluoride complexes mimicking the substrate state, such as $NDP:BeF_3$ and $NDP:AlF_3$, and 247 sites contained $NDP:AlF_4^-$ (204), $NDP:MgF_3^-$ (10) and $ADP:VO_4^-$ (33) thought to be TS analogs [35, 40, 42-44].

2.2.1. Stabilization of the O^{2G} atom of γ-phosphate by HN^{K-3} of the Walker A motif.

For the MnmE GTPase, we have shown that the insertion of a K^+ ion and its simultaneous interaction with O^{2A}, O^{3B}, and O^{3G} atoms triggered the twisting of γ-phosphate, leading to the formation of a new H-bond between the O^{2G} atom and **HN** of Asn226^{K-3}, see Fig. 1D, S2 and [70]. Generally, the position of $\mathbf{HN}^{K-3}$ in the vicinity of the O^{2G} atom (or the corresponding atom of an NTP analog) is structurally conserved across P-loop NTPases, being determined by the highly conserved H-bond of $\mathbf{HN}^{K-3}$ with the bridging O^{3B} oxygen (see Fig. 1C,D).

To assess the possibility of a transient H-bond formation between $\mathbf{HN}^{K-3}$ and O^{2G} of γ-phosphate, as found in MD simulations of MnmE GTPases [70], we measured corresponding distances in the available structures of P-loop NTPases with bound substrates or their analogs as described and depicted in Methods. The data obtained are presented in Table S1 and Fig. 2, where the H-bond-compatible distance range is highlighted in amber. For simplicity, we used the same threshold of 3.4 Å for the H-F and H-O bonds. On the one hand, this value is somewhat lower than the threshold of 3.5 Å, as suggested for H-bonds in protein structures by Martz [81]. On the other hand, this distance corresponds to the longest F-H-N bond reported for crystalized L-cysteine-hydrogen fluoride [82]. Still, this threshold is rather arbitrary; according to Jeffrey, weak H-bonds in proteins can have donor-acceptor distances up to 4.0 Å long [83].

In all groups of complexes, there is a fraction with distances shorter than 3.4 Å between $\mathbf{HN}^{K-3}$ and the nearest O^G atom or its structural analog (hereafter $\mathbf{HN}^{K-3}$ - O^G distance). For the ATP- and GTP-containing structures, this fraction makes 31% of complexes (326 of 1043 binding sites); for the non-hydrolyzable analogs, it makes 24% of complexes (392 out of 1612 binding sites). Among complexes with NDP:BeF$_3$, the $\mathbf{HN}^{K-3}$ - O^G distance is < 3.4 Å in 20.5% of

cases(35 catalytic sites out of 171), whereas in the case of structures containing NDP:AlF$_3$ this

fraction makes 28.5% of catalytic sites.

However, the distances between **HN^{K-3}** and O^G analog are shorter than 3.4 Å in most structures

with TS analogs as shown in Fig. 2 where the data for the three TS analogs are plotted

separately for clarity. Distances of < 3.4 Å are observed in 61% of complexes with ADP:VO$_4^{3-}$

(out of 33), in seven out of ten NDP:MgF$_3^-$ complexes and in 80% of NDP:AlF$_4^-$ complexes

(163 out of 204). These structures are thought to be TS-like and usually have stimulators

inserted into the catalytic site. Hence, the TS-like structures of catalytic sites correlate with a

H-bond compatible distance between **HN^{K-3}** and O^G analog.

246
Figure 2. Distances between HN^{K-3} and the O^{2G} atom or its analog in the catalytic sites of
P-loop fold NTPases.
For each type of complexes, distances are visualized as a kernel density estimate (KDE) plot,
a boxplot, and individual data points, each point representing one catalytic site in one structure.
For ADP/GDP:MgF$_3^-$ complexes the density plot is not shown because of scarcity of data. The
range of H-bond-compatible lengths is highlighted in amber.

2.2.2. Pre-catalytic configurations in NTP-containing structures.

While H-bond-compatible distances between $\mathbf{HN}^{K-3}$ and O^G or its analog in the case of TS-like structures (Fig. 2) confirmed our earlier suggestion on the importance of this H-bond for stabilization of the TS [70], we were rather surprised to see that the $\mathbf{HN}^{K-3}$–O^G distances were H-bond compatible also in 31% of ATP- and GTP-containing structures (Fig. 2, Table S1). Therefore, we inspected the top 100 high-resolution NTP-containing structures with H-bond compatible $\mathbf{HN}^{K-3}$–O^{2G} distances to clarify their origin.

In principle, an ATP or GTP molecule could be crystalized within an NTPase only if the latter is inactive. Not surprisingly, the ATP or GTP-containing P-loop NTPases which were crystalized in the absence of their cognate activators made the majority in the inspected set. Although in these proteins the $\mathbf{HN}^{K-3}$–O^{2G} distances were indeed H-bond compatible, we did not explore these structures further because they could hardly help in clarifying the mechanisms of hydrolysis stimulation.

Still, we could identify a set of structures where NTP molecules remained not hydrolyzed despite the presence of activating partner(s). In many such complexes, the interaction of W_{cat} with γ-phosphate was hindered by the incompleteness of W_{cat} ligands, specifically caused by mutations, so that the NTP-binding sites were trapped in precatalytic configurations. Some such structures are shown in Fig. 3A-C. In these enzymes, the stimulatory fingers are inserted into the catalytic sites, γ-phosphates are twisted, and the triphosphate chains are in a configuration like that observed upon the MD simulations of MnmE GTPase [70], which is superimposed as a dark-red contour (cf Fig. S2). One can see from Fig. 3A-C that H-bond compatible $\mathbf{HN}^{K-3}$-O^{2G} distances correlate with linking of O^{2A} and O^{3G} atoms by the stimulator, twisting of γ-phosphate, and a more eclipsed configuration of the triphosphate chain.

Figure 3. **Conformation of native ATP molecules in selected crystal structures as compared to MD simulations of the MnmE GTPase.**

Crystal structures of P-loop ATPases in complex with ATP molecules are shown in lime or green colors; they are superposed with conformations of GTP bound to the MnmE GTPase, as sampled from MD simulations of inactive monomer (blue) and active, K^+-bound dimer (red), see also Fig. S2 and [70]. All distances are given in ångströms. **A**. N-ethylmaleimide-sensitive factor (PDB ID 1NSF, [84]). **B**. ATPase MinD (PDB ID 3Q9L, [85]). **C**. Chromosome segregation protein Soj (PDB ID 2BEK, [86]).

2.2.3. Geometry of the ADP:AlF₃ complex in a P-loop NTPase

It is noteworthy that a particularly flat distribution of distances between NH^{K-3} and O^G analog is observed with the AlF_3-containing complexes (Fig. 2), which could be due to the suggested presence of $NDP:MgF_3^-$ complexes in some of the structures deposited in the PDB as $NDP:AlF_3$-containing structures [35, 44, 58]. Earlier it was shown that, depending on pH, aluminum fluorides can make complexes with NDP in two forms, yielding $NDP:AlF_4^-$ or $NDP:AlF_3$ complexes [54]. However, after identification of $NDP:MgF_3^-$ as one more TS analog [42], Blackburn and colleagues argued that all $NDP:AlF_3$ complexes from previously determined structures are, in fact, misassigned $NDP:MgF_3^-$ complexes [35, 44, 58]. They proposed that the low $Al(OH)_3$ solubility above pH 7.5 would trigger the substitution of Al^{3+} for Mg^{2+} that is usually present in the crystallization solution of NTPases in high amounts to promote the NTP binding [35, 44]. Since the atomic numbers of Al and Mg atoms are very similar, specific methods, such as proton-induced X-ray emission spectroscopy (PIXE), are required to determine whether the structure contains $NDP:AlF_3$ or a TS-analog $NDP:MgF_3^-$.

Application of these methods to the crystals that were studied many years ago is hardly
possible.

We manually inspected all the moieties denoted as AlF_3 in the sampled structures of P-loop
NTPases (Table S1). Upon the inspection, we identified an AlF_3:ADP-containing structure of
the Zika Virus helicase (PDB ID 5Y6M, [87]) with a resolution of 2Å, see Fig. 4A. This
structure could not contain a MgF_3^- ion instead of AlF_3 because the crystallization solution
contained no Mg^{2+}, but only Mn^{2+} ions. No manganese fluoride complexes with more than two
F^- ions in the Mn^{2+} coordination sphere could be found in the PDB (as of 25.06.2020). Also,
the inspection of electron density (Fig. 4B) ruled out the possibility of a misassigned AlF_4^-.

Figure 4. Zika virus helicase in complex with an ADP:AlF_3 moiety**.**

A. Nucleotide-binding pocket of Zika virus helicase with a ADP:AlF_3 moiety bound (PDB ID
5Y6M [2]). Water molecules are shown as spheres. Mn^{2+} ion is shown as a purple sphere; the
conserved Lys200[WA] is shown in green, the Gly197 k-and the amino acid residue three residues
before it is highlighted in magenta. The conserved Asp residue of Walker B motif is shown in
pale red; the Arg fingers are shown in cyan. **B**. 2fo-fc electron density for AlF_3 moiety
contoured at 1σ and 2σ.
318

The high resolution of this helicase structure allowed us to determine the geometry of an

Mn^{2+}:ADP:AlF_3 complex in a P-loop ATPase (Fig. 4). The F^3-F^1-F^2-Al dihedral angle is 17.5°

and the distance between the Al atom and W_{cat} is almost 3 Å, as compared with about 2.5 Å in

MgF_3^--containing complexes and about 2.0 Å in AlF_4^--containing complexes [35, 44]. The

HN^{K-3}-O^{2G} distance was 3.97Å, i.e. longer than, on average, in complexes with MgF_3^- and

AlF_4^- (cf. Fig. 2). This difference is unlikely to be due to the replacement of Mg^{2+} by Mn^{2+}

since the radii of the two ions are similar.

Hence, our analysis indicates that P-loop-bound NDP:AlF_3 complexes are present in the PDB.

Their AlF_3 moiety appear to have a substrate-like geometry that differs from that of planar, TS-

like MgF_3^- moieties. Hence, the mixing of NDP:AlF_3 and NDP:MgF_3^- might indeed contribute

to the flatness of distribution of distances to **HN**$^{K-3}$ in the case of AlF_3-containing structures in

Fig. 2.

2.2.4. Different modes of AlF_4^- interaction with the Mg^{2+} ion

Also we noticed the breadth of distance distributions as measured in NDP:AlF_4^{--}-containing

structures (Fig. 2). . Of course, the nucleotide-binding pockets of diverse of P-loop NTPases

could differ somewhat; the main classes of these enzymes, as shown in Fig. S1, split even

before the LUCA [7, 8, 13, 15, 19-23, 88]. Also, the possible artifacts of structure determination

(as addressed in the Supplementary File 1) and uncertainties of such determination should be

accounted for. Still, the manual examination of the NDP:AlF_4^--containing structures revealed

one more reason for the structural differences. It turned out that AlF_4^- moieties can interact

with Mg^{2+} in two different ways, at least, see Fig. 5A-D and Supplementary Table 3 (hereafter

Table S3). In most structures (77% of AlF_4^- complexes), only one fluorine atom interacts with

Mg^{2+}, like its structural counterpart, the O^{1G} atom of γ-phosphate. In this case, the next closest

fluorine atom is on > 3.0 Å from Mg^{2+}, see Fig. 5A, C and Table S3. In some structures,

however, the two fluoride atoms are at similar distances of 2.0-2.7 Å from the Mg^{2+} ion and

both appear to interact with it, which is only possible when the AlF_4^- is rotated by approx. 45°

around the O^{3B}-Al bond, see Fig. 5B, D and Table S3. This interaction is non-physiological

because it usually prevents Mg^{2+} from interaction with one of its physiological ligands (see

Fig. 5B and Table S3). Our manual inspection of the AlF_4^- structures showed that, in general,

structures with a second fluorine atom found within 3 Å from Mg^{2+} (these include structures

with different degrees of AlF_4^- rotation) usually have distortions in the Mg^{2+} coordination

sphere, namely missing ligands, incorrect bonding to β-phosphate or a missing bond to Ser^{K+1}.

Only two catalytic sites out of 14 where two F—Mg distances are shorter than 2.7 Å appear

not to have additional distortions (see Table S3). One of these two structures has a large Ca^{2+}

ion instead of Mg^{2+} so that interactions in the coordination sphere are preserved due to a larger

ionic radius of Ca^{2+} and its ability to bind up to 8 ligands (Fig. 5D).

Accordingly, all structures with two bonds between fluoride atoms and Mg^{2+} (indicated in

Table S3) appear to be suspicious as TS-state analogs because of the non-physiological

coordination of the Mg^{2+} ion; this point is addressed in the Discussion section and also the

accompanying article [18].

Still, the comparison of the same or closely related NTPases with differently bound $NDP:AlF_4^-$

complexes has proven useful. Fig. 5A, B shows two structures of the Family 2 helicases (ASCE

division, SF1/SF2 class, see Fig. S1) with different coordination of the AlF_4^- moiety. One can

see that the non-physiological coordination in Fig. 5B is achieved via counterclockwise rotation

of the more "physiological" configuration of the AlF_4^- moiety in Fig. 5A, whereby the

interactions of AlF_4^- with Lys^{WA} and the stimulatory Arg residues are retained in both

structures. The two stimulatory fingers retain their H-bonds with AlF_4^- in both configurations.

This comparison shows that the residues which bind γ-phosphate appear to be adapted to the

counterclockwise rotation of γ-phosphate by 30-40°. Similar counterclockwise rotation of γ-

phosphate can be inferred from the comparison of G-α protein structures (Kinase-GTPase

division, TRAFAC class, see Fig. S1) in Fig. 5C, D.

Figure 5. Variation in the AlF$_4^-$ interaction with Mg^{2+} ions.

371
Residues are highlighted as in Fig. 1C,D. The Switch I motif is colored yellow, Mg^{2+} ions are
shown as bright green spheres, Ca^{2+} ion is shown as a teal-colored sphere. The distances that
are too long for H-bonds are indicated by red dashed lines. All distances are given in ångströms.
A. The ADP:AlF$_4^-$ binding in the structure of HCV (Hepatitis C Virus) NS3 helicase, PDB
5E4F, chain A [89]. The Mg^{2+} ion is coordinated by the O^{2B} atom of ADP, one fluorine atom
of AlF$_4^-$, the conserved Ser residue of Walker A motif, and three water molecules. **B.** The
ADP:AlF$_4^-$ binding in the structure of double-stranded RNA (dsRNA)-dependent helicase
LGP2 from chicken (PDB 5JAJ [90]). Here, AlF$_4^-$ is rotated counterclockwise and makes two

bonds with the Mg^{2+} ion; the Ser^{K+1} residue is no longer directly coordinating Mg^{2+} and,

instead, is forming an unusual bond with the O^{1A} atom of ADP. **C.** The AlF_4^- binding in a

structure of guanine nucleotide-binding protein $G\alpha_{i3}$ complexed with the regulator of G-protein

signaling 8 (PDB 2ODE [91]). The Mg^{2+} ion is coordinated by the O^{2B} atom of GDP, one

fluorine atom of AlF_4^-, the conserved Ser residue of Walker A motif, the conserved Thr of

Switch I, and two water molecules. **D.** In the structure of transducin $G_{t\alpha}$ complexed with Ca^{2+}

instead of Mg^{2+} (PDB 1TAD [29]), AlF_4^- is rotated and forms two bonds with Ca^{2+}, but all

interactions in the coordination sphere are preserved due to a larger ionic radius of Ca^{2+} and its

ability to bind up to 8 ligands. However, the catalytic site is more open, as evidenced by a

distance of 3.9 Å between Ser^{K+1} and Asp^{WB}.

The data on all found non-physiologically bound AlF_4^- moieties are highlighted pink in Table

S1 and separately summarized in Table S3.

In sum, the data in Fig. 2, 3 and 5 indicate that the insertion of a stimulator and linking the O^{2A}

and O^{3G} atoms leads to the twisting of γ-phosphate and shortening of the $\mathbf{HN}^{K-3}$-O^{2G} distance

in diverse NTP-containing P-loop NTPases in support of our earlier MD simulation data [70].

2.2.5. Patterns of interaction between stimulatory moieties and the phosphate chain

In most families of P-loop NTPases, Arg, Lys, or Asn residues serve as stimulatory moieties.

We used computational approach to inspect the patterns of their interactions with phosphate

chain atoms (or their analogs) in the PDB-deposited structures of P-loop NTPases.

For this purpose, we analyzed, as described in the Methods section, the same 3136 PDB

structures of catalytic sites that contain complexes of Mg^{2+} ions with NTPs or NTP-like

molecules. For each complex, we measured the distances between oxygen atoms of the

triphosphate chain (or their structural counterparts in the NTP analogs) and the amino groups

of Arg, Lys, and Asn side chains within 4 Å radius, see the Methods section for further details

and the scheme of the analysis pipeline. The distances were measured towards the

NE/NH1/NH2 atoms of Arg residues; the NZ atom of Lys residues; and the ND2 atoms of Asn

residues (see Fig. 6 and 7 for the atom naming scheme). The data on all atom pairs and

corresponding distances are summarized in Table S1.

We have found that more than half of analyzed Mg-NTP complexes (60%) have none of the

inspected residue types within the 4Å radius around the β-phosphate oxygen atoms. Those are

structures of P-loop NTPases that were crystallized in the absence of their activating partners

or are stimulated by moieties other than Asn/Arg/Lys (e.g. a monovalent cation or the signature

motif of ABC-ATPases, see below).

In the remaining 1380 catalytic sites of P-loop NTPases, at least one Arg, Asn, or Lys residue

(other than the reference Lys residue of the Walker A motif) was found in the proximity of the

phosphate chain and categorized as a stimulator. The analysis of interactions between Arg, Lys,

and Asn fingers and the phosphate chains has revealed several distinct types of configurations,

which hereafter are called "stimulatory patterns". The Table S2 shows how many proteins were

assigned to each stimulatory pattern.

Our provisional screening of structural information on different classes of P-loop NTPases has

identified Arg, Lys and Asn residues, monovalent cations, and stimulatory polypeptide loops

of ABC-NTPases as "main" stimulators, see [92] and the accompanying article [18]. Below we

describe their interaction patterns.

In addition, we quantitatively evaluated the interaction patterns of Arg, Lys and Asn residues

as main stimulators, as well as interactions of amino acid residues that serve as "auxiliary

residues/stimulators", see the Methods section and Table S1. The interactions of monovalent

cations were quantified in our previous paper [70]. For ABC-NTPases, there was not much to

quantify because their only two available TS-like structures were obtained for the same protein

[93]. We saw that histidine residues may serve as stimulators in several NTPase families (as

discussed in the Supplementary File 1 of the accompanying article [18]); however, for these

families there are no TS-like structures so that we cannot state this for certain. We could not

identify a glutamine residue as a dedicated stimulator in any structure, although Gln residues

serve as auxiliary, W_{cat}-stabilizing "fingers" in many P-loop NTPases, see Fig. 1C, 3, 5 and

Table S1. We discuss this conundrum in the Discussion section.

2.2.5.1. Stimulatory patterns of Arg fingers

Arginine fingers are the most widespread stimulatory moieties among P-loop NTPases. In an

arginine side chain, the positive charge is distributed over three nitrogen atoms of the

guanidinium group. In principle, each of these atoms can interact with the phosphate chain.

Consequently, we observed a variety of interactions for the Arg fingers.

In most cases, the type of stimulatory pattern was assigned automatically based on the H-bond

compatibility of distances between the NTP molecule or its analog and nearby Arg residue(s).

Here we relied on Jeffrey who categorized H-bonds with donor-acceptor distances of 2.2-2.5

Å as "strong, mostly covalent", those with distances of 2.5-3.2 Å as "moderate, mostly

electrostatic", and H-bonds of 3.2-4.0 Å as "weak, electrostatic" [83].

Due to inconsistencies in the atom numbering and differences among NTP analogs, we

measured the distances from the Arg side chain nitrogen atoms to the nearest oxygen atom of

α-phosphate (hereafter Oα) and the nearest oxygen of γ-phosphate (hereafter Oγ) or the

corresponding atom in NTP analogs. Hereafter, for simplicity, we will use "γ-phosphate" both

for γ-phosphate proper and for its analogs.

Several sets of criteria were applied consecutively, with each following criterion applied only

to the cases that did not match any of the previous criteria:

1) If both distances NH1-Oα and NH1-Oγ do not exceed 3.2Å, the interaction type "NH1" was

assigned, meaning that NH1 atom forms H-bonds with both α- and γ-phosphates. Similarly,

"NH2" interaction type was assigned if both distances NH2-Oα and NH2-Oγ were less than

3.2Å.

2) If both distances NH1-Oα and NH1-Oγ do not exceed 4Å, whereas both distances NH2-Oα

and NH2-Oγ are longer than 4Å, the interaction type "NH1 weak" was assigned, meaning that

NH1 atom forms weak interactions with both α- and γ-phosphates. Analogous criteria were

used to assign the "NH2 weak" interaction type.

3) If at least one of the distances NH1-Oγ and NH2-Oγ do not exceed 3.2Å, whereas both

distances NH1-Oα and NH2-Oα are longer than 4Å, the interaction type "only gamma" was

assigned. Similarly, if at least one of the distances NH1-Oγ and NH2-Oγ do not exceed 4Å,

whereas both distances NH1-Oα and NH2-Oα are longer than 4Å, the interaction type "Only

gamma weak" was assigned.

4) If all distances between NH1/NH2 atoms and the nearest oxygen (or fluorine) atoms of α-

and γ-phosphates exceed 4Å, the Arg residue was considered not to be a stimulatory finger

(interaction type "none").

The remaining cases, which did not match any of these criteria, were inspected manually (see

below). After the interaction types were assigned to all structures under investigation, the

interaction types were attributed to particular stimulatory patterns and their frequencies were

assessed, see Table S3.

Arg residues in the proximity of the phosphate chain were identified as stimulatory moieties in

981 cases. Majority of Arg fingers link α- and γ-phosphates by their NH1 or NH2 groups and

fall into NH1, "NH1 weak", NH2, "NH2 weak" interaction types, which together are grouped

into the stimulatory pattern "AG" seen in the case of 63% of all identified Arg fingers (Tables

S1, S2, Fig. 6A-C). Among the structures with TS analogs, the fraction of this stimulatory

pattern reaches 94%. In contrast, in complexes with ATP or GTP molecules, only 56% of

interactions could be categorized in this way.

In most remaining structures, Arg fingers show interaction types "only gamma"/"only gamma

weak" and interact only with oxygen atom(s) of γ-phosphate or their analogs (Fig. 6D,

stimulatory pattern "G"). This stimulatory pattern was identified in 39% of complexes with

ATP/GTP, 47% of complexes with nonhydrolyzable NTP analogs and only 6% of complexes

with TS analogs (Table S2).

The remaining 33 complexes, which account for 3% of all Arg fingers, did not match any of

these patterns. In these 33 cases, one NH1/NH2 atom of the Arg residue forms an H-bond with

α-phosphate, whereas the other NH2/NH1 atom forms another H-bond with γ-phosphate.

Further, we refer to such Y-shaped interactions as "Y-interactions" or "Y-patterns". Since such

Y-interactions are seen only in a small fraction of catalytic sites, we had inspected each of these

sites manually; the results of this inspection are presented in the Supplementary File 1. As

argued and illustrated in Supplementary File 1, there are reasons to consider all cases of Y-

interactions as structure determination/crystallization artifacts of diverse nature.

Figure 6. Examples of different interaction types/stimulatory patterns involving Arg residues. Protein fragments including P-loop domain are shown as grey cartoons, other subunits carrying Arg/Lys fingers are shown in cyan; functionally relevant residues are shown as sticks: P-loop Lys residue is shown in green, Arg and Lys fingers are shown in cyan; Mg cations are shown as green spheres. All distances are given in ångströms.

A. Both α and γ phosphates are coordinated by the NH2 atom, an additional H-bond is formed by NE atom (stimulatory pattern AG); the structure of the Ras-like GTPase RhoA (PDB ID 5HPY, chain B [94]). **B**. Both α and γ phosphates are coordinated by NH1 atom, an additional H-bond is formed by the NH2 atom (stimulatory pattern AG); the structure of GTP-binding protein G(q) subunit α, (PDB ID 5DO9, chain A [95]). **C.** Both α and γ phosphates are coordinated by the NH1 atom, an additional H-bond is formed by NH2 atom (stimulatory pattern AG). An additional Arg residue provides more interactions with γ phosphate. Notably, this residue also interacts with the backbone atom of the α-subunit that provides the main Arg finger; the structure of bovine mitochondrial F_1-ATPase, β-subunit (PDB 1H8E, chain D [41]). **D.** Only the γ-phosphate is coordinated by the NH2 atom of the Arg finger, the NE atom and Lys residue provide additional H-bonds (stimulatory pattern G_{multi}); the structure of circadian clock protein KaiC, (PDB 4TL8, chain C [96]).

Since the guanidinium group of Arg residues can donate several H-bonds, further H-bonds are seen between amino groups of the Arg finger and the oxygen atoms of the γ-phosphate (or its mimicking group). There are two types of such additional bonds: formed by the NE atom and formed by NH1/NH2 groups not involved in the main stimulatory interaction, as exemplified by Fig. 6.

The NE atoms of Arg fingers are often located at the H-bond distances from the γ-phosphate. Such interactions are documented for 10% of all Arg fingers, both for those Arg fingers that interact only with the γ-phosphate (Fig. 6D), and for those fingers that coordinate both α- and γ-phosphate with the NH2 atom (Fig. 6A).

An additional H-bond can be formed also by an NH1/NH2 atom which is not involved in the main stimulatory interaction. Usually, this occurs when one NH1/NH2 atom coordinates both α- and γ-phosphates; in 51% of such complexes, the other atom (NH2/NH1 correspondingly) forms a H-bond with γ-phosphate (Fig. 6B, C). This interaction is particularly common in complexes with TS analogs (77%). Finally, when the Arg finger interacts only with γ-phosphate, it can accept H-bonds both from NH1 and NH2 atoms, as observed in 13% of complexes with such interaction pattern (Table S2). In these cases, the longer H-bond was categorized as the "auxiliary" interaction, see Tables S1 and S2 for the complete data set.

Overall, the NH2/NH1 groups of Arg fingers that interact only with γ-phosphate or its analog are often assisted – in 40% of such complexes additional bonds are provided by the second NH1/NH2 atom, NE atom, or an additional Arg/Lys finger (Fig. 6D). In this case, one can speak about the stimulatory pattern G_{multi}. The Arg fingers in the AG position can also receive assistance (Fig 6C). For example, in the FtsK DNA translocase structure (PDB ID 6T8B, chain C [97]), the Arg residue interacts with α- and γ-phosphates via the NH2 atom, while the Lys finger reaches the γ-phosphate. Arg fingers reaching only γ-phosphate often contact residues involved in the coordination of W_{cat} (see Fig. 6C, D).

2.2.5.2. Stimulatory patterns of Lys fingers

Lys residues were assumed to be present in AG site if both NZ-Oα and NZ-Oγ distances are

shorter than 4Å (stimulatory pattern "AG" see Fig. 7A and Table S1) and to interact only with

γ-phosphate if only the second distance met the criteria (stimulatory pattern "G" in Table S1).

Otherwise, no interaction with a Lys finger was presumed (pattern "None"), see Table S1.

Lys fingers were identified in 141 structures. One typical pattern is with the NZ atom of Lys

interacting with both α- and γ-phosphates, similarly to a K$^+$ ion in K$^+$-dependent P-loop

NTPases, cf Fig. 1D and [70]. Although a Lys finger interacts both with α- and γ-phosphates

in 22% of all cases (we categorize these cases as stimulatory patterns AG, Fig. 7A), the fraction

of such interactions was as high as 84% in complexes with TS analogs (Tables S1, S2). When

the NZ atom of Lys interacts only with the γ-phosphate (pattern "G"), another Arg residue is

also often involved in the interaction with the γ-phosphate (in 78% of cases, see Fig. 6D). Six

of inspected structures had Lys finger coordinating both α- and γ-phosphate and an additional

Arg residue in the proximity of γ-phosphate. All these structures are subunits of the Large T

antigen (PDB ID 1SVM, in complex with ATP [98]), see Table S1 for details.

2.2.5.3. Interaction patterns of Asn fingers

Asn residues were classified as stimulatory fingers when both ND2-Oα and ND2-Oγ distances

were shorter than 4Å (stimulatory pattern "AG", see Fig. 7B, in Table S1). The Asn residues

were found to be in contact with both α- and γ-phosphate groups in 67 complexes. All these

structures belong to myosin or kinesin families (PF00063, PF00225).

More common are auxiliary Asn residues, which were found in the proximity of γ-phosphates

in 248 catalytic sites, in addition to the "main" stimulators, these auxiliary Asn residues are

indicated in Table S1.

562
563 Figure 7. Examples of stimulatory interactions involving moieties other than arginine
564 residues as stimulators. Residues are highlighted as in Fig. 6.
A. Interaction of a Lys finger, as provided by another subunit of the homodimer, with α
and γ phosphates; the structure of ATPase GET3, (PDB 2WOJ, chain B [99]). **B**. Asn finger
in myosin II from *Dictyostelium discoideum* (PDB 1W9J). The The Asn[S-4] finger is provided
by the Switch I motif which is shown in yellow. **C.** Na[+] ion as a stimulator in dynamin, (PDB
2X2E [100]). The switch I/K-loop motif is shown in lilac. Notably, a Gly residue coordinating
Na[+] is also contacting the O[2A] atom of GDP. **D.** The LSGGQ motif in a structure of ABC-
ATPase MBP-Maltose transporter MalK, (PDB 3PUW, chain A [93]). While Ser and Gly
residues are coordinating γ-phosphate via the backbone HN atom and the OG atom of the
conserved serine residue, the Gln residue is contacting the O2' atom of GDP**.**

2.2.5.4. Summary on quantitative analysis of stimulatory interactions of Arg, Lys, and Asn

fingers in P-loop NTPases

As summarized in Tables S1, S2 and shown in Fig. 6 and 7A, B, most of the analyzed P-loop

NTPase complexes with stimulatory Arg, Lys, or Asn residue(s) positioned next to the

triphosphate chain (or its structural analog) possess either a residue providing an amino group

to interact with both α- and γ-phosphates (stimulatory pattern AG, 56,6% with Y-interactions),

or (2) an amino-group-providing residue(s) forming multiple bonds with γ-phosphate

(stimulatory pattern G_{multi}, 25,6%). In the rest of the cases (17,8%), only one amino group is

contacting γ-phosphate (stimulatory pattern G_{lone}).

In the case of TS analogs, the fraction of catalytic sites with stimulatory pattern AG is

remarkably high: all structures with $ADP:VO_4^{3-}$ and $NDP:MgF_3^-$ that possess any kind of

stimulatory residue have it interacting with both α- and γ-phosphates. The same interaction is

observed in 75% of $NDP:AlF_4^-$ complexes whereas 22% of them display the stimulatory pattern

G_{multi}. Only 3% of TS analogs show stimulatory pattern G_{lone} with a single amino group

contacting γ-phosphate.

The complete data on diverse kinds of auxiliary residues additionally stabilizing the negatively

charged γ-phosphate and/or W_{cat} in all the studied structures are presented in Tables S1 and S2.

2.2.5.5. Stimulation by monovalent cations

In at least two clades of P-loop NTPases, monovalent cations serve as stimulators, as already

elucidated in a separate article [70]. First, many TRAFAC class GTPases are stimulated by K^+-

ions; the formation of a cation-binding site requires a particular positioning of the Switch I loop

(dubbed K-loop in K^+-dependent NTPases [27]), which is achieved either by the specific

interaction of the P-loop domain with its activating partner(s) - protein and/or RNA molecules

[70], or can be induced by binding of a TS analog, such as GDP:AlF$_4^-$ in Fig. 1D. The K$^+$ ion

in MnmE GTPase is coordinated by O^{2A}, O^{3B}, and O^{3G} atoms of the triphosphate chain, two

CO groups of the K-loop, and the side chain of Asn^{K-3} (Fig. 1F, S2). [27, 33]). In the unique

eukaryotic protein family of dynamins, the NTP hydrolysis can be stimulated by either K$^+$ or

Na$^+$ ions, see Fig. 7C [100]. Here a Na$^+$ or a K$^+$ ion interacts only with the O^{3B} and O^{3G} atoms

but does not reach the O^{2A} atom, see [70] for further details. We did not quantify the

interactions with monovalent cations here because we have done it already in our previous

paper [70].

Second, in archaeal and eukaryotic RadA/Rad51-like recombinases of the RecA/F$_1$ class, the

positions that are taken by terminal groups of stimulatory Lys/Arg residues in other proteins of

this class are occupied by two K$^+$ ions, which might interact either only with γ-phosphate, or

with the γ-phosphate and W$_{cat}$, as could be inferred from available structures with substrate

analogs bound, see Fig. 6D in the accompanying article [18] and [70, 101, 102]. No TS-like

structures are available for these K$^+$-dependent enzymes.

2.2.5.6. Stimulatory interactions in ABC-NTPases

ABC (*ATP-binding cassette*) NTPases are multidomain proteins that usually operate as

homo- or heterodimers [103]. Members of the ABC class make several families named

alphabetically from A to I [104]. Most of these families contain members that possess

transmembrane domains and operate as genuine ATP-driven membrane transporters where the

P-loop domains hydrolyze ATP. However, the members of ABCE and ABCF families have no

transmembrane domain(s) [105].

As discussed in more detail in the accompanying article [18], activation of ABC-transporter

ATPases (ABC-ATPases) is triggered by the transported substrate and accompanied by

dimerization of P-loop domains. Upon dimerization, each monomer, instead of an amino acid

"finger", inserts a whole signature motif LSGGQ into the catalytic pocket of the other monomer

(Fig. 7D). Some soluble ABC-NTPases have a non-canonical signature motif (e.g. CSAGQ in

Rad50 [106] and xSTFx in MutS [107]). In some cases, the last residue of the motif is Glu or

even Trp [108]. Thus, the serine residue is the most conserved member of the signature motif.

Two structures with ADP:VO$_4$ and ADP:AlF$_4^-$ as TS analogs bound are available for the

maltose transporter complex [93], see Fig. 7D here and Fig. 5D in the accompanying article

[18]. In both structures, the side chain of serine and the backbone **HN** of the second glycine

residue of the signature motif L$\underline{S}$G$\underline{G}$Q interact with the O^{3G} atom of γ-phosphate. The side

chain of serine is located between the α- and γ-phosphates, approximately in the position of the

Na$^+$ ion in dynamin-like proteins, cf Fig. 7C with Fig. 7D.

In sum, P-loop NTPases exhibit many various stimulatory patterns (see Fig.1-7 and Tables S1,

S2) involving diverse atoms of stimulators, different phosphate groups and, in some cases, also

the W$_{cat}$ molecules (Fig. 1C). These interactions can play an instrumental role both in twisting

of the γ-phosphate group upon catalytic transition and, as discussed in more details in the

accompanying paper [18], in constricting the catalytic site and the catalytic proton transfer.

3. Discussion

Here we report the results of a comparative structural analysis of 3136 catalytic sites of P-loop

NTPases with nucleoside triphosphates or their analogs bound. The aim of the analysis was to

find common features in their stimulatory patterns and to use this information for elucidating

the general mechanism of these enzymes.

Our comparative structural analysis with emphasis on TS-like structures showed that highly

diverse stimulatory moieties interact with the triphosphate chain only in two distinct ways that

may complement each other, as discussed below.

3.1. Linking of α-and γ-phosphates by the stimulator

In most classes of P-loop NTPases, at least one stimulator, provided either by the same P-loop

domain or by another domain/protein, gets inserted between α-and γ-phosphates, which implies

the possibility of simultaneous interaction with O^{2A} and O^{3G} atoms (stimulatory pattern AG).

As seen in Fig. 1, 3-7, in many such cases, the stimulator in the AG site is complemented by a

second auxiliary stimulator (finger) that interacts with γ-phosphate, see also Tables S1, S2. The

AG pattern was observed in 56.6% of cases.

Simultaneous interaction of a single amino group or a K^+ ion with O^{2A} and O^{3G} atoms is

possible either if the triphosphate chain bends (as it is observed with ATP or GTP molecules

in water in the presence of Na^+ ions [70]), or when the γ-phosphate group twists. The bending

of the triphosphate chain of a P-loop-bound NTP molecule is not possible because of its

multiple bonds with the protein, see Fig. 1C, D, the accompanying article [18] and [70]. At the

same time, the examination of available structures revealed several cases which can be

considered as evidence of γ-phosphate twisting in the TS (see Fig. 4, 5), in support of our earlier

predictions from MD simulations [70].

Quantification of the interaction types for 3136 computationally analyzed catalytic sites

(Tables S1, S2) showed that the AG stimulatory pattern dramatically prevailed in the structures

containing TS analogs. This pattern was observed in all structures with $ADP:VO_4^{3-}$ and

$NDP:MgF_3^-$ and in 75% of $NDP:AlF_4^-$-containing structures. The fractions of this stimulatory

pattern in structures containing ATP, GTP, or their non-hydrolyzable analogs are smaller

(Table S2). In many cases, the Arg residue is in the AG-site in the presence of a TS analog but

"outside" when the substrate or its analog are bound. Apparently, catalytic sites constrict

additionally in the transition state; we elaborate on this point in the accompanying article [18].

The notable feature is the apparent scarcity – if not complete absence – of Y-patterns with

NH1 and NH2 groups of an Arg finger separately interacting with α- and γ-phosphates,

respectively (see Table S1). The Y-pattern is not observed in a single structure with a bound

TS analog, and it is such structures that enable us to judge with certainty the stimulatory pattern

in a particular ATPase. Our analysis has shown that the few structures with the Y-pattern are

likely to be artifacts either of crystallization or of structure determination, as substantiated in

the Supplementary File 1.

Outside of P-loop NTPases, however, the Y-pattern of Arg interaction is very common,

especially in protein-DNA complexes, where one Arg residue often donates its NH1 and NH2

groups to neighboring phosphate groups of the DNA backbone [109, 110]. In the case of P-

loop NTPases, however, a Y-linked Arg residue would fix the O^{2A} and O^{3G} atoms approx. 6.1

Å apart and prevent the γ-phosphate from twisting. The apparent absence of Y-patterns in the

examined structures of P-loop NTPases can be seen as further evidence in favor of the γ-

phosphate twisting as the key precatalytic configuration change in P-loop NTPases.

Earlier, our MD simulations showed that γ-phosphate, after its twisting by the stimulatory K^+

ion, was stabilized by a new H-bond between HN^{K-3} and O^{2G} [70]. Our structural analysis

strongly indicates that the counter-clockwise twisting of γ-phosphate correlates with the

formation of a new H-bond between the backbone of the P-loop and O^{2G} indeed. Specifically,

H-bond-compatible distances between $\mathbf{HN}^{K-3}$ and the nearest oxygen/fluorine atom are seen in

most structures with TS analogs bound (Fig. 2, Table S1). Twisted γ-phosphates and H-bond

compatible $\mathbf{HN}^{K-3}$ - O^{2G} distances are also seen in diverse NTP-containing proteins which were

crystalized after being trapped in their pre-transition configurations (because of mutated W_{cat}-

stabilizing residues and/or their impaired interaction with W_{cat}, see Fig. 3A-C, Table S1).

Hence, linking the O^{2A} and O^{3G} atoms by the stimulator, independently on whether it is an Arg,

Lys, Asn, residue or a K^+ ion, causes a counter-clockwise twisting of γ-phosphate that

correlates with the formation of a new H-bond between the O^{2G} atom (or its counterpart in NTP

analogs) and the backbone HN^{K-3} group of the P-loop.

699   *3.2. Interaction of the stimulator only with γ-phosphate*

In the remaining 43,3% of P-loop NTPase structures with determined stimulatory pattern, the

stimulators interact only with γ-phosphate (stimulatory pattern G). In most such structures

(25,6%), γ-phosphate is involved in several interactions with distinct amino groups of an Arg

finger and/or with auxiliary stimulators (stimulatory pattern G_{multi}), see Fig. 6D, 7D and [111].

In 17.7% of structures, our computational analysis has reported only one H-bond between the

stimulatory residue and γ-phosphate (stimulatory pattern G_{lone}). However, in many cases, the

crystal structure does not contain all the partners involved in the activation, or additional

stimulatory fingers are present but are too remote because of crystallization artifacts. In fact,

many catalytic sites exhibiting a G_{lone} pattern have counterparts with a richer network of H-

bonds around γ-phosphate in other subunits of homooligomers of the same protein (47.6%) or

in proteins belonging to the same Pfam domain (75%). Consequently, the value of 17.7%

should be at least halved.

While linking of O^{2A} and O^{3G} atoms by a stimulator enforces a counterclockwise twisting of γ-

phosphate, it is not clear yet, what conformational changes are caused by stimulators that

interact only with γ-phosphate. There is indirect evidence that γ-phosphate may be twisted

clockwise in RecA NTPases, see [112, 113]. Also, in the structure of the ABC ATPase of the

maltose transporter (PDB ID 3PUW, [93]) the AlF_4^- moiety is slightly twisted clockwise (Fig.

7D).

Even interacting only with γ-phosphate, the stimulator is often located between α- and γ-

phosphates, as in dynamins (Fig. 7C) or ABC transporters (Fig. 7D) and connected to the

"head" of the NTP molecule. For instance, the Na^+ ion in dynamins, while not reaching the α-

phosphate directly, is connected to it via two noncovalent bonds (Fig. 7C). The signature motif

of ABC-NTPases is H-bonded via conserved Ser and Gln residues to the O2' atom of the ribose

(Fig. 7D). Such connectivity may strengthen the mechanistic impact on γ-phosphate.

In sum, our structural analysis shows that in all cases where stimulators reach only γ-phosphate,

these stimulators, independently on whether they are Arg or Lys residues, K^+ or Na^+ ions, or

signature motifs of ABC-ATPases, are in position to twist or turn the γ-phosphate group.

3.3. Role of mechanistic bonding in the common stimulation mechanism of P-loop NTPases

Looking together at all types of identified stimulatory patterns provides some additional clues

about the mechanisms of hydrolysis stimulation. Without challenging the previously proposed

tentative stimulation effects as referred to in the Introduction section, our structure analysis

indicates that none of so far suggested mechanisms is common to all P-loop NTPases. Indeed,

it is beyond doubt that the positive charges of Arg/Lys fingers or K^+/Na^+ ions, by interacting

with oxygen atoms of γ-phosphate, would make the P^G atom more prone to the nucleophilic

attack, as suggested by Warshel and colleagues [60-62] and as calculated by Rudack et al. [67].

The positive charge of stimulators could also compensate for the negative charge that develops

at β-phosphate upon the breakaway of γ-phosphate [49, 66]. Nevertheless, the absence of a

positive charge on the stimulatory signature motifs of ABC ATPases (Fig. 7D) does not stop

them from triggering ATP hydrolysis. Also, expelling water molecules out of the catalytic

pocket by stimulatory Arg fingers may provide an entropic gain, as suggested by Kotting and

colleges [63]. However, such effects are hardly to be expected when K^+ or Na^+ ions act as

stimulators and immobilize water molecules in the catalytic pocket (Fig. 1D, 7C). Reorientation

of the W_{cat} molecule into the attack position and its polarization, as suggested by Jin and

colleagues [44, 64, 65], can be realized by those Arg or Lys fingers that reach W_{cat} (Fig. 6D),

but not by most other stimulators.

Our analysis points to the importance of mechanistic interaction of stimulators with the γ-

phosphate group, which is the only feature shared by all inspected stimulators. Importance of

this interaction is exemplified by NTPases that are stimulated by moieties with only a minute

positive charge. These are ABC-NTPases, where γ-phosphate binds the signature LSGG[Q/E]

motif via the side chain of its serine residue and the backbone **HN** of the second glycine residue,

see Fig. 7D and [93]. Other examples are the kinesin and myosin families, where the Asn finger

inserts its NH group between α- and γ-phosphates, see Fig. 7B and [114-116]. It is unlikely that

small partial electric charges of Ser or Asn side chains could be decisive for catalysis in these

cases; rather, their mechanistic H-bonding to the O^{3G} atom appears to be the key.

The mechanistic nature of the stimulatory interaction is consistent with the predominance of

Arg residues as stimulators (Table S1-S3). First, a guanidinium group could donate up to three

H-bonds for interaction with the oxygens of triphosphate. Second, the strength of H-bonds

between guanidinium groups and phosphate anions has been shown to be comparable to that

of covalent bonds [117-119].

The importance of mechanistic binding rationalizes the preference for multiple stimulatory

fingers (Table S2), as well as the choice of the stimulatory signature motif by omnipresent

ABC ATPases. This motif is electrically neutral but donates several H-bonds to the O^{3G} atom.

Consequently, the common denominator of stimulatory patterns in diverse P-loop NTPases is

the mechanistic interaction of stimulators with the γ-phosphate group in the transition state;

this interaction is observed in all analyzed TS-like structures (Fig. 1C,D, 3-7, Table S1, S2)

and can be inferred from many other structures, specifically those that can be related to post-

transition states, see [113] and the accompanying paper [18].

Mechanistic interaction with the γ-phosphate group may promote hydrolysis in different ways;

for instance, it may destabilize the $O^{3B} - P^G$ bond and/or make the triphosphate chain almost

fully eclipsed, and/or facilitate the inversion of γ-phosphate, see the discussion in [70] and the

accompanying article [18]. Notably, any rotation of γ-phosphate inevitably disturbs the

coordination sphere of Mg^{2+}, since the O^{1G} atom of γ-phosphate is one of the Mg^{2+} ligands

(Fig. 1B-D). In P-loop NTPases, the O^{1G} atom is negatively charged, so that its displacement,

by affecting the proton affinity of the other five Mg^{2+} ligands, may trigger the deprotonation of

W_{cat}. In more detail, we address this point in the accompanying article [18] where we further

use comparative structure analysis to reconstruct those steps of the catalytic transition that

follow the interaction with the stimulators.

3.4. The puzzling absence of glutamine residues as stimulators

Glutamine residues are involved in stabilization of W_{cat} in many P-loop NTPases, see e.g. Fig.

1C, 3, 4, 6A,B and the accompanying paper [18]. Still, Gln residues, unlike Asn residues (Fig.

7B), could not be identified as actual stimulators in any of catalytic site structures. Notably,

however, Gln residues occupy stimulator-like positions in non-catalytic sites of F_1-ATPases

(see a plethora of entries in Table S1). These sites, however, are non-functional (see also

Supplementary File 1). It is tempting to speculate that the excessive, as compared to Asn,

flexibility of the glutamate side chains does not allow twisting of γ-phosphate.

3.5. Geometry of the AlF₃ moiety in the NDP:AlF₃ complexes.

As a side result, we determined the geometry of an Mn^{2+}:ADP:AlF$_3$ complex in a P-loop

ATPase (Fig. 4) and thus offered a solution to the long-standing controversy as to whether

NDP:AlF$_3$ complexes could form in the catalytic site of P-loop NTPases under appropriate

conditions or whether all such complexes are misassigned NDP:MgF$_3^-$ complexes, see [35, 42,

44, 58]. The Mn^{2+}:ADP:AlF$_3$ complex was crystallized in the absence of Mg and therefore

could not contain a MgF$_3^-$ moiety. The identified AlF$_3$ moiety has substrate-like geometry of

the AlF$_3$ group, unlike the planar, TS-like MgF$_3^-$ moieties. This non-planarity may help to

discriminate such complexes from NDP:MgF$_3^-$ complexes in earlier obtained crystal structures.

Hence, it is realistic to sort out the P-loop-bound NDP:AlF$_3$ complexes, as assigned in the PDB,

into misassigned NDP:MgF$_3^-$ complexes and genuine NDP:AlF$_3$ complexes.

796 *3.6. Unwelcome effects of AlF$_4^-$ binding*

Quite unexpectedly, we found that the AlF$_4^-$ moieties are bound in some catalytic sites in such

a way that two fluorine atoms interact with the Mg^{2+} ion. In most of these cases, the

coordination bond between Mg^{2+} ion and its ligand #4, [Ser/Thr]$^{K+1}$, is lost, leading to a

distortion of the catalytic site (Fig. 5, Tables S1, S3). This finding is alarming because fluoride

complexes are deservedly vaunted as powerful TS analogues, and structures containing them

are commonly interpreted as TS-like [35, 40, 44]. Still, our data show that the integrity of Mg^{2+}

coordination in the presence of NDP: AlF$_4^-$ should be evaluated separately for each enzymatic

structure before linking the corresponding structural data to the catalytic mechanism. Notably,

NDP:AlF$_4^-$ complexes are also used as TS analogues in other enzyme superfamilies [35, 44].

It is important to check whether a similar distortion of catalytic sites by NDP:AlF$_4^-$ complexes

is possible in enzymes other than P-loop NTPases.

Although this discovery was rather unpleasant as such, it provided some useful information.

The "wrong" binding of AlF$_4^-$ was accompanied by its counterclockwise rotation as compared

to "properly bound" AlF$_4^-$ moieties, whereby the interactions of the fluoride atoms with Lys^{WA}

and the stimulator were preserved. It could be inferred that the ligands of γ-phosphate oxygens

are adapted to twisting of the γ-phosphate group.

4. Conclusions and a Flashback

Here, we performed the first, to the best of our knowledge, comparative structural analysis of

the interactions between the stimulatory moieties and substrate molecules in P-loop NTPases

of the major classes. After screening over 3,100 available structures of catalytic sites with

bound Mg-NTPs or their analogs, we found that seemingly different interactions between

completely distinct stimulatory moieties (Arg/Asn/Lys residues, or K^+/Na^+ ions, or LSGGQ/E

motifs) come down to only two stimulatory patterns. In most cases, at least one stimulator links

the O^{2A} atom of α-phosphate with the O^{3G} atom of γ-phosphate, which requires

counterclockwise twisting of the γ-phosphate group. In other cases, stimulators only interact

with the γ-phosphate group. In general, the only shared feature of all the identified stimulators

seems to be the ability to enter a mechanistic interaction with the γ-phosphate group, which

may enable its twisting/turning.

We provide the relevant outlook on the future studies of P-loop NTPases in the last section of

the accompanying paper [18]. Here, instead, we would like to give a flashback.

In 1998, Wittinghofer and colleagues published a seminal review with the telling title "GTPase-

activating proteins: helping hands to complement an active site," where they viewed a variety

of activating molecules as hands with "fingers" [31]. In this context, our data show that

removing γ-phosphate from an NTP molecule resembles plucking an apple from a tree - the

"fingers" seem to need to twist the γ-phosphate group before they can "rip it off".

5. Methods

Structures were selected among those PDB entries that matched the following criteria: (1) the entry is assigned to InterPro record IPR027417; (2) contains an ATP/GTP molecule, or non-hydrolyzable analog of NTP, or a transition-state analog; (3) contains at least one Mg^{2+}, Mn^{2+} or Ca^{2+} ion. We considered X-ray structures as well as cryo-EM and NMR structures. This search yielded 1474 structures with 3666 catalytic sites in them. For the NMR structures, only the first frame containing an NTP analog bound to the P-loop region was included. For X-ray and cryo-EM structures an additional criterion was applied: structures with resolution less than 5 Å were excluded. In the case of structures containing multiple subunits or multiple copies of the same protein, each interaction between a protein and an NTP-like molecule was treated separately, as an individual complex. The pipeline that was applied to each complex is depicted in Fig. 8.

To identify structures with a substrate or its analog bound to the P-loop motif, we applied the following filter: an NTP(-like) molecule was considered bound to the P-loop if both $Mg^{2+}/Ca^{2+}/Mn^{2+}$ cation and the NZ atom of Lys^{WA} can be found within 5Å of the oxygen atoms of the β-phosphate. Furthermore, Lys^{WA} must be followed in the sequence by a Ser or Thr residue. Complexes with molecules annotated as nonhydrolyzable NTP analogs but lacking β- and/or γ-phosphate mimicking groups were not included in the analysis.

In total, we had analyzed 3136 complexes in 1383 structures with various substrates: ATP and GTP, nonhydrolyzable analogs of ATP and GTP, and ADP or GDP molecules associated with γ-phosphate-mimicking moieties (see Fig. 14).

In each catalytic site under consideration, the K-3 residue was identified by its position relative to the P-loop Lys residue. Distances (in Å) were measured from the **HN** of the K-3 residue to

the nearest oxygen or fluorine atom of the γ-phosphate or its mimic. We accounted for

differences in phosphate oxygen atoms numeration among studied structures.

Putative [Asp/Glu]^{WB} residue was identified as follows: distances from all Asp and Glu residues

to [Ser/Thr]$^{K+1}$ were measured, and the closest residue that was preceded by at least three non-

ionizable residues (Glu, Asp, Ser, Thr, Tyr, Lys, Arg and His were considered as ionizable)

was chosen as the partner of [Ser/Thr]$^{K+1}$. Residues located further than 5Å were not

considered.

We have also measured distances from [Ser/Thr]$^{K+1}$ to Mg^{2+}, to ensure the correct binding of

the Mg^{2+} and general reliability of the structure resolution at the binding site (i.e, very long

distance would indicate a disturbed catalytic site or resolution at the site that is insufficient for

our purposes of comparative analysis), and from [Asp/Glu]^{WB} to Mg^{2+}, to identify cases of

direct coordination of Mg^{2+} by the acidic residue (short distances) or disassembled binding

sites (long distances).

We had also inspected the presence of positively charged stimulatory residues near the

phosphate chains in all complexes. To identify such residues, we considered Arg and Lys

residues (excluding the P-loop Lys) nearest to the β-phosphate group oxygen atoms or its

structural analogs. Distances were then measured from the guanidinium group nitrogen atoms

(NE, NH1, NH2) of the Arg residue or from the NZ atom of the Lys residue to the closest

oxygen atom of α-phosphate moiety and to the nearest fluorine or oxygen atom of γ-phosphate

or its mimicking group. Similarly, we checked possible interactions of the phosphate chain

with the ND2 atom of Asn. Possible additional interactions of γ-phosphate oxygens with N

atoms of protein backbone and side chains in vicinity were also listed for each complex. For

all these interactions the distance threshold was 4 Å.

Figure 8. Automated comparative structural analysis of P-loop NTPases. The examination pipeline, as applied to the structure of each selected nucleotide-binding site, is shown. We checked for (i) the presence of the $HN^{K-3} - O^{2G}$ bond, (ii) the length of the $[Asp/Glu]^{WB}$-$[Ser/Thr]^{K+1}$ H-bond and (iii) the type of interaction of stimulatory residues. Scripts used to download and analyze the structures are available at github.com/servalli/pyploop

In the systematic analysis of all available structures, the patterns of Arg finger binding were

assigned automatically, based on the composition of H-bonds between the Arg residue and the

substrate molecule. We had considered each pair of possible donor and acceptor atoms, where

donors were NE/NH1/NH2 atoms of Arg residue (see Fig. 6) and the acceptors were

oxygen/fluorine atoms of the substrate. The presence of an H-bond was inferred from the

atomic distance: an H-bond was stated at distances less or equal 3.2Å, a weak H-bond stated at

distances between 3.2 and 4Å, and no possibility of a H-bond was stated for distances over 4

Å [82, 120, 121]).

To assign the Arg finger types, several sets of criteria were applied consecutively, as described

in the Results. Structures that did not fit any of the criteria were additionally inspected and Arg

binding patterns assigned manually. After all the Arg fingers were categorized, the frequency

of each interaction type was determined together for the automatically and manually assigned

types.

The lysine residues were assumed to be present in AG site if both NZ-Oα and NZ-Oγ distances

are shorter than 4Å (finger type "AG"), and to interact only with γ-phosphate if only the second

distance met the criteria (finger type "G"). Otherwise, no interaction with a Lys finger was

presumed (finger type "None").

For each complex, AG site was interpreted as occupied by an Arg residue if the closest Arg

residue was assigned interaction type "NH1", "NH1 weak", "NH2", "NH2 weak" or "Y-

TYPE", or by a Lys residue if the closest Lys was assigned type "AG".

All other Arg or Lys residues located within 4Å from γ-phosphate/γ-mimic were listed as

present in "G-site".

Proteins were assigned to major classes of P-loop NTPases according to membership in Pfam

families that was determined from PDB-to-Pfam mapping (retrieved 10.10.2020 from

ftp://ftp.ebi.ac.uk/pub/databases/Pfam/mappings/pdb_pfam_mapping.txt). Each chain was

treated separately and only the first Pfam domain included in Pfam clan CL0023 ("P-

loop_NTPase") was used for the assignment. Since many Pfam domains describing P-loop

NTPases were described before a coherent classification of P-loop NTPases was developed [7,

8], some domain names may not be accurate.

Visualization: Structure superposition, manual distance measurements, manual inspection and

structures visualization were performed by Mol* Viewer [122] and PyMol v 2.5.0 [123].

Data availability:

Descriptions of each binding site are available in Table S1. Scripts used to generate and

annotate the data and quickly visualize selected sites listed in Table S1 are available from

github.com/servalli/pyploop.

6. Acknowledgements

Very fruitful discussions with Drs. D.A. Cherepanov, M.Y. Galperin, A. V. Golovin,

Y.Kalaidzidis, E.V. Koonin, B.H. Meier, A. Lupas, N. Voskoboynikova, T. Wiegand and M.

Zereal are highly appreciated. The authors are thankful to Prof. H.-J. Steinhoff for useful

suggestion on improving the manuscript. We also are thankful to Alexander Mulkidzhanyan

for his help during the initial stage of the project. The research was supported by DFG, DAAD,

and the Osnabrueck University (the EvoCell Program and Open Access Publishing Fund).

References

[1] J.E. Walker, M. Saraste, M.J. Runswick, N.J. Gay, Distantly related sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold, EMBO J 1(8) (1982) 945-51.

[2] M. Saraste, P.R. Sibbald, A. Wittinghofer, The P-loop – a common motif in ATP- and GTP-binding proteins, Trends Biochem. Sci. 15(11) (1990) 430-4.

[3] A.E. Gorbalenya, E.V. Koonin, Helicases: amino acid sequence comparisons and structure-function relationships, Curr Opin Struc Biol 3(3) (1993) 419-429.

[4] C.A. Smith, I. Rayment, Active site comparisons highlight structural similarities between myosin and other P-loop proteins, Biophys J 70(4) (1996) 1590-602.

[5] A.F. Neuwald, L. Aravind, J.L. Spouge, E.V. Koonin, AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes, Genome Res 9(1) (1999) 27-43.

[6] E. Muneyuki, H. Noji, T. Amano, T. Masaike, M. Yoshida, F(0)F(1)-ATP synthase: general structural features of 'ATP-engine' and a problem on free energy transduction, Biochim Biophys Acta 1458(2-3) (2000) 467-81.

[7] D.D. Leipe, Y.I. Wolf, E.V. Koonin, L. Aravind, Classification and evolution of P-loop GTPases and related ATPases, J Mol Biol 317(1) (2002) 41-72.

[8] D.D. Leipe, E.V. Koonin, L. Aravind, Evolution and classification of P-loop kinases and related proteins, J Mol Biol 333(4) (2003) 781-815.

[9] V. Anantharaman, L. Aravind, E.V. Koonin, Emergence of diverse biochemical activities in evolutionarily conserved structural scaffolds of proteins, Curr. Opin. Chem. Biol. 7(1) (2003) 12-20.

[10] L.M. Iyer, D.D. Leipe, E.V. Koonin, L. Aravind, Evolutionary history and higher order classification of AAA+ ATPases, J. Struct. Biol. 146(1-2) (2004) 11-31.

[11] L.M. Iyer, K.S. Makarova, E.V. Koonin, L. Aravind, Comparative genomics of the FtsK-HerA superfamily of pumping ATPases: implications for the origins of chromosome segregation, cell division and viral capsid packaging, Nucleic Acids Res 32(17) (2004) 5260-79.

[12] A. Wittinghofer, I.R. Vetter, Structure-function relationships of the G domain, a canonical switch motif, Annu. Rev. Biochem. 80 (2011) 943-971.

[13] A.M. Burroughs, L. Aravind, The Origin and Evolution of Release Factors: Implications for Translation Termination, Ribosome Rescue, and Quality Control Pathways, Int J Mol Sci 20(8) (2019).

[14] L.M. Longo, J. Jablonska, P. Vyas, M. Kanade, R. Kolodny, N. Ben-Tal, D.S. Tawfik, On the emergence of P-Loop NTPase and Rossmann enzymes from a Beta-Alpha-Beta ancestral fragment, Elife 9 (2020).

[15] A. Krishnan, A.M. Burroughs, L.M. Iyer, L. Aravind, Comprehensive classification of ABC ATPases and their functional radiation in nucleoprotein dynamics and biological conflict systems, Nucleic Acids Res 48(18) (2020) 10045-10075.

[16] R.D. Schaeffer, Y. Liao, H. Cheng, N.V. Grishin, ECOD: new developments in the evolutionary classification of domains, Nucleic Acids Res 45(D1) (2017) D296-D302.

[17] R.D. Finn, P. Coggill, R.Y. Eberhardt, S.R. Eddy, J. Mistry, A.L. Mitchell, S.C. Potter, M. Punta, M. Qureshi, A. Sangrador-Vegas, G.A. Salazar, J. Tate, A. Bateman, The Pfam protein families database: towards a more sustainable future, Nucleic Acids Res. 44(D1) (2016) D279-D285.

[18] M.I. Kozlova, D.N. Shalaeva, A.Y. Mulkidjanian, Common mechanism of activated catalysis in P-loop fold nucleoside 1 triphosphatases - in varietate concordia, submitted (2022).

[19] A.N. Lupas, C.P. Ponting, R.B. Russell, On the evolution of protein folds: are similar
motifs in different protein folds the result of convergence, insertion, or relics of an ancient
peptide world?, J Struct Biol 134(2-3) (2001) 191-203.
[20] C.P. Ponting, R.R. Russell, The natural history of protein domains, Annual review of
biophysics and biomolecular structure 31 (2002) 45-71.
[21] J. Söding, A.N. Lupas, More than the sum of their parts: on the evolution of proteins from
peptides, BioEssays 25(9) (2003) 837-46.
[22] J.A. Ranea, A. Sillero, J.M. Thornton, C.A. Orengo, Protein superfamily evolution and
the last universal common ancestor (LUCA), J. Mol. Evol. 63(4) (2006) 513-25.
[23] V. Alva, J. Soding, A.N. Lupas, A vocabulary of ancient peptides at the origin of folded
proteins, eLife 4 (2015) e09410.
[24] M. Kanade, S. Chakraborty, S.S. Shelke, P. Gayathri, A Distinct Motif in a Prokaryotic
Small Ras-Like GTPase Highlights Unifying Features of Walker B Motifs in P-Loop NTPases,
J Mol Biol 432(20) (2020) 5544-5564.
[25] G.M. Blackburn, J. Cherfils, G.P. Moss, N.G.J. Richards, J.P. Waltho, N.H. Williams, A.
Wittinghofer, How to name atoms in phosphates, polyphosphates, their derivatives and mimics,
and transition state analogues for enzyme-catalysed phosphoryl transfer reactions (IUPAC
Recommendations 2016), Pure Appl. Chem. 89(5) (2017) 653-675.
[26] X. Zhou, W. Ren, S.R. Bharath, X. Tang, Y. He, C. Chen, Z. Liu, D. Li, H. Song, Structural
and Functional Insights into the Unwinding Mechanism of Bacteroides sp Pif1, Cell Rep 14(8)
(2016) 2030-9.
[27] A. Scrima, A. Wittinghofer, Dimerisation-dependent GTPase reaction of MnmE: how
potassium acts as GTPase-activating element, EMBO J 25(12) (2006) 2940-51.
[28] D.E. Coleman, A.M. Berghuis, E. Lee, M.E. Linder, A.G. Gilman, S.R. Sprang, Structures
of active conformations of Gi alpha 1 and the mechanism of GTP hydrolysis, Science
265(5177) (1994) 1405-12.
[29] J. Sondek, D.G. Lambright, J.P. Noel, H.E. Hamm, P.B. Sigler, GTPase mechanism of
Gproteins from the 1.7-A crystal structure of transducin alpha-GDP-AIF-4, Nature 372(6503)
(1994) 276-9.
[30] K. Scheffzek, M.R. Ahmadian, W. Kabsch, L. Wiesmuller, A. Lautwein, F. Schmitz, A.
Wittinghofer, The Ras-RasGAP complex: structural basis for GTPase activation and its loss in
oncogenic Ras mutants, Science 277(5324) (1997) 333-8.
[31] K. Scheffzek, M.R. Ahmadian, A. Wittinghofer, GTPase-activating proteins: helping
hands to complement an active site, Trends Biochem Sci 23(7) (1998) 257-62.
[32] T. Ogura, S.W. Whiteheart, A.J. Wilkinson, Conserved arginine residues implicated in
ATP hydrolysis, nucleotide-sensing, and inter-subunit interactions in AAA and AAA+
ATPases, J. Structural Biology 146(1-2) (2004) 106-12.
[33] M.-R. Ash, M.J. Maher, J.M. Guss, M. Jormakka, The cation-dependent G-proteins: in a
class of their own, FEBS Lett. 586(16) (2012) 2218-24.
[34] P. Wendler, S. Ciniawsky, M. Kock, S. Kube, Structure and function of the AAA+
nucleotide binding pocket, Biochim. Biophys. Acta 1823(1) (2012) 2-14.
[35] Y. Jin, R.W. Molt, Jr., G.M. Blackburn, Metal fluorides: Tools for structural and
computational analysis of phosphoryl transfer enzymes, Top. Curr. Chem. (Cham) 375(2)
(2017) 36.
[36] R. Gasper, F. Wittinghofer, The Ras switch in structural and historical perspective, Biol
Chem 401(1) (2019) 143-163.
[37] I.R. Vetter, A. Wittinghofer, Signal transduction - The guanine nucleotide-binding switch
in three dimensions, Science 294(5545) (2001) 1299-1304.

[38] K. Nam, J. Pu, M. Karplus, Trapping the ATP binding state leads to a detailed
understanding of the F1-ATPase mechanism, Proc Natl Acad Sci U S A 111(50) (2014) 17851-
6.
[39] P. Llinas, T. Isabet, L. Song, V. Ropars, B. Zong, H. Benisty, S. Sirigu, C. Morris, C.
Kikuti, D. Safer, H.L. Sweeney, A. Houdusse, How actin initiates the motor activity of Myosin,
Dev Cell 33(4) (2015) 401-12.
[40] A. Wittinghofer, Signaling mechanistics: aluminum fluoride for molecule of the year,
Curr. Biol. 7(11) (1997) R682-R685.
[41] R.I. Menz, J.E. Walker, A.G. Leslie, Structure of bovine mitochondrial F_1-ATPase with
nucleotide bound to all three catalytic sites: implications for the mechanism of rotary catalysis,
Cell 106(3) (2001) 331-41.
[42] D.L. Graham, P.N. Lowe, G.W. Grime, M. Marsh, K. Rittinger, S.J. Smerdon, S.J.
Gamblin, J.F. Eccleston, MgF(3)(-) as a transition state analog of phosphoryl transfer, Chem
Biol 9(3) (2002) 375-81.
[43] D.R. Davies, W.G. Hol, The power of vanadate in crystallographic investigations of
phosphoryl transfer enzymes, FEBS Lett. 577(3) (2004) 315-21.
[44] Y. Jin, N.G. Richards, J.P. Waltho, G.M. Blackburn, Metal fluorides as analogues for
studies on phosphoryl transfer enzymes, Angew. Chem. Int. Ed. Engl. 56(15) (2017) 4110-
4128.
[45] J.R. Knowles, Enzyme-catalyzed phosphoryl transfer reactions, Annu Rev Biochem 49
(1980) 877-919.
[46] F.H. Westheimer, Why Nature Chose Phosphates, Science 235(4793) (1987) 1173-1178.
[47] Z.A. Shabarova, A.A. Bogdanov, Advanced Organic Chemistry of Nucleic Acids, VCH,
Weinheim, 1994.
[48] M.W. Bowler, M.J. GCliff, J.P. Waltho, G.M. Blackburn, Why did Nature select
phosphate for its dominant roles in biology?, New Journal of Chemistry 34 (2010) 784-794.
[49] J.K. Lassila, J.G. Zalatan, D. Herschlag, Biological phosphoryl-transfer reactions:
understanding mechanism and catalysis, Annu Rev Biochem 80 (2011) 669-702.
[50] T. Higashijima, K.M. Ferguson, P.C. Sternweis, E.M. Ross, M.D. Smigel, A.G. Gilman,
The effect of activating ligands on the intrinsic fluorescence of guanine nucleotide-binding
regulatory proteins, J Biol Chem 262(2) (1987) 752-6.
[51] M. Chabre, Aluminofluoride and beryllofluoride complexes: new phosphate analogs in
enzymology, Trends in biochemical sciences 15(1) (1990) 6-10.
[52] B. Antonny, J. Bigay, M. Chabre, A novel magnesium-dependent mechanism for the
activation of transducin by fluoride, FEBS Lett 268(1) (1990) 277-80.
[53] B. Antonny, M. Sukumar, J. Bigay, M. Chabre, T. Higashijima, The mechanism of
aluminum-independent G-protein activation by fluoride and magnesium. 31P NMR
spectroscopy and fluorescence kinetic studies, J Biol Chem 268(4) (1993) 2393-402.
[54] I. Schlichting, J. Reinstein, pH influences fluoride coordination number of the AlFx
phosphoryl transfer transition state analog, Nat Struct Biol 6(8) (1999) 721-3.
[55] D.L. Graham, J.F. Eccleston, C.W. Chung, P.N. Lowe, Magnesium fluoride-dependent
binding of small G proteins to their GTPase-activating proteins, Biochemistry 38(45) (1999)
14981-7.
[56] M. Chaney, R. Grande, S.R. Wigneshwaraj, W. Cannon, P. Casaz, M.T. Gallegos, J.
Schumacher, S. Jones, S. Elderkin, A.E. Dago, E. Morett, M. Buck, Binding of transcriptional
activators to sigma 54 in the presence of the transition state analog ADP-aluminum fluoride:
insights into activator mechanochemical action, Genes Dev 15(17) (2001) 2282-94.
[57] L. Gremer, B. Gilsbach, M.R. Ahmadian, A. Wittinghofer, Fluoride complexes of
oncogenic Ras mutants to study the Ras-RasGap interaction, Biol. Chem. 389(9) (2008) 1163-
71.

[58] N.J. Baxter, G.M. Blackburn, J.P. Marston, A.M. Hounslow, M.J. Cliff, W. Bermel, N.H.
Williams, F. Hollfelder, D.E. Wemmer, J.P. Waltho, Anionic charge is prioritized over
geometry in aluminum and magnesium fluoride transition state analogs of phosphoryl transfer
enzymes, J Am Chem Soc 130(12) (2008) 3952-8.
[59] N. Zhang, M. Buck, Formation of MgF3 (-)-dependent complexes between an AAA(+)
ATPase and sigma(54.), FEBS Open Bio 2 (2012) 89-92.
[60] T.M. Glennon, J. Villa, A. Warshel, How does GAP catalyze the GTPase reaction of Ras?:
A computer simulation study, Biochemistry 39(32) (2000) 9641-9651.
[61] B.R. Prasad, N.V. Plotnikov, J. Lameira, A. Warshel, Quantitative exploration of the
molecular origin of the activation of GTPase, Proc. Natl. Acad. Sci. USA 110(51) (2013)
20509-20514.
[62] S.C. Kamerlin, P.K. Sharma, R.B. Prasad, A. Warshel, Why nature really chose phosphate,
Q. Rev. Biophys. 46(1) (2013) 1-132.
[63] C. Kotting, A. Kallenbach, Y. Suveyzdis, A. Wittinghofer, K. Gerwert, The GAP arginine
finger movement into the catalytic site of Ras increases the activation entropy, Proc. Natl.
Acad. Sci. USA 105(17) (2008) 6260-5.
[64] Y. Jin, R.W. Molt, Jr., J.P. Waltho, N.G. Richards, G.M. Blackburn, [19]F NMR and DFT
analysis reveal structural and electronic transition state features for RhoA-catalyzed GTP
hydrolysis, Angew. Chem. Int. Ed. Engl. 55(10) (2016) 3318-22.
[65] R.W. Molt, Jr., E. Pellegrini, Y. Jin, A GAP-GTPase-GDP-Pi Intermediate Crystal
Structure Analyzed by DFT Shows GTP Hydrolysis Involves Serial Proton Transfers,
Chemistry 25(36) (2019) 8484-8488.
[66] K.A. Maegley, S.J. Admiraal, D. Herschlag, Ras-catalyzed hydrolysis of GTP: a new
perspective from model studies, Proc. Natl. Acad. Sci. USA 93(16) (1996) 8160-6.
[67] T. Rudack, F. Xia, J. Schlitter, C. Kotting, K. Gerwert, Ras and GTPase-activating protein
(GAP) drive GTP into a precatalytic state as revealed by combining FTIR and biomolecular
simulations, Proc. Natl. Acad. Sci. USA 109(38) (2012) 15295-300.
[68] D. Mann, C. Teuber, S.A. Tennigkeit, G. Schroter, K. Gerwert, C. Kotting, Mechanism of
the intrinsic arginine finger in heterotrimeric G proteins, Proc. Natl. Acad. Sci. USA 113(50)
(2016) E8041-E8050.
[69] K. Gerwert, D. Mann, C. Kotting, Common mechanisms of catalysis in small and
heterotrimeric GTPases and their respective GAPs, Biol. Chem. 398(5-6) (2017) 523-533.
[70] D.N. Shalaeva, D.A. Cherepanov, M.Y. Galperin, A.V. Golovin, A.Y. Mulkidjanian,
Evolution of cation binding in the active sites of P-loop nucleoside triphosphatases in relation
to the basic catalytic mechanism, Elife 7 (2018).
[71] K. Yamanaka, J. Hwang, M. Inouye, Characterization of GTPase activity of TrmE, a
member of a novel GTPase superfamily, from *Thermotoga maritima*, J. Bacteriol. 182(24)
(2000) 7078-82.
[72] S. Meyer, S. Bohme, A. Kruger, H.-J. Steinhoff, J.P. Klare, A. Wittinghofer, Kissing G
domains of MnmE monitored by X-ray crystallography and pulse electron paramagnetic
resonance spectroscopy, PLoS Biol. 7(10) (2009) e1000212.
[73] S. Bohme, S. Meyer, A. Kruger, H.J. Steinhoff, A. Wittinghofer, J.P. Klare, Stabilization
of G domain conformations in the tRNA-modifying MnmE-GidA complex observed with
double electron electron resonance spectroscopy, J. Biol. Chem. 285(22) (2010) 16991-7000.
[74] B. Anand, P. Surana, B. Prakash, Deciphering the catalytic machinery in 30S ribosome
assembly GTPase YqeH, PLoS One 5(4) (2010) e9944.
[75] J.A. Ballestros, H. Weinstein, Integrated methods for the construction of three-
dimensional models and computational probing of structure-function relations in G protein-
coupled receptors, Methods Neurosci. 25 (1995) 366-428.

[76] M.V. Milburn, L. Tong, A.M. deVos, A. Brunger, Z. Yamaizumi, S. Nishimura, S.H. Kim,
Molecular switch for signal transduction: structural differences between active and inactive
forms of protooncogenic ras proteins, Science 247(4945) (1990) 939-45.
[77] M. Frech, J. John, V. Pizon, P. Chardin, A. Tavitian, R. Clark, F. McCormick, A.
Wittinghofer, Inhibition of GTPase activating protein stimulation of Ras-p21 GTPase by the
Krev-1 gene product, Science 249(4965) (1990) 169-71.
[78] H.M. Berman, J. Westbrook, Z. Feng, G. Gilliland, T.N. Bhat, H. Weissig, I.N.
Shindyalov, P.E. Bourne, The Protein Data Bank, Nucleic Acids Res 28(1) (2000) 235-42.
[79] S.K. Burley, C. Bhikadiya, C. Bi, S. Bittrich, L. Chen, G.V. Crichlow, C.H. Christie, K.
Dalenberg, L. Di Costanzo, J.M. Duarte, S. Dutta, Z. Feng, S. Ganesan, D.S. Goodsell, S.
Ghosh, R.K. Green, V. Guranovic, D. Guzenko, B.P. Hudson, C.L. Lawson, Y. Liang, R.
Lowe, H. Namkoong, E. Peisach, I. Persikova, C. Randle, A. Rose, Y. Rose, A. Sali, J. Segura,
M. Sekharan, C. Shao, Y.P. Tao, M. Voigt, J.D. Westbrook, J.Y. Young, C. Zardecki, M.
Zhuravleva, RCSB Protein Data Bank: powerful new tools for exploring 3D structures of
biological macromolecules for basic and applied research and education in fundamental
biology, biomedicine, biotechnology, bioengineering and energy sciences, Nucleic Acids Res
49(D1) (2021) D437-D451.
[80] M. Blum, H.Y. Chang, S. Chuguransky, T. Grego, S. Kandasaamy, A. Mitchell, G. Nuka,
T. Paysan-Lafosse, M. Qureshi, S. Raj, L. Richardson, G.A. Salazar, L. Williams, P. Bork, A.
Bridge, J. Gough, D.H. Haft, I. Letunic, A. Marchler-Bauer, H. Mi, D.A. Natale, M. Necci,
C.A. Orengo, A.P. Pandurangan, C. Rivoire, C.J.A. Sigrist, I. Sillitoe, N. Thanki, P.D. Thomas,
S.C.E. Tosatto, C.H. Wu, A. Bateman, R.D. Finn, The InterPro protein families and domains
database: 20 years on, Nucleic Acids Res 49(D1) (2021) D344-D354.
[81] E. Martz, Help, Index & Glossary for Protein Explorer,
http://www.umass.edu/microbio/chime/pe_beta/pe/protexpl/igloss.htm?q=microbio/chime/ex
plorer/igloss.htm, 2001.
[82] V.S. Minkov, V.V. Ghazaryan, E.V. Boldyreva, A.M. Petrosyan, Unusual hydrogen
bonding in L-cysteine hydrogen fluoride, Acta Crystallogr C Struct Chem 71(Pt 8) (2015) 733-
41.
[83] G.A. Jeffrey, An introduction to hydrogen bonding, Oxford University Press1997.
[84] R.C. Yu, P.I. Hanson, R. Jahn, A.T. Brunger, Structure of the ATP-dependent
oligomerization domain of N-ethylmaleimide sensitive factor complexed with ATP, Nat Struct
Biol 5(9) (1998) 803-11.
[85] W. Wu, K.T. Park, T. Holyoak, J. Lutkenhaus, Determination of the structure of the MinD-
ATP complex reveals the orientation of MinD on the membrane and the relative location of the
binding sites for MinE and MinC, Mol Microbiol 79(6) (2011) 1515-28.
[86] T.A. Leonard, P.J. Butler, J. Lowe, Bacterial chromosome segregation: structure and DNA
binding of the Soj dimer--a conserved biological switch, EMBO J 24(2) (2005) 270-82.
[87] X. Yang, C. Chen, H. Tian, H. Chi, Z. Mu, T. Zhang, K. Yang, Q. Zhao, X. Liu, Z. Wang,
X. Ji, H. Yang, Mechanism of ATP hydrolysis by the Zika virus helicase, Faseb J 32(10) (2018)
5250-5257.
[88] C.A. Orengo, J.M. Thornton, Protein families and their evolution-a structural perspective,
Annu. Rev. Biochem. 74 (2005) 867-900.
[89] M. Gu, C.M. Rice, The Spring alpha-Helix Coordinates Multiple Modes of HCV
(Hepatitis C Virus) NS3 Helicase Action, J Biol Chem 291(28) (2016) 14499-509.
[90] E. Uchikawa, M. Lethier, H. Malet, J. Brunel, D. Gerlier, S. Cusack, Structural Analysis
of dsRNA Binding to Anti-viral Pattern Recognition Receptors LGP2 and MDA5, Mol Cell
62(4) (2016) 586-602.
[91] M. Soundararajan, F.S. Willard, A.J. Kimple, A.P. Turnbull, L.J. Ball, G.A. Schoch, C.
Gileadi, O.Y. Fedorov, E.F. Dowler, V.A. Higman, S.Q. Hutsell, M. Sundstrom, D.A. Doyle,

D.P. Siderovski, Structural diversity in the RGS domain and its interaction with heterotrimeric G protein alpha-subunits, Proc Natl Acad Sci U S A 105(17) (2008) 6457-62.

[92] D.N. Shalaeva, D.A. Cherepanov, M.Y. Galperin, A.Y. Mulkidjanian, Comparative analysis of active sites in P-loop nucleoside triphosphatases suggests an ancestral activation mechanism, BioRxiv 439992 [preprint] (2018).

[93] M.L. Oldham, J. Chen, Snapshots of the maltose transporter during ATP hydrolysis, Proc Natl Acad Sci U S A 108(37) (2011) 15152-6.

[94] F. Yi, R. Kong, J. Ren, L. Zhu, J. Lou, J.Y. Wu, W. Feng, Noncanonical Myo9b-RhoGAP Accelerates RhoA GTP Hydrolysis by a Dual-Arginine-Finger Mechanism, J Mol Biol 428(15) (2016) 3043-57.

[95] V.G. Taylor, P.A. Bommarito, J.J. Tesmer, Structure of the Regulator of G Protein Signaling 8 (RGS8)-Galphaq Complex: MOLECULAR BASIS FOR Galpha SELECTIVITY, J Biol Chem 291(10) (2016) 5138-45.

[96] J. Abe, T.B. Hiyama, A. Mukaiyama, S. Son, T. Mori, S. Saito, M. Osako, J. Wolanin, E. Yamashita, T. Kondo, S. Akiyama, Circadian rhythms. Atomic-scale origins of slowness in the cyanobacterial circadian clock, Science 349(6245) (2015) 312-6.

[97] N.L. Jean, T.J. Rutherford, J. Lowe, FtsK in motion reveals its mechanism for double-stranded DNA translocation, Proc Natl Acad Sci U S A 117(25) (2020) 14202-14208.

[98] D. Gai, R. Zhao, D. Li, C.V. Finkielstein, X.S. Chen, Mechanisms of conformational change for a replicative hexameric helicase of SV40 large tumor antigen, Cell 119(1) (2004) 47-60.

[99] A. Mateja, A. Szlachcic, M.E. Downing, M. Dobosz, M. Mariappan, R.S. Hegde, R.J. Keenan, The structural basis of tail-anchored membrane protein recognition by Get3, Nature 461(7262) (2009) 361-6.

[100] J.S. Chappie, S. Acharya, M. Leonard, S.L. Schmid, F. Dyda, G domain dimerization controls dynamin's assembly-stimulated GTPase activity, Nature 465(7297) (2010) 435-40.

[101] X. Qian, Y. He, Y. Wu, Y. Luo, Asp302 determines potassium dependence of a RadA recombinase from *Methanococcus voltae*, J. Mol. Biol. 360(3) (2006) 537-47.

[102] D.V. Dibrova, M.Y. Galperin, E.V. Koonin, A.Y. Mulkidjanian, Ancient systems of sodium/potassium homeostasis as predecessors of membrane bioenergetics, Biochemistry. Biokhimiia 80(5) (2015) 495-516.

[103] D.C. Rees, E. Johnson, O. Lewinson, ABC transporters: the power to change, Nat Rev Mol Cell Biol 10(3) (2009) 218-27.

[104] M. Dean, A. Rzhetsky, R. Allikmets, The human ATP-binding cassette (ABC) transporter superfamily, Genome Res 11(7) (2001) 1156-66.

[105] I.D. Kerr, Sequence analysis of twin ATP binding cassette proteins involved in translational control, antibiotic resistance, and ribonuclease L inhibition, Biochem Biophys Res Commun 315(1) (2004) 166-73.

[106] K.P. Hopfner, A. Karcher, D.S. Shin, L. Craig, L.M. Arthur, J.P. Carney, J.A. Tainer, Structural biology of Rad50 ATPase: ATP-driven conformational control in DNA double-strand break repair and the ABC-ATPase superfamily, Cell 101(7) (2000) 789-800.

[107] J.A. Eisen, A phylogenomic study of the MutS family of proteins, Nucleic Acids Res 26(18) (1998) 4291-300.

[108] A. Decottignies, A. Goffeau, Complete inventory of the yeast ABC proteins, Nat Genet 15(2) (1997) 137-45.

[109] K.A. Schug, W. Lindner, Noncovalent binding between guanidinium and anionic groups: focus on biological- and synthetic-based arginine/guanidinium interactions with phosph[on]ate and sulf[on]ate residues, Chem Rev 105(1) (2005) 67-114.

[110] B.J. Calnan, B. Tidor, S. Biancalana, D. Hudson, A.D. Frankel, Arginine-mediated RNA recognition: the arginine fork, Science 252(5009) (1991) 1167-71.

[111] A.V. Afonin, I.V. Sterkhova, A.V. Vashchenko, M.V. Sigalov, Estimating the energy of
intramolecular bifurcated (three-centered) hydrogen bond by X-ray, IR and 1H NMR
spectroscopy, and QTAIM calculations, J Mol Struct 1163 (2018) 185-196.
[112] D. Lacabanne, T. Wiegand, N. Wili, M.I. Kozlova, R. Cadalbert, D. Klose, A.Y.
Mulkidjanian, B.H. Meier, A. Bockmann, ATP Analogues for Structural Investigations: Case
Studies of a DnaB Helicase and an ABC Transporter, Molecules 25(22) (2020).
[113] A.A. Malaer, N. Wili, L.A. Volker, M.I. Kozlova, R. Cadalbert, A. Dapp, M.E. Weber,
J. Zehnder, G. Jeschke, H. Eckert, A. Bockmann, D. Klose, A.Y. Mulkidjanian, B.H. Meier, T.
Wiegand, Spectroscopic glimpses of the transition state of ATP hydrolysis trapped in a
bacterial DnaB helicase, Nat Commun 12(1) (2021) 5293.
[114] A.J. Fisher, C.A. Smith, J.B. Thoden, R. Smith, K. Sutoh, H.M. Holden, I. Rayment, X-
ray structures of the myosin motor domain of Dictyostelium discoideum complexed with
MgADP.BeFx and MgADP.AlF4, Biochemistry 34(28) (1995) 8960-72.
[115] M.A. Geeves, Review: The ATPase mechanism of myosin and actomyosin, Biopolymers
105(8) (2016) 483-91.
[116] R.A. Cross, Review: Mechanochemistry of the kinesin-1 ATPase, Biopolymers 105(8)
(2016) 476-82.
[117] A.S. Woods, S. Ferre, Amazing stability of the arginine-phosphate electrostatic
interaction, J Proteome Res 4(4) (2005) 1397-402.
[118] L. Shimoni, J.P. Glusker, Hydrogen bonding motifs of protein side chains: descriptions
of binding of arginine and amide groups, Protein Sci 4(1) (1995) 65-74.
[119] C.A. Seipp, N.J. Williams, M.K. Kidder, R. Custelcean, CO2 Capture from Ambient Air
by Crystallization with a Guanidine Sorbent, Angew Chem Int Ed Engl 56(4) (2017) 1042-
1045.
[120] I.K. McDonald, J.M. Thornton, Satisfying hydrogen bonding potential in proteins, J Mol
Biol 238(5) (1994) 777-93.
[121] B. van Beusekom, W.G. Touw, M. Tatineni, S. Somani, G. Rajagopal, J. Luo, G.L.
Gilliland, A. Perrakis, R.P. Joosten, Homology-based hydrogen bond information improves
crystallographic structures in the PDB, Protein Sci 27(3) (2018) 798-808.
[122] D. Sehnal, S. Bittrich, M. Deshpande, R. Svobodova, K. Berka, V. Bazgier, S. Velankar,
S.K. Burley, J. Koca, A.S. Rose, Mol* Viewer: modern web app for 3D visualization and
analysis of large biomolecular structures, Nucleic Acids Res 49(W1) (2021) W431-W437.
[123] W.L. DeLano, The PyMOL Molecular Graphics System, Version 1.7.2.1, Schrödinger,
LLC., 2010.
1260

1261

1262

1263

Supplementary Materials

1264

Supplementary Figures

1265

Supplementary File 1. Detailed analysis of structures, depicting atypical "Y-type"-like

1266

1267 interactions between Arg finger and the phosphate chain region of an NTP-mimicking

1268 molecule.

Supplementary Table 1. Results of the computational analysis of all available structures of

1269

1270 the P-loop proteins in complex with NTPs and NTP-like molecules. Excel table contains the

1271 list of all analyzed structures, together with residues, identified as the P-loop K and K-3

1272 residues, Arg or Lys fingers; and distances from (1) the respective atoms of NTPs/their analogs

1273 to K-3 residues and Arg/Lys fingers, as well as (2) from $[Asp/Glu]^{WB}$ to $[Ser/Thr]^{K+1}$.

Supplementary Table 2. Relative occurrence of activation patterns for Arg, Lys, and Asn

1274

1275 fingers.

Supplementary Table 3. Coordination of the Mg^{2+} ion in AlF_4^--containing structures of P-

1276

1277 loop NTPases

1278

1279