Social learning mechanisms shape transmission pathways through replicate
great tit social networks

Kristina B. Beck[1*], Ben C. Sheldon[1], Josh A. Firth[1]

[1]Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford, Oxford
OX1 3SZ, United Kingdom

*Correspondence:
kbbeck.mail@gmail.com

10
11
Abstract

The emergence and spread of novel behaviours via social learning can lead to rapid population-level
changes whereby the social connections between individuals shape information flow. However,
behaviours can spread via different mechanisms and little is known about how information flow
depends on the underlying social learning rule individuals employ. Here, comparing four different
learning mechanisms, we simulated behavioural spread on replicate empirical social networks of wild
great tits and explored the relationship between individual sociality and the order of acquisition of
seeded behaviours. Our results reveal that, for learning rules dependent on the sum and strength of
social connections to informed individuals, social centrality was strongly related to the order of
acquisition, with more social individuals adopting new behaviours faster than less social individuals.
However, when behavioural adoption depends on the ratio of an individuals' social connections to
informed versus uninformed individuals, social centrality was not related to the order of acquisition.
Finally, we show how specific learning mechanisms may limit behavioural spread within networks.
These findings have important implications for understanding whether and how diverse behaviours
are likely to spread across social systems, the relationship between individuals' sociality and
behavioural acquisition, and therefore the costs and benefits of sociality.

28
29

Introduction
Social learning, in which individuals learn from others, is widespread in the animal kingdom, and
enables individuals to acquire novel behaviours facilitating phenotypic change (Heyes 1994; Hoppitt
and Laland 2013; Whiten 2021). Socially-induced changes in behaviour can spread through a
population and social networks provide the pathways along which behaviour can spread (Hasenjager
et al. 2021). Research increasingly shows how the structure of social networks and individual sociality
together can influence information flow (Voelkl and Noë 2008; Aplin et al. 2012; Kulahci et al. 2016;
Romano et al. 2018). However, information can spread via various social learning mechanisms (Nunn
et al. 2009; Firth 2020; Cantor et al. 2021) and we know surprisingly little about how the relationship
between sociality and information flow depends on the social learning mechanisms at play.

By definition, the social spread of behaviour to a focal individual requires contact with at least one
knowledgeable individual. Frequently, it is intuitively assumed that the extent (i.e. number and
duration) of social contacts to knowledgeable others predicts the likelihood of adoption (Hasenjager
et al. 2021). In this way, behavioural spread is predicted to follow a similar logic to the transmission

of diseases. However, in contrast to disease spread, individuals may employ 'social learning rules'
where they can actively shape how to act on acquired novel information. For example, for more costly
behaviours such as the usage of a novel food type, an individual may only change its behaviour after
the majority of its social contacts consumes the novel food. Consequently, behavioural spread may
require exposure to multiple sources (rather than just one) and depends on the ratio of connections
to both informed and uninformed individuals (rather than just the connections to informed others)
(Centola and Macy 2007; Guilbeault et al. 2018; Firth et al. 2020). Therefore, the type of behaviour
considered and its underlying learning rule can fundamentally influence whether and how behaviour
spreads through a social network (Centola and Macy 2007; Firth et al. 2020).

In fact, previous studies have revealed several social learning strategies that suggest a range of
different underlying social learning mechanisms (Hoppitt and Laland 2013; Kendal et al. 2018). For
instance, an increasingly-reported learning mechanism is conformist learning in which individuals
disproportionally adopt the behaviour performed by the majority of their social connections (Pike and
Laland 2010; Van de Waal et al. 2013; Aplin et al. 2015; Danchin et al. 2018). Further, individuals often
only learn from specific individuals (e.g. depending on status (Canteloup et al. 2020), relatedness (Wild
et al. 2019) or conspecifics (Farine et al. 2015)) or adopt behaviours only once the social connections
to informed others surpass a certain threshold (Rosenthal et al. 2015).

To date, research on social learning in animals usually assumes that more social individuals (i.e. with
more social connections and central network positions) have a higher probability to adopt new
behaviours because they are more likely to hold connections to knowledgeable others compared to
less social individuals (Aplin et al. 2012; Claidiere et al. 2013; Kulahci and Quinn 2019). This link
between individual sociality and behavioural adoption can be expected if the learning rule depends
on the sum and strength of social connections to knowledgeable others. However, this relationship
may change when learning rules rely on both the connections to informed and uninformed individuals
(Firth 2020; Firth et al. 2020). For instance, in the case of conformist learning, we may expect that the
most social individuals will be less likely to adopt (because it may take longer until the majority of their
social connections becomes informed). Hence, predictions of how individual sociality relates to the
probability of acquiring novel behaviour, and the resulting transmission pathways, can change
fundamentally depending on the social learning mechanism at play.

Examining and comparing the transmission pathways of behaviours that follow different learning
mechanisms in the wild is challenging, if not impossible. Therefore, research investigating the
relationship between social structure and information flow often simulates behavioural spread (Voelkl
and Noë 2008; Nunn et al. 2009; Cantor et al. 2021; Evans et al. 2021). However, only a handful of
studies have considered the impact of different social learning rules (Nunn et al. 2009; Cantor et al.
2021). In addition, behavioural simulations are often performed on artificial social networks with pre-
defined structure and size (Voelkl and Noë 2008; Nunn et al. 2009; Cantor et al. 2021; Evans et al.
2021; but see: Naug 2008; Romano et al. 2018), and may thus represent unrealistic social structures,
failing to capture the social behaviour observed in real animal social networks. Further, for behavioural
spread, an individual's social connections at a relatively small temporal and spatial scale will predict
subsequent transmission (Somveille et al. 2018). Thus, it is important to study real animal social
networks on a meaningful spatiotemporal scale to better understand whether and how different types
of behaviours spread through social networks.

In this study, we explore by simulation how novel behaviours, transmitted according to different social
learning mechanisms, spread through replicated empirical social networks of great tits (*Parus major*).

Great tits are small songbirds that forage in mixed-species flocks during winter (Ekman 1989) and
frequently use social information (e.g. to find novel food (Aplin et al. 2012; Firth et al. 2016), to access
novel food (Aplin et al. 2015), prey avoidance (Thorogood et al. 2018; Hämäläinen et al. 2020)) which
makes them an ideal study species. Here, we create weekly social networks from the birds' foraging
associations at distinct locations to capture the social structure at a relatively small spatiotemporal
scale. Subsequently, we simulate behavioural spread on these weekly, local, networks using four
different social learning mechanisms. The first learning mechanism follows the omnipresent concept
which assumes that the probability of adopting depends on the number and strength of connections
to informed individuals (i.e. simple adoption rule). The other three learning mechanisms imply more
complex adoption rules: (i) a threshold rule, (ii) a proportion rule, and (iii) a conformity adoption rule.
Here, the probability of adopting the novel behaviour depends on: (i) the connections to informed
individuals surpassing a given threshold; (ii) the proportion of connections to informed individuals
(rather than the sum); and (iii), the behaviour that the majority of connections performs. We then
investigate the transmission pathways by comparing how the social behaviour of individual great tits
relates to the order in which they acquire novel behaviour under the four different social learning
rules.

Individual variation in sociality - the number of social connections and centrality within the network -
may influence the access to information and thus behavioural adoption. We infer individuals' sociality
by extracting three commonly used weighted social network metrics: the weighted degree (i.e. sum
of their social connections to others), weighted clustering coefficient (propensity for their associates
to be associated with one-another), and weighted betweenness (propensity to act as a 'bridge' within
the network). We predicted that the relationship between individual sociality and behavioural
adoption would differ depending on the social learning mechanism. Specifically, if the likelihood to
adopt a behaviour depends on the number and strength of connections to informed individuals such
as in the case for the simple and threshold rule, we predicted that more social individuals (i.e. high
degree and betweenness and low clustering coefficient) should be faster in adopting the novel
behaviour due to being more likely to be connected to at least one informed conspecific. In contrast,
if the likelihood of adopting a behaviour depends on the ratio of an individual's informed and
uninformed connections, such as in the case for the proportion or conformity rule, we expected that
less social individuals (i.e. low degree and betweenness and high clustering coefficient) should be
faster in adopting the novel behaviour because the majority of their social connections should become
informed faster.

133
134
Figure 1. Overview of the simulation procedure. First, a weekly social network of one of the 65
locations (shown as black dots) in the study site was selected. Second, behavioural spread was
simulated on the selected network using four different social learning rules (i.e. simple, threshold,
proportion, conformity). The starting point (i.e. the first individual performing the new behaviour)
was randomly chosen. Then, at each timestep (t1-tn), a naïve individual became knowledgeable with
a given probability of the adoption event being from social learning (dependent on the social
learning rule at play; see methods for further details) until all individuals in the network were
knowledgeable. Third, we calculated a correlation coefficient (Spearman's rank correlation
coefficient) between three individual social network metrics (i.e. weighted clustering coefficient,
weighted degree and weighted betweenness) and the order (i.e. timestep) in which individuals
acquired the novel behaviour. Finally, we repeated this process 100 times for each of our 1345
weekly, local social networks.

147
148

Results

The data in this study were simulated across 1345 empirically-derived social networks, inferred from
recordings of 1764 individual great tits at 62 feeder locations across 39 weekends and 3 years. Social
networks varied in size (right skewed distribution towards smaller networks, see Figure S1) and
consisted of an average of 21.7 individuals (min:10, max:77, sd=10.0).

Relationship between individual social behaviour and order of acquisition

Simulating behavioural spread under four different social learning rules revealed different
transmission pathways across social networks. For both the simple and the threshold rule, the
weighted clustering coefficient was positively related to the order of acquisition, with more clustered

individuals adopting the seeded behaviour later than less clustered individuals (Figure 2). Weighted
degree and betweenness were negatively related to the order of acquisition. Thus, individuals with
higher weighted degree and weighted betweenness adopted the novel behaviour faster than
individuals with a lower weighted degree and betweenness (Figure 2).

In addition, we show the relationship between mean individual network metrics and the standardized
order of acquisition across all network sizes (Figure S2). For the simple learning rule, the relationship
between network metric and order of acquisition only changed when the majority of individuals in a
network had already adopted the behaviour (with approximately 75% of individuals knowledgeable;
Figure S2). Behavioural spread under the threshold rule showed a small 'hump' shortly after the start
of behavioural spread (Figure 2, S2), before the relationship between mean network metric and order
of acquisition increased (for weighted clustering coefficient) or decreased (for weighted degree and
betweenness, Figure 2, S2). This 'hump' indicates that initially more social individuals (i.e. lower
weighted clustering coefficient, higher weighted degree and betweenness) adopted the behaviour
first, before the relationship between mean network metric and order of acquisition turned (Figure 2,
S2). In contrast to the simple and threshold rules, there was little or no relationship between individual
social network metrics and the order of acquisition under both the proportion and conformity learning
rules (Figure 2, S2).

Figure 2. Relationship between individual network metric and the order of acquisition for each social learning rule. Each column shows a different network metric (left to right: weighted clustering coefficient, weighted degree and weighted betweenness). Each row represents one of the four spreading rules (top to bottom: simple, threshold, proportion and conformity). Lines plot the average network metric for each order of acquisition and network size from the 100 simulations of each network. Colour represents network size with darker colour indicating smaller networks.

Social network size and behavioural spreading

Assessing the relationship between each individual's network metric and the order of acquisition across all social networks and simulation runs, revealed substantial variation in relationship strength (Figure 3). Across network metrics, there were on average the strongest correlations under the simple

rule (Figure 3), and coefficients for the threshold, proportion and conformity model were lower. The mean average network metric for later adopters (i.e. larger values for the order of acquisition) increased (for weighted clustering coefficient) and decreased (for weighted degree and betweenness) with increasing network size under the simple and threshold rule (Figure 2). The correlation between individual sociality and their order of acquisition were predicted by network size (Figure 4, Table S1). For the simple and threshold model, behavioural spread on larger networks led to more positive (for weighted clustering coefficient) and negative correlations (for weighted degree and betweenness) between individual network metric and order of acquisition (Figure 4, Table S1). The predicted effects of network size on mean correlation coefficient inferred under the proportion and conformity rule suggest contrasting directions (Figure 4, Table S1).

Social spread under the threshold rule may be limited in smaller networks (see also section on *'Probability of social* spread') causing the reduced correlations between individual network metrics and order of acquisition. Repeating the simulations under the threshold rule, but with a threshold location proportional to network size (in contrast to a fixed threshold location across all networks, see methods), revealed no strong change in the mean individual network metric and order of acquisition with increasing network size (Figure S3). Similar patterns were detected when repeating simulations under the threshold model, with a threshold location smaller (a=3, Figure S4) and larger (a=7, Figure S4) than the location set for the main analyses. When using a smaller threshold, the negative relationship between mean network metric and order of acquisition was already evident in smaller networks (Figure S4). However, when setting a larger threshold, the relationship between network metric and order of acquisition did not reveal a distinct direction (Figure S4). Further, for the threshold rule, the limited ability of social spread and the prominent 'hump' at the initial stages of the spread, may have led to overall low correlation coefficients between individual network metrics and order of acquisition (Figure 3), even though the relationship between mean network metric and order of acquisition (Figure 2) indicated otherwise.

Figure 3. **Distribution of average correlation coefficients for each social learning rule and network metric.** Violin and boxplots show the distribution of the average correlation coefficients across 100 simulations from each network for each of the four social learning rules (i.e. simple, proportion, conformity and threshold). Each plot shows one of the individual network metrics (weighted clustering coefficient, weighted degree, weighted betweenness).

Figure 4. **Relationship between correlation coefficient and network size across the four social learning rules.** Each plot shows one of the individual network metrics. Average correlation coefficients across each 100 simulations per network are plotted as dots, lines represent the effects generated from linear mixed-effect models (LMM) and ribbon represents the 95% CI (see Table S1 for model results).

Probability of social spread

For each spreading simulation, an initial random individual was seeded with the behaviour, and then at each timestep a new individual adopted the seeded behaviour, whereby each time each individual has a probability of adopting the behaviour through either asocial learning (which is constant across all individuals with no connections to informed individuals) or through social learning which ranges from 0 (asocial acquisition only, i.e. an individual has no connections to informed individuals and cannot socially learn the behaviour under the set learning rule) to 1 (obligate social learning, i.e. when an individual is well connected to informed individuals so the individual will only learn under the set learning rule).

Across the different social learning mechanisms, the average probability that an individual socially adopted the seeded behaviour increased with increasing timestep (i.e. order of acquisition). This is because with an increasing number of individuals becoming knowledgeable, the probability to be connected to informed others and thus to reach the set learning criterion increased (Figure 5). However, under certain spreading processes, social learning was limited (Figure 5). This was particularly the case for smaller networks when transmission followed the threshold rule. For instance, in networks consisting of less than approximately 20 individuals, the set threshold for socially adopting the behaviour was almost never met (average probability of social learning is close to 0; Figure 5 Threshold). Repeating the simulations under the threshold rule, but with a threshold location proportional to network size, removed these patterns (Figure S5). For the conformity rule, larger networks limited social spread, but only at the initial phase of transmission (Figure 5 Conformity). In larger social networks, individuals have on average more social connections and thus at the initial stages of behavioural spread, it is less likely that the majority of an individual's social connections are already informed.

Behaviours spreading under the simple and proportion rule, on average always had a non-zero probability to socially spread (Figure 5). This is the case because both the simple and proportion rule, did not require to surpass a given threshold (such as set for the threshold rule or reaching the majority

of an individual's connections as set for the conformity rule). Therefore, one connection to an
informed conspecific was sufficient for the seeded behaviour to spread socially.

For both the simple and threshold model, the mean probability of social spread initially increased but
then dropped towards the later adoptions (Figure 5). This may be caused by a few individuals that are
not well connected in the social network, and thus have in general a low probability to socially learn
under the simple and threshold rule. For instance, some great tits may only be connected to one other
individual and thus, even if the number of knowledgeable individuals in the social network increases
over time, it is very unlikely to reach the required threshold for social learning. For the proportion and
conformity rule, the average probability of social learning peaked after a given percentage of
individuals being knowledgeable (Figure 5).

Figure 5. Relationship between the probability of an individual socially adopting the seeded
behaviour and the number of informed individuals within the network. Panels show the results for
each of the four social learning rules (simple, threshold, proportion, conformity). The x-axis describes
the number of informed individuals within a social network. The simulations are set so that at each
timestep one individual changes its status from 'uninformed' to 'informed'. At each timestep a new
individual adopted the seeded behaviour, whereby each time each individual has a probability of
adopting the behaviour through social learning (y-axis) given the set learning rule. Lines plot the
average probability for each timestep and network size from the 100 simulations. Colour represents
network size with darker colour indicating smaller networks.

Discussion

We show how the underlying social learning rules individuals employ strongly influences the transmission pathways of behaviours across social networks derived from real-world data. Under learning rules that rely purely on the extent of social connections to informed others, we found that more social individual great tits were likely to adopt the seeded behaviour faster, in line with common expectations of the benefits of sociality for gaining information. However, if the likelihood of adopting a behaviour depended on the ratio of connections to informed and uninformed others, such as conformist learning, social network centrality was not strongly related to the order in which individuals acquired the seeded behaviour. Notably, this contrasts with the widely proposed prediction that more social individuals may be more likely to adopt new information. Thus, our results show how the relationship between individual sociality and information acquisition can change fundamentally with the type of social learning mechanism at play. Finally, we reveal that the probability of social spread under certain social learning rules is predicted to be limited in certain real-world settings, particularly in networks of a very small or large size.

Individuals differ in the quantity and quality of social connections to others which impacts several aspects of life history (McDonald 2007; Formica et al. 2012; MacIntosh et al. 2012; Farine and Sheldon 2015; Alberts 2019; Beck et al. 2021), including the social transmission of information (Aplin et al. 2012; Kulahci and Quinn 2019). Generally, individuals that hold many social connections to others, and occupy central network positions, are expected to be more likely to acquire information (Hoppitt and Laland 2011; Aplin et al. 2012; Claidiere et al. 2013; Kulahci et al. 2016). In line with these assumptions, we find that more social great tits (i.e. with a lower clustering coefficient, higher degree and betweenness) were more likely to adopt the seeded behaviour faster when the underlying spreading mechanism depended on the extent of connections to knowledgeable others (Figure 2). However, when rules were more complex, for example when behavioural adoption depended on the ratio of connections to knowledgeable and naïve individuals, sociality was not strongly related to the order of acquisition (Figure 2), matching the expectations from studies on complex contagions in humans (Guilbeault et al. 2018). These findings suggest that under certain spreading mechanisms, the extent of social connections can reduce social spread. Thus, the concept that more social and central individuals are especially important in acquiring and subsequently spreading information (Kulahci and Quinn 2019) cannot be generalized across different types of behaviours, and may even lead to erroneous conclusions. For instance, if no evidence is found that individual sociality is related to the probability of behavioural adoption, one might be led to conclude the absence of social learning where, in fact more complex social learning rules may be in operation. Therefore, our findings have important consequences for our understanding of the relationship between sociality and behavioural acquisition, and also for what may constitute an 'optimal' social structure for efficient social transmission (Pasquaretta et al. 2014; Romano et al. 2018).

Furthermore, our results have implications for our understanding of the costs and benefits of individual sociality. While increased access to information is one of the postulated key benefits of sociality, simply holding more connections to others may in fact hinder the adoption of novel behaviours under certain social learning rules. For instance, if a novel behaviour follows a conformist learning mechanism, highly social individuals with lots of connections may be exposed to the new behaviour sooner than less social individuals because they are more likely to be connected to at least

one informed conspecific, but despite this will be less likely to adopt the novel behaviour if it requires
the majority of their social connections to become informed first. Hence, considering the social
learning mechanism at play, and differentiating between access to information and behavioural
adoption (Carter et al. 2016), are crucial for our understanding of the potential costs and benefits of
individual sociality.

Examining behavioural innovation and its subsequent spread in nature is challenging (Whiten and
Mesoudi 2008; Klump et al. 2021). This is because new behaviours are often only detected once the
majority of individuals in a population are already knowledgeable. Alternatively, behavioural
innovations may occur much more frequently but remain undetected if the behaviour does not spread
far (e.g. because the behaviour is mechanically challenging for individuals (Gajdon et al. 2006) or when
the carryover of older, outdated behaviour hinders the spread of novel and more adaptive behaviours
(Aplin et al. 2017; Barrett et al. 2019)). Here we used simulations on real-world empirical networks,
and demonstrate that the ability of a behaviour to socially spread depends on its underlying social
learning mechanism, which is consistent with other studies that investigated different spreading
processes (Nunn et al. 2009; Cantor et al. 2021). Across the four models, the probability that individual
great tits socially adopted the seeded behaviour increased with increasing timestep (Figure 5). This is
expected as at each timestep in simulations a new individual becomes knowledgeable and remains in
this state thereby increasing the number of knowledgeable individuals within the network. However,
social transmission under certain social learning mechanisms was limited. For instance, behaviours
that needed to surpass a given threshold of social connections to informed others (i.e. threshold or
conformity rule) required more asocial learning events in the initial spreading phase to be able to
subsequently transmit via social learning (Figure 5). In contrast, for the simple and proportion rules,
social transmission was always possible, even in the initial spreading phase (Figure 5). This
demonstrates that some behaviours may have a much higher likelihood to spread socially, whereas
other behaviours, following more complex processes, may rely on a larger extent of asocial learning
events or social reinforcement. Therefore, the majority of empirical research on behavioural spread
in animals may in fact examine more simple social transmission processes (because they are easier to
detect and observe) which may bias our picture of the existing social learning mechanisms of animals
and behavioural innovations.

We found that network size impacted behavioural spread across the four transmission models. The
strength of the relationship between individual network metric and the order of acquisition increased
with increasing network size under the simple and threshold rule (Figure 4). The weaker correlation
between individual sociality and order of acquisition in smaller networks might be caused by the
relatively reduced likelihood of social spread (Figure 5). For instance, for the threshold rule, individuals
in larger networks have an overall higher number of social connections (compared to individuals in
smaller networks) which facilitates reaching the set threshold for social learning and thus increases
the likelihood of social spread. This is supported by our results when simulating spread using different
threshold values (see Figure S4). Past studies have investigated behavioural spread on relatively large
networks (using simulations: (Voelkl and Noë 2008; Nunn et al. 2009; Cantor et al. 2021; Evans et al.
2021) or natural observations: (Allen et al. 2013; Aplin et al. 2015)). However, for behavioural spread,
social connections at a relatively small spatiotemporal scale may predict an individual's decision to
adopt a novel behaviour. Our findings that network size can impact behavioural spread thus have
important consequences for our understanding of the influence of wider society structure on when
and where behaviours may emerge, and how to interpret empirical results. As such, it is important to

improve our understanding of the factors that give rise to different social network sizes and shapes
and subsequently influence information flow. For instance, spatiotemporal variation in environmental
features (such as the availability and distribution of resources) may influence population densities and
subsequently local social network size, and as such facilitating or hindering social spread.

Our study suggests further questions for future research. First, our models assume that each individual
adopts a seeded behaviour under the same 'learning rule'. However, individuals may differ in the
extent of social information use, which is often context and state-dependent (Rendell et al. 2011;
Penndorf and Aplin 2020). For instance, dominance rank (Krueger et al. 2014) and sex (Aplin et al.
2013) have been shown to be related to variation in social information use. Further, individuals may
only copy behaviour from certain individuals, based on familiarity or kin (Kavaliers et al. 2005; Boogert
et al. 2018). Thus, future simulation or empirical studies could explore how heterogeneity in learning
rules impacts information flow. Second, using simulations did not allow us to test changes in the social
network resulting from behavioural adoption: individuals may occupy more central social network
positions once performing a new behaviour (Kulahci et al. 2018; Kulahci and Quinn 2019). This can be
the case if individuals preferentially associate with knowledgeable others (Kulahci et al. 2018), or if
individuals change their behaviour in response to information acquisition which can also lead to an
increased in social connections (Kulahci and Quinn 2019). Such dynamics are not reflected in our study,
and would require the investigation of natural behavioural spread ideally under experimental
conditions in the wild. Third, our models assume that once individuals become 'informed', they cannot
return to an 'uninformed' state. In natural conditions, however, individuals may return again to an
'uninformed' state if the novel behaviour was not rewarding. Future studies incorporating these
aspects could provide further new insights into the patterns of social transmission and its link to
sociality.

From copying the majority (Van de Waal et al. 2013; Aplin et al. 2015; Danchin et al. 2018) to learning
from specific tutors (Wild et al. 2019; Canteloup et al. 2020), individuals use a large range of different
learning strategies (Hoppitt and Laland 2013; Kendal et al. 2018). While, an increasing number of
studies show how social network structure can influence social transmission (using simulations:
(Voelkl and Noë 2008; Romano et al. 2018; Evans et al. 2021) and empirical data: (Naug 2008; Firth et
al. 2016; Romano et al. 2018)), our research highlights the importance of the underlying social learning
mechanism in shaping the transmission pathways across social networks. We demonstrate that the
common assumption that sociality is linked to a higher likelihood in acquiring information and
adopting new behaviours cannot be generalized for behavioural spread in real world networks. This
also sheds new light on our current understanding of the costs and benefits of individuals sociality and
asks to focus more on the social learning mechanism at play, and to differentiate between access to
information and behavioural adoption. In addition, we reveal that social transmission can be limited
under certain adoption rules (such as the threshold rule), and social networks of particular size. Our
findings thus have important consequences for our understanding of whether and how behaviours
spread across different social networks, and subsequently the establishment of traditions and
cultures. Further, differences in spreading mechanisms alter predictions of what may constitute
optimal social structures for the transmission of information, and how selection may act on sociality.

Materials and methods

Study system

The empirical data used in this study were collected over three years (December 2011 – March 2014) in a population of great tits located in Wytham Woods, Oxfordshire, UK (51° 46' N, 01° 20' W, approx. 385ha). Great tits are hole-nesting songbirds that form socially monogamous pairs, and establish territories during the breeding season (March – June). During the non-breeding season (September – February), great tits forage with other species in loose foraging flocks (Hinde 1952; Ekman 1989) and frequently use social information (Aplin et al. 2012; Farine et al. 2015; Firth et al. 2016).

The woodland contains 1017 nest-boxes hosting breeding great tits and 65 bird feeders that were deployed during the winter months in an evenly spaced grid (see Figure 1). Each feeder contained two access holes of which both were equipped with radio-frequency identification (RFID) antennas. For the duration of the study, the feeders collected data on the bird visits from pre-dawn Saturday morning until after dusk on Sunday evening. In each of the three winters, the feeders were in place from December to February and the location of each feeder was consistent.

All birds were caught in either a nest-box or a mist-net and were fitted with a uniquely numbered metal leg ring (British Trust for Ornithology). In addition, each bird was also fitted with a uniquely coded passive integrated transponder (PIT) tag enclosed in a plastic ring fitted to the other leg. This allowed us to record each visit of a PIT-tagged bird when it came close to the RFID antenna of a feeder (approximately 3 cm). At every detection, the bird's unique PIT tag code, and the date and time were saved to a data logger.

Social networks

We created social networks based on the foraging associations of PIT-tagged great tits at each feeder and each of the 13 weekends across the three years. We created temporal and locally restricted networks because we wanted to generate a large number of different social networks (rather than just one network from the whole population) and because we expect networks from such a small time-window (i.e. one weekend) to be most meaningful in capturing the social connections relevant for the spread of a novel behaviour (i.e. social networks spanning a large time-window may contain many connections not relevant at the time a novel behaviour was introduced).

An "association" was defined as two birds foraging together within the same flock. Flock membership was identified using Gaussian Mixture Models (Psorakis et al. 2012; Psorakis et al. 2015) from the R package "asnipe" (Farine 2013). This method detects events of increased feeding activity in the spatio-temporal data, clusters these into non-overlapping gathering events (i.e. flocking events), and assigns each individual detection to the event it most likely belonged to. This provided us with information about which individuals co-occurred in the same flock (Psorakis et al. 2012; Psorakis et al. 2015). From the pattern of co-occurrences, we then inferred the strength of associations for each dyad. We calculated association strength using the simple ratio index (SRI, (Cairns and Schwager 1987)). The SRI describes the proportion of observations of two individuals in which they were seen together, ranging from 0 (never observed in the same flock) to 1 (always observed in the same flock). We created undirected social networks with edges weighted by the SRI for each sampling weekend (in total: 39 weekends across 3 years) and feeder location (in total: 65).

For each of the weekly, local networks, we then inferred for each individual three social network
metrics: the weighted clustering coefficient (following (Barrat et al. 2004)), the weighted degree and
the weighted betweenness. Network metrics were calculated using the R package 'igraph' (Csardi and
Nepusz 2006) and describe respectively: how tightly a focal individual is clustered to form distinct
cliques, the total interaction rate for a foal individual with all other individuals, and the number of
shortest paths from all individuals to all other individuals that pass through the focal individual. Last,
we standardized the individual metrics within each network to allow comparisons between networks.
Social networks including less than ten individuals and exhibiting no variation in individual network
metrics were excluded from further analyses resulting in the final sample size of 1345 social networks
(generated from 62 locations).
476

Simulations

On each network, we simulated behavioural spread using four different social learning rules, as
described in (Firth et al. 2020), using the R package 'complexNBDA'. A brief explanation is provided
below but full detailed description of each can be found in Firth et al. 2020 and all resources are freely
available at https://github.com/whoppitt/complexNBDA.

(1) Simple rule. This transmission rule follows the logic of the classic NBDA framework. Here, the
probability that an uninformed individual adopts a new behaviour is directly dependent on the
number and strength of connections to informed others that are currently displaying the behaviour.
Consequently, the more connections to informed individuals, the more likely an individual is to adopt
the behaviour. The social transmission strength s was set to 1.

(2) Threshold rule: This transmission rule assumes that the probability of adopting a new behaviour
remains low until an individual's social connections to informed individuals reaches a given threshold.
We set the threshold value a to 5 for all networks. We set the sharpness of the threshold effect, b, to
3 (reflecting a sharp threshold). The social transmission strength s was set to 30. In addition, we
repeated simulations by setting a threshold in relation to the network size, a threshold of 3 and a
threshold of 7 (see supplementary material).

(3) Proportion rule: This transmission rule defines the learning rate to be proportional to the ratio of
connections that an individual holds to informed others. As such, the individual with the highest
proportion is most likely to learn and assumes additional influence from individuals' uninformed
connections rather than just considering the sum of connections to informed individuals. The social
transmission strength s was set to 30.

(4) Conformity rule: Finally, the fourth transmission rule assumes that individuals are
disproportionately more likely to copy the majority of the population (i.e. frequency dependent).
Thus, an individual is expected to adopt a new behaviour if it is perceived as being performed by the
majority of its' social connections. Similar to the proportional rule, the conformity rule considers an
individual's informed and uninformed connections. The value f describes the probability of adopting
the new behaviour, whereby $f>1$ means that the individual adopts the behaviour performed by the
majority. For our simulations, we set f to 3 and the social transmission strength s to 30.

For each simulation the individual 'initiating' the behaviour (i.e. the demonstrator) was randomly
chosen but was consistent across the four transmission models. We simulated behavioural spread

across the entire network. At each timestep within each simulation, a new individual was chosen to acquire the behaviour, which is dependent on the likelihoods of acquisition assigned to each uninformed individual at that timestep based on the probability of any individual 'asocially' acquiring the behaviour, and the probability of each individual socially acquiring the behaviour which is derived from their social connections' status (informed or uninformed) and the learning rule at play. Full details for this method, and associated tests, are available in the complexNBDA package (Firth et al. 2020) and follows the principles of NBDA transmissions (Hasenjager et al. 2021). Once individuals adopted the novel behaviour, they remained 'informed' within each simulation run. For each network, we repeated the simulations 100 times to minimize the influence of the identity of the randomly selected demonstrator on the subsequent transmission pathways. We selected 100 simulation runs because this was enough in acquiring a relatively stable mean correlation coefficient between network metric and order of acquisition (Figure S6). As such, as total of 538000 simulations (1345 weekly networks x 4 different learning rules x 100 simulation runs) were carried out across all the variations of the reported analyses.

Data summary statistics

To assess the relationship between each individual's network metric and the order of acquisition, we calculated Spearman's rank correlation coefficients. After every simulation, for each of the networks and transmission models, we calculated the Spearman's rank correlation coefficient between the order in which individuals adopted the behaviour (always excluding the demonstrator) and each of the three individual network metrics (i.e. the weighted clustering coefficient, the weighted degree and the weighted betweenness). For an overview of the transmission process, see Figure 1. For each network and model, we then calculated the average correlation coefficient for each network metric across the 100 simulations (Figure 3). To additionally assess the relationships between the size of the network, and the correlation coefficient between individuals' centralities and their acquisition order, we used linear mixed-effect models (LMM). We set the average correlation coefficient as the dependent variable and network size in interaction with the learning rule as the predictor variable. Location identity and week nested in year were set as random effects to factor in these differences when assessing this relationship (see Table S1 for results).

Acknowledgments

We thank the large number of contributors to the data collection. The study has been supported by grants from NERC (NE/S010335/1) and BBSRC (BB/S009752/1).

References

Alberts SC. 2019. Social influences on survival and reproduction: Insights from a long-term study of wild baboons. J Anim Ecol. 88(1):47–66.

Allen J, Weinrich M, Hoppitt W, Rendell L. 2013. Network-based diffusion analysis reveals cultural
transmission of lobtail feeding in humpback whales. Science (80-). 340(6131):485–488.

Aplin LM, Farine DR, Morand-Ferron J, Cockburn A, Thornton A, Sheldon BC. 2015. Experimentally
induced innovations lead to persistent culture via conformity in wild birds. Nature. 518(7540):538–
541.

Aplin LM, Farine DR, Morand-Ferron J, Sheldon BC. 2012. Social networks predict patch discovery in
a wild population of songbirds. Proc R Soc London B Biol Sci. 279(1745):4199–4205.

Aplin LM, Sheldon BC, McElreath R. 2017. Conformity does not perpetuate suboptimal traditions in a
wild population of songbirds. Proc Natl Acad Sci. 114(30):7830–7837.

Aplin LM, Sheldon BC, Morand-Ferron J. 2013. Milk bottles revisited: social learning and individual
variation in the blue tit, Cyanistes caeruleus. Anim Behav. 85(6):1225–1232.

Barrat A, Barthelemy M, Pastor-Satorras R, Vespignani A. 2004. The architecture of complex
weighted networks. Proc Natl Acad Sci. 101(11):3747–3752.

Barrett B, Zepeda E, Pollack L, Munson A, Sih A. 2019. Counter-culture: Does social learning help or
hinder adaptive response to human-induced rapid environmental change? Front Ecol Evol. 7:183.

Beck KB, Farine DR, Kempenaers B. 2021. Social network position predicts male mating success in a
small passerine. Behav Ecol. 32(5):856–864. doi:10.1093/beheco/arab034.
https://doi.org/10.1093/beheco/arab034.

Boogert NJ, Lachlan RF, Spencer KA, Templeton CN, Farine DR. 2018. Stress hormones, social
associations and song learning in zebra finches. Philos Trans R Soc B Biol Sci. 373(1756):20170290.

Cairns SJ, Schwager SJ. 1987. A comparison of association indices. Anim Behav. 35(5):1454–1469.

Canteloup C, Hoppitt W, van de Waal E. 2020. Wild primates copy higher-ranked individuals in a
social transmission experiment. Nat Commun. 11(1):1–10.

Cantor M, Chimento M, Smeele SQ, He P, Papageorgiou D, Aplin LM, Farine DR. 2021. Social network
architecture and the tempo of cumulative cultural evolution. Proc R Soc B. 288(1946):20203107.

Carter AJ, Tico MT, Cowlishaw G. 2016. Sequential phenotypic constraints on social information use
in wild baboons. Elife. 5:e13125.

Centola D, Macy M. 2007. Complex contagions and the weakness of long ties. Am J Sociol.
113(3):702–734.

Claidiere N, Messer EJE, Hoppitt W, Whiten A. 2013. Diffusion dynamics of socially learned foraging
techniques in squirrel monkeys. Curr Biol. 23(13):1251–1255.

Csardi G, Nepusz T. 2006. The igraph software package for complex network research, InterJournal,
Complex Systems 1695. 2006. http://igraph.org.

Danchin E, Nöbel S, Pocheville A, Dagaeff A-C, Demay L, Alphand M, Ranty-Roby S, Van Renssen L,
Monier M, Gazagne E. 2018. Cultural flies: Conformist social learning in fruitflies predicts long-lasting
mate-choice traditions. Science (80-). 362(6418):1025–1030.

Ekman J. 1989. Ecology of non-breeding social systems of Parus. Wilson Bull.:263–288.

Evans JC, Hodgson DJ, Boogert NJ, Silk MJ. 2021. Group size and modularity interact to shape the
spread of infection and information through animal societies. Behav Ecol Sociobiol. 75(12):1–14.

Farine DR. 2013. Animal social network inference and permutations for ecologists in R using asnipe.

Methods Ecol Evol. 4(12):1187–1194.

Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015. Interspecific social networks promote
information transmission in wild songbirds. Proc R Soc B Biol Sci. 282(1803):20142804.

Farine DR, Sheldon BC. 2015. Selection for territory acquisition is modulated by social network
structure in a wild songbird. J Evol Biol. 28(3):547–556.

Firth JA. 2020. Considering complexity: animal social networks and behavioural contagions. Trends
Ecol Evol. 35(2):100–104.

Firth JA, Albery GF, Beck KB, Jarić I, Spurgin LG, Sheldon BC, Hoppitt W. 2020. Analysing the Social
Spread of Behaviour: Integrating Complex Contagions into Network Based Diffusions. arXiv Prepr
arXiv201208925.

Firth JA, Sheldon BC, Farine DR. 2016. Pathways of information transmission among wild songbirds
follow experimentally imposed changes in social foraging structure. Biol Lett. 12(6):20160144.

Formica VA, Wood CW, Larsen WB, Butterfield RE, Augat ME, Hougen HY, BRODIE III ED. 2012.
Fitness consequences of social network position in a wild population of forked fungus beetles
(Bolitotherus cornutus). J Evol Biol. 25(1):130–137.

Gajdon GK, Fijn N, Huber L. 2006. Limited spread of innovation in a wild parrot, the kea (Nestor
notabilis). Anim Cogn. 9(3):173–181.

Guilbeault D, Becker J, Centola D. 2018. Complex contagions: A decade in review. Complex spreading
Phenom Soc Syst.:3–25.

Hämäläinen L, Mappes J, Rowland HM, Teichmann M, Thorogood R. 2020. Social learning within and
across predator species reduces attacks on novel aposematic prey. J Anim Ecol. 89(5):1153–1164.

Hasenjager MJ, Leadbeater E, Hoppitt W. 2021. Detecting and quantifying social transmission using
network-based diffusion analysis. J Anim Ecol. 90(1):8–26.

Heyes CM. 1994. Social learning in animals: categories and mechanisms. Biol Rev. 69(2):207–231.

Hinde RA. 1952. The Behaviour of the Great Tit (parus Major) and Some Other Related Species. Brill.

Hoppitt W, Laland KN. 2011. Detecting social learning using networks: a users guide. Am J Primatol.
73(8):834–844.

Hoppitt W, Laland KN. 2013. Social learning. Princeton University Press.

Kavaliers M, Colwell DD, Choleris E. 2005. Kinship, familiarity and social status modulate social
learning about "micropredators"(biting flies) in deer mice. Behav Ecol Sociobiol. 58(1):60–71.

Kendal RL, Boogert NJ, Rendell L, Laland KN, Webster M, Jones PL. 2018. Social learning strategies:
Bridge-building between fields. Trends Cogn Sci. 22(7):651–665.

Klump BC, Martin JM, Wild S, Hoersch JK, Major RE, Aplin LM. 2021. Innovation and geographic
spread of a complex foraging culture in an urban parrot. Science.

Krueger K, Farmer K, Heinze J. 2014. The effects of age, rank and neophobia on social learning in
horses. Anim Cogn. 17(3):645–655.

Kulahci IG, Ghazanfar AA, Rubenstein DI. 2018. Knowledgeable lemurs become more central in social
networks. Curr Biol. 28(8):1306–1310.

Kulahci IG, Quinn JL. 2019. Dynamic relationships between information transmission and social
connections. Trends Ecol Evol. 34(6):545–554.

Kulahci IG, Rubenstein DI, Bugnyar T, Hoppitt W, Mikus N, Schwab C. 2016. Social networks predict
selective observation and information spread in ravens. R Soc open Sci. 3(7):160256.

MacIntosh AJJ, Jacobs A, Garcia C, Shimizu K, Mouri K, Huffman MA, Hernandez AD. 2012. Monkeys
in the middle: parasite transmission through the social network of a wild primate. PLoS One.
7(12):e51144.

McDonald DB. 2007. Predicting fate from early connectivity in a social network. Proc Natl Acad Sci.
104(26):10910–10914.

Naug D. 2008. Structure of the social network and its influence on transmission dynamics in a
honeybee colony. Behav Ecol Sociobiol. 62(11):1719–1725.

Nunn CL, Thrall PH, Bartz K, Dasgupta T, Boesch C. 2009. Do transmission mechanisms or social
systems drive cultural dynamics in socially structured populations? Anim Behav. 77(6):1515–1524.

Pasquaretta C, Levé M, Claidiere N, Van de Waal E, Whiten A, MacIntosh AJJ, Pelé M, Bergstrom ML,
Borgeaud C, Brosnan SF. 2014. Social networks in primates: smart and tolerant species have more
efficient networks. Sci Rep. 4(1):1–8.

Penndorf J, Aplin L. 2020. Environmental and life history factors, but not age, influence social
learning about food: a meta-analysis. Anim Behav. 167:161–176.

Pike TW, Laland KN. 2010. Conformist learning in nine-spined sticklebacks' foraging decisions. Biol
Lett. 6(4):466–468.

Psorakis I, Roberts SJ, Rezek I, Sheldon BC. 2012. Inferring social network structure in ecological
systems from spatio-temporal data streams. J R Soc Interface. 9(76):3055–3066.

Psorakis I, Voelkl B, Garroway CJ, Radersma R, Aplin LM, Crates RA, Culina A, Farine DR, Firth JA,
Hinde CA. 2015. Inferring social structure from temporal data. Behav Ecol Sociobiol. 69(5):857–866.

Rendell L, Fogarty L, Hoppitt WJE, Morgan TJH, Webster MM, Laland KN. 2011. Cognitive culture:
theoretical and empirical insights into social learning strategies. Trends Cogn Sci. 15(2):68–76.

Romano V, Shen M, Pansanel J, MacIntosh AJJ, Sueur C. 2018. Social transmission in networks: global
efficiency peaks with intermediate levels of modularity. Behav Ecol Sociobiol. 72(9):1–10.

Rosenthal SB, Twomey CR, Hartnett AT, Wu HS, Couzin ID. 2015. Revealing the hidden networks of
interaction in mobile animal groups allows prediction of complex behavioral contagion. Proc Natl
Acad Sci. 112(15):4690–4695.

Somveille M, Firth JA, Aplin LM, Farine DR, Sheldon BC, Thompson RN. 2018. Movement and
conformity interact to establish local behavioural traditions in animal populations. bioRxiv.:1–19.
doi:10.1101/338657.

Thorogood R, Kokko H, Mappes J. 2018. Social transmission of avoidance among predators facilitates
the spread of novel prey. Nat Ecol Evol. 2(2):254–261.

Voelkl B, Noë R. 2008. The influence of social structure on the propagation of social information in
artificial primate groups: a graph-based simulation approach. J Theor Biol. 252(1):77–86.

Van de Waal E, Borgeaud C, Whiten A. 2013. Potent social learning and conformity shape a wild
primate's foraging decisions. Science (80-). 340(6131):483–485.

Whiten A. 2021. The burgeoning reach of animal culture. Science (80-). 372(6537).

Whiten A, Mesoudi A. 2008. Establishing an experimental science of culture: animal social diffusion
experiments. Philos Trans R Soc B Biol Sci. 363(1509):3477–3488.

Wild S, Allen SJ, Krützen M, King SL, Gerber L, Hoppitt WJE. 2019. Multi-network-based diffusion
analysis reveals vertical cultural transmission of sponge tool use within dolphin matrilines. Biol Lett.
15(7):20190227.