Social learning mechanisms shape transmission pathways through replicate
local social networks of wild birds

Kristina B. Beck[1*], Ben C. Sheldon[1], Josh A. Firth[1]

[1]Edward Grey Institute of Field Ornithology, Department of Biology, University of Oxford, Oxford
OX1 3SZ, United Kingdom

*Correspondence:
kbbeck.mail@gmail.com

10
11

Abstract

The emergence and spread of novel behaviours via social learning can lead to rapid population-level
changes whereby the social connections between individuals shape information flow. However,
behaviours can spread via different mechanisms and little is known about how information flow
depends on the underlying learning rule individuals employ. Here, comparing four different learning
mechanisms, we simulated behavioural spread on replicate empirical social networks of wild great tits
and explored the relationship between individual sociality and the order of behavioural acquisition.
Our results reveal that, for learning rules dependent on the sum and strength of social connections to
informed individuals, social connectivity was related to the order of acquisition, with individuals with
increased social connectivity and reduced social clustering adopting new behaviours faster. However,
when behavioural adoption depends on the ratio of an individuals' social connections to informed
versus uninformed individuals, social connectivity was not related to the order of acquisition. Finally,
we show how specific learning mechanisms may limit behavioural spread within networks. These
findings have important implications for understanding whether and how behaviours are likely to
spread across social systems, the relationship between individuals' sociality and behavioural
acquisition, and therefore for the costs and benefits of sociality.

28
29

Introduction
Social learning, in which individuals learn from others, is widespread in the animal kingdom, and
enables individuals to acquire novel behaviours facilitating phenotypic change (Heyes 1994; Hoppitt
and Laland 2013; Whiten 2021). Socially-induced changes in behaviour can spread through a
population and social networks provide the pathways along which behaviour can spread (Hasenjager
et al. 2021). Research increasingly shows how the structure of social networks and individual sociality
can together influence information flow (Voelkl and Noë 2008; Aplin et al. 2012; Kulahci et al. 2016;
Romano et al. 2018; Evans et al. 2021). However, information can spread via various social learning
mechanisms (Nunn et al. 2009; Firth 2020; Cantor et al. 2021; Evans et al. 2021) and we know
surprisingly little about how the relationship between sociality and information flow depends on the
social learning mechanisms at play.

By definition, the social spread of behaviour to a focal individual requires contact with at least one
knowledgeable individual. Frequently, it is intuitively assumed that the extent (i.e. number and
duration) of social contacts to knowledgeable others predicts the likelihood of adoption (Coussi-Korbel

and Fragaszy 1995; Franz and Nunn 2009; Hasenjager et al. 2021). In this way, behavioural spread is
predicted to follow a similar pattern to the transmission of many diseases. However, in contrast to
disease spread, individuals may employ 'social learning rules' where they can actively shape how to
act on acquired novel information. For example, for more costly behaviours such as the usage of a
novel food type, an individual may only change its behaviour after the majority of its social contacts
consumes the novel food. Consequently, behavioural spread may require exposure to multiple
sources (rather than just one) and depends on the ratio of connections to both informed and
uninformed individuals (rather than just the connections to informed others) (Centola and Macy 2007;
Hodas and Lerman 2014; Guilbeault et al. 2018; Firth et al. 2020). Therefore, the type of behaviour
considered and its underlying learning rule can fundamentally influence whether and how behaviour
spreads through a social network (Centola and Macy 2007; Firth et al. 2020).

In sociology, research increasingly demonstrates that the spread of various behaviours, from
innovations, to health, and political movements (Guilbeault et al. 2018), follow diverse and more
complex learning rules compared to the assumptions of many disease models (Centola and Macy
2007). In contrast, research in animal systems has rarely explored how the diffusion dynamics of
behaviours may be altered by learning rules (but see: Nunn et al. 2009; Cantor et al. 2021; Evans et al.
2020; Evans et al. 2021), which is somewhat surprising given that previous studies have revealed
several social learning strategies in animals that suggest a range of different underlying social learning
mechanisms (Hoppitt and Laland 2013; Kendal et al. 2018). For instance, an increasingly reported
learning mechanism is conformist learning in which individuals disproportionally adopt the behaviour
performed by the majority of their social connections (e.g. stickelbacks: Pike and Laland 2010;
chimpanzees: Haun et al. 2012; vervet monkeys: Van de Waal et al. 2013; great tits: Aplin et al. 2015a;
fruit flies: Danchin et al. 2018). Further, individuals often only learn from specific individuals (e.g.
depending on status (Canteloup et al. 2020), relatedness (Wild et al. 2019) or conspecifics (Farine et
al. 2015)) or adopt behaviours only once the social connections to informed others surpass a certain
threshold (Rosenthal et al. 2015).

Research on social learning in animal social networks has frequently assumed that more social
individuals (i.e. with more social connections and central network positions) have a higher probability
to adopt new behaviours because they are more likely to hold connections to knowledgeable others
compared to less social individuals (Aplin et al. 2012; Claidiere et al. 2013; Kulahci and Quinn 2019).
This link between individual sociality and behavioural adoption can be expected if the learning rule
depends on the sum and strength of social connections to knowledgeable others. However, this
relationship may change when learning rules rely on both the connections to informed and
uninformed individuals (Centola and Macy 2007; Firth 2020). For instance, in the case of conformist
learning, we may expect that the most social individuals will be less likely to adopt (because it may
take longer until the majority of their social connections becomes informed). Such patterns have been
reported in humans, where highly connected individuals required stronger social signals in order to
act on information (Hodas and Lerman 2012; Hodas and Lerman 2014) and poorly connected
individuals may utilise information sooner (González-Avella et al. 2011). Hence, predictions of how
individual sociality relates to the probability of acquiring novel behaviour, and the resulting
transmission pathways, can change fundamentally depending on the social learning mechanism at
play (Centola and Macy 2007; Firth et al. 2020).

Examining and comparing the transmission pathways of behaviours that follow different learning
mechanisms in wild animals is challenging. Therefore, research investigating the relationship between

social structure and information flow often simulates behavioural spread (Voelkl and Noë 2008; Nunn
et al. 2009; Evans et al. 2020; Cantor et al. 2021; Evans et al. 2021). For instance, studies compared
the transmission speed (number or proportion of individuals informed at a given timestep) of simple
versus conformity learning (Evans et al. 2020; Evans et al. 2021) or 'prestige' (subordinates copy
dominants) versus conformity learning (Nunn et al. 2009). While these studies show that on the
population level, different learning mechanisms, together with the social network structure, can
fundamentally impact the diffusion dynamics (e.g. how quickly a behaviour can spread), we know little
on how learning mechanisms impact the relationship between individual sociality and the probability
of behavioural adoption.

In addition, behavioural simulations are often performed on artificial social networks with pre-defined
structure and size (Voelkl and Noë 2008; Nunn et al. 2009; Cantor et al. 2021; Evans et al. 2021), and
may thus represent unrealistic social structures, failing to capture the social behaviour observed in
real animal social networks (but see Naug 2008; Romano et al. 2018). Further, simulated social
networks are often relatively large including 100 or more individuals. However, for behavioural
spread, an individual's social connections at a relatively small temporal scale may predict subsequent
transmission (Aplin et al. 2015a; Aplin et al. 2015b; Somveille et al. 2018). Many animal species live in
non-stable social groups such as fission-fusion societies (e.g. various species of birds (Silk et al. 2014),
primates (Amici et al. 2008), and fish (Wilson et al. 2014; Papastamatiou et al. 2020)) where group
composition and size frequently change. As a result, social connections between individuals can
change over time, and social networks generated over weeks/months may overestimate the social
connections of an individual at the time a new behaviour emerges. Thus, in addition to pure
computational studies, it is important to examine real animal social networks on a meaningful
temporal scale (which will be study species dependent) to better understand whether and how
different types of behaviours spread through social networks and to test whether general findings
from pure computational studies can be replicated using real animal networks.

In this study, we explore by simulation how novel behaviours, transmitted according to different social
learning mechanisms, spread through replicated empirical social networks of great tits (*Parus major*).
Great tits are small songbirds that forage in fission-fusion mixed-species flocks during winter (Ekman
1989) and frequently use social information (e.g. to find novel food (Aplin et al. 2012; Firth et al. 2016),
to access novel food (Aplin et al. 2015a), and for prey avoidance (Thorogood et al. 2018; Hämäläinen
et al. 2020)) which makes them an ideal study species. Here, we create social networks from empirical
data on birds' foraging associations at distinct locations sampled on two days each week to capture
the social structure at a relatively small spatiotemporal scale. Subsequently, we simulate behavioural
spread on these weekly, local, networks using four different social learning mechanisms and compare
how the social behaviour of individual great tits relates to the order in which they acquire novel
behaviour under the four different mechanisms.

The first learning mechanism follows the omnipresent concept of simple contagion, which is mainly
inspired by models on disease spread and was first formulated in the field of sociology (Guilbeault et
al. 2018). Simple contagion assumes that the probability of adopting a novel behaviour depends on
the number and strength of connections to informed individuals (thereafter simple rule (Coussi-Korbel
and Fragaszy 1995; Franz and Nunn 2009; Hasenjager et al. 2021)). The other three learning
mechanisms imply more complex adoption rules (Centola and Macy 2007) where behavioural
adoption requires more social reinforcement: (i) a threshold rule, (ii) a proportion rule, and (iii) a

conformity adoption rule. Here, the probability of adopting the novel behaviour depends on: (i) the connections to informed individuals surpassing a given threshold; (ii) the proportion of connections to informed individuals (rather than the sum); and (iii), the behaviour that the majority of connections performs. Threshold-based learning rules have been studied frequently in sociology and network sciences (Granovetter 1978; Watts 2002; González-Avella et al. 2011), but have rarely been considered in animals (Rosenthal et al. 2015). In contrast, conformity learning, where individuals are disproportionally more likely to copy the behaviour performed by the majority, has received much attention both in humans (Boyd and Richerson 1988; Haun et al. 2012; Toyokawa and Gaissmaier 2022) and animals (Van de Waal et al. 2013; Aplin et al. 2015a; Danchin et al. 2018). The proportion rule assumes that the transmission rate is proportional to the ratio of informed and uninformed individuals (rather than disproportional as in the conformity rule) and has rarely been considered (Rosenthal et al. 2015; Centola 2018; Firth 2020).

Individual variation in sociality - the number and strength of social connections and centrality within the network - may influence the access to information and thus behavioural adoption. We infer individuals' sociality by extracting three commonly used weighted social network metrics: the weighted degree (i.e. sum and strength of their social connections to others), weighted clustering coefficient (propensity for their associates to be associated with one-another), and weighted betweenness (propensity to act as a 'bridge' within the network). We predicted that the relationship between individual sociality and behavioural adoption would differ depending on the social learning mechanism. Specifically, if the likelihood to adopt a behaviour depends on the number and strength of connections to informed individuals such as in the case for the simple and threshold rule, we predicted that individuals with high degree and betweenness and low clustering coefficient should be faster in adopting the novel behaviour due to being more likely to be connected to at least one informed conspecific. In contrast, if the likelihood of adopting a behaviour depends on the ratio of an individual's informed and uninformed connections, such as in the case for the proportion or conformity rule, we expected that individuals with low degree and betweenness and high clustering coefficient should be faster in adopting the novel behaviour because the majority of their social connections should become informed faster.

Materials and methods

Study system

The empirical data used in this study were collected over three years (December 2011 – March 2014) in a population of great tits located in Wytham Woods, Oxfordshire, UK (51° 46' N, 01° 20' W, approx. 385ha). Great tits are short-lived (mean lifespan of 1.9 years (Bulmer and Perrins 1973)) hole-nesting songbirds that form socially monogamous pairs, and establish territories during the breeding season (March – June). During the non-breeding season (September – February), great tits forage with other species in loose fission-fusion flocks that differ in size and composition (Hinde 1952; Ekman 1989) and consist of mainly unrelated individuals (annual population turnover of about 50% and less than 1.5% of social foraging associations are between first-order relatives (Firth and Sheldon 2016)). Great tits frequently use social information in foraging contexts (Aplin et al. 2012; Farine et al. 2015; Firth et al. 2016; Thorogood et al. 2018).

The woodland contains 1017 nest-boxes hosting breeding great tits and 65 bird feeders that were
deployed during the winter months in an evenly spaced grid (see Figure 1). Each feeder contained two
access holes of which both were equipped with radio-frequency identification (RFID) antennas. The
feeders were in place from December to February across three winters (2011-12, 2012-2013, 2013-
2014) and collected data on the bird visits 2 days each week (from pre-dawn Saturday morning until
after dusk on Sunday evening) resulting in 13 sampling periods each year. At other times feeders were
closed. For the duration of the study, the location of each feeder was consistent.

All birds were caught in either a nest-box or a mist-net and were fitted with a uniquely numbered
metal leg ring (British Trust for Ornithology). In addition, each bird was also fitted with a uniquely
coded passive integrated transponder (PIT) tag enclosed in a plastic ring fitted to the other leg. This
allowed us to record each visit of a PIT-tagged bird when it came close to the RFID antenna of a feeder
(approximately 3 cm). At every detection, the bird's unique PIT tag code, and the date and time were
saved to a data logger. Breeding surveys and frequent trapping allowed to fit almost all individuals
with metal rings and PIT tags (>90%, Aplin, Farine, et al. 2013)

Social networks

We created social networks based on the foraging associations of PIT-tagged great tits at each feeder
and each of the 13 weekends across the three years. We created temporal and locally restricted
networks because we wanted to generate a large number of different social networks (rather than
just one network from the whole population) and because we expect networks from such a small time-
window (i.e. one weekend) to be most meaningful in capturing the social connections relevant for the
spread of a novel behaviour. For instance, (Aplin et al. 2015b) showed that individuals
disproportionally copied the behaviour of the majority of individuals in the social group that preceded
the focal individual's first successful solve. Across the winter, individual great tits may move between
locations and new individuals arrive at different times to the study site. Therefore, generating a social
network spanning the whole study period will contain many connections not relevant at the time a
novel behaviour emerges. Further, when examining the relationship between individual sociality and
the probability of adoption, generating a social network from the whole population would add
considerable spatial noise. For instance, within a sub-population where a new behaviour emerges,
individuals with high connectivity may be faster in adopting the behaviour. However, if examined on
the population-level, such a relationship may be obscured by spatial effects, because an individual's
probability of behavioural adoption will be considerably predicted by its' spatial proximity to the
location of behavioural emergence. Further, creating social networks for each feeder location
provided a comparable spatial unit and did not require to draw arbitrary spatial boundaries across the
study site.

All analyses were conducted in R 4.0.5 (R Core Team 2020). An "association" was defined as two birds
foraging together within the same flock. Flock membership was identified using Gaussian Mixture
Models (Psorakis et al. 2012; Psorakis et al. 2015) from the R package "asnipe" (Farine 2013). This
method detects events of increased feeding activity in the spatio-temporal data, clusters these into
non-overlapping gathering events (i.e. flocking events), and assigns each individual detection to the
event it most likely belonged to. This provided us with information about which individuals co-
occurred in the same flock (Psorakis et al. 2012; Psorakis et al. 2015). From the pattern of co-

occurrences, we then inferred the strength of associations for each dyad. We calculated association strength using the simple ratio index (SRI, (Cairns and Schwager 1987)). The SRI describes the proportion of observations of two individuals in which they were seen together, ranging from 0 (never observed in the same flock) to 1 (always observed in the same flock). We inferred a proportional measure for the association strength between individuals (rather than just a measure for the total number of times two individual were observed together) because we have an unequal number of observations for each individual (Farine and Whitehead 2015; Hoppitt and Farine 2018). Further, the SRI provides a more representative measure for the social relationship between two individuals across multiple contexts (e.g. while not foraging at the feeder). We created undirected social networks with edges weighted by the SRI for each sampling weekend (in total: 39 weekends across 3 years) and feeder location (in total: 65).

For each of the weekly, local networks, we then inferred for each individual three social network metrics: the weighted clustering coefficient, the weighted degree and the weighted betweenness. All network metrics were calculated using the R package 'igraph' (Csardi and Nepusz 2006). The weighted clustering coefficient was calculated following (Barrat et al. 2004). It represents the proportion of the sum of edge weights of all direct connections of a focal individual i over the sum of weights of all connections of individual i that form a triangle (i.e. where two direct connections of individual i are themselves connected). The weighted degree describes the total interaction rate for a focal individual i with all other individuals, defined as the sum of all the focal individual's edge weights. The weighted betweenness describes the number of weighted, shortest paths from all individuals to all other individuals that pass through the focal individual i and measures an individuals' propensity to move between groups. Here weights were added by considering the inverse of an individuals' edge weights.

Finally, we standardized the individual metrics within each network to allow comparisons between networks. Social networks including fewer than ten individuals and exhibiting no variation in individual network metrics were excluded from further analyses resulting in the final sample size of 1343 social networks (generated from 62 locations). The three individual network metrics are moderately correlated, with weighted clustering coefficient being negatively correlated to weighted degree and weighted betweenness, weighted degree and weighted betweenness were positively correlated (see Table S1). In addition, we provide example networks with individual great tits colour-coded based on their different weighted network metrics (Figure S1) and calculated four global network measures (network density, average path length, average edge weight, modularity) to provide a general overview of the weekly, local great tit social structures (Figure S2, details on how these metrics were calculated can be found in the figure legend).

Simulations

On each network, we simulated behavioural spread using four different social learning rules, as described in (Firth et al. 2020), using the R package 'complexNBDA'. A brief explanation is provided below but full detailed description and tests of each can be found in Firth et al. (2020) and all resources are freely available at https://github.com/whoppitt/complexNBDA. The R code and data to replicate the simulations used in this manuscript can be found at https://osf.io/6jrhz/.

(1) Simple rule: This transmission rule follows the logic of the classic NBDA framework:

$$256 \qquad \lambda_i(t) = \lambda_o(t)\left(s\sum_{j=1}^{N} a_{ij}z_j(t) + 1\right)(1 - z_i(t))$$

Here $\lambda_i(t)$ represents the rate at which individual *i* acquires a novel behaviour as a function of time.
$\lambda_o(t)$ represents a baseline rate function (i.e. the rate of asocial learning at time *t*) and *s* determines
the strength of social transmission. $z_i(t)$ is the 'status' of individual *i* at time *t*, (1=informed; 0=naïve),
and N is the number of individuals in the network. The rate at which an individual acquires a novel
behaviour through social learning is proportional to $\sum_{j=1}^{N} a_{ij}z_j(t)$, the total connections to informed
individuals at time *t*. Therefore, *s* gives the rate of transmission per unit connection relative to the rate
of asocial learning of the novel behaviour. For example, when s=2, an increase of 1 in an individuals'
edge weights to informed individuals will increase the rate of social learning by 2 times the baseline
rate. $(1 - z_i(t))$ ensures that only naïve individuals acquire the behaviour. Consequently, the more
and stronger connections to informed individuals, the more likely an individual is to adopt the
behaviour. The social transmission strength *s* was set to 5.

Following, we define three more complex rules that generalise the classic NBDA model as:

$$269 \qquad \lambda_i(t) = \lambda_o(t)(T(a_i, z(t)) + 1)(1 - z_i(t))$$

270   Here a_i represents the connections individual *i* has to all others in the network, $z(t)$ gives the status
271   of each individual in the network at time *t*, and $T(a_i, z(t))$ is a transmission function determining how
272   the rate of transmission is determined by a_i and $z(t)$.

273   **(2) Threshold rule**: This transmission rule is defined as:

$$274 \qquad T(a_i, z(t)) = \left(\frac{c}{1 - \frac{1}{1 + exp(ab)}}\right)\left(\frac{1}{1 + exp\left(-b(\sum_j a_{ij}z_j(t) - a)\right)} - \frac{1}{1 + exp(ab)}\right)$$

Similar to the classic NBDA, the rate of social transmission is zero when the total connections to
informed individuals, $\sum_j a_{ij}z_j(t)$=0. However, the rate of transmission increases suddenly as the
threshold, *a*, is approached, to a maximum value of *c*. The parameter *b* determines how sharp the
threshold effect is. Our aim here is to generate a model with a clear sharp threshold, so we set *b*=3 for
our simulations (for details and other parameter settings for *b*, see Firth et al. 2020). We set the
threshold value *a* to 5 for all networks. This means that an individual's weighted connections to
informed individuals need to be ≥ 5 before an individual's behavioural adoption is likely to stem from
social learning. For example, if an individual with two connections with weights of 2 each to informed
others adopts the new behaviour, the probability of this adoption event stemming from social learning
is low. In contrast, if the individual's two connections have a weight of 3 each, the behavioural
adoption likely stemmed from social learning under the set threshold rule (see also Firth et al. 2020
for more details). The social transmission strength *s* was set to 5.

(3) Proportion rule: This transmission rule is defined as:

$$288 \qquad T(a_i, z(t)) = s\frac{\sum_j a_{ij}z_j(t)}{\sum_j a_{ij}}$$

Here the learning rate is proportional to the ratio of connections that an individual i holds to informed
others. As such, the individual with the highest proportion is most likely to learn and assumes
additional influence from individuals' uninformed connections rather than just considering the sum of
connections to informed individuals such as in the simple and threshold model. The social transmission
strength s was set to 5.

(4) Conformity rule: Finally, the fourth transmission rule assumes that individuals are
disproportionately more likely to copy the majority of the population (i.e. frequency dependent):

296
$$T\big(a_i, z(t)\big) = s \frac{\big(\sum_j a_{ij} z_j(t)\big)^f}{\big(\sum_j a_{ij} z_j(t)\big)^f + \big(\sum_j a_{ij} \big(1 - z_j(t)\big)\big)^f}$$

Here the frequency dependence parameter is $f \geq 1$, and $s > 0$. When $f=1$ this model reduces to the
proportional model above, and as f increases the strength of conformity bias increases. Thus, an
individual is expected to adopt a new behaviour if it is perceived as being performed by the majority
of its' social connections. Similar to the proportional rule, the conformity rule considers an individual's
informed and uninformed connections. For our simulations, we set f to 5 and the social transmission
strength s to 5.

For each simulation the individual 'initiating' the behaviour (i.e. the demonstrator) was randomly
chosen, and was then used across the four transmission models. We then simulated behavioural
spread across the entire network under each transmission model separately. This means that at each
timestep one new individual adopted the seeded behaviour, whereby each time each individual has a
given probability of adopting the behaviour through social learning which ranges from 0 (asocial
acquisition, i.e. an individual has no connections to informed individuals and cannot socially learn the
behaviour under the set learning rule) to 1 (social acquisition, that is when an individual has the
maximum probability of learning socially under the set learning rule). Therefore, at each timestep, the
new individual adopting the behaviour was stochastically chosen based on their probability of
adopting the behaviour in the previous timestep (i.e. where the one most likely to be chosen was the
one most likely to adopt the behaviour under the given social learning rule). For instance, under the
proportion rule, the next individual adopting the behaviour (i.e. from t1 → t2, see Figure 1) would
most likely be the one with the highest proportion of connections to informed others. Following, we
inferred the probability of this adoption event stemming from social learning. For instance, for an
individual that adopted the novel behaviour with a proportion of connections to informed others of
0.2, the probability of the behavioural adoption stemming from social learning would be lower
compared to the individual having a proportion of 0.7.

Once individuals adopted the novel behaviour, they remained 'informed' within each simulation run.
For each network, we repeated the simulations 100 times to minimize the influence of the identity of
the randomly selected demonstrator on the subsequent transmission pathways. We selected 100
simulation runs because this was enough in acquiring a relatively stable mean correlation coefficient
between network metric and order of acquisition (Figure S3). We repeated our simulations testing
various parameter combinations for s (1, 5, 10), f (3, 5, 7), and a (3, 5, 7) which we consider appropriate
for our study system (i.e. the average strength of an individuals' connections is approximately 2.5). In
the main text we present results for $s=5$, $f=5$, $a=5$ and as such a total of 537200 simulations (1343
weekly, local networks x 4 different learning rules x 100 simulation runs). Results for all other

parameters can be found in the supplementary material and with the code provided other parameter
combinations can be tested.

Data summary statistics

To assess the relationship between each individual's network metric and the order of acquisition, we
calculated Spearman's rank correlation coefficients. After every simulation, for each of the networks
and transmission models, we calculated the Spearman's rank correlation coefficient between the
order in which individuals adopted the behaviour (always excluding the demonstrator) and each of
the three individual network metrics (i.e. the weighted clustering coefficient, the weighted degree and
the weighted betweenness). For an overview of the transmission process, see Figure 1. For each
network and model, we then calculated the average correlation coefficient for each network metric
across the 100 simulations (Figure 1). To additionally assess the general relationships between the size
of the network, and the correlation coefficient between individuals' centralities and their acquisition
order, we used linear mixed-effect models (LMM) using the 'lme4' package (Bates et al. 2015). For
each model separately, we set the average correlation coefficient as the dependent variable and
network size as the predictor variable. Location identity and week nested in year were set as random
effects to factor in these differences when assessing this relationship (Table S2). We examined mode
assumptions and fit using graphical methods (e.g. qq plot of residuals, fitted values versus residual
plots, (Korner-Nievergelt et al. 2015)).

Figure 1. Schematic overview of the simulation procedure. First, a weekly social network of one of
the feeder locations (shown as black dots) in the study site was selected. Second, behavioural spread
was simulated on the selected network using four different social learning rules (i.e. simple, threshold,
proportion, conformity). The starting point (i.e. the first individual performing the new behaviour) was
randomly chosen. Then, at each timestep (t1-tn), a naïve individual adopted the novel behaviour with

a given probability of the adoption event being from social learning (dependent on the social learning rule at play; see methods for further details) until all individuals in the network had adopted the novel behaviour. Third, we calculated a correlation coefficient (Spearman's rank correlation coefficient) between three individual social network metrics (i.e. weighted clustering coefficient, weighted degree and weighted betweenness) and the order (i.e. timestep) in which individuals adopted the novel behaviour. Finally, we repeated this process 100 times for each weekly, local social network.

Results

The data in this study were simulated across 1343 empirically-derived social networks, inferred from recordings of 1774 individual great tits at 62 feeder locations across 39 weekends and 3 years. Social networks varied in size (right skewed distribution towards smaller networks, see Figure S4) and consisted of an average of 21.7 individuals (min=10, max=77, sd=10.0; note that we excluded social networks smaller than 10 from the analysis (see methods) resulting in the minimum network size of 10). For each location, we included on average 21.7 networks into the analysis (min=1, max=39, sd=11.9) and each individual was part of on average 16.5 networks (min=1, max=88, sd=13.2). Individuals on average visited 1.3 different feeder locations on a weekend (min=1, max=10, sd=0.7) and from 21036 occasions where individuals were recorded on a given weekend, individuals had visited only one location in 14888 occasions (71%).

Relationship between individual social behaviour and order of acquisition

Simulating behavioural spread under four different social learning rules revealed different transmission pathways across social networks. For both the simple and the threshold rule, the weighted clustering coefficient was on average positively related to the order of acquisition, with more clustered individuals adopting the seeded behaviour later than less clustered individuals across different network sizes (Figure 2). Weighted degree and betweenness were on average negatively related to the order of acquisition. Thus, individuals with higher weighted degree and weighted betweenness adopted the novel behaviour on average faster than individuals with a lower weighted degree and betweenness (Figure 2). In addition, we show the relationship between mean individual network metrics and the standardized order of acquisition across network sizes (Figure S5). For the simple learning rule, the relationship between network metric and order of acquisition only changed when the majority of individuals in a network had already adopted the behaviour (with approximately 75% of individuals knowledgeable; Figure S5). Behavioural spread under the threshold rule showed a small 'hump' shortly after the start of behavioural spread (Figure 2, S5), before the relationship between mean network metric and order of acquisition increased for weighted clustering coefficient or decreased for weighted degree and betweenness (Figure 2, S5). This 'hump' indicates that initially individuals with lower weighted clustering coefficient, higher weighted degree and betweenness adopted the behaviour first, before the relationship between mean network metric and order of acquisition turned (Figure 2, S5). In contrast to the simple and threshold rules, there was little or no relationship between individual social network metrics and the order of acquisition under both the proportion and conformity learning rules (Figure 2, S5).

Figure 2. Relationship between individual network metric and the order of acquisition for each social learning rule. Each column shows a different network metric (left to right: weighted clustering coefficient, weighted degree and weighted betweenness). Each row represents one of the four spreading rules (top to bottom: simple, threshold, proportion and conformity). Lines plot the average network metric for each order of acquisition and ribbons show the 95% confidence interval from the 100 simulations for each binned group of network sizes. Colour represents network size with darker colour indicating smaller networks. The social transmission rate, the threshold location and the frequency dependence parameter were set to 5.

Assessing the relationship between each individuals' network metric and the order of acquisition
across all social networks and simulation runs, revealed substantial variation in relationship strength
(Figure 3). Across network metrics, there were on average the strongest positive (Figure 3: weighted
clustering coefficient) and negative (Figure 3: weighted degree and betweenness) correlations under
the simple rule, and coefficients for the threshold, proportion and conformity model were lower. Even
though Figure 2 indicates a clear relationship between average network metric and order of
acquisition under the threshold model (Figure 2), correlation coefficients were very small (Figure 3).
This may be because of the prominent 'hump' at the initial stages of the spread under the threshold
rule (Figure 2, S5), leading to overall low correlation coefficients between individual network metrics
and order of acquisition (Figure 3).

The direction of the relationship between average network metrics and order of acquisition remained
unchanged when setting lower or higher parameters for the social transmission strength 's' (Figure
S6, S7). However, for the simple rule, the correlation coefficients became on average stronger under
larger social transmission rates (Figure S8). Further, different values for the frequency dependence 'f'
under the conformity rule, and different values for the threshold location 'a' under the threshold rule
did not change the general direction of the relationship between average network metrics and order
of acquisition (Figure S9, S10, S11). However, under the threshold model, increasing the threshold
location on average reduced the correlation between network metrics and order of acquisition (Figure
S10, S11).

Figure 3. **Distribution of average correlation coefficients for each social learning rule and network**
metric. Violin and boxplots show the distribution of the average correlation coefficients across 100
simulations from each network for each of the four social learning rules (i.e. simple, proportion,
conformity and threshold). Each plot shows one of the individual network metrics (weighted clustering
coefficient, weighted degree, weighted betweenness).

Relationship between social network size and pathways of behavioural diffusion

The direction and magnitude of the correlation between individual sociality and their order of
acquisition were partly predicted by network size (Figure 4, Table S2). For the simple and threshold
model, behavioural spread on larger networks led to more positive correlations between individual
network metric and order of acquisition for weighted clustering coefficient and more negative
correlations for weighted degree and betweenness (Figure 4, Table S2). The predicted effects of

network size on mean correlation coefficient inferred under the proportion and conformity rule suggest contrasting directions or no relationship with network size (Figure 4, Table S2).

Overall, the predicted effects were small, particularly for more complex contagions (Figure 4, Table S2). The relationship between correlation coefficient and network size was modulated by the parameters set for the social transmission rate (Figure S12). For the simple model, the direction of the relationship did not change across different 's' parameters but there were on average more negative correlations for larger 's' vales (Figure S8, S12). The slope remained relatively constant across social transmission rates, except for weighted betweenness, the relationship between correlation coefficients and network size was absent under a strong transmission rate (i.e. s=10; Figure S12). For the threshold model, the direction of the relationship did not change across different 's' parameters but the relationship was strongest (i.e. steepest slope) for larger social transmission rates (Figure S12). Similar patterns are present for weighted degree and weighted betweenness under the proportion and conformity model (Figure S12). For the weighted clustering coefficient, the slope indicated opposing directions for different transmission rates under the proportion and conformity model but were generally small and non-significant (Figure S12). The relationship between correlation coefficient and network size was also modulated by the parameters set for the threshold location 'a' (Figure S13). Here, the relationship between correlation coefficient and network size was strongest for small threshold locations (Figure S13). The frequency dependence parameter had no effect on the predicted relationship (Figure S13).

Figure 4. **Relationship between correlation coefficient and network size across the four social learning rules.** Each row shows one of the individual network metrics (top to bottom: weighted clustering coefficient, weighted degree and weighted betweenness) and each column a different social learning rule (left to right: simple, threshold, proportion, conformity). Average correlation coefficients across the 100 simulations per network are plotted as count dots (larger dots indicate more values for the respective value), lines represent the predicted effects generated from linear mixed-effect models (LMM) and ribbons represent the 95% confidence intervals (see Table S2 for model results).

Probability of social spread

Across the different social learning mechanisms, the average probability that an individual socially adopted the seeded behaviour increased with increasing timestep (i.e. order of acquisition, Figure 5). This is because with an increasing number of individuals becoming knowledgeable, the probability to be connected to informed others and thus to reach the set learning criterion increased (Figure 5). Behaviours were most likely to socially spread under the simple and proportion model (on average a non-zero probability to socially spread, Figure 5). That is because both rules did not require to surpass a given threshold (such as set for the threshold rule or reaching the majority of an individual's connections as set for the conformity rule). Therefore, one connection to an informed conspecific was

sufficient for the seeded behaviour to spread socially. However, under more complex spreading processes, social learning was limited (Figure 5). This was particularly the case for smaller networks when transmission followed the threshold rule. For instance, in networks consisting of less than 20 individuals, the set threshold for socially adopting the behaviour was almost never met (average probability of social learning close to 0; Figure 5 Threshold). For the conformity rule, larger networks limited social spread, but only at the initial phase of transmission (Figure 5 Conformity). In larger social networks, individuals have on average more social connections and thus at the initial stages of behavioural spread, it is less likely that the majority of an individual's social connections are already informed.

For both the simple and threshold model, the mean probability of social spread initially increased but then dropped towards the later adoptions (Figure 5). This may be caused by a few individuals that are not well connected in the social network, and thus have in general a low probability to socially learn under the simple and threshold rule. For instance, some great tits may only be connected to one other individual and thus, even if the number of knowledgeable individuals in the social network increases over time, it is very unlikely to reach the required threshold for social learning. For the proportion and conformity rule, the average probability of social learning peaked after a given percentage of individuals being knowledgeable (Figure 5).

The probability to socially adopt a behaviour was modulated by the model parameters chosen. Across all four models, the probability of social learning increased with an increasing social transmission rate 's' (Figure S14). Further, for the threshold model, the likelihood of social spread decreased with increasing threshold location 'a', especially in smaller networks (Figure S15). For the conformity model, changing the frequency dependence parameter 'f' did only slightly affect the probability of social learning in larger networks whereby with higher parameters social spread at the initial stage was more limited (Figure S15).

Figure 5. Relationship between the probability of an individual socially adopting the seeded behaviour and the number of informed individuals within the network. Panels show the results for each of the four social learning rules (simple, threshold, proportion, conformity). The x-axis describes the number of informed individuals within a social network. The simulations are set so that at each timestep a new individual adopts the behaviour, whereby each time each individual has a probability of adopting the behaviour through social learning (y-axis) given the set learning rule. Lines plot the average probability for each timestep and ribbons show the 95% confidence intervals from the 100 simulations across the binned groups for different network sizes. Colour represents network size with darker colour indicating smaller networks. The social transmission rate, the threshold location and the frequency dependence parameter were set to 5.

Discussion

Using simulations on large numbers of empirical great tit social networks, we show how the underlying social learning rules individuals employ strongly influence the transmission pathways of behaviours across social networks derived from real-world data. Under learning rules that rely purely on the extent of social connections to informed others, we found that individual great tits with a higher weighted degree and betweenness, and lower clustering coefficients were likely to adopt the seeded behaviour faster, in line with common expectations of the benefits of sociality for gaining information. However, if the likelihood of adopting a behaviour depended on the ratio of connections to informed and uninformed others, such as conformist learning, social connectivity was not strongly related to the order in which individuals acquired the seeded behaviour. Notably, this contrasts with the widely proposed prediction that more social individuals may be more likely to adopt new information. Thus,

our results show how the relationship between individual sociality and behavioural acquisition can
change fundamentally with the type of social learning mechanism at play. Finally, we reveal that the
probability of social spread under certain social learning rules is predicted to be limited in certain real-
world settings, particularly in networks of a very small or large size.

Individuals differ in the quantity and quality of social connections to others which impacts several
aspects of life history (McDonald 2007; Formica et al. 2012; MacIntosh et al. 2012; Farine and Sheldon
2015; Alberts 2019; Beck et al. 2021), including the social transmission of information (Aplin et al.
2012; Kulahci and Quinn 2019). Generally, individuals that hold many social connections to others,
and occupy central network positions, are expected to be more likely to acquire information (Hoppitt
and Laland 2011; Aplin et al. 2012; Claidiere et al. 2013; Kulahci et al. 2016). In line with these
assumptions, we find that great tits with lower weighted clustering coefficients, and higher weighted
degree and betweenness were more likely to adopt the seeded behaviour faster when the underlying
spreading mechanism depended on the extent of connections to knowledgeable others (Figure 2).
However, when rules were more complex, for example when behavioural adoption depended on the
ratio of connections to knowledgeable and naïve individuals, sociality was not strongly related to the
order of acquisition (Figure 2), matching the expectations from studies on complex contagions in
humans (Guilbeault et al. 2018). For instance, in humans individuals with more social connections
require stronger exposure to identify 'useful' information from the noise received from all their
associates (Hodas and Lerman 2012; Hodas and Lerman 2014) and individuals with fewer social
connections may utilize information sooner (González-Avella et al. 2011). These findings suggest that,
under certain spreading mechanisms, the extent of social connections can reduce social spread. Thus,
the concept that more social and central individuals are especially important in acquiring and
subsequently spreading information (Kulahci and Quinn 2019) cannot be generalized across different
types of behaviours, and may even lead to erroneous conclusions. For instance, if no evidence is found
that individual sociality is related to the probability of behavioural adoption, one might be led to
conclude the absence of social learning where, in fact more complex social learning rules may be in
operation.

These findings have important consequences for our understanding of the relationship between
sociality and behavioural acquisition, and also for what may constitute an 'optimal' social structure
for efficient social transmission (Pasquaretta et al. 2014; Romano et al. 2018; Cantor et al. 2021). In
many social structures, individuals differ in their social connectivity, ranging from highly to less
connected individuals (i.e. heterogenous degree distribution). In networks with higher variation in
connectivity, simple contagions may spread more efficiently because highly connected individuals can
act as 'hubs' (Evans et al. 2020; Xue et al. 2020). In contrast, more complex contagions may spread
more slowly on networks with heterogenous degree distribution (Evans et al. 2020; Xue et al. 2020).
Many animals, live in fission-fusion societies (Amici et al. 2008; Silk et al. 2014; Wilson et al. 2014)
where individuals frequently join, leave, and re-join groups which can result in large individual
differences in social connectivity (Sah et al. 2018). In contrast, some animals form highly stable groups
(e.g. many primates and carnivores; Kappeler and van Schaik 2002; Holekamp et al. 2007) with lower
individual variation in social connectivity (Sah et al. 2018). In social networks with heterogenous
degree distribution, an individual's social network position may, under certain behavioural contagions,
have a strong impact on the probability of behavioural acquisition whereas in groups with
homogenous degree distribution, an individuals' position within the network may not be as important.
Ultimately, the 'optimal' social structure for information transmission will highly depend on the

behaviour and the underlying learning mechanism. For future work it will be interesting to test how
different behavioural contagions spread on social networks of different species, ideally with
contrasting social structures.

Our results have implications for our understanding of the costs and benefits of individual sociality.
While increased access to information is one of the postulated key benefits of sociality, simply holding
more connections to others may in fact hinder the adoption of novel behaviours under certain social
learning rules. For instance, if a novel behaviour follows a conformist learning mechanism, highly social
individuals with lots of connections may be exposed to the new behaviour sooner than less social
individuals because they are more likely to be connected to at least one informed conspecific, but
despite this will be less likely to adopt the novel behaviour if it requires the majority of their social
connections to become informed first (Firth 2020) or if many social connections make the detection
of useful information more difficult (Hodas and Lerman 2012; Hodas and Lerman 2014). In addition,
our results suggest that occupying more peripheral network positions may be costlier in terms of
behavioural adoption than occupying central network positions is beneficial, but that this is dependent
on the learning rule. For instance, the proportional rule and conformity rule did not show that
individuals with lower social connectivity were likely to adopt very late, yet the simple and threshold
learning rule showed that the average network centrality of late adopters was very low (after
approximately 75% of individuals being informed; Figure S5) but remained relatively unchanged for
earlier adoptions (before 75%, Figure S5). This suggests that in various cases, poorly connected
individuals adopt the behaviour last and have in general a very low probability of socially adopting
behaviour (see decrease in social learning probability for simple and threshold rule, Figure 5) but that
this can be negated when certain social learning rules (e.g. the proportional and conformity rules) are
in play. We speculate that the relationship between network position and order of acquisition at the
initial stage (i.e. before approx. 75%), and the large variation in correlation coefficients (Figure 3) is
also highly determined by the demonstrators' network position (i.e. the starting positing of spread
(Banerjee et al. 2013)) and the underlying network structure (Romano et al. 2018; Evans et al. 2020;
Cantor et al. 2021) which may warrant future work. Finally, how biologically meaningful differences in
the order of acquisition are will ultimately depend on the behaviour and context, and the actual
observed variation in adoption times. Therefore, quantifying real individual variation in the timing of
behavioural acquisition in the wild will be crucial for our understanding of the potential costs and
benefits of individual sociality.

Examining behavioural innovation and its subsequent spread in nature is challenging (Whiten and
Mesoudi 2008; Klump et al. 2021). This is because new behaviours are often only detected once the
majority of individuals in a population are already knowledgeable. Alternatively, behavioural
innovations may occur much more frequently but remain undetected if the behaviour does not spread
far (e.g. because the behaviour is mechanically challenging for individuals (Gajdon et al. 2006) or when
the carryover of older, outdated behaviour hinders the spread of novel and more adaptive behaviours
(Aplin et al. 2017; Barrett et al. 2019)). Here we used simulations on real-world empirical networks,
and demonstrate that the ability of a behaviour to socially spread depends on its underlying social
learning mechanism. Our findings are thus consistent with other studies that investigated different
spreading processes (Nunn et al. 2009; Evans et al. 2020; Cantor et al. 2021; Evans et al. 2021). For
instance, Evans et al. (2020) simulated a simple and conformity contagion on different social structures
and show that disease/information spreads faster (i.e. time until a certain number of individuals had
been infected/informed) under a simple contagion compared to a conformity contagion. Our study

supplements past research by specifically focusing on the diffusion dynamics on the individual rather
than the population level. Across the four models, the probability that individual great tits socially
adopted the seeded behaviour increased with increasing timestep (Figure 5). This is expected as at
each timestep in simulations a new individual becomes knowledgeable and remains in this state
thereby increasing the number of knowledgeable individuals within the network. However, social
transmission under certain social learning mechanisms was limited. For instance, behaviours that
needed to surpass a given threshold of social connections to informed others (i.e. threshold or
conformity rule) required more asocial learning events in the initial spreading phase to be able to
subsequently transmit via social learning (Figure 5, Figure S13). In contrast, for the simple and
proportion rules, the probability of social transmission was always higher, even in the initial spreading
phase and when considering different transmission rates (Figure 5, Figure S14). This demonstrates
that in the initial stage some behaviours may have a higher likelihood to spread socially, whereas other
behaviours, following more complex processes, may rely on a larger extent of asocial learning events
or social reinforcement. Therefore, the majority of empirical research on behavioural spread in
animals may in fact examine more simple social transmission processes (because they are easier to
detect and observe) which may bias our picture of the existing social learning mechanisms of animals
and behavioural innovations.

We found that network size impacted behavioural spread across the four transmission models. The
strength of the relationship between individual network metric and the order of acquisition increased
with increasing network size under the simple and threshold rule (Figure 4) and was mediated by the
social transmission rate (Figure S12) and the threshold location (Figure S13). The weaker correlation
between individual network metrics and order of acquisition in smaller networks might be caused by
the relatively reduced likelihood of social spread (Figure 5). For instance, for the threshold rule,
individuals in larger networks have an overall higher number of social connections (compared to
individuals in smaller networks) which facilitates reaching the set threshold for social learning and
thus increases the likelihood of social spread. This is supported by our results when simulating spread
using different threshold values where behaviours were most likely to socially spread across different
network sizes under small thresholds (see Figure S10). In addition, our findings may suggest that in
smaller networks, the order of acquisition is mainly predicted by who an individual is connected to
(e.g. whether it is directly connected to the demonstrator) compared to the number and extent of
social connections it has. In contrast, in larger networks, the probability of behavioural acquisition
may be strongly influenced by the number and extent of connections an individual has. Therefore, the
importance of 'who' you know versus 'how many you know' may differ in networks of varying size.
Past studies have investigated behavioural spread on relatively large and static networks (using
simulations: (Voelkl and Noë 2008; Nunn et al. 2009; Cantor et al. 2021; Evans et al. 2021) or natural
observations: (Allen et al. 2013; Aplin et al. 2015a)). However, in many species social associations can
be highly dynamic and only the social connections at a relatively small temporal scale (e.g. at the time
of emergence) may predict an individuals' decision to adopt a novel behaviour. Our findings that
network size can impact behavioural spread thus have important consequences for our understanding
of the influence of wider society structure on when and where behaviours may emerge, and how to
interpret empirical results. As such, it is important to improve our understanding of the factors that
give rise to different social network sizes and structures and to consider networks on an appropriate
spatiotemporal scale or dynamic versus static networks (Hobaiter et al. 2014; Hasenjager et al. 2021).
For instance, spatiotemporal variation in environmental features (such as the availability and

distribution of resources) may influence population densities and subsequently local social network
size and structure across space and time, and as such may influence local social spread.

Our study suggests further questions for future research. Our models assume that each individual
adopts a seeded behaviour under the same 'learning rule'. While variation in individual learning rules
and their impact on behavioural contagions have been widely examined in humans (McCullen et al.
2013; Melnik et al. 2013; Muthukrishna et al. 2016; Aral and Nicolaides 2017), its' investigation in
animals remains scarce. However, individuals may differ in the extent of social information use and
the thresholds required for behavioural adoption (Chimento et al. 2022), which may be context and
state-dependent (Rendell et al. 2011; Penndorf and Aplin 2020). For instance, dominance rank
(Krueger et al. 2014) and sex (Aplin, Sheldon, et al. 2013) have been shown to be related to variation
in social information use and individuals may only copy behaviour from certain individuals, based on
familiarity or kin (Kavaliers et al. 2005; Boogert et al. 2018). Therefore, learning rules may differ within
individuals (e.g. with changes in age or dominance) and between individuals (e.g. sex, relatedness).
Future simulation and empirical studies could explore how heterogeneity in learning rules, within and
between individuals, and variation in acquisition versus adoption, impact information flow (Chimento
et al. 2022). In addition, we only explored learning rules which might be relevant for our study system.
However, future research could test the same and different rules on species with different social
structures. For instance, in species with more stable social groups such as in primates and many
carnivores (Kappeler and van Schaik 2002; Holekamp et al. 2007), kin-based or dominance-based
learning rules may be more applicable.

Furthermore, using simulations did not allow us to test changes in the social network resulting from
behavioural adoption: individuals may occupy more central social network positions once performing
a new behaviour (Kulahci et al. 2018; Kulahci and Quinn 2019). This can be the case if individuals
preferentially associate with knowledgeable others (Kulahci et al. 2018), or if individuals change their
behaviour in response to information acquisition which can also lead to an increase in social
connections (Kulahci and Quinn 2019). Such dynamics are not reflected in our study, and would
require the investigation of natural behavioural spread, ideally under experimental conditions in the
wild. In addition, our models assume that once individuals become 'informed', they cannot return to
an 'uninformed' state. In natural conditions, however, individuals may return again to an 'uninformed'
state if the novel behaviour was not rewarding. Future studies incorporating these aspects could
provide further new insights into the patterns of social transmission and its link to sociality. Finally,
we used empirical networks to capture the fine-scale social association patterns between wild birds
and to explore how different behavioural contagions spread on them. While it is important to test
predictions on real networks, such an approach also has additional considerations. For instance,
network metrics are often correlated and dependent on one another (Table S1) which makes it difficult
to tease apart the direct effects on the probability of behavioural adoption for each metric alone. In
studies using simulated networks, interdependencies can be controlled for in certain ways and the
effects of each metric could be teased apart (e.g. by using sensitivity analysis). Therefore, we suggest
that, while both empirical and simulation studies can provide valuable information on their own,
considering both complementary to each other will improve our understanding of how behaviours
spread on different networks and how different social network metrics relate independently to
behavioural adoption.

From copying the majority (Van de Waal et al. 2013; Aplin et al. 2015a; Danchin et al. 2018) to learning from specific tutors (Wild et al. 2019; Canteloup et al. 2020), individuals use a large range of different learning strategies (Hoppitt and Laland 2013; Kendal et al. 2018). While, an increasing number of studies show how social network structure can influence social transmission (using simulations: (Voelkl and Noë 2008; Romano et al. 2018; Evans et al. 2021) and empirical data: (Naug 2008; Firth et al. 2016; Romano et al. 2018)), our research highlights the importance of the underlying social learning mechanism in shaping the transmission pathways across social networks. We demonstrate that the common assumption that sociality is linked to a higher likelihood in acquiring information and adopting new behaviours cannot be generalized for behavioural spread in real world networks. This also sheds new light on our current understanding of the costs and benefits of individuals sociality and asks to focus more on the social learning mechanism at play, and to differentiate between access to information and behavioural adoption (Chimento et al. 2022). In addition, we reveal that social transmission can be limited under certain adoption rules (such as the threshold rule), and social networks of particular size. Our findings thus have important consequences for our understanding of whether and how behaviours spread across different social networks, and subsequently the establishment of traditions and cultures. Further, differences in spreading mechanisms alter predictions of what may constitute optimal social structures for the transmission of information, and how selection may act on sociality.

Acknowledgments

We thank the large number of contributors to the data collection. The study has been supported by grants from NERC (NE/S010335/1 & NE/V013483/1), ERC (AdG 250164) and BBSRC (BB/S009752/1).

Competing interests

The authors declare to have no competing interests.

References

Alberts SC. 2019. Social influences on survival and reproduction: Insights from a long-term study of wild baboons. J Anim Ecol. 88(1):47–66.

Allen J, Weinrich M, Hoppitt W, Rendell L. 2013. Network-based diffusion analysis reveals cultural transmission of lobtail feeding in humpback whales. Science (80-). 340(6131):485–488.

Amici F, Aureli F, Call J. 2008. Fission-fusion dynamics, behavioral flexibility, and inhibitory control in primates. Curr Biol. 18(18):1415–1419.

Aplin LM, Farine DR, Morand-Ferron J, Cockburn A, Thornton A, Sheldon BC. 2015a. Experimentally induced innovations lead to persistent culture via conformity in wild birds. Nature. 518(7540):538–541.

Aplin LM, Farine DR, Morand-Ferron J, Cockburn A, Thornton A, Sheldon BC. 2015b. Counting conformity: evaluating the units of information in frequency-dependent social learning. Anim Behav.

110:e5–e8. https://doi.org/10.1016/j.anbehav.2015.09.015.

Aplin LM, Farine DR, Morand-Ferron J, Sheldon BC. 2012. Social networks predict patch discovery in
a wild population of songbirds. Proc R Soc London B Biol Sci. 279(1745):4199–4205.

Aplin LM, Farine DR, Morand-Ferron J, Cole EF, Cockburn A, Sheldon BC. 2013. Individual
personalities predict social behaviour in wild networks of great tits (Parus major). Ecol Lett.
16(11):1365–1372.

Aplin LM, Sheldon BC, McElreath R. 2017. Conformity does not perpetuate suboptimal traditions in a
wild population of songbirds. Proc Natl Acad Sci. 114(30):7830–7837.

Aplin LM, Sheldon BC, Morand-Ferron J. 2013. Milk bottles revisited: social learning and individual
variation in the blue tit, Cyanistes caeruleus. Anim Behav. 85(6):1225–1232.

Aral S, Nicolaides C. 2017. Exercise contagion in a global social network. Nat Commun. 8(1):1–8.

Banerjee A, Chandrasekhar AG, Duflo E, Jackson MO. 2013. The diffusion of microfinance. Science
(80-). 341(6144):1236498.

Barrat A, Barthelemy M, Pastor-Satorras R, Vespignani A. 2004. The architecture of complex
weighted networks. Proc Natl Acad Sci. 101(11):3747–3752.

Barrett B, Zepeda E, Pollack L, Munson A, Sih A. 2019. Counter-culture: Does social learning help or
hinder adaptive response to human-induced rapid environmental change? Front Ecol Evol. 7:183.

Bates D, Maechler M, Bolker B, Walker S. 2015. lme4: Linear mixed-effects models using Eigen and
S4. R package version 1.1–7. 2014.

Beck KB, Farine DR, Kempenaers B. 2021. Social network position predicts male mating success in a
small passerine. Behav Ecol. 32(5):856–864. doi:10.1093/beheco/arab034.
https://doi.org/10.1093/beheco/arab034.

Boogert NJ, Lachlan RF, Spencer KA, Templeton CN, Farine DR. 2018. Stress hormones, social
associations and song learning in zebra finches. Philos Trans R Soc B Biol Sci. 373(1756):20170290.

Boyd R, Richerson PJ. 1988. Culture and the evolutionary process. University of Chicago press.

Bulmer MG, Perrins CM. 1973. Mortality in the great tit Parus major. Ibis (Lond 1859). 115(2):277–
281.

Cairns SJ, Schwager SJ. 1987. A comparison of association indices. Anim Behav. 35(5):1454–1469.

Canteloup C, Hoppitt W, van de Waal E. 2020. Wild primates copy higher-ranked individuals in a
social transmission experiment. Nat Commun. 11(1):1–10.

Cantor M, Chimento M, Smeele SQ, He P, Papageorgiou D, Aplin LM, Farine DR. 2021. Social network
architecture and the tempo of cumulative cultural evolution. Proc R Soc B. 288(1946):20203107.

Centola D. 2018. How behavior spreads: The science of complex contagions. Princeton University
Press Princeton, NJ.

Centola D, Macy M. 2007. Complex contagions and the weakness of long ties. Am J Sociol.
113(3):702–734.

Chimento M, Barrett BJ, Kandler A, Aplin LM. 2022. Cultural diffusion dynamics depend on
behavioural production rules. Proc R Soc B. 289(1980):20221001.

Claidiere N, Messer EJE, Hoppitt W, Whiten A. 2013. Diffusion dynamics of socially learned foraging
techniques in squirrel monkeys. Curr Biol. 23(13):1251–1255.

Coussi-Korbel S, Fragaszy DM. 1995. On the relation between social dynamics and social learning.
Anim Behav. 50(6):1441–1453. doi:10.1016/0003-3472(95)80001-8.

Csardi G, Nepusz T. 2006. The igraph software package for complex network research, InterJournal,
Complex Systems 1695. 2006. http://igraph.org.

Danchin E, Nöbel S, Pocheville A, Dagaeff A-C, Demay L, Alphand M, Ranty-Roby S, Van Renssen L,
Monier M, Gazagne E. 2018. Cultural flies: Conformist social learning in fruitflies predicts long-lasting
mate-choice traditions. Science (80-). 362(6418):1025–1030.

Ekman J. 1989. Ecology of non-breeding social systems of Parus. Wilson Bull.:263–288.

Evans JC, Hodgson DJ, Boogert NJ, Silk MJ. 2021. Group size and modularity interact to shape the
spread of infection and information through animal societies. Behav Ecol Sociobiol. 75(12):1–14.

Evans JC, Silk MJ, Boogert NJ, Hodgson DJ. 2020. Infected or informed? Social structure and the
simultaneous transmission of information and infectious disease. Oikos. 129(9):1271–1288.

Farine DR. 2013. Animal social network inference and permutations for ecologists in R using asnipe.
Methods Ecol Evol. 4(12):1187–1194.

Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015. Interspecific social networks promote
information transmission in wild songbirds. Proc R Soc B Biol Sci. 282(1803):20142804.

Farine DR, Sheldon BC. 2015. Selection for territory acquisition is modulated by social network
structure in a wild songbird. J Evol Biol. 28(3):547–556.

Farine DR, Whitehead H. 2015. Constructing, conducting and interpreting animal social network
analysis. J Anim Ecol. 84(5):1144–1163.

Firth JA. 2020. Considering complexity: animal social networks and behavioural contagions. Trends
Ecol Evol. 35(2):100–104.

Firth JA, Albery GF, Beck KB, Jarić I, Spurgin LG, Sheldon BC, Hoppitt W. 2020. Analysing the Social
Spread of Behaviour: Integrating Complex Contagions into Network Based Diffusions. arXiv Prepr
arXiv201208925.

Firth JA, Sheldon BC. 2016. Social carry-over effects underpin trans-seasonally linked structure in a
wild bird population. Ecol Lett. 19(11):1324–1332.

Firth JA, Sheldon BC, Farine DR. 2016. Pathways of information transmission among wild songbirds
follow experimentally imposed changes in social foraging structure. Biol Lett. 12(6):20160144.

Formica VA, Wood CW, Larsen WB, Butterfield RE, Augat ME, Hougen HY, BRODIE III ED. 2012.
Fitness consequences of social network position in a wild population of forked fungus beetles
(Bolitotherus cornutus). J Evol Biol. 25(1):130–137.

Franz M, Nunn CL. 2009. Network-based diffusion analysis: a new method for detecting social
learning. Proc R Soc B Biol Sci. 276(1663):1829–1836.

Gajdon GK, Fijn N, Huber L. 2006. Limited spread of innovation in a wild parrot, the kea (Nestor
notabilis). Anim Cogn. 9(3):173–181.

González-Avella JC, Eguíluz VM, Marsili M, Vega-Redondo F, San Miguel M. 2011. Threshold learning
dynamics in social networks. PLoS One. 6(5):e20207.

Granovetter M. 1978. Threshold models of collective behavior. Am J Sociol. 83(6):1420–1443.

Guilbeault D, Becker J, Centola D. 2018. Complex contagions: A decade in review. Complex spreading

Phenom Soc Syst.:3–25.

Hämäläinen L, Mappes J, Rowland HM, Teichmann M, Thorogood R. 2020. Social learning within and
across predator species reduces attacks on novel aposematic prey. J Anim Ecol. 89(5):1153–1164.

Hasenjager MJ, Leadbeater E, Hoppitt W. 2021. Detecting and quantifying social transmission using
network-based diffusion analysis. J Anim Ecol. 90(1):8–26.

Haun DBM, Rekers Y, Tomasello M. 2012. Majority-biased transmission in chimpanzees and human
children, but not orangutans. Curr Biol. 22(8):727–731.

Heyes CM. 1994. Social learning in animals: categories and mechanisms. Biol Rev. 69(2):207–231.

Hinde RA. 1952. The Behaviour of the Great Tit (parus Major) and Some Other Related Species. Brill.

Hobaiter C, Poisot T, Zuberbühler K, Hoppitt W, Gruber T. 2014. Social network analysis shows direct
evidence for social transmission of tool use in wild chimpanzees. PLoS Biol. 12(9):e1001960.

Hodas NO, Lerman K. 2012. How visibility and divided attention constrain social contagion. In: 2012
International Conference on Privacy, Security, Risk and Trust and 2012 International Confernece on
Social Computing. IEEE. p. 249–257.

Hodas NO, Lerman K. 2014. The simple rules of social contagion. Sci Rep. 4(1):1–7.

Holekamp KE, Sakai ST, Lundrigan BL. 2007. Social intelligence in the spotted hyena (Crocuta
crocuta). Philos Trans R Soc B Biol Sci. 362(1480):523–538.

Hoppitt W, Laland KN. 2011. Detecting social learning using networks: a users guide. Am J Primatol.
73(8):834–844.

Hoppitt W, Laland KN. 2013. Social learning. Princeton University Press.

Hoppitt WJE, Farine DR. 2018. Association indices for quantifying social relationships: how to deal
with missing observations of individuals or groups. Anim Behav. 136:227–238.

Kappeler PM, van Schaik CP. 2002. Evolution of primate social systems. Int J Primatol. 23(4):707–740.

Kavaliers M, Colwell DD, Choleris E. 2005. Kinship, familiarity and social status modulate social
learning about "micropredators"(biting flies) in deer mice. Behav Ecol Sociobiol. 58(1):60–71.

Kendal RL, Boogert NJ, Rendell L, Laland KN, Webster M, Jones PL. 2018. Social learning strategies:
Bridge-building between fields. Trends Cogn Sci. 22(7):651–665.

Klump BC, Martin JM, Wild S, Hoersch JK, Major RE, Aplin LM. 2021. Innovation and geographic
spread of a complex foraging culture in an urban parrot. Science.

Korner-Nievergelt F, Roth T, Von Felten S, Guélat J, Almasi B, Korner-Nievergelt P. 2015. Bayesian
data analysis in ecology using linear models with R, BUGS, and Stan. Academic Press.

Krueger K, Farmer K, Heinze J. 2014. The effects of age, rank and neophobia on social learning in
horses. Anim Cogn. 17(3):645–655.

Kulahci IG, Ghazanfar AA, Rubenstein DI. 2018. Knowledgeable lemurs become more central in social
networks. Curr Biol. 28(8):1306–1310.

Kulahci IG, Quinn JL. 2019. Dynamic relationships between information transmission and social
connections. Trends Ecol Evol. 34(6):545–554.

Kulahci IG, Rubenstein DI, Bugnyar T, Hoppitt W, Mikus N, Schwab C. 2016. Social networks predict
selective observation and information spread in ravens. R Soc open Sci. 3(7):160256.

MacIntosh AJJ, Jacobs A, Garcia C, Shimizu K, Mouri K, Huffman MA, Hernandez AD. 2012. Monkeys
in the middle: parasite transmission through the social network of a wild primate. PLoS One.
7(12):e51144.

McCullen NJ, Rucklidge AM, Bale CSE, Foxon TJ, Gale WF. 2013. Multiparameter models of
innovation diffusion on complex networks. SIAM J Appl Dyn Syst. 12(1):515–532.

McDonald DB. 2007. Predicting fate from early connectivity in a social network. Proc Natl Acad Sci.
104(26):10910–10914.

Melnik S, Ward JA, Gleeson JP, Porter MA. 2013. Multi-stage complex contagions. Chaos An
Interdiscip J Nonlinear Sci. 23(1):13124.

Muthukrishna M, Morgan TJH, Henrich J. 2016. The when and who of social learning and conformist
transmission. Evol Hum Behav. 37(1):10–20.

Naug D. 2008. Structure of the social network and its influence on transmission dynamics in a
honeybee colony. Behav Ecol Sociobiol. 62(11):1719–1725.

Nunn CL, Thrall PH, Bartz K, Dasgupta T, Boesch C. 2009. Do transmission mechanisms or social
systems drive cultural dynamics in socially structured populations? Anim Behav. 77(6):1515–1524.

Papastamatiou YP, Bodey TW, Caselle JE, Bradley D, Freeman R, Friedlander AM, Jacoby DMP. 2020.
Multiyear social stability and social information use in reef sharks with diel fission–fusion dynamics.
Proc R Soc B. 287(1932):20201063.

Pasquaretta C, Levé M, Claidiere N, Van de Waal E, Whiten A, MacIntosh AJJ, Pelé M, Bergstrom ML,
Borgeaud C, Brosnan SF. 2014. Social networks in primates: smart and tolerant species have more
efficient networks. Sci Rep. 4(1):1–8.

Penndorf J, Aplin L. 2020. Environmental and life history factors, but not age, influence social
learning about food: a meta-analysis. Anim Behav. 167:161–176.

Pike TW, Laland KN. 2010. Conformist learning in nine-spined sticklebacks' foraging decisions. Biol
Lett. 6(4):466–468.

Psorakis I, Roberts SJ, Rezek I, Sheldon BC. 2012. Inferring social network structure in ecological
systems from spatio-temporal data streams. J R Soc Interface. 9(76):3055–3066.

Psorakis I, Voelkl B, Garroway CJ, Radersma R, Aplin LM, Crates RA, Culina A, Farine DR, Firth JA,
Hinde CA. 2015. Inferring social structure from temporal data. Behav Ecol Sociobiol. 69(5):857–866.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rendell L, Fogarty L, Hoppitt WJE, Morgan TJH, Webster MM, Laland KN. 2011. Cognitive culture:
theoretical and empirical insights into social learning strategies. Trends Cogn Sci. 15(2):68–76.

Romano V, Shen M, Pansanel J, MacIntosh AJJ, Sueur C. 2018. Social transmission in networks: global
efficiency peaks with intermediate levels of modularity. Behav Ecol Sociobiol. 72(9):1–10.

Rosenthal SB, Twomey CR, Hartnett AT, Wu HS, Couzin ID. 2015. Revealing the hidden networks of
interaction in mobile animal groups allows prediction of complex behavioral contagion. Proc Natl
Acad Sci. 112(15):4690–4695.

Sah P, Mann J, Bansal S. 2018. Disease implications of animal social network structure: a synthesis
across social systems. J Anim Ecol. 87(3):546–558.

Silk MJ, Croft DP, Tregenza T, Bearhop S. 2014. The importance of fission–fusion social group

dynamics in birds. Ibis (Lond 1859). 156(4):701–715.

Somveille M, Firth JA, Aplin LM, Farine DR, Sheldon BC, Thompson RN. 2018. Movement and
conformity interact to establish local behavioural traditions in animal populations. bioRxiv.:1–19.
doi:10.1101/338657.

Thorogood R, Kokko H, Mappes J. 2018. Social transmission of avoidance among predators facilitates
the spread of novel prey. Nat Ecol Evol. 2(2):254–261.

Toyokawa W, Gaissmaier W. 2022. Conformist social learning leads to self-organised prevention
against adverse bias in risky decision making. Elife. 11:e75308.

Voelkl B, Noë R. 2008. The influence of social structure on the propagation of social information in
artificial primate groups: a graph-based simulation approach. J Theor Biol. 252(1):77–86.

Van de Waal E, Borgeaud C, Whiten A. 2013. Potent social learning and conformity shape a wild
primate's foraging decisions. Science (80-). 340(6131):483–485.

Watts DJ. 2002. A simple model of global cascades on random networks. Proc Natl Acad Sci.
99(9):5766–5771.

Whiten A. 2021. The burgeoning reach of animal culture. Science (80-). 372(6537).

Whiten A, Mesoudi A. 2008. Establishing an experimental science of culture: animal social diffusion
experiments. Philos Trans R Soc B Biol Sci. 363(1509):3477–3488.

Wild S, Allen SJ, Krützen M, King SL, Gerber L, Hoppitt WJE. 2019. Multi-network-based diffusion
analysis reveals vertical cultural transmission of sponge tool use within dolphin matrilines. Biol Lett.
15(7):20190227.

Wilson ADM, Krause S, James R, Croft DP, Ramnarine IW, Borner KK, Clement RJG, Krause J. 2014.
Dynamic social networks in guppies (Poecilia reticulata). Behav Ecol Sociobiol. 68(6):915–925.

Xue X, Pan L, Zheng M, Wang W. 2020. Network temporality can promote and suppress information
spreading. Chaos An Interdiscip J Nonlinear Sci. 30(11):113136.