Title

A latitudinal pattern of plant leaf-associated bacterial community assembly

Authors

Zihui Wang[1,2], Yuan Jiang[4], Minhua Zhang[2], Chengjin Chu[4], Yongfa Chen[4], Shuai Fang[10],

Guangze Jin[5], Mingxi Jiang[6], Juyu Lian[7], Yanpeng Li[11], Yu Liu[2], Keping Ma[8], Xiangcheng Mi[8],

Xiujuan Qiao[6], Xihua Wang[9], Xugao Wang[10], Han Xu[11], Wanhui Ye[7], Li Zhu[8], Yan Zhu[8],

Fangliang He[2, 3*] and Steven W. Kembel[1,2*]

Affiliations

1. Département des sciences biologiques, Université du Québec à Montréal, Montréal, Québec,

H2X 1Y4, Canada

2. ECNU-Alberta Joint Lab for Biodiversity Study, Tiantong Forest Ecosystem National

Observation and Research Station, School of Ecology and Environmental Sciences, East China

Normal University, Shanghai, 200241, China

3. Department of Renewable Resources, University of Alberta, Edmonton, Alberta, T6G 2H1,

Canada

4. State Key Laboratory of Biocontrol, School of Life Sciences / School of Ecology, Sun Yat-sen

University, Guangzhou, 510275, China

5. Center for Ecological Research, Northeast Forestry University, Harbin, 150040, China

6. Key Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical Garden,

Chinese Academy of Sciences, Wuhan, 430074, China

7. Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China

8. State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing, 100093, China

9. Zhejiang Tiantong Forest Ecosystem National Observation and Research Station, School of Ecology and Environmental Sciences, East China Normal University, Shanghai, 200241, China

10. CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang, 110016, China

11. Research Institute of Tropical Forestry, Chinese Academy of Forestry, Guangzhou, 510520, China

Abstract

Plant-associated microbes are essential for promoting plant well-being, maintaining biodiversity,

and supporting ecosystem function. However, little is known about the geographic distribution of

plant-microbe symbioses and how they are formed and change along latitudinal gradients. Here

we identified leaf bacteria for 328 plant species sampled from 10 forests along a tropical to

temperate gradient in China. We analyzed the diversity and composition of plant leaf-associated

bacteria and quantified the contributions of hosts, habitats, and neighborhood plants to the plant-

bacterial symbiosis. We found a strong latitudinal gradient in leaf bacterial diversity and

composition. Bacterial assemblages on leaves were most strongly selected by host plants, and the

selection pressure increased with latitude. In contrast, at low latitudes and at large geographical

scales multiple factors were found to jointly regulate bacterial community composition. Our

result also showed that plant-bacteria symbiotic networks were structured by network hub

bacteria taxa with high co-occurrence network centrality, and the abundance of temperate hub

taxa was more influenced by host plants than that in tropical forests. For the first time, we

documented a previously unrecognized latitudinal gradient in plant-bacterial symbioses that was

regulated by a joint effect of multiple factors at low latitudes but mostly by host selection at high

latitudes, implying that leaf microbiomes are likely to respond differently to global change along

the latitudinal gradient.

Main text

All animals and plants are colonized by microorganisms, which are critical for ecosystem function and global biogeochemical cycles[1]. As one of the most common types of biotic interactions, plant-microbial symbioses are essential for plant growth, health and stress resilience, and also for supporting ecosystem functions including enhancing productivity, biogeochemical cycling, and diversity maintenance[2,3]. However, our knowledge about plant-microbial symbioses is highly variable among symbiosis types, with belowground symbioses such as nitrogen-fixation symbioses and root mycorrhizal symbioses being extremely well studied while aboveground plant-microbe symbioses have not been explored as deeply[4,5]. An example of aboveground symbioses is microbial assemblages on leaves – a metasymbiosis that is formed not by individual microbes or a specific group of microbes (i.e. *Bradyrhizobium* bacteria or glomeromycete mycorrhizal fungi in the case of roots) but by the diverse metacommunity of microorganisms available to colonize leaves[6]. How such leaf-microbe associations are formed and maintained is a critical but unanswered question, leading to a major concern that the future role of plant microbiomes in supporting plant well-being and sustaining ecosystem functioning may be compromised in the face of global change[7,8]. To address this question, it is necessary to understand the geographic variation of the phyllosphere metasymbiosis across a broad geographic scale in order to gain insights into how leaf-microbe associations are assembled and how they vary across different habitat conditions and host compositions. This understanding of geographic patterns of phyllosphere microbial diversity is indispensable for predicting the responses of plant microbiomes to global change[7,8].

Bacteria are the most abundant microbes in the phyllosphere with an estimated 10^7 bacterial cells per cm^2 of leaf surface[9]. Considering the total area of plant leaf surfaces on Earth,

the global bacterial population in the phyllosphere is enormous (up to 10^{26} cells), as is their

impact on plant communities and ecosystem functions through effects including altering plant

immunity, modifying plant hormone production and facilitating global carbon and nitrogen

cycling[10–12]. The assembly of this ubiquitous bacterial leaf symbiosis is governed by a range of

factors that influence the community assembly processes such as selection and dispersal[13]. For

example, host plant attributes including species identity, functional traits and phylogeny shape

bacterial community assembly through selection processes[14–16]. Neighborhood plants also select

for and create a bacterial metacommunity that alters the composition of local communities

through dispersal[17,18]. Additionally, abiotic environments such as temperature and microhabitats

which are often spatially autocorrelated have been observed to affect leaf bacterial diversity[19–21].

However, these drivers of leaf bacterial assembly have commonly been examined separately in

previous studies. As a result, there is poor understanding of the relative importance of these

different drivers of plant-bacterial symbioses. Furthermore, current knowledge on the drivers of

plant-microbe associations is almost exclusively based on individual sites; we understand very

little how assembly processes of bacteria community vary along latitudinal gradients, in stark

contrast with our knowledge about latitudinal diversity gradients of higher plants and

animals[22,23].

Studies of microbial co-occurrence networks have revealed hub taxa that occupy central

positions within microbial interaction networks, and have demonstrated that these hub taxa play

key roles driving microbial community structure and function[24,25]. At the metacommunity level,

hub taxa connect local communities via dispersal and gene flow and influence local community

assembly by microbe-microbe and plant-microbe interactions, through which their population

dynamic could impact large-scale patterns of plant-bacterial interactions[26–28]. Identifying the hub

taxa in plant-bacterial association at the metacommunity level is a pressing need given their

importance in mediating eco-evolutionary feedbacks, but has been limited to date by a lack of

large-scale datasets on plant-microbe associations[29]. Furthermore, knowledge about the factors

that control the geographic distribution of these hub bacterial taxa is essential for predicting

future change in microbiome structure due to gain or loss of hub taxa, but is currently

unavailable for plant-associated microbes[30].

Here, we addressed these knowledge gaps by investigating geographic patterns of

phyllosphere bacterial diversity, their network structure and assembly mechanisms along a large

latitudinal gradient for a network of stem-mapped forest sites across China. We quantified

bacterial communities using high-throughput sequencing of the 16S rRNA gene for a total of

1453 leaf samples from 328 plant species from 10 forest sites across major forest biomes in

China. Specifically, we aimed to (1) document latitudinal patterns of leaf-associated bacterial

diversity, (2) disentangle the effects of hosts, environments, neighborhood plants and spatial

factors on the community assembly of phyllosphere bacteria along the latitudinal gradient and

across geographic scales, and (3) construct plant-bacterial symbiosis networks and identify the

metacommunity-level hub taxa that are instrumental to plant-bacteria associations; we further

modeled the distribution of these bacterial hub taxa along the latitudinal gradient.

Results

We identified 49254 16S rRNA gene amplicon sequence variants (ASV) from 1453 leaf

samples, belonging to 237 bacterial orders and 386 families. *Proteobacteria* were the dominant

phylum*,* accounting for 70% of the total abundance of sequences across all samples.

Beijerinckiaceae (*Rhizobiales*) were most abundant in tropical and subtropical regions while

Sphingomonadaceae (*Sphingomonadales*) were most abundant in temperate forests (Fig. S1).

ASV richness of phyllosphere bacteria decreased with latitude (Fig. 1a). There was strong

evidence that ASV richness was correlated positively with the diversity of neighborhood plants,

negatively with slope, and was linked with elevation and annual temperature in quadratic

relationships (Fig. S2). We used global non-metric multidimensional scaling ordinations to

visualize the variation in bacterial community composition and show clear clustering of bacterial

communities by study sites and host taxonomy (Fig. 1b&S3). Permutational multivariate analysis

of variance revealed strong evidence that climate, soil conditions, topography, neighborhood

plant diversity and host traits all impact bacterial community assembly (Table S1). There was a

higher similarity of phyllosphere bacterial communities on phylogenetically closely related

plants and spatially closer samples, leading to host phylogenetic structure and spatial

autocorrelation effects on bacterial community assembly at both local and large scales (Fig. S4).

We quantified the variation in bacterial ASV richness and composition explained by

multiple variables that have been proposed as drivers of host-associated microbial diversity using

generalized linear models and redundancy analysis respectively. The variables included host-

related factors (phylogenetic eigenvectors + plant traits), abiotic environment (soil conditions +

topography + climate), diversity and abundance of neighborhood plants, and spatial eigenvectors

(see Methods). At the local scale of individual sites, these models explained 26%-76% of the

variation in ASV richness and 19%-36% of the variation in community composition of

phyllosphere communities. At regional levels, these models explained 27%-68% of the variation

in species richness, and 33%-42% of the variation in community composition respectively. At a

'global' scale (defined as the scale including all study sites), the model explained 43% of the

total variation in phyllosphere bacterial species richness and 47% of the variation in community

composition across all sites (Fig. 2&S5).

We used variation partitioning to disentangle the independent and joint effect of these

variables and quantify their relative importance in structuring bacterial diversity and community

assembly. Generally, the independent effect of host was the most important in explaining both

phyllosphere bacterial ASV richness and community composition, accounting for 15-89% of the

explained variance, followed by abiotic environments and spatial eigenvectors (contributions of

up to 29% and 25% respectively), while the attributes of plant neighborhoods were less

important (only contributing 9% of explained variance). There was a strong joint effect of

different variables, which accounted for 8-72% of the explained variation in bacterial community

composition (Fig. 2). The relative importance of these factors varied across geographic scales

and was highly latitude dependent; for example, the independent effect of host was positively

correlated with latitude and was less important at large scale, while the joint effect of these four

variables decreased with latitude and was more pronounced at large scales, and the overall

contribution of host-related factors increased while that of other factors such as environment

decreased with latitude (Fig. 3).

We constructed metacommunity-level networks of plant-bacterial associations at regional

and global scales and quantified the topological positions of ASVs within each co-occurrence

network. Hub ASVs with high network betweenness centrality were identified for tropical,

subtropical, and temperate regional networks and a global network respectively (see Methods).

Hub bacterial ASVs differed across regions; in subtropical and tropical regions, bacteria hubs

were primarily ASVs belonging to the *Rhizobiales* genera *1174-901-12, Methylocella* and

Methylobacterium, while in temperate networks the hubs included *Sphingomonas*

(*Sphingomonadales*) and *Hymenobacter* (*Cytophagales*) (Fig. S6). These bacterial ASVs connect

with most of the nodes (i.e. plant species) in the association networks and structure plant-

bacterial association networks at metacommunity levels; plant-phyllosphere bacteria association

networks at local sites were interconnected by hub bacterial ASVs to form regional-scale

networks, and these regional networks were further linked together through hub ASVs at global

scales (Fig. 4).

The distribution of phyllosphere bacterial hubs was clustered along latitudinal gradients

(Fig. 1c). To determine the relative importance of different factors structuring the biogeography

of these hub ASVs, we applied a joint species distribution modeling approach where nine habitat

variables were taken as fixed effects and two distance matrices representing the spatial and host

phylogenetic structure as random effects (see Methods). The models explained up to 90% and

73% of the variation in presence-absence and abundance of hub bacterial ASV, respectively.

Overall, climate (including temperature, precipitation and isothermality) was the most important

factor explaining the distribution of phyllosphere bacterial hub ASVs (Fig. 5&S7). There was

also a major contribution of the random effects of spatial factors and host phylogeny on hub

ASV abundance, highlighting the importance of taking spatial and host phylogenetic structure

into account for unbiased inference and to improve the prediction of the distribution of plant

microbiomes. Host plant attributes (leaf traits and phylogeny) were more important in

determining the abundance of hub ASVs in temperate regions, while spatial-related effects were

more important in influencing tropical hub ASVs (Fig. 6). Additionally, we found a strong

phylogenetic signal in the responses of ASVs to habitat covariance, indicating that

phylogenetically closely related ASVs have similar habitat preferences (Fig. S8).

Discussion

In this study, we revealed for the first time a strong latitudinal pattern in the diversity,

community composition and distribution of plant-leaf associated bacteria, highlighting the

varying strength of community assembly processes along the latitudinal gradient in shaping plant

leaf-microbial symbioses. It is remarkable that we found a shift from strong host filtering in

bacterial community assembly in temperate forests, to weak host filtering but a strong joint effect

of multiple drivers in the tropics. These results suggest that plant-associated bacteria on leaves

are formed following different rules across latitudes, with an increased complexity of different

processes acting to explain bacterial symbioses at lower latitudes. The latitude-dependent

community assembly processes were consistently evident in driving bacterial ASV richness,

community composition and the abundance of metacommunity-level hub bacteria taxa, which

together provide the most comprehensive understanding to date of the patterns and mechanisms

of plant-bacterial associations at large geographical scales. Our findings imply responses of

plant-bacteria symbioses to host plant composition and environmental change differ in tropical

vs. temperate regions, serving as a baseline for predicting and managing plant-microbe

symbioses under global change.

Although host attributes, environment, climate, plant neighborhoods and spatial factors

have been shown to influence the diversity of phyllosphere bacteria in previous studies [14,16,18–20],

most of these factors were analyzed individually in these studies. As such, their relative

importance was unknown, which made it difficult to infer the processes underlying the observed

host-microbe associations[31]. By simultaneously evaluating a broad range of factors across spatial

scales, we have shown that host-related factors including plant phylogeny and leaf traits were the

most important drivers of phyllosphere bacterial community assembly, especially in temperate

regions. This result suggested that phylogenetically conserved host attributes including specific

leaf area and leaf nutrient concentrations could select for leaf-associated bacteria[14,15]. Our results

of independent effects of hosts on bacterial diversity at large spatial scales indicated consistent

matching between host plant and phyllosphere bacteria regardless of climates, which represents a

potential consequence of coevolutionary feedbacks between plant hosts and their associated

microbes[15,32]. The importance of host plant selection mediated by phylogenetically conserved

traits could further lead to host specialization of plant-associated microbes, which may allow

microbes to coexist and explain their incredible diversity on leaves, and also contribute to the

host-specific effects of plant-associated microbes in structuring plant communities[33–35].

While host selection was among the most important factor in explaining phyllosphere

bacterial community assembly, the joint effect of multiple other factors also played a critical role

particularly in tropical forests. Many of these factors including spatial structure and neighboring

plant communities are likely related to the importance of stochastic processes including dispersal

for community assembly of plant-associated bacteria, suggesting a latitudinal patterns in the

relative contribution of deterministic and stochastic process to shaping plant-bacteria association.

This result concurred with the prediction of the niche-neutral continuum hypothesis that the

importance of niche-based processes such as selection are stronger in low diversity temperate

regions while the importance of species-neutral processes such as dispersal are more important in

high diversity tropical regions [22,36]. Previous studies have shown that the processes structuring

plant community assembly increasingly deviated from neutrality with latitude, and here we

provided the first evidence that this hypothesis was supported for microbes, suggesting that

ecological hypotheses developed based on evidence from macroorganisms also apply to explain

differences in microbial community assembly along latitudinal gradients[22].

Although the composition of phyllosphere bacterial communities varied greatly across

study sites along the latitudinal gradient, the different bacterial communities were interconnected

by a small proportion of hub bacterial ASVs to form large-scale plant-bacterial cooccurrence

networks. These hub ASVs were associated with a wide range of plant species in a broad

geographic space and include taxa such as *Methylorubrum* and *Sphingomonas* that have been

shown to play key roles in promoting plant health and productivity[10]. Additionally, the hub

bacterial taxa at the metacommunity level could affect large-scale plant-bacteria interactions: the

movement of hub taxa across local communities could mediate otherwise discrete ecological

processes within a metacommunity. For example, although plant species differing in growth-

mortality strategies hosted distinct leaf bacteria communities, the bacteria on one plant species

can affect that of other plant species through their influence on the metacommunity of shared

hub bacteria taxa[14,26]. Likewise, hub taxa could synchronize coevolutionary processes and

control the pace of host-bacteria coevolution at metacommunity scales through gene flow[37].

Thus, the spatial distribution of hub bacteria taxa should have a major impact on large-scale

patterns of plant-bacteria associations[26]. The distribution of hub bacterial ASVs was clustered

along the latitudinal gradient and was mainly controlled by climate, host attributes and spatial

factors. This suggests that climate change and plant species range shifts could change plant-

bacteria interactions through affecting the distribution of those hub taxa. Moreover, the influence

of host plant attributes on bacterial abundance was more important for hub taxa in temperate than

that in tropical forests, while spatial-related factors were more important for tropical than

temperate hub taxa, consistent with the observation of varying strengths of different processes

structuring bacterial diversity and community assembly along the latitudinal gradient. This result

emphasizes the importance of host selection in temperate regions and dispersal in the tropics to

explain the spatial distribution of hub bacteria taxa.

Understanding the response of plant microbiomes to global change and the consequent

impact on plant communities and ecosystem function requires large-scale studies on the diversity

and biogeography of plant-associated microbes[7,30,38]. Our results show for the first time that the

community assembly of plant leaf-associated bacterial communities is driven by processes

whose importance varies along a latitudinal gradient. A corollary of this finding is that global

change is likely to have different effects on plant leaf-microbe associations along the latitudinal

gradient, with tropical associations being more sensitive to climate warming and environmental

changes and temperate associations being more sensitive to changes in host plant abundance and

plant species range shifts. The direction of the global change effect could even reverse from

negative to positive from low latitudes to high latitudes, as recently shown for plant soil

microbes[39,40]. Host selection could result in distinct microbiome compositions that exert plant

species-specific impacts on host fitness. Although our study showed varying strength of host

selection across leaf microbiomes from temperate to tropical forests, it remains to be understood

whether the impact of plant leaf microbiomes on plant communities differs along latitude

gradients. Additional broad-scale comparative studies that take multiple factors into account will

be required to reveal the role of plant microbiomes in mediating the impact of global change on

plant communities and ecosystem function.

Methods

Study sites

We conducted the study at 10 forest sites across China with latitude ranging from 18.7°N

to 51.8°N (Table S2). The sites are part of a global network of stem-mapped forest plots of the

Forest Global Earth Observatory (ForestGEO http://forestgeo.si.edu/). For each site, we selected

approximately 1/3 of the tree species and around 3 individuals per species. The choice of these

species considered both shade tolerant and intolerant, early- and late-successional species, in

order to have a gradient in host attributes that may explain the variation in bacterial community

assemblages. In total, 1453 trees from 328 plant species of 148 genera and 59 families were

sampled.

Sampling and processing

For each tree individual, 50-100g leaves from the sub-canopy (2-10 m above ground)

were collected. Leaves were placed in a sterile plastic bag and washed using 100 mL of a 1:50

dilution of Redford buffer (1M Tris, 500 mM EDTA, 1.2% CTAB) within 24 hours after

sampling[14]. The suspension was collected and centrifuged at 3300 g for 25 minutes and the pellet

was added into an extraction tube to store in -20 °C until DNA extraction. Microbial DNA was

extracted following the manufacturer's protocol (QIAGEN Powersoil DNA extraction kit) and

the V5-V6 region of bacteria 16S rRNA genes was amplified using primers 799F and 1115R to

determine the composition of the phyllosphere bacterial community. PCR products were

normalized using a SequalPrep Normalization kit (Thermo Fisher Scientific), pooled and then

purified using AMPure to remove contaminants. The DNA library was prepared by mixing

equimolar concentrations of DNA from each sample, and then sequenced using Illumina MiSeq

Reagent Kit v3 for 300 bp paired-end sequencing.

Bioinformatics

We used DADA2 to identify the amplicon sequence variants (ASV) observed in each

sample[41]. The base quality scores (Q) were calculated, and the nucleotides at 19-240 bp and 16-

240 bp positions were kept for forward and reverse DNA sequences respectively. Then we

inferred error rates and inferred ASVs using DADA2 default parameters[41]. Paired read ends were

merged with a minimum overlap of 20. Non-target sequences and chimeras were removed.

Taxonomy was assigned to each ASV by comparison with the SILVA SSU r138 database.

We revealed around 116 million high-quality 16S rRNA gene sequences and identified

55693 amplicon sequence variants (ASVs) from 1453 samples. Sequence abundance per sample

ranged from 15485 to 522574, which could bias estimates of ASV diversity (Fig. S9a). Thus, we

randomly rarefied the ASV table to 15000 sequences per sample using the R package 'vegan',

and 49254 ASVs were retained after rarefaction. This eliminated only a small proportion of

extremely rare ASVs and had little impact on beta diversity (e.g. community dissimilarity; Fig.

S9b). Most ASVs (82%) were identified to the family level, 58% were classified to genus and

only 4% were identified to the species level.

Metadata

We collected metadata describing host plant attributes, abiotic environments,

neighborhood plants and spatial location. Plant phylogenetic tree was derived from a phylogeny

database for vascular plants according to the species identity using the *V.phyloMaker* function in

R [42]. Host plant identity and functional traits and abiotic environments including topographic and

edaphic variables were derived from the base data of each site. Climate variables including

annual mean temperature, annual precipitation, and precipitation seasonality were collected from

WorldClim database based on the geographical coordinates of sites. We calculated the richness,

abundance, mean phylogenetic distance and mean nearest neighbor distance of neighborhood

plants present within 10 meters around the sample, and included the local abundance of host

plant species in the metadata (Table S3).

Statistical analysis

We used R (version 3.6) for all statistical analyses. The diversity of phyllosphere bacteria

was modelled as a function of variables of host, environment, neighborhood plants and spatial

factors. To prepare these variables, Moran's Eigenvector Maps (MEM) was used to decompose

geographic distance matrices into a series of orthogonal eigenvectors representing the spatial

relationships among samples, and principal coordinates of neighbor matrices (PCNM) was

applied on the phylogenetic distance matrix among host plants to generate a series of variables

that representing the position of host plant in a multi-dimensional phylogenetical space. Both

phylogenetic and spatial eigenvectors can be used in regression to take phylogenetic/spatial

autocorrelation into account [43]. Environment, host traits and diversity of plant neighborhoods

variables were rescaled using the formula $(x - x_{min})/(x_{max} - x_{min})$ to standardize the effects of the

variables. Pairwise Pearson's correlation coefficients were calculated to test the correlation

among all variables, and for predictors with correlation coefficient > 0.9, one predictor was

randomly removed to reduce multicollinearity.

Species richness was calculated as the number of ASVs with abundance > 1, which

follows a negative binomial distribution (NBD). To model the effects of variables on richness,

we used generalized linear model with NBD as the error structure, implemented with function

glm.nb in the R package MASS. We first built a candidate set of generalized linear models to

represent the individual and joint linear contributions for types of mechanisms (host,

environment, neighborhoods and spatial proximity) using the predictors for each mechanism,

resulting in a total of 15 models. For each model, a stepwise selection procedure was used for

model selection based on Akaike's information criterion corrected for small sample size (AICc).

Non-significant variables ($P < 0.05$) were further excluded using the *update* function. Variance

inflation factor (VIF) was calculated and variables with VIF > 5 were also removed to avoid

multicollinearity. To quantify the individual effects of four types of variables and their

interactions on bacterial richness, we used variation partitioning where the independent effect of

each factor was evaluated as the difference in R^2 between models with and without this factor,

and the interactive effect was calculated by subtracting the independent effect from overall

effects. We used a subset of samples to evaluate the effect of host plant traits as the trait data are

available for most but not all samples (Table S3). The effect of plant traits was highly corrected

with that of plant phylogeny (Fig. S10). Thus, we only included plant phylogenetic eigenvectors

as host variables in the variation partitioning analysis.

We repeated the same variance partition analysis on bacterial composition. To do that,

the ASV table was Hellinger transformed to represent the community composition of bacteria.

We used redundancy analysis (RDA) to model the effect of those variables on community

composition. Variable selection was done for each model using forward selection implemented

with the function *forward.sel* in R package packfor following the recommended stopping rules,

and variation partitioning was implemented by *varpart* function in the 'vegan' package [44].

The analyses of richness and community composition were conducted at local scales for

the 10 sites, at regional scales for tropical, subtropical and temperate forests, and at the global

scale by combining all the sites. Specifically, Heishiding (HSD) and Dinghushan (DHS) sites

(located at 23.27°N, 23.17°N respectively) were on the northern boundary of the tropics.

Although classified as subtropical sites, the two forests are more similar in vegetation

composition and climate conditions with the tropical forest Jianfengling (JFL, 18.73°N) than

those typical subtropical forests (Table S6). We thus included HSD and DHS as 'tropical' sites,

resulting in 3 tropical, 4 subtropical and 3 temperate sites. The relative importance of each factor

and their interactive effects were evaluated as the proportion of explained variance, which was

further related to latitude using a linear regression.

To construct networks that represent plant-bacteria associations at different geographical

scales, we first transformed ASV abundances to presence-absence with a threshold of 15 reads

per sample (0.1% of the total read count of each sample) to be considered present, in order to

minimize the effects of extreme rare species and sequencing errors on network construction. The

occurrence of ASV in samples was then converted to occurrences on plant species according to

the plant host identity of each sample, where a cell entry represented the number of leaf samples

that were observed to have the ASV occurring on a plant species. In this step, the same plant

species from different sites were represented in different rows as we were interested in how

metacommunity bacteria interconnect local plant-bacteria networks across sites. The species-

ASV table comprises 490 rows and 12065 columns, and the binary (presence/absence) data of

this table were used to construct the network. We constructed 10 local networks representing

plant-bacteria associations at site level, 3 regional networks for temperate, subtropical and

tropical regions, and a global network using 'igraph' package in R. The regional and global

networks were visualized using program GePhi with the 'ForceA-tlas2' layout algorithm, where

plant species from different sites were highlighted using warm colors. For these networks, we

identified hub ASVs in core positions of the networks by calculating the betweenness centrality

score of each node/ASV. Betweenness centrality was standardized using the formula (x –

x_{min})/(x_{max} – x_{min}) to make it range from 0 to 1, and ASVs with scaled betweenness above 0.5

were recognized as hubs[26]. A total of 115 ASVs were identified as metacommunity-level hubs at

regional and/or global scales.

The biogeography of the hub ASVs was first explored using a heatmap that mapped the

frequency of ASVs (the proportion of samples in which the ASVs occurred) for each site. We

then applied a joint species distribution modeling approach, called Hierarchical Modelling of

Species Composition (HMSC)[45], to predict the abundance of the 115 hub ASVs at the global

scale. We selected a set of habitat variables that indicate important mechanisms potentially

structuring bacterial community assembly, including 3 soil variables (total nitrogen, total

phosphorus and water content), 3 climate variables (temperature, precipitation and

isothermality), and 3 plant leaf traits (nitrogen, phosphorus and specific leaf area). As plant trait

data were not available for all samples, 1242 out of 1453 samples with available plant trait data

were used in this analysis. The 9 variables were treated as fixed terms to evaluate the effect of

habitat conditions on ASVs, and two distance matrices representing the spatial and host

phylogenetic relationships between samples were used as random terms to control for the effects

of spatially and phylogenetically clustered data.

We applied a hurdle model, i.e., one model for presence-absence and another one for

abundance conditional on presence, in order to quantify the drivers of both presence-absence and

abundance of hub ASVs. We applied probit regression in the presence-absence model, and linear

regression for transformed count data in the abundance model. The count data were transformed

by declaring zeros as missing data, log-transforming, and then scaling the data to zero mean and

unit variance within each species[45]. We fitted the HMSC model with the R-package Hmsc

assuming the default prior distributions. We sampled the posterior distribution with five Markov

Chain Monte Carlo (MCMC) chains, each of which was run for 37,500 iterations, of which the

first 12,500 were removed as burn-in. We examined MCMC convergence by examining the

potential scale reduction factors of the model parameters. The MCMC convergence of our model

was satisfactory: the potential scale reduction factors for the β-parameters (that measures the

responses of the species to environmental covariates) were on average 1.013 and 1.04 for

abundance and presence-absence model respectively. To compare the species distribution

patterns of ASV among regions, we calculated the relative importance of each mechanism: the

fixed effects of host traits, soil condition and climate, and the random effects of space and host

phylogeny, which were compared among temperate, subtropical and tropical hubs using paired

samples Wilcoxon tests.

References

1. Gilbert, J. A. & Neufeld, J. D. Life in a world without microbes. *Plos Biol* **12**, e1002020 (2014).

2. Berendsen, R. L., Pieterse, C. M. J. & Bakker, P. A. H. M. The rhizosphere microbiome and plant health. *Trends Plant Sci* **17**, 478–486 (2012).

3. Liu, H., Brettell, L. E. & Singh, B. Linking the phyllosphere microbiome to plant health. *Trends Plant Sci* **25**, 841–844 (2020).

4. Linde, S. van der *et al.* Environment and host as large-scale controls of ectomycorrhizal fungi. *Nature* **558**, 243–248 (2018).

5. Martin, F. M., Uroz, S. & Barker, D. G. Ancestral alliances: Plant mutualistic symbioses with fungi and bacteria. *Science* **356**, (2017).

6. Vorholt, J. A. Microbial life in the phyllosphere. *Nat Rev Microbiol* **10**, 828–840 (2012).

7. Cavicchioli, R. *et al.* Scientists' warning to humanity: microorganisms and climate change. *Nat Rev Microbiol* **17**, 569–586 (2019).

8. Perreault, R. & Laforest-Lapointe, I. Plant-microbe interactions in the phyllosphere: facing challenges of the anthropocene. *Isme J* **16**, 339–345 (2022).

9. Lindow, S. E. & Leveau, J. H. J. Phyllosphere microbiology. *Curr Opin Biotech* **13**, 238–243 (2002).

10. Chen, T. *et al.* A plant genetic network for preventing dysbiosis in the phyllosphere. *Nature* **580**, 653–657 (2020).

11. Zamioudis, C. & Pieterse, C. M. J. Modulation of host immunity by beneficial microbes. *Mol Plant-microbe Interactions* **25**, 139–150 (2012).

12. Moyes, A. B. *et al.* Evidence for foliar endophytic nitrogen fixation in a widely distributed subalpine conifer. *New Phytol* **210**, 657–668 (2016).

13. Vacher, C. *et al.* The phyllosphere: microbial jungle at the plant–climate interface. *Annu Rev Ecol Evol Syst* **47**, 1–24 (2016).

14. Kembel, S. W. *et al.* Relationships between phyllosphere bacterial communities and plant functional traits in a neotropical forest. *Proc National Acad Sci* **111**, 13715–13720 (2014).

15. Lajoie, G. & Kembel, S. W. Plant-bacteria associations are phylogenetically structured in the phyllosphere. *Mol Ecol* **30**, 5572–5587 (2021).

16. Laforest-Lapointe, I., Messier, C. & Kembel, S. W. Host species identity, site and time drive
temperate tree phyllosphere bacterial community structure. *Microbiome* **4**, 27 (2016).

17. Laforest-Lapointe, I., Paquette, A., Messier, C. & Kembel, S. W. Leaf bacterial diversity
mediates plant diversity and ecosystem function relationships. *Nature* **546**, 145–147 (2017).

18. Lajoie, G. & Kembel, S. W. Host neighborhood shapes bacterial community assembly and
specialization on tree species across a latitudinal gradient. *Ecol Monogr* (2021)
doi:10.1002/ecm.1443.

19. Leducq, J.-B. *et al.* Fine-scale adaptations to environmental variation and growth strategies
drive phyllosphere methylobacterium diversity. *Mbio* **13**, e03175-21 (2022).

20. Finkel, O. M. *et al.* Distance-decay relationships partially determine diversity patterns of
phyllosphere bacteria on tamrix trees across the sonoran desert. *Appl Environ Microb* **78**, 6187–
6193 (2012).

21. Magan, N. & Baxter, E. Effect of increased CO_2 concentration and temperature on the
phyllosphere mycoflora of winter wheat flag leaves during ripening. *Ann Appl Biol* **129**, 189–195
(1996).

22. Qiao, X., Jabot, F., Tang, Z., Jiang, M. & Fang, J. A latitudinal gradient in tree community
assembly processes evidenced in Chinese forests. *Global Ecol Biogeogr* **24**, 314–323 (2015).

23. Dyer, L. A. *et al.* Host specificity of Lepidoptera in tropical and temperate forests. *Nature*
448, 696–699 (2007).

24. Banerjee, S., Schlaeppi, K. & Heijden, M. G. A. van der. Keystone taxa as drivers of
microbiome structure and functioning. *Nat Rev Microbiol* **16**, 567–576 (2018).

25. Agler, M. T. *et al.* Microbial hub taxa link host and abiotic factors to plant microbiome
Variation. *Plos Biol* **14**, e1002352 (2016).

26. Toju, H. *et al.* Species-rich networks and eco-evolutionary synthesis at the metacommunity
level. *Nat Ecol Evol* **1**, 0024 (2017).

27. Leibold, M. A. *et al.* The metacommunity concept: a framework for multi-scale community
ecology. *Ecol Lett* **7**, 601–613 (2004).

28. Thompson, J. N. The Geographic mosaic of coevolution. (2005)
doi:10.7208/chicago/9780226118697.001.0001.

29. Toju, H., Tanabe, A. S. & Sato, H. Network hubs in root-associated fungal metacommunities.
Microbiome **6**, 116 (2018).

30. Martiny, J. B. H. *et al.* Microbial biogeography: putting microorganisms on the map. *Nat Rev*
Microbiol **4**, 102–112 (2006).

31. Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. H. Beyond
biogeographic patterns: processes shaping the microbial landscape. *Nat Rev Microbiol* **10**, 497–
506 (2012).

32. Jr, P. R. G., Jordano, P. & Thompson, J. N. Evolution and coevolution in mutualistic
networks. *Ecol Lett* **14**, 877–885 (2011).

33. Wang, Z. *et al.* Effects of host phylogeny, habitat and spatial proximity on host specificity
and diversity of pathogenic and mycorrhizal fungi in a subtropical forest. *New Phytol* **223**, 462–
474 (2019).

34. Lajoie, G. & Parfrey, L. W. Beyond specialization: re-examining routes of host influence on
symbiont evolution. *Trends Ecol Evol* **37**, 590–598 (2022).

35. Semchenko, M. *et al.* Deciphering the role of specialist and generalist plant–microbial
interactions as drivers of plant–soil feedback. *New Phytol* (2022) doi:10.1111/nph.18118.

36. Gravel, D., Canham, C. D., Beaudet, M. & Messier, C. Reconciling niche and neutrality: the
continuum hypothesis. *Ecol Lett* **9**, 399–409 (2006).

37. Urban, M. C. *et al.* The evolutionary ecology of metacommunities. *Trends Ecol Evol* **23**,
311–317 (2008).

38. Brunel, C. *et al.* Towards unraveling macroecological patterns in rhizosphere microbiomes.
Trends Plant Sci **25**, 1017–1029 (2020).

39. Bachelot, B. *et al.* Altered climate leads to positive density-dependent feedbacks in a tropical
wet forest. *Global Change Biol* **26**, 3417–3428 (2020).

40. Liu, Y. & He, F. Warming intensifies soil pathogen negative feedback on a temperate tree.
New Phytol **231**, 2297–2307 (2021).

41. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat Methods* **13**, 581–583 (2016).

42. Jin, Y. & Qian, H. V.PhyloMaker: an R package that can generate very large phylogenies for
vascular plants. *Ecography* **42**, 1353–1359 (2019).

43. Dray, S., Legendre, P. & Peres-Neto, P. R. Spatial modelling: a comprehensive framework
for principal coordinate analysis of neighbour matrices (PCNM). *Ecol Model* **196**, 483–493
(2006).

44. Peres-Neto, P. R., Legendre, P., Dray, S. & Borcard, D. Variation partitioning of species data
matrices: estimation and comparison of fractions. *Ecology* **87**, 2614–2625 (2006).

45. Ovaskainen, O. *et al.* How to make more out of community data? A conceptual framework
and its implementation as models and software. *Ecol Lett* **20**, 561–576 (2017).

520 Author contribution

521 Z.W., F.H. and S.W.K. conceived the project; Z.W. conducted field sampling and phyllosphere

522 data collection; Y.J., M.Z., C.C., Y.C., S.F., G.J., M.J., J.L., Y.L., Y.L., K.M., X.M., X.Q., X.W.,

523 X.W., H.X., W.Y., L.Z., Y.Z. collected and collated the metadata and assisted with fieldwork;

524 Z.W. analyzed data with supported from S.W.K.; Z.W. drifted the manuscript with input from

525 S.W.K. and F.H.; All authors contributed to editing of the manuscript.

527 Competing interests

528 The authors declare no competing interests.

530 Materials & Correspondence

Correspondence and requests for materials should be addressed to Zihui Wang

(wang.zihui@courrier.uqam.ca), Fangliang He (fhe@ualberta.ca) or Steven. W. Kembel

(kembel.steven_w@uqam.ca)

Figure 1 The latitude patterns of phyllosphere bacterial (a) diversity, (b) community

composition and (c) occurrence of metacommunity-level hub ASVs. First- and second order

polynomial fits are shown in black (dashed) and red (solid) respectively in (a), Bacterial

community composition was visualized using nonmetric multidimensional scaling ordination

with each ellipse representing site in (b), where arrows inside indicate significant variables

correlated with ordination scores and the explained power (R^2) of the correlation. Heatmap in (c)

show the relative frequency of hub ASVs clustering regarding to (1) temperate hubs (2)

subtropical hubs (3) tropical hubs, (4) hubs that connect temperate and subtropical, (5) hubs that

connect subtropical and tropical sites, and (6) global hubs that connect all sites.

Figure 2 Map of study sites indicating the percentage of variation in community composition

explained by host, abiotic environment, neighborhood plants and spatial factors at local scales

for each site, regional scales for temperate, subtropical, and tropical sites, and a 'global' scale

that includes all sites. The Venn diagrams show the absolute variance explained by each factor

and the histograms on the right of Venn diagrams show the relative contribution of the

independent and shared effects of four variables to the explained variance. Explained variation

less than 1% was not shown in the Veen diagram. Abbreviation; H: host, E: abiotic

environments; S: spatial variables; N: neighborhood plant communities.

Figure 3 The relative importance of different mechanisms in structuring phyllosphere bacterial

community assembly varies across latitude gradients (*a-d*) and across geographic scales (*e*).

Coefficients of linear regression between the relative importance of drivers and latitude were

reported in *a-d*, and Kruskal-Wallis rank sum test was applied to test the variation across three

scales in *e*, showing moderate evidence (p=0.046) for increased confounding effects at larger

scales (sample size: 10, 3 and 1 for local, regional and global scales).

Figure 4 Global and regional networks representing the associations between plants and

phyllosphere bacteria. Circles represent either bacteria ASVs (dark blue) or plant species (warm

colors). A link was established if the plant species was associated with more than 15 reads of an

ASV (see Methods). Size of the circles indicates the betweenness centrality of nodes. Plant

species were colored according to latitude of site, including tropical: JFL, DHS, HSD;

subtropical: BSZ, GTS, BDGS, TT; and temperate sites: DLS, CBS, LS.

Figure 5 The results of joint species distribution modeling, showing (a) the absolute variation in

bacterial abundance explained by each factor, the relative contribution of (b) host plant traits and

(c) spatial factors on determining the abundance of ASVs in different regions. 115 ASVs were

included in the model, and 9 habitat covariates and two distance matrices representing spatial

and host phylogenetic relationship among samples were used as fixed and random term

respectively. Each bar represents a hub bacterial ASV sorted according to the total variance

explained, and the caption above the bar plot shows the averaged variation explained by each

factor.

Figure 6 The contribution of host attributes (a) and spatial-related factors (b) to the abundance of

hub taxa varies among regions. Numbers above the paired boxes symbolize the p values of

pairwise Wilcoxin tests.