1   **The Functional Landscape of SARS-CoV-2 3CL Protease**

4   Sho Iketani[1,2]*, Seo Jung Hong[3]*, Jenny Sheng[3,4]*, Farideh Bahari[5]*, Bruce

5   Culbertson[4,6], Fereshteh Fallah Atanaki[5], Arjun K. Aditham[7], Alexander F. Kratz[3,4],

6   Maria I. Luck[1,8], Ruxiao Tian[2], Stephen P. Goff[1,2,9], Hesam Montazeri[5], Yosef Sabo[1,8],

7   David D. Ho[1,2,8], Alejandro Chavez[3]^

9
[1] Aaron Diamond AIDS Research Center, Columbia University Vagelos College of

Physicians and Surgeons, New York, NY, USA

[2] Department of Microbiology and Immunology, Columbia University Vagelos College of

Physicians and Surgeons, New York, NY, USA

[3] Department of Pathology and Cell Biology, Columbia University Vagelos College of

Physicians and Surgeons, New York, NY, USA

[4] Integrated Program in Cellular, Molecular, and Biomedical Studies, Columbia

University Irving Medical Center, New York, NY, USA

[5] Department of Bioinformatics, Institute of Biochemistry and Biophysics, University of

Tehran, Tehran, Iran

[6] Medical Scientist Training Program, Columbia University Irving Medical Center, New

York, NY, USA

[7] Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

[8] Division of Infectious Diseases, Department of Medicine, Columbia University Vagelos

College of Physicians and Surgeons, New York, NY, USA

[9] Department of Biochemistry and Molecular Biophysics, Columbia University Vagelos

College of Physicians and Surgeons, New York, NY, USA

27
28   *These authors contributed equally

29   ^Address correspondence to Alejandro Chavez (ac4304@cumc.columbia.edu, 212-305-

30   1150, 650 West 168th St, BB1208, New York, NY 10032)

31
32   Character count: 26614
33
34

Summary

SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) as the etiologic agent of COVID-19 (coronavirus disease 2019) has drastically altered life globally. Numerous efforts have been placed on the development of therapeutics to treat SARS-CoV-2 infection. One particular target is the 3CL protease (3CL^pro), which holds promise as it is essential to the virus and highly conserved among coronaviruses, suggesting that it may be possible to find broad inhibitors that treat not just SARS-CoV-2 but other coronavirus infections as well. While the 3CL protease has been studied by many groups for SARS-CoV-2 and other coronaviruses, our understanding of its tolerance to mutations is limited, knowledge which is particularly important as 3CL protease inhibitors become utilized clinically. Here, we develop a yeast-based deep mutational scanning approach to systematically profile the activity of all possible single mutants of the SARS-CoV-2 3CL^pro, and validate our results both in yeast and in authentic viruses. We reveal that the 3CL^pro is highly malleable and is capable of tolerating mutations throughout the protein, including within the substrate binding pocket. Yet, we also identify specific residues that appear immutable for function of the protease, suggesting that these interactions may be novel targets for the design of future 3CL^pro inhibitors. Finally, we utilize our screening results as a basis to identify E166V as a resistance-conferring mutation against the therapeutic 3CL^pro inhibitor, nirmatrelvir, in clinical use. Collectively, the functional map presented herein may serve as a guide for further understanding of the biological properties of the 3CL protease and for drug development for current and future coronavirus pandemics.

Main text

SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) has transformed daily

life around the globe and caused widespread socioeconomic damage (Wu et al., 2020;

Zhou et al., 2020). To curtail the ongoing pandemic, researchers have intensively

studied SARS-CoV-2 in hopes of identifying effective avenues for therapeutic

intervention. To that end, several antibodies and antivirals have been identified and

have consequently been authorized for use in COVID-19 (coronavirus disease 2019)

patients (Beigel et al., 2020; Chen et al., 2021b; Consortium et al., 2021; Gupta et al.,

2021; Hammond et al., 2022; Jayk Bernal et al., 2022; Weinreich et al., 2021). Current

treatments are directed towards three viral targets: the spike glycoprotein (monoclonal

antibodies), the RNA-dependent RNA polymerase (remdesivir and molnupiravir), and

the 3-chymotrypsin-like protease ($3CL^{pro}$) (nirmatrelvir). Despite the promise of these

therapies, previous experiences with the rapid evolution of viruses suggest that

resistance is likely to quickly arise (Strasfeld and Chou, 2010). Indeed, spike variants

with enhanced resistance against antibody neutralization underscores this unfortunate

possibility (Iketani et al., 2022; Liu et al., 2022b; Planas et al., 2021; Wang et al., 2021a;

Wang et al., 2021b).

Furthermore, studies have demonstrated that adaptation of the virus to

remdesivir-mediated polymerase inhibition can be readily achieved both *in vitro* and *in*

vivo, suggesting that despite the conserved nature of a viral target, resistance can still

arise (Gandhi et al., 2022; Stevens et al., 2022; Szemiel et al., 2021). While no reports

yet exist for SARS-CoV-2 $3CL^{pro}$ resistance, examples of resistance to protease

inhibitors in the case of HIV-1 and HCV foreshadow this likely event for the coronavirus

(Clavel and Hance, 2004; Strasfeld and Chou, 2010).

The 3CL^{pro} is an attractive therapeutic target as it is essential for viral replication

and there is a history of using protease inhibitors to successfully treat other viral

illnesses (Chen and Njoroge, 2009; Ghosh et al., 2016; Jin et al., 2020a). To date,

numerous SARS-CoV-2 3CL^{pro} inhibitors have been reported, with a few advancing into

clinical trials, and one (nirmatrelvir; utilized in combination with ritonavir and marketed

as PAXLOVID™ by Pfizer) being authorized for clinical use to mitigate against severe

disease and deaths (Amporndanai et al., 2021; Boras et al., 2021; Dai et al., 2020; Fu et

al., 2020; Hammond et al., 2022; Hattori et al., 2021; Iketani et al., 2021; Jin et al.,

2020a; Jin et al., 2020b; Ma et al., 2020; Owen et al., 2021; Qiao et al., 2021; Vuong et

al., 2020; Zhang et al., 2020). Such drug discovery efforts are enticing, particularly as

the 3CL^{pro} exhibits significant conservation across coronaviruses, suggesting that

broad-spectrum or ready-made "off-the-shelf" antivirals may be achievable for this and

future epidemics. Indeed, several 3CL^{pro} inhibitors developed for other coronaviruses

have been found to also hold activity against SARS-CoV-2 3CL^{pro} (Iketani et al., 2021;

Resnick et al., 2021; Vuong et al., 2020). However, the tolerance of the 3CL^{pro} to

mutations remains unknown; such knowledge could allow for the rational design of

inhibitors with improved potency and breadth, as well as providing insight into the

biology of the enzyme.

Here, we sought to systematically evaluate the plasticity of SARS-CoV-2 3CL^{pro}

by interrogating the activity of all possible point mutants. We speculated that such a

comprehensive analysis would reveal the malleability of the enzyme, as well as uncover

conserved sites that could be used to rationally design inhibitors that are resistant to

viral escape. Such extensive profiling of mutants of a gene, also known as deep

mutational scanning (DMS), has been conducted successfully for the spike gene of

SARS-CoV-2 by several groups, revealing the impact of mutations on expression,

binding to its cognate receptor angiotensin converting enzyme 2 (ACE2), and binding by

antibodies, underscoring the applicability of this approach (Garrett et al., 2021; Garrett

et al., 2022; Greaney et al., 2021a; Greaney et al., 2021b; Starr et al., 2021; Starr et al.,

2020).

To conduct DMS, a scalable assay that can report on the properties under study

must first be designed (Fowler and Fields, 2014). We chose to work with

Saccharomyces cerevisiae (budding yeast) due to their ease of manipulation,

successful use in previous SARS-CoV-2 DMS studies, and similarity to humans with

regard to their folding, post-translational modification, and turnover of proteins (Botstein

and Fink, 2011; Greaney et al., 2021a; Greaney et al., 2021b; Starr et al., 2021; Starr et

al., 2020). Previous reports have identified that exogenous expression of viral proteases

in cells can lead to cellular toxicity (Blanco et al., 2003; Resnick et al., 2021). We

speculated that a similar approach could be used with SARS-CoV-2 3CL^{pro} in yeast, and

so we cloned the SARS-CoV-2 3CL^{pro} gene into an inducible yeast expression vector

and generated a yeast strain containing this construct. As compared to cells containing

a non-toxic control protein, enhanced yellow fluorescent protein (EYFP), cells

expressing the SARS-CoV-2 3CL^{pro} showed a significant growth deficiency (**Fig. S1A**).

To determine if the observed decrease in growth was due to the activity of the protease

or a cryptic function performed by the exogenously expressed protease, we generated a

catalytically inactive C145A mutant (Lee et al., 2020; Resnick et al., 2021). When

expressed in yeast, the C145A mutant grew similarly to the non-toxic EYFP control,

suggesting that the observed toxicity was a direct effect of 3CL^{pro} activity (**Fig. S1A**).

This growth defect upon expressing the 3CL^{pro} in yeast formed the basis of our

assay and suggested that we could profile libraries of 3CL^{pro} mutants by growing them

within a mixed pool and analyzing their relative abundances before and after inducing

their expression. In this paradigm, we expected active variants to deplete and inactive

variants to enrich within the pool upon induction. To generate our library of 3CL^{pro}

variants, we used degenerate oligonucleotides coupled with a site-directed mutagenesis

protocol (see **Methods** for details). Saturation mutagenesis of each residue was

conducted by designing a single primer containing an NNK stretch, which generates all

possible amino acid codings for the targeted residue. We also introduced into each

primer a degenerate codon(s) upstream and/or downstream of the NNK stretch. This

additional site(s) of degeneracy enabled the same amino acid mutation to be

represented by a much larger number of DNA sequences than in conventional

mutagenesis approaches, allowing us to average across the various DNA codings and

by doing so better control for the noise inherent with assays performed at this scale

(Schmierer et al., 2017; Zhu et al., 2019) (**Fig. S1B**). Once the mutant plasmids were

generated, they were transformed into yeast, and cells were grown with or without

induction of protease expression, followed by analysis of the abundance of the various

mutants under each condition.

We conducted the DMS in biological duplicate for each of the induced and

uninduced conditions. Upon observing robust correlation between the replicates, we

proceeded to aggregate the effects between the two replicates, as done before (**Fig.**

S1C) (Starr et al., 2020). The representation of mutations at each residue was as

expected, with mostly non-synonymous mutations created by the site-directed

mutagenesis method employed (**Fig. S1D**). With regard to overall coverage, we

observed most variants, with 94.3% (6060 of 6426 possibilities) of all possible variants

assessed for activity.

To determine the activity of each SARS-CoV-2 3CL^{pro} variant, we compared its

enrichment in the induced versus uninduced conditions, and calculated an activity score

(see **Methods** for details). We note that these scores by definition incorporate not just

the enzymatic activity of the protease, but additional factors, such as their stability and

expression. These activity scores were then normalized to the wild-type and stop coding,

set at 0 and -1, respectively, and visualized as a heatmap (**Fig. 1A**). Significant

heterogeneity in tolerance to mutation across the protein was observed, with some

specific residues and regions appearing highly constrained and intolerant to mutation

(i.e., mutations resulted in loss of activity).

As an initial validation of our data, we first examined known characteristics of the

3CL^{pro}, either in SARS-CoV-2 or in its close relative, SARS-CoV. The 3CL^{pro} can be

structurally separated into the N-finger (residues 1-7), domains I (residues 8-100) and II

(residues 101-183), a substrate binding loop (SBL) connecting domains II and III

(residues 184-199), and domain III (residues 200-306) (Tahir Ul Qamar et al., 2020).

Looking at specific residues, mutations to the catalytic dyad, His41 and Cys145,

resulted in loss of activity, as would be expected given their essential role in protease

activity (Kneller et al., 2020; Lee et al., 2020). Additional critical residues within the

active site were also generally conserved. These included Asp187, which stabilizes

His41, and Arg40, which forms a salt bridge with Asp187 to stabilize its position (Kneller

et al., 2020; Suarez and Diaz, 2020). Residues previously shown to directly bind one of

the natural substrates of the 3CL^{pro}, such as Phe140, Glu143, His163, His164, and

Gln192, were mostly conserved and intolerant to mutations, although other substrate

binding residues such as Glu166 and Gln189 were not (Lee et al., 2020; Zhao et al.,

2022). These latter nonconserved residues have previously been implicated in

conferring substrate flexibility for SARS-CoV 3CL^{pro}, and therefore may tolerate

mutation (Muramatsu et al., 2016; Zhao et al., 2022). These observations suggested

that our data recapitulated known interactions and could also reveal complexities in the

properties of the protease.

To further validate the data, we curated a list of mutations within the 3CL^{pro} that

have been identified within patient samples and mapped them onto our DMS data

(Chen et al., 2021a; Shu and McCauley, 2017). We expected that clinical variants

should have high activity scores within our dataset, as viruses with inactive proteases

would not be expected to replicate sufficiently to be found in clinical isolates. We

identified 932 single mutants from clinical samples (with ≥3 occurrences), and found

that all of these mutants clustered around the wild-type score of 0, suggesting that our

data were in line with what is observed in natural infections (**Fig. 1B**).

Although the 3CL^{pro} is largely conserved, there is still variability that can be

observed throughout coronavirus 3CL proteases. We speculated that residues that are

less conserved across evolution should overlap with mutation-tolerant sites as denoted

by our DMS data. To investigate this, we utilized the ConSurf server to determine the

rate of evolution for each residue, with residues that were poorly conserved being

scored as having a high rate of evolution (Glaser et al., 2003; Landau et al., 2005). We

then grouped residues as having a high, medium, or low rate of evolution, and plotted

the average activity score for each of the residues in the various groups. As expected,

residues with high or low rates of evolution were generally tolerant or intolerant to

mutation, respectively (**Fig. 1C**). Of note, while we do observe an association between

these two metrics, there are additional factors that influence evolutionary conservation,

such as compensatory or enabling mutations that were not considered in our screening

and likely limit the strength of our association.

Finally, we considered whether our data aligned with the known structural

features of SARS-CoV-2 3CL^{pro}. When we mapped the average activity score at each

position onto the crystal structure of the protein, we observed that surface-exposed

residues were significantly more tolerant to mutations than buried residues (**Fig. 1D**).

Furthermore, regions of importance, such as those forming the catalytic pocket or the

dimerization interface, strongly stood out as regions that were intolerant of mutations.

Collectively, the above analyses suggest that our data are robust and reflect true

biological properties of the protease.

We next sought to further validate our data by investigating specific mutants

indicated to be active or inactive by our screen. We chose 24 mutants throughout the

gene with varying scores (14 predicted to be active and 10 predicted to be inactive), and

produced isogenic yeast strains for each mutation. Each mutant was then individually

tested for protease activity, and all of the mutations were found to behave as predicted

from our DMS (**Fig. 2A**). Unexpectedly, various C-terminal truncations, which lack

residues that are believed to be critical for protein dimerization, exhibited activity (Hsu et

al., 2005). To verify this observation, we also generated an independent strain with a

truncated 3CL^pro variant (deletion of residues 301-306). In agreement with our screening

data, this truncated mutant showed protease activity (**Fig. S2A**). We note that such a

mutant is likely inviable in nature in which the C-terminal residues are critical for release

of the 3CL protease from the polyprotein, whereas the direct expression of the protease

in our system circumvents the need for polyprotein processing.

While the correlation of our DMS data with clinical sequencing of SARS-CoV-2

isolates suggests that our high-throughput approach captures the true tolerance of the

protease to mutation, as a further validation, we tested several mutants, as well as WT

and C145A as controls, in the context of authentic SARS-CoV-2 virus. To generate our

viral mutants, we used an established reverse genetics system, and introduced into it a

subset of our active and inactive variants and attempted to infect Vero E6 cells (Ye et al.,

2020). We found that testing within the context of the virus recapitulated the screening

results, with the mutations predicted to retain an active protease resulting in a fully

viable virus, whereas the mutations predicted to inactivate the protease led to inviable

virus, with the exception of one mutant, E288F (**Fig. 2B**). This mutant was scored to be

inactive in the screen and validated as such within yeast, yet exhibited activity within the

recombinant virus. This difference in phenotype may be due to the slightly different

environmental conditions between yeast and mammalian cells as well as the differences

in substrates in the two contexts. Nevertheless, 14 of the 15 SARS-CoV-2 strains tested

led to comparable results between the yeast and virus assays, suggesting that our

screening largely recapitulates what is observed in a functional SARS-CoV-2 virus (**Fig.**

2A-B).

For the 10 inactive protease variants that were independently validated, we

wondered if a loss of 3CL^{pro} dimerization, protein instability, or deficient catalytic activity

might explain their lack of function. To address the question of dimerization, we

designed a yeast two-hybrid assay to test whether the mutations were still competent

for dimerization and found that seven of the 10 could still dimerize at varying levels,

suggesting that a loss of enzymatic activity, reduction in dimerization efficiency, or a

combination of both could explain the observed phenotypes (**Fig. S2B**). For the three

that no longer seemed to be able to dimerize, we tested protein expression levels by

immunoblotting, observing that two of the three (L250G and L268N) were poorly

detected, suggesting that they formed unstable proteins. The final variant, D295G, was

well-expressed, and hence its lack of signal in the yeast two-hybrid assay suggests it to

be a dimerization-defective mutant, although additional experiments such as analytical

ultracentrifugation are required to further investigate this mutant's dimerization

properties (**Fig. S2C**). These results suggest that despite the multiple mechanisms by

which the protease can be inactivated, our DMS robustly reflects the protease activity,

regardless of the nature of the mutation.

Altogether, we believe that the activity values we derived for each mutant are

reflective of their true properties given the correlations we observed with clinical data,

evolutionary data, and structure, in combination with the validation of individual

mutations, both in yeast and authentic viruses. Therefore, we speculated that by

examining residues defined to be invariant in our screen, we could identify critical

interactions within the protease that could serve as targets for the next generation of

protease inhibitors.

As numerous sites appeared to have varying tolerance to mutation, we focused

our subsequent structural analysis on the small subset that were highly intolerant to

mutations. Upon analyzing these residues within a crystal structure of SARS-CoV-2

3CL^{pro} complexed with its physiological substrate (the C-terminal autocleavage

sequence, Ser301 to Gln306), we found that these sites formed critical interactions with

their adjacent neighbors in three-dimensional space, both within the same or different

domains of the protein, or across protease protomers (Lee et al., 2020). Within domain I,

several residues likely contribute to the stabilization of the protein structure. As shown in

Fig. 2C, the side chains of Asn95 and Cys16 form H-bonds with the backbones of

Trp31 and Lys12, respectively. Backbone hydrogen bonds are also identified between

Val13 and Cys16, and Met17 and Trp31. In addition, the side chains of Asn95 and

Trp31 interact favorably via the π electrons of their amide and aromatic moieties,

respectively. As such, Lys12, Val13, Cys16, Met17, Trp31, and Asn95 contribute to the

critical intra-domain interactions to stabilize the tertiary structure within domain I, which

may explain the loss of activities when these residues are mutated. Examining the

interface between domain I and domain II (**Fig. 2D**), the side chains of Val13, Leu115,

Phe150, Pro122, and Pro9' of the opposing protomer form strong hydrophobic

interactions. Mutating these residues to larger or more polar ones causes a reduction in

protease activity, suggesting they are important inter-domain anchors. At the dimer

interface between domain I and I' (**Fig. 2E**), an extensive hydrogen bond network is

observed among Ser10, Gly11, Glu14 and Ser10', Gly11', Glu14' from the second

protomer of the homodimer. These same residues have been shown to be critical for

dimerization in SARS-CoV 3CL^{pro} (Chen et al., 2008a; Chen et al., 2008b). Looking at

the catalytic domain II with a focus on the P1 pocket of the substrate binding site (**Fig.**

2F), besides the catalytic Cys145, residues including Ser144 and His163 form a

multitude of hydrogen bond interactions with the substrate. The side chains of Phe140

and His163 also form a π-π stacking interaction. Moreover, Phe140 interacts with S1' in

the N-finger domain of the neighboring protomer via hydrogen bonds. This network of

interactions affords the structural stability near the substrate binding site, which could

explain the sensitivities of the enzymatic activities to the mutations in this region.

Consistently, these residues have previously been shown to be important for SARS-

CoV 3CL^{pro} catalytic activity (Barrila et al., 2006; Cheng et al., 2010; Hu et al., 2009;

Huang et al., 2004; Tan et al., 2005). During close examination of our DMS data and the

interface between domain II and domain III, we observed a novel interaction which our

data suggest may help to stabilize this interface (**Fig. 2G**). Specifically, residues Gly109

and Thr111 of domain II form hydrogen bonds with Asn203 and Asp295 of domain III,

respectively. Previous studies have implicated the importance of the N-finger in

dimerization, and we observed similar results (Chen et al., 2008b; Chou et al., 2004; Hu

et al., 2009; Wei et al., 2006). The N-finger folds between domains II, III, and II' of the

abutting protomer (**Fig. 2H-I**). The aromatic side chain of Phe3 is completely

encapsulated in the hydrophobic pocket composed of side chains of Trp207, Ala210,

Phe291, and Val296 (**Fig. 2H**). As such, the biological activity of SARS-CoV-2 3CL^{pro} is

largely preserved when Phe3 is mutated to other hydrophobic residues including

isoleucine, leucine, methionine, and tryptophan. However, mutations to residues of

polar side chains completely abolish protein activity, corroborating the structural role of

the hydrophobic encapsulation. The residue Ala7 also engages interactions at the

interface of N-finger, domain II and II' (**Fig. 2I**). Hydrogen bonds are observed between

the backbones of Ala7 and Val125'. The enzymatic activities are similar to wild-type

when Ala7 is mutated to valine, cysteine, serine, or threonine, but considerably lower for

other mutants, suggesting the volume of the pocket is spatially constrained. In general,

the structural insights support the results from our mutagenesis study, and begin to

explain the mechanism by which various residues are recalcitrant to mutation.

A recent study, which explored the binding interactions of all compounds co-

crystallized with 3CL^{pro} to date, identified that 185 from a total of 233 unique molecules

bind within the active site, engaging the following 13 residues: Thr25, His41, Met49,

Asn142, Ser144, Cys145, His163, His164, Glu166, Pro168, His172, Gln189, Ala191

(Cho et al., 2021). Our analysis (**Fig. 1A**) suggests that while some of these positions

are unlikely to evolve (e.g. Ser144, His163, His172), others are more flexible (e.g.

Met49, Asn142, Glu166, Pro168, Gln189, Ala191) and may therefore harbor the

potential to confer resistance against many of the existing therapeutic candidates and

lead compounds. This is of particular concern as the only authorized 3CL^{pro} inhibitor,

nirmatrelvir (PF-07321332), is such a therapeutic, targeting the catalytic Cys145 and

nestling within the substrate binding pocket (Owen et al., 2021). While most of the

contact sites of this compound are intolerant to mutation, several demonstrate

significant plasticity in our data. These include Glu166 at the S1 subsite, which through

its side chain forms a hydrogen bond with the lactam ring of nirmatrelvir at the P1

position. From our observations, Glu166 variants that convert the acidic residue into

aliphatic, basic, sulfur-containing, and even aromatic amino acids retain wild type-like

activity. Residues Met49 and Gln189, which contribute to the formation of the lipophilic

S2 subsite via their side chains and engage in hydrophobic interactions with the P2

dimethylcyclopropylproline group of nirmatrelvir, are also notably flexible, in accordance

with other studies (Zhao et al., 2022). These analyses suggest that resistance to

nirmatrelvir may readily arise.

To demonstrate this possibility, we generated a recombinant SARS-CoV-2 virus

harboring the E166V mutation in 3CL^{pro} and tested its inhibition by nirmatrelvir and

remdesivir as a control. As Vero E6 cells express high levels of the efflux transporter P-

glycoprotein, making them unsuitable for such assays, Huh7-ACE2 cells were utilized

for these assays (Owen et al., 2021). Relative to the wild-type SARS-CoV-2, inhibition

of the E166V mutant by nirmatrelvir was significantly hampered, with a 265-fold loss in

EC$_{50}$ (**Fig. 2J and 2L**). In contrast, inhibition by remdesivir was unaffected, suggesting

that this mutation specifically enables escape from the 3CL protease inhibitor (**Fig. 2K-**

L). These data underscore the risk of resistance developing against nirmatrelvir as it

becomes widely used and further highlight the value of our screen as a resource for

drug design.

In summation, we have mapped the full functional landscape of the SARS-CoV-2

3CL^{pro} using a deep mutational scanning approach. Our work reveals that despite the

strongly conserved nature of the gene across coronaviruses, the enzyme is highly

malleable and can tolerate a plethora of mutations, even within the catalytic pocket,

without significant loss of activity. These empirical results differ from bioinformatic

predictions based on available sequences, emphasizing the value of this approach

(Krishnamoorthy and Fakhro, 2021). Consequently, these data suggest that resistance

to 3CL protease inhibitors may readily arise, and affirms the need for combinations of

therapeutics to help reduce the rate of viral escape. Yet, despite this plasticity, we were

able to identify highly conserved regions within the protein, both previously known and

novel. Our results suggest that many of these sites are indispensable for catalytic

activity or the overall structural integrity of the protein. Given their critical importance, it

is natural to speculate that such residues may serve as ideal anchor points for the next

generation of 3CL protease inhibitors developed to address the current and future

coronavirus pandemics.

Figure Legends

Figure 1. Activity of SARS-CoV-2 3CL^{pro} mutants. (**A**) Heatmap delineating protease

activity of all point mutants in SARS-CoV-2 3CL^{pro}. Wild-type at each residue is denoted

with a black circle, asterisk designates the insertion of a stop codon at the residue. The

activity score is bounded from -2 to 2. NF: N-finger; SBL: substrate binding loop. (**B**)

Overlay of histograms of all activity scores in the screen and for those observed in

clinical isolates. (**C**) Correlation between protease activity and evolutionary conservation

of residues. Data are shown as a box and whiskers plot with whiskers denoting

minimum to maximum. Dashed line is shown at 0. Statistical significance was

determined by Kruskal-Wallis test followed by Dunn's multiple comparisons test. **p <

0.01; ****p < 0.0001. (**D**) Tolerance of residues within SARS-CoV-2 3CL^{pro} for mutations

overlaid onto the crystal structure.

Figure 2. Validation of mutations and critical interactions identified in this study. (**A**)

Validation in yeast of individual SARS-CoV-2 3CL^{pro} mutants that scored as active or

inactive in the screen. Strains were grown for 48 h and then growth was quantified by

measurement of OD at 595 nm. Data are shown as mean ± s.e.m. of technical

triplicates. (**B**) Testing of recombinant live infectious virus with selected individual 3CL^{pro}

mutations through a reverse genetics system. Generated mutants were used to infect

Vero E6 cells for 72 h, and then infection was quantified by measurement of luciferase

activity. Dashed line is shown at mean value of mock. Data are shown as mean ± s.e.m.

of biological triplicates. (**C-J**) Conserved residues identified in the screen and their

possible interactions and roles in 3CL^{pro} activity. The following colors are used to denote

specific regions: yellow (domain I), pink (domain II), green (domain III), gray (N-finger),

and cyan (substrate binding loop). The substrate is shown in dark blue. A previously

solved structure of the 3CL^{pro} with its native substrate is shown (PDB: 7KHP). Residues

within the other protomer are denoted with a prime symbol ('). (**J-L**) Testing of inhibition

of wild-type and E166V live infectious virus by nirmatrelvir and remdesivir. Viruses were

used to infect Huh7-ACE2 cells for 24 h, and infection was quantified by measurement

of luciferase activity. Data are shown as mean ± s.e.m. of technical triplicates.

Supplemental Figure 1 – Related to Figure 1. Screening system and properties of the

screen. (**A**) Toxicity of SARS-CoV-2 3CL^{pro} expression in yeast. A growth defect is

observed in yeast upon expression, which is dependent on the catalytic activity of the

3CL^{pro}. Strains were grown in uninduced or induced conditions for 48 h and then growth

was quantified by measurement of OD at 595 nm. Data are shown as mean ± s.e.m. of

technical quadruplicates. (**B**) Cumulative distribution plot of codings per residue in the

screen. (**C**) Correlation between biological replicates from the screen. The Pearson

correlation coefficient is denoted in the top left. Axes are bounded from -2 to 2 as in

Figure 1A, and outliers are therefore not shown. (**D**) Proportion of types of mutations

observed at each residue in the screen.

408
Supplemental Figure 2 – Related to Figure 2. Characterization of SARS-CoV-2 3CL^{pro}

mutants. (**A**) 3CL^{pro} mutant with a truncated C-terminus retains activity in this system.

Strains were grown in uninduced or induced conditions for 48 h and then growth was

quantified by measurement of OD at 595 nm. Data are shown as mean ± s.e.m. of

technical triplicates. (**B**) Yeast two-hybrid assay to test dimerization capability of inactive

3CL^{pro} mutants. Mated strains were grown for 38 h and then growth was quantified by

measurement of OD at 595 nm. Data are shown as mean ± s.e.m. of technical

triplicates. (**C**) Western blot to determine if dimerization defective mutants were truly

dimerization defective or if they were unable to be expressed. The band corresponding

to 3CL^{pro} is indicated by the red arrow.

 STAR☐Methods

 Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal GAL4 antibody (Clone RK5C1)	Santa Cruz Biotechnology	Catalog #sc-510, RRID: AB_627655
Anti-mouse IgG Kappa light chain binding protein HRP-conjugated	Santa Cruz Biotechnology	Catalog #sc-516102, RRID: AB_2687626
Bacterial and virus strains		
pBeloBAC11 bacterial artificial chromosome containing the SARS-Cov-2 genome with the ORF7a replaced with NanoLuc luciferase reporter, with the following mutations as independent constructs: WT, C145A, T25C, Q69C, C160F, P168R, H172Y, N180G, G258Q, T292M, V114W, L250G, L268N, E288F, D295G, or E166V	(Ye et al., 2020); this manuscript	N/A
Recombinant SARS-CoV-2 NanoLuc reporter viruses (with ORF7a deletion), with the following mutations as independent viruses: WT, C145A, T25C, Q69C, C160F, P168R, H172Y, N180G, G258Q, T292M, V114W, L250G, L268N, E288F, D295G, or E166V	(Ye et al., 2020); this manuscript	N/A
10-beta Electrocompetent E. coli	New England Biolabs	Catalog #C3020K
MDS42 E. coli	(Umenhoffer et al., 2010)	N/A
Chemicals, peptides, and recombinant proteins		
PF-07321332 (nirmatrelvir)	Aobius	Catalog #AOB14800
Remdesivir	Selleckchem	Catalog #S8932
Critical commercial assays		
Nucleobond Xtra Midi kit	Macherey-Nagel	Catalog #740410.50
Q5® High-Fidelity DNA Polymerase	New England Biolabs	Catalog #M0491L
SuperSignal™ West Femto Maximum Sensitivity Substrate	Thermo Fisher Scientific	Catalog #34094
Nucleobond BAC 100 kit	Takara Bio	Catalog #740579
TRIzol™ Reagent	Thermo Fisher Scientific	Catalog #15596026
TURBO™ DNase	Thermo Fisher Scientific	Catalog #AM2238
TaqMan™ Fast Virus 1-Step Master Mix	Thermo Fisher Scientific	Catalog #4444432
TaqMan™ 2019nCoV Assay Kit v1	Thermo Fisher Scientific	Catalog #A47532
Nano-Glo® Luciferase Assay System	Promega	Catalog #N1130
Deposited data		
Raw sequencing data	NCBI Short Read Archive	BioProject ID PRJNA813935
Experimental models: Cell lines		
HEK293T	ATCC	Catalog #CRL-3216, RRID: CVCL_0063

Vero E6	ATCC	Catalog #CRL-1586, RRID: CVCL_0574
Huh7-ACE2	(Liu et al., 2022a)	N/A
Experimental models: Organisms/strains		
Yeast, BY4741 strain (S288C background)	(Brachmann et al., 1998)	N/A
Yeast, PJ69-4a/□ strain	(James et al., 1996)	N/A
Oligonucleotides		
Oligos for mutagenizing SARS-CoV-2-3CL	This manuscript	Supplemental Table 1
Oligos for amplifying SARS-CoV-2 3CL amplicons	This manuscript	Supplemental Table 2
Recombinant DNA		
pAG416-GAL-SARS-CoV-2-3CL	This manuscript	N/A
Software and algorithms		
Pipeline for calculation of activity scores of each 3CL^{pro} variant	This manuscript	https://github.com/alejandrochavezlab/3CL_protease_DMS
BWA mem	(Li and Durbin, 2009)	N/A
SAMtools	(Li et al., 2009)	N/A
ConSurf Server	(Glaser et al., 2003); (Landau et al., 2005)	N/A
GraphPad Prism v9.2	GraphPad Software	N/A

Resource availability

Lead contact

Further information and requests for materials should be directed to and will be fulfilled

by the lead contact, Alejandro Chavez (ac4304@cumc.columbia.edu).

Materials Availability

Materials used in this study will be made available under an appropriate Materials

Transfer Agreement.

Data and Code Availability

All raw data tables and code for analysis are available at

https://github.com/alejandrochavezlab/3CL_protease_DMS. The activity scores are also

available as **Supplementary Table 3**. The raw sequencing data is available for

download from the NCBI Short Read Archive under BioProject ID PRJNA813935.

Acknowledgements

This work was supported by funding from the Jack Ma Foundation to Y.S., S.P.G.,

D.D.H., and A.C, and by funding from the JPB Foundation, Andrew and Peggy Cherng,

Samuel Yin, Carol Ludwig, and David and Roger Wu to D.D.H. A.C. is also supported

by a Career Awards for Medical Scientists from the Burroughs Wellcome Fund and a

pilot grant from the Columbia HICCC. S.I. was supported by NIH grant T32AI106711.

We thank Luis Martinez-Sobrido and Chengjin Ye for the bacterial artificial chromosome

system to generate recombinant SARS-CoV-2, Jesse D. Bloom, Li Xing, and Fang-Yu

Lin for help with analysis of the data, Sagi Shapira and Barry Honig for useful

discussions, and Ayako Matsuda for assistance with preparation of Figure 2.

Author contributions

A.C. conceived this project. S.I. and S.J.H. conducted the deep mutational scan. J.S.,

R.T., and A.C. conducted initial studies to develop the screening method. F.F., F.B., and

H.M. developed and conducted the computational analyses of the screening data. J.S.

and A.K.A. provided support to improve the computational analyses. S.I., S.J.H., B.C.,

M.I.L., Y.S., and A.C. developed the method to generate recombinant SARS-CoV-2

variants and tested those viruses. S.I., S.J.H., J.S., B.C., A.F.K., and A.C. analyzed the

results. S.P.G. and D.D.H. contributed to discussions of the data and analysis. H.M.,

Y.S., D.D.H., and A.C. directed and supervised the project. S.I., S.J.H., B.C., and A.C.

wrote the manuscript with input from all authors.

Declaration of interests

S.I., D.D.H., and A.C. are inventors on patent applications related to the development of

inhibitors against the SARS-CoV-2 3CL protease.

Experimental Model and Subject Details

Yeast.

The yeast strains used for activity profiling were constructed in a BY4741 (S288C

background) (Brachmann et al., 1998) and contained additional drug sensitizing

mutations (*Δpdr1 Δprd3 Δsnq2*). For yeast two-hybrid studies, we used the PJ69-4a/□

background without the addition of the drug sensitizing mutations (James et al., 1996).

All yeast were grown at 30 °C in a rotary shaker (for culture tubes) or in a plate shaker

at 1000 RPM (for 96 deep well-plates) in growth media as appropriate.

The following yeast media were used in this study: yeast extract-peptone-

dextrose medium (YPD), synthetic complete medium deficient in uracil with glucose (SC

-ura GLU), synthetic complete medium deficient in uracil with galactose (SC -ura GAL),

synthetic complete medium deficient in leucine and tryptophan with glucose (SC -leu -

trp GLU), synthetic complete medium deficient in leucine, histidine, and tryptophan with

glucose (SC -leu –his -trp GLU). YPD was formulated with 10 g/L yeast extract, 20 g/L

peptone, and 20 g/L D-(+)-glucose. Synthetic complete media were formulated with 1.5

g/L yeast synthetic drop-out mix without yeast nitrogen base (US Biological), 1.7 g/L

yeast nitrogen base without amino acids, carbohydrate, and without ammonium sulfate

(US Biological), 5 g/L ammonium sulfate, 20 g/L D-(+)-glucose or galactose, and

supplemented with the appropriate amino acids at the following concentrations: 90 mg/L

histidine, 180 mg/L leucine, 90 mg/L lysine, 90 mg/L methionine, 90 mg/L adenine, and

18 mg/L uracil. For plates, the same recipe was used with the addition of 2% (w/v) agar

and 600 µL of 5 M NaOH/L to help the plates solidify.

For yeast transformations, a modified lithium acetate-heat shock method was

used (Gietz and Schiestl, 2007). Briefly, the appropriate parental strain was first grown

to saturation in YPD, then diluted 1:100 in YPD (~1 mL per reaction) and grown for 4 h

at 30 °C to achieve log-phase growth. A plate mix was prepared by combining the

following reagents per reaction: 71 μL 50% (w/v) PEG 3350, 8.8 μL 10X TE, 1.47 μL

freshly boiled salmon sperm ssDNA (5 mg/mL), 8.8 μL 10X LiAc, 8.8 μL DMSO, and 1

μg of plasmid DNA to be transformed. The yeast were centrifuged at 4000 RPM for 5

min, the supernatant discarded, and then resuspended in 1X LiAc. The yeast were

centrifuged again and then resuspended in 10 μL 1X LiAc per reaction. The plate mix

and yeast were then combined into individual wells of a 96 well-plate. The plate was

sealed and incubated for 20 min at 42 °C in a thermocycler. The plate was centrifuged

at 4000 RPM for 5 min, the supernatant discarded, and then each pellet was

resuspended in 20 μL sterile PBS. The yeast were then plated onto appropriate plates

and grown for 48 h at 30 °C.

Mammalian cells.

HEK293T cells and Vero E6 cells were obtained from ATCC (Catalog #CRL-3216 and

#CRL-1586, respectively). Huh7-ACE2 cells (Huh7 cells overexpression human ACE2)

were generated previously by transduction of Huh7 cells with lentivirus encoding human

ACE2, packaged using pLEX307-ACE2-blast (Addgene plasmid #158449), pMD2.G

(Addgene plasmid #12259, gift of Didier Trono), and psPAX2 (Addgene plasmid #12260,

gift of Didier Trono), and selecting for stable expression with 5 μg/mL blasticidin (Liu et

al., 2022a). Morphology was visually confirmed prior to use and all cell lines tested

mycoplasma negative.

Method Details

Measurement of SARS-CoV-2 3CL protease-induced toxicity.

The full-length SARS-CoV-2 3CL protease gene (Genbank ID: MN908947) was

synthesized (Twist Biosciences) and cloned into the yeast expression vector pAG416-

GAL (Alberti et al., 2007) by Gateway cloning (Thermo Fisher Scientific). The N-

terminus of the gene has the addition of a Met for translation initiation and the C-

terminus is native.

Mutations were generated by site-directed mutagenesis (further detailed below).

The plasmids were then transformed into the parental strain, and grown to saturation in

SC -ura GLU. The yeast were then diluted 1:500 into 1 mL of SC -ura GLU or SC -ura

GAL and grown at 30 °C with shaking. Growth was quantified by removing 100 µL from

wells after 48 h and measuring OD at 595 nm with a spectrophotometer.

Deep mutational scanning (DMS) of SARS-CoV-2 3CL protease.

For the deep mutational scanning, libraries were prepared as biological duplicates by

conducting all steps with two independent replicates.

The yeast expression vector pAG416-GAL-SARS-CoV-2-3CL was freshly

prepared by Midiprep (Macherey-Nagel), and variants were introduced by a modified

single primer site-directed mutagenesis method (Shenoy and Visweswariah, 2003).

Oligos were designed to introduce the degenerate codon NNK at the residue of interest

to introduce all possible amino acids. In addition, each oligo was designed to introduce

four or more synonymous mutations at the codon(s) immediately preceding or following

the residue of interest to be used as genetic redundancies to average observed effects,

analogous to screening approaches in which unique molecular identifiers (UMIs) have

been used (Schmierer et al., 2017; Zhu et al., 2019) (see the following **Methods**

sections for further details and **Supplemental Table 1** for all oligos). Mutagenesis was

conducted by mixing the following for each reaction: 150 ng template DNA, 1.25 µL 10

µM primer, 0.5 µL 10 mM dNTPs, 5 µL 5X Q5® Reaction Buffer, 0.25 µL Q5®

polymerase, and H_2O to 25 µL. All mutagenesis reactions were conducted in technical

duplicate. The following cycling conditions were used:

1. 98 °C, 45 s

2. 98 °C, 15 s

3. 57 °C, 15 s

4. 72 °C, 4 min 30 s

5. Return to step #2 for 29 additional cycles

6. 72 °C, 2 min

7. Hold at 4 °C

Each reaction was then digested by adding 1 µL DpnI and incubating at 37 °C for

1 h. Technical duplicates were then combined. Multiple reactions were then further

combined into sets that could be sequenced together in one amplicon (see

Supplemental Table 2 for sets). The combined reactions were then cleaned and

concentrated by a PCR purification column.

Each set of mutagenesis reactions was then transformed into 10-beta

electrocompetent E. coli (New England Biolabs) according to the manufacturer's

instructions. Transformations were conducted in quadruplicate to ensure adequate

coverage. Following recovery and outgrowth, the bacteria was directly inoculated into

liquid culture, again in quadruplicate for each transformation to avoid bottlenecking (e.g.

16 culture tubes were inoculated per set). Simultaneously, some of the bacteria was

serially plated on agar plates to check transformation efficiency and to send for Sanger

sequencing to confirm that mutagenesis was successful. Typical transformations

resulted in 3 to 4 million colonies and 10 to 40% mutation rate when 30 colonies were

sequenced.

The bacterial cultures were combined for each set and the plasmid libraries

purified from these cultures by Midiprep. The libraries were transformed into the

parental yeast strain. To ensure appropriate coverage, each set was independently

transformed 96 times. The transformed yeast were then combined, plated on SC -ura

GLU agar plates, and grown at 30 °C for 48 h. Serial dilutions were performed to check

transformation efficiency. Typical transformations resulted in approximately 10,000

colonies per transformation, producing on average a million colonies across all the

transformations for a given set of mutants.

Yeast were then scraped off all plates into sterile PBS. The concentration of

yeast was determined by measuring OD at 595 nm with a spectrophotometer and

diluted to be equivalent to a saturated culture (~40,000 cells/µL). The yeast were then

diluted 1:1000 into 1 mL of SC -ura GLU or SC -ura GAL, and grown at 30 °C for 48 h.

To control against "jackpotting" events, each outgrowth condition was repeated across

24 replicates.

Following growth, DNA was extracted from the yeast using a lithium-acetate

(LiOAc)-SDS lysis method (Looke et al., 2011). Briefly, plates were centrifuged at 4000

RPM for 5 min and the supernatant discarded. Pellets were resuspended in 200 µL of

200 mM LiOAc + 1% SDS, then incubated at 70 °C for 20 min. Then, 600 µL of 100%

ethanol was added, and mixed well. Plates were centrifuged at 4000 RPM for 10 min,

the supernatant discarded, and then allowed to air-dry until the residual ethanol

evaporated. Pellets were resuspended in 200 µL of 1X TE, incubated at 42 °C for 30

min, then centrifuged again at 4000 RPM for 10 min. The supernatant, containing the

extracted DNA, was removed and transferred to a new plate. DNA was stored at -20 °C

until further processing.

Each set was independently amplified and indexed (see **Supplemental Table 1**

for primers used for amplification of each set) and then combined and sequenced. DNA

from the 24 replicates of each condition were pooled together and used as the template

for first-round PCR. The following mix was used for each reaction: 0.5 µL template DNA,

0.1 µL 100 µM forward primer, 0.1 µL 100 µM reverse primer, 0.4 µL 10 mM dNTPs, 4

µL 5X Q5® Reaction Buffer, 0.1 µL Q5® polymerase, 14.8 µL H_2O. All PCRs were

conducted with six technical replicates to reduce sampling bias. The following cycling

conditions were used:

599   1. 98 °C, 45 s

600   2. 98 °C, 15 s

601   3. 57 °C, 20 s

602   4. 72 °C, 1 min

603   5. Return to step #2 for 27 additional cycles

604   6. 72 °C, 3 min

605   7. Hold at 4 °C

PCR products were run on a gel to confirm size, and then technical replicates

were combined and purified with AMPure XP beads (Beckman Coulter) according to the

manufacturer's instructions. The purified products were used as template for second-

round PCR with the following mix: 0.5 µL template DNA, 0.1 µL 100 µM forward primer,

0.1 µL 100 µM reverse primer, 0.4 µL 10 mM dNTPs, 2 µL 10X Taq Reaction Buffer, 0.1

µL Taq polymerase, 16.8 µL H_2O. All second-round PCRs were conducted in technical

duplicate. The following cycling conditions were used:

613   1. 94 °C, 3 min

614   2. 94 °C, 30 s

615   3. 57 °C, 20 s

616   4. 72 °C, 30 s

617   5. Return to step #2 for 11 additional cycles

618   6. 72 °C, 3 min

619   7. Hold at 4 °C

PCR products were gel purified and sequenced on an Illumina NextSeq system

with 75 bp single-end reads, except for Sets #8 - 10 which were sequenced on an

Illumina NextSeq system with 150 bp single-end reads along with residues 113, 114,

180, 187, 196, 198, 199, and 233.

Determination of catalytic activity of 3CL^{pro} variants from DMS.

The computational pipeline for the calculation of activity scores of each 3CL^{pro} variant is

available at https://github.com/alejandrochavezlab/3CL_protease_DMS. First, the raw

FASTQ files were preprocessed to determine the abundances of mutations compatible

with the designed oligos under each of the conditions. This was achieved by aligning

the reads to the original SARS-CoV-2 3CL^{pro} sequence using BWA mem (version

0.7.17-r1188) (Li and Durbin, 2009), converting the aligned reads to bam files with

SAMtools view, sort, and index (version 1.7-2) (Li et al., 2009), and then determining the

counts for sequences consistent with the oligos used for mutagenesis. The raw counts

for both replicates in both conditions (induced and non-induced) are available at

https://github.com/alejandrochavezlab/3CL_protease_DMS.

After the preprocessing, activity scores were calculated for each variant. We first

filtered out counts corresponding to sequences that differed only by one nucleotide from

the native wild-type SARS-CoV-2 3CL^{pro} sequence as these counts have a possibility of

being artificially inflated due to PCR or sequencing error. Additionally, we applied a

count threshold of 10 for the glucose condition for each variant (no threshold for the

galactose condition). Following the filtering, a pseudocount of 1 was added to all raw

counts, and the resulting counts were normalized to the native wild-type read counts of

each set to determine the relative abundance values of all mutant variants within the

pool. Using these values, the log fold change (LFC) between the glucose and galactose

conditions for each variant was calculated and an activity score determined by

performing a one-sided t-test that compared the set of LFC values representing a single

amino acid variant against the set of LFC values representing the recoded wild-type

variant of the corresponding set. The t-statistic of the t-test was taken to be the activity

score of each mutant. This t-statistic represents the difference between the LFC values

of a variant and those of the wild-type variants divided by the estimated pooled standard

deviation of two sets of LFC values. These obtained scores were then normalized set by

set, such that wild-type = 0 and stop codon = -1. We note that the reported wild-type

activity score at each residue (**Supplementary Table 3**) is derived only from the

synonymous codings of the wild-type protease, and does not use the native wild-type

DNA sequence which was only used for raw count normalization. The scores are

available at https://github.com/alejandrochavezlab/3CL_protease_DMS and also

provided as **Supplemental Table 3**.

Identification of clinical single nucleotide 3CL^{pro} variants.

To obtain a list of 3CL^{pro} variants derived from clinical specimens we made use of the

COVID-CG database (Chen et al., 2021a; Shu and McCauley, 2017). The database

was accessed on 3/25/2022 and used to download all amino acid variants from all

geographic regions within the 3CL^{pro}, corresponding to nucleotides 10,055 to 10,972. At

the time the database was accessed, there were a total of 10,441,890 3CL^{pro}

sequences within the database. As it is possible for interactions between mutations to

convert a variant that alone is detrimental into one that is well tolerated, we curated the

data provided by COVID-CG to identify mutations that occur in isolation within the

3CL^{pro} (≥ 3 occurrences), as these mutants are most similar to the variants we tested

within our high-throughput study. Insertions are not considered by COVID-CG so did not

need to be removed from the processed list. Deletions were removed as there is no

equivalent in our data.

Analysis of residue conservation within 3CL^{pro}.

Residue level evolutionary conservation was estimated by using the ConSurf server, run

with the default settings on 3/25/2022 (Glaser et al., 2003; Landau et al., 2005). PDB

7JST was used as the input sequence. MAFFT was used as the MSA method.

Homologs were collected from the UNIREF90 database using HMMER with settings of

E-value 0.0001, 3 iterations, 95% maximal %ID between sequences, and 35%

minimal %ID for homologs. 150 sequences closest to the query were chosen for

sequence selection. Rate calculation was performed using the Bayesian method. Best

fit was used as the model of substitution for proteins. The conservation score from 1 to

9 for each residue as calculated by ConSurf was used to group residues, with low

conservation being scores 1 to 3, medium conservation being scores 4 to 6, and high

conservation being scores 7 to 9.

Yeast two-hybrid to examine protease dimerization.

Yeast two-hybrid experiments were conducted by generating haploid PJ69-4a and

PJ69-4☐ strains expressing SARS-CoV-2 3CL^{pro} C145A with or without additional

mutations and fused at the N-terminus with the Gal4 activation domain or DNA binding

domain, respectively. The wild-type 3CL^{pro} could not be used for these experiments as it

is toxic to the yeast when expressed (**Fig. S1A**). The MATa and MAT☐ haploid strains

were mated on a YPD agar plate overnight to generate MATa/MAT☐ diploids, and then

grown to saturation in liquid SC -leu -trp GLU media. Yeast were subsequently diluted

1:500 into 1 mL SC -leu -trp -his GLU (selection media), and grown at 30 °C. Growth

was quantified by removing 100 μL from wells after 38 h and measuring OD at 595 nm

to assess the dimerization capability of the mutants.

To confirm that these proteases were expressed and that the proteins were not

unstable, each of the strains were also examined by immunoblotting. Strains were

grown to saturation to equilibrate input cell number and then proteins were extracted

from 1 mL of culture by a tricholoroacetic acid (TCA) method (Chun et al., 2019;

McCullough et al., 2015). Briefly, cultures were spun down, washed once with 20% TCA,

snap frozen and thawed, and then resuspended in 20% TCA with glass beads (acid-

washed, 425-600 μm, Sigma). After incubation on ice for 5 min, the mixture was

repeatedly vortexed and left on ice to disrupt cells, and then the pellet of extracted

proteins were washed with 5% TCA. The pellet was then washed with ice-cold 100%

ethanol, and then the pellet was resuspended in 1 M Tris-HCl pH 8.0. Samples were run

on SDS-PAGE and transferred to PVDF membranes. Membranes were then washed,

blocked with Blotting-Grade Blocker (Bio-Rad), washed, incubated overnight at 4 °C

with mouse monoclonal GAL4 antibody (1:100 dilution, Clone #RK5C1, Santa Cruz

Biotechnology), washed, incubated with anti-mouse IgG kappa light chain binding

protein HRP-conjugated (1:1000 dilution, Santa Cruz Biotechnology) for 1 h at RT,

washed, and then developed using SuperSignal™ West Femto Maximum Sensitivity

Substrate (Thermo Fisher Scientific).

Validation of 3CL^{pro} mutations in a SARS-CoV-2 reverse genetics system.

Individual 3CL^{pro} mutations were introduced into the pBeloBAC11 bacterial artificial

chromosome containing the SARS-CoV-2 genome in which ORF7a was replaced with a

NanoLuc luciferase reporter as previously described (Ye et al., 2020). The BAC

containing the wild-type 3CL^{pro} was kindly provided by Luis Martinez-Sobrido, and was

propagated in the MDS42 *E. coli* strain to prevent insertion of transposable elements

within the viral genome (Umenhoffer et al., 2010). To introduce 3CL^{pro} mutants, the

intact BAC was digested with PacI and MluI to remove the WT 3CL^{pro} sequence,

overlapping fragments were amplified using primers encoding the desired mutation

(**Supplementary Table 1**), and the mutant BAC was generated by Gibson assembly

(New England Biolabs). The resulting mutant BAC was electroporated into NEB 10-beta

cells and purified using the Nucleobond BAC 100 kit (Takara).

These BACs were transfected into HEK293T cells using Lipofectamine 3000®

(Thermo Fisher Scientific) in 12 well-plates (2 µg DNA per well) and incubated at 37 °C

under 5% CO_2 for 6-8 h. A mock transfection in which DNA was not added was included

with each set of transfections. The media was changed to fresh complete media (DMEM

+ 10% FCS + P/S), and further incubated for an additional 40-42 h. These cells were

then co-cultivated with naive Vero E6 cells in media with reduced serum (DMEM + 2%

FCS + P/S) and incubated for 7 days. The supernatant was then collected, clarified by

centrifugation, and aliquoted and stored at -80 °C until further use.

To normalize the amount of virus used during our subsequent assays quantifying

the replication of each mutant, RNA was first extracted from an aliquot from each viral

sample using TRIzol™ Reagent (Thermo Fisher Scientific) according to the

manufacturer's instructions. The extracted RNA was treated with TURBO™ DNase

(Thermo Fisher Scientific), and then quantified by qRT-PCR using the TaqMan™ Fast

Virus 1-Step Master Mix (Thermo Fisher Scientific) with the TaqMan™ 2019nCoV

Assay Kit v1 (Thermo Fisher Scientific). These Ct values were then utilized for

normalization of all of the viruses to equilibrate the input volume by RNA, such that

adequate signal could be observed by the wild-type SARS-CoV-2 infection. The

determined normalized volumes were used to infect naive Vero E6 cells in complete

media in a 24 well-plate. After 72 h, cells were lysed and the luciferase activity was

quantified using the Nano-Glo® Luciferase Assay System (Promega) according to the

manufacturer's instructions.

Inhibition assay with recombinant SARS-CoV-2.

The E166V mutant was generated as described above. Viruses were first titrated to

equilibrate input. One day prior to infection, Huh7-ACE2 cells were seeded at a density

of 20,000 cells per well in 96 well-plates in complete media. The following day, serial

five-fold dilutions of nirmatrelvir or remdesivir were prepared in complete media, and

then added to cells in triplicate. Cells were then infected with 0.05 MOI of either wild-

type or E166V. After 24 h, cells were lysed and the luciferase activity was quantified

using the Nano-Glo® Luciferase Assay System according to the manufacturer's

instructions. Inhibition was calculated by comparison to uninfected and infected but

untreated cells. EC_{50} values were determined by nonlinear regression using GraphPad

Prism v9.2.

Statistics.

P values between groups in **Fig. 1C** were determined by Kruskal-Wallis test followed by

Dunn's multiple comparisons test using GraphPad Prism v9.2.

References

Alberti, S., Gitler, A.D., and Lindquist, S. (2007). A suite of Gateway cloning vectors for

high-throughput genetic analysis in Saccharomyces cerevisiae. Yeast *24*, 913-919.

Amporndanai, K., Meng, X., Shang, W., Jin, Z., Rogers, M., Zhao, Y., Rao, Z., Liu, Z.J.,

Yang, H., Zhang, L.*, et al.* (2021). Inhibition mechanism of SARS-CoV-2 main protease

by ebselen and its derivatives. Nat Commun *12*, 3061.

Barrila, J., Bacha, U., and Freire, E. (2006). Long-range cooperative interactions

modulate dimerization in SARS 3CLpro. Biochemistry *45*, 14908-14916.

Beigel, J.H., Tomashek, K.M., Dodd, L.E., Mehta, A.K., Zingman, B.S., Kalil, A.C.,

Hohmann, E., Chu, H.Y., Luetkemeyer, A., Kline, S.*, et al.* (2020). Remdesivir for the

Treatment of Covid-19 - Preliminary Report. N Engl J Med.

Blanco, R., Carrasco, L., and Ventoso, I. (2003). Cell killing by HIV-1 protease. J Biol

Chem *278*, 1086-1093.

Boras, B., Jones, R.M., Anson, B.J., Arenson, D., Aschenbrenner, L., Bakowski, M.A.,

Beutler, N., Binder, J., Chen, E., Eng, H.*, et al.* (2021). Preclinical characterization of an

intravenous coronavirus 3CL protease inhibitor for the potential treatment of COVID19.

Nat Commun *12*, 6055.

Botstein, D., and Fink, G.R. (2011). Yeast: an experimental organism for 21st Century

biology. Genetics *189*, 695-704.

Brachmann, C.B., Davies, A., Cost, G.J., Caputo, E., Li, J., Hieter, P., and Boeke, J.D.

(1998). Designer deletion strains derived from Saccharomyces cerevisiae S288C: a

useful set of strains and plasmids for PCR-mediated gene disruption and other

applications. Yeast *14*, 115-132.

Chen, A.T., Altschuler, K., Zhan, S.H., Chan, Y.A., and Deverman, B.E. (2021a).

COVID-19 CG enables SARS-CoV-2 mutation and lineage tracking by locations and

dates of interest. Elife *10*.

Chen, K.X., and Njoroge, F.G. (2009). A review of HCV protease inhibitors. Curr Opin

Investig Drugs *10*, 821-837.

Chen, P., Nirula, A., Heller, B., Gottlieb, R.L., Boscia, J., Morris, J., Huhn, G., Cardona,

J., Mocherla, B., Stosor, V.*, et al.* (2021b). SARS-CoV-2 Neutralizing Antibody LY-

CoV555 in Outpatients with Covid-19. N Engl J Med *384*, 229-237.

Chen, S., Hu, T., Zhang, J., Chen, J., Chen, K., Ding, J., Jiang, H., and Shen, X.

(2008a). Mutation of Gly-11 on the dimer interface results in the complete

crystallographic dimer dissociation of severe acute respiratory syndrome coronavirus

3C-like protease: crystal structure with molecular dynamics simulations. J Biol Chem

283, 554-564.

Chen, S., Zhang, J., Hu, T., Chen, K., Jiang, H., and Shen, X. (2008b). Residues on the

dimer interface of SARS coronavirus 3C-like protease: dimer stability characterization

and enzyme catalytic activity analysis. J Biochem *143*, 525-536.

Cheng, S.C., Chang, G.G., and Chou, C.Y. (2010). Mutation of Glu-166 blocks the

substrate-induced dimerization of SARS coronavirus main protease. Biophys J *98*,

1327-1336.

Cho, E., Rosa, M., Anjum, R., Mehmood, S., Soban, M., Mujtaba, M., Bux, K., Moin,

S.T., Tanweer, M., Dantu, S.*, et al.* (2021). Dynamic Profiling of beta-Coronavirus 3CL

M(pro) Protease Ligand-Binding Sites. J Chem Inf Model *61*, 3058-3073.

Chou, C.Y., Chang, H.C., Hsu, W.C., Lin, T.Z., Lin, C.H., and Chang, G.G. (2004).

Quaternary structure of the severe acute respiratory syndrome (SARS) coronavirus

main protease. Biochemistry *43*, 14958-14970.

Chun, Y., Joo, Y.J., Suh, H., Batot, G., Hill, C.P., Formosa, T., and Buratowski, S.

(2019). Selective Kinase Inhibition Shows That Bur1 (Cdk9) Phosphorylates the Rpb1

Linker In Vivo. Mol Cell Biol *39*.

Clavel, F., and Hance, A.J. (2004). HIV drug resistance. N Engl J Med *350*, 1023-1035.

Consortium, W.H.O.S.T., Pan, H., Peto, R., Henao-Restrepo, A.M., Preziosi, M.P.,

Sathiyamoorthy, V., Abdool Karim, Q., Alejandria, M.M., Hernandez Garcia, C., Kieny,

M.P.*, et al.* (2021). Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity

Trial Results. N Engl J Med *384*, 497-511.

Dai, W., Zhang, B., Su, H., Li, J., Zhao, Y., Xie, X., Jin, Z., Liu, F., Li, C., Li, Y.*, et al.*

(2020). Structure-based design of antiviral drug candidates targeting the SARS-CoV-2

main protease. Science.

Fowler, D.M., and Fields, S. (2014). Deep mutational scanning: a new style of protein

science. Nat Methods *11*, 801-807.

Fu, L., Ye, F., Feng, Y., Yu, F., Wang, Q., Wu, Y., Zhao, C., Sun, H., Huang, B., Niu, P.*,*

et al. (2020). Both Boceprevir and GC376 efficaciously inhibit SARS-CoV-2 by targeting

its main protease. Nat Commun *11*, 4417.

Gandhi, S., Klein, J., Robertson, A.J., Pena-Hernandez, M.A., Lin, M.J., Roychoudhury,

P., Lu, P., Fournier, J., Ferguson, D., Mohamed Bakhash, S.A.K.*, et al.* (2022). De novo

emergence of a remdesivir resistance mutation during treatment of persistent SARS-

CoV-2 infection in an immunocompromised patient: a case report. Nat Commun *13*,

1547.

Garrett, M.E., Galloway, J., Chu, H.Y., Itell, H.L., Stoddard, C.I., Wolf, C.R., Logue, J.K.,

McDonald, D., Weight, H., Matsen, F.A.t.*, et al.* (2021). High-resolution profiling of

pathways of escape for SARS-CoV-2 spike-binding antibodies. Cell *184*, 2927-2938

e2911.

Garrett, M.E., Galloway, J.G., Wolf, C., Logue, J.K., Franko, N., Chu, H.Y., Matsen,

F.A.t., and Overbaugh, J.M. (2022). Comprehensive characterization of the antibody

responses to SARS-CoV-2 Spike protein finds additional vaccine-induced epitopes

beyond those for mild infection. Elife *11*.

Ghosh, A.K., Osswald, H.L., and Prato, G. (2016). Recent Progress in the Development

of HIV-1 Protease Inhibitors for the Treatment of HIV/AIDS. J Med Chem *59*, 5172-5208.

Gietz, R.D., and Schiestl, R.H. (2007). High-efficiency yeast transformation using the

LiAc/SS carrier DNA/PEG method. Nat Protoc *2*, 31-34.

Glaser, F., Pupko, T., Paz, I., Bell, R.E., Bechor-Shental, D., Martz, E., and Ben-Tal, N.

(2003). ConSurf: identification of functional regions in proteins by surface-mapping of

phylogenetic information. Bioinformatics *19*, 163-164.

Greaney, A.J., Loes, A.N., Crawford, K.H.D., Starr, T.N., Malone, K.D., Chu, H.Y., and

Bloom, J.D. (2021a). Comprehensive mapping of mutations in the SARS-CoV-2

receptor-binding domain that affect recognition by polyclonal human plasma antibodies.

Cell Host Microbe *29*, 463-476 e466.

Greaney, A.J., Starr, T.N., Gilchuk, P., Zost, S.J., Binshtein, E., Loes, A.N., Hilton, S.K.,

Huddleston, J., Eguia, R., Crawford, K.H.D.*, et al.* (2021b). Complete Mapping of

Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape Antibody

Recognition. Cell Host Microbe *29*, 44-57 e49.

Gupta, A., Gonzalez-Rojas, Y., Juarez, E., Crespo Casal, M., Moya, J., Falci, D.R.,

Sarkis, E., Solis, J., Zheng, H., Scott, N.*, et al.* (2021). Early Treatment for Covid-19

with SARS-CoV-2 Neutralizing Antibody Sotrovimab. N Engl J Med *385*, 1941-1950.

Hammond, J., Leister-Tebbe, H., Gardner, A., Abreu, P., Bao, W., Wisemandle, W.,

Baniecki, M., Hendrick, V.M., Damle, B., Simon-Campos, A.*, et al.* (2022). Oral

Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19. N Engl J Med.

Hattori, S.I., Higashi-Kuwata, N., Hayashi, H., Allu, S.R., Raghavaiah, J., Bulut, H., Das,

D., Anson, B.J., Lendy, E.K., Takamatsu, Y.*, et al.* (2021). A small molecule compound

with an indole moiety inhibits the main protease of SARS-CoV-2 and blocks virus

replication. Nat Commun *12*, 668.

Hsu, M.F., Kuo, C.J., Chang, K.T., Chang, H.C., Chou, C.C., Ko, T.P., Shr, H.L., Chang,

G.G., Wang, A.H., and Liang, P.H. (2005). Mechanism of the maturation process of

SARS-CoV 3CL protease. J Biol Chem *280*, 31257-31266.

Hu, T., Zhang, Y., Li, L., Wang, K., Chen, S., Chen, J., Ding, J., Jiang, H., and Shen, X.

(2009). Two adjacent mutations on the dimer interface of SARS coronavirus 3C-like

protease cause different conformational changes in crystal structure. Virology *388*, 324-

334.

Huang, C., Wei, P., Fan, K., Liu, Y., and Lai, L. (2004). 3C-like proteinase from SARS

coronavirus catalyzes substrate hydrolysis by a general base mechanism. Biochemistry

43, 4568-4574.

Iketani, S., Forouhar, F., Liu, H., Hong, S.J., Lin, F.Y., Nair, M.S., Zask, A., Huang, Y.,

Xing, L., Stockwell, B.R.*, et al.* (2021). Lead compounds for the development of SARS-

CoV-2 3CL protease inhibitors. Nat Commun *12*, 2016.

Iketani, S., Liu, L., Guo, Y., Liu, L., Chan, J.F., Huang, Y., Wang, M., Luo, Y., Yu, J.,

Chu, H.*, et al.* (2022). Antibody evasion properties of SARS-CoV-2 Omicron

sublineages. Nature.

James, P., Halladay, J., and Craig, E.A. (1996). Genomic libraries and a host strain

designed for highly efficient two-hybrid selection in yeast. Genetics *144*, 1425-1436.

Jayk Bernal, A., Gomes da Silva, M.M., Musungaie, D.B., Kovalchuk, E., Gonzalez, A.,

Delos Reyes, V., Martin-Quiros, A., Caraco, Y., Williams-Diaz, A., Brown, M.L.*, et al.*

(2022). Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized Patients. N Engl

J Med *386*, 509-520.

Jin, Z., Du, X., Xu, Y., Deng, Y., Liu, M., Zhao, Y., Zhang, B., Li, X., Zhang, L., Peng, C.*,*

et al. (2020a). Structure of M(pro) from SARS-CoV-2 and discovery of its inhibitors.

Nature.

Jin, Z., Zhao, Y., Sun, Y., Zhang, B., Wang, H., Wu, Y., Zhu, Y., Zhu, C., Hu, T., Du, X.*,*

et al. (2020b). Structural basis for the inhibition of SARS-CoV-2 main protease by

antineoplastic drug carmofur. Nat Struct Mol Biol *27*, 529-532.

Kneller, D.W., Phillips, G., O'Neill, H.M., Jedrzejczak, R., Stols, L., Langan, P.,

Joachimiak, A., Coates, L., and Kovalevsky, A. (2020). Structural plasticity of SARS-

CoV-2 3CL M(pro) active site cavity revealed by room temperature X-ray

crystallography. Nat Commun *11*, 3202.

Krishnamoorthy, N., and Fakhro, K. (2021). Identification of mutation resistance

coldspots for targeting the SARS-CoV2 main protease. IUBMB Life *73*, 670-675.

Landau, M., Mayrose, I., Rosenberg, Y., Glaser, F., Martz, E., Pupko, T., and Ben-Tal,

N. (2005). ConSurf 2005: the projection of evolutionary conservation scores of residues

on protein structures. Nucleic Acids Res *33*, W299-302.

Lee, J., Worrall, L.J., Vuckovic, M., Rosell, F.I., Gentile, F., Ton, A.T., Caveney, N.A.,

Ban, F., Cherkasov, A., Paetzel, M.*, et al.* (2020). Crystallographic structure of wild-type

SARS-CoV-2 main protease acyl-enzyme intermediate with physiological C-terminal

autoprocessing site. Nat Commun *11*, 5877.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-

Wheeler transform. Bioinformatics *25*, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,

Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The

Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.

Liu, H., Iketani, S., Zask, A., Khanizeman, N., Bednarova, E., Forouhar, F., Fowler, B.,

Hong, S.J., Mohri, H., Nair, M.S.*, et al.* (2022a). Development of optimized drug-like

small molecule inhibitors of the SARS-CoV-2 3CL protease for treatment of COVID-19.

Nat Commun *13*, 1891.

Liu, L., Iketani, S., Guo, Y., Chan, J.F., Wang, M., Liu, L., Luo, Y., Chu, H., Huang, Y.,

Nair, M.S.*, et al.* (2022b). Striking antibody evasion manifested by the Omicron variant

of SARS-CoV-2. Nature *602*, 676-681.

Looke, M., Kristjuhan, K., and Kristjuhan, A. (2011). Extraction of genomic DNA from

yeasts for PCR-based applications. Biotechniques *50*, 325-328.

Ma, C., Sacco, M.D., Hurst, B., Townsend, J.A., Hu, Y., Szeto, T., Zhang, X., Tarbet, B.,

Marty, M.T., Chen, Y.*, et al.* (2020). Boceprevir, GC-376, and calpain inhibitors II, XII

inhibit SARS-CoV-2 viral replication by targeting the viral main protease. Cell Res *30*,

678-692.

McCullough, L., Connell, Z., Petersen, C., and Formosa, T. (2015). The Abundant

Histone Chaperones Spt6 and FACT Collaborate to Assemble, Inspect, and Maintain

Chromatin Structure in Saccharomyces cerevisiae. Genetics *201*, 1031-1045.

Muramatsu, T., Takemoto, C., Kim, Y.T., Wang, H., Nishii, W., Terada, T., Shirouzu, M.,

and Yokoyama, S. (2016). SARS-CoV 3CL protease cleaves its C-terminal

autoprocessing site by novel subsite cooperativity. Proc Natl Acad Sci U S A *113*,

12997-13002.

Owen, D.R., Allerton, C.M.N., Anderson, A.S., Aschenbrenner, L., Avery, M., Berritt, S.,

Boras, B., Cardin, R.D., Carlo, A., Coffman, K.J.*, et al.* (2021). An oral SARS-CoV-2

M(pro) inhibitor clinical candidate for the treatment of COVID-19. Science *374*, 1586-

1593.

Planas, D., Veyer, D., Baidaliuk, A., Staropoli, I., Guivel-Benhassine, F., Rajah, M.M.,

Planchais, C., Porrot, F., Robillard, N., Puech, J.*, et al.* (2021). Reduced sensitivity of

SARS-CoV-2 variant Delta to antibody neutralization. Nature.

Qiao, J., Li, Y.S., Zeng, R., Liu, F.L., Luo, R.H., Huang, C., Wang, Y.F., Zhang, J., Quan,

B., Shen, C.*, et al.* (2021). SARS-CoV-2 M(pro) inhibitors with antiviral activity in a

transgenic mouse model. Science *371*, 1374-1378.

Resnick, S.J., Iketani, S., Hong, S.J., Zask, A., Liu, H., Kim, S., Melore, S., Lin, F.Y.,

Nair, M.S., Huang, Y.*, et al.* (2021). Inhibitors of Coronavirus 3CL Proteases Protect

Cells from Protease-Mediated Cytotoxicity. J Virol *95*, e0237420.

Schmierer, B., Botla, S.K., Zhang, J., Turunen, M., Kivioja, T., and Taipale, J. (2017).

CRISPR/Cas9 screening using unique molecular identifiers. Mol Syst Biol *13*, 945.

Shenoy, A.R., and Visweswariah, S.S. (2003). Site-directed mutagenesis using a single

mutagenic oligonucleotide and DpnI digestion of template DNA. Anal Biochem *319*,

335-336.

Shu, Y., and McCauley, J. (2017). GISAID: Global initiative on sharing all influenza data

- from vision to reality. Euro Surveill *22*.

Starr, T.N., Greaney, A.J., Addetia, A., Hannon, W.W., Choudhary, M.C., Dingens, A.S.,

Li, J.Z., and Bloom, J.D. (2021). Prospective mapping of viral mutations that escape

antibodies used to treat COVID-19. Science *371*, 850-854.

Starr, T.N., Greaney, A.J., Hilton, S.K., Ellis, D., Crawford, K.H.D., Dingens, A.S.,

Navarro, M.J., Bowen, J.E., Tortorici, M.A., Walls, A.C.*, et al.* (2020). Deep Mutational

Scanning of SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding

and ACE2 Binding. Cell *182*, 1295-1310 e1220.

Stevens, L.J., Pruijssers, A.J., Lee, H.W., Gordon, C.J., Tchesnokov, E.P., Gribble, J.,

George, A.S., Hughes, T.M., Lu, X., Li, J.*, et al.* (2022). Mutations in the SARS-CoV-2

RNA dependent RNA polymerase confer resistance to remdesivir by distinct

mechanisms. Sci Transl Med, eabo0718.

Strasfeld, L., and Chou, S. (2010). Antiviral drug resistance: mechanisms and clinical

implications. Infect Dis Clin North Am *24*, 809-833.

Suarez, D., and Diaz, N. (2020). SARS-CoV-2 Main Protease: A Molecular Dynamics

Study. J Chem Inf Model *60*, 5815-5831.

Szemiel, A.M., Merits, A., Orton, R.J., MacLean, O.A., Pinto, R.M., Wickenhagen, A.,

Lieber, G., Turnbull, M.L., Wang, S., Furnon, W.*, et al.* (2021). In vitro selection of

Remdesivir resistance suggests evolutionary predictability of SARS-CoV-2. PLoS

Pathog *17*, e1009929.

Tahir Ul Qamar, M., Alqahtani, S.M., Alamri, M.A., and Chen, L.L. (2020). Structural

basis of SARS-CoV-2 3CL(pro) and anti-COVID-19 drug discovery from medicinal

plants. J Pharm Anal *10*, 313-319.

Tan, J., Verschueren, K.H., Anand, K., Shen, J., Yang, M., Xu, Y., Rao, Z., Bigalke, J.,

Heisen, B., Mesters, J.R.*, et al.* (2005). pH-dependent conformational flexibility of the

SARS-CoV main proteinase (M(pro)) dimer: molecular dynamics simulations and

multiple X-ray structure analyses. J Mol Biol *354*, 25-40.

Umenhoffer, K., Feher, T., Baliko, G., Ayaydin, F., Posfai, J., Blattner, F.R., and Posfai,

G. (2010). Reduced evolvability of Escherichia coli MDS42, an IS-less cellular chassis

for molecular and synthetic biology applications. Microb Cell Fact *9*, 38.

Vuong, W., Khan, M.B., Fischer, C., Arutyunova, E., Lamer, T., Shields, J., Saffran, H.A.,

McKay, R.T., van Belkum, M.J., Joyce, M.A.*, et al.* (2020). Feline coronavirus drug

inhibits the main protease of SARS-CoV-2 and blocks virus replication. Nat Commun *11*,

4282.

Wang, P., Casner, R.G., Nair, M.S., Wang, M., Yu, J., Cerutti, G., Liu, L., Kwong, P.D.,

Huang, Y., Shapiro, L.*, et al.* (2021a). Increased resistance of SARS-CoV-2 variant P.1

to antibody neutralization. Cell Host Microbe *29*, 747-751 e744.

Wang, P., Nair, M.S., Liu, L., Iketani, S., Luo, Y., Guo, Y., Wang, M., Yu, J., Zhang, B.,

Kwong, P.D.*, et al.* (2021b). Antibody resistance of SARS-CoV-2 variants B.1.351 and

B.1.1.7. Nature *593*, 130-135.

Wei, P., Fan, K., Chen, H., Ma, L., Huang, C., Tan, L., Xi, D., Li, C., Liu, Y., Cao, A.*, et*

al. (2006). The N-terminal octapeptide acts as a dimerization inhibitor of SARS

coronavirus 3C-like proteinase. Biochem Biophys Res Commun *339*, 865-872.

Weinreich, D.M., Sivapalasingam, S., Norton, T., Ali, S., Gao, H., Bhore, R., Musser,

B.J., Soo, Y., Rofail, D., Im, J.*, et al.* (2021). REGN-COV2, a Neutralizing Antibody

Cocktail, in Outpatients with Covid-19. N Engl J Med *384*, 238-251.

Wu, F., Zhao, S., Yu, B., Chen, Y.M., Wang, W., Song, Z.G., Hu, Y., Tao, Z.W., Tian,

J.H., Pei, Y.Y.*, et al.* (2020). A new coronavirus associated with human respiratory

disease in China. Nature *579*, 265-269.

Ye, C., Chiem, K., Park, J.G., Oladunni, F., Platt, R.N., 2nd, Anderson, T., Almazan, F.,

de la Torre, J.C., and Martinez-Sobrido, L. (2020). Rescue of SARS-CoV-2 from a

Single Bacterial Artificial Chromosome. mBio *11*.

Zhang, L., Lin, D., Sun, X., Curth, U., Drosten, C., Sauerhering, L., Becker, S., Rox, K.,

and Hilgenfeld, R. (2020). Crystal structure of SARS-CoV-2 main protease provides a

basis for design of improved alpha-ketoamide inhibitors. Science *368*, 409-412.

Zhao, Y., Zhu, Y., Liu, X., Jin, Z., Duan, Y., Zhang, Q., Wu, C., Feng, L., Du, X., Zhao,

J.*, et al.* (2022). Structural basis for replicase polyprotein cleavage and substrate

specificity of main protease from SARS-CoV-2. Proc Natl Acad Sci U S A *119*,

e2117142119.

Zhou, P., Yang, X.L., Wang, X.G., Hu, B., Zhang, L., Zhang, W., Si, H.R., Zhu, Y., Li, B.,

Huang, C.L.*, et al.* (2020). A pneumonia outbreak associated with a new coronavirus of

probable bat origin. Nature *579*, 270-273.

Zhu, S., Cao, Z., Liu, Z., He, Y., Wang, Y., Yuan, P., Li, W., Tian, F., Bao, Y., and Wei,

W. (2019). Guide RNAs with embedded barcodes boost CRISPR-pooled screens.

Genome Biol *20*, 20.

1017

1018

1019   **Figure 1**

1020

1021   Activity Score

1022

Figure 2

1025 **Supplemental Figure 1**

1026

1027

1028

1029 **Supplemental Figure 2**

1030

1031