1 **The slowly evolving genome of the xenacoelomorph worm *Xenoturbella bocki***
2

3 Philipp H. Schiffer[1,2,*], Paschalis Natsidis[1], Daniel J. Leite[1,3], Helen E. Robertson[1],
4 François Lapraz[1,4], Ferdinand Marlétaz[1], Bastian Fromm[5], Liam Baudry[6], Fraser
5 Simpson[1], Eirik Høye[7,8], Anne-C. Zakrzewski[1,9], Paschalia Kapli[1], Katharina J.
6 Hoff[10,11], Steven Mueller[1,12], Martial Marbouty[13], Heather Marlow[14], Richard R.
7 Copley[15], Romain Koszul[13], Peter Sarkies[16], Maximilian J. Telford[1,*]
8
9 *Corresponding authors: p.schiffer@uni-koeln.de, m.telford@ucl.ac.uk
10
1 Center for Life's Origins and Evolution, Department of Genetics, Evolution and Environment,
University College London, London WC1E 6BT, UK
2 Institute for Zoology, University of Cologne, 50674 Cologne, Germany
3 Department of Biosciences, Durham University, Durham DH1 3LE, UK
4 Université Côte D'Azur, CNRS, Inserm, iBV, Nice, France
5 The Arctic University Museum of Norway, UiT – The Arctic University of Norway, Tromsø, Norway
6 Collège Doctoral, Sorbonne Université, F-75005 Paris, France
7 Department of Tumor Biology, Institute for Cancer Research, The Norwegian Radium Hospital, Oslo
University Hospital, Oslo, Norway
8 Institute of Clinical Medicine, Medical Faculty, University of Oslo, Oslo, Norway
9 Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Invalidenstr. 43,
10115 Berlin, Germany
10 University of Greifswald, Institute for Mathematics and Computer Science, Greifswald, Germany
11 University of Greifswald, Center for Functional Genomics of Microbes, Greifswald, Germany
12 Royal Brompton Hospital, Guy's and St Thomas' NHS Foundation Trust
13 Institut Pasteur, Université de Paris, CNRS UMR3525, Unité Régulation Spatiale des Génomes, F-
75015 Paris, France
14 The University of Chicago, Division of Biological Sciences, Chicago, IL 60637, USA
15 Laboratoire de Biologie du Développement de Villefranche-sur-mer (LBDV), Sorbonne Université,
CNRS, 06230 Villefranche-sur-mer, France
16 Department of Biochemistry, University of Oxford, Oxford, UK
32

Abstract

The evolutionary origins of Bilateria remain enigmatic. One of the
more enduring proposals highlights similarities between a
cnidarian-like planula larva and simple acoel-like flatworms. This idea is based in part
on the view of the Xenacoelomorpha as an outgroup to all other bilaterians which are
themselves designated the Nephrozoa (protostomes and deuterostomes). Genome
data can help to elucidate phylogenetic relationships and provide important
comparative data. Here we assemble and analyse the genome of the simple, marine
xenacoelomorph *Xenoturbella bocki*, a key species for our understanding of early
bilaterian and deuterostome evolution. Our highly contiguous genome assembly of *X.*

bocki has a size of ~11<u>1</u> Mbp in 18 chromosome like scaffolds, with repeat content

and intron, exon and intergenic space comparable to other bilaterian invertebrates.

We find *X. bocki* to have a similar number of genes to other bilaterians and to have

retained ancestral metazoan synteny. Key bilaterian signalling pathways are also

largely complete and most bilaterian miRNAs are present. We conclude that *X. bocki*

has a complex genome typical of bilaterians, in contrast to the apparent simplicity of

its body plan. Overall, our data do not provide evidence supporting the idea that

Xenacoelomorpha are a primitively simple outgroup to other bilaterians and gene

presence/absence data support a relationship with Ambulacraria.

Introduction

Xenoturbella bocki (Fig 1) is a morphologically simple marine worm first described

from specimens collected from muddy sediments in the Gullmarsfjord on the West

coast of Sweden. There are now 6 described species of *Xenoturbella* - the only genus

in the higher-level taxon of Xenoturbellida[1]. *X. bocki* was initially included as a species

within the Platyhelminthes[2], but molecular phylogenetic studies have shown that

Xenoturbellida is the sister group of the Acoelomorpha, a second clade of

morphologically   simple   worms   also   originally   considered   Platyhelminthes<u>:</u>

Xenoturbellida   and   Acoelomorpha   constitute   their   own   phylum,   the

Xenacoelomorpha[3,4]. The monophyly of Xenacoelomorpha is convincingly supported

by their sharing unique amino acid signatures in their Caudal genes[3] and Hox4/5/6

gene[5]. In the present work we analyse our data in this phylogenetic framework of a

monophyletic taxon Xenacoelomorpha.

66   The simplicity of xenacoelomorph species compared to other bilaterians is a

central feature of discussions over their evolution. While Xenacoelomorpha are clearly

monophyletic, their phylogenetic position within the Metazoa has been controversial

for a quarter of a century. There are two broadly discussed scenarios: a majority of

studies have supported a position for Xenacoelomorpha as the sister group of all other

Bilateria (the Protostomia and Deuterostomia, collectively named Nephrozoa)[4,6–8];

work we have contributed to[1,3,9,10], has instead placed Xenacoelomorpha within the

Bilateria as the sister group of the Ambulacraria (Hemichordata and Echinodermata)

to form a clade called the Xenambulacraria[9].

Figure 1: (a) Schematic drawings of *Xenoturbella bocki* showing the simple body organisation of the marine vermiform animal. Abbreviations: ant - anterior, post - posterior, lf - lateral furrow, rf - ring furrow, m - mouth opening. (b) A comparison of total length of exons, introns, and intergeneic space in the *X. bocki* genome with other metazoans (data from ref 20). *X. bocki* does not appear to be an outlier in any of theses comparisons.

Xenoturbella bocki has neither organized gonads nor a centralized nervous

system. It has a blind gut, no body cavities and lacks nephrocytes[11]. If

Xenacoelomorpha is the sister group to Nephrozoa these character absences can be

parsimoniously interpreted as representing the primitive state of the Bilateria.

According to advocates of the Nephrozoa hypothesis, these and other characters

absent in Xenacoelomorpha must have evolved in the lineage leading to Nephrozoa

after the divergence of Xenacoelomorpha. More generally there has been a tendency

to interpret Xenacoelomorpha (especially Acoelomorpha) as living approximations of

Urbilateria[12].

An alternative explanation for the simple body plan of xenaceolomorphs is

that it is derived from that of more complex urbilaterian ancestors through loss of

morphological characters. The loss or remodelling of morphological complexity is a

common feature of evolution in many animal groups and is typically associated with

unusual modes of living[13,14] – in particular the adoption of a sessile (sea squirts,

barnacles) or parasitic (neodermatan flatworms, orthonectids) lifestyle, extreme

miniaturization (e.g. tardigrades, orthonectids), or even neoteny (e.g. flightless

hexapods).

In the past some genomic features gleaned from analysis of various

Xenacoelomorpha have been used to test these evolutionary hypotheses. For

example, the common ancestor of the protostomes and deuterostomes has been

reconstructed with approximately 8 Hox genes but only 4 have been found in the

Acoelomorpha (Nemertoderma) and 5 In *Xenoturbella*. This has been interpreted as

a primary absence with the full complement of 8 appearing subsequent to the

divergence of Xenacoelomorpha and Nephrozoa. Similarly, analysis of the microRNAs

(miRNAs) of an acoelomorph, *Symsagittifera roscoffensis*, found that many bilaterian

miRNAs were absent from its genome[15]. Some of the missing bilaterian miRNAs,

however, were subsequently observed in *Xenoturbella*[9].

The few xenacoelomorph genomes available to date are from the acoel

Hofstenia miamia[16] – like other Acoelomorpha it shows accelerated sequence

evolution relative to *Xenoturbella*[3] – and from two closely related species

Praesagittifera naikaiensis[17] and *Symsagittifera roscoffensis*[18]. The analyses of gene

content of *Hofstenia* showed similar numbers of genes and gene families to other

bilaterians[16], while an analysis of the neuropeptide content concluded that most
bilaterian neuropeptides were present in Xenacoelomorpha[19].

In order to infer the characteristics of the ancestral xenacoelomorph genome,
and to complement the data from the Acoelomorpha, we describe a highly-scaffolded
genome of the slowly evolving xenacoelomorph *Xenoturbella bocki*. This allows us to
contribute knowledge of Xenacoelomorpha and *Xenoturbella* in particular of genomic
traits, such as gene content and genome-structure and to help reconstruct the genome
structure and composition of the ancestral xenacoelomorph.

Results

Assembly of a draft genome of *Xenoturbella bocki*.

We collected *Xenoturbella bocki* specimens (Fig. 1) from the bottom of the
Gullmarsfjord close to the biological field station in Kristineberg (Sweden). These adult
specimens were starved for several days in tubes with artificial sea water, and then
sacrificed in lysis buffer. We extracted high molecular weight (HMW) DNA from single
individuals for each of the different sequencing steps below.

We assembled a high-quality draft genome of *Xenoturbella bocki* using one short
read Illumina library and one TruSeq Synthetic Long Reads (TSLR) Illumina library.
We used a workflow based on a primary assembly with SPAdes (Methods; [20]). The
primary assembly had an N50 of 8.5kb over 37,880 contigs with a maximum length of
206,709bp. After using the redundans pipeline[21] this increased to an N50 of ~62kb
over 23,094 contigs and scaffolds spanning ~121Mb, and a longest scaffold of
960,978kb (supplementary Table 1).

The final genome was obtained with Hi-C scaffolding using the program
instaGRAAL (Methods, see supplementary for contact map;[22]). The scaffolded
genome has a span of 111 Mbp (117 Mbp including small fragments unincorporated
into the HiC assembly) and an N50 of 2.7 Mbp (for contigs >500bp). The assembly
contains 18 megabase-scale scaffolds encompassing 72 Mbp (62%) of the genomic
sequence, with 43% GC content. The original assembly indicated a repeat content of
about 25% after a RepeatModeller based RepeatMasker annotation (Methods). As
often seen in non-model organisms, about 2/3 of the repeats are not classified.

We used BRAKER1[23,24] with extensive RNA-Seq data, and additional single-cell

UTR enriched transcriptome sequencing data to predict 15,154 gene models. 9,575

gene models (63%) are found on the 18 large scaffolds (which represent 62% of the

total sequence). 13,298 of our predicted genes (88%) have RNA-Seq support.

Although this proportion is at the low end of bilaterian gene counts, we note that our

RNA-seq libraries were all taken from adult animals and thus may not represent the

true complexity of the gene complement. We consider our predicted gene number to

be a lower bound estimate for the true gene content.

The predicted *X. bocki* genes have a median coding length of 873 nt and a mean

length of 1330 nt. Median exon length is 132 nt (mean 212 nt) and median intron length

is 131 nt (mean 394 nt). Genes have a median of 4 exons and a mean of 8.5 exons.

2,532 genes have a single exon and, of these, 1,381 are supported as having a single

exon by RNA-Seq (TPM>1). A comparison of the exon, intron, and intergenic

sequence content in *Xenoturbella* with those described in other animal genomes[25]

show that *X. bocki* falls within the range of other similarly sized metazoan genomes

(Fig. 1b) for all these measures.

The genome of a co-sequenced *Chlamydia* species

We recovered the genome of a marine *Chlamydia* species from Illumina data obtained

from one *X. bocki* specimen and from Oxford Nanopore data from a second specimen

supporting previous microscopic analyses and single gene PCRs suggesting that *X.*

bocki is host to a species in the bacterial genus *Chlamydia*. The bacterial genome was

found as 5 contigs spanning 1,906,303 bp (N50 of 1,237,287 bp) which were

assembled into 2 large scaffolds. Using PROKKA[26], we predicted 1,738 genes in this

bacterial genome, with 3 ribosomal RNAs, 35 transfer RNAs, and 1 transfer-

messenger RNA. The genome is 97.5% complete for bacterial BUSCO[27] genes,

missing only one of the 40 core genes.

Marine chlamydiae are not closely related to the group of human pathogens[28]

and we were not able to align the genome of the *Chlamydia*-related symbioint from *X.*

bocki to the reference strain *Chlamydia trachomatis* F/SW4, nor to *Chlamydophila*

pneumoniae TW-183. To investigate the phylogenetic position of the species co-

occurring with *Xenoturbella*, we aligned the 16S rRNA gene from the *X. bocki*-hosted

Chlamydia with orthologs from related species including sequences of genes amplified

from DNA/RNA extracted from deep sea sediments. The *X. bocki*-hosted *Chlamydia*

belong to a group designated as Simkaniaceae in[28], with the sister taxon in our

phylogenetic tree being the *Chlamydia* species previously found in *X. westbladi* (*X.*

westbladi is almost certainly a synonym of *X. bocki*)[7] (Fig. 2a).

To investigate whether the *X. bocki*-hosted *Chlamydia* might contribute to the

metabolic pathways of its host, we compared the completeness of metabolic pathways

in KEGG for the *X. bocki* genome alone and for the *X. bocki* genome in combination

with the bacteria. We found only slightly higher completeness in a small number of

pathways involved in carbohydrate metabolism, carbon fixation, and amino acid

metabolism (see supplementary material) suggesting that the relationship is likely to

be commensal or parasitic rather than a true symbiosis.

A second large fraction of bacterial reads, annotated as Gammaproteobacteria,

were identified and filtered out during the data processing steps. These bacteria were

also previously reported as potential symbionts of *X. bocki*[29]. However, these

sequences were not sufficiently well covered to reconstruct a genome and we did not

investigate them further.

A phylogenetic gene presence/absence matrix supports Xenambulacraria

The general completeness of the *X. bocki* gene set allowed us to use the presence

and absence of genes identified in our genomes as a source of information to find the

best supported phylogenetic position of the Xenacoelomorpha. We conducted two

separate phylogenetic analyses of gene presence/absence data: one including the

fast-evolving Acoelomorpha and one without. In both analyses the best tree grouped

Xenoturbella with the Ambulacraria (Fig. 2b). The analysis including acoels, however,

placed the acoels as the sister-group to Nephrozoa separate from *Xenoturbella* (Fig.

2c). Because other data have shown the monophyly of Xenacoelomorpha to be robust,

we interpret this result as being the result of systematic error caused by a high rate of

gene loss or by orthologs being incorrectly scored as missing due to higher rates of

sequence evolution in acoelomorphs[30].

The *X. bocki* molecular toolkit is typical of bilaterians.

One of our principal aims was to ask whether the *Xenoturbella* genome lacks

characteristics otherwise present in the Bilateria. We found that for the Metazoa gene

2a

Figure 2a: *Xenoturbella bocki* harbours a marine Chlamydiae species as potential symbiont. In the phylogenetic analysis of 16S rDNA (ML: GTR+F+R7; bootstrap values included) the bacteria in our *X. bocki* isolate (arrow) are sister to a previous isolate from *X. westbaldi*. *X. westbaldi* is most likely a mis-identification of *X. bocki*. (b/c) A phylogeny based on presence and absence of genes calculated with OMA. Both analysis (b) and (c) confirm Xenambularcraria, i.e. Xenoturbellida in a group with Echinoderms and Hemichordates. Inclusion of the acoel flat worms places these as sister to all other Bilateria (b). This placement appears an artefact due to the very fast evolution in this taxon, in particular as good evidence exists for uniting Xenoturbellida and Acoela refs 5 and 6.

2b

2c

set in BUSCO (v5) the *X. bocki* proteome translated from our gene predictions is
82.5% complete and ~90% complete when partial hits are included (82% and 93%
respectively for the Eukaryote gene set). This estimate is even higher in the acoel
Hofstenia miamia, which was originally reported to be 90%[16], but in our re-analysis
was 95.71%. In comparison, the morphologically highly simplified and fast evolving
annelid *Intoshia linei*[31] has a genome of fewer than 10,000 genes[32] and in our analysis
is only ~64% complete for the BUSCO (v5) Metazoa set. The model nematode
Caenorhabditis elegans is ~79% complete for the same set. Despite the morphological
simplicity of both *Xenoturbella*, and *Hofstenia,* these Xenacoelomorpha are missing
few core genes compared to other bilaterian lineages that we perceive to have
undergone a high degree of morphological evolutionary change (such as the evolution
of miniaturisation, parasitism, sessility etc).
Using our phylogenomic matrix of gene presence/absence (see above) we identified
all orthologs that could be detected both in Bilateria (in any bilaterian lineage) and in
any non-bilaterian; ignoring horizontal gene transfer and other rare events, these
genes must have existed in Urbilateria (and, of less interest to us, in Urmetazoa). The
absence of any of these bilaterian genes in any lineage of Bilateria must therefore be
explained by loss of the gene. All individual bilaterian genomes were missing many
of these orthologs but Xenacoelomorphs and some other bilaterians lacked more of
these than did other taxa. The average numbers of these genes present in bilaterians
= 7577; *Xenoturbella* = 5459; *Hofstenia* = 5438; *Praesagittifera* = 4280; *Drosophila* =
4844; *Caenorhabditis* = 4323.
To better profile the *Xenoturbella* and xenacoelomorph molecular toolkit, we
used OrthoFinder to conduct orthology searches in a comparison of 155 metazoan
and outgroup species, including the transcriptomes of the sister species *X. profunda*
and an early draft genome of the acoel *Paratomella rubra* we had available*,* as well as
the *Hofstenia* and *Praesagittifera* proteomes (Supplementary online material). For
each species we counted, in each of the three Xenacoelomorphs, the number of
orthogroups for which a gene was present. The proportion of orthogroups containing
an *X. bocki* and *X. profunda* protein (87.4% and 89.2%) are broadly similar to the
proportions seen in other well characterised genomes, for example *S. purpuratus*
proteins (93.8%) or *N. vectensis* proteins (84.3%) (Fig 3a). In this analysis, the fast-

Figure 3: (a) In our orthology screen *X. bocki* shows similar percentages of genes in orthogroups, unassigned genes, and species-specific orthogroups as other well-annotated genomes. (b) The number of family members per species in major gene families (based on Pfam domains), like transcription factors, fluctuates in evolution. The *X. bocki* genome does not appear to contain particularly less or more genes in any of the analysed families. (c) Cell signalling pathways in *X. bocki* are functionally complete, but in comparison to other species contain less genes. The overall completeness is not significantly different to, for example, the nematode *C. elegans* (inset, t-test). Schematic cladograms in b/c drawn by the authors.

t-test

Species 1	Species 2	p-value
X. bocki	*H. miamia*	0.9448
X. bocki	*S. kowalevskii*	**0.0001974**
X. bocki	*S. purpuratus*	**0.0004118**
X. bocki	*C. intestinalis*	**0.0003404**
X. bocki	*B. floridae*	**0.004928**
X. bocki	*C. elegans*	0.3893
X. bocki	*D. melanogaster*	**0.0004194**
X. bocki	*I. linei*	0.2469
X. bocki	*N. vectensis*	**0.00277**
X. bocki	*H. vulgaris*	0.06593
X. bocki	*T. adhaerens*	0.4552
X. bocki	*A. queenslandica*	**0.001896**
X. bocki	*C. owczarzaki*	0.1184
X. bocki	*S. cerevisiae*	**0.007309**

evolving nematode *Caenorhabditis elegans* appears as an outlier, with only ~64% of

its proteins in orthogroups and ~35% unassigned. Both *Xenoturbella* species have an

intermediate number of unassigned genes of ~11-12%. Similarly, the proportion of

species-specific genes (~14% of all genes) corresponds closely to what is seen in

most other species (with the exception of the parasitic annelid *I. linei*, Fig. 3a).

<u>Idiosyncrasies of *Xenoturbella*</u>

In order to identify sets of orthologs specific to the two *Xenoturbella* species we used

the kinfin software[33] and found 867 such groups in the OrthoFinder clustering. We

profiled these genes based on Pfam domains and GO terms derived from

InterProScan. While these *Xenoturbella* specific proteins fall into diverse classes, we

did see a considerable number of C-type lectin, Immunoglobulin-like, PAN, and Kringle

domain containing Pfam annotations. Along with the Cysteine-rich secretory protein

family and the G-protein coupled receptor activity GO terms, these genes and families

of genes may be interesting for future studies into the biology of *Xenoturbella* in its

native environment.

<u>Gene families and signaling pathways are retained in *X. bocki*</u>

In our orthology clustering we did not see an inflation of *Xenoturbella*-specific groups

in comparison to other taxa, but also no conspicuous absence of major gene families

(Fig. 3b). Family numbers of transcription factors like Zinc-fingers or homeobox-

containing genes, as well as, for example, NACHT-domain encoding genes seem to

be neither drastically inflated nor contracted in comparison to other species in our

InterProScan based analysis.

To catalogue the completeness of cell signalling pathways we screened the *X.

bocki* proteome against KEGG pathway maps using GenomeMaple[34]. The *X. bocki*

gene set is largely complete in regard to the core proteins of these pathways, while an

array of effector proteins is absent (Fig. 3c). In comparison to other metazoan species,

as well as a unicellular choanoflagellate and a yeast, the *X. bocki* molecular toolkit has

significantly lower KEGG completeness than morphologically complex animals such

as the sea urchin and amphioxus (t-test; Fig. 3c). *Xenoturbella* is, however, not

significantly less complete when compared to other bilaterians considered to have low

Figure 4: (a) *X. bocki* has 5 HOX genes, which are located in relatively close proximity on one of our chromosome size scaffolds. Similar clusters exist for the ParaHox and "pharyngeal" genes. Numbers between genes are distance (below) and number of genes between (below). Colours indicate gene families. Red box marks the position of a partial Hox gene. The "?" gene has an unresolved homeodomain identity. (b) We found a specific prokineticin ligand signature sequence in *X. bocki*, which was previously reported for Ecdysozoa and Chordata, as well as a "K/R-RFP-K/R", sequence shared only by ambulacrarians and *X. bocki*. Signature previously reported for Ecdysozoa and Chordata, as well as new signatures we found in Spiralia and Cnidaria is absent from ambulacrarian and *X. bocki* prokineticin ligand sequences. The inset cladogram depicts the evolutionary origin of sequences in accordance with our analysis: **Bilat**erian, **Prot**ostomia, **Chor**date, **X**enacoelomorpha + **A**mbulacraria last common ancestor respectively. (c) The revised microRNA complement of *X. bocki* has a near complete set of metazoan, bilaterian and deuterostome families and genes. Presence (color) and absence (black) of microRNA families (column), paralogue numbers (values & heatmap coloring) organized in node-specific blocks in a range of representative protostome and deuterostome species compared with *Xenoturbella* (species from MirGeneDB 2.1 - Fromm et al 2021). The bottom row depicts 2011 complement by Philippe et al 2011 (blue numbers on black depict detected miRNA reads, but lack of genomic evidence). Red "x" in pink box highlights the lack of evidence for an ambulacraria-specific microRNA in *X. bocki*.

morphological complexity and which have been shown to have reduced gene content,

such as *C. elegans*, the annelid parasite *Intoshia linei*, or the acoel *Hofstenia miamia*

(Fig. 3c).

Clustered homeobox genes in the *X. bocki* genome

Acoelomorph flatworms possess three unlinked HOX genes, orthologs of anterior

(Hox1), central (Hox4/5 or Hox5) and posterior Hox (HoxP). In contrast, previous

analysis of *X. bocki* transcriptomes identified one anterior, three central and one

posterior Hox genes. We identified clear evidence of a syntenic Hox cluster with four

Hox genes (centHox1, postHox, centHox3, and antHox1) in the *X. bocki* genome (Fig.

4). There was also evidence of a fragmented annotation of centHox2, split between

the 4 gene Hox cluster and a separate scaffold (Fig. 4). In summary, this suggests that

all five Hox genes form a Hox cluster in the *X. bocki* genome, but that there are

possible unresolved assembly errors disrupting the current annotation. We also

identified other homeobox genes on the Hox cluster scaffold, including Evx (Fig. 4a).

Along with the Hox genes, we surveyed other homeobox genes that are

typically clustered in Bilateria. The canonical bilaterian paraHox cluster contains three

genes Cdx, Xlox (=Pdx) and Gsx. We identified Cdx and a new Gsx annotation on the

same scaffold, as well as a previously reported Gsx paralog on a separate scaffold.

This indicates partial retention of the paraHox cluster in *X. bocki* along with a

duplication of Gsx. On both of these paraHox containing scaffolds we observed other

homeobox genes.

Hemichordates and chordates have a conserved cluster of genes involved in

patterning their pharyngeal pores - the so-called 'pharyngeal cluster'. The homeobox

genes of this cluster (Msxlx, Nk2-1/2/4/8) were present on a single *X. bocki* scaffold.

Another pharyngeal cluster transcription factor, the Forkhead containing Foxa, and

'bystander' genes from that cluster including Egln, Mipol1 and Slc25a21 are found in

the same genomic region. Different sub-parts of the cluster are found in non-bilaterians

and protostomes and the cluster may well be plesiomorphic for the Bilateria rather

than a deuterostome synapomorphy[35].

The *X. bocki* neuropeptide complement is larger than previously thought

A catalogue of acoelomorph neuropeptides was previously described using

transcriptome data[36]. We have discovered 12 additional neuropeptide genes and 39
new neuropeptide receptors in *X. bocki* adding 6 bilaterian peptidergic systems to the
Xenoturbella catalogue (NPY-F ; MCH/Asta-C ; TRH ; ETH ; CCHa/Nmn-B ; Np-
S/CCAP), and 6 additional bilaterian systems to the Xenacoelomorpha catalogue
(Corazonin ; Kiss/GPR54 ; GPR83 ; 7B2 ; Trunk/PTTH ; NUCB2) making a total of 31
peptidergic systems (Fig. 4, Supplementary).

Among the ligand genes, we identified 6 new repeat-containing sequences. One
of these, the LRIGamide-peptide, had been identified in Nemertodermatida and
Acoela and its loss in *Xenoturbella* had been proposed[36]. We also identified the first
7B2 neuropeptide and NucB2/Nesfatin genes in Xenacoelomorpha. Finally, we
identified 3 new *X. bocki* insulin-like peptides, one of them sharing sequence similarity
and an atypical cysteine pattern with the Ambulacrarian octinsulin, constituting a
potential synapomorphy of Xenambulacraria (see Supplementary).

Our searches also revealed the presence of components of the arthropod
moulting pathway components (PTTH/trunk, NP-S/CCAP and Bursicon receptors),
which have recently been shown to be of ancient origin (de Oliveira et al., 2019). We
further identified multiple paralogs for, e.g the Tachykinin, Rya/Luquin, tFMRFa,
Corazonin, Achatin, CCK, and Prokineticin receptor families. Two complete *X. bocki*
Prokineticin ligands were also found in our survey (see Supplementary).

Chordate Prokineticin ligands possess a conserved N-terminal "AVIT" sequence
required for the receptor activation[37]. This sequence is absent in arthropod Astakine,
which instead possess two signature sequences within their Prokineticin domain [38].
To investigate Prokineticin ligands in Xenacoelomorpha we compared the sequences
of their prokineticin ligands with those of other bilaterians (Fig. 4b, Supplementary).
Our alignment reveals clade specific signatures already reported in Ecdysozoa and
Chordata sequences, but also two new signatures specific to Lophotrochozoa and
Cnidaria sequences, as well as a very specific "K/R-RFP-K/R" signature shared only
by ambulacrarian and *Xenoturbella bocki* sequences. The shared
Ambulacrarian/Xenacoelomorpha signature is found at the same position as the
Chordate sequence involved in receptor activation - adjacent to the N-terminus of the
Prokineticin domain (Fig. 4b).

The *X. bocki* genome contains most bilaterian miRNAs reported missing from acoels.

microRNAs have previously been used to investigate the phylogenetic position of the acoels and *Xenoturbella.* The acoel *Symsagittifera roscoffensis* lacks protostome and bilaterian miRNAs and this lack was interpreted as supporting the position of acoels as sister-group to the Nephrozoa. Based on shallow 454 microRNA sequencing (and sparse genomic traces) of *Xenoturbella*, some of the bilaterian miRNAs missing from acoels were found - 16 of the 32 expected metazoan (1 miRNA) and bilaterian (31 miRNAs) microRNA families – of which 6 could be identified in genome traces[9].

By deep sequencing two independent small RNA samples, we have now identified the majority of the missing metazoan and bilaterian microRNAs and identified them in the genome assembly (Fig. 4c). Altogether, we found 23 out of 31 bilaterian microRNA families (35 genes including duplicates); the single known Metazoan microRNA family (MIR-10) in 2 copies; the Deuterostome-specific MIR-103; and 7 *Xenoturbella*-specific microRNAs giving a total of 46 microRNA genes. None of the protostome-specific miRNAs were found. We could not confirm in the RNA sequences or new assembly a previously identified, and supposedly xenambulacrarian-specific MIR-2012 ortholog.

The *X. bocki* genome retains ancestral metazoan linkage groups.

The availability of chromosome-scale genomes has made it possible to reconstruct 24 ancestral linkage units broadly preserved in bilaterians[39]. In fast-evolving genomes, such as those of nematodes, tunicates or platyhelminths, these ancestral linkage groups (ALGs) are often dispersed and/or extensively fused (Supplementary). We were interested to test if the general conservation of the gene content in *X. bocki* is reflected in its genome structure.

We compared the genome of *Xenoturbella* to several other metazoan genomes and found that it has retained most of these ancestral bilaterian units: 12 chromosomes in the *X. bocki* genome derive from a single ALG, five chromosomes are made of the fusion of two ALGs, and one *Xenoturbella* chromosome is a fusion of three ALGs, as highlighted with the comparison of ortholog content with amphioxus, the sea urchin and the sea scallop (Fig. 5 and Supplementary).

One ancestral linkage group that has been lost in chordates but not in

Figure 5: (a) A comparison of scaffolds in the *X. bocki* genome with other Metazoa. 17 of the 18 large scaffolds in the *X. bocki* genome are linked via synteny to distinct chromosomal scaffolds in these species. (b) Phylostratigraphic age distribution of genes on all major scaffolds in the *X. bocki* genome. One scaffold (c1896), which showed no synteny to a distinct chromosomal scaffold in the other metazoan species also had a divergent gene age structure in comparison to other *X. bocki* scaffolds.

ambulacrarians nor in molluscs (ALG R in sea urchin and sea scallop) is detectable in

X. bocki (Fig. 5), while *X. bocki* does not show the fusions that are characteristic of

lophotrochozoans.

We also attempted to detect some pre-bilaterian arrangement of ancestral

linkage: for instance, ref [40] predicted that several pre-bilaterian linkage groups

successively fused in the bilaterian lineage to give ALGs A1, Q and E. These ALGs

are all represented as single units in *X. bocki* in common with other Bilateria. None of

the inferred pre-bilaterian chromosomal arrangements that could have provided

support for the Nephrozoa hypothesis were found *in X. bocki* although of course this

does not rule out Nephrozoa.

<u>One *X. bocki* chromosomal fragment appears aberrant</u>

The smallest of the 18 large scaffolds in the *X. bocki* genome did not show strong 1:1

clustering with any scaffold/chromosome of the bilaterian species we compared it to.

To exclude potential contamination in the assembly as a source for this contig we

examined the orthogroups to which the genes from this scaffold belong. We found that

Xenoturbella profunda[41]*,* for which a transcriptome is available, was the species that

most often occurred in the same orthogroup with genes from this scaffold (41 shared

orthogroups), suggesting the scaffold is not a contaminant.

We did observe links between the aberrant scaffold and several scaffolds from

the genome of the sponge *E. muelleri* in regard to synteny, but could not detect distinct

synteny relationships to a single scaffold in another species. In line with this, genes

on the scaffold show a different age structure compared to other scaffolds, with both

more older genes (pre bilaterian) and more *Xenoturbella* specific genes (Fig. 5b;

supported by Ks statistics, Supplementary). This aberrant scaffold also had

significantly lower levels of methylation than the rest of the genome (Supplementary).

DISCUSSION

The phylogenetic positions of *Xenoturbella* and the Acoelomorpha have been

controversial since the first molecular data from these species appeared over twenty

five years ago. Today we understand that they constitute a monophyletic group of

morphologically simple worms[1,9,42], but there remains a disagreement over whether

they represent a secondarily simplified sister group of the Ambulacraria or a primitively

simple sister group to all other Bilateria.

Previous analyses of the content of genomes, especially of Acoela, have been

used to bolster the latter view, with the small number of Hox genes and of microRNAs

of acoels being interpreted as representing an intermediate stage on the path to the

~8 Hox genes and 30 odd microRNAs of the Nephrozoa. A strong version of the

Nephrozoa idea would go further than these examples and anticipate, for example, a

genome-wide paucity of bilaterian genes, GRNs and biochemical pathways and/or an

arrangement of chromosomal segments intermediate between those of the

Eumetazoa and the Nephrozoa.

One criticism of the results from analyses of acoel genomes is that the

Acoelomorpha have evolved rapidly (their long branches in phylogenetic trees

showing high rates of sequence change). This rapid evolution might plausibly be

expected to correlate with other aspects of rapid genome evolution such as higher

rates of gene loss and chromosomal rearrangements leading to significant differences

from other Bilateria. The more normal rates of sequence evolution observed in

Xenoturbella therefore recommend it as a more appropriate xenacoelomorph to study

with fewer apomorphic characters expected.

We have sequenced, assembled, and analysed a draft genome of *Xenoturbella*

bocki. To help with annotation of the genome we have also sequenced miRNAs and

small RNAs as well as using bisulphite sequencing, Hi-C and Oxford nanopore. We

compared the gene content of the *Xenoturbella* genome to species across the

Metazoa and its genome structure to several other high-quality draft animal genomes.

We found the *X. bocki* genome to be fairly compact, but not unusually reduced

in size compared to many other bilaterians. It appears to contain a similar number of

genes (~15,000) as other animals, for example from the model organisms *D.*

melanogaster (>14,000) and *C. elegans* (~20,000). The BUSCO completeness, as

well as a high level of representation of *X. bocki* proteins in the orthogroups of our 155

species orthology screen indicates that we have annotated a near complete gene set.

Surprisingly, there are fewer genes than in the acoel *Hofstenia* (>22,000; BUSCO_v5

score ~95%). This said, of the genes found in Urbilateria (orthogroups in our

presence/absence analysis containing a member from both a bilaterian and an

outgroup) *Xenoturbella* and *Hofstenia* have very similar numbers (5459 and 5438

respectively). Gene, intron and exon lengths all also fall within the range seen in many
other invertebrate species[25]. It thus seems that basic genomic features in *Xenoturbella*
are not anomalous among Bilateria. Unlike some extremely simplified animals, such
as orthonectids, we observe no extreme reduction in gene content.

All classes of homeodomain transcription factors have previously been reported
to exist in Xenacoelomorpha[43]. We have identified 5 HOX-genes in *X. bocki* and at
least four, and probably all five of these are on one chromosomal scaffold within 187
Kbp. *X. bocki* also has the parahox genes Gsx and Cdx; while Xlox/pdx is not found,
it is present in Cnidarians and must therefore have been lost[44]. If block duplication
models of Hox and Parahox evolutionary relationships are correct, the presence of a
complete set of parahox genes implies the existence of their Hox paralogs in the
ancestor of Xenacoelomorphs suggesting the xenacoelomorph ancestor also
possessed a Hox 3 ortholog. If anthozoans also have an ortholog of bilaterian Hox 2[45],
this must also have been lost from Xenacoelomorphs. The minimal number of Hox
genes in the xenacoelomorph stem lineage was therefore probably 7 (AntHox1, lost
Hox2, lost Hox 3, CentHox 1, CentHox 2, CentHox 3 and postHoxP).

Based on early sequencing technology and without a reference genome
available, it was thought that Acoelomorpha lack many bilaterian microRNAs. Using
deep sequencing of small RNAs and our high-quality genome, we have shown that
Xenoturbella shows a near-complete bilaterian set of miRNAs including the single
deuterostome-specific miRNA family (MIR-103) (Figure X). The low number of
differential family losses of *Xenoturbella* (8 of 31 bilaterian miRNA families) inferred is
the same as the number lost in the flatworm *Schmidtea,* and substantially lower than
the number lost in the rotifer *Brachionus* (which has lost 14 bilaterian families). It is
worth mentioning that *X. bocki* shares the absence of one miRNA family (MIR-216)
with all Ambulacrarians although if Deuterostomia are paraphyletic this could be
interpretable as a primitive state[35].

The last decade has seen a re-evaluation of our understanding of the evolution
of the neuropeptide signaling genes[46,47]. The peptidergic systems are thought to have
undergone a diversification that produced approximately 30 systems in the bilaterian
common ancestor[46,47]. Our study identified 31 neuropeptide systems in *X.bocki* and
for all of these either the ligand, receptor, or both are also present in both protostomes

and deuterostomes indicating conservation across Bilateria. It is likely that more
ligands (which are short and variable) remain to be found with better detection
methods. It appears that the *Xenoturbella* genome contains a nearly complete
bilaterian neuropeptide complement with no signs of simplification but rather signs of
expansions of certain gene families. Our analyses also reveal a potential
synapomorphy linking Xenacoelomorpha with Ambulacraria (Fig 4 and
Supplementary).

We have used the predicted presence and absence of genes across a selection
of metazoan genomes as characters for phylogenetic analyses. Our trees re-confirm
the findings of recent phylogenomic gene alignment studies in linking *Xenoturbella* to
the Ambulacraria. We also used these data to test different bilaterians for their
propensity to lose otherwise conserved genes (or for our inability to identify
orthologs[30]). While the degree of gene loss appears similar between *Xenoturbella* and
acoels, the phylogenetic analysis shows longer branches leading to the acoels, most
likely due to faster evolution, gain of lineages specific genes, and some degree of
gene loss in the branch leading to the Acoelomorpha. Recent work has shown the
tendency of rapidly evolving genes (such as those belonging to rapidly evolving
species) to be missed by orthology detection software[48,49].

This pattern of conservation of evolutionarily old parts of the Metazoan genome
is further reinforced by the retention in *Xenoturbella* of linkage groups present from
sponges to vertebrates. It is interesting to note that *X. bocki* does not follow the pattern
seen in other morphologically simplified animals such as nematodes and
platyhelminths, which have lost and/or fused these ancestral linkage groups. We
interpret this to be a signal of comparably slower genomic evolution in *Xenoturbella* in
comparison to some other bilaterian lineages. The fragmented genome sequence of
Hofstenia prevents us from asking whether the ancient linkage groups have also been
preserved in the Acoelomorpha.

One of the chromosome-scale scaffolds in our assembly showed a different
methylation and age signal, with both older and younger genes, and no clear
relationship to metazoan linkage groups. By analyzing orthogroups of genes on this
scaffold for their phylogenetic signal and finding *X. bocki* genes to cluster with those

of *X. profunda* we concluded that the scaffold most likely does not represent a

contamination. It remains unclear whether this scaffold is a fast-evolving chromosome,

or a chromosomal fragment or arm. Very fast evolution on a chromosomal arm has for

example been shown in the zebrafish[50].

Apart from DNA from *X. bocki* we also obtained a highly contiguous genome of

a species related to marine *Chlamydia* species (known from microscopy to exist in *X.*

bocki); a symbiotic relationship between *Xenoturbella* and the bacteria has been

thought possible[51]. The large gene number and the completeness of genetic pathways

we found in the chlamydial genome do not support an endosymbiotic relationship.

Overall, we have shown that, while *Xenoturbella* has lost some genes - in

addition to the reduced number of Hox genes previously noted, we observe a reduction

of some signaling pathways to the core components - in general, the *X. bocki* genome

is not strikingly simpler than many other bilaterian genomes.   We   do   not   find

support for a strong version of the Nephrozoa hypothesis which would predict many

missing bilaterian genes. Bilaterian Hox and microRNA absent from Acoelomorpha

are found in *Xenoturbella* eliminating the impact of two character types that were

previously cited in support of Nephrozoa. The *Xenoturbella* genome has also largely

retained the ancestral linkage groups found in other bilaterians and does not represent

a structure intermediate between Eumetazoan and bilaterian ground states. Overall,

while we can rule out a strong version of the Nephrozoa hypothesis with many

Bilaterian characteristics missing in xenacoelomorphs, our analysis of the

Xenoturbella genome cannot distinguish between a weak version of Nephrozoa and

the Xenambulacraria topology; nevertheless, our phylogenetic analysis of gene

presence and absence supports the latter.

Methods

Genome Sequencing, Assembly, and Scaffolding

We extracted DNA from individual *Xenoturbella* specimens with a standard and

additionally worked with a Phenol-Chloroform protocol specifically developed to

extract HMW DNA (dx.doi.org/10.17504/protocols.io.mrxc57n). The extracted DNA

was quality controlled with a Nanodrop instrument in our laboratory and subsequently

a TapeStation at the sequencing center. Worms were first starved and kept in

repeatedly replaced salt water, reducing the likelihood of food or other contaminants

in the DNA extractions. First, we sequenced Illumina short paired-end reads and mate

pair libraries (see ref [3] for details). As the initial paired-read datasets were of low

complexity and coverage, we later complemented these data with an Illumina HiSeq

2000/2500 series paired-end dataset with ~700 bp insert size and 250bp read lengths,

yielding ~354 Million reads. Additionally, we generated ~40 Million

Illumina TruSeq Synthetic Long Reads (TSLR) for high confidence primary

scaffolding.

After read cleaning with Trimmomatic v.0.38[52] we conducted initial test assemblies

using the clc assembly cell v.5 and ran the blobtools pipeline[53] to screen for

contamination (Supplementary). Not detecting any significant numbers of reads from

suspicious sources in the HiSeq dataset we used SPAdes v. 3.9.0[20] to correct and

assemble a first draft genome. We also tried to use dipSPAdes but found the runtime

to exceed several weeks without finishing. We submitted the SPAdes assembly to the

redundans pipeline to eliminate duplicate contigs and to scaffold with all available mate

pair libraries. The resulting assembly was then further scaffolded with the aid of

assembled transcripts (see below) in the BADGER pipeline[54]. In this way we were able

to obtain a draft genome with ~60kb N50 that could be scaffolded to chromosome

scale super-scaffolds with the use of 3C data.

We also used two remaining specimens to extract HMW DNA for Oxford Nanopore

PromethION sequencing in collaboration with the Loman laboratory in Birmingham.

Unfortunately, the extraction failed for one individual with the DNA appearing to be

contaminated with a dark coloured residue. We were able to prepare a ligation and a

PCR library for DNA from the second specimen and obtain some genomic data.

However, due to pore blockage on both flow cells the combined data amounted to only

about 0.5-fold coverage of the genome and was thus not useful in scaffolding. We

suspect that the dark colouration of the DNA indicates a natural modification to be

present in *X. bocki* DNA that inhibits sequencing with the Oxford Nanopore method.

Library preparation for genome-wide bisulfite sequencing was performed as

previously described[55]. The resulting sequencing data were aligned to the *X. bocki*

draft genome using Bismark in non-directional mode to identify the percentage

methylation at each cytosine genome-wide. Only sites with >10 reads mapping were

considered for further analysis.

<u>Preparation of the Hi-C libraries</u>

The Hi-C protocol was adapted at the time from (Lieberman-Aiden et al., 2009; Sexton

et al., 2012 and Marie-Nelly et al., 2014). Briefly, an animal was chemically cross-

linked for one hour at room temperature in 30 mL of PBS 1X added with 3%

formaldehyde (Sigma – F8775 - 4x25 mL). Formaldehyde was quenched for 20 min

at RT by adding 10 ml of 2.5 M glycine. The fixed animal was recovered through

centrifugation and stored at -80°C until use. To prepare the proximity ligation library,

the animal was transferred to a VK05 Precellys tubes in 1X DpnII buffer (New England

Biolabs; 0.5mL) and the tissues were disrupted using the Precellys Evolution

homogenizer (Bertin-Instrument). SDS was added (0.3% final) to the lysate and the

tubes were incubated at 65°C for 20 minutes followed by an incubation at 37°C for 30

minutes and an incubation of 30 minutes after adding 50 μL of 20% triton-X100. 150

units of the DpnII restriction enzyme were then added and the tubes were incubated

overnight at 37°C. The endonuclease was inactivated 20 min at 65°C and the tubes

were then centrifuged at 16,000 x g during 20 minutes, supernatant was discarded

and pellets were re-suspended in 200 μl NE2 1X buffer and pooled. DNA ends were

labeled using 50 μl NE2 10X buffer, 37.5 μl 0.4 mM dCTP-14-biotin, 4.5 μl 10mM

dATP-dGTP-dTTP mix, 10 μl klenow 5 U/μL and incubation at 37°C for 45 minutes.

The labeling mix was then transferred to ligation reaction tubes (1.6 ml ligation buffer;

160 μl ATP 100 mM; 160 μl BSA 10 mg/mL; 50 μl T4 DNA ligase (New England

Biolabs, 5U/μl); 13.8 ml H2O) and incubated at 16°C for 4 hours. A proteinase K mix

was added to each tube and incubated overnight at 65°C. DNA was then extracted,

purified and processed for sequencing as previously described[22]. Hi-C libraries were

sequenced on a NextSeq 500 (2 × 75 bp, paired-end using custom made

oligonucleotides as in Marie-Nelly et al., 2014). Libraries were prepared separately on

two individuals in this way but eventually merged. Note that more recent version of the

HI-C protocol than the one used here have been described elsewhere[56].

 InstaGRAAL assembly pre-processing

The primary Illumina assembly contains a number of very short contigs, which are

disruptive when computing the contact distribution needed for the instaGRAAL

proximity ligation scaffolding (pre-release version, see[57] and[22] for details). Testing

several Nx metrics we found a relative length threshold, that depends on the scaffolds'

length distribution, to be a good compromise between the need for a low-noise contact

distribution and the aim of connecting most of the genome. We found N90 a suitable

threshold and excluded contigs below 1,308 bp. This also ensured no scaffolds shorter

than three times the average length of a DpnII restriction fragment (RF) were in the

assembly. In this way every contig contained enough RFs for binning and were

included in the scaffolding step.

Reads from both libraries were aligned with bowtie2 (v. 2.2.5)[58] against the

DpnII RFs of the reference assembly using the hicstuff pipeline

(https://github.com/koszullab/hicstuff) and in paired-end mode (with the options: -fg-

maxins 5 -fg-very-sensitive-local), with a mapping quality >30. The pre-processed

genome was reassembled using instaGRAAL. Briefly, the program uses a Markov

Chain Monte Carlo (MCMC) method that samples DNA segments (or bins) of the

assembly for their best relative 1D positions with respect to each other. The quality of

the positions is assessed by fitting the contact data first on a simple polymer model,

then on the plot of contact frequency according to the genomic distance law computed

from the data. The best relative position of a DNA segment with respect to one of its

most likely neighbours consists in operations such as flips, swaps, merges or a split

of contigs. Each operation is either accepted or rejected based on the computed

likelihood, resulting in an iterative progression toward the 1D structure that best fits

the contact data. Once the entire set of DNA segments is sampled for position (i.e. a

cycle), the process starts over. The scaffolder was run independently for 50 cycles,

long enough for the chromosome structure to converge. The corresponding genome

is then considered stable and suitable for further analyses. The scaffolded assemblies

were then refined using instaGRAAL's instaPolish module, to correct small artefactual

inversions that are sometimes a byproduct of instaGRAAL's processing.

Genome Annotation

 <u>Transcriptome Sequencing</u>

We extracted total RNA from a single *X. bocki* individual and sequenced a strand

specific Illumina paired end library. Extraction of total RNA was performed using a

modified Trizol & RNeasy hybrid protocol for which tissue had to be stored in RNAlater.

cDNA transcription reaction/cDNA synthesis was done using the RETROscript kit

(Ambion) using both Oligo(dT) and Random Decamer primers. Detailed extraction and

transcription protocols are available from the corresponding authors. The resulting

transcriptomic reads (deposited under <u>SRX20415651</u>) were assembled with the Trinity

pipeline[59,60] into 103,056 sequences (N50: 705; BUSCO_v5 Eukaryota scores:

C:65.1%, [S:34.1%, D:31.0%], F:22.0%, M:12.9%) for initial control and then supplied

to the genome annotation pipeline (below).

<u>Repeat annotation</u>

In the absence of a repeat library for Xenoturbellida we first used RepeatModeller

v. 1.73 to establish a library *de novo*. We then used RepeatMasker v. 4.1.0

(<u>https://www.repeatmasker.org</u>) and the Dfam library[61,62] to soft-mask the genome.

We mapped the repeats to the instaGRAAL scaffolded genome with RepeatMasker.

<u>Gene prediction and annotation</u>

We predicted genes using Augustus[63] implemented into the BRAKER (v.2.1.0)

pipeline[23,24] to incorporate the RNA-Seq data. BRAKER uses spliced aligned RNA-

Seq reads to improve training accuracy of the gene finder GeneMark-ET[64].

Subsequently, a highly reliable gene set predicted by GeneMark-ET in *ab initio* mode

was selected to train the gene finder AUGUSTUS, which in a final step predicted

genes with evidence from spliced aligned RNA-Seq reads. To make use of additional

single cell transcriptome data allowing for a more precise prediction of 3'-UTRs we

employed a production version of BRAKER (August 2018 snapshot). We had

previously mapped the RNA-Seq data to the genome with gmap-gsnap v. 2018-07-

04[65] and used samtools[66] and bamtools[67] to create the necessary input files. This

process was repeated in an iterative way, visually validating gene structures and

comparing with mappings loci inferred from a set of single-cell RNA-Seq data

(published elsewhere, see: [68]) in particular regarding fused genes. Completeness of

the gene predictions was independently assessed with BUSCO_v5[27] setting the

metazoan and the eukaryote datasets as reference respectively on gVolante[69]. We

used InterProScan v. 5.27-66.0 standalone[70,71] on the UCL cluster to annotate the

predicted *X. bocki* proteins with Pfam, SUPERFAM, PANTHER, and Gene3D

information.

<u>Horizontal Gene Transfer</u>

To detect horizontally acquired genes in the *X. bocki* genome we used a pipeline

available from (https://github.com/reubwn/hgt). Briefly, this uses blasts against the

NCBI database, alignments with MAFFT[72], and phylogenetic inferences with

IQTree[73,74] to infer most likely horizontally acquired genes, while trying to discard

contamination (e.g. from co-sequenced gut microbiota).

<u>Orthology inference</u>

We included 155 metazoan species and outgroups into our orthology analysis. We

either downloaded available proteomes or sourced RNA-Seq reads from online

repositories to then use Trinity v 2.8.5 and Trinnotate v. 3.2.0 to predict protein sets.

In the latter case we implemented diamond v. 2.0.0 blast[75,76] searches against UniProt

and Pfam[77] hmm screens against the Pfam-A dataset into the prediction process. We

had initially acquired 185 datasets, but excluded some based on inferior BUSCO

completeness, while at the same time aimed to span as many phyla as possible.

Orthology was then inferred using Orthofinder v. 2.2.7[78,79], again with diamond as the

blast engine.

Using InterProScan v. 5.27-66.0 standalone on all proteomes we added

functional annotation and then employed kinfin[33] to summarise and analyse the

orthology tables. For the kinfin analysis, we tested different query systems in regard

to phylogenetic groupings (Supplementary).

To screen for inflation and contraction of gene families we first employed

CAFE5[80], but found the analysis to suffer from long branches and sparse taxon

sampling in Xenambulacraria. We thus chose to query individual gene families (e.g.

transcription factors) by looking up Pfam annotations in the InterProScan tables of

high-quality genomes in our analysis.

Through the GenomeMaple online platform we calculated completeness of
signaling pathways within the KEGG database using GhostX as the search engine.

<u>Presence/absence phylogenetics</u>

We used a database of metazoan proteins, updated from ref [81], as the basis for
an OMA analysis to calculate orthologous groups, performing two separate runs, one
including *Xenoturbella* and acoels, and one with only *Xenoturbella*. We converted
OMA gene OrthologousMatrix.txt files into binary gene presence absence matrices in
Nexus format with datatype = restriction. We calculated phylogenetic trees on these
matrices using RevBayes (see https://github.com/willpett/metazoa-gene-content for
RevBayes script), as described in ref 74 with corrections for no absent sites
and no singleton presence, using the reversible, not the Dollo model,
as it is more likely to be able to correct for noise related to
prediction errors [82,78]. For each matrix, two runs were performed and compared and
consensus trees generated with bpcomp from Phylobayes[83].

<u>Hox and ParaHox gene cluster identification and characterisation</u>

Previous work has already used transcriptomic data and phylogenetic inference
to identify the homeobox repertoire in *Xenoturbella bocki*. These annotations were
used to identify genomic positions and gene annotations that correspond to Hox and
ParaHox clusters in *X. bocki*. Protein sequences of homeodomains (Evx, Cdx, Gsx,
antHox1, centHox1, centHox2, cent3 and postHoxP) were used as TBLASTN queries
to identify putative scaffolds associated with Hox and ParaHox clusters. Gene models
from these scaffolds were compared to the full length annotated homeobox transcripts
from[84] using BLASTP, using hits over 95% identity for homeobox classification. There
were some possible homeodomain containing genes on the scaffolds that were not
previously characterised and were therefore not given an annotation.

There were issues concerning the assignment of postHoxP and Evx to gene
models. To ascertain possible CDS regions for these genes, RNA-Seq reads were
mapped with HISAT2 to the scaffold and to previous annotation[84],were assembled
with Trinity and these were combined with BRAKER annotations.

Some issues were also observed with homeodomain queries matching genomic
sequences that were identical, suggesting artefactual duplications. To investigate
contiguity around genes the ONT reads were aligned with Minimap2 to capture long
reads over regions and coverage.

<u>Small RNA Sequencing and Analysis</u>
Two samples of starved worms were subjected to 5' monophosphate dependent
sequencing of RNAs between 15 and 36 nucleotides in length, according to previously
described methods[85]. Using miRTrace[86] 3.3, 18.6 million high-quality reads were
extracted and merged with the 27 635 high quality 454 sequencing reads from Philippe
et al. The genome sequence was screened for conserved miRNA precursors using
MirMachine[87] followed by a MirMiner run that used predicted precursors and
processed and merged reads on the genome[88]. Outputs of MirMachine and MirMiner
were manually curated using a uniform system for the annotation of miRNA genes[89]
and by comparing to MirGeneDB[90].

<u>Neuropeptide prediction and screen</u>
Neuropeptide prediction was conducted on the full set of *X.bocki* predicted
proteins using two strategies to detect neuropeptide sequence signatures. First, using
a custom script detecting the occurrence of repeated sequence patterns:
RRx(3,36)RRx(3,36)RRx(3,36)RR,RRx(2,35)ZRRx(2,35)ZRR,
RRx(2,35)GRRx(2,35)GRR, RRx(1,34)ZGRRx(1,34)ZGRR where R=K or R; x=any
amino acid; Z=any amino acid but repeated within the pattern. Second, using
HMMER3.1[91] (hmmer.org), and a combination of neuropeptide HMM models obtained
from the PFAM database (pfam.xfam.org) as well as a set of custom HMM models
derived from alignment of curated sets of neuropeptide sequences[46,47,92]. Sequences
retrieved using both methods and comprising fewer than 600 amino acids were further
validated. First, by blast analysis: sequences with E-Value ratio "best blast hit versus
ncbi nr database/best blast hit versus curated neuropeptide dataset" < 1e-40 were
discarded. Second by reciprocal best blast hit clustering using Clans[93]
(eb.tuebingen.mpg.de/protein-evolution/software/clans/) with a set of curated

neuropeptide sequences[46]. SignalP-5.0[94] (cbs.dtu.dk/services/SignalP/) was used to

detect the presence of a signal peptide in the curated list of predicted neuropeptide

sequences while Neuropred[95] (stagbeetle.animal.uiuc.edu/cgi-bin/neuropred.py) was

used to detect cleavage sites and post-translational modifications. Sequence

homology of the predicted sequence with known groups was analysed using a

combination of (i) blast sequence similarity with known bilaterian neuropeptide

sequences, (ii) reciprocal best blast hit clustering using Clans and sets of curated

neuropeptide   sequences,   (iii)   phylogeny   using   MAFFT

(mafft.cbrc.jp/alignment/server/), TrimAl[96] (trimal.cgenomics.org/) and IQ-TREE[97]

webserver for alignment, trimming and phylogeny inference respectively. Bilaterian

prokineticin-like sequences were searched in ncbi nucleotide, EST and SRA

databases as well as in the *Saccoglossus kowalevskii* genome assembly[74,98]

(groups.oist.jp/molgenu) using various bilaterian prokineticin-related protein

sequences as query. Sequences used for alignments shown in figures were collected

from ncbi nucleotide and protein databases as well as from the following publications:

7B2[46]; NucB2[92]; Insulin[99]; Prokineticin[37,38,100]. Alignments for figures were created with

Jalview (jalview.org).

<u>Neuropeptide receptor search</u>

§Neuropeptide Receptor sequences for Rhodopsin type GPCR, Secretin type GPCR

and tyrosine and serine/threonine kinase receptors were searched by running

HMMER3.1 on the full set of *X.bocki* predicted proteins using the 7tm_1 (PF00001),

7tm_2 (PF00002) and PK_Tyr_Ser-Thr (PF07714) HMM models respectively which

were obtained from the PFAM database (pfam.xfam.org). Sequences above the

significance threshold were then aligned with sequences from the curated dataset,

trimmed and phylogeny inference was conducted using same method as for the

neuropeptide. A second alignment and phylogeny inference was conducted after

removal of all *X.bocki* sequences having no statistical support for grouping with any of

the known neuropeptide receptors from the curated dataset. Curated datasets were

collected from the following publications: Rhodopsin type GPCR beta and gamma and

Secretin type GPCR[100]; Rhodopsin type GPCR delta (Leucine-rich repeat-containing

G-protein coupled Receptors)[101]; Tyrosine kinase receptors[102,103]; and were

complemented with sequences from NCBI protein database.

Synteny

Ancestral linkage analyses rely on mutual-best-hits computed using Mmseqs2[104]

between pairs of species in which chromosomal assignments to ancestral linkage

groups (ALG) was previously performed, such as *Branchiostoma floridae* or *Pecten*

maximus[39]. Oxford dotplots were computed by plotting reciprocal positions of indexed

pairwise orthologs between two species as performed previously[39,40]. The significance

of ortholog enrichment in pairs of chromosomes was assessed using a fisher test.

We also used a Python implementation of MCscanX[105] (Haibao Tang and available

on https://github.com/tanghaibao/jcvi/wiki/MCscan-(Python-version)) to compare *X.*

bocki to *Euphydtia muelleri*, *Trichoplax adhearens*, *Branchiostoma floridae*,

Saccoglossus kowalevskii, *Ciona intestinalis*, *Nematostella vectensis*, *Asteria rubens,*

Pecten maximus, *Nemopilema nomurai*, *Carcinoscorpius rotundicauda* (see

Supplementary). Briefly, the pipeline uses high quality genomes and their annotations

to infer syntenic blocks based on proximity. For this an all vs. all blastp is performed

and synteny extended from anchors identified in this way. Corresponding heatmaps

(see Supplementary) were plotted with Python in a Jupyter notebooks instance.

Chlamydia assembly and annotation

We identified a highly contiguous *Chlamydia* genome in the *X. bocki* genome

assembly using blast. We then used our Oxford Nanopore derived long-reads to

scaffold the *Chlamyida* genome with LINKS[106] and annotated it with the automated

PROKKA pipeline. To place the genome on the *Chlamydia* tree we extracted the 16S

ribosomal RNA gene sequence, aligned it with set of *Chlamydia* 16S rRNA sequences

from[28] using MAFFT, and reconstructed the phylogeny using IQ-TREE 2[73] We

visualized the resulting tree with Figtree (http://tree.bio.ed.ac.uk/).

Acknowledgements

We thank Josh Quick and Nick Loman for help with the generation of ONT long-read

data. Analyses were conducted mainly on the UCL Cluster, with some computations

also run on the CHEOPS cluster at the University of Cologne. We are grateful to Kevin

J. Peterson for his comments on the manuscript, the miRNA section in particular. We

thank the Kristineberg Center for Marine Research and Innovation for their essential
support in sampling *Xenoturbella*.
816

Conflict of interest
817
818   The authors declare no conflict of interest.
819

Data availability
821
All read sets (RNA and DNA derived) used in this study will be made available with
the publication of this manuscript on the SRA database under the BioProject ID
PRJNA864813. Hi-C reads are deposited under SAMN30224387, RNA-Seq under
SAMN35083895. The genome assemblies of *X. bocki* (ERS12565994,
ERA16814408) and the *Chlamydia* sp. (ERS12566084, ERA16814775) are
deposited under PRJEB55230 at ENA.
828

Funding
829
PHS was funded by an ERC grant (ERC-2012-AdG 322790) to MJT, which also
supported HR, ACZ, SM. PHS was also funded through an Emmy-Noether grant
(434028868) to himself. Part of this work was funded by BBSRC grant
BB/R016240/1 (M.J.T./P.K.), by a Leverhulme Trust Research Project Grant RPG-
2018-302 (M.J.T./D.J.L.), and by the European Union's Horizon 2020 research and
innovation program under the Marie Skłodowska-Curie grant agreement no 764840
IGNITE (M.J.T./P.N.).

References

1. Telford, M. J. Xenoturbellida: the fourth deuterostome phylum and the diet of worms. *Genesis (New York, N.Y. : 2000)* 46, 580–586 (2008).

2. Westblad, E. *Xenoturbella bocki* n. g., n. sp., a peculiar, primitive Turbellarian type. *Arkiv för Zoologi* 3–29 (1949).

3. Philippe, H. *et al.* Mitigating Anticipated Effects of Systematic Errors Supports Sister-Group Relationship between Xenacoelomorpha and Ambulacraria. *Current Biology* 29, 1818-1826.e6 (2019).

4. Cannon, J. T. *et al.* Xenacoelomorpha is the sister group to Nephrozoa. *Nature* 530, 89–93 (2016).

5. Ueki, T., Arimoto, A., Tagawa, K. & Satoh, N. Xenacoelomorph-Specific Hox Peptides: Insights into the Phylogeny of Acoels, Nemertodermatids, and Xenoturbellids. *Zool Sci* 36, 395–401 (2019).

6. Hejnol, A. *et al.* Assessing the root of bilaterian animals with scalable phylogenomic methods. *Proceedings. Biological sciences / The Royal Society* 276, 4261–4270 (2009).

7. Rouse, G. W., Wilson, N. G., Carvajal, J. I. & Vrijenhoek, R. C. New deep-sea species of *Xenoturbella* and the position of Xenacoelomorpha. *Nature* 530, 94–97 (2016).

8. Srivastava, M., Mazza-Curll, K. L., Wolfswinkel, J. C. van & Reddien, P. W. Whole-body acoel regeneration is controlled by Wnt and Bmp-Admp signaling. *Current Biology* 24, 1107–1113 (2014).

9. Philippe, H. *et al.* Acoelomorph flatworms are deuterostomes related to *Xenoturbella*. *Nature* 470, 255–258 (2011).

10. Bourlat, S. J. *et al.* Deuterostome phylogeny reveals monophyletic chordates and the new phylum Xenoturbellida. *Nature* 444, 85–88 (2006).

11. Nakano, H. What is *Xenoturbella*? *Zoological Letters* 1, 1 (2015).

12. Hejnol, A. & Martindale, M. Q. Acoel development supports a simple planula-like urbilaterian. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 363, 1493–1501 (2008).

13. Martynov, A. *et al.* Multiple paedomorphic lineages of soft-substrate burrowing invertebrates: parallels in the origin of *Xenocratena* and *Xenoturbella*. *Plos One* 15, e0227173 (2020).

14. Westheide, W. Progenesis as a principle in meiofauna evolution. *J Nat Hist* 21, 843–854 (1987).

15. Sempere, L. F., Cole, C. N., McPeek, M. A. & Peterson, K. J. The phylogenetic
distribution of metazoan microRNAs: insights into evolutionary complexity and constraint.
306, 575–588 (2006).

16. Gehrke, A. R. *et al.* Acoel genome reveals the regulatory landscape of whole-body
regeneration. *Science* 363, 1–9 (2019).

17. Arimoto, A. *et al.* A draft nuclear-genome assembly of the acoel flatworm *Praesagittifera*
naikaiensis. *Gigascience* 8, giz023 (2019).

18. Martinez, P. *et al.* Genome assembly of the acoel flatworm *Symsagittifera roscoffensis*, a
model for research on body plan evolution and photosymbiosis. *G3 Genes Genomes Genetics*
13, jkac336 (2022).

19. Moroz, L. L., Romanova, D. Y. & Kohn, A. B. Neural versus alternative integrative
systems: molecular insights into origins of neurotransmitters. *Philosophical Transactions*
Royal Soc Lond Ser B Biological Sci 376, 20190762 (2021).

20. Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *www.liebertpub.com* 19, 455–477 (2012).

21. Pryszcz, L. P. & Gabaldón, T. Redundans: an assembly pipeline for highly heterozygous
genomes. *Nucleic Acids Research* 44, e113–e113 (2016).

22. Baudry, L. *et al.* instaGRAAL: chromosome-level quality scaffolding of genomes using a
proximity ligation-based scaffolder. *Genome Biol* 21, 148 (2020).

23. Hoff, K. J., Lomsadze, A., Borodovsky, M. & Stanke, M. Whole-genome annotation with
BRAKER. *Methods Mol Biology Clifton N J* 1962, 65–95 (2019).

24. Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M. & Stanke, M. BRAKER1:
Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.
Bioinformatics 32, 767–769 (2016).

25. Francis, W. R. & Wörheide, G. Similar ratios of introns to intergenic sequence across
animal genomes. *Genome Biol Evol* 9, evx103- (2017).

26. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinform Oxf Engl* 30, 2068–
9 (2014).

27. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. 31, 3210–3212 (2015).

28. Dharamshi, J. E. *et al.* Marine sediments illuminate Chlamydiae diversity and evolution.
Curr Biol 30, 1032-1048.e7 (2020).

29. Kjeldsen, K. U., Obst, M., Nakano, H., Funch, P. & Schramm, A. Two types of
endosymbiotic bacteria in the enigmatic marine worm *Xenoturbella bocki*. 76, 2657–2662
(2010).

30. Natsidis, P., Kapli, P., Schiffer, P. H. & Telford, M. J. Systematic errors in orthology
inference and their effects on evolutionary analyses. *Iscience* 102110 (2021).

31. Schiffer, P. H., Robertson, H. E. & Telford, M. J. Orthonectids are highly degenerate
annelid worms. *Current Biology* 1–9 (2018).

32. Mikhailov, K. V. *et al.* The genome of *Intoshia linei* affirms orthonectids as highly
simplified spiralians. *Current Biology* 26, 1768–1774 (2016).

33. Laetsch, D. R., Laetsch, D. R., Blaxter, M. L. & Blaxter, M. L. KinFin: Software for
taxon-aware analysis of clustered protein sequences. *G3 (Bethesda, Md.)* 7, 3349–3357
(2017).

34. Takami, H. *et al.* An automated system for evaluation of the potential functionome:
MAPLE version 2.1.0. *Dna Res* 23, 467–475 (2016).

35. Kapli, P. *et al.* Lack of support for Deuterostomia prompts reinterpretation of the first
Bilateria. *Sci Adv* 7, eabe2741 (2021).

36. Thiel, D., Franz-Wachtel, M., Aguilera, F. & Hejnol, A. Xenacoelomorph
neuropeptidomes reveal a major expansion of neuropeptide systems during early bilaterian
evolution. *Molecular Biology And Evolution* 35, 2528–2543 (2018).

37. Negri, L. & Ferrara, N. The Prokineticins: neuromodulators and mediators of
inflammation and myeloid cell-dependent angiogenesis. *Physiol Rev* 98, 1055–1082 (2018).

38. Ericsson, L. & Söderhäll, I. Astakines in arthropods–phylogeny and gene structure. *Dev
Comp Immunol* 81, 141–151 (2018).

39. Simakov, O. *et al.* Deeply conserved synteny resolves early events in vertebrate
evolution. *Nat Ecol Evol* 1–11 (2020).

40. Simakov, O. *et al.* Deeply conserved synteny and the evolution of metazoan
chromosomes. *Sci Adv* 8, eabi5884 (2022).

41. Rouse, G. W., Wilson, N. G., Carvajal, J. I. & Vrijenhoek, R. C. New deep-sea species of
Xenoturbella and the position of Xenacoelomorpha. *Nature* 530, 94–97 (2016).

42. Hejnol, A. Acoelomorpha and Xenoturbellida. in 203–214 (Springer Vienna, 2015).

43. Brauchle, M. *et al.* Xenacoelomorpha survey reveals that all 11 animal homeobox gene
classes were present in the first bilaterians. *Genome Biol Evol* 10, 2205–2217 (2018).

44. Jimenez-Guri, E., Paps, J., Garcia-Fernandez, J. & Salo, E. Hox and ParaHox genes in
Nemertodermatida, a basal bilaterian clade. *Int J Dev Biology* 50, 675–679 (2006).

45. Ryan, J. F. *et al.* The cnidarian-bilaterian ancestor possessed at least 56 homeoboxes:
evidence from the starlet sea anemone, *Nematostella vectensis*. *Genome Biol* 7, R64–R64
(2006).

46. Jekely, G. Global view of the evolution and diversity of metazoan neuropeptide signaling.
Proc National Acad Sci 110, 8702–8707 (2013).

47. Mirabeau, O. & Joly, J.-S. Molecular evolution of peptidergic signaling systems in
bilaterians. *Proc National Acad Sci* 110, E2028–E2037 (2013).

48. Natsidis, P., Kapli, P., Schiffer, P. H. & Telford, M. J. Systematic errors in orthology
inference and their effects on evolutionary analyses. *iScience* 24, 102110 (2021).

49. Weisman, C. M., Murray, A. W. & Eddy, S. R. Many, but not all, lineage-specific genes
can be explained by homology detection failure. *Plos Biol* 18, e3000862 (2020).

50. Howe, K. *et al.* Structure and evolutionary history of a large family of NLR proteins in
the zebrafish. *Open Biology* 6, 160009 (2016).

51. Pillonel, T., Bertelli, C. & Greub, G. Environmental metagenomic assemblies reveal
seven new highly divergent chlamydial lineages and hallmarks of a conserved intracellular
lifestyle. *Front Microbiol* 9, 79 (2018).

52. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30, 2114–2120 (2014).

53. Laetsch, D. R. & Blaxter, M. L. BlobTools: Interrogation of genome assemblies.
F1000research 6, 1287 (2017).

54. Elsworth, B., Jones, M. & Blaxter, M. Badger--an accessible genome exploration
environment. *Bioinformatics* 29, 2788–2789 (2013).

55. Lewis, S. H. *et al.* Widespread conservation and lineage-specific diversification of
genome-wide DNA methylation patterns across arthropods. *Plos Genet* 16, e1008864 (2020).

56. Lafontaine, D. L., Yang, L., Dekker, J. & Gibcus, J. H. Hi-C 3.0: improved protocol for
genome-wide chromosome conformation capture. *Curr Protoc* 1, e198 (2021).

57. Marie-Nelly, H. *et al.* High-quality genome (re)assembly using chromosomal contact
data. *Nat Commun* 5, 5695 (2014).

58. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature
Methods* 9, 357–359 (2012).

59. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-Seq: reference
generation and analysis with Trinity. 8, 1494–1512 (2013).

60. *RNA-Seq De novo Assembly Using Trinity.* 1–7 (2015).

61. Wheeler, T. J. *et al.* Dfam: a database of repetitive DNA based on profile hidden Markov
models. *Nucleic Acids Res* 41, D70–D82 (2013).

62. Hubley, R. *et al.* The Dfam database of repetitive DNA families. *Nucleic Acids Res* 44,
D81–D89 (2016).

63. Stanke, M. & Waack, S. Gene prediction with a hidden Markov model and a new intron
submodel. *Bioinformatics* 19 Suppl 2, ii215-25 (2003).

64. Lomsadze, A., Burns, P. D. & Borodovsky, M. Integration of mapped RNA-Seq reads
into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res* 42, e119–
e119 (2014).

65. Wu, T. D., Reeder, J., Lawrence, M., Becker, G. & Brauer, M. J. GMAP and GSNAP for
Genomic Sequence Alignment: Enhancements to Speed, Accuracy, and Functionality.
Methods Mol Biology Clifton N J 1418, 283–334 (2016).

66. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25,
2078–2079 (2009).

67. Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P. & Marth, G. T.
BamTools: a C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* 27,
1691–1692 (2011).

68. Robertson, H. E. *et al.* Single cell atlas of *Xenoturbella bocki* highlights the limited cell-
type complexity of a non-vertebrate deuterostome lineage. *Biorxiv* 2022.08.18.504214 (2022)
doi:10.1101/2022.08.18.504214.

69. Nishimura, O., Hara, Y. & Kuraku, S. gVolante for standardizing completeness
assessment of genome and transcriptome assemblies. *Bioinformatics* 33, 3635–3637 (2017).

70. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.
Bioinformatics 30, 1236–1240 (2014).

71. Mulder, N. & Apweiler, R. InterPro and InterProScan: tools for protein sequence
classification and comparison. *Methods in molecular biology (Clifton, N.J.)* 396, 59–70
(2007).

72. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol Biol Evol* 30, 772–780 (2013).

73. Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for phylogenetic
inference in the genomic era. *Mol Biol Evol* 37, 1530–1534 (2020).

74. Nguyen, L.-T., Schmidt, H. A., Haeseler, A. von & Minh, B. Q. IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*
32, 268–274 (2015).

75. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. *Nature Methods* 12, 59–60 (2014).

76. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale
using DIAMOND. *Nat Methods* 18, 366–368 (2021).

77. Finn, R. D. *et al.* The Pfam protein families database: towards a more sustainable future.
Nucleic Acids Research 44, D279-85 (2016).

78. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* 20, 238 (2019).

79. Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biology* 16, E9-
13 (2015).

80. Han, M. V., Thomas, G. W. C., Lugo-Martinez, J. & Hahn, M. W. Estimating gene gain
and loss rates in the presence of error in genome assembly and annotation using CAFE 3. *Mol*
Biol Evol 30, 1987–1997 (2013).

81. Leclère, L. *et al.* The genome of the jellyfish *Clytia hemisphaerica* and the evolution of
the cnidarian life-cycle. *Nat. Ecol. Evol.* 1–41 (2019).

82. Pett, W. *et al.* The Role of Homology and Orthology in the Phylogenomic Analysis of
Metazoan Gene Content. *Mol Biol Evol,* 1–7 (2019).

83. Lartillot, N., Lepage, T. & Blanquart, S. PhyloBayes 3: a Bayesian software package for
phylogenetic reconstruction and molecular dating. 25, 2286–2288 (2009).

84. Brauchle, M. *et al.* Xenacoelomorpha survey reveals that all 11 animal homeobox gene
classes were present in the first bilaterians. *Genome biology and evolution* (2018)
doi:10.1093/gbe/evy170.

85. Sarkies, P. *et al.* Ancient and novel small RNA pathways compensate for the loss of
piRNAs in multiple independent nematode lineages. *Plos Biol* 13, e1002061 (2015).

86. Kang, W. *et al.* miRTrace reveals the organismal origins of microRNA sequencing data.
Genome Biol 19, 213 (2018).

87. Umu, S. U. *et al.* Accurate microRNA annotation of animal genomes using trained
covariance models of curated microRNA complements in MirMachine. *Biorxiv*
2022.11.23.517654 (2023).

88. Wheeler, B. M. *et al.* The deep evolution of metazoan microRNAs. *Evol Dev* 11, 50–68
(2009).

89. Fromm, B. *et al.* A uniform system for the annotation of vertebrate microRNA genes and
the evolution of the human microRNAome. *Annu Rev Genet* 49, 213–242 (2015).

90. Fromm, B. *et al.* MirGeneDB 2.1: toward a complete sampling of all major animal phyla.
Nucleic Acids Res 50, D204–D210 (2022).

91. Johnson, L. S., Eddy, S. R. & Portugaly, E. Hidden Markov model speed heuristic and
iterative HMM search procedure. *Bmc Bioinformatics* 11, 431–431 (2010).

92. Zandawala, M. *et al.* Discovery of novel representatives of bilaterian neuropeptide
families and reconstruction of neuropeptide precursor evolution in ophiuroid echinoderms.
Open Biol 7, 170129 (2017).

93. Frickey, T. & Lupas, A. CLANS: a Java application for visualizing protein families based
on pairwise similarity. *Bioinformatics* 20, 3702–3704 (2004).

94. Armenteros, J. J. A. *et al.* SignalP 5.0 improves signal peptide predictions using deep
neural networks. *Nat Biotechnol* 37, 420–423 (2019).

95. Southey, B. R., Rodriguez-Zas, S. L. & Sweedler, J. V. Prediction of neuropeptide
prohormone cleavages with application to RFamides. *Peptides* 27, 1087–1098 (2006).

96. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldon, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25, 1972–1973
(2009).

97. Nguyen, L.-T., Schmidt, H. A., Haeseler, A. von & Minh, B. Q. IQ-TREE: A fast and
effective stochastic algorithm for estimating Maximum-Likelihood phylogenies. *Mol Biol*
Evol 32, 268–274 (2015).

98. Simakov, O. *et al.* Hemichordate genomes and deuterostome origins. *Nature* 527, 459–
465 (2015).

99. Cherif--Feildel, M., Berthelin, C. H., Rivière, G., Favrel, P. & Kellner, K. Data for
evolutive analysis of insulin related peptides in bilaterian species. *Data Brief* 22, 546–550
(2019).

100. Thiel, D., Franz-Wachtel, M., Aguilera, F. & Hejnol, A. Changes in the neuropeptide
complement correlate with nervous system architectures in xenacoelomorphs. 1–57 (2018)
doi:10.1101/265579.

101. Roch, G. J. & Sherwood, N. M. Glycoprotein hormones and their receptors emerged at
the origin of metazoans. *Genome Biol Evol* 6, 1466–79 (2014).

102. Oliveira, A. L. de, Calcino, A. & Wanninger, A. Ancient origins of arthropod moulting
pathway components. *elife* 8, e46113 (2019).

103. Smýkal, V. *et al.* Complex evolution of insect insulin receptors and homologous decoy
receptors, and functional significance of their multiplicity. *Mol Biol Evol* 37, 1775–1789
(2020).

104. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nat Biotechnol* 35, 1026–1028 (2017).

105. Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis of gene
synteny and collinearity. *Nucleic Acids Research* 40, e49–e49 (2012).

106. Warren, R. L. *et al.* LINKS: Scalable, alignment-free scaffolding of draft genomes with
long reads. *Gigascience* 4, 35 (2015).