In Vitro Evaluation and Mitigation of Niclosamide's Liabilities as a COVID-19 Treatment.

Jesse W. Wotring[1], Sean M. McCarty[1], Khadija Shafiq[1], Charles J. Zhang[1], Theophilus Nguyen[2], Sophia R. Meyer[1], Reid Fursmidt[2], Carmen Mirabelli[3], Martin C. Clasby[1], Christiane E. Wobus[3], Matthew J. O'Meara[4], Jonathan Z. Sexton[1,2,5] *

[1] Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, Ann Arbor, MI, 48109, USA.

[2] Department of Internal Medicine, Gastroenterology and Hepatology, Michigan Medicine at the University of Michigan, Ann Arbor, MI, 48109, USA.

[3] Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI, USA.

[4] Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI 48109, USA.

[5] U-M Center for Drug Repurposing, University of Michigan, Ann Arbor, MI, 48109, USA.

* Corresponding author: jzsexton@med.umich.edu

Abstract

Niclosamide, an FDA-approved oral anthelmintic drug, has broad biological activity including anticancer, antibacterial, and antiviral properties. Niclosamide has also been identified as a potent inhibitor of SARS-CoV-2 infection *in vitro,* generating interest in its use for the treatment or prevention of COVID-19. Unfortunately, there are several potential issues with using niclosamide for COVID-19, including low bioavailability, significant polypharmacology, high cellular toxicity, and unknown efficacy against emerging SARS-CoV-2 variants of concern. In this study, we used high-content imaging-based immunofluorescence assays in two different cell models to assess these limitations and evaluate the potential for using niclosamide as a COVID-19 antiviral. We show that despite promising preliminary reports, the antiviral efficacy of niclosamide overlaps with its cytotoxicity giving it a poor *in vitro* selectivity index for anti-SARS-CoV-2 inhibition. We also show that niclosamide has significantly variable potency against the different SARS-CoV-2 variants of concern and is most potent against variants with enhanced cell-to-cell spread including B.1.1.7. Finally, we report the activity of 33 niclosamide analogs, several of which have reduced cytotoxicity and increased potency relative to niclosamide. A preliminary structure-activity relationship analysis reveals dependence on a protonophore for antiviral efficacy, which implicates nonspecific endolysosomal neutralization as a dominant mechanism of action. Further single-cell morphological profiling suggests niclosamide also inhibits viral entry and cell-to-cell spread by syncytia. Altogether, our results suggest that niclosamide is not an ideal candidate for the treatment of COVID-19, but that there is potential for developing improved analogs with higher clinical translational potential in the future.

Key Words: Niclosamide, SARS-CoV-2, COVID-19, Polypharmacology, Drug Repurposing

Introduction

Since it emerged as a novel betacoronavirus in late 2019, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has caused a global pandemic[1]. COVID-19, the disease caused by SARS-CoV-2 infection, presents as varying symptoms with different degrees of severity ranging from dry cough and difficulty breathing to acute cardiac injury and refractory pulmonary failure[2,3]. As of April 2022, COVID-19 has caused the death of over six million individuals worldwide[4] and this death toll continues to increase as new SARS-CoV-2 Variants of Concern (VOCs) emerge with enhanced transmissibility and increased adaptive immune escape[5].

The deadly impact of COVID-19 has created a need to identify potential antiviral treatments for SARS-CoV-2 infection. This has culminated in the development and FDA-authorization/approval of several vaccines[6], and three small-molecule antiviral medications including remdesivir[7], molnupiravir[8], and Paxlovid[9]. Unfortunately, because of limited worldwide vaccine availability[10], the modest clinical efficacy of existing antivirals[8,9,11,12], and the potential resistance of SARS-CoV-2 variants[13,14], additional therapeutics are urgently needed to help stop the spread of the virus.

A promising strategy for identifying new therapies with the potential for rapid deployment is drug repurposing, whereby compounds with already established safety profiles and robust supply chains are used to treat other diseases[15]. Since the start of the pandemic, several large-scale drug repurposing screens have been conducted[16–20] and have identified many different potential candidates for the treatment of COVID-19. One of the repurposed drugs, which had potent anti-SARS-CoV-2 efficacy *in vitro,* was the oral anthelmintic compound niclosamide[17,20].

Traditionally used to treat tapeworm infection, niclosamide has been often repurposed in treating

a wide range of diseases including several cancers, bacterial infections, viral infections, type 2

diabetes, non-alcoholic fatty liver disease, and rheumatoid arthritis[21]. This range of potential uses

is due to the significant polypharmacology of niclosamide, which is known to act on many

different biological targets and is a modulator of the Wnt/b-catenin, mTOR and JAK/STAT3

signaling pathways among others[21]. Niclosamide is also a weakly acidic lipophilic protonophore

that can disrupt pH gradients by shuttling protons across lipid membranes[22] including

mitochondria and lysosomes/endosomes. This physiochemical property is responsible for its

activity as a mitochondrial uncoupler[23] and contributes to its broad activity against viruses, many

of which rely on endosomal-cytoplasmic pH gradients in their life cycle[24].

Niclosamide was also previously identified as a potential antiviral for the related coronavirus

SARS-CoV, where it was shown to inhibit viral replication *in vitro* with low micromolar

potency[25]. The mechanism of action (MOA) for niclosamide against SARS-CoV-2 may be more

complex and multimodal than for SARS-CoV. Niclosamide has been shown to inhibit SARS-

CoV-2 endocytosis mediated entry[26], block viral replication by promoting cellular autophagy[27],

and disrupt spike (S) protein-mediated syncytia formation via inhibition of the host cell calcium-

dependent scramblase TMEM16F[28]. Given the polypharmacology of niclosamide, it is likely that

there are additional factors that contribute to its overall efficacy and complicate its MOA. The

degree to which each of these MOAs plays a role in the antiviral efficacy of niclosamide against

SARS-CoV-2 is unclear.

While niclosamide was clinically effective as an anthelmintic drug, it has substantial limitations

for use as a COVID-19 antiviral including its low oral bioavailability (<10%) and poor water

solubility[29,30]. Oral administration of niclosamide at 5 mg/kg in rats reaches a maximal serum

concentration (C_{max}) of only 354 ± 152 ng/mL[31]. As a result, the concentration of niclosamide in

the lungs would likely be too low to achieve therapeutic effect. Another limitation exists in that

polypharmacology is generally associated with increased adverse effects for repurposed drugs[32].

Finally, the toxicity of niclosamide is a major concern as it has been previously repurposed as a

broad anti-cancer agent and has shown significant cytotoxic/cytostatic effects *in vitro*[33,34]. A

better understanding of the anti-SARS-CoV-2 mechanism of action for niclosamide, including

potential toxicity and activity against emerging variants of concern (VOC), is needed to

effectively evaluate its clinical potential.

The goal of this study was to expand upon the understanding of niclosamide anti-SARS-CoV-2

activity and its potential as a clinical therapeutic using high content fluorescence imaging and

analysis. Herein, we reveal some of the mechanistic and cell morphological characteristics of

niclosamide activity, including an analysis of its cellular toxicity after long-term exposure to a

therapeutic antiviral dose. Additionally, we investigate niclosamide antiviral activity against

SARS-CoV-2 viral VOCs. We hypothesized that because the fusogenicity amongst SARS-CoV-

2 variants is known to be different[35–37], the potency/efficacy of niclosamide may also vary

between strains. Here we reveal the potency of niclosamide against several variants including

WA1 (wildtype), B.1.1.7 (alpha), B.1.351 (beta), P.1 (gamma) and B.1.617.2 (delta). Lastly, we

report the *in vitro* results of a structure-activity relationship (SAR) campaign of 33 niclosamide

analogs against SARS-CoV-2 in two different cell lines (VeroE6 and H1437). We used the

results from this SAR campaign to reveal additional mechanistic details for niclosamide, and aid

with the identification of structural analogs with reduced cellular toxicity. Altogether, our results

suggest that niclosamide itself is not a suitable candidate for the treatment of COVID-19, yet

there is potential for developing analogs with improved properties for future clinical use.

Results

Niclosamide has a poor selectivity index

One major concern regarding the utility of niclosamide as a COVID-19 antiviral is its

cytotoxicity in comparison with its antiviral efficacy. Here, we aimed to determine the selectivity

index (SI) of niclosamide *in vitro* against SARS-CoV-2 in two different cell models for

infection, VeroE6 and the human lung adenocarcinoma cell line H1437. To assess compound-

related toxicity, we evaluated the effects of niclosamide on cells after 72 hours of compound

exposure. We designed and optimized two separate high-content fluorescence imaging assays in

384-well plate format using the different cell lines and measured cell viability and viral

inhibition concurrently. In both assays, we used the detection of viral nucleocapsid (N) protein as

a direct marker for SARS-CoV-2 infection and cell count per well as an indicator of cell

viability. As summarized in the **Figure 1A** workflow, VeroE6 or H1437 cells were preincubated

with a 10-point 2-fold dilution series of niclosamide (N=10 replicates per condition) for 24 hours

and then infected with the SARS-CoV-2 B.1.1.7 variant for an additional 48 hours post-infection

(p.i.). Following infection, cells were fixed, permeabilized, and stained to identify nuclei and

viral N protein. Assay plates were imaged at 10X magnification using a CX5 high content

imaging platform (N=9 fields captured per well) and processed using the image segmentation

and analysis software CellProfiler. Data from the CellProfiler output were used to determine

percent infection and percent viability. Infection data were normalized to the average well-level

% N positive for infected controls (mock) in each cell line (**Supplementary Figure S1A**).

Percent viability was determined by normalizing the average well-level cell counts for the

infected control (**Supplementary Figure S1B**). We found that niclosamide has potent 50%

maximal inhibition (IC_{50}) values of 564 nM for VeroE6 and 261 nM for H1437. However,

niclosamide caused a 50% reduction in cell viability (CC_{50}) at concentrations of 1050 nM and

438 nM for VeroE6 and H1437, respectively, resulting in poor selectivity indices in both cell

lines (1.86 for VeroE6 and 1.67 for H1437). The concentration-response curves for this

experiment are shown in **Figure 1B** along with representative images for infected control, mock,

and 10 μM niclosamide conditions for each cell line. As illustrated in **Figure 2B** and

Supplementary **Figure 1A**, the average percentage of N protein-positive cells in untreated

infected controls after 48 hours of infection was significantly higher in VeroE6 (69 %) than

H1437 (9%) indicating more efficient cell-to-cell spread in the former. In conclusion,

niclosamide has a low SI in two cell lines of fibroblast origin, representing a liability for

therapeutic use.

Niclosamide potency is SARS-CoV-2 variant dependent

Niclosamide has a complex polypharmacology profile against host-cell pathways, which may

contribute to the antiviral efficacy and/or cytotoxicity of the compound and lead to variable

responses across SARS-CoV-2 VOCs that rely differentially on these pathways. To evaluate the

antiviral efficacy of niclosamide against VOCs, we used a modified infection assay in VeroE6.

Exposure to niclosamide was reduced to a 1- hour preincubation and the assay window was

shortened to 24 hours post-infection (**Figure 2A**) to limit compound toxicity. VeroE6 cells were

used as they demonstrated a higher N-protein positivity rate than H1437 cells. We evaluated the

antiviral activity of niclosamide against the WA1 (wildtype), B.1.1.7 (alpha), B.1.351 (beta), P.1

(gamma), and B.1.617.2 (delta) variants in 10-point, 2-fold dilution. 10-Point dose-response

efficacy experiments showed niclosamide had statistically significant differences in efficacy

against VOCs, was most potent against the B.1.1.7 strain (IC_{50} = 298 nM), and least potent

against the WA1 strain (IC_{50} = 1664 nM). The full efficacy data for all variants are shown in

Figures 2B-C. The CC_{50} for niclosamide in this shortened assay was > 10 µM (not shown).

These data demonstrate variant dependent antiviral efficacy of niclosamide.

High-content analysis suggests inhibition of entry and syncytia formation

Cell morphologic analysis of cells infected with VOCs, under the treatment of niclosamide,

revealed several defining characteristics of infection influenced by compound treatment. We

quantified these observations using morphological cell profiling analysis. We used data (B.1.1.7

variant in VeroE6) from the viral control (mock) and three different efficacious concentrations of

niclosamide around its IC_{50} (156 nM, 313 nM, 625 nM) to reanalyze using a more extensive

analysis pipeline that included intensity and area/shape measurements for both nuclear and viral

channels. For this analysis, syncytia/individually infected cells were defined as "viral objects."

We determined that treatment with niclosamide decreased the maximum size of syncytia (**Figure**

3A) consistent with an inhibition of cell-to-cell spread. We also found that treatment reduced the

number of individually infected cells within a well (**Figure 3B**) consistent with an inhibition of

viral entry. Finally, we found that the N protein intensity of remaining viral objects increased

with escalating concentrations of niclosamide (**Figure 3C**). The combination of these

observations suggests multiple MOA including inhibition of viral entry and cell-to-cell spread

resulting in fewer infected cells with dramatically increased cellular viral N protein content.

These results provide support for the polypharmacology of niclosamide that contributes to

multimodal efficacy against SARS-CoV-2. In addition, these results suggest that niclosamide

may not directly inhibit viral replication *in vitro*.

Structure-activity relationship of niclosamide analogs versus SARS-CoV-2 infection

Efficacy and cytotoxicity of 33 previously designed analogs[38] of niclosamide were used to

establish a preliminary structure-activity relationship (SAR) profile for niclosamide for anti-

SARS-CoV-2 activity in VeroE6 and H1437 cell lines. These analogs were also salicylanilides

and had substituent modifications on the nitroaniline and/or chlorosalicyl rings (**Figure 4A**)

intending to improve the selectivity while maintaining antiviral efficacy. Analog structures are

shown in **Supplementary Figure S2**. Analogs were evaluated using the assay described in

Figure 1A. All analogs were tested in 10-point 2-fold dilution series (N=3) from a starting

concentration of 20 μM. VeroE6 analog screening was performed using the B.1.1.7 variant at an

MOI of 0.1, while H1437 screening was performed using the WA1 variant at an MOI of 1. The

results from compound testing are summarized in **Table 1**, which includes IC_{50} and CC_{50} values

for both cell lines. As shown in Table 1, we found that 12 analogs retained IC_{50} values in the

nanomolar or micromolar range in VeroE6, while 8 compounds were efficacious in both VeroE6

and H1437. Four compounds (2, 7, 11, 24) showed improved potency and reduced cytotoxicity

against VeroE6 compared to niclosamide (**Figure 4C-F**). Overall, improvements in cytotoxicity

were less pronounced against H1437. Notably, the variant dependent potency difference was

conserved in H1437 cells and was significantly less potent against WA1 ($IC_{50} = 16770$ nM) than

B.1.1.7 ($IC_{50} = 261$ nM), consistent with results in VeroE6 cells.

In general, we found that the replacement of the nitro group on the nitroaniline ring was well

tolerated (compounds 5, 6, 7, 24 and 34) and improved the selectivity index. We also noted that

modification to the chloro position on the salicylic acid ring (R2) was well tolerated and all

compounds with only this modification retained antiviral efficacy (compounds 2, 3, 4, 11, 12).

Many analogs (compounds 7,8 12-19, 25, 28, 29, and 33) were found to exacerbate infection in

H1437 cells (**Supplementary Figure S3**) and showed inverted concentration-response curves at

high concentrations. Remarkably, the removal of the hydroxyl group (R1) on the salicylic acid

ring (compound 9) resulted in a complete loss of activity in both VeroE6 and H1437 (**Figure**

5A). This hydroxyl group has been previously reported as the protonophore responsible for the

mitochondrial uncoupling activity of niclosamide[23,38].

Given the drastic loss of activity, we evaluated the anti-SARS-CoV-2 efficacy of other

protonophore mitochondrial uncouplers including FCCP[39], 2,4 DNP[40], oxyclozanide[41] and

dicumarol[42]. These compounds were evaluated against WA1 and B.1.1.7 variants in VeroE6 cells

using the 24-hour infection conditions described in **Figure 3A**. Both FCCP and oxyclozanide

showed efficacy in the micromolar range (**Figure 5B-F**). The potency of these compounds was

higher against B.1.1.7 than WA1, however the difference was more pronounced for niclosamide.

These results indicate that the mechanism of action for niclosamide against SARS-CoV-2 is at

least partially due to its physiochemical property as a protonophore, implicating energetic stress

response pathways in SARS-CoV-2 infection.

Discussion

There remains an urgent need for COVID-19 therapeutics, which can be used to prevent or treat

the spread of the virus SARS-CoV-2. The FDA approved oral anthelmintic drug niclosamide has

antiviral activity against SARS-CoV-2 infection *in vitro* and *in* vivo[43], which has generated

interest in its application for the treatment of COVID-19 and resulted in the conductance of

several human clinical trials. However, given its high cytotoxicity, unknown efficacy against

SARS-CoV-2 variants, low systemic bioavailability, and significant polypharmacology, we were

hesitant to consider niclosamide as a promising antiviral option. In this study, we used high-

content imaging of SARS-CoV-2 infected cells to evaluate some of the limitations of

niclosamide as a COVID-19 antiviral. We also extended our studies to structural analogs of

niclosamide, intending to reveal a preliminary structure-activity relationship profile that could be

used for future compound development.

Niclosamide has potent cytotoxic/cytostatic effects when applied directly to cells *in vitro*[34],

suggesting that it may have high acute toxicity *in vivo* with increased systemic exposure.

Clinically, the cytotoxicity is limited by the poor bioavailability of niclosamide, which has low

systemic exposure. To evaluate niclosamide toxicity, we used high-content fluorescence imaging

to determine a selectivity index for niclosamide in two different cell models including VeroE6

and the more physiologically relevant human lung adenocarcinoma cell line H1437. We found

that niclosamide has a very poor selectivity index in both cell lines (SI <2) after 72 hours of

compound exposure, suggesting that it would likely have a small therapeutic window clinically

even if the compound exposure was high enough in the lungs for antiviral efficacy. Longer

durations of exposure to niclosamide at relevant antiviral concentrations are likely to cause

significant side effects, which limits clinical application. Further studies are needed to evaluate

the safety of niclosamide at antiviral concentrations *in vivo*.

Our results are consistent with the results from recent clinical studies of niclosamide. Since it

was identified as an anti-SARS-CoV-2 agent *in vitro*, there have been several clinical studies to

evaluate the antiviral efficacy and safety of niclosamide. Notably, a recent phase 2 clinical trial

using 2g of orally administered niclosamide for 7 days revealed no statistically significant effect

on the duration of the contagious period of SARS-CoV-2[44]. While niclosamide was well

tolerated in this trial, the low efficacy and low adverse event rate are likely because the systemic

exposure is lower than what is required to observe antiviral activity or compound-related

toxicity. To address the poor oral bioavailability, several different formulations have been

developed for niclosamide to improve its exposure to the necessary site of action[45,46]. This has

included a formulation as an inhalable/intranasal powder to increase compound exposure in the

lungs. Unfortunately, a recent phase-1 safety trial using 50 mg over 2.5 days of

inhalable/intranasal niclosamide revealed moderate lung irritation in 59% of participants, which

suggests compound-related toxicity may be playing a significant role at higher local

concentrations in the lungs[47]. Although niclosamide is generally well-tolerated when used as an

anthelmintic drug, this is because it has low bioavailability, poor solubility, and stays within the

GI tract with very low systemic exposure.

A further limitation for using niclosamide as a COVID-19 therapeutic is its unknown efficacy

against the different emerging SARS-CoV-2 VOCs. We determined the efficacy for niclosamide

against the WA1 (wildtype), B.1.1.7 (alpha), B.1.351 (beta), P.1 (gamma) and B.1.617.2 (delta)

variants in VeroE6 cells. We found that there were significant differences in potency ranging

from 298 nM (beta variant) to 1664 nM (wildtype). Interestingly, the trend in potency correlates

with the ACE2 binding affinity for the different variants[48]. Variants, including alpha and beta,

also have higher fusogenicity than the wildtype variant and are more likely to undergo cell-to-

cell spread by syncytia[35], which may help explain the differences in potency.

These results are in contrast with those reported by Weiss *et al.*, which showed no significant

difference in potency amongst variants[49]. However, their study used qRT-PCR of viral RNA to

determine IC_{50} values, which is far less sensitive than a high-content imaging approach and does

not provide information on the clinically relevant endpoint of cell-to-cell spread inhibition.

Given the differences in potency amongst SARS-CoV-2 variants of concern, there arises a

concern for the rapid development or selection of resistant strains that do not respond to

niclosamide treatment. While the emergence of drug resistance is possible for any mechanism of

action inhibiting SARS-CoV-2, the pronounced difference between niclosamide's efficacy

amongst the VOCs makes niclosamide resistance inexorable. Further studies to understand the

mechanistic differences underlying variant-dependent responses to drugs like niclosamide may

ultimately inform *de novo* drug development for COVID-19.

To understand the MOA, we used morphological profiling of B.1.1.7 infected VeroE6 cells to

evaluate the effect of niclosamide treatment on SARS-CoV-2 infection. We found that

niclosamide inhibits the spread of virus to adjacent cells in a concentration-dependent fashion as

indicated by the reduction in size of viral syncytia. We also observed that the total number of

viral objects (individually infected cells or syncytia) decreased with niclosamide treatment,

which is consistent with entry inhibition. For example, if niclosamide were only influencing cell-

to-cell spread, the total number of viral objects would remain constant and only the size of the

syncytia would be affected. While the complete mechanism of action for niclosamide is

complex, our results suggest that both inhibition of cell-to-cell spread and entry inhibition play a

role in its activity (**Figure 6**). The degree to which each of the MOAs contributes to efficacy may

be different for SARS-CoV-2 variants, which could help explain the differences in potency.

The polypharmacology of niclosamide is a major issue for its utility as a COVID-19 antiviral.

Niclosamide is known to influence many different signal transduction pathways and has been

implicated in the treatment of a wide range of diseases including several cancers, bacterial

infections, viral infections, type 2 diabetes, non-alcoholic fatty liver disease, rheumatoid arthritis,

and others. Unfortunately, the mechanism of action for niclosamide remains elusive for the

majority of its biological effects. It is often unclear if there is a direct interaction between

niclosamide and a molecular target, or if there is an indirect mechanism of action at play[21]. An

underlying mechanism for its broad activity may be its ability to act as a protonophore, which

has many different downstream effects in cells including disruption of pH gradients,

mitochondrial uncoupling, and transcriptional modulation of various gene targets[21]. This

mechanistic ambiguity also translates to its antiviral efficacy. It is likely that the antiviral

activities of niclosamide (e.g., inhibition of entry, replication, and syncytia formation) are all

downstream consequences of its activity as a nonspecific protonophore since activity was lost

following removal of the hydroxyl group. If this is the case, then it may be challenging to

separate the undesired off-target effects from the antiviral effects. While our studies suggest that

niclosamide and other mitochondrial uncouplers demonstrate anti-SARS-CoV-2 efficacy, further

studies are warranted to determine if these mechanisms of action are unified by protonophore

activity.

While niclosamide is not an ideal candidate itself, it may represent a promising chemical tool for

the development of more specific SARS-CoV-2 inhibitors. In particular, inhibition of cell-to-cell

spread by syncytia is an extremely attractive mechanism for inhibition. Syncytia, which are

multinucleated bodies resulting from the fusion of adjacent cells, are a key characteristic of

SARS-CoV-2 infection and have been observed in many post-mortem histological samples from

fatal COVID-19 cases[28,50]. Syncytia formation facilitates the rapid spread of the viral genome

between cells[51], which increases the area of infected tissue and may enhance immune system

evasion[52]. An inhibitor of cell-to-cell infection like niclosamide may be clinically useful for the

treatment or prevention of COVID-19, especially when cocktailed with other direct acting

antivirals.

In this study, we also tested the antiviral efficacy of 33 structural analogs of niclosamide to

establish a preliminary structure-activity relationship profile which could aid in the development

of compounds with antiviral efficacy and less off-target effects. We identified eight compounds

(compounds 2, 3, 4, 10, 11, 22, 24, 34) that were efficacious in both VeroE6 and H1437 cell

models and four of which had improved potency and reduced cytotoxicity in VeroE6

(Compounds 2,7,11 and 24). Consequently, we believe there is a potential for designing better

niclosamide analogs with improved properties. Additionally, our structure-activity analysis

revealed some mechanistic features of niclosamide. Most noteworthy, the removal of the

protonophore hydroxyl group resulted in complete loss of activity in both cell models. The

efficacy of analogs strongly relied on their weakly acidic and lipophilic nature. Analogs with

higher predicted acidity (pKa) due to the presence of carboxylic acid substituents were generally

completely inactive. We also determined that other protonophores, including FCCP and

oxyclozanide, also had anti-SARS-CoV-2 efficacy, suggesting that the ability to disrupt pH

gradients is central to the mechanism of action for niclosamide. Niclosamide has been shown to

neutralize endo-lysosomal pH gradients, which is believed to be responsible for its broad-

spectrum antiviral activity[24]. Our results indicate that this nonspecific mechanism of action also

significantly contributes to the activity of niclosamide against SARS-CoV-2.

Overall, the poor selectivity index, low bioavailability, complex polypharmacology, nonspecific

protonophore activity, and variant-dependent potency of niclosamide limit its potential as a

COVID-19 therapeutic. However, our studies have shown that changes to the salicyl and aniline

rings can modulate selectivity and bioavailability while maintaining its activity. Therefore,

niclosamide represents a useful chemical probe that can be leveraged in a large-scale SAR

campaign to design better analogs in the future.

Methods

Compounds

Niclosamide, FCCP, 2,4 DNP, Oxyclozanide and Dicumarol were obtained from Sigma Aldrich and prepared as 10 mM stock solutions in dimethylsulfoxide (DMSO). The 33 structural analogs of niclosamide were obtained from previous studies[38]. Compounds were solubilized at 10 mM in DMSO, and were dispensed onto cells using an HPD300e digital compound dispenser.

Cells and Virus

VeroE6 cells were maintained in Dulbecco's Modified Eagle Medium (DMEM), and H1437 cells were maintained in RPMI 1640 base medium. Both cell lines were supplemented with 10% fetal bovine serum (FBS) and 1X pen-strep solution and were grown at 37°C with 5% CO_2 following standard cell culture procedures. These cell lines were tested for mycoplasma contamination before use and were negative. The following reagents were deposited by the Centers for Disease Control and Prevention and were obtained through BEI resources, NIAID, NIH: SARS-Related Coronavirus 2, Isolate USA-WA1/2020, NR-52281, USA/CA_CDC_5574/2020 (B.1.1.7), NR-54011, USA/MD-HP01542/2021 (Lineage B.1.351), NR-55282, Japan/TY7-503/2021 (Brazil P.1), NR-54982, USA/PHC658/2021 (Lineage B.1.617.2), NR-55611.Viral stocks were grown in VeroE6 and titers were determined by TCID50 using the Reed and Muench method[53]. All the work with live SARS-CoV-2 virus was performed in biosafety level-3 containment lab (BSL3) with the approval of the University of Michigan's Department of Environment and Health and Safety and the Institutional Biosafety Committee.

Anti-SARS-CoV-2 high content bioassays

Assays were adapted from previous work and optimized for H1437 and VeroE6 cell lines[17,54].

For 48-hour infection experiments, VeroE6 and H1437 cells were seeded onto 384 well plates

(6057300, Perkin Elmer) at densities of 3000 and 5000 cells per well, respectively, in 50 µL of

media. After 24 hours of cell attachment at 37°C and 5% CO_2, compounds were dispensed

directly to the cell plates using an HPD300e digital compound dispenser. All wells were

normalized to a constant DMSO concentration of 0.2%, and plates contained both infected and

uninfected control wells. After 24 hours of preincubation with compounds, cells were inoculated

with the indicated SARS-CoV-2 variant at MOIs of 0.1 for VeroE6 and 1 for H1437. Cells were

incubated with virus and compounds for an additional 48 hours and then fixed with 4%

paraformaldehyde for 30 minutes at room temperature. Cells were then permeabilized with 0.3%

Triton-X100 for 15 minutes and stained with anti-nucleocapsid protein primary antibody

(ABIN6952432, Antibodies Online) at a dilution of 1:2000 overnight at +4°C. Following

primary antibody staining, cells were stained with a dye cocktail containing 1:1000 secondary

antibody Alexa-647 (goat anti-mouse, A21235, Thermo Fisher) and 10 µg/mL Hoechst 33342

pentahydrate (bis-benzimide) for nuclear labeling for a total of 30 minutes at room temperature.

Cells were stored in PBS before imaging. For 24-hour infection experiments, the methods were

comparable except that VeroE6 cells were seeded at 5000 cells per well, compounds were

preincubated for 1 hour instead of 24, and the infection window was 24 hours instead of 48. All

other inoculation, fixation and staining procedures were identical.

High Content Imaging

Stained assay plates were imaged using a both a Thermo Fisher CX5 with a 10X/0.45NA

objective lens and a Yokogawa Cell Voyager 8000 (CV8000) microscope with a 20X/1.0NA

water immersion lens. Imaging techniques were followed as described previously for detection

of nuclei and SARS-CoV-2 nucleocapsid protein[17,54]. A total of N=9 fields per well were imaged

for all assay plates, accounting for roughly 80% of the total well area.

Image Processing

Images were processed using the image segmentation and analysis software CellProfiler 4.0[55].

Separate pipelines were developed for H1437 and VeroE6 images. Pipelines were used to

identify nuclei (Hoechst 33342) and viral objects including multinucleated syncytia and

individually infected cells (Alexa Fluor 647) by adaptive otsu thresholding. Similar to previous

work, infected cells were identified using the *relateobjects* module whereby any nucleus

contained within a viral object was defined as infected[54]. For morphological profiling of B.1.1.7

infection vs. niclosamide in VeroE6, additional intensity, textural and spatial features were

measured using CellProfiler 4.0 for both the nuclear and viral channels.

Concentration response analysis and IC_{50}/CC_{50} determination

Field level data were grouped at the well level using Knime[56] and used to determine normalized

percent infection and percent viability scores. Raw percent infection per well was determined by

taking the ratio of infected nuclei to total nuclei and multiplying by 100. Normalized percent

infection was then generated such that "100% infection" was equivalent to the average raw

percent infection of the viral control for each plate. Cell counts for the entire plate were

normalized and 100% viability was based on the average cell count of the infected DMSO

control wells. Concentration-response curves were plotted in GraphPad Prism 9.0 (GraphPad

Software) and fitted using a semi-log 4-parameter variable slope model. IC_{50} and CC_{50} values were

extracted from percent infection curves and percent viability curves, respectively. Selectivity

indices were determined by taking a ratio of the CC_{50} and IC_{50}.

High content imaging analysis of B.1.1.7 infection versus niclosamide

Object level data for nuclei and viral objects (syncytia and individually infected cells) was used to evaluate morphological and phenotypic features of B.1.1.7 infection versus niclosamide including changes in N protein intensity and area of viral objects. Only images for the infected DMSO vehicle control, as well as 3 different concentrations of niclosamide (156 nM, 313 nM and 625 nM) were included in this analysis. The max viral object area reported in Figure 3A represents the largest viral object observed for each condition including all fields and replicate wells. The mean N protein intensity for infected cells was computed at the object level and the results from Figure 3C include data for cells in each condition.

Statistical analysis and hypothesis testing

All statistical analyses and hypothesis testing was performed using GraphPad Prism 9.0 (GraphPad software). Specifics for statistical analyses, including sample sizes and other important data are included within the text of figure legends.

Acknowledgements

We acknowledge funding from the University of Michigan Institute for Clinical and Health Research (MICHR; National Center for Advancing Translational Science Grant UL1TR002240) and the U-M Center for Drug Repurposing. JZS is supported by the National Institute of Diabetes and Kidney Diseases (R01DK120623). CEW is supported in part by The University of Michigan Biological Scholars Science Program. JWW is supported by an American Foundation for Pharmaceutical Education (AFPE) regional award. CM is supported by European funding MSCA actions, GA481247. SRM is supported by a pharmacological sciences training program (PSTP) T32 training grant. We thank Tracey Schultz for thoughtful discussions and assistance

437   with cell culture. Figures were generated using Biorender.com. The authors declare no conflicts

438   of interest.

References

1.   Hu, B., Guo, H., Zhou, P. & Shi, Z. L. Characteristics of SARS-CoV-2 and COVID-19. *Nature Reviews*
Microbiology 2020 19:3 **19**, 141–154 (2020).

2.   Aslan, A., Aslan, C., Zolbanin, N. M. & Jafari, R. Acute respiratory distress syndrome in COVID-19:
possible mechanisms and therapeutic management. *Pneumonia 2021 13:1* **13**, 1–15 (2021).

3.   Mizrahi, B. *et al.* Longitudinal symptom dynamics of COVID-19 infection. *Nature Communications*
2020 11:1 **11**, 1–10 (2020).

4.   Dong, E., Du, H. & Gardner, L. An interactive web-based dashboard to track COVID-19 in real
time. *Lancet Infect Dis* **20**, 533–534 (2020).

5.   Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat Rev Microbiol*
19, 409–424 (2021).

6.   Kyriakidis, N. C., López-Cortés, A., González, E. V., Grimaldos, A. B. & Prado, E. O. SARS-CoV-2
vaccines strategies: a comprehensive review of phase 3 candidates. *npj Vaccines 2021 6:1* **6**, 1–17
(2021).

7.   Cully, M. A tale of two antiviral targets - and the COVID-19 drugs that bind them. *Nat Rev Drug*
Discov **21**, 3–5 (2022).

8.   Whitley, R. Molnupiravir — A Step toward Orally Bioavailable Therapies for Covid-19. *New*
England Journal of Medicine **386**, 592–593 (2022).

9.   Hammond, J. *et al.* Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19. *New*
England Journal of Medicine **386**, 1397–1408 (2022).

10.   Feinmann, J. Covid-19: global vaccine production is a mess and shortages are down to more than
just hoarding. *BMJ* **375**, (2021).

11.   Qomara, W. F., Primanissa, D. N., Amalia, S. H., Purwadi, F. v. & Zakiyah, N. Effectiveness of
Remdesivir, Lopinavir/Ritonavir, and Favipiravir for COVID-19 Treatment: A Systematic Review.
Int J Gen Med **14**, 8557–8571 (2021).

12.   Gottlieb, R. L. *et al.* Early Remdesivir to Prevent Progression to Severe Covid-19 in Outpatients. *N*
Engl J Med **386**, 305–315 (2022).

13.   Chen, R. E. *et al.* Resistance of SARS-CoV-2 variants to neutralization by monoclonal and serum-
derived polyclonal antibodies. *Nature Medicine 2021 27:4* **27**, 717–726 (2021).

14.   Takashita, E. *et al.* Efficacy of Antiviral Agents against the SARS-CoV-2 Omicron Subvariant BA.2.
New England Journal of Medicine **386**, 1475–1477 (2022).

15.   Pushpakom, S. *et al.* Drug repurposing: progress, challenges and recommendations. *Nature*
471   *Reviews Drug Discovery 2018 18:1* **18**, 41–58 (2018).

472   16.   Drayman, N. *et al.* Masitinib is a broad coronavirus 3CL inhibitor that blocks replication of SARS-
473   CoV-2. *Science (1979)* **373**, 931–936 (2021).

474   17.   Mirabelli, C. *et al.* Morphological cell profiling of SARS-CoV-2 infection identifies drug repurposing
475   candidates for COVID-19. *Proc Natl Acad Sci U S A* **118**, (2021).

476   18.   Touret, F. *et al.* In vitro screening of a FDA approved chemical library reveals potential inhibitors
477   of SARS-CoV-2 replication. *Scientific Reports* **10**, 13093 (2020).

478   19.   Xiao, X. *et al.* Identification of Potent and Safe Antiviral Therapeutic Candidates Against SARS-
479   CoV-2. *Frontiers in Immunology* **11**, (2020).

480   20.   Jeon, S. *et al.* Identification of antiviral drug candidates against SARS-CoV-2 from FDA-approved
481   drugs. *Antimicrobial Agents and Chemotherapy* **64**, (2020).

482   21.   Chen, W., Mook, R. A., Premont, R. T. & Wang, J. Niclosamide: Beyond an antihelminthic drug.
483   *Cellular Signalling* **41**, 89–96 (2018).

484   22.   Fonseca, B. D. *et al.* Structure-activity analysis of niclosamide reveals potential role for
485   cytoplasmic pH in control of mammalian target of rapamycin complex 1 (mTORC1) signaling. *J*
486   *Biol Chem* **287**, 17530–17545 (2012).

487   23.   Kumar, R. *et al.* Mitochondrial uncoupling reveals a novel therapeutic opportunity for p53-
488   defective cancers. *Nature Communications 2018 9:1* **9**, 1–13 (2018).

489   24.   Jurgeit, A. *et al.* Niclosamide Is a Proton Carrier and Targets Acidic Endosomes with Broad
490   Antiviral Effects. *PLoS Pathogens* **8**, 1002976 (2012).

491   25.   Wu, C. J. *et al.* Inhibition of Severe Acute Respiratory Syndrome Coronavirus Replication by
492   Niclosamide. *Antimicrobial Agents and Chemotherapy* **48**, 2693 (2004).

493   26.   Prabhakara, C. *et al.* Strategies to target SARS-CoV-2 entry and infection using dual mechanisms
494   of inhibition by acidification inhibitors. *PLOS Pathogens* **17**, e1009706 (2021).

495   27.   Gassen, N. C. *et al.* SARS-CoV-2-mediated dysregulation of metabolism and autophagy uncovers
496   host-targeting antivirals. *Nature Communications 2021 12:1* **12**, 1–15 (2021).

497   28.   Braga, L. *et al.* Drugs that inhibit TMEM16 proteins block SARS-CoV-2 spike-induced syncytia.
498   *Nature 2021 594:7861* **594**, 88–93 (2021).

499   29.   Lin, C. K. *et al.* Preclinical evaluation of a nanoformulated antihelminthic, niclosamide, in ovarian
500   cancer. *Oncotarget* **7**, 8993 (2016).

501   30.   Fan, X., Li, H., Ding, X. & Zhang, Q. Y. Contributions of Hepatic and Intestinal Metabolism to the
502   Disposition of Niclosamide, a Repurposed Drug with Poor Bioavailability. *Drug Metab Dispos* **47**,
503   756–763 (2019).

31. Chang, Y. W. *et al.* Pharmacokinetics of anti-SARS-CoV agent niclosamide and its analogs in rats.
Journal of Food and Drug Analysis **14**, 15 (2020).

32. Reddy, A. S. & Zhang, S. Polypharmacology: drug discovery for the future. *Expert Rev Clin*
Pharmacol **6**, 41–47 (2013).

33. Wieland, A. *et al.* Anticancer effects of niclosamide in human glioblastoma. *Clin Cancer Res* **19**,
4124–4136 (2013).

34. Hamdoun, S., Jung, P. & Efferth, T. Drug repurposing of the anthelmintic niclosamide to treat
multidrug-resistant leukemia. *Frontiers in Pharmacology* **8**, (2017).

35. Rajah, M. M. *et al.* SARS-CoV-2 Alpha, Beta, and Delta variants display enhanced Spike-mediated
syncytia formation. *EMBO J* **40**, (2021).

36. Saito, A. *et al.* Enhanced fusogenicity and pathogenicity of SARS-CoV-2 Delta P681R mutation.
Nature **602**, 300–306 (2022).

37. Zhang, Y. *et al.* SARS-CoV-2 spike L452R mutation increases Omicron variant fusogenicity and
infectivity as well as host glycolysis. *Signal Transduction and Targeted Therapy 2022 7:1* **7**, 1–3
(2022).

38. Bhagat, H. A. *et al.* N-substituted phenylbenzamides of the niclosamide chemotype attenuate
obesity related changes in high fat diet fed mice. *PLOS ONE* **13**, e0204605 (2018).

39. Park, K. S. *et al.* FCCP depolarizes plasma membrane potential by activating proton and Na+
currents in bovine aortic endothelial cells. *Pflugers Archiv : European journal of physiology* **443**,
344–352 (2002).

40. Blaikie, F. H. *et al.* Targeting dinitrophenol to mitochondria: limitations to the development of a
self-limiting mitochondrial protonophore. *Biosci Rep* **26**, 231–243 (2006).

41. Shrestha, R., Johnson, E. & Byrne, F. L. Exploring the therapeutic potential of mitochondrial
uncouplers in cancer. *Molecular Metabolism* **51**, 101222 (2021).

42. Kotova, E. A. & Antonenko, Y. N. Fifty Years of Research on Protonophores: Mitochondrial
Uncoupling As a Basis for Therapeutic Action. *Acta Naturae* **14**, 4 (2022).

43. Blake, S. *et al.* Salicylanilides Reduce SARS-CoV-2 Replication and Suppress Induction of
Inflammatory Cytokines in a Rodent Model. *ACS Infectious Diseases* **7**, 2229–2237 (2021).

44. Cairns, D. M. *et al.* Efficacy of Niclosamide vs Placebo in SARS-CoV-2 Respiratory Viral Clearance,
Viral Shedding, and Duration of Symptoms Among Patients With Mild to Moderate COVID-19: A
Phase 2 Randomized Clinical Trial. *JAMA Network Open* **5**, e2144942–e2144942 (2022).

45. Parikh, M. *et al.* Phase Ib trial of reformulated niclosamide with abiraterone/prednisone in men
with castration-resistant prostate cancer. *Scientific Reports 2021 11:1* **11**, 1–7 (2021).

46. Lin, C. K. *et al.* Preclinical evaluation of a nanoformulated antihelminthic, niclosamide, in ovarian
cancer. *Oncotarget* **7**, 8993 (2016).

47. Backer, V. *et al.* A randomized, double-blind, placebo-controlled phase 1 trial of inhaled and
intranasal niclosamide: A broad spectrum antiviral candidate for treatment of COVID-19. *The*
Lancet Regional Health - Europe **4**, 100084 (2021).

48. Liu, H., Wei, P., Kappler, J. W., Marrack, P. & Zhang, G. SARS-CoV-2 Variants of Concern and
Variants of Interest Receptor Binding Domain Mutations and Virus Infectivity. *Frontiers in*
Immunology **13**, 50 (2022).

49. Weiss, A. *et al.* Niclosamide shows strong antiviral activity in a human airway model of SARS-CoV-
2 infection and a conserved potency against the Alpha (B.1.1.7), Beta (B.1.351) and Delta variant
(B.1.617.2). *PLOS ONE* **16**, e0260958 (2021).

50. Bussani, R. *et al.* Persistence of viral RNA, pneumocyte syncytia and thrombosis are hallmarks of
advanced COVID-19 pathology. *EBioMedicine* **61**, (2020).

51. Lin, L., Li, Q., Wang, Y. & Shi, Y. Syncytia formation during SARS-CoV-2 lung infection: a disastrous
unity to eliminate lymphocytes. *Cell Death & Differentiation 2021 28:6* **28**, 2019–2021 (2021).

52. Rajah, M. M., Bernier, A., Buchrieser, J. & Schwartz, O. The Mechanism and Consequences of
SARS-CoV-2 Spike-Mediated Fusion and Syncytia Formation. *Journal of Molecular Biology* **434**,
167280 (2022).

53. Reed, L. J. & Muench, H. A simple method of estimating fifty per cent endpoints. *American*
Journal of Epidemiology **27**, 493–497 (1938).

54. Wotring, J. W., Fursmidt, R., Ward, L. & Sexton, J. Z. Evaluating the in vitro efficacy of bovine
lactoferrin products against SARS-CoV-2 variants of concern. *J Dairy Sci* **105**, 2791–2802 (2022).

55. McQuin, C. *et al.* CellProfiler 3.0: Next-generation image processing for biology. *PLOS Biology* **16**,
e2005970 (2018).

56. Berthold, M. R. *et al.* KNIME - the Konstanz information miner. *ACM SIGKDD Explorations*
Newsletter 58–61 (2009) doi:10.1145/1656274.1656280.

Figures

Figure 1. Niclosamide is toxic at antiviral concentrations after long term exposure. A.)

Workflow for high content anti-SARS-CoV-2 bioassay screening to determine infection

inhibition and cytotoxicity. B.) 10-point, 2-fold dilution concentration-response curves for

VeroE6 and H1437 cells with a starting concentration of 10 μM. VeroE6 cells were infected with

SARS-CoV-2 B.1.1.7 variant at a multiplicity of infection (MOI) of 0.1, while H1437 were

infected with SARS-CoV-2 WA1 variant at an MOI =1 to achieve optimal infection at 48 hours

post-infection. Data points represent mean ± SEM for N=10 replicates per condition. Curve

fitting was performed in GraphPad Prism 9.0 using a semi-log 4-parameter variable slope model.

Representative overlay images for mock, vehicle, and 10 μM Niclosamide treatment (infected)

are included (cyan = nuclei, magenta = SARS-CoV-2 N protein).

Figure 2. Niclosamide potency is SARS-CoV-2 variant dependent. A.) Assay timeline for

24-hour infection experiment. The assay window was shortened to reduce niclosamide toxicity.

B.) 10-point 2-fold concentration-response curves for niclosamide against the different SARS-

CoV-2 variants of concern (MOI = 0.1 for each variant) with a top concentration of 10μM.

Curves were fitted with GraphPad Prism 9.0 software using a semi-log 4-parameter variable

slope model. Data for each variant was normalized to the average percent infected of its

respective viral control. Data points represent mean ± SEM for N=3 replicates. C.) IC$_{50}$ values

for niclosamide potency against SARS-CoV-2 variants of concern. Values were extracted from

curve fitting using GraphPad 9.0 and include SEM error bars (WA1: 1664 ± 149 nM, B.1.1.7:

298 ± 23 nM, B.1.351: 440 ± 21 nM, B.1.617.2: 774 ± 58 nM, P.1: 399 ± 34 nM). Significance

was determined using Student's T-tests (* = P<0.05, ** = P<0.01).

Figure 3. Morphological profiling of B.1.1.7 infection versus niclosamide treatment in

VeroE6. Image analysis reveals mechanistic characteristics of niclosamide activity against

SARS-CoV-2 infection. A.) The maximum area of viral objects decreases with increasing

niclosamide concentration. Data is the max area for viral objects in each condition. Viral control:

N=17452, +156 nM niclosamide: N=2425, +313 nM niclosamide: N= 1470, +625 nM

niclosamide: N= 496. B.) Viral objects per well decreases with increasing niclosamide

concentration. Viral objects include single infected cells and syncytia. Replicate values are

indicated on the X axis. C.) Mean pixel intensity for viral objects in each condition. Pixel

intensity increases with increasing niclosamide concentration. D.) Representative images for

each condition including N-protein channel, nuclear channel, an overlayed image and a fire

lookup table (LUT) image of the N-protein channel. Images were taken on a CX5 high content

microscope at 10X magnification. * = P<0.05, ***= P<0.001, ****= P<0.0001.

Figure 4. Niclosamide analogs have improved efficacy and reduced cytotoxicity in VeroE6.

A.) Structure of niclosamide indicating the chlorosalicyl/nitroaniline rings, and analog scaffold with modified substituent positions labeled. B.) IC_{50} vs. CC_{50} plot highlighting efficacious compounds in VeroE6. Compounds with improved potency and cytotoxicity profiles are circled on the plot. C.) 10-point, 2-fold concentration-response curves for the top four niclosamide analogs with a starting concentration of 20μM. Data are shown as the mean ± SEM of N=3 replicate wells per condition. Curves for infection (in red) and cell viability (in black) are included. D.) Representative images of infected cells treated with indicated compounds and viral control (Vehicle). (10X magnification, Cyan = nuclei, magenta = SARS-CoV-2 nucleocapsid protein).

Figure 5. Efficacy of protonophores against SARS-CoV-2. A.) Removal of protonophore

hydroxyl of niclosamide results in a complete loss of efficacy in both VeroE6 and H1437 cells.

B.) Chemical structures for other protonophores evaluated against SARS-CoV-2 infection. C-D.)

10-point, 2-fold concentration response curves for protonophores versus WA1 variant (C.) and

B.1.1.7 variant (D.) with starting concentrations of 100 µM. E.) IC_{50} plot for antiviral

protonophores indicating significantly different potency against variants. F.) Table of IC_{50} and

CC_{50} values for protonophores.

Figure 6. Diagram of niclosamides effect on SARS-CoV-2 entry and spike protein-mediated

syncytia formation. 1.) SARS-CoV-2 binds to the ACE2 receptor of the host cell and enters.

Niclosamide has been shown to inhibit this entry step *in vitro* 2.) Viral replication generates

many copies of the RNA genome. 3.) Infection results in an increased expression of viral spike

(S) protein and host cell TMEM16F at the plasma membrane. 4.) The S protein at the surface on

an infected cell binds to the ACE2 receptor of an adjacent uninfected cell. 5.) Spike-dependent

syncytia formation is mediated by the calcium-dependent lipid scramblase TMEM16F to

generate multinucleated infected cell bodies. Niclosamide, an inhibitor of TMEM16F, has been

shown to block spike-dependent syncytia formation.

Tables

Table 1. SAR table for Niclosamide Analogs. IC_{50} and CC_{50} values for VeroE6 (SARS-CoV-2 B.1.1.7 variant) and H1437 (SARS-CoV-2 WA1 variant). Physical properties including cLogP, pKa and logS were calculated using MOE and included in the table for each compound. Compounds that have efficacy against both cell lines are highlighted in gray.

| Compound # | R1 | R2 | R3 | R4 | R5 | R6 | R7 | Vero-E6 (B.1.1.7) | | H1437 (WA1) | | cLogP | pKa | logS |
								IC$_{50}$ (nM)	CC$_{50}$ (nM)	IC$_{50}$ (nM)	CC$_{50}$ (nM)			
1 (Niclosamide)	OH	Cl	H	Cl	H	NO2	H	564	1050	16770	17060	4.17	7.98	-5.00
2	OH	H	H	Cl	H	NO2	H	296	5590	4915	8403	3.47	7.52	-4.32
3	OH	CH3	H	Cl	H	NO2	H	1254	6277	2595	3056	3.97	7.52	-4.66
4	OH	OCH3	H	Cl	H	NO2	H	1769	10110	10880	8941	3.39	7.65	-4.39
5	OH	Cl	H	Cl	H	H	H	890	8435	>20000	>20000	4.17	8.01	-4.54
6	OH	Cl	H	Cl	H	CH3	H	760	4591	>20000	6048	4.66	8.01	-4.88
7	OH	Cl	H	Cl	H	COOCH3	H	334	8142	Inverted	>20000	4.15	7.98	-4.94
8	OH	Cl	H	Cl	H	COOH	H	>20000	>20000	Inverted	>20000	3.67	4.03	-4.54
9	H	Cl	H	Cl	H	NO2	H	>20000	>20000	>20000	>20000	4.63	14	-5.27
10	OH	Cl	H	Cl	H	OCH3	H	4248	>20000	3097	1921	4.05	8.02	-4.59
11	OH	t-Bu	H	Cl	H	NO2	H	423	3407	16880	7529	5.50	7.51	-5.68
12	OH	OCH3	H	Cl	H	NO2	H	1498	10290	Inverted	>20000	3.39	7.65	-4.39
13	OH	t-Bu	H	Cl	H	COOCH3	H	>20000	14570	Inverted	>20000	5.49	7.50	-5.62
14	OH	t-Bu	H	Cl	H	COOH	H	>20000	14250	Inverted	>20000	5.00	4.93	-5.22
15	OH	t-Bu	H	F	H	COOH	H	>20000	>20000	Inverted	>20000	4.41	4.73	-4.69
16	OH	t-Bu	H	CH3	H	COOH	H	>20000	>20000	Inverted	>20000	4.49	4.95	-4.74
17	OH	t-Bu	H	Cl	H	H	COOH	>20000	>20000	Inverted	>20000	5.00	4.93	-5.22
18	OH	t-Bu	t-Bu	Cl	H	CH3SO2N	H	>20000	>20000	Inverted	>20000	5.98	7.02	-6.39
19	OH	t-Bu	t-Bu	Cl	H	NH2	H	>20000	17590	Inverted	>20000	6.58	7.57	-6.14
20	OH	t-Bu	t-Bu	H	CH3	COOH	H	>20000	>20000	>20000	>20000	6.52	4.96	-6.10
21	OH	Cy	H	Cl	H	COOCH3	H	9910	>20000	>20000	>20000	5.96	7.50	-6.66
22	OH	Cy	H	Cl	H	COOH	H	4511	>20000	7700	>20000	5.48	4.93	-6.26
23	OH	CF3	H	CH3	H	COOCH3	H	>20000	>20000	>20000	>20000	4.97	7.49	-5.15
24	OH	Cl	H	Cl	H	CN	H	348	4720	1070	880	4.27	8.00	-5.03
25	OH	H	t-Bu	Cl	H	COOCH3	H	>20000	>20000	Inverted	>20000	5.22	7.52	-5.52
26	OH	t-Bu	t-Bu	Cl	H	COOH	H	>20000	>20000	>20000	>20000	6.77	4.93	-6.48
27	OH	t-Bu	t-Bu	Cl	H	COOCH3	H	16500	>20000	>20000	>20000	6.52	4.96	-6.10
28	OH	OCH3	H	CH3	H	COOH	H	>20000	>20000	Inverted	>20000	2.38	4.95	-3.46
29	OH	H	t-Bu	Cl	H	COOCH3	H	>20000	>20000	Inverted	>20000	5.22	7.52	-5.52
30	OH	H	t-Bu	Cl	H	COOH	H	>20000	>20000	>20000	>20000	4.74	4.93	-5.12
31	OH	H	t-Bu	Cl	H	NO2	H	>20000	7214	>20000	1110	5.24	7.52	-5.58
32	OH	H	t-Bu	Cl	H	NH2	H	>20000	>20000	>20000	>20000	4.55	7.58	-4.78
33	OH	H	t-Bu	Cl	H	NSO2CH3	H	2879	>20000	Inverted	>20000	3.95	7.03	-5.03
34	OH	Cl	H	Cl	H	Br	H	760	5473	4067	3242	4.97	8.01	-5.36

664