Title: Spatial replication can best advance our understanding of population responses to climate

Abstract

Understanding the responses of plant populations dynamics to climatic variability is frustrated by

the need for long-term datasets. Here, we advocate for new studies that estimate the effects of

climate by sampling replicate populations in locations with similar climate. We first use data analysis

on spatial locations in the conterminous USA to assess how far apart spatial replicates should be

from each other to minimize temporal correlations in climate. We find that on average spatial

locations separated by 316 Km (SD = 149Km) have moderate (0.5) correlations in annual

precipitation. Second, we use simulations to demonstrate that spatial replication can lead to

substantial gains in the range of climates sampled during a given set of years so long as the climate

correlations between the populations are at low to moderate levels. Third, we use simulations to

quantify how many spatial replicates would be necessary to achieve the same statistical power of a

single-population, long-term data set under different strengths and directions of spatial correlations

in climate between spatial replicates. Our results indicate that spatial replication is an untapped

opportunity to study the effects of climate on demography and to rapidly fill important knowledge

gaps in the field of population ecology.

INTRODUCTION

Understanding the responses of biodiversity to climate drivers is necessary to mitigate and adapt to

climate change (Urban et al. 2016). In recent years, the field of ecological forecasting has

experienced substantial theoretical and applied growth (Lewis et al. 2022). These forecasts on

population dynamics are necessary to assess species extinction risk (Mace et al. 2008) and predict

range shifts (Schurr et al. 2012). However, forecasting the effects of climate on populations remains

elusive. This plausibly occurs because it takes 20-25 years of data to sufficiently describe the

relationship between climate and demography (Teller et al. 2016, Tenhumberg et al. 2018). This

large replication is necessary to sample a sufficiently wide range of climatic conditions and to

increase statistical power.

We have important knowledge gaps in population ecology that cannot wait 20-25 years to

be filled. Our recent synthesis showed that knowledge on climate-demography relationships for

plants is particularly poor for the species-rich tropics, and for species with extreme generation times

(Compagnoni et al. 2021b). We need immediate research targeting these locations and plant life

histories. As most plant ecologists are at young career stages, we need to engage Doctoral

Researchers and Postdoctoral Associates in this field of research. However, this will not happen in

case projects last decades.

We propose that new studies should prioritize spatial over temporal replication to assess the

demographic responses of a species to climate. Having both spatial and temporal data allows

collecting a high sample size in a relatively short period of time. Spatial sampling increases our

statistical power by increasing the range of climates sampled per unit time, and allowing us to "see

through the noise" caused by non-climatic factors. The range of climates sampled increases as the

distance between populations decreases the correlation of yearly climatic anomalies among them.

We are not advocating a "space-for-time substitution", but to estimate the short-term

effects of climate by prioritizing spatial versus temporal replication. Space-for-time substitutions

infer the long-term effects of climate on plant populations (Blois et al. 2013). These long-term

effects are typically larger than short-term climate effects, possibly because they include the effects

of indirect effects (e.g. on biophysical conditions, Elmendorf et al., 2012). Instead, short-term

climatic effects are directly relevant to inform the management and conservation of populations

during the upcoming century of rapid climate change (e.g. Compagnoni, Pardini, et al., 2021; Hunter

et al., 2010). Here, we therefore advocate to replicate sampling across populations that occur in

similar climates, and to use this spatial data as *replicates* of the same temporal process. This

recommendation relies on the assumption that in similar environments, plant populations should

respond similarly to climate anomalies. Such assumption has so far motivated much research on the

demography across species ranges (Kleinhesselink & Adler, 2018; Morley et al., 2017). However, to

our knowledge, no study on climate-demography relationships has yet prioritized spatial replication

across sites with similar climatic conditions.

In this manuscript, we show the opportunities of spatial replication in climate-demography

studies through data analysis and simulation. First, we assess how far apart populations must be

from each other to attenuate temporal correlations in climate using gridded climatic data from the

conterminous USA. Second, we consider how sampling design can maximize the range of climates

captured during a study. To do so, we estimate how the range of climates sampled depends on the

climate correlations among the populations and the study duration (between five and 30 years).

Third, we use simulation to quantify the statistical power of a climate-demography relationship

across sampling designs that vary in the number of spatial and temporal replicates, and in the

strength of spatial correlations in climate between populations. In this simulation, we also address

cases in which populations respond differently to climate. Based on these results, we make

recommendations for new demographic data collection efforts.

METHODS

Spatial correlation in climate

To understand how spatial correlation in climate depends on the distance between sites, we

estimate the spatial correlation of annual climate in the conterminous United States of America

(USA), a large and climatically heterogeneous section of terrestrial environments. We downloaded

monthly temperature and precipitation data for the conterminous USA from the CHELSA database

(Karger et al. 2017, 2018). CHELSA data is accurate on varied topographic terrain, and gridded

climatic data generally correlates strongly with weather station data (Behnke et al. 2016). We

downloaded data between 1979 and 2013 following a regular grid of 0.5 degrees, for a total of 3261

locations. For each location, we calculated annual temperature means, annual precipitation sums,

and computed their standardized yearly anomalies (z-scores, henceforth "anomalies").

For each location, we calculated the average distance at which the correlation in yearly climate

anomalies is expected to decrease to 0.5 or lower. To do so, we calculated the correlation between

the 35 annual anomalies at a location, and the same anomalies observed at the remaining 3260

locations. Then, we fit a local polynomial regression using observed correlations as a response

variable, and distance from the original location as predictor. The predictions of this local regression

identified the distance at which the correlation in climate anomalies is expected to be 0.5 or lower.

Finally, we produced heatmaps showing, for each location, the average predicted distance at which

the correlation in climatic anomalies is 0.5 or lower.

To show a sample of the raw data we used for this analysis, we show the data referred to

five reference locations. We picked these locations subjectively, locating them in the Southwestern

coast, Northwestern coast, Northeastern Coast, Southeastern coast, and in the center of the USA.

For each location, we first plotted the correlation between climatic anomalies versus the distance

from the reference location, including the average prediction of the polynomial regression

summarizing this relationship. Then, we produced heatmaps showing how the climatic anomalies

observed across the USA correlated with the reference location.

Range of climates sampled with different sampling designs

We performed simulations to understand how the range of climate anomalies sampled changes

depending on temporal replication, spatial replication, and the spatial correlation of climatic

anomalies. We estimated the range of climatic anomalies sampled at two sites using a multivariate

normal (MVN) distribution:

$$X \sim MVN\left(\mathbf{0}, \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}\right), \qquad (1a)$$

$$\boldsymbol{Range}_x = \frac{Max(X) - Min(X)}{2}, \qquad (1b)$$

where **X** is an *n* by 2 matrix of climate anomalies at two separate sites, *n* is the temporal replication

of the study, *ρ* represents the correlation between the climatic anomalies of the two sites. The

parameters of the MVN distribution show **0** for the means, and diagonal elements of 1 for the

variance to simulate a series of anomalies. When simulating the climate at a single site, we needed a

single series of climate anomalies, so we substituted Eq. (1a) with $X \sim Normal(0,1)$. To obtain

Range_X, the expected range of **X** values (Eq. 1b), we simulated Eq. 1a 1000 times across *n* values

ranging from two to 30 in increments of one, and across a series of *ρ* values of 0, 0.5, 0.95, and one.

Finally, we calculated the mean of **Range_X** across these 1000 replicate simulations.

Statistical power for climate-demography relationship with different sampling designs

We used simulations to quantify the statistical power of the relationship between climate and

population growth rate for different spatio-temporal sampling designs and different spatial

correlations in climate. Starting from a case with two separate sites,

$$X \sim MVN\left(\mathbf{0}, \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}\right), \qquad (2a)$$

$$\hat{Y}_{is} = \beta_0 + \beta_1 X_{is}, \qquad (2b)$$

$$Y_{is} \sim Normal(\hat{Y}_{is}, \varepsilon) \qquad (2c)$$

where X is an n by 2 matrix of normally distributed climatic anomalies, n refers to temporal

replicates, $\hat{Y}_{is}$ is the average prediction of the model referred to site s and year i, β_0 the intercept of

the linear model, β_1 is the slope, and Y_{is} represents the observed log population growth rate. Y_i is a

log population growth rate because we have synthetic estimates of climatic effects on this variable

(Compagnoni et al. 2021b), because it is the central focus of demographic theory (Sibly and Hone

2002), and because it is normally distributed, facilitating simulations and their interpretation. For

this and subsequent simulations we used a β_1 value of 0.05, and a ε value of 0.15 to reflect empirical

estimates from 162 plant populations (Compagnoni et al. 2021b). We simulated the process

described in Eq. 2 1000 times using study durations, n, of three, four, and five years, correlations ρ of

0.95 and 0.5, 0, and -0.5 and a number of spatial replicates of two, 10, 20, 30, 40, and 50. We divided

spatial replicates in two subsets, and assigned each subset to one of the two series of climatic

anomalies simulated by Eq. 2. For example, when spatial replicates were 50, we subdivided these

replicates in two groups of 25 replicates, each group experiencing identical climate. We used low

values for n to reflect that the median length of demographic studies of plants is four years

(Salguero-Gómez et al. 2015). This sampling effort likely reflects the length of many PhD programs.

131   As an alternative, we also simulated assuming a single censused population, using study

132 durations of 20 and 30 years. This simulation was meant to represent the handful of single site, long-

133 term studies found in the literature (e.g. Chu et al. 2016). In these simulations, we substituted Eq. 2a

134 with $X \sim Normal(0,1)$. We calculated power as the proportion of the 1000 simulations for which β_1

135 had a p-value below 0.05. This power analysis based on simulation is a straightforward way to

136 quantify how the uncertainty of model estimates is influenced by the sampling design.

137   The power estimate described above assumed that β_1, in Eq. 2b, were the same for each

138 population. Therefore, we also addressed the sensitivity of our power estimates to spatial variation

139 in β_1. We ran all of our analyses in R version 4.0.2 (R Core Team 2020). We fit polynomial regressions

140 using the R function loess, and we produced spatial plots using package ggmap (Kahle and Wickham

141 2013). We ran the analyses on the spatial correlation in climate using a high-performance computing

142 cluster.

RESULTS

Spatial correlation in climate

In the conterminous USA annual temperature anomalies are strongly correlated even at relatively

large (e.g. 500Km, Fig. S1-5A) distances. The correlation between precipitation anomalies is less

strong, and it decays more rapidly with distance (Fig. S1-5C). In the conterminous USA, our loess

models predict that the distance to reach a correlation of 0.5 is on average 316 Km (SD = 149 Km) for

precipitation and 1460Km (SD = 428Km) for temperature. However, there is substantial

heterogeneity in the decay of correlation for each location (Fig. 1). Considering the 5th and 95th

percentiles of the distances at which the correlation of climate anomalies is predicted to be 0.5,

these range from 52 to 568 Km for precipitation, and from 868 to 2196Km for temperature. Sites

with a predicted correlation of zero are on average always at least 1000Km apart: only 5% of the

predicted distances at which the correlation is zero have a distance equal to or less than 1000 and

1920 Km for precipitation and temperature, respectively.

Range of climates sampled with different sampling designs

Spatial replication can lead to substantial gains in the range of climates sampled during a given set of

years so long as the climate correlations between populations are at low to intermediate levels. To

reach a range of ±2 standard deviation at a single site, one would on average need 27 years of data.

This number of years decreases to 20 when using two sites whose climate has correlation 0.9, and 15

when two sites have correlations 0.5 (Fig. S6).

Statistical power for climate-demography relationship with different sampling designs

Our power analysis indicates that spatial replication greatly increases the power to detect a

relationship between climate and population growth rate (Fig. 2). The statistical power of very long

time series for a single site is comparable to that of datasets with high spatial replication. One site

sampled for 20 and 25 times provides a statistical power of about 30 and 41%, respectively. These

two statistical powers are reached with just three temporal samples distributed across, respectively,

10 and 20 spatial replicates experiencing medium to highly correlated climate (0.5, 0.95). With four

temporal samples and 10 spatial replicates, a statistical power of 40% is reached with climate

correlation 0.95; with five temporal samples and 10 spatial replicates, statistical power always

exceeds 40%. Lowering climatic correlation between populations from 0.95 to 0.5 slightly increases

statistical power: depending on the number of temporal samples and spatial replicates, increases in

statistical power vary from 4 to 8% (Fig. 2).

Variation in the effect of climate on demography (represented by β_1 values) did not

noticeably affect statistical power (Fig. S7). Statistical power remained unchanged, reflecting that is

was influenced by the average β_1, which was still 0.05, rather than its variation.

DISCUSSION

Until recently, understanding the response of plant species to climatic variation has relied on either

long-term monitoring efforts, which are rare (Salguero-Gómez et al. 2015), or on responses of plant

populations to spatial climate gradients ("space-for-time substitutions", Blois et al., 2013) which, are

affected by several confounding factors. Specifically, spatial climatic gradients do not measure the

direct effect of climate on populations, because they encompass substantial changes in community

composition (Whittaker 1970), and biophysical factors (Shaver et al. 2000). Fortunately, our power

analysis shows that we can propel our understanding of species responses to climate using spatial

replicates that come from sites with similar long-term average climates.

We argue that using numerous spatial replicates in studies of demographic responses to

climate is an opportunity that should be exploited urgently by conservation scientists. Our power

analysis suggests that large spatial replication allows obtaining estimates of climate effects in as little

as three years. This short temporal horizon would allow early career researchers or distributed

networks to study the effects of climatic variability on populations. Such abundance of investigators

would facilitate ameliorating the taxonomic, geographic, and life-history biases present in current

data (e.g. Compagnoni et al., 2021). The result would encourage the development and improve the

quality of climate change vulnerability assessments. The IUCN recommends evaluating vulnerability

by focusing on time series of population data (Mace et al. 2008). These data can be readily used to

estimate the effects of climate on population dynamics and, as a result, climate vulnerability (Hunter

et al. 2010, Jenouvrier et al. 2022). However, given that long-term population data are rare in plants

(Salguero-Gómez et al. 2015), large spatial replication provides an opportunity to perform such

population-based climate vulnerability assessments.

We suggest new studies focusing on sites with similar climate located at the leading or rear

edge of species ranges, because these locations are ecologically more important, and because they

should provide higher statistical power. Leading or rear range edges are the locations where

populations are most sensitive to climate (Morley et al. 2017, Amburgey et al. 2018, Kleinhesselink

and Adler 2018); accordingly, range edges also show the highest variability in population growth

rates (Sexton et al. 2009, Csergő et al. 2017, Guyennon et al. 2023). From an ecological perspective,

such high climate sensitivity provides an opportunity to understand range limit formation and to

forecast range shifts (Parmesan 2006, Ehrlén and Morris 2015). From a statistical point of view, a

higher sensitivity to climate increases statistical power.

We believe that the assumptions of our power analysis are likely to hold, as they reflect

current ecological understanding. First, the assumption that populations experiencing the same

average climate respond similarly to climate variation is well supported in the literature (Morley et

al. 2017, Amburgey et al. 2018, Kleinhesselink and Adler 2018). Moreover, our power analysis is

robust to relaxing this assumption (Fig. S6). Second, the assumption that residual variance

(represented by ε in Eq. 2c) is constant across replicates also reflects current knowledge. Specifically,

review (Sexton et al. 2009) and synthesis studies (Csergő et al. 2017, Guyennon et al. 2023) indicate

that the variability of population growth rates should become larger only at range edges. Third, so

long as the spatial replicates are sufficiently spaced (e.g., separated by dispersal), they should meet

the assumption of statistical independence. In principle, statistical independence might not hold in

response to unobserved large-scale drivers such as herbivores with large foraging ranges, or

epidemic outbreaks. For this reason, we recommend testing for the presence of spatial

autocorrelation in the residuals (see below).

To improve ecological inference, we suggest that researchers exploit the covariates that

might mediate demographic responses to climate, such as soil conditions (Nicolè et al. 2011) or

competition (Alexander et al. 2015). In single-species studies, these mediators cannot be taken into

consideration, putting the external validity of the results into question. Spatial replication allows to

quantify and test the importance of mediating factors. However, because the effect of these

mediators is assessed by estimating an interaction, this requires a concomitant increase in sample

size (Ch. 16: Gelman et al., 2020).

To detect spatial autocorrelations that might occur due to unexpected, large-scale processes

such as epidemics, we suggest that spatial replicates should be sampled randomly rather than on a

regular grid (Fortin et al. 1989). Detecting spatial autocorrelation is important, because it would

indicate that the spatial replicates are not fully independent. A large spacing between groups of sites

(e.g. 100 Km) would contribute to minimize such potential autocorrelation, while minimizing the

climatic correlation among sites. This spacing might be possible for a distributed network of

researchers, but it may be prohibitively expensive or logistically difficult for a single doctoral

researcher. We suggest that, given the relatively small effect of climate correlation on statistical

power, smaller spacing between populations might be a viable option in many real world situations.

Our simulations and analyses ignored the importance of microclimate which, however,

might still be leveraged in spatially replicated studies. Recent studies highlight the importance of

microtopography (Scherrer and Körner 2010), exposure (Ackerly et al. 2020), and vegetation type

(Sanczuk et al. 2023) on plant responses to climate. It is unlikely that nearby sites experiencing

different microclimates will also observe substantially different annual climatic anomalies. However,

microclimates experience different absolute climatic conditions (e.g. minimum and maximum

temperatures (Bennie et al. 2008). These conditions can be leveraged in studies that use

physiological climatic predictors, such as growing degree days. While such physiological predictors

require daily weather data, these can be measured via data loggers, or derived from gridded climatic

data (e.g. (Maclean et al. 2019, Kearney and Porter 2020).The spatially replicated sampling we

propose here is a practical solution to estimate climate-demography relationships and rapidly fill

important knowledge gaps in the field of population ecology. However, monitoring 20 or more

populations simultaneously is a large task for a single researcher. Such spatial replication might

become more feasible through collaborative research networks (e.g. Villellas et al., 2021), or through

unmanned aerial vehicles (UAV). UAV might make spatially replicated plant demographic data

increasingly easy to obtain. For example, in both grassland and forest vegetation, UAV can identify

individuals of a focal species (Schmidt et al. 2017, Bogdan et al. 2021, Allen et al. 2023), and quantify

vegetation structure (Zhang et al. 2021, Coverdale and Davies 2023). We believe that the sampling

choices we advocate in this article will contribute to the maturation of population ecology and its

links to conservation science, functional ecology, and macro-ecology.

Acknowledgements. We performed the analyses on the spatial correlation of climate using the High-

Performance Computing Cluster EVE (a joint effort of both the Helmholtz Centre for Environmental

Research - UFZ (http://www.ufz.de/) and the German Centre for Integrative Biodiversity Research

(iDiv) Halle-Jena-Leipzig (http://www.idiv-biodiversity.de/).

Data availability statement. The data and reproducible code for the analyses connected to this

study are available at https://doi.org/10.5281/zenodo.8271041. We downloaded climatic data via

CHELSA, dataset version 1.2, which is available at http://dx.doi.org/doi:10.5061/dryad.kd1d4.

Conflict of interest statement. We declare no conflicts of interest.

Ethics statement. We identified no ethical issues associated with this project.

Author contributions

Aldo Compagnoni: Conceptualization (lead), data curation (lead), formal analysis (lead),

investigation (lead), methodology (lead), resources (lead), software (lead), validation (lead),

visualization (lead), writing – original draft (lead), writing – review & editing (equal). **Sanne Evers**:

Conceptualization, data curation, formal analysis, investigation, methodology, resources, software,

visualization, writing – review & editing (equal). **Tiffany M. Knight**: Conceptualization, funding

acquisition (lead), methodology, project administration (lead), supervision (lead), writing – review &

editing (equal).

REFERENCES

Ackerly, D. D. et al. 2020. Topoclimates, refugia, and biotic responses to climate change. - Frontiers in Ecology and the Environment 18: 288–297.

Alexander, J. M. et al. 2015. Novel competitors shape species' responses to climate change. - Nature 525: 515–518.

Allen, M. J. et al. 2023. Tree species classification from complex laser scanning data in Mediterranean forests using deep learning. - Methods in Ecology and Evolution 14: 1657–1667.

Amburgey, S. M. et al. 2018. Range position and climate sensitivity: The structure of among-population demographic responses to climatic variation. - Global Change Biology 24: 439–454.

Behnke, R. et al. 2016. Evaluation of downscaled, gridded climate data for the conterminous United States. - Ecological Applications 26: 1338–1351.

Bennie, J. et al. 2008. Slope, aspect and climate: Spatially explicit and implicit models of topographic microclimate in chalk grassland. - Ecological Modelling 216: 47–59.

Blois, J. L. et al. 2013. Space can substitute for time in predicting climate-change effects on biodiversity. - PNAS 110: 9374–9379.

Bogdan, A. et al. 2021. Demographic analysis of an Israeli Carpobrotus population. - PLOS ONE 16: e0250879.

Chu, C. et al. 2016. Direct effects dominate responses to climate perturbations in grassland plant communities. - Nat Commun 7: 11766.

Compagnoni, A. et al. 2021a. Increasing temperature threatens an already endangered coastal dune plant. - Ecosphere 12: e03454.

Compagnoni, A. et al. 2021b. Herbaceous perennial plants with short generation time have stronger responses to climate anomalies than those with longer generation time. - Nature Communications 12: 1824.

Coverdale, T. C. and Davies, A. B. 2023. Unravelling the relationship between plant diversity and vegetation structural complexity: A review and theoretical framework. - Journal of Ecology 111: 1378–1395.

Csergő, A. M. et al. 2017. Less favourable climates constrain demographic strategies in plants. - Ecology Letters 20: 969–980.

Ehrlén, J. and Morris, W. F. 2015. Predicting changes in the distribution and abundance of species under environmental change. - Ecology Letters 18: 303–314.

Elmendorf, S. C. et al. 2012. Plot-scale evidence of tundra vegetation change and links to recent summer warming. - Nature Climate Change 2: 453–457.

Fortin, M.-J. et al. 1989. Spatial Autocorrelation and Sampling Design in Plant Ecology. - Vegetatio 83:
209–222.

Gelman, A. et al. 2020. Regression and Other Stories. - Cambridge University Press.

Guyennon, A. et al. 2023. Beyond mean fitness: Demographic stochasticity and resilience matter at
tree species climatic edges. - Global Ecology and Biogeography 32: 573–585.

Hunter, C. M. et al. 2010. Climate change threatens polar bear populations: a stochastic
demographic analysis. - Ecology 91: 2883–2897.

Jenouvrier, S. et al. 2022. Detecting climate signals in populations across life histories. - Global
Change Biology 28: 2236–2258.

Kahle, D. and Wickham, H. 2013. ggmap: Spatial Visualization with ggplot2. - The R Journal 5: 144.

Karger, D. N. et al. 2017. Climatologies at high resolution for the earth's land surface areas. - Sci Data
4: 1–20.

Karger, D. N. et al. 2018. Data from: Climatologies at high resolution for the earth's land surface
areas.: 7266827510 bytes.

Kearney, M. R. and Porter, W. P. 2020. NicheMapR – an R package for biophysical modelling: the
ectotherm and Dynamic Energy Budget models. - Ecography 43: 85–96.

Kleinhesselink, A. R. and Adler, P. B. 2018. The response of big sagebrush (Artemisia tridentata) to
interannual climate variation changes across its range. - Ecology 99: 1139–1149.

Lewis, A. S. L. et al. 2022. The power of forecasts to advance ecological theory. - Methods in Ecology
and Evolution in press.

Mace, G. M. et al. 2008. Quantification of Extinction Risk: IUCN's System for Classifying Threatened
Species. - Conservation Biology 22: 1424–1442.

Maclean, I. M. D. et al. 2019. Microclima: An r package for modelling meso- and microclimate. -
Methods in Ecology and Evolution 10: 280–290.

Morley, J. W. et al. 2017. Marine assemblages respond rapidly to winter climate variability. - Global
Change Biology 23: 2590–2601.

Nicolè, F. et al. 2011. Interdependent effects of habitat quality and climate on population growth of
an endangered plant. - Journal of Ecology 99: 1211–1218.

Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. - Annual Review
of Ecology, Evolution, and Systematics 37: 637–669.

Salguero-Gómez, R. et al. 2015. The COMPADRE Plant Matrix Database: an open online repository for
plant demography (M Rees, Ed.). - Journal of Ecology 103: 202–218.

Sanczuk, P. et al. 2023. Microclimate and forest density drive plant population dynamics under
climate change. - Nat. Clim. Chang. 13: 840–847.

Scherrer, D. and Körner, C. 2010. Infra-red thermometry of alpine landscapes challenges climatic
warming projections. - Global Change Biology 16: 2602–2613.

Schmidt, J. et al. 2017. Adapting a Natura 2000 field guideline for a remote sensing-based
assessment of heathland conservation status. - International Journal of Applied Earth
Observation and Geoinformation 60: 61–71.

Schurr, F. M. et al. 2012. How to understand species' niches and range dynamics: a demographic
research agenda for biogeography. - Journal of Biogeography 39: 2146–2162.

Sexton, J. P. et al. 2009. Evolution and Ecology of Species Range Limits. - Annual Review of Ecology,
Evolution, and Systematics 40: 415–436.

Shaver, G. R. et al. 2000. Global Warming and Terrestrial Ecosystems: A Conceptual Framework for
Analysis: Ecosystem responses to global warming will be complex and varied. Ecosystem
warming experiments hold great potential for providing insights on ways terrestrial
ecosystems will respond to upcoming decades of climate change. Documentation of initial
conditions provides the context for understanding and predicting ecosystem responses. -
BioScience 50: 871–882.

Sheth, S. N. and Angert, A. L. 2018. Demographic compensation does not rescue populations at a
trailing range edge. - PNAS 115: 2413–2418.

Sibly, R. M. and Hone, J. 2002. Population growth rate and its determinants: an overview (RM Sibly, J
Hone, and TH Clutton–Brock, Eds.). - Phil. Trans. R. Soc. Lond. B 357: 1153–1170.

Teller, B. J. et al. 2016. Linking demography with drivers: climate and competition (J Metcalf, Ed.). -
Methods Ecol Evol 7: 171–183.

Tenhumberg, B. et al. 2018. Time-lagged effects of weather on plant demography: drought and
Astragalus scaphoides. - Ecology 99: 915–925.

Urban, M. C. et al. 2016. Improving the forecast for biodiversity under climate change. - Science 353:
aad8466–aad8466.

Villellas, J. et al. 2021. Phenotypic plasticity masks range-wide genetic differentiation for vegetative
but not reproductive traits in a short-lived plant (C Violle, Ed.). - Ecology Letters 24: 2378–
2393.

Whittaker, R. H. 1970. Communities and Ecosystems. - Macmillan, London: Collier Macmillan.

Zhang, X. et al. 2021. Using UAV LiDAR to Extract Vegetation Parameters of Inner Mongolian
Grassland. - Remote Sensing 13: 656.

FIGURE LEGENDS

Figure 1. The correlation between temperature and precipitation anomalies decays slowly with

distance. The heatmaps show for each grid cell, the average distance at which the correlation in

temperature (A) and precipitation (C) anomalies decreases to 0.5 or lower. These average distances

are estimated using a local regression (loess) model.

Figure 2. Spatial replication provides a statistical power similar, or higher, to temporal replication.

Plot showing statistical power on the y-axis against spatial replication on the x-axis. Symbols show

temporal replications of three (circles), four (triangles), five (squares), 20 (cross), and 30 (crossed

square). The color of symbols refers to the correlation among spatial replicates. This correlation is

one for the simulations with a single spatial replicate. The dashed horizontal line denotes a statistical

power of 80%.

Figure 1.

397 **Figure 2**.