Main Manuscript for

Cooperation genes are more pleiotropic than private genes in the bacterium *Pseudomonas aeruginosa*

Trey J. Scott

Department of Biology, Washington University in St. Louis

Corresponding author: Trey J. Scott

Email: tjscott@wustl.edu

Orcid ID: 0000-0001-6609-9638

Author Contributions: TS designed the study, analyzed the data, and wrote the manuscript

Competing Interest Statement: I have no competing interests to disclose.

Classification: Biological Sciences, Evolution

Keywords: Pleiotropy, cooperation, kin selection, Pseudomonas aeruginosa

This file includes:

> Main Text
> Figures 1 to 2

Abstract

Pleiotropy may affect the evolution of cooperation by limiting cheater mutants if such

mutants also lose other important traits. Because pleiotropy limits cheaters, selection may favor

cooperation genes that are more pleiotropic. However, the same should not be true for private

genes with functions unrelated to cooperation. Pleiotropy in cooperative genes has mostly been

studied with single genes and has not been measured on a wide scale or compared to a suitable set

of control genes with private functions. I remedy this gap by comparing genomic measures of

pleiotropy in previously identified cooperative and private gene sets in *Pseudomonas aeruginosa*.

I found that cooperative genes in *P. aeruginosa* tended to be more pleiotropic than private genes

according to the number of protein-protein interactions, the number of gene ontology terms, and

gene expression specificity. These results show that pleiotropy may be a general way to limit

cheating and that cooperation may shape pleiotropy in the genome.

Main Text

Introduction

Many genes are pleiotropic, where a pleiotropic gene is defined as one that affects

multiple traits. Pleiotropy constrains evolution because mutations with beneficial effects on one

trait can have deleterious effects on other traits. Several measures of pleiotropy have been used

and shown to be associated with evolutionary constraint. Three examples are the number of

protein interactions (1), the number of functional annotations (2), and how widely genes are

expressed across tissues (3).

Pleiotropy is thought to limit cheaters and stabilize cooperation (4-7; see 8 for an

alternate way pleiotropy can affect cooperation and 9 for a dissenting view). Explaining how

cheater evolution can be limited is a central question in the study of cooperation (10). Cheaters

benefit from cooperation without paying the costs and are expected to outcompete cooperators.

This cheater advantage can lead to the breakdown of cooperation unless cheaters are constrained

(10).

Pleiotropy can limit cheaters when mutations that cause cheater phenotypes come with

harmful effects on other traits. One example of this comes from the social amoeba *Dictyostelium*

discoideum. D. discoideum has a cooperative stage where 20% of cells sacrifice themselves to

become stalk. The remaining cells become spores that are held up for dispersal by the stalk (10).

This act of cooperation can be exploited by cheaters that contribute less to the stalk and increase

their abundance in spores (10). Amoebae with *dimA* mutations are potential cheaters because they

ignore the signal to become stalk (4). This should increase *dimA* mutant representation as spores.

Instead, *dimA* mutants are excluded from spores when mixed with wildtype cells as a pleiotropic

effect (4). This trade-off between becoming a stalk cell and entering spores could limit the

cheating ability of *dimA* mutants.

Because pleiotropy limits cheaters, it should be higher in cooperative genes relative to

private (non-cooperative) genes (5). It is unknown whether cooperative genes are generally more

pleiotropic than private genes. Prior studies of pleiotropy in cooperative genes mostly focused on

individual genes (4) or on the observation that genes are co-regulated (7). Such studies do not

explicitly quantify pleiotropy or compare between cooperative and private genes. Private genes

are an important control because they should reflect the background pleiotropy that is not affected

by social selection.

47    Here, I take a genomic approach to compare cooperative and private genes as categorized

48 by Belcher et al. (11) in four *P. aeruginosa* gene sets. These gene sets have been constructed so

49 that cooperative and private genes in a set are expressed under similar conditions. Because of this,

50 cooperative and private genes should be similarly exposed to selection, all else being equal within

51 a set (11). These sets of cooperative and private genes are therefore ideal for testing whether

52 cooperative genes are more pleiotropic.

54 **Results**

56    To test whether pleiotropy is higher in cooperation genes compared to private genes, I

57 first used 315 quorum sensing genes (12) that were previously categorized as cooperative (N =

58 41) or private (N = 274) based on gene annotations and experimental data (11). As measures of

59 pleiotropy, I used the number of protein interactions contained in the STRING database (13), the

60 number of biological process Gene Ontology (GO) terms (14), and gene expression pleiotropy for

61 each gene with available data (Figure 1). These pleiotropy measures were not highly correlated

62 and thus represent independent measures of pleiotropy (see appendix).

63    I found that cooperative genes in the quorum sensing pathway were more pleiotropic

64 across all three measures of pleiotropy (Fig. 2A). Cooperative genes had about 65 more STRING

65 interactions on average than private genes (generalized linear model (GLM); p = 0.0162).

66 Cooperative genes had 2 GO terms while private genes tended to have only 1 (GLM; p = 0.0090).

67 Gene expression pleiotropy was around 15% higher in cooperative genes than private genes (beta

68 regression; p = 0.0011).

69    To test whether these results apply beyond the quorum sensing pathway, I tested

70 additional sets of cooperative and private genes in *P. aeruginosa* that were included in Belcher et

al. (11). These gene sets consisted of antibiotic resistance genes (AMR) and pyochelin and

pyoverdine genes that are involved in binding iron. To increase statistical power, I analyzed all

three gene sets together (Fig. 2B; see appendix). Cooperative genes tended to have more STRING

protein interactions than private genes across these three gene sets (GLM; p = 0.0078). However,

cooperative and private genes were not different in terms of GO terms (GLM; p = 0.371) and

expression pleiotropy (beta regression; p = 0.102).

Discussion

Cooperation can break down because of the evolution of cheaters that benefit from

cooperation without helping (10). The advantage of a cheater mutation can be limited if it also

has pleiotropic effects on other important traits (4–7). This may result in high pleiotropy in

cooperative genes to limit cheaters. Prior studies have focused on single genes (4) and

coregulation (7) instead of measuring pleiotropy across many genes.

Using three independent measures of pleiotropy in *P. aeruginosa*, I found that pleiotropy

tended to be higher in cooperative genes than in private genes (Fig. 2). The majority of

comparisons in this study supports the hypothesis that pleiotropy limits cheaters in populations of

cooperators. In two comparisons, I found no difference between cooperative and private genes

(Fig. 2B). However, these comparisons had the smallest number of genes with measures of

pleiotropy and may have lacked power to detect a difference.

Another implication of increased pleiotropy in cooperation genes (Fig. 2) is a

strengthening of the conclusions of Belcher et al. (11), who found evidence of kin selection

because of relaxed selection in cooperation genes relative to private genes. Pleiotropy can also

affect patterns of selection, but it should be in the opposite direction of kin selection. Pleiotropy is

associated with signals of conservation (2) and stabilizing selection (15), which should decrease

genetic diversity and divergence. This means that the signal of kin selection in (11) may be an

underestimate since increased pleiotropy in cooperation genes should dampen the signal of

relaxed selection

Theoretical studies have disagreed about the direction of causality between cooperation

and pleiotropy and whether pleiotropy is able to stabilize cooperation if pleiotropy itself can

evolve (5, 6, 8, 9). While the data in this study cannot resolve these theoretical disagreements,

they show that pleiotropy and cooperation are linked in *P. aeruginosa*. Cooperation may thus

shape patterns of pleiotropy in the genomes of other cooperative organisms through its link with

high pleiotropy.

Materials and Methods

I used cooperation and private genes identified in Belcher et al (11) and generated

pleiotropy measures (see Figure 1, the appendix, and Datasets S1-S3). To test for differences

between cooperative and private genes, I used generalized linear models (GLMs) and beta

regression in R version 4.1.2. More detailed methods can be found in the appendix. Data and R

code for analysis are available at www.gitlab.com/treyjscott/kin_selection_pleiotropy.

Acknowledgments

I thank Laura Walker, Tyler Larsen, Joan Strassmann, and David Queller for comments on the

manuscript. This work was supported by NSF awards DEB 1753743 and IOS-1656756.

References

1.   H. B. Fraser, A. E. Hirsh, L. M. Steinmetz, C. Scharfe, M. W. Feldman, Evolutionary Rate in the Protein Interaction Network. *Science* **296**, 750–752 (2002).

2.   Y. Pritykin, D. Ghersi, M. Singh, Genome-Wide Detection and Analysis of Multifunctional Genes. *PLOS Comput. Biol.* **11**, e1004467 (2015).

3.   J. E. Mank, L. Hultin-Rosenberg, M. Zwahlen, H. Ellegren, Pleiotropic Constraint Hampers the Resolution of Sexual Antagonism in Vertebrate Gene Expression. *Am. Nat.* **171**, 35–43 (2008).

4.   K. R. Foster, G. Shaulsky, J. E. Strassmann, D. C. Queller, C. R. L. Thompson, Pleiotropy as a mechanism to stabilize cooperation. *Nature* **431**, 693–696 (2004).

5.   M. A. Bentley, C. A. Yates, J. Hein, G. M. Preston, K. R. Foster, Pleiotropic constraints promote the evolution of cooperation in cellular groups. *PLOS Biol.* **20**, e3001626 (2022).

6.   A. Frénoy, F. Taddei, D. Misevic, Genetic Architecture Promotes the Evolution and Maintenance of Cooperation. *PLoS Comput. Biol.* **9**, e1003339 (2013).

7.   A. A. Dandekar, S. Chugani, E. P. Greenberg, Bacterial Quorum Sensing and Metabolic Incentives to Cooperate. *Science* **338**, 264–266 (2012).

8.   D. C. Queller, Pleiotropy and synergistic cooperation. *PLOS Biol.* **17**, e3000320 (2019).

9.   M. dos Santos, M. Ghoul, S. A. West, Pleiotropy, cooperation, and the social evolution of genetic architecture. *PLOS Biol.* **16**, e2006671 (2018).

10.   J. E. Strassmann, D. C. Queller, Evolution of cooperation and control of cheating in a social microbe. *Proc. Natl. Acad. Sci.* **108**, 10855–10862 (2011).

11.   L. J. Belcher, A. E. Dewar, M. Ghoul, S. A. West, Kin selection for cooperation in natural bacterial populations. *Proc. Natl. Acad. Sci.* **119**, e2119070119 (2022).

12.   M. Schuster, C. P. Lostroh, T. Ogi, E. P. Greenberg, Identification, Timing, and Signal Specificity of *Pseudomonas aeruginosa* Quorum-Controlled Genes: a Transcriptome Analysis. *J. Bacteriol.* **185**, 2066–2079 (2003).

13.   D. Szklarczyk, *et al.*, The STRING database in 2021: customizable protein–protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Res.* **49**, D605–D612 (2021).

14.   Gene Ontology Consortium, The gene ontology resource: 20 years and still GOing strong. *Nucleic Acids Res.* **47**, D330–D338 (2019).

153   15.   K. McGuigan, J. M. Collet, S. L. Allen, S. F. Chenoweth, M. W. Blows,
154   Pleiotropic Mutations Are Subject to Strong Stabilizing Selection. *Genetics* **197**, 1051–
155   1062 (2014).

158
159   **Figures**
160

Fig. 1. Examples of pleiotropy measures used in this study. (A) Number of protein interactions

(STRING database). Arrows show predicted interactions that are summed to measure pleiotropy

(the numbers on the nodes). (B) Number of biological process Gene Ontology (GO) terms from

the *P. aeruginosa* genome database. Arrows show annotations for an example gene with three GO

terms. (C) Gene expression pleiotropy is calculated from gene expression data across multiple

conditions. Gene 1 is a gene that is expressed in every condition and is maximally pleiotropic.

Gene 2 is a gene that is specialized for a single condition and is minimally pleiotropic.

Fig. 2. Cooperative genes tend to be more pleiotropic than private genes in the (A) quorum

sensing and (B) other gene sets in *P. aeruginosa* (AMR = antibiotic resistance). Panels show the

number of protein-protein interactions in the STRING database (left), the number of gene

ontology (GO) terms (middle), and the gene expression pleiotropy (right). Number of genes with

measures of pleiotropy are shown on the x-axis. Colors are the same as in (10) for easy

comparison. *'s indicate $p \leq 0.05$.