Modelling vasopressin synthesis and storage dynamics during prolonged osmotic challenge and recovery based on activity dependent upregulation of mRNA transcription

Duncan J. MacGregor

Centre for Discovery Brain Sciences, University of Edinburgh, George Square

Edinburgh EH8 9XD, UK

E-mail: duncan.macgregor@ed.ac.uk

Abstract

Hypothalamic vasopressin neurons are neuroendocrine cells which form part of the homeostatic systems that maintain osmotic pressure. In response to synaptic inputs encoding osmotic pressure and changes in plasma volume, they generate spike triggered secretion of peptide hormone vasopressin from axonal terminals in the posterior pituitary. The thousands of neurons' secretory signals generate a summed plasma vasopressin signal acting at the kidneys to regulate water loss. Vasopressin is synthesised in cell bodies, packaged into vesicles, and transported to large stores in the pituitary terminals. Supported by activity-dependent upregulation of synthesis and transport, these stores can maintain a secretion response for several days of elevated osmotic pressure, tested by dehydration or salt loading. However, despite upregulated synthesis, stores gradually decline during sustained challenge, followed by a slow recovery. With no evidence of a store encoding feedback signal, previous modelling explained these synthesis dynamics based on activity-dependent upregulation of transcription and mRNA content. Here this model is adapted and integrated into our existing spiking and secretion model to generate a neuronal population model, able to simulate the secretion, store depletion, and replenishment, response to sustained osmotic challenge, matching the dynamics observed experimentally and making functional predictions for the cell body mechanisms.

Introduction

Magnocellular vasopressin neurons, of the supraoptic and paraventricular nuclei of the hypothalamus, in response to input signals that encode osmotic pressure and plasma volume, synthesise and secrete the antidiuretic hormone vasopressin. Vasopressin in its antidiuretic role is a core element of the homeostatic system that maintains osmotic pressure (water/salt balance), signalling the kidneys to regulate how much water is retained. Acting as a heterogeneous population, these neurons are able to maintain a constantly functioning physiological signal over lifelong periods of time. To sustain such a signal the system must be both very robust and efficient. It must also be able to respond rapidly to large changes in demand.

Vasopressin is synthesised in the neuronal cell bodies and packaged into large dense core vesicles that are transported down the axons to the posterior pituitary, where the vesicles are stored in axon swellings and terminals that form larger reserve and smaller releasable pools. We have previously modelled the spiking and secretion mechanisms of these neurons (MacGregor and Leng, 2012, 2013), including the dynamics of these pools. Here we build and integrate with our existing model, a quantitative model of the synthesis mechanisms, to better understand how the properties of these neurons relate to their function on long timescales.

In normal (basal) conditions mammals drink and ingest sodium (in the diet) intermittently, but constantly lose water through respiration and perspiration. Under homeostatic regulation, osmotic pressure fluctuates around a 'set point'; increases above this will be corrected by increased sodium excretion in urine, by increased thirst, and to compensate lack of availability or intermittent ingestion of water, by secreting vasopressin to concentrate the urine and minimise water loss. Falling below this set point occurs less commonly, when excess water has been consumed, or salt has been lost. Both tonic signalling and a response to perturbations must be maintained, and accordingly there is an almost continuous depletion of the pituitary vasopressin stores, which must be replenished by the synthesis, packaging, and transport of new vasopressin vesicles.

In conscious, normally hydrated rats, as in humans, the basal vasopressin plasma concentration is ~1 pg/ml (Robertson, Shelton and Athar, 1976; Verbalis, Baldwin and Robinson, 1986) and maximal antidiuresis is observed at a concentration

of about 10 pg/ml. At concentrations higher than this, vasopressin continues to have

an important role by its vasoconstrictor actions, which compensate for loss of fluid

volume in conditions of dehydration. The pituitary vasopressin stores are large,

between 1 and 2 µg (Leng and Ludwig, 2008), sufficient to maintain basal levels for

around a month (Jones and Pickering, 1972). These large stores buffer against rapid

increases in demand, but the rate of synthesis is also activity dependent, upregulating

production in response to sustained increase in demand, and downregulating

production in response to sustained low demand (Verbalis, Baldwin and Robinson,

1986). The system thus attempts to match supply and demand, minimising waste

whilst protecting its stores (MacGregor, Clayton and Leng, 2013).

Upregulation of synthesis is limited however, and under conditions of

sustained high demand, such as limited water access, the stores are depleted, falling to

less than 30% after five days (Jones and Pickering, 1969). The activity-dependent

synthesis rate is no longer able to match activity dependent secretion. When demand

and the stimulating osmotic signal has returned to normal, the stores are gradually

replenished, over a course of days (Young and Van Dyke, 1968). At this point the rate

of synthesis must be exceeding the activity-dependent secretion rate.

Building a system, the simplest way to do this would be to have some

feedback signal of store depletion. However we have no evidence for such a signal.

The pituitary stores at the neurons' secretory terminals are distant from the cell body

and highly distributed among thousands of release sites, making the measuring and

transmission of such a signal very difficult. The alternative is that some property of

the synthesis mechanisms forms a memory of the challenge, sufficient to maintain

higher synthesis rates beyond the direct stimulus. The best candidate for this is the

pool of vasopressin mRNA. The mRNA pool increases in size several fold in response

to prolonged challenge (Sherman, McKelvy and Watson, 1986). This mechanism was

extensively investigated using both experimental and modelling work by a group in

Pittsburgh in the late 1980s and early 1990s (Robinson *et al.*, 1989; Fitzsimmons *et

al.*, 1992; Robinson and Fitzsimmons, 1993). They tested several alternative models

(Fitzsimmons *et al.*, 1992) and showed that the best match to observed dynamics of

store depletion and replenishment uses an mRNA pool dependent rate of synthesis,

combined with activity dependent upregulation of transcription. During a prolonged

challenge the simulated mRNA pool increases in size, and following, the pool is

gradually depleted, sustaining increased synthesis sufficient to replenish the stores,

without requiring any feedback signal.

Their model (Fitzsimmons *et al.*, 1992) forms the basis for our work here.

Focussed on testing different models of the mRNA pool and its relation to synthesis

rate, they made the simplification that the synthesis rate always approaches a steady

state equal to the rate of secretion. What limits this response and causes store

depletion (and replenishment) in their model, is that this change in rate uses a long

time constant, dependent on the half-life of mRNA, estimated by them at two days.

They tested alternative models with activity dependent mRNA decay, and with longer

and shorter fixed decay rates, but the best fit to the experimentally observed dynamics

was with this model. The model was fitted to data from several sources (Young and

Van Dyke, 1968; Jones and Pickering, 1969; Zingg, Lefebvre and Almazan, 1986;

Roberts *et al.*, 1991) measuring the changing vasopressin content during a prolonged

osmotic challenge (water deprivation or salt loading through high Na^+ drinking

water), and the following recovery. It was also based on data estimating the rate of

synthesis by measuring the accumulated vasopressin content at the cell bodies with

transport blocked to the peripheral stores (Roberts *et al.*, 1991). Synthesis rates were

estimated to be ~1-3 ng/h at basal, and 10 ng/h under hyper-osmotic conditions (3

days of salt loading). They also showed that the synthesis rates only gradually return

toward basal levels in the days following the osmotic challenge and that the transport

rates (from cell body to peripheral stores) up- and down-regulate in parallel (Roberts

et al., 1991). This prolonged upregulation of synthesis and transport acts to replenish

the peripheral stores.

The Pittsburgh model, which only simulates synthesis and the store, using the

simplification that synthesis always tracks secretion, does not deal with the pathway

between osmotic stimulus and regulation of mRNA. A representation of this pathway

is required for our model, which takes as input a synaptic signal that encodes osmotic

pressure. It is well established that osmotic stimulation increases vasopressin mRNA

content (Sherman, McKelvy and Watson, 1986), and also known that hypo-osmotic

conditions reduce mRNA content (Svane *et al.*, 1995). As well as increasing

transcription rates, prolonged osmotic stimulation increases the length of the mRNA

poly(A) tails (Carrazana, Pasieka and Majzoub, 1988; Zingg, Lefebvre and Almazan,

1988), and the overall changes in content are likely due to a combination. Longer

poly(A) tails are thought to either increase mRNA stability or increase translation

efficiency (Emanuel *et al.*, 1998). The functional effect of either would be to increase

the amount of synthesis per unit of mRNA.

The pathway between osmotic stimulus and transcription is still uncertain. The

major candidate is a pathway via cyclic AMP (Carter and Murphy, 1989; Sladek *et*

al., 1996; Wong *et al.*, 2003) that acts to drive the CREB3L1 transcription promoter

(Greenwood *et al.*, 2015). There is also evidence for a glutamate-NMDA receptor-

Ca^{2+} entry driven pathway (Lake, Corrêa and Müller, 2019). Here we are using a very

simple representation to predict the necessary dynamics rather than any detailed

modelling of the mechanisms.

Our objective here was to integrate, adapt and extend the Pittsburgh model

into our existing integrated spiking and secretion model in order to fully simulate the

pathway from osmotic signal to plasma hormone signal. The challenge we identified

when testing the secretion model (MacGregor and Leng, 2013) is that heterogeneity,

which brings essential benefits to producing a robust signal response, results in widely

varying rates of secretion and store depletion across the population. The synthesis

mechanism must be able to cope with varied demand not only as a population but also

between individual neurons.

The new synthesis modelling has been kept as simple and general as possible,

and should be capable of being adapted to other neuroendocrine cells, but is still able

to produce strong quantitative as well as qualitative matches to the experimental data

on synthesis rates, mRNA content, and depletion and repletion of vasopressin stores

during prolonged osmotic challenge and recovery. However, in designing and fitting

to match experimental data that shows a cycle of depletion and recovery during and

after an osmotic challenge, the synthesis model is essentially constrained to fail at the

task of matching supply to demand. By attempting to fix this in the model we explore

why the stores get depleted; what are the limiting mechanisms, and why these limits

might be necessary.

Results

Osmotic Challenge and Recovery Data

To set targets for fitting and testing the model, an extensive literature survey

was used to gather multiple types of physiological data recorded in rats during a

prolonged osmotic challenge and the following recovery. This extends the examples

of (Fitzsimmons *et al.*, 1992) where they compared multiple sources measuring the

depletion and recovery of pituitary vasopressin stores. The data used here (Figure 1)

includes osmolarity, plasma Na concentration, plasma vasopressin concentration,

hypothalamic vasopressin mRNA content, and pituitary vasopressin content, during

depletion and recovery. The data was extracted and reconstructed mostly by using

graphics software (Adobe Illustrator) to measure points plotted in figures. A full list

of the sources and tables of the data are given in the supplementary material.

The comparisons between multiple sources are imperfect. Experiments use

different breeds and ages of rat, different timings of measurements (which are likely

to have circadian sensitivity), and different assay techniques. In particular plasma Na

is measured both by flame photometry and by electrode based techniques.

Measurements of plasma vasopressin by radioimmunoassay are dependent on varying

sample extraction techniques and assay antibodies. Using multiple sources has

attempted to provide as clear a consensus as can be achieved, providing data to fit and

test the input osmotic stimulus (osmolarity and plasma Na), the internal mRNA

content, and the output plasma vasopressin and pituitary content.

The osmotic stimulus protocols vary between using dehydration (water

deprivation) and salt loading (high Na drinking water) to raise osmolarity. In all the

measurements except for plasma vasopressin these two protocols appear to produce an

equivalent response (Figure 1). The lower plasma vasopressin concentrations

observed under salt loading (~4 pg/ml vs 15 pg/ml under dehydration) are inconsistent

with the similar rates of pituitary content depletion. Content depletion is likely to be a

more robust measure of sustained vasopressin secretion rates, and so the model here

targets the higher and more consistent plasma vasopressin concentrations observed

under dehydration.

Figure 1. Experimental data gathered during prolonged osmotic challenge and recovery
The data here is gathered from multiple published sources where rats have been measured
during a prolonged osmotic challenge consisting of several days of dehydration or salt loading
(high sodium drinking water), and the following recovery period, with normal water access
restored. During the challenge osmolarity and plasma Na (top panels) rise mostly linearly, with
some data showing a reduced rate of rise and plateau towards day 4 and 5. Radioimmunoassay
measured plasma vasopressin concentration (mid left) mostly shows a matching linear rise, but
data is mostly limited to three days, and varies in magnitude between dehydration (higher) and
salt loading (lower) protocols. Vasopressin cell body mRNA content (mid right) shows a mostly
linear rise after one day that eventually plateaus. The data is variable, but the most consistent
experiments, with more time points, suggest a five to eight fold rise in content. The core target
data for the model is the measurements of pituitary store content (bottom). During the challenge
there is a mostly linear fall in pituitary content, which slows after day 3, falling to around 15 to
30% of normal content. During recovery, where osmolarity rapidly (a few hours) returns to
normal, the stores are slowly replenished over about two weeks. The faster recovery shown
here (blue squares) is in rats made hypo-osmotic after the prolonged hyper-osmotic challenge.
Detail on the sources is given in supplementary Figure S1.

 The Spiking and Secretion Models

The spiking model used to generate the results here uses parameters (Table 1)

chosen to simulate a typical magnocellular vasopressin neuron, based on detailed fits

to *in vivo* recordings (MacGregor and Leng, 2012). As the synaptic input rate is

increased, spiking shifts from silence to irregular spiking, phasic patterned spiking

(long bursts and silences), increasing burst durations, and eventually continuous

spiking. Figure 2 shows phasic spiking in the model. The non-linear stimulus-

response properties of the secretory terminals (frequency facilitation and fatigue),

simulated by the secretion model, are such that the phasic pattern is optimal in terms

of secretion per spike. Thus the increase in the rate of secretion slows as the neuron is

driven into less optimal continuous spiking.

Figure 2. Spike activity dependent regulation of transcription
Phasic spiking in a highly stimulated integrate-and-fire based model neuron is both driven by
and generates Ca^{2+} entry, producing an intracellular Ca^{2+} signal that is used to drive the model's
vasopressin mRNA transcription rate. The essential dynamic is that the mechanism translates
the rapidly changing and noisy electrical activity into a sustained slow-changing measure of
activity.

The secretion model is modified from the previously published version as

described in the Methods. It is coupled to a model of plasma diffusion and clearance

which we previously developed to simulate oxytocin plasma concentrations (Maícas-

Royo, Leng and MacGregor, 2018), with parameters adjusted using experimental data

on vasopressin plasma concentrations and clearance rates, again as described in the

Methods.

Synthesis Model Basic Function and Tuning

The synthesis model was initially tested using a single neuron. The secretion

rate is scaled to the number of neurons to maintain comparable secretion and plasma

concentrations independent of population size.

The transcription rate (T) half-life $\lambda_T = 1000$ s and upregulation rate $k_T = 0.33$

were chosen to produce a plausible timescale transcription rate signal, taking with

input rate $I_{re} = 380$ Hz, ~1 h to reach equilibrium $T = \sim 60$ (arbitrary units) from an

initial $T = 0$ (Figure 2). This signal forms a long timescale measure of spike activity

which in turn drives the increase in mRNA content.

Figure 3 illustrates the basic function of the model with steady input rate $I_{re} = $

380 Hz, corresponding to a sustained hyperosmotic state. Vasopressin mRNA content

rises very slowly towards an equilibrium at a rate and level determined by the balance

between the transcription rate and depletion due to translation and synthesis. The rate

of synthesis is directly proportional to the mRNA content (m). The secretion rate and

plasma concentration driven by the single phasic neuron are noisy but sustain steady

levels until the reserve store is depleted. The releasable pool (which is refilled from

the reserve) buffers the secretion response to maintain a steady rate until the reserve

store is very heavily depleted. With no synthesis, the store fully depletes and plasma

concentration falls to zero. With synthesis, the rate is insufficient to match the highly

stimulated secretion rate and the store is still depleted, though at a slower rate. When

it is depleted, secretion and plasma concentration is sustained, at a level purely

dependent on the upregulated synthesis rate. We would not expect to observe this in

the heterogeneous population *in vivo*, but this is what we would predict in a

homogeneous population, assuming a sustained osmotic stimulus.

Figure 3. Single neuron transcription-dependent regulation of mRNA content and synthesis rates.

For illustration, rather than physiological simulation, the model here is initialised with a full store and zero stimulus, switching at time 0 to a sustained highly osmotic input signal. Transcription drives the accumulation of mRNA content, which in turn determines the rate of synthesis which maintains (or slows the depletion of) hormone stores. In the rapid change to a highly stimulated state here, elevated synthesis is not sufficient to match the rate of secretion, and stores are gradually depleted. When the stores are depleted the rate of secretion becomes purely synthesis rate dependent.

The secretion model parameters were fixed by fitting the secretion model to *in vitro*

data as described in the Methods and detailed in (MacGregor and Leng, 2013;

Maícas-Royo, Leng and MacGregor, 2018). Coupled to the plasma model, this allows

the prediction of the rates of secretion that correspond to plasma concentrations

observed *in vivo*. In basal normo-osmotic conditions rat plasma vasopressin *in vivo* is

~ 1 pg/ml. In highly stimulated hyper-osmotic conditions plasma vasopressin *in vivo*

rises to around 20 pg/ml. The left panels of Figure 4 show the single neuron model

sustaining a mean 1 pg/ml plasma concentration with input rate I_{re} = 252 Hz. The

initial value for mRNA content (m = 15), was set using an initial test to find its stable

value at this input rate. The first target for tuning the synthesis model parameters was

for synthesis to match secretion in basal conditions, in order to sustain a stable reserve

store. The reserve store plot shows this achieved by reducing the synthesis rate (s_r =

0.65) compared to the final heterogeneous model parameter (s_r = 1.1), which produces

a small rise in the store, with synthesis exceeding secretion.

Simulating Sustained Osmotic Challenge and Recovery

The second target for tuning the model was to match the experimental data

measuring vasopressin store content in rats during a five day osmotic challenge (no

water access, or salt loading using high Na^+ drinking water) and the following

recovery (Figure 1). Experiments measuring osmolarity (or equivalent plasma Na^+)

and plasma vasopressin during similar protocols (Walters and Hatton, 1974;

Nordmann, 1985; Yue *et al.*, 2008) suggest that the osmotic stimulus rises mostly

linearly during the challenge for at least the first three days before levelling off at

sustained high levels, and rapidly recovering after the challenge period. We simplified

this by using a linear ramp in the input rate to simulate the prolonged osmotic

challenge, illustrated in the right hand panels of Figure 4. The initial input rate 252 Hz

was ramped to 640 Hz over 5 days and then returned to 252 Hz, targeted to match the

store depletion observed in the experimental data.

The transcription rate mostly tracks the osmotic stimulus. The stores decline in

content to ~ 25% before slowly recovering following the challenge, matching the

experimental data and the results with the original Pittsburgh synthesis model. The

mRNA content shows a more non-linear increase, and decline during the recovery

period, as it sustains elevated synthesis rates to replenish the stores. However, the

reduced synthesis rate (s_r = 0.65) used to match secretion at basal levels (Figure 4 left)

Figure 4. Single neuron basal activity and prolonged osmotic challenge and recovery
The plots on the left show basal activity sustaining a mean 1pg/ml plasma vasopressin. The
transcription rate rapidly rises to sustain mRNA content at 15 units. With default synthesis
rate scaling $s_r = 1.1$, the synthesis rate slightly exceeds the secretion rate and the reserve store
increases (blue). Setting $s_r = 0.65$ (green) maintains a stable store. Removing synthesis by
setting $s_r = 0$ shows a depleting store. The plots on the right show a 5 day osmotic challenge
(a linear ramp from basal, simulating progressive dehydration or salt loading) followed by 10
day recovery (input returned to basal). The transcription rate mostly tracks the osmotic
stimulus with some drop off due to non-linearities in the spiking response. The mRNA
content rises non-linearly as the balance shifts between transcription, and depletion due to
synthesis (translation). The synthesis rate increases but fails to track the increasing secretion
rate. The reserve store is gradually depleted to ~27%. Plasma vasopressin increases initially
linearly but then slows as the neurons shift from phasic to continuous spiking. Following the
ramped challenge, secretion falls to basal rates, but elevated synthesis is sustained by the
increased mRNA content, depleting this to recharge the reserve store.

produces more depletion of the stores and an incomplete recovery. As shown below,

the synthesis model was more difficult to tune for a homogeneous than a

heterogeneous population.

Synthesis Response in a Heterogeneous Population

Osmotically stimulated vasopressin neurons recorded *in vivo* show widely

varying spiking rates. We can simulate this heterogeneity by randomly varying the

amount of synaptic input received by each model neuron, applying a varied input

density parameter, as detailed in the Methods. This heterogeneity has a substantial

functional advantage, producing a much more linear plasma vasopressin response to a

changing osmotic input signal than a homogeneous population (MacGregor and Leng,

2013). This matches the response that has been observed experimentally, however it

results in also producing highly heterogeneous secretion rates and store depletion

across the population. Here we tested how the synthesis model would respond to these

varied stimulus and store depletion rates, and how this would affect the summed

population response. The model was set up to record both the summed population and

the individual neuron mRNA content, reserve store, and secretion rates.

A 100 neuron heterogeneous population was randomly generated with a

lognormal distribution, illustrated in the inset of Figure 5. The basal population input

rate (which is modified by each neuron's input density) was set to 207 Hz to produce

a sustained 1 pg/ml plasma vasopressin concentration. The initial values for m were

set for each neuron by running the model with a 207 Hz population input until the

neurons' m values had stabilised, starting with common values of $m = 10$. No other

parameter adjustment was required from the same basal protocol tested with the

homogeneous model (Figure 4). This produced both a stable plasma vasopressin

signal and a stable summed reserve store. As well as the summed population data

Figure 5 shows a sample of three neurons from the low, middle, and high end of the

activity distribution. The low and middle neurons (blue and green) both had very low

spiking and secretion rates. Their mRNA content m was down-regulated to almost

zero, with a matching low synthesis rate. This matches what is observed

experimentally in hypo-osmotic conditions, showing the ability to down-regulate as

well as upregulate from the basal synthesis activity. The high activity neurons

(example here in red) show an increased mRNA content and a sustained secretion rate

much higher than the population mean. They also show a gradual increase of their

Figure 5. Basal activity and prolonged challenge and recovery in a heterogeneous population

The data here show a 100 neuron heterogeneous population following the same protocols as Figure 4. The inset distribution shows the heterogeneous input rates and the colour coded example neurons. In the left hand plots, with default synthesis scaling $s_r = 1.1$, the heterogeneous population sustains a stable reserve store at basal activity (1 pg/ml plasma vasopressin). The majority of the secretion is from the more active neurons (red) and these show gradual increase of individual neuron stores. In the right hand plots, the challenge and recovery protocol shows similar results to the single neuron (Figure 4), but with a more stable and linear plasma signal. The individual neurons show some complex divergence in their store recovery, due to varied non-linearities in secretion and synthesis. The more active cells, even with highly elevated mRNA, show more rapid depletion, but also a more rapid, and even excess recovery.

stores as synthesis exceeds the secretion rate.

Testing the ramped challenge and recovery protocol (initial population input rate 207 Hz ramped to 595 Hz over 5 days then returned to 207 Hz), plasma vasopressin and the summed population data shows very similar results to the homogeneous population. In this highly stimulated protocol the middle activity neuron more closely matches the mean population rates. The secretion rates of

individual neurons are much more non-linear than the population mean, and the

plasma concentration shows a more linear response to the ramped stimulus than the

homogeneous population.

Figure 6. Model challenge and recovery compared to experimental data
The 100 neuron heterogeneous population here simulates 2 days of basal activity, followed by
5 days of dehydration, and 15 days of recovery, compared against experimental data from
Figure 1. Two input signal protocols are compared, a default linear ramp (black) and a non-
linear ramp (red) with a more rapid initial increase in osmotic signal. Within the variability of
the experimental data both ramps are potentially consistent. The linear ramp varies from the
store data in its slower initial decline in content, more closely matched by the non-linear ramp,
which produces a faster increase in secretion than synthesis, resulting in more rapid initial store
depletion. The model was further tested with an added delay between synthesis and store fill
rate (blue), simulating the estimated 24h transport delay. The delay only moderately changes
the population store depletion and recovery profile, however more active neurons become fully
depleted, resulting in a drop off in the plasma signal.

Model Compared to Experimental Data

Figure 6 uses the same heterogeneous population model as Figure 5, with an extended time protocol for comparison with the experimental data, 2 days of basal activity followed by a 5 day osmotic challenge and 15 days of recovery. The results show a population store that falls to ~ 25 % during the 5 day challenge and replenishes to almost full over the 15 day recovery period, very similar to the experimental data in rats. A notable difference however is that the model's store using the linear ramp (black) protocol shows a slower initial decline. The match was improved by using a more non-linear ramp (red) in the osmotic input signal, producing a more rapid initial increase that gradually slows. Experimental evidence for the ramp in osmotic input and plasma vasopressin is variable and limited by temporally sparse measurements but suggests something that lies between these two. The model was further modified (blue) by adding a 24 h delay between the synthesis rate and the store fill rate representing the slow transport of new vesicles from the cell body to the pituitary stores, thought to take up to 24 h depending on osmotic status (Russell, Brownstein and Gainer, 1981). The delay more closely matches the depletion observed experimentally, but its effect is modulatory, and not sufficient to explain store depletion alone.

What Limits the Synthesis Response?

The current model matches the limited upregulation of synthesis, and depletion of stores, observed in the experimental data. Changes to the model, attempting to improve this response, were tested to predict which mechanisms might be responsible for this limited ability to match secretion demand (Figure 7). Two methods were found which were able to produce a much faster upregulation of synthesis while maintaining the ability to function in basal and stimulated states without under- or over-filling the stores. The first method (red in Figure 7) accelerates the upregulation of transcription. This required three parameter changes, increasing the rate of transcription but also compensating the increased amount so that only the speed of the response was changed (k_T 0.33 to 3.3, s_{basal} 0.7 to 7, s_r 1.1 to 0.11). The produces a much faster increase in the mRNA store and corresponding synthesis rate, resulting in a much smaller vasopressin store depletion. However it also predicts a much larger increase in mRNA than is observed experimentally (20-fold compared to ~5 to 8-fold). The second method (blue in Figure 7) increases the rate of translation

Figure 7. Improving synthesis performance by enhancing transcription or translation
Two enhancements are compared to the basic (black) experimental data fitted model (Figure 5 and 6), attempting to predict what limits the synthesis response *in vivo*. Accelerating transcription (red) produces a much faster upregulation of mRNA content and synthesis rate, reducing store depletion but depending on a larger mRNA capacity. Increasing the proportional translation rate (blue) similarly produces a faster upregulation of synthesis, with a more rapid depletion of mRNA content resulting in a lower equilibrium level. This reduces store depletion but likely depends on a translation rate which is beyond the capability of the cells.

(s_{basal} 0.7 to 3), increasing the rate of synthesis in exchange for a faster depletion of the mRNA store. This similarly produces a faster upregulation and reduced vasopressin store depletion but also results in a much smaller increase in the mRNA store, since the mRNA equilibrium level is determined by the balance between transcription and translation. Thus, the model predicts that the main elements responsible for vasopressin store depletion are the lag in upregulation of mRNA, and the maximum mRNA capacity, combined with a limit on the rate of translation. It may be that cells are capable of exceeding these limits, but that it is not efficient to maintain this capacity.

Discussion

Discussion

This study is part of a project aimed at understanding how vasopressin

neurons function as part of a physiological system on very long time scales. On the

surface they appear to perform a very simple signal processing task, producing a

plasma hormone signal that linearly encodes osmotic pressure. However, they have

many complex features, in particular their distinctive phasic firing, its relationship

with the highly non-linear properties of their secretory terminals, and their highly

heterogeneous activity levels. The phasic firing is asynchronous and not reflected in

the functional signal of their plasma summed secretory output. It appears that the

complexity is not about computation, but about being robust, adaptable, and efficient,

and maintaining function over lifelong periods of time. This relationship between

complexity and function is likely to be true of many neuro-physiological systems, and

the experimentally accessible and well-studied vasopressin neurons therefore serve as

a very good model system.

Essential to understanding the long term function of neuroendocrine

neurons (and other endocrine cells) is the dynamics of hormone storage and synthesis

and the focus here has been building and testing a synthesis model to integrate with

our existing spiking and secretion model. The new model is built on the work of

Fitzsimmons et al (Fitzsimmons *et al.*, 1992) which showed that activity-dependent

upregulation of mRNA content could best explain the experimentally observed

dynamics of store depletion and recovery. The challenge here was to integrate and

adapt the model to function without any direct tie between the rates of synthesis and

secretion, and for it to function within individual neuron models as part of a

heterogeneous population. This has been successful, providing further evidence that

the accumulation of mRNA is key to synthesis dynamics. In normal and hypoosmotic

conditions mRNA content functions to measure and service current demand. In

hyperosmotic conditions it serves as a memory of sustained challenge and following

the challenge provides a resource to recover depleted hormone stores.

The mechanisms of the robust new model components presented here are very

simple. The key to this was the strongly quantitative properties of the secretion and

plasma model. The existing vasopressin neuron model was also further developed

here by integrating a new model of hormone diffusion and clearance in plasma, and

by refining the quantitative scaling of the existing secretion model, based on previous

work in oxytocin neurons (Maícas-Royo, Leng and MacGregor, 2018). The

importance of this was in relating rates of secretion to experimentally observed

plasma concentrations, thereby accurately simulating hormone store depletion (and

recovery) and constraining synthesis rate demands. Initial attempts at building the

synthesis model, before the secretion rate scaling had been corrected, and the apparent

synthesis rate demands were much higher, used an additional activity-dependent

component for the translation rate, shown ghosted in Figure 8. This was not robust,

being very sensitive to the balance between parameter values driving the activity

dependent transcription and translation components. The version presented here uses

only a fixed translation rate, proportional to the mRNA content. Thus, making the

model more quantitatively accurate actually reduced the necessary complexity.

The model also uses a simple representation of the relationship between

osmotic stimulus and transcription, making use of the spiking model's Ca^{2+} variable.

Rather than the mechanism necessarily being Ca^{2+}-dependent, the necessary

assumption here is that transcription closely tracks spike activity. This helps the

model to maintain a tracking between the synthesis and secretion rates without any

cross-communication. If transcription was more directly driven by synaptic input then

the complexities of phasic firing would disrupt this tracking.

Where the model's behaviour becomes more complex is in the dynamics of

stores in a heterogeneous population. Heterogeneous activity levels would be

expected to present a big challenge to maintaining the tracking between synthesis and

secretion rates and it was surprising how robust the heterogeneous population proved

to be. This is partly because heterogeneity as well as adding complexity, also removes

some by linearising the relationship between the input and output signals. However,

there is some variation across the heterogenous population in how well stores are

maintained, suggesting that a statically heterogeneous population will gradually

diverge in store content. The simple model tested here puts no limits on the mRNA

content or vasopressin stores in individual neurons. These limits are likely to exist in

some form and would act to reduce the divergence between neurons, but it does

nevertheless seem likely that a static heterogeneous population would struggle to

maintain function over long periods. Thus the model here has developed a tool to

further examine rather than solve the problem of store divergence identified in the

previous work (MacGregor and Leng, 2013).

The alternative is dynamic or regulated heterogeneity. Here we refer to our

neurons as a population, connected only at their functional input and output signals.

However vasopressin neurons have the ability to communicate through dendritic

release of various signals including vasopressin, and potentially act as a network.

There is evidence that these signals act to modulate the activity of neighbouring

neurons (Gouzènes *et al.*, 1998) and it has been proposed that the network might act

to cycle activity levels (Scott and Brown, 2010), letting rested neurons replace those

that have been more active and become depleted. Such a mechanism would also

contribute to the lifetime robustness of the system by compensating for lost neurons.

The question for this that arises from the work here is what measure would regulate

the dendritic signals? For the same reasons that synthesis rates are not thought to be

directly coupled to secretion (distant and distributed stores), it would be difficult to

directly measure store depletion. Do the stores available for dendritic release deplete

sufficiently in parallel to the peripheral stores? Could elements of the synthesis

mechanism also regulate dendritic signalling?

One of the main limitations in interpreting the results here is the highly

simplified simulation of the prolonged osmotic challenge. The linear ramp is based on

experimental data measuring vasopressin concentrations, osmotic pressure, and/or

plasma Na^+ which show mostly linear increases with time during an osmotic

challenge over at least three days. After three days however, these increases tend to

slow, probably as the high sustained vasopressin output, and regulation of other

elements involved in osmotic homeostasis, such as salt excretion, achieve some sort

of new equilibrium. There is also evidence of this in the data measuring store content,

where the rate of depletion appears to fall towards the latter part of the challenge. We

began addressing this here in the model using a non-linear input ramp, but a much

better approach in terms of gaining understanding would be to integrate the neural

population model into a simple system model of osmotic homeostasis, providing

feedback between the vasopressin output and the osmotic input signal.

Another assumption here is the simple linear encoding between osmotic

stimulus and the rate of synaptic input. Recent work in oxytocin neurons (Maícas

Royo, Leng and MacGregor, 2019) modelling osmotic stimulus in more detail, to

simulate experiments in which plasma oxytocin was measured in response to Na^+

injections or infusions, supports this. The linear encoding assumption was sufficient

to closely match experimental plasma concentrations with the model, and it is

reasonable to assume similar in vasopressin neurons. The exceptions to this are likely

to be in special conditions such as pregnancy.

The work here has modelled the activity-dependent hormone synthesis

mechanisms of vasopressin neurons and integrated this into a model of spiking and

secretion, further refined and developed to accurately simulate plasma vasopressin

concentrations in response to dynamic osmotic stimuli. It has shown that the idea of

synthesis driven by the regulation of mRNA content remains robust without any

assumption of synthesis directly coupled to secretion, and within the complexities of

population heterogeneity. The model provides a strong base for future work exploring

the mechanisms that coordinate vasopressin neurons as a network to maintain function

over lifelong periods of time, including investigation of the dysfunction of these

systems.

Methods

The Spiking Model

Many vasopressin neurons when stimulated generate a distinctive phasic

pattern of spiking, alternating between sustained bursts and silences lasting tens of

seconds. This is modelled here using an integrate-and-fire based model (MacGregor

and Leng, 2012) modified to include a set of activity-dependent potentials that

modulate excitability to shape spike patterning and generate an emergent bistability,

matching the observed phasic firing and detailed spike patterning measured using

analysis such as the inter-spike interval (ISI) histogram and hazard function (Sabatier

et al., 2004). Importantly the model also matches the changes in the phasic spiking

that occur in response to a changing input signal.

The excitability modulating potentials include a hyperpolarising afterpotential

(HAP), a fast depolarising afterpotential (DAP), and a slow after hyperpolarisation

(AHP). Each of these is modelled using a single variable that is step incremented with

each spike and decays exponentially. This simple form has proven sufficient to

produce close quantitative matches to experimentally measured spike patterning and

is used here for the activity dependent elements of the model.

The phasic firing mechanism uses a slow DAP based on a Ca^{2+} inactivated

hyperpolarising K$^+$ leak current V_L (i.e. an activity-dependent depolarisation generated by switching off a hyperpolarisation). This is modulated by two opposing step-and-decay variables representing spike generated Ca^{2+} entry, and dendritic dynorphin release, which slowly accumulates to oppose the action of Ca^{2+} and reactivate the K$^+$ leak current, eventually terminating a burst and sustaining the period of inter-burst silence.

With more detail in (MacGregor and Leng, 2012; Maícas Royo *et al.*, 2016), the spiking model is summarised by:

$$V = V_{rest} + V_{syn} - HAP - AHPslow + DAP - V_L$$

where V is the membrane potential, V_{rest} is the resting potential, and V_{syn} is the summed synaptic input signal described below. AHPslow is the renamed AHP of (MacGregor and Leng, 2012) to distinguish it from the medium AHP of (Maícas Royo *et al.*, 2016). The default parameters are given in Table 1.

Table 1: Spiking Model Parameters

Name	Description	Value (units)
I_{re}	excitatory input rate	230 (Hz)
I_{ratio}	inhibitory input ratio	0.75
e_h	EPSP amplitude	3 (mV)
i_h	IPSP amplitude	-3 (mV)
λ_{syn}	PSP half life	7.5 (ms)
k_{HAP}	HAP amplitude per spike	60 (mV)
λ_{HAP}	HAP half life	9 (ms)
k_{DAP}	fast DAP amplitude per spike	1 (mV)
λ_{DAP}	fast DAP half life	150 (ms)
$k_{AHPslow}$	slow AHP activation factor	0.00012 (mV/nM)
$\lambda_{AHPslow}$	slow AHP half life	10000 (ms)
$C_{AHPslow}$	minimum [Ca]$_i$ to activate slow AHP	200 (nM)
C_{rest}	rest [Ca]$_i$	113 (nM)
k_C	[Ca]$_i$ increase per spike	11 (nM)
λ_C	[Ca]$_i$ half life	2500 (ms)
k_D	dynorphin activation per spike	2.693
λ_D	dynorphin half life	7500 (ms)
k_L	K$^+$ leak calcium sensitivity	36 (nM)
g_L	K$^+$ leak maximum voltage	8.5 (mV)
V_{rest}	resting potential	-62 (mV)
V_{thresh}	spike threshold potential	-50 (mV)

Synaptic Input Signal

608
The osmotic stimulus is encoded by a mixed train of excitatory and inhibitory
post-synaptic potentials (EPSPs and IPSPs). Mixed synaptic input contributes to
producing a more linear spiking response to an increasing input signal (Leng *et al.*,
2001; Maícas Royo *et al.*, 2016). This synaptic input signal V_{syn} is simulated using a
Poisson random process to generate small (3 mV) exponentially decaying positive and
negative perturbations to the membrane potential. The proportion of inhibitory to
excitatory PSPs uses a fixed value of 0.75, reduced from the previous 1.0 ratio, based
on more detailed modelling of magnocellular neurons (Leng, Leng and MacGregor,
2017). The strength of the stimulus is represented by the mean EPSP rate I_{re}.

Secretion and Plasma Model

Good quantitative scaling is essential to understanding the qualitative
properties of the neurons, and relating mechanism to function. A necessary element
for understanding the synthesis mechanisms is to be able to relate the input signal and
spiking activity to the output plasma concentration, in order to constrain the rates of
secretion and synthesis. Plasma concentration is the most accessible measure of
secretion in the experimental data. Our previous work developing the vasopressin
secretion model used a simple, single volume estimate of the relation between
secretion rate and plasma concentration (MacGregor and Leng, 2013). More recently
we adapted the secretion model to oxytocin neurons and integrated a new model of
plasma diffusion and clearance (Maícas-Royo, Leng and MacGregor, 2018) which is
able to accurately predict experimental measurements of oxytocin plasma
concentration in response to both an acute stimulus (CCK injection, (Maícas-Royo,
Leng and MacGregor, 2018)) and slower osmotic challenges (Maícas Royo, Leng and
MacGregor, 2019).
The plasma model's volume, clearance, and diffusion rate parameters were
fitted using experimental data testing exogenous infusions of oxytocin (Ginsburg and
Smith, 1959; Fabian *et al.*, 1969). It models peripherally secreted oxytocin as
distributed between plasma and extravascular fluid (EVF) compartments, with
roughly similar volumes (8.5 ml and 9.75 ml respectively for a 250g rat), diffusing
between the two according to the concentration gradient with a time constant
estimated by the experimental data. Clearance is a single component from the plasma,

	2013	2022
α	0.0005	0.00009
c_θ	0.070	0.067
c_n	5	3
e_θ	2.8	3.0

Figure 8. Refinement of the secretion model
The quantitative fit of the published secretion model (MacGregor and Leng, 2013) was improved using more detailed *in vitro* data and better parameter tuning methods based on recent work applying the model to oxytocin neurons (Maícas-Royo, Leng and MacGregor, 2018). The model is fitted to data measuring both frequency facilitation (top panel) and fatigue (lower panel), simulating the in vitro experimental protocols. The changed parameters are given in the table. The major adjustment was to reduce α which scales the rate of secretion to units of pg.

representing the total clearance from the kidneys and other organs. There are no

equivalently detailed data available for vasopressin, however the vasopressin peptide

has a very similar size and transport properties, and we assume that the same volumes

and diffusion can be applied to vasopressin.

There are differences however in the clearance rates, mainly due to the added

component of bound vasopressin cleared at the kidneys. Experiments measuring the

stable plasma concentrations in response to sustained infusions of oxytocin and

vasopressin (Robinson *et al.*, 1989) estimated the total clearance rate of vasopressin as

almost exactly double that of oxytocin. This is consistent with previous experiments

that show higher oxytocin concentrations in response to the same stimulus (Dogterom,

Van Wimersma Greidanus and Swabb, 1977; Windle *et al.*, 1993). Thus we modified

the plasma model by reducing the clearance half-life parameter from 68s to 34s.

Combined with the diffusion component this produces an overall clearance half-life of

~51s. This matches the estimate of (Ginsburg and Heller, 1953) but is shorter than

other estimates of 120s (Czaczkes and Kleeman, 1964).

As well as adding the plasma model we also refitted the existing vasopressin

secretion model (MacGregor and Leng, 2013) using the same technique and

equivalent data as used to fit the oxytocin secretion model (Maícas-Royo, Leng and

MacGregor, 2018). The improved fits (Figure 8) reduced the scaling of secretion per

spike (parameter α) by a factor of seven, consistent with the smaller total functional

volume estimate of the improved plasma model (18 ml reduced from 100ml).

The Synthesis Model

The development of the synthesis model tested many more complex forms

than those presented here. Our aim was to produce a concise and robust model which

is sufficient to make a close qualitative and quantitative match to the available

experimental data. The new model adds only two new variables to the integrated

vasopressin neuron spiking, secretion, and plasma model, representing the rate of

transcription, and the mRNA store (Figure 9).

Transcription is upregulated with the osmotic stimulus. Without any

quantitative data available we have not attempted to make any detailed model of the

proposed cAMP or other messenger dependent pathway. The essential property of this

mechanism is that it needs a sustained activity-dependent signal, acting on a much

slower timescale than the rapidly changing electrical activity of the neuron. Informed

by previous experience of signal transduction across timescales in modelling circadian

and circannual rhythms (Macgregor and Lincoln, 2008) this uses a two-step process.

The spiking model's existing activity-dependent Ca^{2+} variable (C) is used, relative to

basal Ca^{2+} (C_{rest} = 113 nM), to drive the transcription rate (T), which increases in

proportion to C at rate 0.001 k_T units per s, and decays exponentially with half-life λ_T

= 1000s:

$$\frac{dT}{dt} = 0.001\, k_T(C - C_{rest}) - \frac{T}{\tau_T}$$

$$\tau_T = \frac{\lambda_T}{\ln(2)}$$

Figure 9. The integrated synthesis model
The spiking model, stimulated by synaptic input that encodes osmotic pressure, drives the
synthesis model through its Ca^{2+} variable C. This regulates the transcription rate T which
increases the store of vasopressin mRNA m. The synthesis rate s is proportional to m using a
fixed translation rate, and also depletes m. The ghosted link showing activity-dependent
regulation of translation was not used in the results here. The synthesis model is coupled to
the secretion model through the charging of its reserve store r.

The half-life is very approximate, chosen to produce a slowly changing value that
reaches an equilibrium proportional to the osmotic input stimulus (Figure 2). The
0.001 scaling factor produces a more convenient scale for parameter k_T.

The mRNA store (m) accumulates at a rate proportional to T. Contrary to the
Pittsburgh model, it has no explicit exponential decay component, but is depleted in
proportion to the synthesis rate (s). However, m does decay exponentially when
translation is at a fixed rate proportional to m (parameter tl_{basal}), as it is in the results
here.

$$\frac{dm}{dt} = s_{scale}T - s$$

$$s = tl_{basal}s_{scale}m$$

The model includes functional timescales ranging from ms to days, and parameter
s_{scale} is used to scale the rates between the spiking and secretion model and the

720 synthesis model. The results here use a fixed value, but it was convenient for testing

721 to be able to use this parameter to accelerate the synthesis timescale.

722 The reserve store r equation from the secretion model (eqn 8 in (MacGregor

723 and Leng, 2013)) was modified to add the synthesis component:

725
$$\frac{dr}{dt} = s_r s - \beta \frac{r}{r_{max}} \; where \; p < p_{max}; \; s_r s \; otherwise$$

where parameter s_r scales the synthesis units to pg units of stored vasopressin, and

parameter β is the refill rate of the secretion model's releasable pool p. The default

parameters are given in Table 2. To simulate a transport delay between synthesis and

the store, s in this equation was replaced s_{delay}, using the recorded value for s from an

earlier timestep.

Table 2: Default Synthesis Model Parameters

Name	Description	Value (units)
k_T	transcription upregulation rate	0.33 (T units s^{-1})
λ_T	transcription rate half-life	1000 (s)
tl_{basal}	fixed translation rate	0.7
s_{scale}	synthesis time scaling factor	0.000003
s_r	vasopressin scaling factor	1.1 (pg per s unit)

Population Simulation and Heterogeneity

The population was simulated by running in parallel 100 copies of the coupled

spiking, secretion, and synthesis model, with the secretion rate outputs forming a

summed input to the single plasma model. A heterogeneous population is generated

by randomly varying the proportional input rates (synaptic input density I_{syn}) for each

neuron using a lognormal distribution with mean = 0 and standard deviation = 0.25.

This approximates the highly heterogeneous range of spiking rates observed

experimentally (MacGregor and Leng, 2013). The stimulus is then represented by the

population input rate I_{pop} and individual neuron input rates are generated using:

746
$$I_{re} = I_{pop} I_{syn}$$

Implementation

The differential equations were integrated using the first order Euler method. We can do this safely since the step size (1 ms) inherited from the spiking model is much smaller than any of the time constants in the model. Using the same fixed time step makes it simple to couple the synthesis model, and the secretion and plasma models, to the integrate-and-fire based spiking model. The modelling software was developed in C++, using the open source wxWidgets graphical interface library. Each neuron runs as a duplicate integrated spiking, secretion, and synthesis model thread, with secretion rates feeding into a single plasma model thread. A typical run of the full model, simulating 22 days of activity for a population of 100 neurons takes ~18 minutes on an AMD Ryzen 9 5900X 12-core processor.

The model source code, and software, compiled for Windows PC, are available at https://github.com/HypoModel/MagNet/releases.

Acknowledgement

Professor Gareth Leng is gratefully acknowledged for his contribution to discussions on the project and advice on editing the manuscript.

References

Carrazana, E.J., Pasieka, K.B. and Majzoub, J.A. (1988) 'The vasopressin mRNA poly(A) tract is unusually long and increases during stimulation of vasopressin gene expression in vivo', *Molecular and Cellular Biology*, 8(6), pp. 2267–2274. doi:10.1128/mcb.8.6.2267-2274.1988.

Carter, D.A. and Murphy, D. (1989) 'Cyclic nucleotide dynamics in the rat hypothalamus during osmotic stimulation: in vivo and in vitro studies', *Brain Research*, 487(2), pp. 350–356. doi:10.1016/0006-8993(89)90839-1.

Czaczkes, J.W. and Kleeman, C.R. (1964) 'The effect of various states of hydration and the plasma concentration on the turnover of antidiuretic hormone in mammals', *The Journal of Clinical Investigation*, 43, pp. 1649–1658. doi:10.1172/JCI105040.

Dogterom, J., Van Wimersma Greidanus, T.B. and Swabb, D.F. (1977) 'Evidence for the release of vasopressin and oxytocin into cerebrospinal fluid: measurements in plasma and CSF of intact and hypophysectomized rats', *Neuroendocrinology*, 24(2), pp. 108–118. doi:10.1159/000122702.

Emanuel, R.L. *et al.* (1998) 'Vasopressin messenger ribonucleic acid regulation via the protein kinase A pathway', *Endocrinology*, 139(6), pp. 2831–2837. doi:10.1210/endo.139.6.6043.

Fabian, M. *et al.* (1969) 'The release, clearance and plasma protein binding of
oxytocin in the anaesthetized rat', *The Journal of Endocrinology*, 43(2), pp. 175–189.
doi:10.1677/joe.0.0430175.

Fitzsimmons, M.D. *et al.* (1992) 'Models of neurohypophyseal homeostasis', *The*
American journal of physiology, 262(6 Pt 2), pp. R1121-1130.

Ginsburg, M. and Heller, H. (1953) 'The clearance of injected vasopressin from the
circulation and its fate in the body', *The Journal of Endocrinology*, 9(3), pp. 283–291.
doi:10.1677/joe.0.0090283.

Ginsburg, M. and Smith, M.W. (1959) 'The fate of oxytocin in male and female rats',
British Journal of Pharmacology and Chemotherapy, 14, pp. 327–333.
doi:10.1111/j.1476-5381.1959.tb00252.x.

Gouzènes, L. *et al.* (1998) 'Vasopressin regularizes the phasic firing pattern of rat
hypothalamic magnocellular vasopressin neurons', *The Journal of neuroscience: the*
official journal of the Society for Neuroscience, 18(5), pp. 1879–1885.

Greenwood, M. *et al.* (2015) 'Transcription factor CREB3L1 mediates cAMP and
glucocorticoid regulation of arginine vasopressin gene transcription in the rat
hypothalamus', *Molecular Brain*, 8(1), p. 68. doi:10.1186/s13041-015-0159-1.

Jones, C.W. and Pickering, B.T. (1969) 'Changes in the hormone content of the rat
neurohypophysis induced by substituting 2 per cent saline for drinking water', *The*
Journal of Endocrinology, 43(1), p. vi.

Jones, C.W. and Pickering, B.T. (1972) 'Intra-axonal transport and turnover of
neurohypophysial hormones in the rat', *The Journal of Physiology*, 227(2), pp. 553–
564. doi:10.1113/jphysiol.1972.sp010047.

Lake, D., Corrêa, S.A.L. and Müller, J. (2019) 'NMDA receptor-dependent signalling
pathways regulate arginine vasopressin expression in the paraventricular nucleus of
the rat', *Brain Research*, 1722, p. 146357. doi:10.1016/j.brainres.2019.146357.

Leng, G. *et al.* (2001) 'Responses of magnocellular neurons to osmotic stimulation
involves coactivation of excitatory and inhibitory input: an experimental and
theoretical analysis', *The Journal of Neuroscience*, 21(17), pp. 6967–6977.

Leng, G. and Ludwig, M. (2008) 'Neurotransmitters and peptides: whispered secrets
and public announcements', *The Journal of physiology*, 586(Pt 23), pp. 5625–5632.
doi:10.1113/jphysiol.2008.159103.

Leng, T., Leng, G. and MacGregor, D.J. (2017) 'Spike patterning in oxytocin
neurons: Capturing physiological behaviour with Hodgkin-Huxley and integrate-and-
fire models', *PloS One*, 12(7), p. e0180368. doi:10.1371/journal.pone.0180368.

MacGregor, D.J., Clayton, T.F. and Leng, G. (2013) 'Information coding in
vasopressin neurons - The role of asynchronous bistable burst firing', *Bio Systems*
[Preprint]. doi:10.1016/j.biosystems.2013.03.010.

MacGregor, D.J. and Leng, G. (2012) 'Phasic Firing in Vasopressin Cells:

Understanding Its Functional Significance through Computational Models', *PLoS*
Comput Biol, 8(10), p. e1002740. doi:10.1371/journal.pcbi.1002740.

MacGregor, D.J. and Leng, G. (2013) 'Spike Triggered Hormone Secretion in
Vasopressin Cells; a Model Investigation of Mechanism and Heterogeneous
Population Function', *PLoS Comput Biol*, 9(8), p. e1003187.
doi:10.1371/journal.pcbi.1003187.

Macgregor, D.J. and Lincoln, G.A. (2008) 'A physiological model of a circannual
oscillator', *Journal of biological rhythms*, 23(3), pp. 252–264.
doi:10.1177/0748730408316796.

Maícas Royo, J. *et al.* (2016) 'Oxytocin Neurones: Intrinsic Mechanisms Governing
the Regularity of Spiking Activity', *Journal of Neuroendocrinology*, 28(4).
doi:10.1111/jne.12358.

Maícas Royo, J., Leng, G. and MacGregor, D.J. (2019) 'The spiking and secretory
activity of oxytocin neurones in response to osmotic stimulation: a computational
model', *The Journal of Physiology*, 597(14), pp. 3657–3671. doi:10.1113/JP278045.

Maícas-Royo, J., Leng, G. and MacGregor, D.J. (2018) 'A Predictive, Quantitative
Model of Spiking Activity and Stimulus-Secretion Coupling in Oxytocin Neurons',
Endocrinology, 159(3), pp. 1433–1452. doi:10.1210/en.2017-03068.

Nordmann, J.J. (1985) 'Hormone content and movement of neurosecretory granules in
the rat neural lobe during and after dehydration', *Neuroendocrinology*, 40(1), pp. 25–
32. doi:10.1159/000124047.

Roberts, M.M. *et al.* (1991) 'Vasopressin transport regulation is coupled to the
synthesis rate', *Neuroendocrinology*, 53(4), pp. 416–422. doi:10.1159/000125750.

Robertson, G.L., Shelton, R.L. and Athar, S. (1976) 'The osmoregulation of
vasopressin', *Kidney International*, 10(1), pp. 25–37. doi:10.1038/ki.1976.76.

Robinson, A.G. *et al.* (1989) 'Total translation of vasopressin and oxytocin in
neurohypophysis of rats', *The American Journal of Physiology*, 257(1 Pt 2), pp.
R109-117. doi:10.1152/ajpregu.1989.257.1.R109.

Robinson, A.G. and Fitzsimmons, M.D. (1993) 'Vasopressin homeostasis:
coordination of synthesis, storage and release', *Regulatory Peptides*, 45(1–2), pp.
225–230. doi:10.1016/0167-0115(93)90210-y.

Russell, J.T., Brownstein, M.J. and Gainer, H. (1981) 'Time course of appearance and
release of [35S]cysteine labelled neurophysins and peptides in the neurohypophysis',
Brain Research, 205(2), pp. 299–311. doi:10.1016/0006-8993(81)90341-3.

Sabatier, N. *et al.* (2004) 'Phasic spike patterning in rat supraoptic neurones in vivo
and in vitro', *The Journal of Physiology*, 558(Pt 1), pp. 161–180.
doi:10.1113/jphysiol.2004.063982.

Scott, V. and Brown, C.H. (2010) 'State-dependent plasticity in vasopressin neurones:
dehydration-induced changes in activity patterning', *Journal of neuroendocrinology*,

22(5), pp. 343–354. doi:10.1111/j.1365-2826.2010.01961.x.

Sherman, T.G., McKelvy, J.F. and Watson, S.J. (1986) 'Vasopressin mRNA
regulation in individual hypothalamic nuclei: a northern and in situ hybridization
analysis', *The Journal of Neuroscience: The Official Journal of the Society for*
Neuroscience, 6(6), pp. 1685–1694.

Sladek, C.D. *et al.* (1996) 'cAMP regulation of vasopressin mRNA content in
hypothalamo-neurohypophysial explants', *The American Journal of Physiology*,
271(3 Pt 2), pp. R554-560. doi:10.1152/ajpregu.1996.271.3.R554.

Svane, P.C. *et al.* (1995) 'Effect of hypoosmolality on the abundance, poly(A) tail
length and axonal targeting of arginine vasopressin and oxytocin mRNAs in rat
hypothalamic magnocellular neurons', *FEBS letters*, 373(1), pp. 35–38.
doi:10.1016/0014-5793(95)01008-3.

Verbalis, J.G., Baldwin, E.F. and Robinson, A.G. (1986) 'Osmotic regulation of
plasma vasopressin and oxytocin after sustained hyponatremia', *The American*
Journal of Physiology, 250(3 Pt 2), pp. R444-451.
doi:10.1152/ajpregu.1986.250.3.R444.

Walters, J.K. and Hatton, G.I. (1974) 'Supraoptic neuronal activity in rats during five
days of water deprivation', *Physiology & Behavior*, 13(5), pp. 661–667.
doi:10.1016/0031-9384(74)90237-6.

Windle, R.J. *et al.* (1993) 'Patterns of neurohypophysial hormone release during
dehydration in the rat', *The Journal of Endocrinology*, 137(2), pp. 311–319.
doi:10.1677/joe.0.1370311.

Wong, L.-F. *et al.* (2003) 'cAMP-dependent protein kinase A mediation of
vasopressin gene expression in the hypothalamus of the osmotically challenged rat',
Molecular and Cellular Neurosciences, 24(1), pp. 82–90. doi:10.1016/s1044-
7431(03)00123-4.

Young, T.K. and Van Dyke, H.B. (1968) 'Repletion of vasopressin and oxytocin in
the posterior lobe of the pituitary gland of the rat', *The Journal of Endocrinology*,
40(3), pp. 337–342. doi:10.1677/joe.0.0400337.

Yue, C. *et al.* (2008) 'Differential kinetics of oxytocin and vasopressin heteronuclear
RNA expression in the rat supraoptic nucleus in response to chronic salt loading in
vivo', *Journal of Neuroendocrinology*, 20(2), pp. 227–232. doi:10.1111/j.1365-
2826.2007.01640.x.

Zingg, H.H., Lefebvre, D. and Almazan, G. (1986) 'Regulation of vasopressin gene
expression in rat hypothalamic neurons. Response to osmotic stimulation', *The*
Journal of Biological Chemistry, 261(28), pp. 12956–12959.

Zingg, H.H., Lefebvre, D.L. and Almazan, G. (1988) 'Regulation of poly(A) tail size
of vasopressin mRNA', *The Journal of Biological Chemistry*, 263(23), pp. 11041–
11043.