Turicibacter **strains differentially modify bile acids and host lipids**

Jonathan B. Lynch[1][§][*], Erika L. Gonzalez[1], Kayli Choy[1], Kym F. Faull[2,3,4], Talia Jewell[5],

Abelardo Arellano[5], Jennifer Liang[5], Kristie B. Yu[1], Jorge Paramo[1], Elaine Y. Hsiao[1]

[1]Department of Integrative Biology & Physiology, University of California Los Angeles, Los

Angeles, CA 90095, USA

[2]Department of Psychiatry and Biobehavioral Sciences, University of California Los Angeles,

Los Angeles, CA 90095, USA

[3]Jane and Terry Semel Institute for Neuroscience and Human Behavior, University of California

Los Angeles, Los Angeles, CA 90095, USA

[4]Pasarow Mass Spectrometry Laboratory, University of California Los Angeles, Los Angeles,

CA 90095, USA

[5]Isolation Bio, San Carlos, CA 94070, USA

[§]Current address: Department of Biological Chemistry, Johns Hopkins University, Baltimore,

MD , USA

*Correspondence to: jlynch48@jhmi.edu

<u>ABSTRACT</u>

Bacteria from the *Turicibacter* genus are prominent members of the mammalian gut microbiota

and are associated with alterations in dietary fat and body weight, but the specific connections

between these symbionts and host physiology are poorly understood. We genomically and

phenotypically characterized a diverse set of mouse- and human-derived *Turicibacter* strains,

and found they group into three clades that differ in their transformations of bile acids. We

identified *Turicibacter* bile salt hydrolases that confer strain-specific differences in bile

deconjugation. Colonization with individual *Turicibacter* strains led to changes in host bile acid

profiles, generally aligning with those produced *in vitro*. Further, colonizing mice with another

bacterium expressing bile-modifying genes from these strains decreased serum cholesterol and

triglycerides, as well as adipose tissue mass. This work identifies genes that enable diverse

Turicibacter strains to differentially modify host bile acids and lipid metabolism, and positions

multiple *Turicibacter* strains as candidates for altering host fat biology.

MAIN TEXT

Introduction

The gut microbiota forms complex relationships with its host organism, modulating broad

aspects of host physiology including metabolism[1,2] and neurobiology[3,4]. Often, the connections

between the gut microbiota and host physiology are easiest to decipher through presence/absence

of large sectors of the microbial community (examples in [5–7]), but in some cases, specific

microbial features and/or taxa serve important roles in host physiology[8–10].

The bacterium *Turicibacter sanguinis* is the best-studied species of the *Turicibacter*

genus, which is prevalent in the gut microbiota of several animals, including humans[11], mice[7],

cows[12], pigs[13], and chickens[14]. *Turicibacter* can reach relative abundances of >20% in the rodent

small intestine[15] and 0.5% in the human fecal microbiota[16,17]. A study of 1126 sets of twins

found *Turicibacter* was a highly heritable taxon in the human fecal microbiota[18], supporting

associations between *Turicibacter* abundance and host genetics found in Diversity Outbred

mouse collections[19–21]. Long-standing co-evolutionary connections between this taxon and

humans are suggested by the high abundance of *Turicibacter* sequences that have been measured

in ancient and non-industrial human gut microbiotas[17,22–24].

The mammalian gut microbiota has long been associated with obesity[25,26], but studies

often provide strong correlations rather than mechanistic determinants of these relationships,

indicating a further need for fundamental interrogation into connections between the microbiota

and host fat[27]. Numerous microbiota community profiling studies reveal correlations between

Turicibacter and features of host fat metabolism, such as adiposity and dietary lipids[28–33], but the

nature of these correlations varies[34,35]. We recently observed that the type strain of *T. sanguinis,*

MOL361[36,37], broadly alters the host serum lipidome while decreasing serum triglycerides in

mice[38]. This same strain was also reported to modify bile species through deconjugation and

dehydrogenation *in vitro*[21], suggesting at least one potential means by which *Turicibacter* can

influence host lipid status. Based on these findings, we hypothesized that there may be variations

in the functional activity of *Turicibacter* strains that account for differences in host bile and lipid

biology, providing a mechanism to connect this taxon to aspects of host physiology.

Results

Turicibacter *isolates separate into genetically distinguishable strains*

To better understand the diversity within the *Turicibacter* genus, we gathered nine

isolates from the fecal microbiotas of mice and humans that had been identified as *T. sanguinis*

based on their 16S rRNA gene sequence (97% full length 16S rRNA gene sequence similarity

cutoff, **Supplementary Table 1**). Two of these isolates had been previously identified (human-

derived type strain MOL361 and H121, which was derived from contaminated germ-free

mice[36,39]); five had been isolated but not published (human isolates 18F6, T46, and T129, and

mouse isolates 1E2 and TA25); and two were isolated from a human fecal sample specifically

for this study (GALT-E2 and GALT-G1) using an array-based isolation and cultivation platform

(see Methods). We performed shotgun short read sequencing and created draft assemblies of

each isolate genome. Comparisons of the 16S rRNA gene phylogeny (**Fig. 1a**), general genome

characteristics (**Fig. 1b**), or specific genome sequence (**Fig. 1c, d**) revealed that even with this

fairly small sample of the 16S rRNA gene-based "*T. sanguinis*" species designation, there were

at least three distinct subgroups: two from humans (exemplar isolates MOL361 and H121, with

99.3% full length 16S rRNA gene similarity), and one from mice (exemplar strain 1E2, 97.5%

and 97.8% 16S rRNA gene similarity with MOL361 and H121, respectively). Genomic

alignments indicated a substantial amount of shared DNA sequences within members of the

same subgroups (all within group average nucleotide identity [ANI] >98.3%) with the remaining

amount of genetic variation indicating smaller genetic differences between related isolates. These

within-subgroup shared sequences were distinct from members of the other two subgroups

(intergroup ANI: MOL361-H121=76.80%, MOL361-1E2=74.95%, H121-1E2=77.43%). It is

important to note that the H121-group genomically resembles the newly described species

Turicibacter bilis[13] (98.8% ANI), currently the only other named species from this genus.

Overall, these genomic differences suggest distinct evolutionary histories that correspond at least

partially with host origin.

Turicibacter *isolates differ in their impact on host fat biology and circulating metabolome*

Previous findings revealed that monocolonizing mice with *T. sanguinis* MOL361 altered

host fat tissue and circulating lipids[38]. Due to the large genomic variation between our

Turicibacter strains, we predicted that they would vary in their effects on host lipid biology. We

chose representative isolates from each of the distinct phylogenetic subgroups (MOL361, H121

and T129, and 1E2) and measured their effects on circulating metabolites and adipose tissue in

monocolonized mice relative to germ-free (GF) and conventionalized controls (*i. e.* gavaged with

complete microbiota, CONV). Compared to GF mice, CONV mice had decreased levels of

several dicarboxylate fatty acids, long-chain fatty acids, and long-chain acyl carnitines, with a

broad increase in short- and medium-chain acyl carnitines (**Fig. 2a**). Consistent with a previous

report[38], colonization with individual *Turicibacter* strains also induced widespread alterations in

host serum lipids, with several strain-level differences in host lipid alterations (**Fig. 2a,**

Supplementary Table 2). Compared to GF controls, MOL361 increased a subset of long-chain

acyl carnitines, and decreased many long-chain saturated fatty acids. In addition, colonization

with MOL361 elicited trending decreases in host cholesterol to levels below those seen in GF

(**Fig. 2b**). Compared to GF, H121 colonization significantly increased serum levels of several

medium-chain fatty acids, dicarboxylic acids, and short-, medium- and long-chain acyl

carnitines. 1E2 colonization had a smaller overall effect on host lipids, but led to a decrease in

several dicarboxlyate fatty acids (**Fig. 2a**). In addition to differences between GF or CONV mice

and *Turicibacter*-monocolonized animals, there were also alterations in host lipids that showed

some strain-dependency, with trending discrepancies in dicarboxylate and significant differences

in long chain saturated fatty acids (**Fig. 2a**) At the tissue level, two of the four *Turicibacter*

strains stimulated statistically significant increases in epididymal/gonadal white adipose tissue

(e/gWAT) mass in comparison to GF controls, and a third strain elicited similar increases that

were not statistically significant (T129). In contrast, there was no noticeable effect of H121 on

e/gWAT mass (**Fig. 2c, Supplementary Table 3**), distinguishing it from the other *Turicibacter*

strains. Consistent with this, H121 showed the smallest e/gWAT adipocyte size within fat pads

(**Supplementary Fig. 1a-f**). This may be due to lower colonization of H121 in both the small

intestine and the colon (**Supplementary Fig. 1g, h**) The effects of particular *Turicibacter* strains

on serum lipids, cholesterol, and fat mass directed us to investigate molecules that could

potentially connect these biomolecules: bile acids.

Bile acids can affect circulating host lipids by altering fat digestion and systemic

hormonal signaling[40]. They are produced by the host and released into the small intestine, where

they promote digestion of fats by facilitating micelle formation, and can also act through

receptors like the farnesoid X receptor (FXR)[40] and GPBAR1/TGR5[41]. Gut bacteria can modify

bile acids, primarily through transformations like deconjugation. Previous reports found that

MOL361 can broadly modify bile acids *in vitro*[21], so to determine if these abilities allowed

MOL361 and other *Turicibacter* strains to modify host bile acids *in vivo*, we profiled serum and

cecal bile acids in *Turicibacter*-monocolonized mice. Though each strain had some unique

impacts on host serum metabolites, we noted some consistent patterns in bile acids across

monocolonized mice in comparison to either their GF or CONV controls. (Note: bile species

with amino acid conjugants are typically referred to as "bile salts," but for simplicity, we will

herein refer to both conjugated and unconjugated bile species as "bile acids"). Colonization with

all of the *Turicibacter* strains led to a general increase in serum levels of unconjugated primary

bile acids like cholate (CA), chenodeoxycholate (CDCA), and β-muricholate (βMCA) (**Fig. 3a-**

d), and a similar rise in unconjugated secondary bile acids 3-dehydrocholate and 7-

ketodeoxycholate (**Fig. 3e,f**). These responses were highly variable in the case of T129

colonization, leading us to de-emphasize this strain for subsequent experiments. In response to

colonization with MOL361, 1E2, or H121, we also found a similar increase in several

unconjugated bile acids in the cecum (**Supplemental Fig. S2).** In all, these results suggest that

these *Turicibacter* strains are able to impact host bile acids, potentially by deconjugating them in

the gut.

When we investigated potential explanations for increases in unconjugated bile species,

we noticed that the serum levels of conjugated bile acids differed between animals colonized by

different *Turicibacter* strains, with the clearest delineation separating H121 from MOL361 and

1E2. MOL361- and 1E2-colonized animals generally had lower levels of taurine-conjugated

primary bile acids in comparison to H121-colonized animals (**Fig. 3g-j**), whereas H121

colonization led to an increase in glyco-beta-muricholic acid (GβMCA) (**Fig. 3k**). Female mice

displayed the most significant *Turicibacter*-associated changes in circulating bile species, lipids,

and cholesterol, indicating sex-dependent responses to *Turicibacter* colonization

(**Supplementary Fig. 3**). This sex difference has also been reported in C57BL/6 mice

monocolonized with *T. sanguinis* MOL361, albeit with a directionally different change in

adipocyte size[38]. Notably, in the cecum, there was less strain-specific distinction in conjugated

bile acids, with taurine-conjugated bile acids decreased by colonization with any of the three

Turicibacter strains we surveyed (**Supplementary Fig. S2**). The one cecal bile acid that did

display strain dependency was the secondary bile acid lithocholic acid (**Supplementary Fig. S2,**

bottom). Also, the low levels of detected glycine-conjugated bile acids (specifically glycocholic

acid) in both serum and cecal contents was slightly increased by *Turicibacter* colonization (**Fig.**

3k,l; Supplementary Fig. S2c) Overall, these data demonstrate that colonization with

Turicibacter alters serum bile acids, lipids, and cholesterol, as well as host fat mass. Further,

while some changes like increased unconjugated bile acids were conserved across *Turicibacter*

colonizations, levels of specific conjugated bile acids differed between *Turicibacter* strains, with

MOL361 and 1E2 leading to lower levels of taurine-conjugated bile acids than H121

colonization.

Turicibacter *strains differ in their ability to modify host bile acids*

Based on the large genetic variation between the *Turicibacter* strains (**Fig. 1**) and the

differences in aspects of serum lipid and bile acid profiles seen in response to colonization with

different strains (**Fig. 3**), we posited that these strains differ in their ability to modify bile acids.

To test this, we grew each of our nine isolates to stationary phase in rich medium supplemented

with a sub-inhibitory concentration of five bile species[21]: cholic acid, chenodeoxycholic acid,

deoxycholic acid (DCA), taurocholic acid (TCA) and glycochenodeoxycholic acid (GCDCA).

We then used liquid chromatography-mass spectrometry (LC-MS) to characterize the resulting

bile transformations performed by each isolate (**Fig. 4a**). We discovered that the strains not only

differed in their ability to modify this combination of bile species, but also that these capabilities

generally mirrored the groupings identified in genomic comparisons (**Fig. 4b**). MOL361, 18F6,

and GALT-E2 deconjugated both tauro- and glyco-bile acids, and also dehydrogenated CA and

CDCA (**Fig. 4b**). 1E2 and TA25 deconjugated tauro-bile acids, but did not detectably

deconjugate glyco-bile acids or perform dehydrogenation (**Fig. 4b**). H121 and T129

deconjugated glyco-bile acids, but did not readily deconjugate tauro-bile acids nor perform

detectable dehydrogenation (**Fig. 4b**). T46 and GALT-G1 did not have bile-modifying capacity

that mirrored their genetic phylogeny; T46 genomically resembled the MOL361 group but

performed modifications similar to the H121 group (*i.e.* glyco- but not tauro- deconjugation,

minimal dehydrogenation), whereas GALT-G1, which genomically resembled the H121 group,

performed more MOL361-like transformations (*i.e.* glyco- and tauro-deconjugation,

dehydrogenation of CDCA) (**Fig. 1, 4b, Supplementary Fig. 4**). Overall, each strain performed

at least one of three bile transformations, with some showing capacity for all three (**Fig. 4b,**

Supplementary Fig. 4).

To confirm these intragroup distinctions, we chose one isolate from each of the

subgroups (MOL361, H121, 1E2), and grew them in the presence of four primary conjugated

bile acids: TCA, taurochenodeoxycholic acid (TCDCA), glycocholic acid (GCA), and GCDCA.

This supported the same pattern seen above; MOL361 deconjugated both groups of bile acids,

1E2 preferentially deconjugated tauro-conjugates, and H121 preferentially deconjugated glyco-

conjugates (**Fig. 4c**). MOL361 and 1E2 displayed broad deconjugation of tauro-conjugates and

were able to process at least six taurine-conjugated bile acids (**Supplementary Fig. 5**). These

data reveal that while all tested strains are efficient modifiers of bile species, their specific

transformations differ in a strain-dependent manner, potentially reflecting functions that

influence their differential effects on host lipid biology.

Turicibacter *genomes have different repertoires of bile salt hydrolases*

The different bile modification abilities across strain subgroups suggested that each

contained unique repertoires of bile-modifying genes. Certain bacteria from the gut microbiota

dehydrogenate hydroxyl groups from the steroid core of bile acids[42], increasing their polarity and

modulating their affinity for host bile acid receptors[43]. We searched the *Turicibacter* genomes for

homologs of the characterized 7α-hydroxysteroid dehydrogenase (7α-HSDH[21]) from *Clostridium*

absolum[44]. This revealed genes with 57% amino acid identity in MOL361, 18F6, T46, and

GALT-E2, and homologs with 59% amino acid identity in H121 and T129 (**Supplementary Fig.**

6a). Though the H121-derived putative homolog had higher overall sequence identity than the

MOL361-derived homolog, it lacked certain features predicted to be critical for dehydrogenase

activity, such as the analogous Asp38 that is catalytically critical for this reaction[44]. Because *in*

vitro experiments showed only isolates from the MOL361 group performed bile

dehydrogenation, we cloned the putative 7α-HSDH homolog from MOL361 into *E. coli* C41-

pLys and then grew these cells in individual unconjugated bile acids that can be dehydrogenated:

CA, CDCA, and DCA. Indeed, the protein encoded by the gene removed the mass equivalent of

two hydrogens from CA (**Supplementary Fig. 6b, c**). Background transformation by *E. coli*

prevented clear evidence of CDCA dehydrogenation by this putative 7α-HSDH homolog

(**Supplementary Fig. 6d**), but this homolog did not act on DCA (**Supplementary Fig. 6e**),

supporting its annotation as a 7α-HSDH.

Conjugation increases bile acid solubility and emulsification ability[45], while

deconjugation reverses these effects, leading to decreased dietary lipid absorption. To identify

Turicibacter bile salt hydrolase (*bsh*) genes responsible for the strain-specific differences in bile

deconjugation, we first searched our assembled genomes for annotations of "choloylglycine

hydrolase," the broad category that includes these genes. Of these annotated genes, we identified

eight groups of potentially homologous sequences, and found that each *Turicibacter* strain

encodes putative choloylglycine hydrolases from at least two of the eight groups (**Fig. 5a, b**).

Isolates within the same phylogenetic and phenotypic subgroups largely shared similar

sequences (**Fig. 5a, b**). To assay the function of the strain-specific putative *bsh* genes, we cloned

and individually expressed one representative sequence from each of the eight putative

choloylglycine hydrolase groups in *E. coli* C41-pLys and measured the ability of these

engineered bacteria to perform the deconjugations we observed in our native *Turicibacter*. We

cultured the individual *E. coli* strains to stationary phase in the presence of two tauro- (TCA,

TCDCA) or glyco- (GCA, GCDCA) bile acids, then measured their ability to deconjugate these

bile acid pools. We found that *E. coli* expressing four of the eight putative *bsh* gene groups

showed deconjugation activity against at least one of the bile acids (**Fig. 5c-e**). From MOL361,

one BSH (group IV) is tauro-specific, and one (group I) deconjugates both glyco- and tauro-

conjugates (**Fig. 5c, d**). 1E2 shares a tauro-specific BSH with MOL361 (group IV), and has

another BSH (group III) with moderate activity on TCDCA (**Fig. 5c, d**). H121 has a BSH (group

II) with activity on TCDCA and GCDCA (**Fig. 5c, d**), although this was blunted when presented

with the combined four bile acids (**Fig. 5e**), potentially because of competitive inhibition.

Collectively, these findings reveal that *Turicibacter* strains contain a range of bile salt hydrolases

with deconjugation preferences for different bile acids.

Strain- and substrate-specific bile salt hydrolases from Turicibacter *differentially alter host lipid*

composition

Given that *Turicibacter* colonization broadly modified host lipid and bile pools (**Fig. 2**),

and that bile transformations have been previously shown to alter host lipids[46,47], we predicted

that expressing *Turicibacter* bile-modifying genes outside the context of *Turicibacter*

colonization would be sufficient to impact host lipid biology. To measure the individual effects

of these bile acid transformations, we expressed *Turicibacter bsh* genes off of a genomically-

integrated high expression vector[48] in the common gut bacterium *Bacteroides thetaiotaomicron*.

This bacterium was chosen because it stably colonizes the murine gut and, unlike *E. coli* C41-

pLys, contains a homolog of a characterized 7α-HSDH similar to that of *T. sanguinis*

MOL361[49,50], allowing the engineered bacteria to more completely mimic *Turicibacter* bile

transformations. As this *B. thetaiotaomicron* strain has previously been described to have some

BSH activity[46], we decided to employ the parental strain expressing the non-bile modifying

green fluorescent protein (GFP) as a comparison baseline for these experiments (named *Bt*-WT).

These engineered *B. thetaiotaomicron* strains generally transformed tauro- and glyco- conjugated

bile acids as predicted based on the BSH characterization in *E. coli*, including preferential

TCDCA transformation, indicating they were capable of performing *Turicibacter*-like bile

transformations (**Fig. 6a, Supplementary Fig. 7**). However, the *B. thetaiotaomicron* strain that

expressed *bsh* (group III) from strain 1E2 more completely transformed glyco-conjugated bile

acids than the strain expressing *bsh* (group II) from H121, counter to the *E. coli* findings. This

observation led us to not use the *bsh* (group II)-expressing strain further. Additionally, although

we did not notice a significant *in vitro* growth defect in the engineered *B. thetaiotaomicron*

strains (**Supplementary Fig. 8)**, we observed a bile transformation delay in *B. thetaiotaomicron*

expressing *bsh* (group I) from MOL361 that we could compensate for by extending the growth

period (**Fig. 6b**).

We monocolonized mice with the *bsh*-expressing *B. thetaiotaomicron* strains and

assessed their circulating lipid profiles and abdominal WAT mass. We found that the engineered

strains colonized the gnotobiotic mouse gut at least as well as the parental strain, with one strain

(BSH-III-1E2) colonizing the cecum slightly better (**Supplemental Fig. 9**), and that expressing

individual *Turicibacter bsh* genes in the *B. thetaiotaomicron* background was sufficient to

significantly alter host cecal bile levels (**Fig. 6c, Supplementary Fig. S10**) and the absolute

abundance of hundreds of circulating lipid species (**Fig. 6d, Supplementary Table 4**). In

particular, expression of either the group I or group IV *bsh* led to a decrease in triglycerides (**Fig.**

6d, e). Expressing the tauro-specific *bsh* (group IV) from MOL361 also decreased

diacylglycerides (**Fig. 6g**). Expressing the broader specificity *bsh* (group I) from MOL361 also

led to a decrease of phosphatidylglycines, phosphatidylserines, and cholesterol (**Fig. 6d-f, h, i**).

Despite having broad capacity for transformation, *B. thetaiotaomicron* expressing *bsh* (group III)

from 1E2 did not alter host lipid profiles as much as the other strains **(Fig. 6d-i)**. On a tissue-

level, *bsh*-expression also altered WAT storage in the colonized mice, with the broad tauro-

deconjugating BSHs (group I and IV) significantly reducing WAT mass (**Fig. 6j**). Similar to our

findings with *Turicibacter* monocolonization, we also observed a sex difference in BSH

responses, with male mice showing more consistent decrease in triglycerides in response to the

tauro-specific BSH from MOL361, and females showing more consistent triglyceride responses

to the broadly deconjugating BSH (**Supplementary Fig. 12**). Overall, these BSH-associated

lipid and adipose tissues results were consistent with previous findings on MOL361 colonization

in the C57Bl/6 mouse model[38], but differed from MOL361 colonization in our Swiss Webster

mice (**Fig. 2c, Supplementary Fig. S13**).

To further explore potential factors that may drive the cholesterol and WAT alterations in

response to *Turicibacter* colonization and *bsh* expression, we measured liver transcript levels of

farnesoid X receptor, (Fxr), a key nuclear receptor for bile acids; cytochrome P450 Family A

Subfamily A Member 1/Cholesterol 7α Hydroxylase (Cyp7a1), the rate-limiting enzyme for

conversion of cholesterol into bile acids; and glucose-6-phosphatase (G6pase), a key enzyme for

gluconeogenesis. There were no differences in Fxr transcript levels across any of the native and

engineered bacterial colonization conditions (**Supplementary Fig. 14a**). However, we found

similar increases in Cyp7a1 expression (**Supplementary Fig. 14b**) and decreases in G6pase

expression (**Supplementary Fig. 14c**) between the *Turicibacter* and *bsh* colonizations. These

findings suggest that the bile-modifying activities of *Turicibacter* lead to downstream alterations

in liver expression of bile signaling and response elements. In all, these results demonstrate that

expressing strain-specific *bsh* genes from *Turicibacter*, especially those able to process the

abundant taurine-conjugated bile acids present in the murine intestine, is sufficient to alter host

cholesterol, bile, and lipid biology.

Discussion

Results from this study show that several strains of *Turicibacter* bacteria from the

mammalian gut microbiota modulate host bile and lipid compositions. We identified and

characterized five novel *Turicibacter* genes capable of performing bile transformations (four *bsh*,

one 7α-HSDH), and revealed that expression of individual *bsh* genes is sufficient to broadly and

differentially alter host lipid profiles. Further, we found that while bile-transforming genes are

present in all our surveyed *Turicibacter* strains, the specific transformation capacity of BSH

variants differed by strain in a manner consistent with host environment co-evolution: bile acids

in the human gut are a mix of taurine- and glycine-conjugants, whereas murine bile acids are

predominantly taurine-conjugants[51,52], providing different bile environments that are

preferentially processed by *Turicibacter* strains isolated from their respective gastrointestinal

tracts. This close connection between host-specific bile composition and bacterial modifications

may be due in part to the bile sensitivity previously exhibited by MOL361 and/or the high

abundance of *Turicibacter* in the small intestine, causing these bacteria to more strongly

associate with host genes for bile reabsorption and lumenal bile levels than other bile-modifying

gut bacteria[21,53–57].

This work displays metabolic consequences of colonization by specific gut bacteria and

improves the resolution of our understanding connecting specific taxa—in this case, at the strain

level—with host physiology. In rodent and human studies, *Turicibacter* relative abundance often

negatively correlates with dietary fats[29,31,58–62] and host adiposity[28,33,63], but some studies have

shown opposite relationships[30,64,65]. This could be a result of the phenotypic diversity identified

here among *Turicibacter* isolates wherein the host may experience different lipid outcomes

depending on their own specific *Turicibacter* strains, but could also vary with other features such

as host genetics and sex[21,38] (**Supplementary Fig. 3, 11, 12**), leading to discrepancies in lipid

responses between hosts despite similar bile modifications performed in the gut microbiota.

Importantly, effects of *Turicibacter* colonization could also be influenced by biogeographic

organization of an individual's microbiota; in addition to the specific taxonomic membership*,*

Turicibacter positioning in the small and large intestine may affect host consequences from their

respective bile modifications by transforming bile pools in either section of the gut tract. Despite

genomic and localization differences between them, we found that MOL361 and H121 induced

lipid metabolite changes that indicate increases in fatty acid oxidation, suggesting that these

strains at least share features that alter host fatty acid metabolism. While we identified some host

lipids that were differentially modulated by different *Turicibacter* strains, host complexities

likely reduced our power to further resolve these differences, pointing to future research that may

still reveal more strain-specific effects of *Turicibacter* bacteria.

Further research into *bsh* gene regulation in the *Turicibacter* genus, the relationship

between bile and *Turicibacter* colonization and transmission behaviors, as well as the native

functionality of the putative BSH and 7α-HSDH homologs we tested, will further explain how

these bacteria wield their bile modifications in the intestine. In addition, exploring the

discrepencies we observed between bacterially-induced serum and cecal bile profiles could

describe other complexities in these effects, such as differential host reabsorption of modified

bile acids. Several of the *in vitro* bile modulations, such as increases in unconjugated bile acids,

were consistent between our *in vitro* characterizations and *Turicibacter* colonization, but

concurrent *in vivo* increases in expression of bile synthesis genes and levels of glyco-conjugated

bile acids seen during *Turicibacter* colonization and expression of individual *Turicibacter* BSH

indicate the possibility of more complex interactions between host bile production and

Turicibacter activity. The fact that the specificity and activity of the individual BSH homologs

differed when expressed in different bacterial backgrounds indicates that other unknown cellular

or environmental factors influence the way individual BSH act *in vivo*. This may also include

mechanisms that modify the functionality of the other putative BSH homologs we identified

from *Turicibacter* that did not deconjugate the specific conjugated bile acids used in our

experiments. These findings may influence the ways that bile-modifying genes can be employed

to shape host lipid profiles through microbiota engineering, positioning certain microbial species

as more appropriate vectors to impart specific host effects. It will also be informative to

determine what other activities performed by *Turicibacter* lead to WAT gain in colonized

animals, which contrasted findings from colonizing mice with specific *bsh*-overexpressing

strains of *B. thetaiotaomicron*. Given this finding, it is likely that *Turicibacter* also influences

host lipids through other mechanisms in addition to the bile transformations we characterized.

Our work also connects specific *Turicibacter* members and BSH activity with specific

host outcomes. Though some host responses such as broadly decreased triglycerides were

consistent across our BSH-recipient mice, the exact lipid and cholesterol responses differed,

indicating that the type of deconjugations might have differing connections with host physiology.

Further work will continue strengthening the exciting prospect of utilizing *Turicibacter* and/or its

bile modifications to intentionally alter host lipid biology to improve host metabolic and lipid-

associated health[66,67], as has been proposed with other bacteria[54,68]. Beyond lipid biology,

Turicibacter abundance has been positively correlated with diseases such as Parkinson's

disease[69] and depression[70], and selective serotonin reuptake inhibitors (SSRIs) have been found

to negatively affect *Turicibacter* growth and colonization[71], potentially because they inhibit

activity of its unique serotonin transporter[38]. SSRI use is frequently associated with metabolic

side effects like weight gain[72,73], and our findings suggests a hypothesis that connects SSRI use

and these side effects: SSRI use could diminish gut colonization of bacteria like *Turicibacter*,

thus unintentionally altering their impact on host physiology. Future work may develop

strategies to reduce interactions between SSRIs and activity of the microbiota, and minimizing

the side effects of these drugs and improving host outcomes. In all, these associations further

emphasize the importance of understanding mechanisms connecting members of the diverse

Turicibacter genus to host physiology.

Methods

Mouse husbandry

Adult (6-8 week old) germ-free Swiss Webster mice were used for all animal

experiments according to UCLA Institutional Animal Care and Use Committee-approved

protocols. Mice were reared in flexible gnotobiotic isolators on a 12h:12h light dark schedule on

standard chow (Labdiet 5K52, 22.1%: 16.6%: 61.3% protein: fat: carbohydrate by calories), then

were exited to autoclaved filter top cages with autoclaved chow (Labdiet 5010, 28.7%: 13.1%:

58.2% protein: fat: carbohydrate by calories) and water. After one day of cage acclimation, the

noted *Turicibacter* or *Bacteroides thetaiotaomicron* strain was grown in YCFA medium (see

below) overnight, pelleted by centrifugation, and resuspended in 1X PBS. Mice were colonized

by a 200μL gavage containing ~10^6 colony-forming units (CFU) of *Turicibacter* or ~10^8 CFU of

B. thetaiotaomicron. Alternatively, mice were gavaged with the same volume of PBS alone

(referred to as germ-free [GF]) or PBS-suspended fecal slurry from a specific pathogen-free

adult mouse (referred to as conventionalized [CONV]). Colonization was quantified using strain-

specific TuriSERT primers (**Supplementary Table 5**) and quantitative PCR (qPCR) from

weight-normalized contents from the distal small intestine and proximal colon after addition of

Low Abundance Microbiota Standard (Zymo) and extraction using the Zymo DNA Mini kit

(Zymo), standardized to a standard curve of CFU derived from plating of serial dilutions from

cultures of that strain.

Bacterial culturing

Turicibacter isolates and *Bacteroides thetaiotaomicron* strains (**Supplementary Table 1**)

were cultured in a flexible vinyl chamber (Coy) in an anaerobic 85%/10%/5% nitrogen/carbon

dioxide/hydrogen mixture (Airgas). *Turicibacter* was grown on Schaedler's agar (BD

Biosciences) or modified YCFA[74] (pH 7.4, per liter: 100mM MOPS, 10g casitone, 2.5g yeast

extract, 2g glucose, 2g maltose monohydrate, 2g cellobiose, 44mg $MgSO_4$, 68mg $CaCl_2$, 0.9g

NaCl, 10mg hemin, 0.45g K_2HPO_4, 0.45g KH_2PO_4, 4g $NaHCO_3$, 1g cysteine, 1mg resazurin,

1.9mL glacial acetic acid, 0.7mL propionic acid, 90μL isobutyric acid, 100uL isovaleric acid,

100uL valeric acid, 10mL ATCC vitamin mixture, 0.2% Tween-80) at 37°C. Cells were normally

grown without shaking, but when appropriate, *Turicibacter* cultures were anaerobically

transferred to sealed Hungate tubes or 1.7mL microcentrifuge tubes and shaken at 225 RPM at

37°C.

For *B. thetaiotaomicron* growth curves, overnight cultures were grown anaerobically in

BHI-S for ~48 hours at 37°C to ensure culture saturation, then were subcultured 1:50 for 6 hours

at 37°C (final OD_{600}=0.41-0.51). Subcultures were all then diluted to OD_{600}=0.1, and then six

replicates were further diluted 1:10 in 100μL BHI-S in a 96 well plate. Plates were anaerobically

sealed with parafilm and incubated at 37°C. OD_{600} readings were taken every 15 minutes in a

Biotek Synergy H1 microplate reader (Agilent).

Escherichia coli C41-pLys (Lucigen) was used for characterizing putative bile

modification genes, which were expressed off the pET21+ plasmid. *E. coli* was grown

aerobically shaking at 37°C C in Luria Broth (LB, 1% NaCl, 1% tryptone, 0.5% yeast extract)

supplemented with 100μg mL^{-1} ampicillin. Expression of genes was induced by addition of

100μM IPTG.

Bacterial isolation and identification

The frozen stool sample was thawed on ice and diluted 1:10 with PRAS anaerobic

dilution blank medium (Anaerobe Systems). 100μL of the diluted stool was further diluted to

1:1000 with modified YCFA media containing 0.05% bovine bile, 0.2% Tween-80, and 50mM

resorufin and loaded on Prospector® system arrays (Isolation Bio, San Carlos, CA, USA)

following manufacturer's instructions. The fluorescent green signal of the arrays at time 0 was

read on the Prospector® instrument in a Coy anaerobic chamber and the arrays incubated at 37^{o}C

in an Anaerobic Systems AS-580 anaerobic chamber (Anaerobe Systems). At 17 and 41 hours of

incubation the arrays were scanned again and the decrease in green fluorescence from time 0 was

used as an indicator for bacterial growth in the array nanowells. Bacteria from the array were

transferred to 96-well transfer plates containing 200μL per well of modified YCFA media,

without the addition of 50mM resorufin. The transfer plates were sealed with a gas permeable

film and incubated at 37^{o}C in a Mitsubishi AnaeroPack jar with a gas-generating sachet (Remel)

for seven days. After incubation, the contents of 538 wells from the transfer plates with visible

turbidity were consolidated into secondary 96-well plates, preserved with reduced glycerol, and

stored at -80^{o}C until needed. Unless stated otherwise, all stool and isolate manipulations were

conducted anaerobically with a 5% CO_2/5% H_2/90% N_2 atmosphere.

Genomic DNA was extracted in a 96-well format from the consolidated Prospector®

culture plates using the Extract All Kit (Applied Biosystems). 20μL of culture was combined

with 20μL of Lysis Solution and incubated for 10 minutes at 95^{o}C, followed by three minutes at

room temperature. The DNA was stabilized with the addition of 20μL of DNA Stabilizing

Solution and the resulting DNA lysate stored at -20^{o}C until needed.

qPCR screening of novel Turicibacter *isolates*

Genomic DNA from 538 isolates was screened for *Turicibacter* 16S and the *Turicibacter*
TuriSERT[38] gene using a multiplexed primer set (**Supplementary Table 5**). Each 25µL qPCR
reaction mixture had 1µL Extract All lysate, 10µL SYBR Power master mix (Applied
Biosystems), 0.5µL of each of the 10µM primers, and 12µL molecular grade water. The reactions
were run in a QuantStudio 6 Flex (Thermo Fisher) with a 95°C hold followed by 40 cycles of
95°C for 15s, 50°C for 30s, 72°C for 30s. *Turicibacter sanguinis* MOL361 gDNA and water were
used as positive and negative controls, respectively.

Molecular cloning

Turicibacter genes were amplified from template culture lysates with Phusion or Q5
DNA polymerase (NEB) and primers designed to amplify denoted *Turicibacter* genes. pET21- or
pWW3837[48]-derived expression plasmids were assembled using Gibson assembly (see
Supplementary Table 5 for oligos) for expression in *E. coli* or *Bacteroides thetaiotaomicron*,
respectively. Cloned constructs were confirmed through Sanger sequencing prior to functional
characterization. pWW3837-derived constructs were cloned into *B. thetaiotaomicron* VPI-5482
as previously described[48,75]. *bsh*-expressing *B. thetaiotaomicron* was compared to *B.*
thetaiotaomicron containing the original pWW3837 construct (referred to as wild-type *B.*
thetaiotaomicron).

Genome assemblies

Each strain was streaked on Schaedler agar plates and incubated anaerobically, then an
individual colony from each isolate was picked into YCFA medium and grown overnight at
37°C. DNA was extracted using the Zymo DNA mini kit (Zymo), with bead beating used to lyse

cells. Purified genomic DNA was sequenced by MiGS (migscenter.org), and 151bp paired-end

sequences were assembled using CLC Genomics Workbench (Qiagen). Genome assemblies have

been deposited at NCBI at BioProject PRJNA846348.

Whole genome and gene comparisons

anvi'o[76] was used to profile and visualize the different *Turicibacter* strain DNA

sequences to locate putative bile salt hydrolase and 7α-HSDH homologs in contig groups,

generate variability profiles, and measure gene coverage and detection statistics. Average

nucleotide identity (ANI) was calculated using OrthoANIu[77] (available

https://www.ezbiocloud.net/tools/ani).

Sequences comparisons between 16S rRNA and *bsh* genes/BSH amino acid sequences

were performed in CLC Genomics Workbench (Qiagen). 7α-HSDH sequence comparisons were

performed using tblastn[78] using the translated amino acid sequence from *Clostridium absonum*[44].

Assessment of bile transformations

In vitro characterization of bile transformations by engineered *E. coli* or *B.*

thetaiotaomicron strains were performed by growing cells in respective media conditions

described above supplemented with 0.5mM (total combined concentration) of the noted bile

species. Cells were grown to stationary phase (shaking at 37°C), then frozen at -80°C until

further processing. Cells were then thawed, pelleted (5 minutes at 16,000 x g), and the

supernatant was removed to a new microcentrifuge tube. Three volumes of methanol was added,

then the mixture was vigorously mixed for 30-60 seconds and incubated (room temperature, 15

minutes). Mixtures were centrifuged (5 minutes, 16,000 x g), the supernatant removed to a clean

microcentrifuge tube and dried in a vacuum concentrator. The dried residue was treated with

methanol/water/formic acid (50/50/0.1, all by volume) then vigorously mixed and centrifuged as

described above. Supernatants were transferred to polypropylene HPLC vials, capped, and

maintained at $4^{o}C$ while aliquots (typically 5µL) were injected onto a reversed phase HPLC

column (Cadenza CD-C18, 3.0µm, 250 x 2mm, Imtakt) equilibrated in solution A (water/formic

acid, 100/0.1, vol./vol.) and eluted ($0.2mL \text{ minute}^{-1}$) with an increasing concentration of solution

B (acetonitrile/formic acid, 100/0.1, vol./vol.); minute/% B: 0/30, 45/70, 48/100, 50/30, 67/30).

The effluent from the column was passed through an electrospray ion source (capillary voltage

42V, capillary temperature $275^{o}C$, sheath gas flow $15L \text{ min}^{-1}$, spray voltage 5kV, and -15kV

conversion dynode with -1.2kV multipliers) connected to a linear ion trap mass spectrometer

(Thermo LTQ) scanning from m/z 95-1000 in the positive ion mode. Spectra were recorded and

analyzed with instrument-manufacturer supplied software. Confirmation of proposed elemental

compositions was achieved using the same chromatography and ion source configuration with

the spectra recorded by scanning on an orbitrap mass spectrometer (Thermo LTQ XL).

For bile species quantification, an internal spike-in standard of 100mM chenodeoxycholic

acid-D4 (CDCA-D4, Sigma) was added to initial culture supernatants as a normalization

reference. Area under the curve from reconstructed ion chromatograms was used to quantify the

abundances of each species.

Serum metabolite analysis

Mice were euthanized with isoflurane and whole blood was collected via cardiac

puncture. Blood was allowed to clot in SST Vacutainer tubes (BD) on ice, then centrifuged ($4^{o}C$,

1 minute, 1500 x g). The supernatant was removed and snap frozen in liquid nitrogen. Serum

metabolites were analyzed using global metabolomics platform by Metabolon (Morrisville, NC,

USA). Unless otherwise noted, values presented are in arbitrary units (a.u.) for that particular

metabolite, determined by the log-transform of the volume-corrected quantification.

Circulating lipid analysis

Mice were fasted for 4-6 hours, then euthanized as described above. Blood was collected

via cardiac puncture and deposited into anticoagulatory K_2EDTA Vacutainer tubes (BD) on ice.

Blood was centrifuged (4^oC, 15 minutes, 2000 x g), then plasma was collected from supernatant

and snap frozen in liquid nitrogen. Shotgun lipidomics was performed by the UCLA Lipidomics

Core (Los Angeles, CA, USA) with the following protocol. Thawed plasma was pipetted into

glass tubes, a mixture of 70 internal standard lipids (Sciex and Avanti) was added, and lipids

were extracted using a modified Bligh and Dyer extraction[79]. The pooled organic layers from

two extractions were dried in a vacuum concentrator and resuspended in 50/50 (vol./vol.)

methanol/dichloromethane plus 10mM ammonium acetate. After transfer to robovials, samples

were analyzed with a Sciex 5500 with DMS Device (Lipidyzer Platform) using a targeted

acquisition list of 1450 lipid species. The Lipidyzer Differential Mobility device was tuned using

the EquiSPLASH LIPIDOMIX standard mixture (Avanti). Data was analyzed using an in-house

platform using previously described parameters[80], and quantitative values were normalized to

input volume. Statistical significance identification for species to include in heatmap was

performed with uncorrected two-tailed Welch's t-test comparison to Bt-WT (p-value cutoff

<0.05). Both raw and corrected p-values are included in Supplementary Table 4.

Total cecal bile analysis

Mice were colonized as described above and contents from the blind end of the cecum

were snap frozen upon sacrifice. Samples were thawed on ice and sent to Metabolon for analysis

with total bile analysis pipeline. Values shown at "0" were non-quantified and below limit of

detection.

Adipocyte area calculation

After sacrifice, mice epidydimal or gonadal white adipose tissue (e/g WAT) pads were

weighed and placed into 4% paraformaldehyde in 1X PBS for 48 hours at 4^{o}C. Fat pads were

washed twice in 70% ethanol, then submitted to the UCLA Translational Pathology Core

Laboratory (Los Angeles, CA, USA) for paraffin embedding, sectioning, and H&E staining. Ten

adipocyte images from each animal (five from each fat pad) were visualized with a 20X

objective on an EVOS microscope (Thermo). Adipocyte area for all cells contained entirely

within the field of view was automatically measured using the Fiji[81] Adiposoft[82] plug-in (version

1.1.16).

qRT-PCR measurement of liver transcripts

Gnotobiotic mice were colonized with single bacterial strains as described above, and

upon sacrifice, the median lobe of the liver was dissected and either directly snap frozen in liquid

nitrogen (all *Turicibacter* colonized animals) or placed in Trizol, bead beat for one minute, then

frozen in liquid nitrogen (all *B. thetaiotaomicron* colonized animals). All livers were then

transferred to -80^{o}C until further processing. Directly snap frozen livers were thawed overnight

at -20^{o}C in RNALater-ICE (ThermoFisher), then bead beat in Trizol for one minute, after which

all samples were processed in the same manner. RNA was extracted from thawed Trizol samples

using the Direct-Zol RNA Miniprep Kit (Zymo), then cDNA was generated using the qScript

cDNA Synthesis Kit (Quantabio). qPCR was performed using the PowerUp SYBR Green Master

Mix (ThermoFisher) on a QuantStudio5 Real-Time PCR System (ThermoFisher) (primers[83,84]

available in **Supplementary Table 5**) (cycling conditions: 50^{o}C for 2 min, 95^{o}C for 2 min, 50

cycles of 95^{o}C for 15 sec, 55^{o}C for 15 sec, 72^{o}C for 1 min, followed by melt curve). Fold

changes in comparison with sex-matched controls (GF for *Turicibacter* colonizations, *Bt*-WT for

B. thetaiotaomicron colonizations) were calculated using the ΔΔCt method with auto-

thresholded Ct values with *ppia* as the housekeeping gene.

Statistical analysis

585   Statistical calculations were performed in in Prism 9.3.1 (Graphpad) or Microsoft Excel 14.7.1. When appropriate, outliers were detected using the ROUT method with Q=1% threshold and removed from analysis. Dotted lines denote statistical comparison between groups directly below line, solid lines denote pairwise comparison. Unless otherwise noted, all tests were two-sided, *** = p<0.0005, ** = p<0.005, * = p<0.05, written p-value = 0.05 < 0.15. Heatmaps were created using the *pheatmap*[85] package in R (version 3.6.3)[86].

Data availability

594 Metabolomics and lipidomics datasets are attached to this manuscript as supplementary tables. Raw sequencing reads and genome assemblies for *Turicibacter* strains are deposited at NCBI under BioProject ID PRJNA846348.

REFERENCES

1. Schoeler, M. & Caesar, R. Dietary lipids, gut microbiota and lipid metabolism. *Rev. Endocr. Metab. Disord.* **20**, 461–472 (2019).

2. Wahlström, A., Sayin, S. I., Marschall, H.-U. & Bäckhed, F. Intestinal Crosstalk between Bile Acids and Microbiota and Its Impact on Host Metabolism. *Cell Metab.* **24**, 41–50 (2016).

3. Sharon, G., Sampson, T. R., Geschwind, D. H. & Mazmanian, S. K. The Central Nervous System and the Gut Microbiome. *Cell* (2016). doi:10.1016/j.cell.2016.10.027

4. Fung, T. C., Olson, C. A. & Hsiao, E. Y. Interactions between the microbiota, immune and nervous systems in health and disease. *Nature Neuroscience* (2017). doi:10.1038/nn.4476

5. Rabot, S. *et al.* Germ-free C57BL/6J mice are resistant to high-fat-diet-induced insulin resistance and have altered cholesterol metabolism. *FASEB J.* **24**, 4948–4959 (2010).

6. Wikoff, W. R. *et al.* Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc. Natl. Acad. Sci.* **106**, 3698–3703 (2009).

7. Yano, J. M. *et al.* Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell* (2015). doi:10.1016/j.cell.2015.02.047

8. Ivanov, I. I. *et al.* Induction of Intestinal Th17 Cells by Segmented Filamentous Bacteria. *Cell* **139**, 485–498 (2009).

9. Sokol, H. *et al.* Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc. Natl. Acad. Sci.* **105**, 16731–16736 (2008).

10. Storelli, G. *et al.* Drosophila Perpetuates Nutritional Mutualism by Promoting the Fitness of Its Intestinal Symbiont Lactobacillus plantarum. *Cell Metab.* **27**, 362-377.e8 (2018).

11. Kim, C. Y. *et al.* Human reference gut microbiome catalog including newly assembled

genomes from under-represented Asian metagenomes. *Genome Med.* **13**, 134 (2021).

12.   Huang, S. *et al.* The day-to-day stability of the ruminal and fecal microbiota in lactating

dairy cows. *Microbiologyopen* **9**, (2020).

13.   Maki, J. J. & Looft, T. Turicibacter bilis sp. nov., a novel bacterium isolated from the

chicken eggshell and swine ileum. *Int. J. Syst. Evol. Microbiol.* **72**, (2022).

14.   Feng, Y. *et al.* Metagenome-assembled genomes and gene catalog from the chicken gut

microbiome aid in deciphering antibiotic resistomes. *Commun. Biol.* **4**, 1305 (2021).

15.   Mo, J. *et al.* Structural and quantitative alterations of gut microbiota in experimental small

bowel obstruction. *PLoS One* **16**, e0255651 (2021).

16.   Browne, H. P. *et al.* Culturing of 'unculturable' human microbiota reveals novel taxa and

extensive sporulation. *Nature* (2016). doi:10.1038/nature17645

17.   Martínez, I. *et al.* The Gut Microbiota of Rural Papua New Guineans: Composition,

Diversity Patterns, and Ecological Processes. *Cell Rep.* **11**, 527–538 (2015).

18.   Goodrich, J. K. *et al.* Genetic Determinants of the Gut Microbiome in UK Twins. *Cell*

Host Microbe **19**, 731–743 (2016).

19.   Benson, A. K. *et al.* Individuality in gut microbiota composition is a complex polygenic

trait shaped by multiple environmental and host genetic factors. *Proc. Natl. Acad. Sci.*

107, 18933–18938 (2010).

20.   Org, E. *et al.* Genetic and environmental control of host-gut microbiota interactions.

Genome Res. (2015). doi:10.1101/gr.194118.115

21.   Kemis, J. H. *et al.* Genetic determinants of gut microbiota composition and bile acid

profiles in mice. *PLoS Genet.* (2019). doi:10.1371/journal.pgen.1008073

22.   Santiago-Rodriguez, T. M. *et al.* Gut Microbiome of an 11th Century A.D. Pre-Columbian

Andean Mummy. *PLoS One* **10**, e0138135 (2015).

23.   Wibowo, M. C. *et al.* Reconstruction of ancient microbial genomes from the human gut.

Nature **594**, 234–239 (2021).

24.   Yatsunenko, T. *et al.* Human gut microbiome viewed across age and geography. *Nature*

486, 222–227 (2012).

25.   Turnbaugh, P. J. *et al.* An obesity-associated gut microbiome with increased capacity for

energy harvest. *Nature* (2006). doi:10.1038/nature05414

26.   Thaiss, C. A. *et al.* Persistent microbiome alterations modulate the rate of post-dieting

weight regain. *Nature* (2016). doi:10.1038/nature20796

27.   Chadaideh, K. S. & Carmody, R. N. Host-microbial interactions in the metabolism of

different dietary fats. *Cell Metab.* **33**, 857–872 (2021).

28.   Dhakal, S., McCormack, L. & Dey, M. Association of the Gut Microbiota with Weight-

Loss Response within a Retail Weight-Management Program. *Microorganisms* **8**, 1246

(2020).

29.   Liu, W. *et al.* Diet- and Genetically-induced Obesity Produces Alterations in the

Microbiome, Inflammation and Wnt Pathway in the Intestine of Apc +/1638N Mice:

Comparisons and Contrasts. *J. Cancer* **7**, 1780–1790 (2016).

30.   Li, T.-T. *et al.* Polyunsaturated fatty acids from microalgae Spirulina platensis modulates

lipid metabolism disorders and gut microbiota in high-fat diet rats. *Food Chem. Toxicol.*

131, 110558 (2019).

31.   Velázquez, K. T. *et al.* Prolonged high-fat-diet feeding promotes non-alcoholic fatty liver

disease and alters gut microbiota in mice. *World J. Hepatol.* **11**, 619–637 (2019).

32.   Jiao, N. *et al.* Gut microbiome may contribute to insulin resistance and systemic

inflammation in obese rodents: a meta-analysis. *Physiol. Genomics* **50**, 244–254 (2018).

33.   Petersen, C. *et al.* T cell–mediated regulation of the microbiota protects against obesity.

Science (80-.). **365**, (2019).

34.   Golloso-Gubat, M. J. *et al.* Gut Microbiota and Dietary Intake of Normal-Weight and

673  Overweight Filipino Children. *Microorganisms* **8**, 1015 (2020).

674 35. Chen, X. *et al.* Alteration of the gut microbiota associated with childhood obesity by 16S

675  rRNA gene sequencing. *PeerJ* **8**, e8317 (2020).

676 36. Bosshard, P. P., Zbinden, R. & Altwegg, M. Turicibacter sanguinis gen. nov., sp. nov., a

677  novel anaerobic, Gram-positive bacterium. *Int. J. Syst. Evol. Microbiol.* (2002).

678  doi:10.1099/ijs.0.02056-0

679 37. Maki, J. J., Nielsen, D. W. & Looft, T. Complete Genome Sequence and Annotation for

680  Turicibacter sanguinis MOL361 T (DSM 14220). *Microbiol. Resour. Announc.* **9**, (2020).

681 38. Fung, T. C. *et al.* Intestinal serotonin and fluoxetine exposure modulate bacterial

682  colonization in the gut. *Nature Microbiology* (2019). doi:10.1038/s41564-019-0540-4

683 39. Auchtung, T. A. *et al.* Complete genome sequence of Turicibacter sp. strain H121,

684  isolated from the feces of a contaminated germ-free mouse. *Genome Announc.* (2016).

685  doi:10.1128/genomeA.00114-16

686 40. Russell, D. W. The Enzymes, Regulation, and Genetics of Bile Acid Synthesis. *Annu. Rev.*

687  *Biochem.* **72**, 137–174 (2003).

688 41. Guo, C., Chen, W.-D. & Wang, Y.-D. TGR5, Not Only a Metabolic Regulator. *Front.*

689  *Physiol.* **7**, (2016).

690 42. Ridlon, J. M., Kang, D.-J. & Hylemon, P. B. Bile salt biotransformations by human

691  intestinal bacteria. *J. Lipid Res.* **47**, 241–259 (2006).

692 43. Makishima, M. *et al.* Vitamin D Receptor As an Intestinal Bile Acid Sensor. *Science (80-.*

693  *).* **296**, 1313–1316 (2002).

694 44. Lou, D. *et al.* The three-dimensional structure of Clostridium absonum 7α-hydroxysteroid

695  dehydrogenase: new insights into the conserved arginines for NADP(H) recognition. *Sci.*

696  *Rep.* **6**, 22885 (2016).

697 45. Klaver, F. A. & van der Meer, R. The assumed assimilation of cholesterol by Lactobacilli

and Bifidobacterium bifidum is due to their bile salt-deconjugating activity. *Appl.*

Environ. Microbiol. **59**, 1120–1124 (1993).

46.   Yao, L. *et al.* A selective gut bacterial bile salt hydrolase alters host metabolism. *Elife*

(2018). doi:10.7554/elife.37182

47.   Joyce, S. A. *et al.* Regulation of host weight gain and lipid metabolism by bacterial bile

acid modification in the gut. *Proc. Natl. Acad. Sci.* **111**, 7421–7426 (2014).

48.   Whitaker, W. R., Shepherd, E. S. & Sonnenburg, J. L. Tunable Expression Tools Enable

Single-Cell Strain Distinction in the Gut Microbiome. *Cell* **169**, 538-546.e12 (2017).

49.   Hylemon, P. B. & Sherrod, J. A. Multiple forms of 7-alpha-hydroxysteroid dehydrogenase

in selected strains of Bacteroides fragilis. *J. Bacteriol.* **122**, 418–424 (1975).

50.   Sherrod, J. A. & Hylemon, P. B. Partial purification and characterization of NAD-

dependent 7α-hydroxysteroid dehydrogenase from Bacteroides thetaiotaomicron. *Biochim.*

Biophys. Acta - Lipids Lipid Metab. **486**, 351–358 (1977).

51.   Marion, S. *et al.* Biogeography of microbial bile acid transformations along the murine

gut. *J. Lipid Res.* **61**, 1450–1463 (2020).

52.   Hofmann, A. F., Hagey, L. R. & Krasowski, M. D. Bile salts of vertebrates: structural

variation and possible evolutionary significance. *J. Lipid Res.* **51**, 226–246 (2010).

53.   Jones, B. V., Begley, M., Hill, C., Gahan, C. G. M. & Marchesi, J. R. Functional and

comparative metagenomic analysis of bile salt hydrolase activity in the human gut

microbiome. *Proc. Natl. Acad. Sci.* **105**, 13580–13585 (2008).

54.   Begley, M., Hill, C. & Gahan, C. G. M. Bile Salt Hydrolase Activity in Probiotics. *Appl.*

Environ. Microbiol. **72**, 1729–1738 (2006).

55.   Prete, R. *et al.* Beneficial bile acid metabolism from Lactobacillus plantarum of food

origin. *Sci. Rep.* **10**, 1165 (2020).

56.   Tanaka, H., Hashiba, H., Kok, J. & Mierau, I. Bile Salt Hydrolase of Bifidobacterium

longum —Biochemical and Genetic Characterization. *Appl. Environ. Microbiol.* **66**,

2502–2512 (2000).

57. Kim, G.-B., Miyamoto, C. M., Meighen, E. A. & Lee, B. H. Cloning and Characterization

of the Bile Salt Hydrolase Genes (bsh) from Bifidobacterium bifidum Strains. *Appl.*

Environ. Microbiol. **70**, 5603–5612 (2004).

58. Ma, D. *et al.* Ketogenic diet enhances neurovascular function with altered gut microbiome

in young healthy mice. *Sci. Rep.* **8**, 6670 (2018).

59. Gao, X. *et al.* Correlations between α-Linolenic Acid-Improved Multitissue Homeostasis

and Gut Microbiota in Mice Fed a High-Fat Diet. *mSystems* **5**, (2020).

60. Everard, A. *et al.* Microbiome of prebiotic-treated mice reveals novel targets involved in

host response during obesity. *ISME J.* **8**, 2116–2130 (2014).

61. Jung, M.-J. *et al.* Chronic Repression of mTOR Complex 2 Induces Changes in the Gut

Microbiota of Diet-induced Obese Mice. *Sci. Rep.* **6**, 30887 (2016).

62. Huang, J. *et al.* Supplementation of Bacillus sp. DU-106 reduces hypercholesterolemia

and ameliorates gut dysbiosis in high-fat diet rats. *Appl. Microbiol. Biotechnol.* **105**, 287–

299 (2021).

63. Golloso-Gubat, M. J. *et al.* Gut microbiota and dietary intake of normal-weight and

overweight filipino children. *Microorganisms* (2020).

doi:10.3390/microorganisms8071015

64. Caesar, R., Tremaroli, V., Kovatcheva-Datchary, P., Cani, P. D. & Bäckhed, F. Crosstalk

between Gut Microbiota and Dietary Lipids Aggravates WAT Inflammation through TLR

Signaling. *Cell Metab.* **22**, 658–668 (2015).

65. Martinez-Guryn, K. *et al.* Small Intestine Microbiota Regulate Host Digestive and

Absorptive Adaptive Responses to Dietary Lipids. *Cell Host Microbe* **23**, 458-469.e5

(2018).

66.   Jones, M. L., Martoni, C. J. & Prakash, S. Cholesterol lowering and inhibition of sterol

749   absorption by Lactobacillus reuteri NCIMB 30242: a randomized controlled trial. *Eur. J.*

750   *Clin. Nutr.* **66**, 1234–1241 (2012).

751   67.   Song, *Z. et al.* Taxonomic profiling and populational patterns of bacterial bile salt

752   hydrolase (BSH) genes based on worldwide human gut microbiome. *Microbiome* **7**, 9

753   (2019).

754   68.   Russell, B. J. *et al.* Intestinal Transgene Delivery with Native E.

755   coli Chassis Allows Persistent Physiological Changes. *bioRxiv*

756   2021.11.11.468006 (2021). doi:10.1101/2021.11.11.468006

757   69.   Jin, M. *et al.* Analysis of the Gut Microflora in Patients With Parkinson's Disease. *Front.*

758   *Neurosci.* **13**, 1184 (2019).

759   70.   Barandouzi, Z. A., Starkweather, A. R., Henderson, W. A., Gyamfi, A. & Cong, X. S.

760   Altered Composition of Gut Microbiota in Depression: A Systematic Review. *Front.*

761   *Psychiatry* **11**, (2020).

762   71.   Jackson, M. A. *et al.* Gut microbiota associations with common diseases and prescription

763   medications in a population-based cohort. *Nat. Commun.* (2018). doi:10.1038/s41467-

764   018-05184-7

765   72.   Ferguson, J. M. SSRI Antidepressant Medications. *Prim. Care Companion J. Clin.*

766   *Psychiatry* **03**, 22–27 (2001).

767   73.   Cascade, E., Kalali, A. H. & Kennedy, S. H. Real-world data on SSRI antidepressant side

768   effects. *Psychiatry* (2009).

769   74.   Duncan, S. H. Growth requirements and fermentation products of Fusobacterium

770   prausnitzii, and a proposal to reclassify it as Faecalibacterium prausnitzii gen. nov., comb.

771   nov. *Int. J. Syst. Evol. Microbiol.* **52**, 2141–2146 (2002).

75.   Lynch, J. B. & Sonnenburg, J. L. Prioritization of a plant polysaccharide over a mucus

carbohydrate is enforced by a Bacteroides hybrid two-component system. *Mol. Microbiol.*

(2012). doi:10.1111/j.1365-2958.2012.08123.x

76.   Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat.*

Microbiol. **6**, 3–6 (2021).

77.   Yoon, S.-H., Ha, S., Lim, J., Kwon, S. & Chun, J. A large-scale evaluation of algorithms

to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* **110**, 1281–1286

(2017).

78.   Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: A new generation of protein

database search programs. *Nucleic Acids Research* (1997). doi:10.1093/nar/25.17.3389

79.   Hsieh, W.-Y., Williams, K. J., Su, B. & Bensinger, S. J. Profiling of mouse macrophage

lipidome using direct infusion shotgun mass spectrometry. *STAR Protoc.* **2**, 100235

(2021).

80.   Su, B. *et al.* A DMS Shotgun Lipidomics Workflow Application to Facilitate High-

Throughput, Comprehensive Lipidomics. *J. Am. Soc. Mass Spectrom.* **32**, 2655–2663

(2021).

81.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

82.   Galarraga, M. *et al.* Adiposoft: automated software for the analysis of white adipose tissue

cellularity in histological sections. *J. Lipid Res.* **53**, 2791–2796 (2012).

83.   Sayin, S. I. *et al.* Gut Microbiota Regulates Bile Acid Metabolism by Reducing the Levels

of Tauro-beta-muricholic Acid, a Naturally Occurring FXR Antagonist. *Cell Metab.* **17**,

225–235 (2013).

84.   Takahashi, S. *et al.* Cyp2c70 is responsible for the species difference in bile acid

metabolism between mice and humans. *J. Lipid Res.* **57**, 2130–2137 (2016).

85.   Kolde, R. pheatmap: Pretty Heatmaps. (2019).

86.   Team, R. C. R: A language and environment for statistical computing. (2020).

87.   Šidák, Z. Rectangular Confidence Regions for the Means of Multivariate Normal

800   Distributions. *J. Am. Stat. Assoc.* **62**, 626–633 (1967).

Acknowledgments

Turicibacter isolates H121, TA25, T46 and T129 were a gift from Thomas Auchtung. 18F6 and

1E2 were a gift from Kenya Honda. pWW3837 and *B. thetaiotaomicron* VPI-5482 was a gift

from Fatima Enam and Justin Sonnenburg. We would like to thank Alexander Yoon, Gazmend

Elezi, and Lauren Seaman for technical assistance with the LC-MS operation and protocols, and

Nathan Hwangbo for statistical guidance. This work was supported by funding from the Packard

Foundation (E.Y.H.) and Ford Foundation (J.B.L.). E.Y.H. is a New York Stem Cell Foundation

– Robertson Investigator. This research was supported in part by The New York Stem Cell

Foundation.

Author Contributions

J. B. L., E. L. G, K. C., K. B. Y., J. P. and E. Y. H. conceptualized and planned bacterial and

mouse experiments. J. B. L. and K. F. F. designed and performed LC-MS experiments. J. B. L.,

T. J., A. A., and J. L. performed isolation and characterization of novel *Turicibacter* isolates. J.

B. L. and E. Y. H. wrote manuscript. All authors read and edited final manuscript.

Competing Interests

The authors declare no competing interests. Findings regarding the host effects of *Turicibacter*

reported in the manuscript are the subject of provisional patent application 63/288980, owned by

UCLA.

Availability of materials

Bacterial strains from this work may be obtained through reasonable request to the

corresponding author. Certain materials may be restricted by MTA requirements.

Fig. 1. Genomic comparisons reveal distinct subgroups of *Turicibacter*. a, Phylogenetic tree

comparing full-length 16S rRNA gene sequences from noted *Turicibacter* isolates. Circles

indicate human-derived isolates, triangles indicate mouse-derived isolates, and square indicates a

mouse-derived contaminating isolate. **b**, Association between guanine-cytosine % (GC%) and

calculated genome size in megabases (Mb) for shotgun-assembled genomes of *Turicibacter*

isolates from **a**. **c**, Full genome sequence comparisons across *Turicibacter* strains. Position of

predicted bile-modification gene homologs are noted outside of rings, with the color of the gene

name denoting the genome family that gene is found in. Each ring represents sequence blocks in

one genome. **d**, Average nucleotide identity (ANI) between noted *Turicibacter* genomes.

Number denotes ANI, white-green-maroon scale represents 100%-75% ANI scale.

Fig. 2. Colonization with different *Turicibacter* strains alters host lipids . a, Heatmap of

relative abundance of serum lipids from gnotobiotic mice monocolonized with noted

Turicibacter strains. Heatmap values (Z-score) represent mean abundance of each detected lipid

species from labeled lipid categories scaled across all the means of that individual lipid species.

Black (p<0.05) and grey (0.05<p<0.15) rectangles indicate statistically significant differences of

that metabolite between: left, GF and MOL361 monocolonized mice; middle, CONV and

MOL361 monocolonized mice; and right, between MOL361, 1E2, and H121 monocolonized

mice. **b**, Serum cholesterol concentrations of mice colonized by noted *Turicibacter* strains. **c**,

Sex and litter-matched relative epidydimal/gonadal white adipose tissue (e/g WAT) mass of mice

monocolonized with noted *Turicibacter* strains. Shapes indicate value for individual mouse,

dotted bar represents ANOVA statistic for groups below the line. Metabolite and cholesterol

analysis n=6-10. WAT analysis n=6-26, Mann-Whitney test for MOL361-GF and MOL361-

CONV comparisons, Kruskal-Wallis for intra-*Turicibacter* comparison, Šidák correction for

multiple comparisons[87]. *p<0.05.

Fig. 3. *Turicibacter* colonization alters circulating host bile species in a strain-specific

manner. Serum concentrations of **a-d**, primary unconjugated bile acids; **e-f**, secondary

unconjugated bile acids; or **g-l**, primary conjugated bile acids. Points indicate log-transformed

value for individual mouse with shapes and colors matching **Fig. 1**, error bars represent mean +/-

SEM. Kruskal-Wallis test across all noted colonizations with Dunn's multiple comparisons to

GF for **a-f**. Kruskal-Wallis test between noted *Turicibacter* strains and multiple comparisons to

H121 for **g-l**. Mann-Whitney test used to compare GF and MOL361 in g-l, and p-values are

noted above GF data points. n=6-10 for each group. Dotted bar represents ANOVA statistic for

groups below the line. *p<0.05, **p<0.005.

Fig. 4: *Turicibacter* isolates differ in their bile-modifying abilities. a, Schematic for types of

bile transformations found to be performed by *Turicibacter* isolates. **b**, inset: 16S rRNA gene-

based phylogenic tree from Fig. 1a. Liquid chromatograms of individual *Turicibacter* isolates

grown for 24 hours in media with sub-inhibitory concentrations of five bile acids: taurocholic

acid (TCA), cholic acid (CA), glycochenodeoxycholic acid (GCDCA), chenodeoxycholic acid

(CDCA), and deoxycholic acid (DCA). Shaded regions indicate expected retention time of each

bile species. **c**, Percent remaining (compared to cultures at time=0) of conjugated bile acids

(TCA, taurochenodeoxycholic acid [TCDCA], glycocholic acid [GCA], GCDCA) after 24

growth with noted *Turicibacter* isolate. Yellow=glycine-conjugated bile acids, orange=taurine-

conjugated bile acids. n=4 independent cultures. Values not shown were below 0.1% remaining.

Statistical analysis performed by one sample t-test, annotations of legend denotes strains with

significant difference from 100% remaining for each bile acid.

Fig. 5. *Turicibacter* **isolates differ in their genetic capacity to modify bile species. a**,
Phylogenetic tree of amino acid sequences for each predicted bile salt hydrolase (BSH) sequence
from *Turicibacter* strains, with observed bile species specificity noted in boxes. We did not

detect bile salt hydrolase activity in sequences without boxes, representing groups V-VIII. **b**,

Presence (+) or absence of sequence homologs with potential BSH activity in *Turicibacter*

strains. **c**, Liquid chromatograms of media after 24 hour cultures of *E. coli* expressing individual

predicted *bsh* genes from each sequence grouping and grown in with TCA and TCDCA. Control

is *E. coli* with same expression vector but expressing non-bile-modifying gene. **d**, same as **c**, but

with GCA and GCDCA instead of tauro-bile acids. **e**, Quantification of percent remaining

(compared to media controls) of conjugated bile acid (TCA, TCDCA, GCA, GCDCA) after 24h

growth with *E. coli* expressing the noted *Turicibacter bsh* gene. n=3 independent cultures,

annotations in legend indicate *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$ using one sample t-test

comparison with 100% remaining. BSH nomenclature indicates homolog group (*e.g.* III) and

isolate of origin (*e.g.* MOL361).

Fig. 6: *Turicibacter bsh* expression is sufficient to alter host lipidome and health-associated lipid markers. a, Percent remaining of noted bile acids after 24 hour growth with *Bacteroides thetaiotaomicron* expressing noted *bsh* genes. n=4 cultures per strain. **b**, Same as **a**, but with 48 hour growth with noted *B. thetaiotaomicron* strains. n=3 cultures per strain. For **a** and **b**, points represent individual comparison with media control, legend annotations denote strains with

statistical significance for each bile acid using one sample t-test comparison with 100%

remaining. **c**, Quantification of combined taurine-conjugated cecal bile acids (BA) from mice

colonized with *bsh*-expressing *B. thetaiotaomicron.* Statistical analysis was performed with

Kruskal-Wallis test with Dunn's multiple comparisons test. n=6 animals per colonization. **d**,

Heatmap of circulating lipid species significantly altered by expression of at least one

Turicibacter bsh in *B. thetaiotaomicron*. Heatmap Z-score values represent mean abundance of

each detected lipid species from labeled lipid categories scaled across all the means of that

individual lipid species. Colors on left correspond to lipid class, each column represents one

animal. **e-i**, Relative combined circulating concentrations of **e**, triglycerides (TG), **f**, cholesterol

esters (CE), **g**, diacylglycerides (DG), **h**, phosphotidylglycines (PG), or **i**, phosphotidylserines

(PS) of mice monocolonized with *bsh*-expressing *B. thetatiotaomicron.* **j**, Relative white adipose

tissue weight of mice monocolonized with *bsh*-expressing *B. thetatiotaomicron*. All values in **e-j**

normalized to sex-matched littermates, n=4-6 animals per colonization, statistical analysis

performed by Welch's ANOVA with Dunnett's multiple comparisons to GF, dotted bar in **c, e-j**

represents ANOVA statistic for groups below the line. $*p<0.05$, $**p<0.005$, $***p<0.0005$.

 SUPPLEMENTARY FIGURES

Supplementary Fig. 1. Strain-dependent variation in colonization and adipocyte size. a-e,

Representative images of adipose tissue from *Turicibacter* monocolonized mice. **f**, Sex- and

litter- matched relative adipocyte area of mice monocolonized with individual *Turicibacter*

strains. Each point represents mean of 10 images of adipocyte area per animal. **g, h**, Colony-

forming units (CFU) equivalents per gram of contents for distal small intestine (**g**) or proximal

colon (**h**). Each dot represents sample from one animal, per colonization n=10-15 for qPCR,

n=10-14 for adipocyte area calculation. Statistics for adipocyte area performed by Welch's

ANOVA with Dunnet's multiple comparisons, dotted bar represents ANOVA statistic for each

group versus the combined experimental mean.

Supplementary Fig. 2. *Turicibacter* colonization alters cecal bile acids. Cecal concentrations

of conjugated and unconjugated species of **a-c,** cholic acid, **d-e**, chenodeoxycholic acid, **f-g**,

alpha-muricholic acid, **h-i**, beta-muricholic acid, **j-k**, ursodeoxycholic acid, and the unconjugated

secondary bile acid **l**, lithocholic acid, in GF or *Turicibacter* monocolonized mice. Each point

represents one animal, n=6 animals per condition. Statistical comparisons between *Turicibacter*

930   isolates performed with Kruskal-Wallis test with Dunn's multiple comparisons, GF-MOL361

931   comparisons with Mann-Whitney test. Dotted line represents combined intra-*Turicibacter*

932   ANOVA p-value, value written above GF represents GF-MOL361 p-value, *p<0.05, **p<0.005.

Supplementary Fig. S3. Host effects of *Turicibacter* colonization varies with sex. a-c, Heatmap of mean relative **a)** bile species, **b)** sterols, or **c)** lipids. Column labels represent colonization state and sex, blue=male, red=female. n=3-6 per group, except female T129 (n=1).

Turicibacter strain	Presence (+) or absence (-) of peak							Deconjugation ability
	TCA	Dehydrogenated CA	CA	GCDCA	Dehydrogenated CDCA	CDCA	DCA	
MOL361	-	+	+	-	+	+	+	Tauro- and glyco-deconjugation; dehydrogenation
18F6	-	+	+	-	+	+	+	
GALT-E2	-	+	+	-	+	+	+	
T46	+	-	+	-	-	+	+	Glyco-deconjugation
H121	+	-	+	-	-	+	+	
T129	+	-	+	-	-	+	+	
GALT-G1	-	-	+	-	+	+	+	Tauro- and glyco-deconjugation; some dehydrogenation
1E2	-	-	+	+	-	+	+	Tauro- deconjugation
TA25	-	-	+	+	-	+	+	

Red = outliers from genomic group

938 **Supplementary Fig. 4. Summary table of bile transformations performed by *Turicibacter***

939 **isolates.** Table indicating presence/absence of noted bile species after growth in mixture of bile

940 acids described in **Fig. 4b**.

Supplementary Fig. 5. *Turicibacter* **strains MOL361 and 1E2 can deconjugate at least six**

taurine-conjugated bile acids. Chromatograms (left) and unconjugated/conjugated bile acid

ratios (right) of *Turicibacter* MOL361 or 1E2 grown for 24 hours in YCFA + 0.5mM of

individual taurine-conjugated bile acids: **a**, TCA; **b**, TCDCA; **c**, tauroursodeoxycholic acid

946   (TUDCA); **d**, taurolithocholic acid (TLCA); **e**, taurohyodeoxycholic acid (THDCA); **f**,

947   taurodeoxycholic acid (TDCA). Chromatograms are concatenated reconstructed chromatograms

948   for conjugated and unconjugated bile acid. Each trace and each point represents one biological

949   replicate, n=3 cultures. Statistics were performed by Mann-Whitney test, #p≤0.1 (Note: this p-

950   value is the minimum for this test and our experimental parameters).

Supplementary Fig. 6. Some strains of *Turicibacter* encode a functional 7alpha-HSDH gene.
a, Table depicting amino acid similarity and sequence coverage between closest predicted 7alpha-HSDH homolog in noted isolate genome and 7alpha-HSDH gene from *Clostridium absonum*. **b**, Chromatograms of cholic acid (CA) or cholic acid with 2 hydrogens removed (CA-2H) from *E. coli* cultures expressing the MOL361 HSDH homolog (7alpha-HSDH) or non-bile modifying gene sequence in the same plasmid (cell control). Chromatograms from independent triplicate cultures are shown, dotted boxes indicate each bile species. **c**, Quantification of CA/CA-2H ratios determined from reconstructed areas under the curve in **b**. Statistical comparison performed with Welch's t-test, n=3 cultures, bars indicate mean +/- SEM. **d**, Same as **b**, but with CDCA instead of CA. **e**, same as **b**, but with DCA instead of CA. **d** and **e** triplicate chromatograms are stacked on the same axis.

Bile salt hydrolase (*bsh*) gene	Presence (+) or absence (-) of peak				Deconjugation ability
	TCA	TCDCA	GCA	GCDCA	
I-MOL361	-	-	-	-	Tauro- and glyco-deconjugation
II-H121	+	-	+	-	-CDCA deconjugation
III-1E2	+	-	+	+	TCDCA deconjugation
IV-MOL361	-	-	+	+	Tauro- deconjugation

964 **Supplementary Fig. 7. Bile transformations performed by *B. thetaiotaomicron* expressing**

965 ***Turicibacter bsh* genes.** Table indicating presence/absence of noted bile species after growth in

966 mixture of bile acids described in **Fig. 6a, b**.

Supplementary Fig. 8. Expressing *bsh* genes from *Turicibacter* does not impart significant

***in vitro* growth defect of *B. thetaiotaomicron*.** OD_{600} readings of noted *B. thetaiotaomicron*

strains in BHI-S medium. Each point represents mean +/- SEM for 6 independent cultures.

Supplementary Fig. 9. *bsh* expression does not alter *B. thetaiotaomicron* colonization of the

murine gut. CFU/mL quantifications of the noted *B. thetaiotaomicron* strains in the **a,** distal

small intestine; **b,** cecum and; **c,** proximal colon of gnotobiotic mice. n=7-12/group, comparisons

with Welch's ANOVA test and Dunnet's multiple comparisons.

Supplementary Fig. 10. Engineered *B. thetaiotaomicron* transforms host bile acids *in vivo*.
Cecal abundance of unconjugated and conjugated species of primary bile acids **a-c,** cholic acid,
d-e, chenodeoxycholic acid, **f-g**, alpha-muricholic acid, **h-i**, beta-muricholic acid, **j-k**,
ursodeoxycholic acid, and the unconjugated secondary bile acid **l**, lithocholic acid. n=3 (3 males,
3 females) for each condition. Statistics performed with Kruskal-Wallis test with Dunn's

982 multiple comparisons. Dotted horizontal line represents total ANOVA statistic across all four

983 groups, $*p<0.05$, $**p<0.005$, $***p<0.0005$.

Supplementary Fig. 11. ***Turicibacter bsh*** **expression by** ***B. thetaiotaomicron*** **is sufficient to drive broad scale changes in circulating host lipids in male and female mice.** Related to **Fig. 6e-j**, circulating absolute concentrations of specific lipid categories in mice colonized by *B. thetaiotaomicron* expressing the noted *Turicibacter bsh* homologs. Each point represents one animal, red points indicate female, blue represent male. n=4-5 per colonization, error bars represent mean +/- SEM. Statistical comparison done with Welch's ANOVA with Dunnett's multiple comparisons, dotted bar represents ANOVA statistic for all four groups, *p<0.05.

males **females**

Supplementary Fig. 12. Sex differences in lipidomic response to *Turicibacter bsh* expression

in *B. thetaiotaomicron*. Similar to **Fig. 5b**, but animal lipidome analysis separated into **a)** males

and **b)** females. Note that lipid species presented were found to be significantly altered by

996   expression of at least one *bsh* in males and females combined (i.e. all lipid species shown across

997   the three analyses are the same). Each column represents one animal.

Supplementary Fig. 13. *T. sanguinis* **MOL361 colonization leads to increased triglycerides and diacylglycerides, consistent with adipose tissue increases.** Plasma levels of combined **a,** cholesterol esters, **b,** triglycerides, and **c,** diacylglycerides. **i:** relative to sex-matched GF littermates, **ii**: absolute quantification. n=4-5 (1-2 females, 3 males per condition). red=female, blue=male. Statistical comparisons Student's two tailed t-test with Welch's correction, *p<0.05.

1004

Supplementary Fig. 14. *Turicibacter* colonization and *bsh* expression induce similar gene

expression patterns in the liver. qRT-PCR analysis of liver transcript levels of **a,** Fxr; **b,**

Cyp7a1; and **c,** G6pase after colonization with the noted *Turicibacter* strains or *B.*

thetaiotaomicron expressing the noted *bsh* gene. Data are displayed as fold change relative to

appropriate control (GF for *Turicibacter* colonizations, *Bt*-WT for *bsh* colonizations). Each point

represent data from a single animal, statistics performed with Kruskal-Wallis test with Dunn's

multiple comparisons. Dotted horizontal line represents ANOVA statistic for that comparison of

groups below the line, *p<0.05.

1013

1014   **Supplementary Table 1. List of strains used in this work.**
1015

1016   **Supplementary Table 2. Serum metabolomics from GF, CONV, and *Turicibacter***
1017   **monocolonized mice.** Volume-adjusted log-transformed levels of listed serum metabolites from
1018   mice from different colonization states. Raw p-values generated from Mann-Whitney U test (GF-
1019   MOL361 and MOL361-CONV) or Kruskal-Wallis (MOL361-1E2-H121) test. Adjusted p-values
1020   represent p-values with Šidák correction for number of tests of that metabolite.

1021

1022   **Supplementary Table 3. Raw and normalized e/g WAT values for *Turicibacter* and *B.***
1023   ***thetaiotaomicron*-colonized mice.**

1024

1025   **Supplementary Table 4. Absolute quantification of lipid species from plasma of mice**
1026   **colonized with *B. thetaiotaomicron* engineered to express *Turicibacter bsh* genes, GF mice,**
1027   **or MOL361 monocolonized mice.** Raw p-values generated from Student's t-test comparison
1028   with Bt-WT for *Bacteroides* conditions or comparison between GF and MOL361. Adjusted p-
1029   values represent p-values with Šidák correction for number of tests of that lipid.

1030

1031   **Supplementary Table 5. List of oligos used in the work.**