Extravascular spaces are the primary reservoir of antigenic diversity in *Trypanosoma brucei* infection

Authors: Alexander K. Beaver[1], Nathan P. Crilly[1], Gracyn Y. Buenconsejo[2], Jill M.C. Hakim[2], Bailin Zhang[2], Bryce Bobb[2], Filipa Rijo-Ferreira[3,4,5], Luisa M. Figueiredo[3], and Monica R. Mugnier[2†].

Affiliations:

[1]Department of Pathology, Johns Hopkins School of Medicine; Baltimore, Maryland, United States of America

[2]Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health; Baltimore, Maryland, United States of America

[3]Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

[4]Department of Neuroscience, Peter O'Donnell Jr. Brain Institute, University of Texas Southwestern Medical Center; Dallas, Texas, United States.

[5]Division of Infectious Diseases and Vaccinology, Berkeley Public Health Molecular and Cell Biology Department; Berkeley, California, United States of America

† Corresponding author. Email: mmugnie1@jhu.edu

Abstract: The protozoan parasite *Trypanosoma brucei* evades clearance by the host immune system through antigenic variation of its dense variant surface glycoprotein (VSG) coat, periodically "switching" expression of the VSG using a large genomic repertoire of VSG-encoding genes[1–6]. Studies of antigenic variation in vivo have focused near exclusively on parasites in the bloodstream[5,7,8], but recent work has shown that many, if not most, parasites reside in the interstitial spaces of tissues[9–13]. We sought to explore the dynamics of antigenic variation in extravascular parasite populations using VSG-seq[7], a high-throughput sequencing approach for profiling VSGs expressed in populations of *T. brucei*. Here we show that tissues, not the blood, are the primary reservoir of antigenic diversity during *T. brucei* infection, with more than 85% of VSGs found exclusively within extravascular spaces. We found that this increased diversity is correlated to slower parasite clearance in tissue spaces. Together, these data support a model in which the slower immune response in extravascular spaces provides more time to generate the antigenic diversity needed to maintain a chronic infection. Our findings reveal the important role that extravascular spaces can play in pathogen diversification.

Main: Every pathogen must contend with the adaptive immune response of its host. *Trypanosoma brucei,* a protozoan parasite and causative agent of human and animal African Trypanosomiasis, has evolved a sophisticated strategy to evade this highly flexible and specific host response[14]. Transmitted by the bite of the tsetse fly, *T. brucei* lives extracellularly in the blood, lymph, and interstitial tissue spaces of its mammalian host[15]. To escape clearance by a continuous onslaught of host antibodies, the parasite periodically "switches" expression of its immunogenic variant surface glycoprotein (VSG) coat to new, antigenically distinct variants[1]. With a genomic repertoire of thousands of different VSG-encoding genes[2–4,6] and the ability to

generate novel VSGs through recombination[5,16], the parasite has an enormous capacity for
altering its antigenic profile.

Studies examining *T. brucei* antigenic variation in vivo have focused nearly exclusively on
parasites in the blood, revealing complex VSG expression dynamics[5,7,8]. However, it has recently
become clear that many, if not most, *T. brucei* parasites inhabit extravascular spaces during both
experimental and natural infections[9–13]. Though research has shown that tissue-resident parasites
adapt to these environments[11], cause tissue-specific symptoms[15], and are associated with
increased disease severity[13], it remains unclear why parasites invade tissue spaces and what role
these populations might play in infection.

Two older studies using low-throughput approaches point towards a role for tissue-resident
parasites in antigenic variation. These studies found that extravascular parasite populations were
antigenically distinct from those in the blood[17,18], suggesting that extravascular spaces could be a
site for antigenic variation or, alternatively, that certain VSGs influence the infiltration of or
adaptation to tissue spaces, leading to distinct VSG expression in extravascular spaces. Surface
proteins involved in antigenic variation influence tissue tropism in other pathogens. In the
malaria parasite *Plasmodium falciparum,* for example, the expression of a single *var* gene,
var2csa, allows for parasites to be sequestered in the placenta[19,20], while antigenic variants of
PilE in *Neisseria gonorrhoeae* influence adhesion to different host tissues[21–24]. It is thus plausible
that VSG could serve a similar function. Here, we use VSG-seq[7], a targeted RNA-sequencing
approach for profiling the VSGs expressed in *T. brucei* populations, to characterize the VSGs
expressed by extravascular *T. brucei* parasites. Our results show that extravascular spaces are
major reservoirs of antigenic diversity during *T. brucei* infection and that this parasite niche is
central to the parasite's ability to continuously outmaneuver the immune system.

Extravascular spaces contain most of the VSG diversity

To investigate how tissue-resident parasites contribute to antigenic variation in vivo, we
intravenously infected 12 mice, each with ~5 pleomorphic *T. brucei* EATRO1125 90-13
parasites[25]. We collected blood, then perfused mice with PBS-glucose (Supplementary Fig. 2)
and harvested the heart, lungs, gonadal fat, subcutaneous fat, brain, and skin at 6, 10, and 14 days
post-infection. For each sample, we extracted RNA and quantified VSG expression using VSG-
seq[7]. A single "initiating" VSG (either AnTat1.1 or EATRO1125 VSG-421) dominated
expression in both the blood and tissues on day 6 (Fig. 1a). This suggests that *T. brucei* invades
tissues before significant VSG switching occurs, which is in line with previous observations[26,27].
At later time points, VSG expression dynamics became more complex, with more VSGs
expressed, a unique composition of VSGs in each tissue, and no single dominating variant (Fig.
1a). While we found no evidence for tissue-specific VSGs or VSG sequence motifs
(Supplementary Fig. 4), the diversity in tissue spaces was, on average, two to four times higher
than the blood (Fig. 1c). This did not correlate with parasite load (Supplementary Fig. 1,3) and
was not driven by any specific tissue (Fig. 1d). The overall contribution of tissue-resident
parasites to antigenic diversity in a single infection was large: at any time, ~87% of expressed
VSGs in any individual infection were found exclusively within extravascular spaces (Fig. 1b).
Thus, tissues, not the blood, are the primary reservoirs of antigenic diversity during *T. brucei*
infection.

Fig. 1. Extravascular parasites harbor most of the antigenic diversity in an infection. (**a**) The percentage of parasites expressing each VSG within a space. The 11 VSGs with the highest overall expression are colored, and all other VSGs are in grey as "other". Gon. fat = gonadal fat, s.c. fat = subcutaneous fat. (**b**) Stacked bar graphs from each infected mouse representing the percentage of VSGs that were found exclusively within the blood (red), exclusively within tissue spaces (blue), or shared by both the blood and at least one tissue (green). (**c**) Quantification of the number of VSGs found within the blood (red) or tissue spaces (blue) at each time point (Shapiro-Wilk normality test followed by a two-tailed Student's t-test BH corrected). (**d**) The number of VSGs in each tissue space (Shapiro-Wilk normality test followed by a two-tailed Dunnett's test) (For statistical tests *P<0.05, ***P<0.001).

Frequently expressed VSGs first appear in tissue populations

The high antigenic diversity observed in tissues could serve to maintain a chronic infection. If
antigenic variation occurs relatively rarely in the bloodstream, then parasites from extravascular
spaces might serve as a source of new, antigenically distinct, VSGs. In line with this, tissue
spaces contain more "unique VSGs", those VSGs that are expressed exclusively within that

space in an infection, than the blood (Fig. 2a,b). To examine the potential for tissue-resident
VSGs to contribute to antigenic variation systemically, we identified VSGs only expressed in
tissues on day 6 post-infection and analyzed whether they later appeared within the blood. The
majority (74%) of these VSGs were expressed within the blood on day 10 or 14. Analysis of
individual VSGs revealed that rare VSGs expressed at low levels exclusively in tissue spaces
also have the potential to become ubiquitously expressed within a host (Fig. 2c). In addition to a
model in which tissue spaces provide new VSGs to re-seed the blood, it is possible that tissue-
resident parasites undergo antigenic variation before blood-resident populations. Therefore, these
data could be explained by either a trigger within the tissue environment that induces parasite
switching or a differential selective pressure imposed by the tissue environment.

Fig. 2. Tissue-resident parasites express a unique repertoire of VSGs during infection.
(**a**) We define "unique" VSGs as those VSGs solely found within a specific space in a mouse
(Created with BioRender.com). (**b**) The percentage of VSGs that were unique to one space
within a mouse (Shapiro-Wilk normality test followed by a two-tailed Dunnett's test,
*$P<0.05$). Day 6 samples were excluded from this analysis because few VSGs are expressed
at this point. Gon. fat = gonadal fat, s.c. fat = subcutaneous fat. (**c**) The expression of three
representative VSGs (cluster 294, 504, and 1831) within all samples on days 6, 10, and 14.
"n.d" indicates that the VSG was not detected.

99
VSG-specific parasite clearance is delayed in tissues compared to the blood

Indeed, our data suggest that the environment within tissue spaces is distinct from the blood,
with parasite clearance occurring at different rates in each space. While parasites expressing the
initiating VSG were cleared from the blood by day 10 post-infection, they were not cleared from
tissues until at least day 14 (Fig. 3a). This suggests that that the clearance of parasites within
extravascular spaces is delayed, but not abolished, compared to the blood. We confirmed this
observation by flow cytometry using the tdTomato-expressing "triple reporter" *T. brucei*
EATRO1125 AnTat 1.1E cell line[28] (Fig. 3b and 3c). Similar to the increase in antigenic
diversity we observed in every tissue space, this delay in clearance was not tissue-specific. Thus,
the immune mechanisms influencing extravascular parasite clearance appear to be general
features of extravascular spaces.

Fig. 3. VSG-specific parasite clearance is slower in tissues than in the blood. (**a**) The percentage of parasites expressing the initiating VSG (AnTat1.1 or VSG-421) at days 6, 10, and 14 post-infection. Tissue samples were grouped together (blue) and compared to blood samples (red) (two-tailed Wilcoxon test, ns = not significant, **P<0.01). "n.d" = not detected, gon. fat = gonadal fat, s.c. fat = subcutaneous fat. (**b**) Quantification of the number of parasites that were tdTomato positive and stained positive for AnTat1.1 by flow cytometry (n = 5 per group). The horizontal dotted line represents the limit of detection for VSG-seq. "n.d" indicates that the VSG was not detected. (**c**) Representative flow cytometry plots from tissues collected from mice infected with chimeric triple marker parasites that express tdTomato constitutively in their cytoplasm. Parasites were stained with anti-AnTat1.1 antibody.

Delayed clearance of parasites leads to increased VSG diversity

The increased antigenic diversity in extravascular spaces could be explained by the distinct
clearance dynamics we observe, as prolonged survival in tissues could provide more time for

parasites to switch. To test whether there was a link between parasite survival and increased
diversity, we sought to interrupt parasite clearance in tissues. Because parasite clearance in the
blood (days 8-10) coincides with the appearance of anti-VSG IgM, and parasite clearance in the
tissues (days 10-14) coincides with the anti-VSG IgG response[1], we hypothesized that clearance
in tissues is dependent on the anti-VSG IgG response. Thus, the loss of IgG might abrogate the
clearance of tissue-resident parasites. To test this hypothesis, we infected activation-induced
cytidine deaminase (AID) knockout (AID Cre) mice[29], which only produce IgM antibodies
(Supplementary Fig. 5), and analyzed blood and tissues from days 6 and 14 post-infection by
VSG-seq. As expected, clearance of the initiating VSG was severely delayed in tissues on day 14
in AID$^{-/-}$ mice, suggesting that IgG is important, if not critical, for the clearance of extravascular
parasites. This could be explained by the fact that IgM, a bulky pentamer, does not diffuse
efficiently into tissue spaces[30,31], while IgG, a monomer, readily diffuses. We also observed a
defect in the clearance of blood-resident parasites in AID$^{-/-}$ mice (Fig. 4a). In both blood and
tissues where clearance was delayed, more VSGs were detected on day 14 post-infection
compared to wild-type (Fig. 4b), revealing a direct relationship between the timing of parasite
clearance and VSG diversity. Regardless of their local environment (intra- or extravascular),
longer-lived parasite populations generated more diverse sets of VSGs.

Fig. 4. Delayed parasite clearance correlates with an increase in VSG diversity. (a) The
percentage of parasites expressing the initiating VSG (AnTat1.1 or VSG-421) in both WT and
AID$^{-/-}$ mice. "n.d" indicates that the VSG was not detected. **(b)** The number of VSGs
expressed within the blood (red) and tissues (blue) of WT and AID$^{-/-}$ mice (Shapiro-Wilk
normality test followed by a two-tailed Student's t-test, **P<0.01).

Discussion

134 Previous in vivo studies of antigenic variation in *T. brucei* have focused mainly on blood-
135 resident populations. Here we show that extravascular spaces are the primary reservoir of VSGs,
136 accounting for the vast majority of antigenic diversity in any individual infection. The number of
137 VSGs we detected in the blood corroborates previous estimates[5,7,8], while the diversity in tissues
138 is two to four times higher. Our data suggest that this is likely the result of slower clearance

dynamics in extravascular spaces and highlight the role that tissue spaces can play in the generation of genetic diversity during infection.

Although extravascular parasite populations were highly diverse, we saw no evidence of tissue-specific VSG expression. Because parasites invade tissues efficiently before significant VSG switching has occurred, it appears unlikely that any specific VSG is required for tissue invasion. Whether VSGs could influence parasite fitness in specific host spaces is less clear. We measured VSG expression up to day 14 post-infection, at which point tissue-resident populations are just beginning to diverge from one another and the blood. It is therefore possible that, as these populations further evolve, there may be selection for VSGs better adapted to certain tissue spaces.

The striking diversity observed in tissue-resident *T. brucei* populations could be important for the maintenance of natural infections, particularly in wild animals, where pre-existing anti-VSG immunity is more likely to exist. A high switch rate may be required to ensure some parasites successfully evade the host's existing anti-VSG antibody repertoire. In addition, infections in the wild can last for months to years[14,32]. During these long infections, the large reservoirs of VSG diversity found in tissues may be essential for both the establishment and maintenance of a chronic *T. brucei* infection.

Indeed, our data support the idea that the large reservoirs of antigenic diversity in extravascular spaces contribute to systemic infection when parasites re-enter the blood after switching: rare VSGs expressed exclusively in tissues at early time points are expressed in the blood and other spaces later. There is another intriguing explanation for this observation, however. Switching in *T. brucei* is known to be semi-predictable[7,33], and it is possible that tissue-resident parasites simply switch earlier, or more frequently, than those in the blood. In this case, the same VSGs would arise independently in every population without any movement between spaces. While the high vascular permeability observed after the initial stages of infection[13] indicates that parasites likely move back and forth between the vasculature and extravascular spaces, it is exciting to speculate that some aspect of the extravascular environment supplies a molecular or physical stimulus that promotes VSG switching as well.

While the timing of VSG switching could be the result of an environmental trigger, our data suggest that the higher antigenic diversity in extravascular spaces compared to the blood can be explained by the dynamics of the immune response to *T. brucei* in each space. In tissue spaces, we observed slower VSG-specific clearance of parasites than in the blood, potentially providing these populations more time to undergo antigenic variation. Additionally, newly switched parasites are still vulnerable to immune clearance by antibodies against their previous VSG for ~29 hours, so moderate delays in immune clearance could allow more switched parasites to survive[34]. The direct relationship we observed between the timing of parasite clearance and antigenic diversity in AID[-/-] mice strongly supports this model, with even small delays in clearance rate showing major effects on parasite VSG diversity. Our results show that the production of IgG plays a key role in clearing parasites from tissues, which we propose could be related to the ready diffusion of this molecule within extravascular spaces facilitating parasite clearance. Of course, the local immune response to *T. brucei* is likely complex, with multiple features likely to play a role in parasite detection and clearance. More research will be required to fully understand the nature of the extravascular immune response that results in differential *T. brucei* clearance in tissues.

Altogether, our results outline a model in which *T. brucei* parasites "hide" in extravascular
spaces to generate new antigenic variants capable of exiting tissues and aiding in systemic
immune evasion. Coupled with other recent studies[9–13,35], this suggests a new framework for the
progression and pathogenesis of *T. brucei* infections. Instead of being the primary parasite
reservoir, the blood may represent a transient population that is regularly re-seeded by
extravascular parasites. The vasculature, then, might act as a highway system for movement
between the tissue spaces and for eventual transmission back into the tsetse fly.

Interfering with the egress from or establishment within tissue spaces might be a strategy for
treating *T. brucei* infections or other infections with pathogens that rely on the distinct features of
the extravascular environment to diversify. In line with this, one recent study showed that partial
inhibition of *T. brucei* tissue invasion using antibodies against P- and E-selectins results in
prolonged survival in mice[13]. It is possible that without the proper establishment of parasite
tissue reservoirs, overall antigenic diversity is lowered, limiting the parasite's capacity for
immune evasion and leading to a decrease in parasite virulence. Defining the dynamics and
variation of parasites both within and between spaces, as well as the unique host environment
within each tissue space, will be central to understanding how *T. brucei* consistently avoids
immune clearance and harnessing this mechanism for disease control.

More broadly, these data demonstrate how the different environmental and immune pressures
within a host can influence pathogen diversification. This production of genetic heterogeneity
within an infection is important for many pathogen virulence processes, including establishing
and maintaining infection, facilitating immune evasion, generating drug resistance, and adapting
to different host environments. The extravascular environment plays a unique role in promoting
pathogen evolution, and *T. brucei* serves as a valuable model for understanding this aspect of the
host-pathogen interface.

Methods:
<u>Mouse infections and sample collection</u>
Female C57Bl/6J (WT, strain# 000664 Jackson Laboratory) or B6.129P2-
Aicda^(tm1(cre)Mnz)/J (AID[-/-], strain# 007770 Jackson Laboratory)[29] between 7-10 weeks old were
each infected by intravenous tail vein injection with ~5 pleiomorphic EATRO 1125 AnTat1.1E
90-13 *T. brucei* parasites[25]. Blood parasitemia was counted by tail bleed every 2 days starting on
day 4 post-infection (PI) by hemocytometer. Blood (25uL) was collected by a submandibular
bleed on days 6, 10, and 14 PI and placed into TRIzol LS. For WT mice, 4 Mice were
anesthetized and perfused at days 6, 10, and 14 PI. Infected AID[-/-] mice were anesthetized and
perfused at days 6 (2 mice) and 13 (3 mice) PI. Mice were perfused with 50mL of PBS-Glucose
(0.055M D-glucose) with heparin. After perfusion, tissues were dissected and placed
immediately into 1mL of RNA later. The heart, lungs, gonadal fat, subcutaneous fat, brain, and
skin (ear) were collected.
For flow cytometry and immunofluorescence experiments, 7-10 week old female
C57Bl/6J mice were infected by intravenous tail vein injection with ~5 AnTat1.1E chimeric
triple reporter *T. brucei* parasites which express tdTomato[28]. Blood was collected by a
submandibular bleed at each time point. Mice were anesthetized and perfused on days 6 and 13
P.I. as discussed above and the gonadal fat and lungs were harvested. All animal studies were
approved by the Johns Hopkins Animal Care and Use Committee (protocol # MO22H163).
227

 VSG-seq sample and library preparation

RNA was isolated from blood samples stored in TRIzol LS (ThermoFisher, 10296010)
by phenol/chloroform extraction. Tissue samples were weighed and homogenized in TRIzol, and
then RNA was isolated by phenol/chloroform extraction. RNA from each sample was DNase
treated using Turbo DNase and cleaned up with Mag-Bind® TotalPure NGS beads (Omega Bio-
Tek M1378-00). First-strand cDNA synthesis was performed using SuperScript III Reverse
Transcriptase and a primer that binds to the conserved VSG 14-mer in the 3′-UTR (5'-
GTGTTAAAATATATC-3'). Products were cleaned up using Mag-Bind® TotalPure NGS beads
(Omega Bio-Tek, M1378-01). Next, a VSG-specific PCR with Phusion polymerase
(ThermoFisher, F530L) was performed using primers for the spliced leader (5'-
ACAGTTTCTGTACTATATTG-3') and SP6-VSG 14-mer sequences (5'-
GATTTAGGTGACACTATAGTGTTAAAATATATC-3') for 25 cycles. VSG-PCR products
were cleaned up using Mag-Bind® TotalPure NGS beads and quantified using the QuBit HS
DNA kit (Life Technologies). Finally, sequencing libraries were prepared with the Nextera XT
DNA Sample Prep Kit (Illumina) using the manufacturer's guidelines, and libraries were
sequenced with 100bp single-end reads on an Illumina HiSeq 2500.

Tissue QPCR
First-strand synthesis was performed with SuperScript III Reverse Transcriptase (Thermo
Fisher Scientific, 18080051) and random hexamers primers on tissue RNA samples. QPCR was
performed in triplicate using SYBR Green qPCR Master Mix (Invitrogen, 4309155). ZFP3
primers were used to estimate parasite load in tissue samples (FW: 5'-
CAGGGGAAACGCAAAACTAA-3'; RV: 5'-TGTCACCCCAACTGCATTCT-3'). CT values
were averaged between the triplicates and parasite load per mg of tissue were estimated using a
standard curve of values from RNA isolated from known numbers of cultured parasites.

VSG-seq analysis
Analysis of sequencing results was performed following the method we reported
previously[7], with two changes: no mismatches were allowed for bowtie alignments and each
sample was analyzed (assembly, alignment, and quantification) separately. To compare
expressed VSG sets between samples, all assembled VSGs were clustered using CD-HIT-EST[36].
VSGs with >98% identity to one another were conservatively treated as one VSG. VSGs were
then identified by their Cluster number for further analysis. Samples that had less than 100,000
successfully aligning reads to VSGs were excluded from further analysis. Four samples, 3 brain
and 1 heart, were discarded because fewer than 100,000 reads aligned to VSG (Supplemental
Figure 1A). Downstream analysis of expression data and generation of figures was performed in
R.

Analysis of VSG sequence motifs
To identify whether there were tissue-specific VSG sequence motifs, the similarity of N
terminal sequences from all assembled VSGs were compared. N terminal sequences were
identified using a HMMEr scan against a database curated by Cross et al[2,37]. All N termini were
compared in an all vs all blast using default parameters. All VSG pairwise comparisons with an
e-value higher than 1E-3 were considered sufficiently similar to one another for further analysis.
VSGs that were found in a given tissue were binned into that tissue group, and the distribution of

the BLAST bitscores in a given compartment was compared against the total population of
similar VSGs.
275

<u>Flow Cytometry</u>
Once mice were perfused, tissues were dissected and washed with HBSS (Hanks
balanced salt solution, ThermoFisher Scientific 14175095). Tissue samples were minced and
placed in DMEM (ThermoFisher Scientific, 11995065) containing either 1 mg/mL collagenase
type 1 (ThermoFisher Scientific, 17100017) for adipose fat or 2 mg/mL collagenase type 2
(ThermoFisher Scientific, 17101015) for lung samples. These were then incubated in a 37°C
water bath for 1 hour and briefly vortexed every 10 minutes. Next, samples were passed through
a 70μM filter and centrifuged at 2600 x g for 8 mins at 4 C, and the cell pellet was taken for
antibody staining.
Blood samples were collected by submandibular bleed and red blood cells were depleted
by magnetic-activated cell sorting (MACS) with anti-Ter-119 MicroBeads (Miltenyi Biotech,
130-049-901) following the manufacturer's protocol. Cells were pelleted and washed with HMI-
9 media.
All samples, both blood and tissues, were stained with Zombie Aqua™ dye at 1:100 in
PBS and washed with PBS following the manufacturer's protocol (BioLegend, 423101).
Samples were then stained for 10 minutes at 4°C with a rabbit anti-AnTat1.1 polyclonal antibody
diluted 1:15,000 in HMI-9 media and washed once with HMI-9 (antibody courtesy of Jay
Bangs). Then, secondary antibody staining was performed for 10 minutes at 4°C with Anti-
Rabbit IgG conjugated to Alexa Fluor® 488 fluorescent dye (Cell Signaling Technology,
4412S). Finally, samples were washed with cold PBS and resuspended in PBS for flow
cytometry analysis. Samples were run on a Beckton Dickenson A3 Symphony flow cytometer
and analysis was performed using FlowJo (version 10.6.1).
298

<u>Immunofluorescence</u>
Mice infected with AnTat1.1E chimeric triple reporter *T. brucei* parasites that express
tdTomato were sacrificed and perfused as previously described at days 6 and 13 PI. Lung, heart,
and gonadal fat were collected and fixed in 4% paraformaldehyde in PBS for 12 hours at 4 C.
Post-fixation, tissues were frozen, embedded in O.C.T. Compound (Tissue-Tek), and cut by
cryostat microtome into 10μm sections.
The following antibodies were applied to sections: rat anti-mouse CD31 (PECAM-1)
(SCBT) with goat anti-rat Fluor 488 (CST). Coverslips were mounted using ProLong Gold (Life
technologies). Tissues were imaged with 4x, 10x, and 20x objectives using a Nikon Eclipse 90i
fluorescence microscope (Nikon) and X-Cite 120 fluorescent lamp (Excelitas) with an ORCA-
ER digital CCD camera (Hammamatsu) and ImageJ v1.53 image analysis software. Image
collection and analysis followed published guidelines for rigor and reproducibility[38].
311

<u>Serum antibody ELISA quantification</u>
Blood (25uL) was collected by submandibular bleed on days 0, 6, 10, and 14 PI from
mice infected with ~5 pleiomorphic EATRO 1125 AnTat1.1E 90-13 *T. brucei* parasites. Serum
was isolated using serum separator tubes (BD Microtainer SST tubes, 365967). IgM and IgG
were quantified by ELISA using ThermoFisher IgM and IgG kits following manufacturer
protocols (IgG cat# 88-50400-88, IgM cat# 88-50470-88).
318

Statistics and figures

Normality was tested for all Students t-tests and Dunnet's tests analyses and can be found in the code on the repository at https://github.com/mugnierlab/Beaver2022. Nearly all VSG diversity measurements were found to be normally distributed, except for some samples with low VSG counts or sample numbers. We thus assumed normality for all VSG diversity measurements. All reported P-values have been corrected for multiple comparisons using the Benjamini–Hochberg procedure. For all figures with boxplots, the box represents the first (25%) and third (75%) quartiles with a line at the median. Extending lines represent the maximum and minimum values not including outliers that are further than 1.5 times the interquartile range from either end of the box.

Data availability:
Code and data for generating the analysis and figures in this paper are available at https://github.com/mugnierlab/Beaver2022. Raw sequencing data are available in National Center for Biotechnology Information (NCBI) Sequence Read Archive under accession number PRJNA858046.

Acknowledgments: We would like to thank Jay Bangs for supplying us with the anti-AnTat1.1 antibody and Brice Rotureau for providing us with the chimeric triple reporter cell line. We thank Tricia Nilles and Worod Allak at the Becton Dickinson Immunology and Flow Cytometry core at Johns Hopkins Bloomberg School of Public Health for training, support, and technical assistance using the BD FACS Symphony A3. The facility is supported in part by CFAR: 5P30AI094189-04 (Chaisson). Data analysis was carried out at the Advanced Research Computing at Hopkins (ARCH) core facility (rockfish.jhu.edu), which is supported by the National Science Foundation (NSF) grant number OAC 1920103.

Author contributions:

Conceptualization: AB, MRM, LMF, FRF

Methodology: AB, NC, GYB, JH, BZ, BB

Investigation: AB, NC, GYB, JH

Visualization: AB, NC, JH

Funding acquisition: MRM, LMF

Project administration: MRM

Supervision: MRM

Writing – original draft: AB, MRM

Writing – review & editing: AB, GYB, BZ, MRM, FRF, LF

Competing interests: The authors declare that they have no competing interests.

Materials and correspondence: Correspondence and requests for materials should be addressed to Monica Mugnier (mmugnie1@jhu.edu).

References:

1.   Magez, S. *et al.* The Role of B-cells and IgM Antibodies in Parasitemia, Anemia, and
VSG Switching in Trypanosoma brucei–Infected Mice. *PLoS Pathog.* **4**, e1000122 (2008).

2.   Cross, G. A. M., Kim, H. S. & Wickstead, B. Capturing the variant surface glycoprotein
repertoire (the VSGnome) of Trypanosoma brucei Lister 427. *Mol. Biochem. Parasitol.*
195, 59–73 (2014).

3.   Müller, L. S. M. *et al.* Genome organization and DNA accessibility control antigenic
variation in trypanosomes. *Nat. 2018 5637729* **563**, 121–125 (2018).

4.   Cosentino, R., Brink, B. & Siegel, T. A phased genome assembly for allele-specific
analysis in Trypanosoma brucei. *bioRxiv* 2021.04.13.439624 (2021).
doi:10.1101/2021.04.13.439624

5.   Hall, J. P. J., Wang, H. & David Barry, J. Mosaic VSGs and the Scale of Trypanosoma
brucei Antigenic Variation. *PLoS Pathog.* **9**, (2013).

6.   Hertz-Fowler, C. *et al.* Telomeric Expression Sites Are Highly Conserved in
Trypanosoma brucei. *PLoS One* **3**, e3527 (2008).

7.   Mugnier, M. R., Cross, G. A. M. & Papavasiliou, F. N. The in vivo dynamics of antigenic
variation in Trypanosoma brucei. *Science* **347**, 1470–3 (2015).

8.   Jayaraman, S. *et al.* Application of long read sequencing to determine expressed antigen
diversity in Trypanosoma brucei infections. *PLoS Negl. Trop. Dis.* **13**, e0007262 (2019).

9.   Capewell, P. *et al.* The skin is a significant but overlooked anatomical reservoir for
vector-borne African trypanosomes. *Elife* **5**, (2016).

10.   Camara, M. *et al.* Extravascular Dermal Trypanosomes in Suspected and Confirmed Cases
of gambiense Human African Trypanosomiasis. *Clin. Infect. Dis. An Off. Publ. Infect. Dis.*
Soc. Am. **73**, 12 (2021).

11.   Trindade, S. *et al.* Trypanosoma brucei Parasites Occupy and Functionally Adapt to the
Adipose Tissue in Mice. *Cell Host Microbe* (2016). doi:10.1016/j.chom.2016.05.002

12.   Carvalho, T. *et al.* Trypanosoma brucei triggers a marked immune response in male
reproductive organs. *PLoS Negl. Trop. Dis.* (2018). doi:10.1371/journal.pntd.0006690

13.   De Niz, M. *et al.* Organotypic endothelial adhesion molecules are key for Trypanosoma
brucei tropism and virulence. *Cell Rep.* **36**, 109741 (2021).

14.   WHO. Control and surveillance of human African trypanosomiasis: Report of a WHO
Expert Committee. *WHO Tech. Rep. Ser.* 1–237 (2013).

15.   Crilly, N. P. & Mugnier, M. R. Thinking outside the blood: Perspectives on tissue-resident
Trypanosoma brucei. *PLOS Pathog.* **17**, e1009866 (2021).

16.   Kamper, S. M. & Barbet, A. F. Surface epitope variation via mosaic gene formation is
potential key to long-term survival of Trypanosoma brucei. *Mol. Biochem. Parasitol.* **53**,
33–44 (1992).

17.   Seed, J. R. & Eefron, H. G. Simultaneous presence of different antigenic populations of
Trypanosoma brucei gambiense in Microtus montanus. *Parasitology* **66**, 269–278 (1973).

18.   SEED, J. R., EDWARDS, R. & SECHELSKI, J. The ecology of antigenic variation. *J.*

Protozool. **31**, 48–53 (1984).

19.   Salanti, A. *et al.* Evidence for the Involvement of VAR2CSA in Pregnancy-associated
Malaria. *J. Exp. Med.* **200**, 1197–1203 (2004).

20.   Duffy, P. E. & Fried, M. Plasmodium falciparum adhesion in the placenta. *Curr. Opin.*
Microbiol. **6**, 371–376 (2003).

21.   Jonsson, A. -B, Ilver, D., Falk, P., Pepose, J. & Normark, S. Sequence changes in the pilus
subunit lead to tropism variation of Neisseria gonorrhoeae to human tissue. *Mol.*
Microbiol. **13**, 403–416 (1994).

22.   Nassif, X. *et al.* Antigenic variation of pilin regulates adhesion of Neisseria meningitidis
to human epithelial cells. *Mol. Microbiol.* **8**, 719–725 (1993).

23.   Rudel, T., van Putten, J. P. M., Gibbs, C. P., Haas, R. & Meyer, T. F. Interaction of two
variable proteins (PilE and PilC) required for pilus-mediated adherence of Neisseria
gonorrhoeae to human epithelial cells. *Mol. Microbiol.* **6**, 3439–3450 (1992).

24.   Virji, M. & Heckels, J. E. The role of common and type-specific pilus antigenic domains
in adhesion and virulence of gonococci for human epithelial cells. *J. Gen. Microbiol.* **130**,
1089–1095 (1984).

25.   Engstler, M. & Boshart, M. Cold shock and regulation of surface protein trafficking
convey sensitization to inducers of stage differentiation in Trypanosoma brucei. *Genes*
Dev. **18**, 2798 (2004).

26.   Turner, C. M., Hunter, C. A., Barry, J. D. & Vickerman, K. Similarity in variable antigen
type composition of Trypanosoma brucei rhodesiense populations in different sites within
the mouse host. *Trans. R. Soc. Trop. Med. Hyg.* **80**, 824–830 (1986).

27.   Turner, C. M. R. & Barry, J. D. High frequency of antigenic variation in Trypanosoma
brucei rhodesiense infections. *Parasitology* **99 Pt 1**, 67–75 (1989).

28.   Calvo-Alvarez, E., Cren-Travaillé, C., Crouzols, A. & Rotureau, B. A new chimeric triple
reporter fusion protein as a tool for in vitro and in vivo multimodal imaging to monitor the
development of African trypanosomes and Leishmania parasites. *Infect. Genet. Evol.* **63**,
391–403 (2018).

29.   Robbiani, D. F. *et al.* Activation Induced Deaminase is required for the chromosomal
breaks in c-myc that lead to c-myc/IgH translocations. *Cell* **135**, 1028 (2008).

30.   Hector, R. F., Collins, M. S. & Pennington, J. E. Treatment of Experimental Pseudomonas
aeruginosa Pneumonia with a Human IgM Monoclonal Antibody. *J. Infect. Dis.* **160**, 483–
489 (1989).

31.   BARTH, W. F., WOCHNER, R. D., WALDMANN, T. A. & FAHEY, J. L. Metabolism
of Human Gamma Macroglobulins. *J. Clin. Invest.* **43**, 1036 (1964).

32.   Mehlitz, D. & Molyneux, D. H. The elimination of Trypanosoma brucei gambiense?
Challenges of reservoir hosts and transmission cycles: Expect the unexpected. *Parasite*
Epidemiol. Control **6**, e00113 (2019).

33.   Morrison, L. J., Majiwa, P., Read, A. F. & Barry, J. D. Probabilistic order in antigenic
variation of Trypanosoma brucei. *Int. J. Parasitol.* **35**, 961–972 (2005).

34.   Pinger, J., Chowdhury, S. & Papavasiliou, F. N. Variant surface glycoprotein density
defines an immune evasion threshold for African trypanosomes undergoing antigenic
variation. *Nat. Commun.* (2017). doi:10.1038/s41467-017-00959-w

35.   Trindade, S. *et al.* Slow growing behavior in African trypanosomes during adipose tissue
colonization. *Nat. Commun. 2022 131* **13**, 1–13 (2022).

36.   Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150 (2012).

37.   So, J. *et al.* Trypanosomal variant surface glycoprotein expression in human African
trypanosomiasis patients. *bioRxiv* 2021.09.09.459620 (2021).
doi:10.1101/2021.09.09.459620

38.   Lee, J.-Y. & Kitaoka, M. A beginner's guide to rigor and reproducibility in fluorescence
imaging experiments. *https://doi.org/10.1091/mbc.E17-05-0276* **29**, 1519–1525 (2018).

449