Extravascular spaces are the primary reservoir of antigenic diversity in *Trypanosoma brucei* infection

Authors: Alexander K. Beaver[1,2], Zhibek Keneskhanova[3,4], Raúl O. Cosentino[3,4], Brian L. Weiss[5], Erick O. Awuoche[5], Gretchen M. Smallenberger[5], Gracyn Y. Buenconsejo[2], Nathan P. Crilly[1,2], Jaclyn E. Smith[2], Jill M.C. Hakim[2], Bailin Zhang[2], Bryce Bobb[2], Filipa Rijo-Ferreira[6,7,8], Luisa M. Figueiredo[6], Serap Aksoy[5], T. Nicolai Siegel[3,4], and Monica R. Mugnier[2†].

Affiliations:

[1]Department of Pathology, Johns Hopkins School of Medicine; Baltimore, Maryland, United States of America

[2]Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health; Baltimore, Maryland, United States of America

[3]Division of Experimental Parasitology, Faculty of Veterinary Medicine, Ludwig-Maximilians-Universität München, Munich, Germany

[4]Biomedical Center, Division of Physiological Chemistry, Faculty of Medicine, Ludwig-Maximilians-Universität München, Munich, Germany

[5]Department of Epidemiology of Microbial Diseases, Yale School of Public Health; New Haven, Connecticut, United States of America

[6]Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

[7]Department of Neuroscience, Peter O'Donnell Jr. Brain Institute, University of Texas Southwestern Medical Center; Dallas, Texas, United States of America

[8]Division of Infectious Diseases and Vaccinology, Berkeley Public Health Molecular and Cell Biology Department; Berkeley, California, United States of America

† Corresponding author. Email: mmugnie1@jhu.edu

Summary paragraph: The protozoan parasite *Trypanosoma brucei* evades clearance by the host immune system through antigenic variation of its dense variant surface glycoprotein (VSG) coat, periodically "switching" expression of the VSG using a large genomic repertoire of VSG-encoding genes[1–6]. Recent studies of antigenic variation in vivo have focused near exclusively on parasites in the bloodstream[4,7,8], but research has shown that many, if not most, parasites reside in the interstitial spaces of tissues[9–13]. We sought to explore the dynamics of antigenic variation in extravascular parasite populations using VSG-seq[7], a high-throughput sequencing approach for profiling VSGs expressed in populations of *T. brucei*. Here we show that tissues, not the blood, are the primary reservoir of antigenic diversity during both needle- and tsetse bite-initiated *T. brucei* infections, with more than 75% of VSGs found exclusively within extravascular spaces. We found that this increased diversity is correlated with slower parasite clearance in tissue spaces. Together, these data support a model in which the slower immune response in extravascular spaces provides more time to generate the antigenic diversity needed to maintain a chronic infection. Our findings reveal the important role that extravascular spaces can play in pathogen diversification.

Main: Every pathogen must contend with the adaptive immune response of its host.
Trypanosoma brucei, a protozoan parasite and causative agent of human and animal African
Trypanosomiasis, has evolved a sophisticated strategy to evade this highly flexible and specific
host response[14]. Transmitted by the bite of the tsetse fly, *T. brucei* lives extracellularly in the
blood, lymph, and interstitial tissue spaces of its mammalian host[15]. To escape clearance by a
continuous onslaught of host antibodies, the parasite periodically "switches" expression of its
immunogenic variant surface glycoprotein (VSG) coat to new, antigenically distinct variants[1].
With a genomic repertoire of thousands of different VSG-encoding genes[2,3,5,6] and the ability to
generate novel VSGs through recombination[4,8,16], the parasite has an enormous capacity for
altering its antigenic profile.

Studies examining *T. brucei* antigenic variation in vivo have focused nearly exclusively on
parasites in the blood, revealing complex VSG expression dynamics[4,7,8]. However, it has recently
become clear that many, if not most, *T. brucei* parasites inhabit extravascular spaces during both
experimental and natural infections[9–13]. Though research has shown that tissue-resident parasites
adapt to these environments[11], cause tissue-specific symptoms[15], and are associated with
increased disease severity[13], it remains unclear why parasites invade tissue spaces and what role
these populations might play in infection.

Several older studies suggested a role for tissue-resident parasites in antigenic variation. These
studies found that brain- and lymph-resident parasite populations were antigenically distinct
from those in the blood[17–20], and that antigenic types detectable in the lymphatic fluid could be
detected in the blood at later timepoints during infection[19]. This led these researchers to
hypothesize that extravascular spaces might be a site for antigenic variation, with antigenic
variants generated in extravascular spaces contributing to systemic immune evasion. However, a
later study focusing on very early timepoints post-infection found no antigenic difference
between *T. brucei* populations within the blood and several extravascular spaces[21], and the
community consequently ruled out a role for extravascular parasites in antigenic variation[22,23].
These early investigations were limited by the methodology available at the time, which relied
on the use of VSG-specific antisera to analyze parasite antigenic diversity. Modern high-
throughput sequencing methods allow VSG expression to be measured accurately and in high
resolution[7]. Given the mounting evidence that *T. brucei* parasites persist in and adapt to
extravascular spaces, there is a clear need to reinvestigate the dynamics of VSG expression
within tissue spaces.

Here, we use VSG-seq[7], a targeted RNA-sequencing approach for profiling the VSGs expressed
in *T. brucei* populations, to characterize the VSGs expressed by extravascular *T. brucei* parasites.
Our results show that extravascular spaces are major reservoirs of antigenic diversity during *T.*
brucei infection and that this parasite niche is central to the parasite's ability to continuously
outmaneuver the immune system.

Extravascular spaces contain most of the VSG diversity

To investigate how tissue-resident parasites contribute to antigenic variation in vivo, we
intravenously (IV) infected 12 mice, each with ~5 pleomorphic *T. brucei* EATRO1125 90-13
parasites[24]. We collected blood, then perfused mice with PBS-glucose (Extended Data Fig. 1)
and harvested the heart, lungs, gonadal fat, subcutaneous fat, brain, and skin at 6, 10, and 14 days
post-infection. For each sample, we extracted RNA and quantified *T. brucei* VSG expression
using VSG-seq[7]. A single "initiating" VSG (either AnTat1.1 or EATRO1125 VSG-421)

dominated expression in both the blood and tissues on day 6 (Fig. 1a), in line with previous
observations[21,25]. At later time points, VSG expression dynamics became more complex, with
more VSGs expressed, a unique composition of VSGs in each tissue, and no single dominating
variant (Fig. 1a). Although tissue-specific expression of variant surface proteins is a feature in
other organisms that use antigenic variation[26–31], we found no evidence for tissue-specific VSGs
or VSG sequence motifs (Extended Data Fig. 2). Instead, we observed an increase in antigenic
diversity in extravascular spaces. The number of detectable VSGs in tissue spaces was, on
average, two to four times higher than the blood (Fig. 1c). This did not correlate with parasite
load (Extended Data Fig. 3 and Extended Data Fig. 4ab) and was not driven by any specific
tissue (Fig. 1d). In addition, parasite differentiation to the non-dividing tsetse infective form via
quorum sensing, which is marked by expression of the PAD1 gene[32], did not correlate with VSG
diversity in either the blood or tissues at the population level (Extended Data Fig. 4c). The
overall contribution of tissue-resident parasites to antigenic diversity in a single infection was
large: at any time, ~87% of expressed VSGs in any individual infection were found exclusively
within extravascular spaces (Fig. 1b).

VSG-seq is a bulk measure of VSG expression. To be sure that the increased VSG diversity we
observed in tissues was not due to the derepression of silent VSGs within individual cells, we
performed single-cell RNA sequencing using the SL-Smart-seq3xpress platform to quantify VSG
expression in single cells from blood and tissue samples[33]. Because *T. brucei* can form new
"mosaic" VSGs through recombination[4,8,16,34], which are, by definition, absent from reference
genomes, we initially used our VSG-Seq pipeline to *de novo* assemble VSGs in each cell. This
approach accounts for the possibility that expressed VSGs might be absent from the reference
genome, either as a result of mosaic VSG formation or as a result of an incomplete genome
assembly, potentially affecting quantification and/or read mapping. Only one VSG assembled in
most cells (94.2%; Extended Data Fig. 5a). We also mapped sequencing reads to the
EATRO1125 genome, which could reveal more subtle signatures of derepression. By this
analysis, there was no obvious difference between the blood and the tissues in the number of
expressed VSGs in each cell (Extended Data Fig. 5b) or in the relative expression of the most
abundant VSG in each cell (Extended Data Fig. 5c).

To estimate the number of cells maintaining monogenic expression, we defined a cell as
maintaining monogenic expression if 80% of UMIs mapping to VSGs mapped to a single VSG.
Although mapping to the genome suggested that most cells (62.7%) maintained monogenic
expression based on this threshold, the proportion of cells estimated to maintain monogenic
expression was lower than estimated by *de novo* assembly. Comparison of the two analyses
revealed that of those cells expressing >1 VSG by genome alignment, 85% were found to
express only one VSG by mapping to the *de novo* assembled VSGs and using the same 80%
threshold for defining monogenic expression. Further investigation revealed that in most of these
cases (95.3%) the alignment to multiple genomic VSGs was an artifact, where the assembled
VSG was not well represented within the annotated sequences of the EATRO1125 genome or
there were several VSGs with high similarity to the assembled VSG, leading to inaccurate VSG
expression quantification (Extended Data Fig. 5d). We estimate that 93.4% of cells were likely
expressing only one VSG, with no bias for multigenic VSG expression in tissue spaces
(Extended Data Fig. 5e, shades of green and Extended Data Table 2). In 5.2% of cells, reads
mapped to multiple EATRO1125 VSGs, but no VSG could be assembled (grey). While it is
impossible to distinguish between multi- and monogenic VSG expression in this set of cells, the
proportion of cells in this category did not differ between blood and tissues. The few cells that

appear to express multiple VSGs (1.05% of cells in the blood and 1.03% of cells in the tissues)
could indicate sorting doublets or, more interestingly, could represent cells mid-switch
(Extended Data Fig. 5e, red and purple). Overall, these data suggest that VSG monogenic
expression is maintained by most cells in both extravascular spaces and the blood and that the
increased VSG diversity we observe in tissue spaces is unlikely to be due to the specific
derepression of silent VSGs in tissue populations.

Frequently expressed VSGs first appear in tissue populations

The high antigenic diversity observed in tissues could serve to maintain a chronic infection. If
antigenic variation occurs relatively rarely in the bloodstream, then parasites from extravascular
spaces might serve as a source of new, antigenically distinct, VSGs. In line with this, tissue
spaces contain more "unique VSGs", those VSGs that are expressed exclusively within one
space in an infection, than the blood (Fig. 2ab). To examine the potential for tissue-resident
VSGs to contribute to antigenic variation systemically, we identified VSGs only expressed in
tissues on day 6 post-infection and analyzed whether they later appeared within the blood. The
majority (74%) of these VSGs were expressed within the blood on day 10 or 14. Analysis of
individual VSGs revealed that rare VSGs expressed at low levels exclusively in tissue spaces
also have the potential to become ubiquitously expressed within a host (Fig. 2c). In addition to a
model in which tissue spaces provide new VSGs to re-seed the blood, it is possible that tissue-
resident parasites undergo antigenic variation before blood-resident populations. Therefore, these
data could be explained by either a trigger within the tissue environment that induces parasite
switching or a differential selective pressure imposed by the tissue environment.

VSG-specific parasite clearance is delayed in tissues compared to the blood

Indeed, our data suggest that the environment within tissue spaces is distinct from the blood,
with parasite clearance occurring at different rates in each space. While parasites expressing the
initiating VSG were cleared from the blood by day 10 post-infection, they were not cleared from
tissues until at least day 14 (Fig. 3a). This suggests that VSG-specific parasite clearance from
extravascular spaces is delayed, but not abolished, compared to the blood. VSG-seq is a measure
of VSG expression at the transcript level, however. To confirm this observation at the protein
level, we performed flow cytometry on *T. brucei* cells from the blood, lungs, and gonadal fat,
using the tdTomato-expressing "triple reporter" *T. brucei* EATRO1125 AnTat1.1E cell line[35]
(Fig. 3bc). The flow cytometry analysis showed a detectable AnTat1.1-expressing tissue parasite
population at day 13 post-infection, a timepoint at which AnTat1.1$^+$ parasites were undetectable,
or nearly undetectable, in the blood. Similar to the increase in antigenic diversity we observed in
every tissue space, this delay in clearance, observed at both the RNA and protein levels, was not
tissue-specific. Thus, the immune mechanisms influencing extravascular parasite clearance
appear to be general features of extravascular spaces.

Infections initiated by tsetse fly bite also show increased diversity and delayed clearance in extravascular spaces

A benefit of starting infections with a small intravenous inoculum is that it creates convenient
and reproducible infections. In nature, however, a tsetse fly bite introduces thousands of parasites
into the skin, each expressing a single metacyclic VSG (mVSG) from a limited repertoire[36]. To
ensure that our observations held true in this more complex context, we repeated our infections
using a more natural tsetse bite infection model. Infections were initiated in 5 mice by tsetse bite
using flies infected with RUMP 503 *T. brucei* parasites[37]. We used VSG-seq to quantify VSG

expression in the blood on day 5 post-infection and the blood and tissues on day 14 post-
infection. In line with our previous observations, we found that in tsetse-initiated infections
~80% of VSGs were exclusively expressed within extravascular spaces (Fig. 4a) and tissue
populations harbored more VSGs than the blood (Fig. 4b). This demonstrates that extravascular
spaces are the primary reservoir of antigenic diversity, even when infections are initiated by fly
bite.

Parasite populations were more diverse at early time points in tsetse infections than IV
infections, likely due to the larger and more heterogeneous inoculum delivered by the fly. To
analyze VSG-specific parasite clearance in tissues, we quantified expression of the most
abundantly expressed VSG in the blood of each mouse on day 5 as well as the known mVSG
repertoire from this parasite strain, which should represent the repertoire of VSGs expressed at
the start of an infection. In both cases, we found that on day 14, tissue spaces still contained
parasites expressing the most abundant VSG and/or mVSGs, while these VSGs were expressed
by no or very few parasites in the blood (Fig. 4c,d). This suggests that in tsetse-bite-initiated
infections, as we observed in IV infections, tissue parasite populations are cleared at a slower
rate than parasites in the blood.

Delayed clearance of parasites leads to increased VSG diversity

The increased antigenic diversity in extravascular spaces could be explained by the distinct
clearance dynamics we observe, as prolonged survival in tissues could provide more time for
parasites to switch. To test whether there was a link between parasite survival and increased
diversity, we sought to interrupt parasite clearance in tissues. Because parasite clearance in the
blood coincides with the appearance of anti-VSG IgM, between days 8 and 10, and parasite
clearance in the tissues coincides with the anti-VSG IgG response, between days 10 and 14[1,12,38–
41], we hypothesized that clearance in tissues is dependent on the anti-VSG IgG response. Thus,
the loss of IgG might abrogate the clearance of tissue-resident parasites. To test this hypothesis,
we infected activation-induced cytidine deaminase (AID) knockout (AID Cre) mice[42], which
only produce IgM antibodies (Extended Data Fig. 6), and analyzed blood and tissues from days 6
and 14 post-infection by VSG-seq. As expected, clearance of the initiating VSG was severely
delayed in tissues on day 14 in AID$^{-/-}$ mice, suggesting that IgG is important, if not critical, for
the clearance of extravascular parasites. This could be explained by the fact that IgM, a bulky
pentamer, does not diffuse efficiently into tissue spaces[43,44], while IgG, a monomer, readily
diffuses. We also observed a defect in the clearance of blood-resident parasites in AID$^{-/-}$ mice
compared to the blood of WT mice (Fig. 5a). In both the blood and tissues of AID$^{-/-}$ mice, where
clearance was delayed, more VSGs were detected on day 14 post-infection compared to wild-
type (Fig. 5b), revealing a direct relationship between the timing of parasite clearance and VSG
diversity. Regardless of their local environment (intra- or extravascular), longer-lived parasite
populations generated more diverse sets of VSGs.

Discussion

The idea that antigenic variation might occur outside of the bloodstream, with extravascular
populations contributing to immune evasion within the bloodstream, is not a new one. Here,
using modern high-resolution techniques, we provide evidence for this long-standing hypothesis.
In both needle- and tsetse bite-initiated infections, we find that extravascular spaces are the
primary reservoir of VSGs, accounting for the vast majority of antigenic diversity in any
individual infection. The number of VSGs we detected in the blood matches previous

estimates[4,7,8], while the diversity in tissues is two to four times higher. Our data suggest that this
is at least partially due to slower clearance dynamics in extravascular spaces and highlight the
role that tissue spaces can play in pathogen diversification.

Although extravascular parasite populations were highly antigenically diverse, we saw no
evidence of tissue-specific VSG expression. Because parasites invade tissues efficiently before
much VSG switching has occurred, it appears unlikely that any specific VSG is required for
tissue invasion. Whether VSGs could influence parasite fitness in specific host spaces is less
clear. We measured VSG expression up to day 14 post-infection, at which point tissue-resident
populations are just beginning to diverge from one another and the blood. It is therefore possible
that, as these populations further evolve, there may be a selection for VSGs better adapted to
certain tissue spaces.

It has previously been shown that in the bloodstream alone *T. brucei* expresses more VSGs than
appear to be required for immune evasion[7]. Here we find that the expressed VSG diversity within
host tissues is even greater. While on the surface it could appear to be disadvantageous for *T.*
brucei to use so many different antigens this quickly, this striking diversity could be important
for the parasite. In natural infections, particularly in wild animals where pre-existing anti-VSG
immunity is more likely to exist, a high switch rate may be required to ensure some parasites
successfully evade the host's existing antibody repertoire. Moreover, infections in the wild can
last for months to years[14,45]. During these long infections, the large reservoirs of VSGs found in
tissues may be essential for the maintenance of a chronic infection.

Indeed, our data support the idea that the large reservoirs of antigenic diversity in extravascular
spaces contribute to systemic infection when parasites re-enter the blood after switching: rare
VSGs expressed exclusively in tissues at early time points are expressed in the blood and other
spaces later. This is also in line with another recent study which showed that blood-resident
parasites are largely non-replicative, indicating that tissue-resident parasites may be required to
re-seed the blood[46]. There is another intriguing explanation for this observation, however.
Switching in *T. brucei* is known to be semi-predictable[7,47], and it is possible that tissue-resident
parasites simply switch earlier, or more frequently, than those in the blood. In this case, the same
VSGs would arise independently in every population without any parasite movement between
spaces. While the high vascular permeability observed after the initial stages of infection
indicates that parasites likely move back and forth between the vasculature and extravascular
spaces, the fact that tissue-resident populations contain many unique VSGs suggests that blood
re-entry may represent a bottleneck for the parasite. An increased rate of switching in tissues
could be explained by a higher proportion of dividing slender form parasites in these spaces, as
has been observed in the adipose tissue[13], but we found no correlation at the population level
between PAD1 expression and increased VSG diversity. Notably, increased diversity is still
observed when the overall parasite load in tissues is lower than the blood (Extended Data Fig
3c). It is therefore exciting to speculate that some aspect of the extravascular environment
supplies a molecular or physical stimulus that promotes VSG switching.

While the timing of VSG switching could be the result of an environmental trigger, our data
suggest that the higher antigenic diversity in extravascular spaces compared to the blood can also
be explained, at least in part, by the dynamics of the immune response to *T. brucei* in each space.
In tissue spaces, we observed slower VSG-specific clearance of parasites than in the blood,
potentially providing these populations more time to undergo antigenic variation. Additionally,
newly switched parasites are still vulnerable to immune clearance by antibodies against their

previous VSG for ~29 hours, so moderate delays in immune clearance could allow more
switched parasites to survive[48]. The direct relationship we observed between the timing of
parasite clearance and antigenic diversity in AID$^{-/-}$ mice supports this model, with even small
delays in clearance showing major effects on parasite VSG diversity in both the tissues and the
blood.

Our results show that the production of IgG plays a key role in clearing parasites from tissues,
which we propose could be related to the ready diffusion of this molecule within extravascular
spaces facilitating parasite clearance. It is important to note, however, that while the difference in
timing between the anti-VSG IgM and IgG responses inspired the hypothesis that IgG might be
important for parasite clearance in tissues, our study does not prove that this difference in timing
explains the delayed clearance in tissue spaces. The local immune response to *T. brucei* is
complex, with multiple mechanisms likely to play a role in parasite detection and clearance; our
results suggest that IgG is an important player in this response. The increased VSG diversity we
observe is certainly multifactorial and could be influenced by parasite factors, such as
metabolism, division, motility, and antibody internalization, and host factors, such as
extravascular environmental stresses, the local immune response, and vascular permeability.
More research will be required to fully understand the complex nature of the *T. brucei* host-
pathogen interaction within the extravascular niche.

Altogether, our results outline a model in which *T. brucei* parasites "hide" in extravascular
spaces to generate new antigenic variants capable of exiting tissues and aiding in systemic
immune evasion. Coupled with other recent studies[9–13,46,49], this suggests a framework for the
progression and pathogenesis of *T. brucei* infections where, instead of being the primary parasite
reservoir, the blood may represent a transient population that is regularly re-seeded by
extravascular parasites. The vasculature, then, might act as a highway system for movement
between the tissue spaces and for eventual transmission back into the tsetse fly.

Interfering with the egress from or establishment within tissue spaces might be a strategy for
treating *T. brucei* infections or other infections with pathogens that rely on the distinct features of
the extravascular environment. In line with this, one recent study showed that partial inhibition
of *T. brucei* tissue invasion using antibodies against P- and E-selectins results in prolonged
survival in mice[13]. This also fits with data from another group that found immotile *T. brucei*
parasites, likely unable to invade tissues, were no longer infectious[50]. It is possible that without
the proper establishment of parasite tissue reservoirs, overall antigenic diversity is lowered,
limiting the parasite's capacity for immune evasion and leading to a decrease in parasite burden.
Defining the dynamics and variation of parasites both within and between spaces, as well as the
unique host environment within each tissue space, will be central to understanding how *T. brucei*
consistently avoids immune clearance and harnessing this mechanism for disease control.

More broadly, these data demonstrate how different environmental and immune pressures within
a host can influence pathogen diversification. This production of genetic heterogeneity within an
infection is important for many pathogen virulence processes, including establishing and
maintaining infection, facilitating immune evasion, generating drug resistance, and adapting to
different host environments. The extravascular environment plays a unique role in promoting
pathogen evolution, and *T. brucei* serves as a valuable model for understanding this aspect of the
host-pathogen interface.

References:

1. Magez, S. *et al.* The Role of B-cells and IgM Antibodies in Parasitemia, Anemia, and VSG Switching in Trypanosoma brucei–Infected Mice. *PLoS Pathog.* **4**, e1000122 (2008).

2. Cross, G. A. M., Kim, H. S. & Wickstead, B. Capturing the variant surface glycoprotein repertoire (the VSGnome) of Trypanosoma brucei Lister 427. *Mol. Biochem. Parasitol.* **195**, 59–73 (2014).

3. Müller, L. S. M. *et al.* Genome organization and DNA accessibility control antigenic variation in trypanosomes. *Nat. 2018 5637729* **563**, 121–125 (2018).

4. Hall, J. P. J., Wang, H. & David Barry, J. Mosaic VSGs and the Scale of Trypanosoma brucei Antigenic Variation. *PLoS Pathog.* **9**, (2013).

5. Hertz-Fowler, C. *et al.* Telomeric Expression Sites Are Highly Conserved in Trypanosoma brucei. *PLoS One* **3**, e3527 (2008).

6. Cosentino, R. O., Brink, B. G. & Nicolai Siegel, T. Allele-specific assembly of a eukaryotic genome corrects apparent frameshifts and reveals a lack of nonsense-mediated mRNA decay. *NAR Genomics Bioinforma.* **3**, (2021).

7. Mugnier, M. R., Cross, G. A. M. & Papavasiliou, F. N. The in vivo dynamics of antigenic variation in Trypanosoma brucei. *Science* **347**, 1470–3 (2015).

8. Jayaraman, S. *et al.* Application of long read sequencing to determine expressed antigen diversity in Trypanosoma brucei infections. *PLoS Negl. Trop. Dis.* **13**, e0007262 (2019).

9. Capewell, P. *et al.* The skin is a significant but overlooked anatomical reservoir for vector-borne African trypanosomes. *Elife* **5**, (2016).

10. Camara, M. *et al.* Extravascular Dermal Trypanosomes in Suspected and Confirmed Cases of gambiense Human African Trypanosomiasis. *Clin. Infect. Dis. An Off. Publ. Infect. Dis. Soc. Am.* **73**, 12 (2021).

11. Trindade, S. *et al.* Trypanosoma brucei Parasites Occupy and Functionally Adapt to the Adipose Tissue in Mice. *Cell Host Microbe* (2016). doi:10.1016/j.chom.2016.05.002

12. Carvalho, T. *et al.* Trypanosoma brucei triggers a marked immune response in male reproductive organs. *PLoS Negl. Trop. Dis.* (2018). doi:10.1371/journal.pntd.0006690

13. De Niz, M. *et al.* Organotypic endothelial adhesion molecules are key for Trypanosoma brucei tropism and virulence. *Cell Rep.* **36**, 109741 (2021).

14. WHO. Control and surveillance of human African trypanosomiasis: Report of a WHO Expert Committee. *WHO Tech. Rep. Ser.* 1–237 (2013).

15. Crilly, N. P. & Mugnier, M. R. Thinking outside the blood: Perspectives on tissue-resident Trypanosoma brucei. *PLOS Pathog.* **17**, e1009866 (2021).

16. Kamper, S. M. & Barbet, A. F. Surface epitope variation via mosaic gene formation is potential key to long-term survival of Trypanosoma brucei. *Mol. Biochem. Parasitol.* **53**, 33–44 (1992).

17. Seed, J. R. & Eefron, H. G. Simultaneous presence of different antigenic populations of Trypanosoma brucei gambiense in Microtus montanus. *Parasitology* **66**, 269–278 (1973).

18. SEED, J. R., EDWARDS, R. & SECHELSKI, J. The ecology of antigenic variation. *J.*
Protozool. **31**, 48–53 (1984).

19. Barry, J. D. & Emery, D. L. Parasite development and host responses during the
establishment of Trypanosoma brucei infection transmitted by tsetse fly. *Parasitology* **88 (**
Pt 1), 67–84 (1984).

20. Tanner, M., Jenni, L., Hecker, H. & Brun, R. Characterization of Trypanosoma brucei
isolated from lymph nodes of rats. *Parasitology* **80**, 383–391 (1980).

21. Turner, C. M., Hunter, C. A., Barry, J. D. & Vickerman, K. Similarity in variable antigen
type composition of Trypanosoma brucei rhodesiense populations in different sites within
the mouse host. *Trans. R. Soc. Trop. Med. Hyg.* **80**, 824–830 (1986).

22. Vickerman, K. Trypanosome sociology and antigenic variation. *Parasitology* **99 Suppl**,
S37–S47 (1989).

23. Barry, J. D. & Turner, C. M. R. The dynamics of antigenic variation and growth of
African trypanosomes. *Parasitol. Today* **7**, 207–211 (1991).

24. Engstler, M. & Boshart, M. Cold shock and regulation of surface protein trafficking
convey sensitization to inducers of stage differentiation in Trypanosoma brucei. *Genes*
Dev. **18**, 2798 (2004).

25. Turner, C. M. R. & Barry, J. D. High frequency of antigenic variation in Trypanosoma
brucei rhodesiense infections. *Parasitology* **99 Pt 1**, 67–75 (1989).

26. Salanti, A. *et al.* Evidence for the Involvement of VAR2CSA in Pregnancy-associated
Malaria. *J. Exp. Med.* **200**, 1197–1203 (2004).

27. Duffy, P. E. & Fried, M. Plasmodium falciparum adhesion in the placenta. *Curr. Opin.*
Microbiol. **6**, 371–376 (2003).

28. Jonsson, A. -B, Ilver, D., Falk, P., Pepose, J. & Normark, S. Sequence changes in the pilus
subunit lead to tropism variation of Neisseria gonorrhoeae to human tissue. *Mol.*
Microbiol. **13**, 403–416 (1994).

29. Nassif, X. *et al.* Antigenic variation of pilin regulates adhesion of Neisseria meningitidis
to human epithelial cells. *Mol. Microbiol.* **8**, 719–725 (1993).

30. Rudel, T., van Putten, J. P. M., Gibbs, C. P., Haas, R. & Meyer, T. F. Interaction of two
variable proteins (PilE and PilC) required for pilus-mediated adherence of Neisseria
gonorrhoeae to human epithelial cells. *Mol. Microbiol.* **6**, 3439–3450 (1992).

31. Virji, M. & Heckels, J. E. The role of common and type-specific pilus antigenic domains
in adhesion and virulence of gonococci for human epithelial cells. *J. Gen. Microbiol.* **130**,
1089–1095 (1984).

32. Dean, S., Marchetti, R., Kirk, K. & Matthews, K. R. A surface transporter family conveys
the trypanosome differentiation signal. *Nature* **459**, 213 (2009).

33. McWilliam, K. R. *et al.* High-resolution scRNA-seq reveals genomic determinants of
antigen expression hierarchy in African Trypanosomes. *bioRxiv* 2024.03.22.586247
(2024). doi:10.1101/2024.03.22.586247

34. Smith, J. E. *et al.* DNA damage drives antigen diversification through mosaic VSG

formation in Trypanosoma brucei. *bioRxiv* 2024.03.22.582209 (2024).
doi:10.1101/2024.03.22.582209

35. Calvo-Alvarez, E., Cren-Travaillé, C., Crouzols, A. & Rotureau, B. A new chimeric triple
reporter fusion protein as a tool for in vitro and in vivo multimodal imaging to monitor the
development of African trypanosomes and Leishmania parasites. *Infect. Genet. Evol.* **63**,
391–403 (2018).

36. Hutchinson, S. *et al.* The establishment of variant surface glycoprotein monoallelic
expression revealed by single-cell RNA-seq of Trypanosoma brucei in the tsetse fly
salivary glands. *PLOS Pathog.* **17**, e1009904 (2021).

37. Savage, A. F. *et al.* Transcript Expression Analysis of Putative Trypanosoma brucei GPI-
Anchored Surface Proteins during Development in the Tsetse and Mammalian Hosts.
PLoS Negl. Trop. Dis. **6**, 1708 (2012).

38. Schopf, L. R., Filutowicz, H., Bi, X. J. & Mansfield, J. M. Interleukin-4-Dependent
Immunoglobulin G1 Isotype Switch in the Presence of a Polarized Antigen-Specific Th1-
Cell Response to the Trypanosome Variant Surface Glycoprotein. *Infect. Immun.* **66**, 451
(1998).

39. Liu, G. *et al.* Distinct contributions of CD4+ and CD8+ T cells to pathogenesis of
trypanosoma brucei infection in the context of gamma interferon and interleukin-10.
Infect. Immun. **83**, 2785–2795 (2015).

40. Reinitz, D. M. & Mansfield, J. M. T-cell-independent and T-cell-dependent B-cell
responses to exposed variant surface glycoprotein epitopes in trypanosome-infected mice.
Infect. Immun. **58**, 2337–42 (1990).

41. Radwanska, M. *et al.* Comparative analysis of antibody responses against HSP60,
invariant surface glycoprotein 70, and variant surface glycoprotein reveals a complex
antigen-specific pattern of immunoglobulin isotype switching during infection by
Trypanosoma brucei. *Infect. Immun.* **68**, 848–860 (2000).

42. Robbiani, D. F. *et al.* Activation Induced Deaminase is required for the chromosomal
breaks in c-myc that lead to c-myc/IgH translocations. *Cell* **135**, 1028 (2008).

43. Hector, R. F., Collins, M. S. & Pennington, J. E. Treatment of Experimental Pseudomonas
aeruginosa Pneumonia with a Human IgM Monoclonal Antibody. *J. Infect. Dis.* **160**, 483–
489 (1989).

44. BARTH, W. F., WOCHNER, R. D., WALDMANN, T. A. & FAHEY, J. L. Metabolism
of Human Gamma Macroglobulins. *J. Clin. Invest.* **43**, 1036 (1964).

45. Mehlitz, D. & Molyneux, D. H. The elimination of Trypanosoma brucei gambiense?
Challenges of reservoir hosts and transmission cycles: Expect the unexpected. *Parasite*
Epidemiol. Control **6**, e00113 (2019).

46. Larcombe, S. D., Briggs, E. M., Savill, N., Szoor, B. & Matthews, K. The developmental
hierarchy and scarcity of replicative slender trypanosomes in blood challenges their role in
infection maintenance. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2306848120 (2023).

47. Morrison, L. J., Majiwa, P., Read, A. F. & Barry, J. D. Probabilistic order in antigenic
variation of Trypanosoma brucei. *Int. J. Parasitol.* **35**, 961–972 (2005).

48. Pinger, J., Chowdhury, S. & Papavasiliou, F. N. Variant surface glycoprotein density
defines an immune evasion threshold for African trypanosomes undergoing antigenic
variation. *Nat. Commun.* (2017). doi:10.1038/s41467-017-00959-w

49. Trindade, S. *et al.* Slow growing behavior in African trypanosomes during adipose tissue
colonization. *Nat. Commun. 2022 131* **13**, 1–13 (2022).

50. Shimogawa, M. M. *et al.* Parasite motility is critical for virulence of African
trypanosomes. *Sci. Rep.* (2018). doi:10.1038/s41598-018-27228-0

51. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150 (2012).

52. So, J. *et al.* VSGs Expressed during Natural T. b. gambiense Infection Exhibit Extensive
Sequence Divergence and a Subspecies-Specific Bias towards Type B N-Terminal
Domains. *MBio* **13**, (2022).

53. Lee, J.-Y. & Kitaoka, M. A beginner's guide to rigor and reproducibility in fluorescence
imaging experiments. *https://doi.org/10.1091/mbc.E17-05-0276* **29**, 1519–1525 (2018).

54. Wirtz, E., Leal, S., Ochatt, C. & Cross, G. A. M. A tightly regulated inducible expression
system for conditional gene knock-outs and dominant-negative genetics in Trypanosoma
brucei. *Mol. Biochem. Parasitol.* **99**, 89–101 (1999).

55. Shanmugasundram, A. *et al.* TriTrypDB: An integrated functional genomics resource for
kinetoplastida. *PLoS Negl. Trop. Dis.* **17**, e0011058 (2023).

56. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res.* **44**, W160–W165 (2016).

57. Moloo, S. K. An artificial feeding technique for Glossina. *Parasitology* **63**, 507–512
(1971).

58. MacLeod, E. T., Maudlin, I., Darby, A. C. & Welburn, S. C. Antioxidants promote
establishment of trypanosome infections in tsetse. *Parasitology* **134**, 827–831 (2007).

59. Barnett, S. A. THE SKIN AND HAIR OF MICE LIVING AT A LOW
ENVIRONMENTAL TEMPERATURE. *Q. J. Exp. Physiol. Cogn. Med. Sci.* **44**, 35–42
(1959).

Methods:

<u>Intravenous mouse infections and sample collection</u>

Female C57Bl/6J (WT, strain# 000664 Jackson Laboratory) or B6.129P2-
Aicda$^{tm1(cre)Mnz}$/J (AID$^{-/-}$, strain# 007770 Jackson Laboratory)[42] between 7-10 weeks old were
each infected by intravenous tail vein injection with ~5 pleiomorphic EATRO 1125 AnTat1.1E
90-13 *T. brucei* parasites[24]. Blood parasitemia was counted by tail bleed every 2 days starting on
day 4 post-infection (PI) by hemocytometer with a limit of detection of 2.22×10^5 parasites/mL.
Blood (25uL) was collected by a submandibular bleed on days 6, 10, and 14 PI and placed into
TRIzol LS. For WT mice, 4 Mice were anesthetized and perfused at days 6, 10, and 14 PI.
Infected AID$^{-/-}$ mice were anesthetized and perfused at days 6 (2 mice) and 13 (3 mice) PI. Mice
were perfused with 50mL of PBS-Glucose (0.055M D-glucose) with heparin. After perfusion,

tissues were dissected and placed immediately into 1mL of RNA Later. The heart, lungs, gonadal
fat, subcutaneous fat, brain, and skin (ear) were collected.
For flow cytometry, immunofluorescence experiments, and single-cell sorting
experiments, 7-10 week old female C57Bl/6J mice were infected by intravenous tail vein
injection with ~5 AnTat1.1E chimeric triple reporter *T. brucei* parasites which express
tdTomato[35]. Blood was collected by a submandibular bleed at designated timepoints. For flow
cytometry, mice were anesthetized and perfused on days 6 and 13 P.I. as discussed above and the
gonadal fat and lungs were harvested. For immunofluorescence experiments, perfused tissues
were collected at day 13 P.I. For single-cell sorting, blood and tissues were only collected on day
14 P.I. All animal studies were approved by the Johns Hopkins Animal Care and Use Committee
(protocol # MO22H163).
479
VSG-seq sample and library preparation
RNA was isolated from blood samples stored in TRIzol LS (ThermoFisher, 10296010)
by phenol/chloroform extraction. Tissue samples were weighed and homogenized in TRIzol, and
then RNA was isolated by phenol/chloroform extraction. RNA from each sample was DNase
treated using Turbo DNase and cleaned up with Mag-Bind® TotalPure NGS beads (Omega Bio-
Tek M1378-00). First-strand cDNA synthesis was performed using SuperScript III Reverse
Transcriptase and a primer that binds to the conserved VSG 14-mer in the 3′-UTR (5'-
GTGTTAAAATATATC-3'). Products were cleaned up using Mag-Bind® TotalPure NGS beads
(Omega Bio-Tek, M1378-01). Next, a VSG-specific PCR with Phusion polymerase
(ThermoFisher, F530L) was performed using primers for the spliced leader (5'-
ACAGTTTCTGTACTATATTG-3') and SP6-VSG 14-mer sequences (5'-
GATTTAGGTGACACTATAGTGTTAAAATATATC-3') for 25 cycles. VSG-PCR products
were cleaned up using Mag-Bind® TotalPure NGS beads and quantified using the QuBit HS
DNA kit (Life Technologies). Finally, sequencing libraries were prepared with the Nextera XT
DNA Sample Prep Kit (Illumina) using the manufacturer's guidelines, and libraries were
sequenced with 100bp single-end reads on an Illumina HiSeq 2500.
496
Tissue-load and PAD1 QPCRs
First-strand synthesis was performed with SuperScript III Reverse Transcriptase (Thermo
Fisher Scientific, 18080051) and random hexamers primers on tissue RNA samples. QPCR was
performed in triplicate using SYBR Green qPCR Master Mix (Invitrogen, 4309155). tbZFP3
primers were used to estimate parasite load in tissue samples (FW: 5'-
CAGGGGAAACGCAAAACTAA-3'; RV: 5'-TGTCACCCCAACTGCATTCT-3'). CT values
were averaged between the triplicates and parasite load per mg of tissue were estimated using a
standard curve of values from RNA isolated from known numbers of cultured parasites
(Standard curves can be found in Extended Data Fig. 3d).
For PAD1 expression quantification, RNA extraction, first-strand synthesis, and QPCR
were performed following the same methods as above. PAD1 expression was quantified by
normalizing to tbZFP3 as a control gene (same primers as above) (PAD1 primers; FW: 5'-
CAGCGGCGATTATTGCATTGG-3'; RV: 5'- AGGAAGAAGGTTCCTTTGGTC-3'). CT
values were averaged between the triplicates and samples were compared using the delta-CT
between PAD1 and tbZFP3.
512
VSG-seq analysis

Analysis of sequencing results was performed following the method we reported
previously[7], with two changes: no mismatches were allowed for bowtie alignments and each
sample was analyzed (assembly, alignment, and quantification) separately. To compare
expressed VSG sets between samples, all assembled VSGs were clustered using CD-HIT-EST[51].
VSGs with >98% identity to one another were conservatively treated as one VSG. VSGs were
then identified by their Cluster number for further analysis. Samples that had less than 100,000
successfully aligning reads to VSGs were excluded from further analysis. Four samples, 3 brain
and 1 heart, were discarded because fewer than 100,000 reads aligned to VSG (Extended Data
Fig. 3a). Downstream analysis of expression data and generation of figures was performed in R.
523
Analysis of VSG sequence motifs
To identify whether there were tissue-specific VSG sequence motifs, the similarity of N
terminal sequences from all assembled VSGs were compared. N terminal sequences were
identified using a HMMEr scan against a database curated by Cross et al[2,52]. All N termini were
compared in an all vs all blast using default parameters. All VSG pairwise comparisons with an
e-value higher than 1E-3 were considered sufficiently similar to one another for further analysis.
VSGs that were found in a given tissue were binned into that tissue group, and the distribution of
the BLAST bitscores in a given compartment was compared against the total population of
similar VSGs.
533
Flow Cytometry
Once mice were perfused, tissues were dissected and washed with HBSS (Hanks
balanced salt solution, ThermoFisher Scientific 14175095). Tissue samples were minced and
placed in DMEM (ThermoFisher Scientific, 11995065) containing either 1 mg/mL collagenase
type 1 (ThermoFisher Scientific, 17100017) for adipose fat or 2 mg/mL collagenase type 2
(ThermoFisher Scientific, 17101015) for lung samples. Hearts were dissociated using 2 mg/mL
collagenase type 2, 50U/mL DNase I, and 20U/mL Hyaluronidase. These were then incubated in
a 37°C water bath for 1 hour and briefly vortexed every 10 minutes. Next, samples were passed
through a 70μM filter and centrifuged at 2600 x g for 8 mins at 4 C, and the cell pellet was taken
for antibody staining.
Blood samples were collected by submandibular bleed and red blood cells were depleted
by magnetic-activated cell sorting (MACS) with anti-Ter-119 MicroBeads (Miltenyi Biotech,
130-049-901) following the manufacturer's protocol. Cells were pelleted and washed with HMI-
9 media.
All samples, both blood and tissues, were stained with Zombie Aqua™ dye at 1:100 in
PBS and washed with PBS following the manufacturer's protocol (BioLegend, 423101).
Samples were then stained for 10 minutes at 4°C with a rabbit anti-AnTat1.1 polyclonal antibody
diluted 1:15,000 in HMI-9 media and washed once with HMI-9 (antibody courtesy of Jay
Bangs). Then, secondary antibody staining was performed while shaking for 10 minutes at 4°C
with Anti-Rabbit IgG (H+L), F(ab')2 Fragment conjugated to Alexa Fluor® 488 fluorescent dye
(Cell Signaling Technology, 4412S). Finally, samples were washed with cold PBS and
resuspended in PBS for flow cytometry analysis. Samples were run on a Beckton Dickenson A3
Symphony flow cytometer and analysis was performed using FlowJo (version 10.6.1) (see
Extended Data Fig. 7a for gating strategy).
558
Immunofluorescence

Mice infected with AnTat1.1E chimeric triple reporter *T. brucei* parasites that express
tdTomato were sacrificed and perfused as previously described at days 6 and 13 PI. Lung, heart,
and gonadal fat were collected and fixed in 4% paraformaldehyde in PBS for 12 hours at 4 C.
Post-fixation, tissues were frozen, embedded in O.C.T. Compound (Tissue-Tek), and cut by
cryostat microtome into 10μm sections.
The following antibodies were applied to sections: rat anti-mouse CD31 (PECAM-1)
(SCBT) with goat anti-rat Fluor 488 (CST). Coverslips were mounted using ProLong Gold (Life
technologies). Tissues were imaged with 4x, 10x, and 20x objectives using a Nikon Eclipse 90i
fluorescence microscope (Nikon) and X-Cite 120 fluorescent lamp (Excelitas) with an ORCA-
ER digital CCD camera (Hammamatsu) and ImageJ v1.53 image analysis software. Image
collection and analysis followed published guidelines for rigor and reproducibility[53].
571
Serum antibody ELISA quantification
Blood (25uL) was collected by submandibular bleed on days 0, 6, 10, and 14 PI from
mice infected with ~5 pleiomorphic EATRO 1125 AnTat1.1E 90-13 *T. brucei* parasites. Serum
was isolated using serum separator tubes (BD Microtainer SST tubes, 365967). IgM and IgG
were quantified by ELISA using ThermoFisher IgM and IgG kits following manufacturer
protocols (IgG cat# 88-50400-88, IgM cat# 88-50470-88).
578
Serum sample flow cytometry on *T. brucei*
Blood was collected from 2 mice infected with AnTat1.1E chimeric triple reporter *T.*
brucei parasites, which initially express the VSG AnTat1.1, by cheek bleed. Serum was isolated
by spinning blood samples at 10,000xg for 5 mins and pipetting off the top serum layer. EATRO
1125 AnTat1.1E 90-13 *T. brucei* parasites[24] expressing the VSG AnTat1.1 and Monomorphic
Single Marker Lister427 VSG221 TetR T7RNAP bloodstream form (NR42011; LOT:
61775530)[54], which express VSG-2, were used for flow cytometry. 10^6 Parasites were stained in
duplicate while shaking for 10 minutes at 4°C using mouse serum diluted 1:100 in PBS. As a
positive control, a rabbit anti-AnTat1.1 polyclonal antibody diluted 1:15,000 in PBS was used
following the same staining procedure (antibody courtesy of Jay Bangs). Then, secondary
antibody staining was performed while shaking for 10 minutes at 4°C with Anti-mouse IgG
(H+L), F(ab')2 Fragment conjugated to Alexa Fluor® 647 fluorescent dye (Cell Signalling
Technology, 4410S) or Anti-Rabbit IgG (H+L), F(ab')2 Fragment conjugated to Alexa Fluor®
647 fluorescent dye (Cell Signaling Technology, 4414S). Finally, samples were washed with
cold PBS and resuspended in PBS for flow cytometry analysis. Samples were run on an Attune
Nxt flow cytometer (Invitrogen) and analysis was performed using FlowJo (version 10.6.1).
595
Single-cell sorting and RNA-seq library preparation
Blood and tissue samples (heart, gonadal fat, and lung) from two mice were collected
after perfusion as described above. For sorting parasites, the same tissue dissociation and flow
cytometry was performed as described above, with the exception of live/dead staining, which
was performed using propidium iodide instead of Zombie Aqua™. Samples were kept on ice as
much as possible through this process. Single live, tdTomato$^+$ *T. brucei* cells were sorted into
chilled 384-well plates for SL-Smart-seq3xpress library preparation containing lysis buffer and
an RNA spike-in control using a Beckman Coulter MoFlo™ XDP cell sorter (see Extended Data
Fig. 7b for gating strategy). For each blood and tissue sample a single plate of parasites were
sorted for a total of 370 cells per sample.

The SL-Smart-seq3xpress library preparation approach was followed as described in
McWilliam et al[33]. Briefly, single cells were lysed by incubation at 72°C for 10 minutes in 0.3 µl
of lysis buffer. Reverse transcription was done by adding 0.1µl of reverse transcription mix to
each well and incubation at 42°C for 90 minutes, followed by ten cycles of 50°C for 2 minutes
and 42°C for 2 minutes, with a final incubation at 85°C for 5 minutes. Preamplification was done
by adding 0.6µl of a mix containing primers annealing to the Spliced-Leader sequence and to a
conserved sequenced added by the oligodT primer during retrotranscription, with the following
cycling conditions: 95°C for 1 minute, 16 cycles of : 98°C for 10 seconds, 65°C for 30 seconds,
68°C for 4 minutes; and final 72°C for 10 minutes. Following, the amplified cDNA was diluted
by adding 9µl of water per well. Next, 1 µl of each well was transferred to a new plate, 1µl of
tagmentation mix was added and the plate incubated at 55°C for 10 minutes for tagmentation.
The reaction was stopped by adding 0.5µl of 0.2% SDS to each well and incubating for 5
minutes. The final index PCR was done by adding 1µl of specific index primer combinations to
each well and 1.5µl of PCR mix. The following cycling conditions were used: 72° for 3 minutes,
95°C for 30 seconds, 14 cycles of: 95°C for 10 seconds, 55°C for 30 seconds, 72°C for 1 minute;
followed by 72°C for minutes. Single-cell libraries were then pooled and purified using AMPure
XP beads at a ratio of 1:0.7. Libraries were run on a 4% non-denaturing PAGE gel and purified
according to standard polyacrylamide gel purification protocols. Purified libraries from multiple
plates were pooled and sequenced on a NextSeq 1000 sequencing platform to produce paired-end
reads of 101nt (cDNA) and 19nt (TAG+UMI read), and 8nt for the index reads.
626
Single-cell RNA-seq primary processing and VSG de novo assembly
The primary processing of the sequencing data was as described in McWilliam et al[33].
Briefly, the two reads containing the indexes (8nt each) and the one containing the TAG+UMI
(19nt) were concatenated into a 35nt read. Artifact reads containing the TAG sequence (or its
reverse complement) in the cDNA reads were filtered out with Cutadapt.
For analysis of derepression by *de novo* assembly of VSGs, the filtered reads were sorted
into individual files for each cell, and these read files were run through our VSG-Seq analysis
pipeline individually using the same parameters as described above for bulk VSG-Seq. Using
this approach, VSG open reading frames (ORFS) were assembled and quantified for each cell
individually. VSG ORFs were then clustered among all single cells (VSG clusters were not
related to previous VSG clusters from bulk VSG-seq and cannot be compared to the previous
analysis based on cluster names).
For analysis of derepression by alignment to the genome, filtered reads were mapped
with STAR (version 2.7.10a) to a hybrid fasta file combining the *T. brucei* EATRO 1125 strain
genome assembly (version 67, downloaded from TriTrypDB[55]) and the set of 10 sequences used
as RNA spike-in. The count matrix obtained was then corrected with the index hopping filtering
pipeline "scSwitchFilter" (https://github.com/colomemaria/scSwitchFilter). Only cells with at
least 500 genes detected, 1000 gene UMI transcript counts, 30 spike-in UMI counts, and 10 VSG
UMI counts were used for downstream analyses. A total of 1216 total cells fit these criteria out
of 2960 total cells sequenced. For each tissue and blood sample a single plate (370 cells) was
sequenced. For assessment of potential derepression (Extended Data Figure 5b,e), VSGs with >1
UMI count were considered expressed in a cell. Cells were considered to be monogenically
expressing a VSG (Extended Data Figure 5e) if the VSG represented ≥80% of VSG UMI counts.
For evaluating read coverage of VSGs and VSG clusters, Bowtie indexes were created
for each reference sequence then reads from a single cell were aligned using Bowtie1. Read

coverage was calculated using deepTools[56] to convert BAM alignment files to bigWig coverage
tracks. Coverage was then visualized using the ggcoverage package in R
(https://github.com/showteeth/ggcoverage).
655
<u>Tsetse fly husbandry and fly bite-inoculated mouse infections</u>
Glossina morsitans morsitans were maintained in the Yale School of Public Health
insectary at 25°C with 65-70% relative humidity under a 12h:12h light:dark photoperiod. All
flies received defibrinated sheep blood (Lampire Biologicals) every 48 hours through an artificial
membrane feeding system[57]. Newly eclosed adult female flies were administered *per os* an initial
blood meal containing 1×10^6/ml of *Trypanosoma brucei brucei* (strain RUMP 503; previously
expanded in rats) and cysteine (10µM; to increase the infection prevalence; [58]). After this single
parasite challenge, flies were maintained on normal blood every other day.
Thirty-five days post-challenge (the time it takes *T. b. brucei* to complete their
developmental cycle within the tsetse fly and become infectious to a new vertebrate host), six to
eight-week-old female C57Bl/6J mice were exposed to the bite of individual, trypanosome
challenged flies 72 hrs after the flies had taken their last blood meal. Following the consumption
of mouse blood, individual flies were microscopically dissected to confirm that their salivary
glands were infected with vertebrate-infectious metacyclic stage *T. b. brucei* (if not, another fly
was allowed to feed on the mouse until it was confirmed that an infectious fly had taken a blood
meal). Five mice were infected using this method. All experiments using mice were performed in
strict accordance with the Yale University Institutional Animal Care and Use Committee policies
(Protocol 2014–07266 renewed on March 2023).
Once mice were infected, blood parasitemia was counted by tail bleed every 2 days
starting on day 4 post-infection (PI) by hemocytometer with a limit of detection of 2.22×10^5
parasites/mL. Blood (25uL) was collected by a submandibular bleed on days 6, 10, and 14 PI and
placed into TRIzol LS. Five Mice were anesthetized and perfused at day 14 PI. Mice were
perfused with 50mL of PBS-Glucose (0.055M D-glucose) with heparin. After perfusion, tissues
were dissected and placed immediately into 1mL of RNA Later. The heart, lungs, gonadal fat,
subcutaneous fat, brain, and skin (ear) were collected. Sequencing libraries were prepared and
analyzed following the methodology described above.
To quantify the mVSG repertoire of RUMP 503, we also collected a pool of tsetse saliva
containing RUMP 503 *T. brucei* parasites. This sample was stored in TRIzol LS, RNA was
extracted, and the sample was prepared for sequencing as described above. The VSG-Seq
pipeline was used to quantify mVSG expression in the sample.
686
<u>Statistics and figures</u>
Normality was tested for all Students t-tests and Dunnet's tests analyses and can be found
in the code on the repository at https://github.com/mugnierlab/Beaver2022. Nearly all VSG
diversity measurements were found to be normally distributed, except for some samples with low
VSG counts or sample numbers. We thus assumed normality for all VSG diversity
measurements. All reported P-values have been corrected for multiple comparisons using the
Benjamini–Hochberg procedure. For all figures with boxplots, the box represents the first (25%)
and third (75%) quartiles with a line at the median. Extending lines represent the maximum and
minimum values not including outliers that are further than 1.5 times the interquartile range from
either end of the box.
697

Data availability:

Code and data for generating the analysis and figures in this paper are available at https://github.com/mugnierlab/Beaver2022. Raw sequencing data are available in National Center for Biotechnology Information (NCBI) Sequence Read Archive under accession number PRJNA858046.

Methods references:

2. Cross, G. A. M., Kim, H. S. & Wickstead, B. Capturing the variant surface glycoprotein repertoire (the VSGnome) of Trypanosoma brucei Lister 427. *Mol. Biochem. Parasitol.* **195**, 59–73 (2014).

7. Mugnier, M. R., Cross, G. A. M. & Papavasiliou, F. N. The in vivo dynamics of antigenic variation in Trypanosoma brucei. *Science* **347**, 1470–3 (2015).

25. Engstler, M. & Boshart, M. Cold shock and regulation of surface protein trafficking convey sensitization to inducers of stage differentiation in Trypanosoma brucei. *Genes Dev.* **18**, 2798 (2004).

28. Calvo-Alvarez, E., Cren-Travaillé, C., Crouzols, A. & Rotureau, B. A new chimeric triple reporter fusion protein as a tool for in vitro and in vivo multimodal imaging to monitor the development of African trypanosomes and Leishmania parasites. *Infect. Genet. Evol.* **63**, 391–403 (2018).

29. Robbiani, D. F. *et al.* Activation Induced Deaminase is required for the chromosomal breaks in c-myc that lead to c-myc/IgH translocations. *Cell* **135**, 1028 (2008).

33. McWilliam, K. R. *et al.* High-resolution scRNA-seq reveals genomic determinants of antigen expression hierarchy in African Trypanosomes. *bioRxiv* 2024.03.22.586247 (2024). doi:10.1101/2024.03.22.586247

36. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**, 3150 (2012).

37. So, J. *et al.* Trypanosomal variant surface glycoprotein expression in human African trypanosomiasis patients. *bioRxiv* 2021.09.09.459620 (2021). doi:10.1101/2021.09.09.459620

38. Lee, J.-Y. & Kitaoka, M. A beginner's guide to rigor and reproducibility in fluorescence imaging experiments. *https://doi.org/10.1091/mbc.E17-05-0276* **29**, 1519–1525 (2018).

49. Shanmugasundram, A. *et al.* TriTrypDB: An integrated functional genomics resource for kinetoplastida. *PLoS Negl. Trop. Dis.* **17**, e0011058 (2023).

50. Moloo, S. K. An artificial feeding technique for Glossina. *Parasitology* **63**, 507–512 (1971).

51. MacLeod, E. T., Maudlin, I., Darby, A. C. & Welburn, S. C. Antioxidants promote establishment of trypanosome infections in tsetse. *Parasitology* **134**, 827–831 (2007).

Acknowledgments: We would like to thank Jay Bangs for supplying us with the anti-AnTat1.1 antibody and Brice Rotureau for providing us with the chimeric triple reporter cell line. We thank Tricia Nilles and Worod Allak at the Becton Dickinson Immunology and Flow Cytometry

core at Johns Hopkins Bloomberg School of Public Health for training, support, and technical
assistance using the BD FACS Symphony A3. We would like to also thank Hao Zhang and
Joseph Margolick from the Flow Cytometry Cell Sorting Core Facility at Bloomberg School of
Public Health, Johns Hopkins University for doing FACS sorting. The facility was supported by
CFAR: 5P30AI094189-04 (Chaisson), 1S10OD016315-01 and 1S10RR13777001. Data analysis
was carried out at the Advanced Research Computing at Hopkins (ARCH) core facility
(rockfish.jhu.edu), which is supported by the National Science Foundation (NSF) grant number
OAC 1920103. AKB was supported by NIH T32AI007417. NPC was supported by NIH T32
OD011089. FR-F is supported by NIH NIGMS 1K99GM132557-01 and is an Investigator in the
Chan Zuckerberg Biohub. MRM, AKB, JES, NPC, JMCH, BB, and BZ were supported by
Office of the Director, NIH (DP5OD023065). In addition, this study was funded by the German
Research Foundation [SI 1610 / 2-2] and an ERC Consolidator Grant (SwitchDecoding
101044320) awarded to TNS. ZK was supported by a MSCA ITN Cell2Cell fellowship.

Author contributions:

Conceptualization: AKB, MRM, LMF, FRF, TNS, ROC, SA, BLW

Methodology: AKB, ROC, ZK, BLW, EOA, GMS, NPC, GYB, JMCH, JES, BZ, BB

Investigation: AKB, ROC, ZK, BLW, NPC, GYB, JH, JES

Visualization: AKB, NPC, JMCH

Funding acquisition: MRM, LMF, TNS, SA

Project administration: MRM

Supervision: MRM

Writing – original draft: AKB, MRM

Writing – review & editing: AKB, JES, GYB, BZ, MRM, FRF, LMF, TNS, ROC

Competing interests: The authors declare that they have no competing interests.

Materials and correspondence: Correspondence and requests for materials should be addressed to Monica Mugnier (mmugnie1@jhu.edu).

Fig. 1. Extravascular parasites harbor most of the antigenic diversity in an infection. (**a**)
The percentage of parasites expressing each VSG within a space. The 11 VSGs with the highest
overall expression are colored, and all other VSGs are in grey as "other". Gon. fat = gonadal fat,
s.c. fat = subcutaneous fat, P.I. = post-infection. (**b**) Stacked bar graphs from each infected
mouse representing the percentage of VSGs that were found exclusively within the blood (red),
exclusively within tissue spaces (blue), or shared by both the blood and at least one tissue
(green). (**c**) Quantification of the number of VSGs found within the blood (red) or tissue spaces
(blue) at each time point (Shapiro-Wilk normality test followed by a two-tailed Student's t-test
BH corrected). (**d**) The number of VSGs in each tissue space (Shapiro-Wilk normality test
followed by a two-tailed Dunnett's test) (For statistical tests *P<0.05, ***P<0.001).

Fig. 2. Tissue-resident parasites express a unique repertoire of VSGs during infection. (**a**)
We define "unique" VSGs as those VSGs solely found within a specific space in a mouse
(Created with BioRender.com). (**b**) The percentage of VSGs that were unique to one space
within a mouse (Shapiro-Wilk normality test followed by a two-tailed Dunnett's test, *P<0.05).
Day 6 samples were excluded from this analysis because few VSGs are expressed at this point.
Gon. fat = gonadal fat, s.c. fat = subcutaneous fat. (**c**) The expression of three representative
VSGs (cluster 294, 504, and 1831) within blood and tissue samples on days 6, 10, and 14. "n.d"
indicates that the VSG was not detected.

Fig. 3. VSG-specific parasite clearance is slower in tissues than in the blood. (**a**) The percentage of parasites expressing the initiating VSG (AnTat1.1 or VSG-421) at days 6, 10, and 14 post-infection. Tissue samples were grouped together (blue) and compared to blood samples (red) (two-tailed Wilcoxon test, ns = not significant, **P<0.01). "n.d" = not detected, gon. fat = gonadal fat, s.c. fat = subcutaneous fat. (**b**) Quantification of the number of parasites that were tdTomato positive and stained positive for AnTat1.1 by flow cytometry (n = 5 mice). The horizontal dotted line represents the limit of detection for VSG-seq. "n.d" indicates that the VSG was not detected. (**c**) Representative flow cytometry plots from tissues collected from mice infected with chimeric triple marker parasites that express tdTomato constitutively in their cytoplasm. Parasites were stained with anti-AnTat1.1 antibody.

Fig. 4. Tsetse bite-initiated infections show increased antigenic diversity and delayed immune clearance in extravascular spaces. Data from five mice infected with RUMP 503 parasites from a tsetse fly bite. (**a**) Bar graphs representing the percentage of VSGs in each mouse that were found exclusively within the blood (red), exclusively within tissue spaces (blue), or shared by both the blood and at least one tissue (green). (**b**) Quantification of the number of VSGs found within the blood (red) or tissue spaces (blue) on day 14 post-infection (Shapiro-Wilk normality test followed by a two-tailed Student's t-test BH corrected, ***P<0.001). (**c**) The percentage of parasites within each mouse expressing the most abundant VSG from the day 5 blood. (**d**) The percentage of parasites expressing one of the mVSGs found to be expressed by RUMP 503 parasites in the salivary gland of a tsetse fly.

Fig. 5. Delayed parasite clearance correlates with an increase in VSG diversity. (**a**) The percentage of parasites expressing the initiating VSG (AnTat1.1 or VSG-421) in both WT and AID[-/-] mice. "n.d" indicates that the VSG was not detected. (**b**) The number of VSGs expressed within the blood (red) and tissues (blue) of WT and AID[-/-] mice (Shapiro-Wilk normality test followed by a two-tailed Student's t-test, **P<0.01).

     Extended data:

Day 13 tissues

Heart

Lung

Gon. Fat

Hoechst   CD31   tdTomato   Merge

Extended Data Fig. 1. Immunofluorescence images of tdTomato expressing parasites (red) from cross sections of perfused tissues stained with hoechst (blue) and anti-CD31 antibody (green). TdTomato-positive parasites localized separately from CD31 lined spaces, showing that parasites are extravascular.

828 **Extended Data Fig. 2.** Distribution of bitscores across tissue compartments. The similarity of
829 VSGs detected in each tissue as measured by bitscore, a sequence similarity metric normalized to
830 the database size, allows for comparing tissue compartments with different numbers of total
831 VSGs expressed in each tissue. No statistical significance was found.
832

Extended Data Fig. 3. (a) Parasitemia of 12 mice infected with AnTat1.1E *T. brucei* counted
from tail blood by hemocytometer ("n.d." = not detectable, limit of detection of 2.22×10^5
parasites/mL). **(b)** Estimated parasite load per gram of tissue using QPCR. tbZFP3 was used as
the control and RNA from known quantities of parasites was used to make standard curves. **(c)**
The approximate total number of parasites represented in each organ. This was calculated using
the estimated number of parasites from QPCR and the recorded organ mass. For the blood and
skin, it was assumed that each mouse had 1.5mL of blood and 2.73 grams of skin[59] to estimate
the total load within these organs. **(d)** The qPCR standard curves used for each plate of samples.
These were used to estimate the number of parasites represented in each of our tissue samples
based on RNA from known parasite concentrations (cultured parasites counted using a
hemocytometer).

Extended Data Fig. 4. (**a**) A comparison of the number of reads successfully aligned in a sample and the number of VSGs observed. (**b**) A comparison of the total number of parasites and the number of VSGs found in each sample. (**c**) The correlation between PAD1 expression relative to the housekeeping gene tbZFP3 and the number of VSGs expressed at the population level for each sample.

Extended Data Fig. 5. (**a**) The number of VSG open reading frames (ORFs) detected per cell by *de novo* assembly with the VSG-Seq pipeline. In this figure, all cells sequenced were evaluated for VSG ORF assembly without any filtering (1377 total cells had a *de novo* assembled VSG). (**b**) The number of EATRO1125 VSGs detected per single cell. Cells were evaluated for VSG

expression only if they had at least 500 genes detected, 1000 gene UMI counts, 30 spike-in UMI
counts, and 10 VSG UMI counts (1216 cells fit these criteria out of 2960 total sequenced cells).
Only VSGs with >1 UMI count were considered for quantification of VSG expression. (**c**)
Alignment to the EATRO1125 genome assembly was used to quantify the fraction of VSG UMI
counts coming from the most abundant VSG gene within each cell. The dashed line represents
the 0.8 fraction of total VSG UMI counts (80%) threshold set to define monogenic expression
with one dominant VSG in a cell. Only cells that had at least 500 genes detected, 1000 gene UMI
transcript counts, 30 spike-in UMI counts, and 10 VSG UMI counts were evaluated for this
analysis (1216 total cells out of 2960 total cells sequenced). (**d**) Representative histograms of
coverage for reads from one cell. Read coverage is shown for the *de novo* assembled VSG
cluster 59 and two genomic VSG ORFs, VSG-60 and VSG-72, which represent a common
scenario that creates ambiguous VSG expression if reads are only aligned to the EATRO1125
genome. Many cells in the Mouse 1 lung sample expressed this VSG and exhibited this mapping
problem. (**e**) The VSG expression classification for each cell using both methods (alignment to
the EATRO1125 genome and *de novo* assembly). In both alignment and assembly, if a VSG
represented 80% or more of the VSG UMI counts in a cell (for genomic mapping) or 80% of the
population (for VSG-seq analysis), that cell was considered to be expressing only the dominant
VSG. Only cells that had at least 500 genes detected, 1000 gene UMI transcript counts, 30 spike-
in UMI counts, and 10 VSG UMI counts were evaluated for this analysis (1216 cells fit these
criteria out of 2960 total cells sequenced). Only VSGs with >1 UMI count were considered for
quantification by alignment to the EATRO1125 genome.
879

Extended Data Fig. 6. (a) Quantification of serum IgM and IgG concentrations in infected AID⁻
/⁻ and wildtype (WT) mice by ELISA (n = 2). (**b**) AnTat1.1 EATRO cells (expressing VSG
AnTat1.1) stained with serum, followed by an anti-IgG secondary antibody that cross-reacts with
IgM and other isotypes, from day 6 and day 14 of two mice infected with AnTat1.1 Triple-
marker cells. A polyclonal rabbit anti-AnTat1.1 antibody on cells known to be expressing
AnTat1.1 was used as a positive control. (**c**) Single-marker cells (expressing VSG-2) stained
with day 14 serum from two mice infected with AnTat1.1 Triple-marker cells. These were used
as negative controls to show that serum from infected mice specifically binds AnTat1.1 and no
other VSGs.

Extended Data Fig. 7. (**a**) Example gating strategy for flow cytometry experiments from Figure
3. Samples were first gated for tdTomato-positive cells, which represent *T. brucei* parasites
expressing tdTomato in their cytoplasm. Then live cells were gated based on live/dead Zombie
Aqua™ staining. Finally, quadrants were placed around AnTat1.1-A488 positive and negative
cells. (**b**) Gating strategy for single-cell sorting into 384-well plates for the SL-Smart-seq3xpress
platform. This example is from the blood of mouse 2. Singlet parasites were selected and then
tdTomato positive, PI (propidium iodide) negative *T. brucei* cells were sorted into single cells.

Summary table of the flow cytometry counts for tdTomato positive cells stained with an anti-AnTat1.1 antibody

Biological experiment	Mouse	Tissue	Day	Count of AnTat1.1 + cells	Count of AnTat1.1 - cells	Total number of cells	Percent of cells AnTat1.1+
1	1	blood	6	20394	486	20880	97.67%
1	1	gon. fat	6	411	32	443	92.78%
1	1	lung	6	817	49	866	94.34%
1	2	blood	6	42969	1458	44427	96.72%
1	2	blood	13	1	19922	19923	0.01%
1	2	gon. fat	13	0	250	250	0.00%
1	2	lung	13	573	4460	5033	11.38%
2	3	blood	6	19737	270	20007	98.65%
2	3	gon. fat	6	220	4	224	98.21%
2	3	lung	6	1464	55	1519	96.38%
2	4	blood	6	39373	592	39965	98.52%
2	4	gon. fat	6	379	10	389	97.43%
2	4	lung	6	676	30	706	95.75%
2	5	blood	6	49406	600	50006	98.80%
2	5	blood	13	1	19885	19886	0.005%
2	5	gon. fat	13	31	4982	5013	0.62%
2	5	lung	13	318	29889	30207	1.05%
2	6	blood	6	10317	31	10348	99.70%
2	6	blood	13	0	1001	1001	0.00%
2	6	gon. fat	13	3	781	784	0.38%
2	6	lung	13	265	29733	29998	0.88%
3	7	blood	6	65135	249	65384	99.62%
3	7	gon. fat	6	69	3	72	95.83%
3	7	lung	6	275	182	457	60.18%
3	8	blood	6	49239	632	49871	98.73%
3	8	gon. fat	6	71	0	71	100.00%
3	8	lung	6	770	137	907	84.90%
3	9	blood	6	49758	147	49905	99.71%
3	9	blood	13	3	5042	5045	0.06%
3	9	gon. fat	13	0	1537	1537	0.00%
3	9	lung	13	403	18944	19347	2.08%
3	10	blood	6	49286	642	49928	98.71%
3	10	blood	13	14	20204	20218	0.07%
3	10	gon. fat	13	33	7034	7067	0.47%
3	10	lung	13	187	38096	38283	0.49%

900 **Extended Data Table 1.** Flow cytometry counts of tdTomato-positive parasites from infections
901 with triple-marker *T. brucei* parasites. Parasites were collected from the blood and dissociated
902 tissues of 10 mice on either day 6 or 13 post-infection and stained with an anti-AnTat1.1
903 polyclonal antibody. Here we report the raw number of live tdTomato-positive parasites that
904 were Antat1.1 positive and negative in each sample. These data were used to generate Fig. 3b.
905

Summary table of single cell sequencing results

Mouse	Tissue	Total cells sequenced	Cells passing QC cutoffs	Average successfully aligning UMI counts per cell	Cells with a de novo assembled VSG ORF	1 VSG by alignment and 1 VSG by de novo assembly	>1 VSG by alignment and 1 VSG by de novo assembly	1 VSG by alignment and no VSG by de novo assembly	1 VSG by alignment and >1 VSG by de novo assembly	>1 VSG by alignment and >1 VSG by de novo assembly	>1 VSG by alignment and no VSG by de novo assembly
						Number of single cells					
	blood	370	238	5736	227	93	91	16	4	1	33
Mouse 1	gon. fat	370	227	4275	228	148	50	13	4	0	12
	heart	370	162	5375	201	125	36	1	0	0	0
	lung	370	123	2952	156	26	88	2	1	0	6
	blood	370	92	3572	148	50	35	5	1	0	1
Mouse 2	gon. fat	370	151	3997	133	106	16	22	1	1	5
	heart	370	156	4699	188	99	46	3	1	2	5
	lung	370	67	5097	96	42	23	0	0	0	2
Totals		2960	1216	4462	1377	689	385	62	12	4	64

Extended Data Table 2. Summary of single cell sequencing results. *T. brucei* single cells were
sorted from blood and tissues on day 14 post-infection from 2 mice and sequenced using the SL-
Smart-seq3xpress platform. For *de novo* assembly of VSGs, all cells were considered without
filters, but for all other columns only cells that had at least 500 genes detected, 1000 gene UMI
counts, 30 spike-in UMI counts, and 10 VSG UMI counts were considered. Additionally, only
VSGs that had >1 UMI counts when aligning to the EATRO1125 genome were considered to be
detected.
913

Supplementary information:

Supplementary Data 1: A summary of VSG expression in each cell by both genome alignment
and *de novo* assembly, including categorization for Extended Data Figure 5e.

Supplementary Data 2: The EATRO1125 VSG read alignment in each cell that meets QC
cutoffs (at least 500 genes detected, 1000 gene UMI counts, 30 spike-in UMI counts, and 10
VSG UMI counts) and has >1 VSG UMI count. Raw alignment count tables and VSG count
tables that include unfiltered cells can also be found at
https://github.com/mugnierlab/Beaver2022.

Supplementary Data 3: The results from *de novo* assembly of VSGs in each cell. The Trinity
name for each ORF and the top VSG blast hit for each assembled VSG ORF is reported,
including the proportion of VSG expression accounted for by that VSG. For each cell, a
dominant VSG is identified if it represented over 80% of the VSG expressed in the cell and
"NA" denotes that there is not a dominant VSG in that cell. VSG assembly was attempted in all
cells without any previous QC filtering. Here, all VSG assemblies are reported without further
filtering and the sequences can be found at https://github.com/mugnierlab/Beaver2022.

932