Group B Streptococcus Adaptation Promotes Survival in a Hyperinflammatory Diabetic Wound Environment

Rebecca A. Keogh[1], Amanda L. Haeberle[1], Christophe J. Langouët-Astrié[2], Jeffrey S. Kavanaugh[1], Eric P. Schmidt[2], Garrett D. Moore[4], Alexander R. Horswill[1,3]* and Kelly S. Doran[1]*.

[1]Department of Immunology and Microbiology, University of Colorado Anschutz, Aurora, CO, USA

[2]Department of Medicine- Pulmonary Sciences and Critical Care, University of Colorado Anschutz, Aurora, CO, USA

[3]Department of Veterans Affairs Eastern Colorado Healthcare System, Aurora, Colorado, USA.

[4]Department of Orthopedics, University of Colorado, Anschutz Medical Campus, Aurora, CO, USA

*Corresponding authors: Alexander R. Horswill and Kelly S. Doran, Department of Immunology and Microbiology, University of Colorado School of Medicine, Anschutz Medical Campus 12800 East 19th Avenue Aurora, CO, USA. Phone: (303) 724-3539, Fax: (303) 724-4226, E-mail: kelly.doran@cuanschutz.edu

Abstract

Diabetic wounds have poor healing outcomes due to the presence of numerous pathogens and a dysregulated immune response. Group B *Streptococcus* (GBS) is commonly isolated from diabetic wound infections, but the mechanisms of GBS virulence during these infections have not been investigated. Here, we develop a murine model of GBS diabetic wound infection, and using dual RNA-sequencing, demonstrate that GBS infection triggers an inflammatory response. GBS adapts to this hyperinflammatory environment by upregulating virulence factors including those known to be regulated by the two-component system *covRS*, such as the surface protein *pbsP*, and the *cyl* operon which is responsible for hemolysin/pigmentation production. We recover hyperpigmented/hemolytic GBS colonies from the murine diabetic wound which we determined encode mutations in *covR*. We further demonstrate that GBS mutants in *cylE* and *pbsP* are attenuated in the diabetic wound. This foundational study provides insight into the pathogenesis of GBS diabetic wound infections.

37   **Teaser**: The Fight for Survival by the Bacterium Group B Streptococcus in the Diabetic Wound.

Introduction

Diabetes is a multifaceted metabolic disease that is estimated to affect more than 500 million individuals worldwide by 2035 (1). One of the most common complications in diabetic individuals is the development of wounds such as foot ulcers, which an estimated 19-34% of all patients will develop in their lifetime (2). Normal wound healing progresses in four overlapping stages; (i) hemostasis, (ii) inflammation, (iii) proliferation and (iv) remodeling (3). However, diabetic wounds are often stalled in a state of chronic inflammation, and cannot progress through the later stages of healing (4–6). A hallmark characteristic of chronic inflammation is excess infiltration of neutrophils and macrophages. Leukocyte abundance can be harmful to wound healing due to high production of reactive oxygen species and serine proteases, which degrade structural proteins of the extracellular matrix therefore limiting proliferation and remodeling (7). In addition, diabetic wounds are colonized with numerous bacterial pathogens that trigger leukocyte influx and exacerbate poor healing outcomes (8).

Some of the most common species isolated from chronic wounds are *Staphylococcus aureus, Pseudomonas aeruginosa* and *Streptococcus agalactiae*, also known as Group B *Streptococcus* (GBS) (9–12). *S. aureus* is responsible for 76% of all skin and soft tissue infections and is found in wounds of both non-diabetic and diabetic individuals (11, 13). Thurlow et al. has shown that elevated glucose in hyperglycemic abscesses leads to increased virulence potential of *S. aureus*, and is dependent on two glucose transporters (14). In addition, specific *S. aureus* factors such as the proteases ClpX, ClpP and ClpC were shown to contribute to *S. aureus* skin infection in diabetic mice (15). *P. aeruginosa* is also a commonly isolated bacterium from wounds of both non-diabetic and diabetic individuals (11, 16). *P. aeruginosa* infection of chronic wounds has been linked to enhanced biofilm formation and activation of matrix metalloproteases that degrade

extracellular matrix molecules (17, 18). Conversely, GBS is rarely identified from non-diabetic

wounds, but is being increasingly identified in diabetic individuals (19, 20). While some work has

characterized *S. aureus* and *P. aeruginosa* in diabetic wound infections, no previous studies have

investigated GBS pathogenesis in the diabetic wound or on the skin in general. In addition, little

is known about the mechanisms by which bacteria survive inflammation in the diabetic wound

environment.

GBS is an opportunistic pathogen that is a leading cause of neonatal invasive infections.

Approximately 20-30% of pregnant women are colonized with GBS in the vaginal tract, and can

transmit this bacterium to the neonate *in utero* or during birth (21, 22). Neonatal infections can

lead to the development of invasive diseases like sepsis, pneumonia and meningitis, with a

mortality rate of 10-15% (23). In addition to neonatal infections, GBS infection of non-pregnant

adults, including the immunocompromised is on the rise, with over two-thirds of invasive GBS

disease occurring in adults (24). Of these individuals, the most common comorbidity is diabetes,

which is present in 20-25% of nonpregnant adults with GBS (24). Despite this, the majority of

GBS studies have focused on vaginal colonization and neonatal disease; and the mechanisms of

GBS pathogenesis in diabetic infection have not been examined.

Numerous reports have identified mechanisms of GBS colonization that contribute to

vaginal persistence, as well as mechanisms of invasive infection such as systemic disease and

meningitis. GBS produces multiple virulence factors, including the β-hemolysin/carotenoid

pigment (encoded by *cylE*), various surface adhesins such as pili, serine-rich repeat (Srr) proteins

and BspC that bind host extracellular matrix proteins (ECMs) and receptors to promote adherence

and invasion, as well as immune evasion factors such as the polysaccharide capsule, that allows

the bacterium to resist phagocytic killing and persist in the human host (25–29). Ten capsular

serotypes have been identified, with serotypes Ia, Ib, II, III and V being the most common

serotypes associated with disease (30). The transition from a commensal lifestyle to infection is

controlled by multiple regulators such as transcription factors and two-component systems (TCS)

that coordinate the expression of these virulence factors in the host. TCS typically encode a sensor

histidine kinase which recognizes a stimulus and a cognate response regulator which can be

activated to control gene expression (31). GBS encodes approximately 21 unique TCS, which is

more than other Streptococcal species such as *Streptococcus pneumoniae* and *Streptococcus*

pyogenes, suggesting the importance of these regulators in this opportunistic pathogen (32, 33).

Perhaps the most well-studies TCS in GBS is CovRS, encoding a sensor histidine kinase (*covS*)

and a response regulator (*covR*). CovR represses the expression of multiple GBS virulence factors

including fibrinogen and plasminogen binding proteins and the β-hemolysin (34). CovR regulation

has been shown to be important for GBS vaginal colonization and development of invasive disease

(35, 36).

Here, we describe for the first time the development of a murine model of GBS diabetic

wound infection to begin to assess host-pathogen interactions in this environment. We find that

GBS infection in the diabetic wound promotes a hyperinflammatory response with increased

cytokine production and neutrophil degranulation. GBS responds to the inflammatory environment

with the upregulation of numerous virulence factors such as *pbsP* and *cylE,* which are necessary

for full GBS virulence in diabetic infection and acquiring mutations in *covR* during diabetic wound

infection. Taken together, these results present numerous strategies by which GBS adapts to the

hyperinflammatory diabetic wound environment and provides insight into the pathogenic

mechanisms of GBS diabetic wound infection.

Results

Development of a Murine Model of GBS Diabetic Wound Infection.

We developed a murine model of GBS diabetic wound infection using $Lepr^{db}$ mice. These

mice harbor a mutation in the leptin receptor gene which results in obesity and mimics type 2

diabetes (37, 38). Mice were wounded with a 6 mm biopsy punch on their backs, infected with 1

x 10^7 CFU of GBS and wrapped with an adhesive to allow efficient bacterial inoculation. Once

the adhesive was removed, and animals were left for additional time before sacrifice, tissue

collection and bacterial enumeration (**Fig 1A**). All mice exhibited non-fasting glucose levels above

250 mg/dL to be considered diabetic and levels under 200 mg/dL for non-diabetic controls (**Fig

1B**). Following GBS infection, diabetic animals had significantly larger wounds at the time of

sacrifice than wild-type (WT) C57Bl/6J controls with the average wound size of non-diabetic mice

being .097 cm^2 and diabetic being .403 cm^2 (p-value= .<.0001, standard deviations .047 and .177

respectively) (**Fig 1C,D**). In addition, we utilized three clinically relevant GBS strains representing

the three of the five most prominent serotypes associated with disease worldwide; A909 (serotype

Ia), COH1 (serotype III) and CJB111 (serotype V) (30). Regardless of strain, we found that even

at an early time point of infection (one day after adhesive removal, four days after initial infection)

there were significantly more bacteria recovered from the wounds of diabetic animals in

comparison to the non-diabetic (33.9-fold for A909, 21.6-fold for COH1, and 2.6-fold for CJB111)

(**Fig 1E**). We also performed experiments on both male and female mice and saw no significant

differences in CFU recovered between sexes (**Supplemental Fig 1**), thus all subsequent infections

were performed in female mice. In addition to GBS, we have recovered colonies of *Enterococcus*

faecalis and *Staphylococcus xylosus* from wound homogenates. These species were only found in

select mice and at low levels (> 10^2 CFU/g wound). These data support clinical findings that GBS

establishes infection in the diabetic wound environment (19, 39), and provide a relevant model for

further investigation.

Further, we obtained 27 clinical isolates of GBS recovered from diabetic wounds of human

patients, and determined the molecular serotype of each isolate using primers specific to *cps* loci

(40). Serotypes Ia, Ib, II and III were confirmed by flow cytometry as described by Burcham et al.

using monoclonal antibodies against capsule (41). The most prominent serotype was Ia, which

represented one third of all isolates recovered (33.33%). The remaining isolates were serotype V

(22.22%), II (22.22%), III (11.11%) and Ib (7.41%) with one isolate that was unable to be typed

by either PCR or flow cytometry (**Supplemental Table 1**).

The hyperinflammatory wound environment in diabetic mice is more susceptible to GBS

persistence.

To determine the host response during GBS infection, we performed dual RNA-sequencing

of both diabetic and non-diabetic infected animals in comparison to animals inoculated with a PBS

control. Principal component analysis (PCA) plots demonstrate that diabetic and non-diabetic

animals clustered more similarly with infected vs. uninfected animals also clustering closely (**Fig**

2A). We utilized Reactome pathway analysis as well as gene set enrichment analysis (GSEA) to

link upregulated genes to known cellular pathways (42). Transcriptomics performed on uninfected

wounds support the current literature that *lepr*^{db} mice promote upregulation of genes and pathways

involving inflammation such as the interferon alpha response, interferon gamma response and

inflammatory response in comparison to C57Bl/6J controls (**Supplemental Fig 2**) (43, 44).

We next examined the transcriptomes of diabetic and non-diabetic mice infected with GBS

to determine if the host response was altered in diabetic infection. A total of 326 transcripts were

significantly upregulated, and 275 were significantly downregulated in diabetic infection in

comparison to non-diabetic (Log2 fold-change > 2, FDR adjusted p-value< .05). Diabetic animals

infected with GBS significantly upregulated genes involved in neutrophil degranulation, activation

of matrix metalloproteases and antimicrobial peptides while downregulating genes involved in

keratinization (**Fig 2B**, **Supplemental Table 2**).Pathway analysis of these genes revealed that GBS

infection of diabetic wounds leads to the significant upregulation of numerous pathways including

the interferon gamma response, the inflammatory response, and the reactive oxygen species

pathway (**Fig 2C**). Enzyme-linked immunosorbent assays (ELISAs) were then performed on tissue

homogenates from diabetic and non-diabetic animals infected with GBS. Wound tissues from

diabetic mice had significantly higher abundance of the neutrophil chemoattractant KC

(homologous to human CXCL1) and myeloperoxidase (MPO) and a marked increase in elastase

(ELANE) in comparison to non-diabetic (**Fig 2D-F**).

Diabetic animals upregulate inflammatory pathways and immune cell recruitment during

GBS infection.

After confirming that GBS survives better in the diabetic wound, we next sought to

determine how infection with GBS affected gene expression specifically in diabetic mice. When

comparing the transcriptome of diabetic animals infected with GBS compared to PBS controls,

146 transcripts were significantly upregulated, and 63 were significantly downregulated (Log2

fold-change > 2, FDR adjusted p-value< .05) with some of the most highly upregulated transcripts

including the subunits of calprotectin (s1008a and s100a9), myeloperoxidase (*mpo),* the

cathelicidin antimicrobial peptide (*camp*) and elastase (*elane*) (**Fig 3A**). Significantly upregulated

pathways in GSEA included interferon gamma response, TNF alpha signaling via NFkB,

complement, inflammatory response and reactive oxygen species (**Fig 3B**). NFkB activation has

been linked to an amplified inflammatory state in chronic wounds due to increased production of

inflammatory cytokines (7). We then ran our significantly dysregulated genes through Reactome

and found some of the most highly upregulated pathways were neutrophil degranulation, IL-10

signaling and the formation of a fibrin clot (**Supplemental Table 3**). Interestingly, there were no

pathways significantly downregulated in GSEA analysis, however, striated muscle contraction and

myogenesis were both enriched in Reactome analysis (**Fig 3B, Supplemental Table 3**).

The upregulation of inflammatory pathways led us to hypothesize that GBS triggers

increased inflammation in the diabetic wound. Mice infected with GBS lost significantly more

weight over the course of infection than uninfected controls, and none of the uninfected mice had

any GBS recovered from wound tissue (**Fig 3C,D**). As mentioned above we used ELISA to

determine that GBS infected wounds had significantly greater abundance of the neutrophil

chemoattractant KC further supporting that GBS infection triggers the recruitment of neutrophils

to the site of infection (**Fig 3E**). Finally, we determined that the abundance of neutrophil

components elastase, myeloperoxidase and a subunit of calprotectin are significantly higher during

GBS infection (**Fig 3F-H**). H&E staining of wound tissue sections demonstrate a gross difference

in tissue architecture from GBS infected mice including thickening of the epithelial layer and

inflammatory infiltrate (**Fig 3I**). Collectively, these data suggest that GBS promotes inflammation

in the already highly inflammatory diabetic wound environment and confirms the upregulation of

these pathways and genes observed in our RNA-seq analysis (**Fig 3A,B**, **Supplemental Table 3**).

GBS transcriptome in the diabetic wound.

To characterize the bacterial response in the diabetic wound environment, we performed

RNA-sequencing on GBS isolated from non-diabetic and diabetic wounds in comparison to GBS

grown *in vitro* (input control). PCA plots indicate that each group clusters independently, with the

largest variation being between the input control bacteria and the bacteria isolated from the mice

(**Fig 4A**). A total of 974 transcripts were significantly altered (minimum fold change of 3, FDR

adjusted p-value <.05) when comparing GBS isolated from diabetic infection in comparison to the

input control with 461 being upregulated and 513 downregulated (**Fig 4B**). Further analysis of

genes with altered regulation during diabetic infection revealed multiple housekeeping genes were

downregulated during GBS infection including ribosomal subunit genes *rpsC*, *rplV* and *rpsE*. (**Fig**

4C). In addition, multiple genes with predicted roles in lipid transport were downregulated in

diabetic infection. Conversely, numerous virulence associated genes were upregulated in

comparison to the input control (**Fig 4C**, **Table 1**). Some of the most highly upregulated transcripts

in diabetic infection include the gene encoding the surface plasminogen binding protein PbsP, the

cyl operon encoding GBS hemolysin and pigment, quorum sensing peptide pheromone *shp2*,

nuclease and protease encoding genes such as *nucA* and *clpL*, as well as various predicted effectors

of the type VII secretion system (**Table 1**.) Interestingly, many of these factors are known to be

part of the core regulon of the CovRS TCS (45–47).

We next compared the bacterial transcriptome from GBS recovered from diabetic wounds

vs. non-diabetic wounds. A total of 166 genes had significantly altered expression in this

comparison, with 75 genes being upregulated and 91 being downregulated in diabetic infection in

comparison to non-diabetic (**Fig 4B,D**). Multiple genes involved in sugar transport were

downregulated in bacteria recovered from diabetic wounds, as well as genes involved in metal

transport of iron, zinc, and manganese (**Supplemental Table 4**). Conversely, many genes involved

in iron export were upregulated in bacteria recovered from diabetic wounds (**Supplemental Table**

4).

The CovRS regulon contributes to diabetic wound infection.

We observed when plating GBS recovered from the diabetic wound that many colonies

were hyperpigmented (**Fig 5A**). These phenotypes were stable, and colonies were hyperhemolytic

when plated on sheep's blood agar (**Fig 5B**). Interestingly, we have only observed this phenotype

in colonies recovered from diabetic wounds with ~60% (7/12) of diabetic wound homogenates

containing a hyperpigmented colony compared to 0% (0/12) in non-diabetic wounds. We next

infected mice with GBS and sacrificed either 24 h or 1 week after adhesive removal and noticed a

marked increase in the total number of pigmented colonies recovered over time (**Fig 5C**)

suggesting that *covR* mutations are selected for in the diabetic wound, specifically.

Hyperpigmentation and hyperhemolysis in numerous GBS strains has been attributed to mutations

in the *covRS* TCS, due to the subsequent de-repression of the *cyl* operon which is linked to GBS

hemolysis and pigment production (25, 48–52). We surveyed the *covR* locus from

hyperpigmented/ hyperhemolytic strains recovered from 13 different mice for single nucleotide

polymorphisms (SNPs) and observed multiple SNPs resulting in mutations in the *covR* locus. Of

these, 100% of the colonies encoded *covR* mutations with 8/13 encoded for amino acid

substitutions in the receiver domain of CovR, 3/13 containing single-nucleotide deletions and 2/13

encoding insertions or deletions (**Fig 5C, Supplemental Table 5**). Nonpigmented colonies were

also isolated and sequenced from each mouse containing a hyperpigmented colony and none of

the nonpigmented colonies had mutations in *covR*. We hypothesized due to the upregulation of the

cyl operon in diabetic wound infection and the spontaneous *covR* mutations *in vivo*, that a *ΔcylE*

mutant would be attenuated in diabetic wound infection. In confirmation, we infected diabetic

mice with the *ΔcylE* mutant strain and observed a significant decrease in bacteria recovered from

wound tissue compared to WT infected mice (**Fig 5D**). Further, wounds from *cylE* infected mice

had a significant reduction in the abundance of the chemokine KC (CXCL1) (**Fig 5E**). As

expected, we saw no significant differences in bacterial recovery when comparing a clean deletion

$\Delta covR$ mutant strain to our WT *in vivo* (**Supplemental Fig 3**), likely due to the fact that WT GBS

is acquiring *covR* mutations in the wound.

The surface protein PbsP contributes to diabetic wound formation via adherence to the skin.

The most highly upregulated gene during GBS diabetic wound infection encodes the

surface plasminogen binding protein PbsP (**Table 3**). Interestingly, *pbsP* has been shown to be

repressed by CovR in multiple GBS strains (26, 34, 53). In addition, PbsP is known to bind

multiple extracellular matrix components such as plasminogen and fibrinogen (26) and contribute

to adherence to colon epithelial and brain endothelial cells (26, 54, 55) We therefore hypothesized

that PbsP contributes to diabetic wound infection by promoting adherence to the skin. A $\Delta pbsP$

mutant is significantly attenuated in diabetic wound infection in comparison to WT (**Fig 8A**).

Further, $\Delta pbsP$ mutant infected tissues have significantly less abundance of neutrophil markers

MPO, elastase and calprotectin, which are all upregulated during GBS diabetic wound infection

(**Table 2**) (**Fig 8B-D**). We next tested the ability of the $\Delta pbsP$ mutant to adhere to skin cells *in*

vitro. Using a cell culture line of immortalized keratinocytes (NTERTs), we found that the $\Delta pbsP$

mutant had a ~50% reduction in adherence to skin epithelial cells that could be restored by

complementing the mutant with WT *pbsP* (**Fig. 8E**) (56). Additionally, we tested whether PbsP

or CylE contributes to GBS infection of non-diabetic wounds and found no significant difference

in bacterial recovery (**Supplemental Fig 4**). These data suggest that PbsP contributes to diabetic

wound infection and direct interaction with skin keratinocytes.

Discussion

A hallmark characteristic of diabetic individuals is an altered neutrophil response due to persistent hyperglycemia. Neutrophils from diabetic individuals are often pro-inflammatory, exhibiting increased neutrophil extracellular trap (NET) formation, pro-inflammatory cytokine production and extracellular ROS generation (1, 57). Additionally, decreased apoptosis, neutrophil migration and intracellular ROS production have been shown in diabetic individuals, leading to impaired bacterial killing (58). Persistent inflammation and decreased bacterial clearance are detrimental to diabetic individuals who develop wounds, as resolution of inflammation is paramount to proper wound healing (59). GBS is one of the most commonly isolated pathogens from diabetic wounds, and has been shown to promote inflammation and neutrophil influx during lung infection as well as during sepsis (11, 50, 60). Here, we have utilized dual RNA-sequencing to determine the transcriptional consequences of GBS infection of non-diabetic and diabetic wounds on both the host and pathogen. We demonstrate that GBS exacerbates inflammation in the already hyperinflammatory diabetic wound environment and identify the first mechanisms by which GBS adapts to this hyperinflammation (**Fig. 7**). The data herein are the first to provide insight into GBS pathogenesis in this clinically relevant model of infection.

We demonstrate that GBS burden is significantly greater in diabetic wounds than non-diabetic, and that these data remain consistent regardless of GBS capsular serotype. GBS infection of diabetic wounds resulted in significant upregulation of pathways involved in the inflammatory response and ROS production in comparison to diabetic wounds without GBS. These findings are particularly interesting as diabetic animals already exhibit significant upregulation of these pathways in comparison to non-diabetic controls, suggesting that the presence of GBS exacerbates inflammation, particularly neutrophil recruitment and activation, during diabetes. We speculate

that GBS promotes inflammation as a strategy to stall wound healing and closure. The presence of

microbes has been linked to neutrophil influx in chronic wounds, but studies on specific bacterial

factors that assist in promoting and surviving inflammation are limited.

An additional hypothesis is that the altered neutrophil response in diabetic individuals is a

large reason GBS can survive the diabetic wound environment. Work by Thurlow et al.

demonstrated that diabetic mice infected with *S. aureus* had impaired phagocyte function including

decreased respiratory bursts and increased glucose available for the bacterium (14). While the

contribution of glucose to GBS pathogenesis is not completely understood, it is known that

increased glucose leads to thickening of the GBS capsule (61), which may contribute to virulence

or survival in the wound. It is also possible that additional changes in neutrophil function change

host nutritional immunity with respect to metal availability and sequestration. While we see

increased calprotectin in diabetic wounds infected with GBS, we do not know if it is functional in

metal sequestration. Future studies on metal availability in the diabetic wound will be critical in

understanding how host nutritional immunity may impact GBS persistence.

Our bacterial RNA-sequencing revealed that GBS upregulates numerous virulence factors

during diabetic wound infection in comparison to those grown in laboratory media. The most

highly upregulated GBS gene in diabetic wound infection encodes for the surface plasminogen

binding protein PbsP, which binds to extracellular matrix components such as plasminogen and

fibrinogen (26). Plasminogen binding proteins are known to bind plasminogen, and, in the

presence of a tissue activator, activate plasminogen into plasmin (62). While plasminogen is

crucial in early wound healing to promote inflammation, it's resolution is necessary to progress

tissue healing into the proliferation phase (63, 64). A Δ*pbsP* mutant is attenuated in multiple GBS

models of infection including in vaginal colonization and meningitis (54, 65). Furthermore, Lentini

et al. demonstrated that brain tissue homogenates from mice infected with a Δ*pbsP* mutant had a

significant reduction in MPO, TNFα and IL-1β abundance in comparison to WT infected mice

(54). Here, we demonstrate that PbsP is required for GBS diabetic wound infection, and that Δ*pbsP*

mutant infected wounds have significantly less MPO, elastase and calprotectin than those infected

with WT. We hypothesize that PbsP binding to host plasminogen may be necessary for GBS

mediated inflammation via increased cytokine and chemokine signaling as well as increased

plasmin production that degrades host matrix components necessary for proliferation and

remodeling of the skin. Further work on the exact mechanism of PbsP mediated inflammation is

being elucidated.

We also investigated the GBS hemolysin/ pigment encoded by the *cyl* operon, which was

highly upregulated during diabetic infection and is known to be part of the CovRS regulon (50,

51). Studies on the *cylE* component of this operon have shown that CylE contributes to GBS

pathogenesis in murine models of sepsis and lung infection, and that induction of the neutrophil

chemoattractant IL-8 was significantly reduced in A549 cells infected with a *cylE* mutant (50, 51).

Diabetic wounds from mice infected with a Δ*cylE* mutant had significantly fewer bacteria

recovered than those infected with WT GBS, coupled with significantly less abundance of the

neutrophil chemoattractant KC (CXCL1). CXCL1 is known to attract neutrophils to the site of

infection, and the abundance of CXCL1 in chronic wounds is high in comparison to burn or

surgical wounds (66). We therefore hypothesize that production of the GBS hemolysin/pigment is

necessary for GBS to trigger neutrophil influx into the diabetic wound environment.

Other factors that were highly upregulated during diabetic wound infection included the

nuclease *nucA*, and the quorum sensing (QS) genes *shp2* and *rovS*. NucA is the major GBS

nuclease and has been shown to degrade NETs as well as contribute to virulence in the lung (67).

We speculate that the upregulation of *nucA* might be in response to the hyperinflammatory

environment, and in particular, the increased NET production which is associated with diabetic

wounds (1, 57). Increased nuclease activity could therefore assist GBS in evading NETs and

persisting in the diabetic wound environment, although further experiments are warranted to

determine this. Increased production of *shp2* and *rovS* suggests a need for bacterial cell-to-cell

communication in the diabetic wound environment. The need for QS systems in wounds is well-

documented in both *S. aureus* and *P. aeruginosa*, which both rely on QS systems for full virulence

and persistence in chronic wounds (68, 69). While GBS *shp2* and *rovS* are required for GBS to

persist in the murine liver and spleen (70), no work has investigated whether this QS system is

critical to survival in the wound environment. Additionally, GBS *shp2* and *rovS* have a known role

in interspecies communication which could highlight the importance of these genes in a

polymicrobial diabetic wound environment (71).

When comparing the GBS transcriptome in diabetic vs. non-diabetic infection, we noticed

a striking number of genes involving transport were dysregulated. Some of the most highly

upregulated genes included *htrAB*, encoding the heme-regulated efflux pump and *hssRS*, the two-

component system that regulates *hrtAB* expression. These systems are critical for GBS survival as

they prevent a lethal heme overdose (72). Frank et al. demonstrated that heme accumulates at the

wound edge of rats in early infection, and that this leads to recruitment of leukocytes to infection

(73). We speculate, since the diabetic wound is hyperinflammatory, that there is excess heme in

this environment, and GBS upregulates heme efflux systems to survive potential toxicity. Other

downregulated genes include components of iron, zinc, and sugar transport systems. We

hypothesize that GBS is turning these transport genes off, as there may be excess metals and sugar

in the diabetic wound environment. Thus, it is possible that the diabetic host may not be

successfully sequestering metals in the wound thereby allowing for GBS persistence, but this

warrants further investigation.

Finally, we noticed that GBS colonies recovered from diabetic wounds were often

hyperpigmented and encoded mutations in the TCS *covRS*. CovR represses numerous GBS

virulence factors including the *cyl* operon as well as *pbsP* in multiple GBS strains (25, 35, 46). Of

note, is that spontaneous mutations in the *covRS* locus of GBS have been identified in clinical

isolates from patients with necrotizing fasciitis, sore throat, and prosthetic joint infection (48, 49,

74). In addition, numerous studies of *Streptococcus pyogenes* (GAS) have linked spontaneous

mutations in *covRS* to invasive infection in this pathogen (75, 76). Importantly, we have never

recovered any hyperpigmented colonies from non-diabetic animals. We therefore speculate that

the diabetic environment is promoting the selection of *covR* mutants in the wound. The exact

environmental factors that may promote *covR* mutations in the diabetic wound are currently

unknown. Interestingly, Jorge et al. found that the antimicrobial peptide LL-37 can bind to CovS

in GAS, resulting in the up-regulation of virulence factors (77). It is possible that GBS CovS may

also bind to antimicrobial peptides such as those we have shown are upregulated in the diabetic

wound environment. If this is the case, increased antimicrobial peptides in the diabetic

environment may be driving GBS to increase virulence factor production, and contribute to the

selection of *covR* mutants in the diabetic wound. It will also be of interest in future studies to

examine the possibility that mutations may also occur in *covS*, which could differentially impact

GBS persistence in the diabetic wound environment.

Using our model for GBS diabetic wound infection, we define changes in both the host and

pathogen transcriptome in the murine diabetic wound environment and demonstrate that GBS

promotes inflammation which contributes to the chronic wound environment. As we demonstrate,

this model is an efficient tool to investigate GBS pathogenesis, and potentially other pathogens, in

the context of diabetic wounds. Our results emphasize the importance of further studies on GBS

pathogenesis and the innate immune response at the diabetic wound interface.

Materials and Methods

Bacterial strains and growth conditions. GBS strains A909, COH1 and CJB111 were used in this study. GBS strains were grown in Todd Hewitt Broth (THB; Research Products International, RPI) statically at 37° C. When needed, antibiotic was added to THB at final concentrations of 100 µg/mL spectinomycin. Strains containing the plasmid pDCErm were grown in THB + 5 µg/mL erythromycin. All strains used in this study can be found in **Supplemental Table 6** and primers in **Supplemental Table 7.** The human immortalized keratinocyte cell line N/TERT-2G cells were kindly gifted from Dr. Johann Gudjonsson with permission from Dr. Jim Rheinwald and were maintained up to 25 passages at 37 ° C with 5% CO_2 according to methods described in Dickson et al., 2000 (56).

Ethics statement. Animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at University of Colorado Anschutz Medical Campus protocol #00987 and were per- formed using accepted veterinary standards. The University of Colorado Anschutz Medical Campus is AAALAC accredited; and its facilities meet and adhere to the standards in the "Guide for the Care and Use of Laboratory Animals".

Murine model of wound infection. Female or male, 8-week-old C57Bl/6J mice or *Lepr^{db}* mice were anesthetized with isoflurane the day prior to infection. The backs of the mice were shaved and treated with Nair and tail snips were performed to measure blood glucose levels with a glucometer. The next day mice were weighed and again anesthetized before undergoing wounding procedure. The backs of mice were disinfected with iodine wipes and then wounded with a 6 mm biopsy punch. Following, 1 x 10^7 CFU of GBS or PBS control was added before wrapping the mice in the adhesive Tegaderm^{TM}. Wounds were left for 72 h before removal of the adhesive. After another 24 h, mice were sacrificed via CO_2 inhalation and tissues were removed and homogenized for

bacterial enumeration. Tissues were placed 500 µL of PBS in 2.0 mL conical screw cap tubes

(Fisher) with 1.0 mm diameter zirconia/silica beads [BioSpec Cat. No. 1107911] and homogenized

by bead beating two times for 60 sec in a BioSpec mini bead beater. Tissue homogenates were

plated on GBS CHROMagar [SB282(B)], which allows only for the growth of GBS (in pink) and

Enterococcus spp. (in blue). These experiments were approved by the committee on the use and

care of animals at the University of Colorado-Anschutz Medical Campus in our protocol #00987.

Histology. Mice were wounded as described above. Following sacrifice, half of the wound tissue

was removed and fixed in formalin for 48 h. After, formalin was removed and wounds were placed

into 70% EtOH and sent to the histology core for processing at the Gates Center for Regenerative

Medicine, Dermatology at CU Anschutz. Images were taken with a BZ-X710 microscope

(Keyence).

RNA preparation and RNA-sequencing. Mice were wounded as previously described and tissues

were collected and placed into 500 µL of RNA protect from the Qiagen RNeasy kit in 2.0 mL

conical screw cap tubes with 1.0 mL zirconia beads as described above. Samples were placed into

a bead beater and homogenized three times for 60 sec with five minutes of ice between each bead

beating step. Sample supernatants were then collected and placed into new tubes for centrifugation.

Samples were centrifuged for 10 minutes at 13,000 RPM and the pellet was resuspended in 600

µL of RLT plus β-Mercaptoethanol and homogenized by beat beating for 60 sec with .1 mL

zirconia beads. Resulting homogenates were used for RNA preparation with the Qiagen RNeasy

kit. For mouse samples, RNA was extracted from 2-3 mice before pooling RNA to form triplicate

samples. For bacteria, we utilized three input controls as replicates and again pooled from 2-3 mice

for all bacterial samples. RNA samples were sent for dual RNA-sequencing (or bacterial RNA-

sequencing for input controls), to the Microbial Genome Sequencing Center for Illumina

Sequencing. Samples were rRNA depleted using RiboZero Plus (Illumina) and libraries for the 12

mouse samples and 3 input bacteria cDNA samples were sequenced using the Illumina Stranded

RNA sequencing platform. Raw sequencing reads in fastq format were aligned and annotated to

the mouse reference genome (mm10) or to the Clinical Group B Streptococcal Isolate CJB111

reference genome (CP063198_sRNA) using Qiagen CLC Genomics Workbench default settings

(version 21.0.5): mismatch cost, 2; insertion and deletion cost, 3; length and similarity fraction,

0.8. Normalization and differential expression calculations of uniquely mapped mouse or bacterial

transcripts were performed using R package DESeq2 (v1.34.0, RRID:SCR_015687) (78).

Heatmaps and further comparison of gene expression changes for all mouse and bacterial samples

were generated with Rstudio (v1.3.1073, RRID:SCR_000432). Gene set enrichment analysis

(GSEA) was performed using the fGSEA R package (v1.20.0, RRID:SCR_020938) with 10,000

permutations and the Hallmarks and GO Biological Processes gene set collections from the

Molecular Signatures Database (79).

ELISAs on wound homogenates. Proteins in homogenized tissues was quantified using R&D

systems ELISA kits (catalog # DY453, DY667, DY8596-05, DY4517-05). Protein detected was

normalized to tissue weight and reported as protein (pg) per mg of tissue.

Adherence of GBS to host cells. Adherence assays were performed as previously described (29,

80). Briefly, cell lines were seeded into 24 well plates and grown to a complete monolayer

(approximately 1×10^6 cells/well). GBS was grown to mid-log phase and normalized to 1×10^8

CFU/mL in PBS. 1×10^6 bacteria were added to one well of host cells to achieve a MOI of 1. To

assess adherence of GBS to host cells, bacteria were incubated with host cells for 30 min then the

cells were washed five times with PBS. Host cells were detached with 0.25% trypsin (Thermo

Fisher Scientific) and permeabilized with 0.025% Triton X-100 (Sigma) in PBS, serially diluted,

and plated to quantify all cell-associated bacteria.

Statistical analyses. Statistical analysis was performed using Prism version 9.0.0 (121) for

Windows (GraphPad Software, La Jolla, CA, United States) as described in figure legends.

References

1.   Dowey R, Iqbal A, Heller SR, Sabroe I, Prince LR. 2021. A Bittersweet Response to

Infection in Diabetes; Targeting Neutrophils to Modify Inflammation and Improve Host

Immunity. Front Immunol 12:1–21.

2.   Estelle E, Mathioudakis N. 2018. Update on management of diabetic foot ulcers. Ann N Y

Acad Sci 176:139–148.

3.   Davidson J. 1995. Wound healing. Adv Dent Res 9:8.

4.   Loots MAM, Lamme EN, Zeegelaar J, Mekkes JR, Bos JD, Middelkoop E. 1998.

Differences in cellular infiltrate and extracellular matrix of chronic diabetic and venous

ulcers versus acute wounds. J Invest Dermatol 111:850–857.

5.   Eming SA, Krieg T, Davidson JM. 2007. Inflammation in wound repair: Molecular and

cellular mechanisms. J Invest Dermatol 127:514–525.

6.   Guest, Christopher B. 2008. The implication of proinflammatory cytokines in type 2

diabetes. Front Biosci Volume:5187.

7.   Grice EA, Segre JA. 2012. Interaction of the microbiome with the innate immune

response in chronic wounds. Adv Exp Med Biol 946:55–68.

8.   Singer AJ, Clark RAF. 1999. Cutaneous Wound Healing. N Engl J Med.

9.   Young H, Knepper B, Hernandez W, Shor A, Bruntz M, Berg C, Price CS. 2015.

Pseudomonas aeruginosa An Uncommon Cause of Diabetic Foot Infection 105:125–129.

10. Young H, Miller W, Burnham R, Heard S, Berg C, Jenkins TC. How Do Preoperative

Antibiotics Affect Culture Yield in Diabetic Foot Infections ? 3–6.

11.   Kalan LR, Meisel JS, Loesche MA, Horwinski J, Soaita I, Chen X, Uberoi A, Gardner SE,

Grice EA. 2019. Strain- and Species-Level Variation in the Microbiome of Diabetic

Wounds Is Associated with Clinical Outcomes and Therapeutic Efficacy. Cell Host

Microbe 25:641-655.e5.

12.   Serra R, Grande R, Butrico L, Rossi A, Settimio UF, Caroleo B, Amato B, Gallelli L, De

Franciscis S. 2015. Chronic wound infections: The role of Pseudomonas aeruginosa and

Staphylococcus aureus. Expert Rev Anti Infect Ther 13:605–613.

13.   Moran GJ, Krishnadasan A, Gorwitz RJ, Fosheim GE, McDougal LK, Carey RB, Talan

DA. 2006. Methicillin-Resistant S. aureus Infections among Patients in the Emergency

Department. N Engl J Med 355:666–674.

14.   Thurlow LR, Stephens AC, Hurley KE, Richardson AR. 2020. Lack of nutritional

immunity in diabetic skin infections promotes Staphylococcus aureus virulence.

15.   Jacquet R, LaBauve AE, Akoolo L, Patel S, Alqarzaee AA, Fok Lung TW, Poorey K,

Stinear TP, Thomas VC, Meagher RJ, Parker D. 2019. Dual gene expression analysis

identifies factors associated with Staphylococcus aureus virulence in diabetic mice. Infect

Immun 87:1–18.

16.   Gjødsbøl K, Christensen JJ, Karlsmark T, Jørgensen B, Klein BM, Krogfelt KA. 2006.

Multiple bacterial species reside in chronic wounds : a longitudinal study 3.

17.   Trøstrup H, Lerche CJ, Christophersen LJ, Thomsen K, Jensen PØ, Hougen HP, Høiby N,

Moser C. 2017. Chronic Pseudomonas aeruginosa biofilm infection impairs murine

S100A8/A9 and neutrophil effector cytokines-implications for delayed wound closure?

Pathog Dis 75:1–39.

18. Maeda H, Akaike T, Sakata Y, Maruo K. 1993. Role of Bradykinin in Microbial Infection:

Enhancement of Septicemia by Microbial Proteases and Kinin, p. 159–165. *In* Proteases,

Protease Inhibitors and Protease-Derived Peptides. Birkhäuser Basel, Basel.

19. Hidekatsu Yanai. 2012. Group B streptococcus infection and diabetes: A review. J

Microbiol Antimicrob 4:1–5.

20. Schwartz B, Schuchat A, Oxtoby MJ, Cochi SL, Hightower A, Broome C V. 1991.

Invasive Group B Streptococcal Disease in Adults: A Population-Based Study in

Metropolitan Atlanta. JAMA J Am Med Assoc 266:1112–1114.

21. Regan JA, Klebanoff MA, Nugent RP. 1991. The epidemiology of group B streptococcal

colonization in pregnancy. Vaginal Infections and Prematurity Study Group. Obstet

Gynecol.

22. Wilkinson HW. 1978. Group b streptococcal infection in humans. Annu Rev Microbiol

41–57.

23. Gaschignard J, Levy C, Romain O, Cohen R, Bingen E, Aujard Y, Boileau P. 2011.

Neonatal Bacterial Meningitis. Pediatr Infect Dis J 30:212–217.

24. Skoff TH, Farley MM, Petit S, Craig AS, Schaffner W, Gershman K, Harrison LH,

Lynfield R, Mohle-boetani J, Zansky S, Albanese BA, Stefonek K, Zell ER, Jackson D,

Thompson T, Schrag SJ. 2009. Increasing Burden of Invasive Group B Streptococcal

Disease in Nonpregnant Adults , 1990 – 2007 30333:85–92.

25.   Rosa-Fraile M, Dramsi S, Spellerberg B. 2014. Group B streptococcal haemolysin and

523   pigment, a tale of twins. FEMS Microbiol Rev 38:932–946.

524   26.   Buscetta M, Firon A, Pietrocola G, Biondo C, Mancuso G, Midiri A, Romeo L, Galbo R,

525   Venza M, Venza I, Kaminski PA, Gominet M, Teti G, Speziale P, Trieu-Cuot P, Beninati

526   C. 2016. PbsP, a cell wall-anchored protein that binds plasminogen to promote

527   hematogenous dissemination of group B Streptococcus. Mol Microbiol 101:27–41.

528   27.   Patras KA, Derieux J, Al-Bassam MM, Adiletta N, Vrbanac A, Lapek JD, Zengler K,

529   Gonzalez DJ, Nizet V. 2018. Group B Streptococcus Biofilm Regulatory Protein A

530   Contributes to Bacterial Physiology and Innate Immune Resistance. J Infect Dis

531   218:1641–1652.

532   28.   Armistead B, Oler E, Waldorf KA, Rajagopal L, Disease I. 2019. The double life of

533   Group B Streptococcus: Asymptomatic colonizer and potent pathogen. J Mol Biol

534   431:2914–2931.

535   29.   Deng L, Spencer BL, Holmes JA, Mu R, Rego S, Weston TA, Hu Y, Sanches GF, Yoon

536   S, Park N, Nagao PE, Jenkinson HF, Thornton JA, Seo KS, Nobbs AH, Doran KS. 2019.

537   The group B streptococcal surface antigen I/II protein, BspC, interacts with host vimentin

538   to promote adherence to brain endothelium and inflammation during the pathogenesis of

539   meningitis. PLoS Pathog 15:1–30.

540   30.   Edmond KM, Kortsalioudaki C, Scott S, Schrag SJ, Zaidi AKM, Cousens S, Heath PT.

541   Group B streptococcal disease in infants aged younger than 3 months : systematic review

542   and meta-analysis. Lancet 379:547–556.

31.   Beier D, Gross R. 2006. Regulation of bacterial virulence by two-component systems.

Curr Opin Microbiol 9:143–152.

32.   Jiang S, Cieslewicz MJ, Kasper DL, Wessels MR. 2005. Regulation of Virulence by a

Two-Component System in Group B Streptococcus † 187:1105–1113.

33.   Faralla C, Metruccio MM, De Chiara M, Mu R, Patras KA, Muzzi A, Grandi G, Margarit

I, Doran KS, Janulczyk R. 2014. Analysis of two-component systems in group B

streptococcus shows that RgfAC and the novel FspSR modulate virulence and bacterial

fitness. MBio 5:1–12.

34.   Lamy M, Zouine M, Fert J, Vergassola M, Couve E, Pellegrini E, Glaser P, Kunst F,

Msadek T, Trieu-cuot P, Poyart C. 2004. CovS / CovR of group B streptococcus : a two-

component global regulatory system involved in virulence 54:1250–1268.

35.   Patras KA, Wang NY, Fletcher EM, Cavaco CK, Jimenez A, Garg M, Fierer J, Sheen TR,

Rajagopal L, Doran KS. 2013. Group B Streptococcus CovR regulation modulates host

immune signalling pathways to promote vaginal colonization. Cell Microbiol 15:1154–

1167.

36.   Lembo A, Gurney MA, Burnside K, Banerjee A, De M, Reyes L, Connelly JE, Lin W,

Jewell KA, Vo A, Renken CW, Doran KS, Rajagopal L. 2010. Regulation of CovR

expression in Group B streptococcus impacts blood-brain barrier penetration. Mol

Microbiol 77:431–443.

37.   Mandel MA, Mahmoud AA. 1978. Impairment of cell-mediated immunity in mutation

diabetic mice (db/db). J Immunol 120:1375–1377.

38.   P. HK, M. DM, L. CD. 1966. Diabetes, a New Mutation in the Mouse. Science (80-)

565  153:1127–1128.

566 39. Edwards JM, Watson N, Focht C, Wynn C, Todd CA, Walter EB, Phillips Heine R,

567   Swamy GK. 2019. Group B Streptococcus (GBS) Colonization and Disease among

568   Pregnant Women: A Historical Cohort Study. Infect Dis Obstet Gynecol 2019.

569 40. Poyart C, Tazi A, Réglier-Poupet H, Billoët A, Tavares N, Raymond J, Trieu-Cuot P.

570   2007. Multiplex PCR assay for rapid and accurate capsular typing of group B streptococci.

571   J Clin Microbiol 45:1985–1988.

572 41. Deng L, Schilcher K, Burcham LR, Kwiecinski JM, Johnson PM, Head SR. 2019.

573   Identification of Key Determinants of Staphylococcus aureus Vaginal Colonization 1–19.

574 42. Fabregat A, Sidiropoulos K, Viteri G, Forner O, Marin-Garcia P, Arnau V, D'Eustachio P,

575   Stein L, Hermjakob H. 2017. Reactome pathway analysis: A high-performance in-

576   memory approach. BMC Bioinformatics 18:1–9.

577 43. Zhang H, Potter BJ, Cao JM, Zhang C. 2011. Interferon-gamma induced adipose tissue

578   inflammation is linked to endothelial dysfunction in type 2 diabetic mice. Basic Res

579   Cardiol 106:1135–1145.

580 44. Gao X, Belmadani S, Picchi A, Xu X, Potter BJ, Tewari-Singh N, Capobianco S, Chilian

581   WM, Zhang C. 2007. Tumor necrosis factor-α induces endothelial dysfunction in Lepr db

582   mice. Circulation 115:245–254.

583 45. Jiang SM, Ishmael N, Hotopp JD, Puliti M, Tissi L, Kumar N, Cieslewicz MJ, Tettelin H,

584   Wessels MR. 2008. Variation in the group B Streptococcus CsrRS regulon and effects on

585   pathogenicity. J Bacteriol 190:1956–1965.

586   46.   Mazzuoli MV, Daunesse M, Varet H, Rosinski-Chupin I, Legendre R, Sismeiro O,

587   Gominet M, Kaminski PA, Glaser P, Chica C, Trieu-Cuot P, Firon A. 2021. The CovR

588   regulatory network drives the evolution of Group B Streptococcus virulence. PLoS Genet

589   17:1–27.

590   47.   Rajagopal L. 2009. Understanding the regulation of Group B Streptococcal virulence

591   factors. Future Microbiol 4:201–221.

592   48.   Lupo A, Ruppen C, Hemphill A, Spellerberg B, Sendi P. 2014. Phenotypic and molecular

593   characterization of hyperpigmented group B Streptococci. Int J Med Microbiol 304:717–

594   724.

595   49.   Whidbey C, Burnside K, Martinez RM, Gendrin C, Frando A, Harrell MI, Mcadams R.

596   2015. A Hyperhemolytic/Hyperpigmented Group B Streptococcus Strain with a CovR

597   Mutation Isolated from an Adolescent Patient with Sore Throat. Clin Res Infect Dis 2:1–

598   12.

599   50.   Doran KS, Chang JCW, Benoit VM, Eckmann L, Nizet V. 2002. Group B streptococcal β-

600   hemolysin/cytolysin promotes invasion of human lung epithelial cells and the release of

601   interleukin-8. J Infect Dis 185:196–203.

602   51.   Liu GY, Doran KS, Lawrence T, Turkson N, Puliti M, Tissi L, Nizet V. 2004. Sword and

603   shield: Linked group B streptococcal β-hemolysin/cytolysin and carotenoid pigment

604   function to subvert host phagocyte defense. Proc Natl Acad Sci U S A 101:14491–14496.

605   52.   Armistead B, Herrero-Foncubierta P, Coleman M, Quach P, Whidbey C, Justicia J, Tapia

606   R, Casares R, Millán A, Haidour A, Granger JR, Vornhagen J, Santana-Ufret V, Merillat

607   S, Adams Waldorf K, Cuerva JM, Rajagopal L. 2020. Lipid analogs reveal features

critical for hemolysis and diminish granadaene mediated Group B Streptococcus infection.

Nat Commun 11:1502.

53.   Papasergi S, Galbo R, Lanza-Cariccio V, Domina M, Signorino G, Biondo C, Pernice I,

Poyart C, Trieu-Cuot P, Teti G, Beninati C. 2013. Analysis of the Streptococcus agalactiae

exoproteome. J Proteomics 89:154–164.

54.   Lentini G, Midiri A, Firon A, Galbo R, Mancuso G, Biondo C, Mazzon E, Passantino A,

Romeo L, Trieu-Cuot P, Teti G, Beninati C. 2018. The plasminogen binding protein PbsP

is required for brain invasion by hypervirulent CC17 Group B streptococci. Sci Rep 8:1–

11.

55.   De Gaetano GV, Pietrocola G, Romeo L, Galbo R, Lentini G, Giardina M, Biondo C,

Midiri A, Mancuso G, Venza M, Venza I, Firon A, Trieu-Cuot P, Teti G, Speziale P,

Beninati C. 2018. The Streptococcus agalactiae cell wall-anchored protein PbsP mediates

adhesion to and invasion of epithelial cells by exploiting the host vitronectin/αvintegrin

axis. Mol Microbiol 110:82–94.

56.   Dickson MA, Hahn WC, Ino Y, Ronfard V, Wu JYd, Weinberg R a, Louis DN, Li FP,

Rheinwald JG. 2000. Human Keratinocytes That Express hTERT and Also Bypass a p16

INK4a -Enforced Mechanism That Limits Life Span Become Immortal yet Retain Normal

Growth and Differentiation Characteristics. Mol Cell Biol 20:1436–1447.

57.   Wong SL, Demers M, Martinod K, Gallant M, Wang Y, Goldfine AB, Kahn CR, Wagner

DD. 2015. Diabetes primes neutrophils to undergo NETosis, which impairs wound

healing. Nat Med 21:815–819.

58.   Hanses F, Park S, Rich J, Lee JC. 2011. Reduced neutrophil apoptosis in diabetic mice

during staphylococcal infection leads to prolonged Tnfα production and reduced

neutrophil clearance. PLoS One 6.

59. Jones SG, Edwards R, Thomas DW. 2004. Inflammation and Wound Healing: The Role

of Bacteria in the Immuno-Regulation of Wound Healing. Int J Low Extrem Wounds

3:201–208.

60. Wennekamp J, Henneke P. 2008. Induction and termination of inflammatory signaling in

group B streptococcal sepsis. Immunol Rev 225:114–127.

61. Baker CJ, Kasper DL. 1976. Microcapsule of type III strains of group B Streptococcus:

production and morphology. Infect Immun 13:189–194.

62. Sumitomo T, Nakata M, Higashino M, Yamaguchi M, Kawabata S. 2016. Group A

Streptococcus exploits human plasminogen for bacterial translocation across epithelial

barrier via tricellular tight junctions. Sci Rep 7:1–8.

63. Guo Y, Li J, Hagström E, Ny T. 2011. Beneficial and detrimental effects of

plasmin(Ogen) during infection and sepsis in mice. PLoS One 6.

64. Shen Y, Guo Y, Mikus P, Sulniute R, Wilczynska M, Ny T, Li J. 2012. Plasminogen is a

key proinflammatory regulator that accelerates the healing of acute and diabetic wounds.

Blood 119:5879–5887.

65. Cook LCC, Hu H, Maienschein-cline M, Federle J. 2018. A Vaginal Tract Signal Detected

by the Group B Streptococcus SaeRS System Elicits Transcriptomic Changes and

Enhances. Infect Immun 1–17.

66. Kroeze K, Gibbs S. 2012. Comparison of cytokine , chemokine and growth factor profiles

in burn wounds , chronic wounds and surgical excision wounds.

67.   Derré-Bobillot A, Cortes-Perez NG, Yamamoto Y, Kharrat P, Couvé E, Da Cunha V,

653   Decker P, Boissier MC, Escartin F, Cesselin B, Langella P, Bermúdez-Humarán LG,

654   Gaudu P. 2013. Nuclease A (Gbs0661), an extracellular nuclease of Streptococcus

655   agalactiae, attacks the neutrophil extracellular traps and is needed for full virulence. Mol

656   Microbiol 89:518–531.

657   68.   Trends C, Directions F. 2021. Wound Healing ResearchWound Healing Research.

658   69.   Hotterbeekx A, Kumar-Singh S, Goossens H, Malhotra-Kumar S. 2017. In vivo and In

659   vitro interactions between Pseudomonas aeruginosa and Staphylococcus spp. Front Cell

660   Infect Microbiol 7:1–13.

661   70.   Pérez-Pascual D, Gaudu P, Fleuchot B, Besset C, Rosinski-Chupin I, Guillot A, Monnet

662   V, Gardan R. 2015. RovS and its associated signaling peptide form a cell-to-cell

663   communication system required for Streptococcus agalactiae pathogenesis. MBio 6.

664   71.   Cook LC, LaSarre B, Federle MJ. 2013. Interspecies communication among commensal

665   and pathogenic streptococci. MBio 4.

666   72.   Joubert L, Dagieu JB, Fernandez A, Aurélie DB, Elise BD, Fleurot I, Gruss A, Lechardeur

667   D. 2017. Visualization of the role of host heme on the virulence of the heme auxotroph

668   Streptococcus agalactiae. Sci Rep 7:1–13.

669   73.   Wagener FADTG, Beurden HE Van, Hoff JWV Den, Adema GJ, Figdor CG. 2003. The

670   heme-heme oxygenase system : a molecular switch in wound healing 102:521–528.

74.   Sendi P, Johansson L, Dahesh S, Van Sorge NM, Darenberg J, Norgren M, Sjölin J, Nizet

V, Norrby-Teglund A. 2009. Bacterial phenotype variánts in group B streptococcal toxic

shock syndrome. Emerg Infect Dis 15:223–232.

75.   Walker MJ, Hollands A, Sanderson-Smith ML, Cole JN, Kirk JK, Henningham A,

McArthur JD, Dinkla K, Aziz RK, Kansal RG, Simpson AJ, Buchanan JT, Chhatwal GS,

Kotb M, Nizet V. 2007. DNase Sda1 provides selection pressure for a switch to invasive

group A streptococcal infection. Nat Med 13:981–985.

76.   Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. 2006. Genome-wide analysis

of group A streptococci reveals a mutation that modulates global phenotype and disease

specificity. PLoS Pathog 2:0041–0049.

77.   Velarde JJ, Ashbaugh M, Wessels MR. 2014. The human antimicrobial peptide LL-37

binds directly to CsrS, a sensor histidine kinase of group a streptococcus, to activate

expression of virulence factors. J Biol Chem 289:36315–36324.

78.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol 15:1–21.

79.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP.

2011. Molecular signatures database (MSigDB) 3.0. Bioinformatics 27:1739–1740.

80.   Spencer BL, Deng L, Patras KA, Burcham ZM, Sanches GF, Nagao PE, Doran KS. 2019.

Cas9 contributes to group b streptococcal colonization and disease. Front Microbiol 10:1–

15.

81.   Flores AR, Galloway-Peña J, Sahasrabhojane P, Saldaña M, Yao H, Su X, Ajami NJ,

Holder ME, Petrosino JF, Thompson E, Ros IMY, Rosini R, Grandi G, Horstmann N,

693   Teatero S, McGeer A, Fittipaldi N, Rappuoli R, Baker CJ, Shelburne SA. 2015. Sequence

694   type 1 group B Streptococcus, an emerging cause of invasive disease in adults, evolves by

695   small genetic changes. Proc Natl Acad Sci U S A 112:6431–6436.

696 82. Madoff LC, Michel JL, Kasper DL. 1991. A monoclonal antibody identifies a protective

697   C-protein alpha-antigen epitope in Group B streptococci. Infect Immun 59:204–210.

698 83. Wilson CB, Weaver WM. 1985. Comparative Susceptibility of Group B Streptococci and

699   Staphylococcus aureus to Killing by Oxygen Metabolites. J Infect Dis 152:323–329.

Acknowledgments

We would like to thank Elizabeth Grice and Sue Gardner for providing clinical GBS isolates, Laura

Cook for providing the Δ*pbsP* mutant, Lakshmi Rajagopal for the Δ*covR* mutant, and Amber

Nguyen for assistance with sequencing. This work was supported by NIH grants R01NS116716,

R01AI153332 to K.S.D., NIHT32DK120521 to R.A.K. and R01AI153185 to A.R.H., and

Department of Veteran Affairs Award BX002711 to A.R.H. The authors declare no other

competing interests.

Data and Materials Availability

711   All data needed to evaluate the conclusions in the paper are present in the paper and/or the

712   Supplementary Materials. Mouse and bacterial RNA-sequencing datasets generated in this study

713   including all raw data files were uploaded to NCBI GEO Depository (GSE201342).

Figures, tables, and legends

Figure 1. Murine Model of GBS Diabetic Wound Infection. (A) Schematic of the murine model

of diabetic wound infection. (B) Non-fasting glucose levels from the blood of mice the day of

infection. (C) Wound area calculated in ImageJ from representative mice following GBS infection.

(D) Representative images of GBS infected wounds from non-diabetic and diabetic mice on the

day of sacrifice. (E) CFU recovered from wounds of non-diabetic and diabetic mice after GBS

infection. All animal infections proceeded for four days with three days under adhesive and

sacrifice 24 h after adhesive removal. Significance determined by Mann–Whitney U test; *p<.05,

**p<.01, ****p<.0001.

Figure 2. The hyper-inflammatory wound environment in diabetic mice is more susceptible to GBS persistence. (A) PCA plot of murine transcriptome from RNA-seq analysis. (B) Volcano plot of differentially expressed genes. (C) GSEA of pathways enriched in diabetic wounds infected with GBS in comparison to non-diabetic. Significant pathways are in bold. Normalized enrichment score (NES) presented as a heat map with highly enriched pathways in red. (D-F) ELISAs on tissue homogenates at time of sacrifice. All concentrations were normalized to tissue weight. Mice were infected for four days with three days under adhesive and sacrifice 24 h after adhesive removal. Significance determined by Mann-Whitney U-test; ***p<.001, # = .0597.

Figure 3. Diabetic animals upregulate inflammatory pathways and immune cell recruitment upon

GBS infection. (A) Volcano plot for differentially expressed genes when comparing diabetic mice

infected with GBS to uninfected controls. (B) Gene set enrichment analysis of diabetic animals

infected with GBS vs. uninfected controls. Significant pathways in bold. NES presented as a heat

map with highly enriched pathways in red. (C) Percent initial body weight of mice after GBS

infection. (D) GBS recovery from wound tissues of uninfected vs. infected mice. (E-H) ELISAs

on tissue homogenates at time of sacrifice. All concentrations were normalized to tissue weight.

(I) Histology of diabetic wound tissue at time of sacrifice. Mice were infected for four days with

three days under adhesive and sacrifice 24 h after adhesive removal. Significance determined by

Mann-Whitney U-test; *p<.05, **p<.01, ****p<.0001.

Figure 4. GBS Transcriptome in Diabetic Wound Infection. (A) PCA plot of bacterial transcriptome in RNA-seq. (B) Venn diagram of differentially expressed genes in each comparison. (C) Volcano plot of differentially expressed genes in diabetic infection vs. input control. (D) Volcano plot of differentially expressed genes in diabetic vs. non-diabetic wound infection.

Gene number (CJB111 genome)	Gene name	Description	Diabetic/ Input	FDR p-value
ID870_07365	*pbsP*	YSIRK signal domain/LPXTG anchor domain surface protein	186.61	8.43E-110
ID870_02195	*shp2*	peptide pheromone SHP2	101.66	3.55E-21
ID870_05830	*nucA*	DNA/RNA non-specific endonuclease	83.14	2.94E-43
ID870_02190	*rovS*	transglutaminase, regulated by shp2 RovS	47.73	2.26E-104
ID870_03115	*clpL*	ATP-dependent Clp protease ATP-binding subunit	26.96	7.49E-50
ID870_07100		type II toxin-antitoxin system RelB/DinJ family antitoxin	25.86	1.94E-33
ID870_07095		toxin-antitoxin system, toxin component, MazF family protein	23.11	8.66E-37
ID870_10045		alpha/beta hydrolase	22.10	5.82E-118
ID870_03740	*tpx*	thiol peroxidase	15.03	8.58E-51
ID870_03230		Ltp family lipoprotein	13.85	1.19E-82
ID870_08245		WXG100 family type VII secretion target	13.12	4.30E-31
ID870_04175		WXG100 family type VII secretion target	11.10	5.76E-26
ID870_06190		type II toxin-antitoxin system death-on-curing family toxin	9.97	3.57E-25
ID870_04170		WXG100 family type VII secretion target	8.94	5.40E-19
ID870_05905	*cylK*	CylK protein	8.91	9.63E-21
ID870_08300		type II toxin-antitoxin system RelE/ParE family toxin	8.53	6.45E-13
ID870_02285		LPXTG cell wall anchor domain-containing protein	8.48	1.63E-45
ID870_06195		type II toxin-antitoxin system Phd/YefM family antitoxin	8.36	1.74E-32
ID870_01270		Asp23/Gls24 family envelope stress response protein	8.20	6.73E-38
ID870_10470	*cfb*	CAMP factor pore-forming toxin Cfb	7.82	6.73E-09
ID870_10565		WXG100 family type VII secretion target	5.93	1.16E-07
ID870_03660		CidA/LrgA family protein	5.05	1.77E-07
ID870_00915		3'-5' exonuclease	4.79	1.59E-12
ID870_05925	*cylE*	cylE protein	3.85	2.41E-16
ID870_05955	*cylD*	CylD	3.82	1.95E-07
ID870_00425		glycoside hydrolase family 88 protein	3.80	3.54E-04
ID870_05890		S8 family serine peptidase	3.44	6.31E-11
ID870_03865		lipase	3.34	1.90E-09

 Table 1. Select GBS Virulence factors upregulated in diabetic infection vs. input control.

Figure 5. The CovRS Regulon Contributes to Diabetic Wound Infection. (A) GBS colonies on THA recovered from diabetic wounds after four days of infection. (B) Hyperpigmented colonies plated on sheep's blood agar; C1-3 from non-pigmented colonies and C4-6 from pigmented colonies (C) Structural model of CovR mutations on VanR structure from *Streptomyces coelicolor*. (D) Percentage of total GBS colonies from diabetic wound tissue 24 h or 1 week after adhesive removal which were hyper-pigmented on Todd-Hewitt agar. (E) CFU of GBS recovered from diabetic wound tissue 24 h after adhesive removal. (F) KC concentration recovered from wound tissue homogenates. All ELISA data is normalized to tissue weight. Significance determined via Mann Whitney U-test; *p<.05, **p<.01.

Figure 6. PbsP contributes to GBS burden and inflammation in the diabetic wound. (A) CFU recovered from diabetic mice infected with A909 or a Δ*pbsP* mutant after four days of infection. (B,C,D) MPO, Elastase, and Calprotectin concentration recovered from wound tissue homogenates. All ELISA data is normalized to tissue weight. (E) GBS adherence to immortalized keratinocytes with either wildtype, mutant or complemented PbsP. Significance determined via Mann Whitney U-test; *p<.05, **p<.01.

Figure 7. Model for GBS pathogenesis in the diabetic wound. Image generated in BioRender.

Supplemental figures and tables, and legends:

Supplemental Figure 1. Murine Model of GBS Diabetic Wound Infection in Female and Male

Mice. (A) CFU recovered from wounds of non-diabetic and diabetic mice after GBS infection. All

animal infections proceeded for four days with three days under adhesive and sacrifice 24 h after

adhesive removal. Significance determined by Mann–Whitney U test; *p<.05.

Serotype	Ia	Ib	II	III	V	Undetermined
Distribution	9/27 (33.33%)	2/27 (7.41%)	6/27 (22.22%)	3/27 (11.11%)	6/27 (22.22%)	1/27 (3.7%)
Strain numbers	112[C]	503[I]	69[C]	64[C]	98[C]	734[I]
	836[I]	1418[I]	85[C]	113[C]	103[C]	
	884[I]		157[C]	361[I]	130[C]	
	982[I]		330[I]		809[I]	
	1052[I]		607[I]		914[I]	
	1394[I]		1962[I]		1101[I]	
	1463[I]					
	1852[I]					
	2000[I]					

Isolates obtained from Colorado are designated with a [C] and isolates from Iowa an [I].

Supplemental Table 1. Clinical isolate serotypes.

Supplemental Figure 2. Murine transcriptome in non-diabetic uninfected vs. diabetic uninfected

comparison. (A) Volcano plot of differentially expressed genes. (B) GSEA of pathways enriched

in diabetic wounds. Significant pathways are in bold. NES presented as a heat map with highly

enriched pathways in red.

Gene ID	DAVID Description	Diabetic/ non-diabetic log2 FC	padj
Neutrophil degranulation			
Ear1	eosinophil-associated, ribonuclease A family, member 1(Ear1)	2.33	1.24E-04
Olfm4	olfactomedin 4(Olfm4)	3.87	6.29E-12
Acaa1b	acetyl-Coenzyme A acyltransferase 1B(Acaa1b)	3.88	6.99E-37
Gpr84	G protein-coupled receptor 84(Gpr84)	2.32	7.57E-06
Hp	haptoglobin(Hp)	2.71	3.74E-23
Slco4c1	solute carrier organic anion transporter family, member 4C1(Slco4c1)	2.12	6.95E-07
Prg2	proteoglycan 2, bone marrow(Prg2)	4.27	1.11E-07
Lrg1	leucine-rich alpha-2-glycoprotein 1(Lrg1)	2.10	4.08E-19
Rab9b	RAB9B, member RAS oncogene family(Rab9b)	2.23	1.85E-02
Mmp8	matrix metallopeptidase 8(Mmp8)	3.82	3.85E-12
Pglyrp1	peptidoglycan recognition protein 1(Pglyrp1)	3.12	2.05E-11
Elane	elastase, neutrophil expressed(Elane)	5.16	1.51E-06
Krt8	keratin 8(Krt8)	3.49	1.23E-04
Prtn3	proteinase 3(Prtn3)	2.93	1.41E-07
Camp	cathelicidin antimicrobial peptide(Camp)	4.49	1.52E-16
Mmp25	matrix metallopeptidase 25(Mmp25)	2.72	2.16E-09
Itgam	integrin alpha M(Itgam)	4.26	1.06E-02
Abca13	ATP-binding cassette, sub-family A (ABC1), member 13(Abca13)	3.64	5.46E-15
Lcn2	lipocalin 2(Lcn2)	3.25	2.54E-30
Cd36	CD36 molecule(Cd36)	3.38	5.74E-49
Itgad	integrin, alpha D(Itgad)	4.75	5.34E-45
Ctsg	cathepsin G(Ctsg)	8.44	3.39E-08
Ms4a3	membrane-spanning 4-domains, subfamily A, member 3(Ms4a3)	5.30	1.12E-03
Mgam	maltase-glucoamylase(Mgam)	2.63	1.45E-07
Epx	eosinophil peroxidase(Epx)	3.31	1.40E-03
Mpo	myeloperoxidase(Mpo)	4.41	4.40E-05
Cd177	CD177 antigen(Cd177)	5.14	3.46E-20
Activation of Matrix Metalloproteases			
Mmp25	matrix metallopeptidase 25(Mmp25)	2.72	2.16E-09
Ctsg	cathepsin G(Ctsg)	8.44	3.39E-08
Mmp8	matrix metallopeptidase 8(Mmp8)	3.82	3.85E-12
Tpsab1	tryptase alpha/beta 1(Tpsab1)	3.69	1.60E-14
Elane	elastase, neutrophil expressed(Elane)	5.16	1.51E-06
Antimicrobial peptides			
Ear1	eosinophil-associated, ribonuclease A family, member 1(Ear1)	2.33	1.24E-04
Prtn3	proteinase 3(Prtn3)	2.93	1.41E-07
Ctsg	cathepsin G(Ctsg)	8.44	3.39E-08
Clec10a	C-type lectin domain family 10, member A(Clec10a)	2.02	3.03E-17
Camp	cathelicidin antimicrobial peptide(Camp)	4.49	1.52E-16
Pglyrp1	peptidoglycan recognition protein 1(Pglyrp1)	3.12	2.05E-11
Elane	elastase, neutrophil expressed(Elane)	5.16	1.51E-06
Lcn2	lipocalin 2(Lcn2)	3.25	2.54E-30
Formation of the cornified envelpoe/ keratinization			
Krt35	keratin 35(Krt35)	-5.82	2.10E-03
Klk13	kallikrein related-peptidase 13(Klk13)	-2.17	1.97E-20
Krt79	keratin 79(Krt79)	-2.14	5.76E-06
Lelp1	late cornified envelope-like proline-rich 1(Lelp1)	-2.25	2.89E-02
Krt71	keratin 71(Krt71)	-7.27	8.43E-06
Sprr3	small proline-rich protein 3(Sprr3)	-2.46	1.29E-02
Lce1b	late cornified envelope 1B(Lce1b)	-2.18	4.53E-05
Krt75	keratin 75(Krt75)	-3.10	7.08E-63
Casp14	caspase 14(Casp14)	-2.53	5.33E-06
Pcsk6	proprotein convertase subtilisin/kexin type 6(Pcsk6)	-2.31	1.07E-21
Lce1f	late cornified envelope 1F(Lce1f)	-2.02	7.08E-05
Krt73	keratin 73(Krt73)	-6.34	2.07E-03
Klk12	kallikrein related-peptidase 12(Klk12)	-2.09	8.60E-13
Krt28	keratin 28(Krt28)	-4.51	1.26E-04
Krt27	keratin 27(Krt27)	-7.76	8.43E-06
Krt25	keratin 25(Krt25)	-8.34	1.47E-04
Lce6a	late cornified envelope 6A(Lce6a)	-2.09	1.60E-03
Krt82	keratin 82(Krt82)	-4.69	9.87E-03
Flg	filaggrin(Flg)	-3.73	1.93E-06
Cdsn	corneodesmosin(Cdsn)	-2.13	2.08E-11
Ivl	involucrin(Ivl)	-3.08	2.16E-11
Rptn	repetin(Rptn)	-3.27	3.10E-14
Klk8	kallikrein related-peptidase 8(Klk8)	-2.12	7.20E-13
Spink5	serine peptidase inhibitor, Kazal type 5(Spink5)	-2.05	3.10E-22
Klk5	kallikrein related-peptidase 5(Klk5)	-2.05	1.41E-07
FGFR1 ligand binding and activation			
Flg	filaggrin(Flg)	-3.73	1.93E-06
Fgf4	fibroblast growth factor 4(Fgf4)	-2.49	3.77E-02

803   **Supplemental Table 2**. Murine transcriptome in diabetic infected vs. non-diabetic infected

804   comparison. Select genes and pathways.

Gene ID	DAVID Description	Diabetic infected/ uninfected log2 FC	padj
Neutrophil Degranulation			
Ear1	eosinophil-associated, ribonuclease A family, member 1(Ear1)	2.17	8.85E-04
Tarm1	T cell-interacting, activating receptor on myeloid cells 1(Tarm1)	2.29	6.11E-05
Olfm4	olfactomedin 4(Olfm4)	3.04	4.43E-07
Gpr84	G protein-coupled receptor 84(Gpr84)	2.70	7.45E-07
S100a8	S100 calcium binding protein A8 (calgranulin A)(S100a8)	2.73	1.61E-09
Slco4c1	solute carrier organic anion transporter family, member 4C1(Slco4c1)	2.05	6.44E-06
S100a9	S100 calcium binding protein A9 (calgranulin B)(S100a9)	2.55	2.91E-10
Prg2	proteoglycan 2, bone marrow(Prg2)	4.50	2.89E-07
Mmp8	matrix metallopeptidase 8(Mmp8)	4.73	3.51E-17
Pglyrp1	peptidoglycan recognition protein 1(Pglyrp1)	3.98	8.18E-17
Elane	elastase, neutrophil expressed(Elane)	5.93	1.45E-06
Prtn3	proteinase 3(Prtn3)	2.77	2.91E-06
Camp	cathelicidin antimicrobial peptide(Camp)	5.72	3.39E-20
Mmp25	matrix metallopeptidase 25(Mmp25)	3.15	1.74E-11
Abca13	ATP-binding cassette, sub-family A (ABC1), member 13(Abca13)	4.35	1.15E-19
Lcn2	lipocalin 2(Lcn2)	4.41	1.71E-54
Gzmb	granzyme B(Gzmb)	2.33	1.59E-07
Cxcl1	chemokine (C-X-C motif) ligand 1(Cxcl1)	2.20	1.79E-06
Fpr1	formyl peptide receptor 1(Fpr1)	2.79	5.57E-07
Cxcr1	chemokine (C-X-C motif) receptor 1(Cxcr1)	2.42	1.66E-05
Ctsg	cathepsin G(Ctsg)	3.08	1.88E-03
Ms4a3	membrane-spanning 4-domains, subfamily A, member 3(Ms4a3)	7.25	5.06E-04
Mgam	maltase-glucoamylase(Mgam)	3.26	3.34E-10
Epx	eosinophil peroxidase(Epx)	4.37	1.09E-03
Mpo	myeloperoxidase(Mpo)	6.35	3.09E-08
Cd177	CD177 antigen(Cd177)	3.91	2.95E-11
Signaling by Interleukins			
Nos2	nitric oxide synthase 2, inducible(Nos2)	4.69	2.21E-10
Il12b	interleukin 12b(Il12b)	2.50	1.12E-02
Ccl5	chemokine (C-C motif) ligand 5(Ccl5)	3.29	2.22E-10
Il12a	interleukin 12a(Il12a)	2.68	3.60E-04
Il27	interleukin 27(Il27)	2.77	1.32E-06
Saa1	serum amyloid A 1(Saa1)	3.61	2.01E-21
Prtn3	proteinase 3(Prtn3)	2.77	2.91E-06
Csf2	colony stimulating factor 2 (granulocyte-macrophage)(Csf2)	3.04	2.99E-02
Lcn2	lipocalin 2(Lcn2)	4.41	1.71E-54
Ebi3	Epstein-Barr virus induced gene 3(Ebi3)	2.58	8.05E-09
Mpl	myeloproliferative leukemia virus oncogene(Mpl)	2.06	2.97E-02
Gzmb	granzyme B(Gzmb)	2.33	1.59E-07
Cxcl1	chemokine (C-X-C motif) ligand 1(Cxcl1)	2.20	1.79E-06
Fpr1	formyl peptide receptor 1(Fpr1)	2.79	5.57E-07
Ctsg	cathepsin G(Ctsg)	3.08	1.88E-03
Formation of a fibrin clot			
Gp9	glycoprotein 9 (platelet)(Gp9)	2.12	2.36E-04
Fga	fibrinogen alpha chain(Fga)	6.12	1.58E-03
Prtn3	proteinase 3(Prtn3)	2.77	2.91E-06
Cd177	CD177 antigen(Cd177)	3.91	2.95E-11
Striated muscle contraction			
Myh8	myosin, heavy polypeptide 3, skeletal muscle, embryonic(Myh3)	-2.79	4.54E-07
Myh3	myosin, heavy polypeptide 8, skeletal muscle, perinatal(Myh8)	-2.65	3.00E-06
Actc1	actin, alpha, cardiac muscle 1(Actc1)	-2.86	3.17E-04
Myogenesis			
Ctnna2	catenin (cadherin associated protein), alpha 2(Ctnna2)	-2.24	3.59E-06
Myf5	myogenic factor 5(Myf5)	-2.41	5.88E-03

Supplemental Table 3. Murine transcriptome in diabetic infected vs. diabetic uninfected

comparison. Select genes and pathways.

Locus Tag (CJB111)	Gene name	Description	Diabetic/ Non-diabetic	padj
ID870_08840	hrtB	FtsX-like permease family protein	254.88	2.45E-08
ID870_01095		universal stress protein	47.59	2.19E-30
ID870_08835	hrtA	ABC transporter ATP-binding protein	37.58	2.69E-04
ID870_05815	fetB	iron export ABC transporter permease subunit FetB	14.81	4.20E-06
ID870_02600	pilB	PI-2a pilus major subunit PilB	12.90	3.04E-03
ID870_03655		LrgB family protein	10.85	7.81E-03
ID870_08490	lrgA	CidA/LrgA family protein	10.12	4.31E-10
ID870_08485	lrgB	antiholin-like protein LrgB	8.67	1.03E-07
ID870_03740	tpx	thiol peroxidase	8.55	3.59E-25
ID870_08825	hssS	HAMP domain-containing histidine kinase	6.67	1.68E-12
ID870_05980		serine hydrolase	6.48	2.03E-02
ID870_08830	hssR	response regulator transcription factor	6.40	2.29E-04
ID870_02595	pilA	PI-2a pilus adhesin PilA	5.77	8.47E-03
ID870_06030		SpaH/EbpB family LPXTG-anchored major pilin	5.53	1.61E-02
ID870_03660		CidA/LrgA family protein	4.71	2.12E-02
ID870_00130	phoP	response regulator transcription factor	-3.16	2.26E-03
ID870_02090	nikE	ABC transporter ATP-binding protein	-3.17	4.95E-04
ID870_00205		sensor histidine kinase	-3.18	2.51E-08
ID870_08645	adcC	metal ABC transporter ATP-binding protein	-3.41	2.16E-08
ID870_05910	cylJ	cylJ protein	-3.64	2.34E-08
ID870_00210		response regulator	-3.67	6.26E-05
ID870_04175		WXG100 family type VII secretion target	-3.72	4.44E-07
ID870_02080	nikC	ABC transporter permease	-3.76	1.52E-06
ID870_02660		iron ABC transporter permease	-3.79	6.61E-03
ID870_04785		pneumococcal-type histidine triad protein	-4.00	6.37E-07
ID870_07205		S8 family serine peptidase	-4.09	4.67E-09
ID870_02700		ferredoxin	-4.13	3.45E-12
ID870_02075	nikB	ABC transporter permease	-4.39	1.18E-06
ID870_03375	hylB	hyaluronate lyase	-4.42	5.51E-12
ID870_02070	nikA	nickel ABC transporter, nickel/metallophore periplasmic binding protein	-4.67	7.88E-07
ID870_05425		LPXTG cell wall anchor domain-containing protein	-5.04	2.40E-07
ID870_10190		bacteriocin immunity protein	-5.22	3.65E-08
ID870_00255	shtII	pneumococcal-type histidine triad protein	-5.56	4.44E-07
ID870_09365		sugar ABC transporter permease	-5.60	5.04E-06
ID870_03205	lmb	metal ABC transporter substrate-binding lipoprotein/laminin-binding adhesin Lmb	-5.79	1.43E-10
ID870_00250	adcAII	zinc ABC transporter substrate-binding protein	-5.96	1.54E-05
ID870_03210	sht	pneumococcal-type histidine triad protein	-6.47	1.26E-18
ID870_07705		PTS sugar transporter subunit IIB	-11.84	2.62E-09
ID870_07700		PTS sugar transporter subunit IIC	-14.29	8.15E-40
ID870_00580		PTS sugar transporter subunit IIC	-19.83	2.09E-44

811 **Supplemental Table 4**. Select GBS transcripts with altered regulation in diabetic wounds vs.
812 non-diabetic.

Category	AA Position	Nucleotide position	Substitution
AA Substitution	G38		Gly-to-Glu
	A96		Ala-to-Val
	R66		Arg-to-Cys
	G61		Gly-to-Asp
	D53		Asp-to-His
	R67		Arg-to-Pro
	R67		Arg-to-Cys
	A96		Ala-to-Val
Nucleotide deletion		387	Deletion of A
		169	Deletion of T
	V31		Deletion of Val
Insertion/deletion		123	Multiple nucleotides deleted
		35	Multiple nucleotides inserted

Supplemental Table 5. Mutations in *covR* from hyperpigmented colonies recovered from the

diabetic wound.

Supplemental Figure 3. Analysis of a Δ*covR* mutant in diabetic wound infection. CFU recovered

from wounds of diabetic mice after GBS infection. Animal infections proceeded for four or 11

days with three days under adhesive and sacrifice 24 h or 1 week after adhesive removal.

Significance determined by Mann–Whitney U test.

Supplemental Figure 4. Analysis of a Δ*pbsP and* Δ*cylE* mutant in non-diabetic wound infection.

CFU recovered from wounds of diabetic mice after GBS infection. Animal infections proceeded

for four days with three days under adhesive and sacrifice 24 h after adhesive removal.

Significance determined by One-way ANOVA with comparison to A909.

Strain	strain type	Antibiotic resistance	Reference
CJB111	clinical isolate		81
A909	clinical isolate		82
COH1	clinical isolate		83
A909 ΔpbsP	mutant	Cm 3	65
A909 ΔpbsP::pbsP	complement	Cm 3, Spec 100	65
A909 ΔcylE	mutant	Cm 2	51
A909 ΔcovR	mutant	Cm 3	36

832 **Supplemental Table 6.** Strains used in this study

Primer Name	Sequence 5' → 3'
Serotype Ia Forward	GGTCAGACTGGATTAATGGTATGC

Serotype Ia Reverse	GTAGAAATAGCCTATATACGTTGAATGC
Serotype Ib Forward	TAAACGAGAATGGAATATCACAAACC
Serotype Ib Reverse	GAATTAACTTCAATCCCTAAACAATATCG
Serotype II Forward	GCTTCAGTAAGTATTGTAAGACGATAG
Serotype II Reverse	TTCTCTAGGAAATCAAATAATTCTATAGGG
Serotype III Forward	TCCGTACTACAACAGACTCATCC
Serotype III Reverse	AGTAACCGTCCATACATTCTATAAGC
Serotype IV Forward	GGTGGTAATCCTAAGAGTGAACTGT
Serotype IV Reverse	CCTCCCCAATTTCGTCCATAATGGT
Serotype V Forward	GAGGCCAATCAGTTGCACGTAA
Serotype V Reverse	AACCTTCTCCTTCACACTAATCCT

835 **Supplemental Table 7.** Primers used in this study

Diabetic Wound Infection

Mpo
Elane
S100a8
S100a9

pbsP
cylE
nucA

Neutrophil   GBS   GBS (*ΔcovR*)