Title

AGO1 and HSP90 buffer different genetic variants in *Arabidopsis thaliana*

Authors

Tzitziki Lemus[1,2], G. Alex Mason[1,3], Kerry L. Bubb[1], Cristina M. Alexandre[1], Christine Queitsch[1*], and Josh T. Cuperus[1*]

To whom correspondence should be addressed: cuperusj@uw.edu, queitsch@uw.edu

Affiliations
1. Department of Genome Sciences, University of Washington, William H. Foege Hall, 3720 15th Ave NE, Seattle, WA 98105.

2. Present address: Department of Human Genetics, University of California Los Angeles. Gonda (Goldschmied) Building, 695 Charles E. Young Drive S. Los Angeles CA 90095

3. Present address: Department of Plant Biology and Genome Center, University of California, Davis. 1 Shields Ave. Davis, CA 95616

Running head: AGO1 and HSP90 buffer different genetic variants

Keywords: AGO1, HSP90, HUA2, Buffering, Capacitors, Epistasis

Article Summary
Argonaute 1 (AGO1), a key player in plant development, interacts with the chaperone HSP90 which buffers environmental and genetic variation. We found that *AGO1* buffers environmental and genetic variation in the same traits; however, AGO1-dependent and HSP90-dependent loci do not overlap. Detailed analysis of a buffered locus found that a non-functional *HUA2* allele decouples days to flowering and rosette leaf number in an AGO1-dependent manner, suggesting the AGO1-dependent buffering acts at the network level.

Abstract
Argonaute 1 (AGO1), the principal protein component of microRNA-mediated regulation, plays a key role in plant growth and development. AGO1 physically interacts with the chaperone HSP90, which buffers cryptic genetic variation in plants and animals. We sought to determine whether genetic perturbation of *AGO1* in *Arabidopsis thaliana* would also reveal cryptic genetic variation, and if so, whether *AGO1*-dependent loci overlap with those dependent on HSP90. To address these questions, we introgressed a hypomorphic mutant allele of *AGO1* into a set of mapping lines derived from the commonly used *Arabidopsis* strains Col-0 and L*er*. Although we identified several cases in which AGO1 buffered genetic variation, none of the *AGO1*-dependent loci overlapped with those buffered by HSP90 for the traits assayed. We focused on one buffered locus where *AGO1* perturbation uncoupled the traits days to flowering and rosette leaf number, which are otherwise closely correlated. Using a bulk segregant approach, we identified a non-functional L*er hua2* mutant allele as the causal AGO1-buffered polymorphism. Introduction of a non-functional *hua2* allele into a Col-0 *ago1* mutant background recapitulated the L*er*-dependent *ago1* phenotype, implying that coupling of these traits involves different molecular players in these closely related strains. Taken together, our findings demonstrate that even though AGO1 and HSP90 buffer genetic variation in the same traits, these robustness regulators interact epistatically with different genetic loci, suggesting that higher-order epistasis is uncommon.

Introduction
Genetic networks rely on various types of feedback loops, redundancy, and other mechanisms like chaperones and small RNAs to ensure phenotypic robustness in spite of environmental or genetic perturbations (Rutherford and Lindquist 1998; Queitsch *et al.* 2002; Masel and Siegal 2009; Whitacre 2012; Lempe *et al.* 2013; Lachowiec *et al.* 2018; Zabinsky *et al.* 2019). Network disruptions decrease environmental and developmental robustness and, dependent on their nature, increase phenotypic variation in a trait or affect organismal phenotypes more broadly. For example, perturbation of the essential chaperone HSP90 broadly increases phenotypic variation in plants, fungi, and animals, with many organismal traits affected in a background-specific manner (Rutherford and Lindquist 1998; Queitsch *et al.* 2002; Yeyati *et al.* 2007; Sangster *et al.* 2008b; a; Jarosz and Lindquist 2010; Rohner *et al.* 2013; Karras *et al.* 2017; Zabinsky *et al.* 2019). When fully functional, HSP90 chaperones a select but highly diverse group of client proteins, including many kinases, receptors and transcription factors with crucial roles in development (Schopf *et al.* 2017). When chaperone function is perturbed, client proteins encoding genetic variants may fail to mature or fold differently, leading to pathway failure or rewiring (Dorrity *et al.* 2018) and hence altered phenotypes (Zabinsky *et al.* 2019). The phenomenon that HSP90 keeps genetic variation phenotypically silent and HSP90 perturbation allows its expression has become known as phenotypic capacitance (Rutherford and Lindquist 1998; Masel and Siegal 2009) – a different term for epistasis (Zabinsky *et al.* 2019). In contrast

to the traditional definition of epistasis, which describes the non-reciprocal interaction of two
loci, phenotypic capacitance is an epistasis phenomenon in which one locus, *e.g., HSP90*,
interacts with several others. HSP90 perturbation can increase phenotypic variation even in the
absence of genetic variation, presumably because subtle differences in microenvironment,
developmental stage, or cell state lead to inhibition of different client proteins among individuals
in a seemingly stochastic manner (Zabinsky *et al.* 2019)..
84
Another important source of developmental and environmental robustness is post-transcriptional
regulation by small RNAs. Small RNAs regulate the expression of their target genes in a
sequence-specific manner. In plants, most endogenous post-transcriptional gene regulation is
mediated by AGO1 loaded with microRNAs (miRNAs, MIR) (Axtell 2013; Bologna and
Voinnet 2014). In animals, some microRNAs are known to buffer stochastic (Hilgers *et al.*
2010), environmental (Li *et al.* 2009), and genetic variation (Cassidy *et al.* 2013). MicroRNAs
play major roles throughout plant development, including in the onset of flowering, an
irreversible developmental transition of outsized effect on reproductive success in annual plants
(Dong *et al.* 2022). In particular, the MIR156 and MIR172 gene families are essential for fine-
tuning expression of the complex gene network that determines the number of days until
flowering is initiated and the number of rosette leaves at this developmental stage. Misregulation
or mutation of their gene targets alters both traits in the crucifer model *Arabidopsis thaliana*
(Aukerman and Sakai 2003; Yamaguchi *et al.* 2009; Wu *et al.* 2009).
98
In *Arabidopsis*, the traits days to flowering (*i.e.*, flowering time, onset of flowering) and rosette
leaf number are so closely linked that the traits are often used interchangeably. This close
correlation reflects the need for sufficient vegetative tissue (*i.e.*, rosette leaves) to produce the
resources for flowering and seed development. Because of the irreversible nature of the transition
from the vegetative to the reproductive stage in *Arabidopsis*, the coupling of these traits is crucial
for reproductive success. *Arabidopsis* mutants that flower with as few as three or four adult
leaves develop very few seeds and often show weakened growth habits. Uncoupling of flowering
time and rosette leaf number occurs in some early and late flowering time mutants (Pouteau *et al.*
2004) and in response to treatment with nitrogen dioxide (Takahashi and Morikawa 2014);
however, the mechanistic underpinnings for this uncoupling remain unknown.
Studies in several organisms suggest that AGO proteins are chaperoned by HSP90. HSP90
physically interacts with AGO proteins in yeast (Wang *et al.* 2013; Okazaki *et al.* 2018), flies
(Iwasaki *et al.* 2010; Miyoshi *et al.* 2010; Gangaraju *et al.* 2011), humans (Johnston *et al.* 2010;
Gangaraju *et al.* 2011), *Tetrahymena* (Woehrer *et al.* 2015), and plants (Iki *et al.* 2010, 2012).
Because miRNAs buffer environmental and genetic perturbations and AGO1 interacts with
HSP90, we set out to investigate the extent to which *AGO1* perturbation affects phenotypic
variation in isogenic *Arabidopsis* seedlings and buffers genetic variation in divergent
backgrounds, and *AGO1*-dependent loci overlap with HSP90-dependent loci. We find that *AGO1*
perturbation can significantly increase phenotypic variation in morphological and quantitative
traits in isogenic seedlings. *AGO1* perturbation also buffers the phenotypic effects of genetic
variation between two divergent backgrounds. However, none of the *AGO1*-buffered loci
overlapped with those buffered by HSP90, consistent with a prevalence of first-order epistatic
interactions relative to higher-order epistasis. Lastly, our detailed investigation of one such
buffered locus reveals that the coupling of the fitness-relevant traits days to flowering and rosette
leaf number relies on different molecular players in these commonly used strains of *Arabidopsis*.

124
Results
Genetic perturbation of *AGO1* increases phenotypic variation in isogenic *Arabidopsis*
seedlings
To determine if perturbation of *AGO1* leads to increased phenotypic variation in isogenic
seedlings, we examined several morphological and quantitative traits of two hypomorphic *ago1*
mutants, *ago1-46* (Smith *et al.* 2009), and *ago1-27* (Morel 2002), the former being a less severe
mutant than the latter. Ten-day old isogenic seedlings of *ago1-46* and *ago1-27* showed increased
phenotypic variation in morphological traits such as lesions in cotyledons (Mason *et al.* 2016),
rosette symmetry, and organ defects compared to isogenic wild-type seedlings (**Figure 1,**
Supplemental Table 1). As expected, the more severe *ago1-27* mutant showed more abnormal
phenotypes than the less severe *ago1-46* mutant. Next, we examined hypocotyl length in the
dark, a quantitative trait that shows increased variation in response to HSP90 perturbation
(Queitsch *et al.* 2002; Sangster *et al.* 2008b). Similar to our previous results (Queitsch *et al.*
2002; Sangster *et al.* 2008b), *ago1-27* dark-grown seedlings showed a different mean value (p <
$2.3e^{-16}$, Wilcoxon test) and significantly greater variance of hypocotyl length than wild-type
seedlings (p = 0.0002, Levene's test) (**Figure 1B, Supplemental Table 1**). The less severe *ago1-*
46 seedlings also showed a different mean value (p-value = $3.6e^{-05}$, Wilcoxon test) and greater
variance of hypocotyl length compared to wild-type seedlings (p-value = 0.00004, Levene's
test). Based on these results, AGO1 maintains phenotypic robustness and buffers developmental
noise among isogenic seedlings in a manner similar to HSP90.
145
AGO1 **buffers genetic variation independent of HSP90**
We next tested whether *AGO1* perturbation could reveal cryptic genetic variation and whether
AGO1-dependent loci overlapped with those buffered by *HSP90*. To do so, we introgressed the
hypomorphic *ago1-27* allele into Col-0 lines with single chromosome substitutions from another,
genetically divergent *Arabidopsis* strain, Landsberg *erecta* (L*er)* (STAIRS, STepped Aligned
Inbred Recombinant Strains, **Figure 2A, Supplemental Table 2**). STAIRS lines have been
generated for chromosomes 1, 3 and 5 (Koumproglou *et al.* 2002). Since *AGO1* is located on
chromosome 1, we excluded these STAIRS lines from our analysis. For chromosomes 3 and 5,
we selected two STAIRS lines each (chr3; N9448 and N9459, chr5; N9472 and N9501). The
introgressed lines were genotyped to confirm the integrity of the respective L*er* segments.
156
We measured hypocotyl and root length, rosette diameter, and the closely correlated traits days
to flowering and rosette leaf number across many individual plants per line using a randomized
experimental design (**Supplemental Table 2**). We selected these traits because they are readily
measurable and show evidence of HSP90-buffered variation in our previous studies of Col-0/L*er*
mapping lines (Sangster et al. 2008a; Sangster et al. 2008b). Specifically, three previously
described *HSP90*-dependent loci within the L*er* segments of the tested STAIRS lines affect the
traits measured here (Sangster *et al.* 2008b; a).
164
AGO1 perturbation may alter the contribution of a cryptic genetic variant to a quantitative trait in
two ways: first, *AGO1* perturbation may reveal a genetic variant by increasing its contribution to
a trait; or second, *AGO1* perturbation may conceal a genetic variant by increasing the relative
contribution of others. Indeed, the phenomenon of revealing and concealing genetic variation has
been previously observed for HSP90 perturbation across many traits in *Arabidopsis* recombinant

inbred lines (Sangster *et al.* 2008a). In addition, genetic variation in the respective L*er* segments
may mask the phenotypic differences observed between Col-0 wild-type and the *ago1-27* mutant
that was generated in the Col-0 background (*i.e.*, L*er* segments may epistatically interact with
ago1-27). We observed all three scenarios of epistasis (**Figure 2B, C**, **Supplemental Figures**
2,3). Despite the strong evidence that HSP90 facilitates AGO1 function in many organisms,
including plants, no overlap of *AGO1*-dependent loci with HSP90-dependent loci was observed.
176
Perturbation of *AGO1* uncouples flowering time and rosette leaf number in a background-
specific manner
One AGO1-buffered locus showed dramatically different effects on the two closely correlated
traits days to flowering and rosette leaf number (**Figure 3A-C**). *Arabidopsis* plants develop
about one rosette leaf per day until flowering is initiated. On average, Col-0 wild-type plants
initiated flowering ~5 days later and have ~5 more leaves than the STAIRS line 9472 that carries
a L*er* segment on chromosome 5 (coordinates 1 – 9,479,000bp). This result was expected
because the L*er* segment in this STAIRS line encompasses the *FLOWERING LOCUS C (FLC)*
gene (**Figure 3D**) which is non-functional in the L*er* strain (Michaels *et al.* 2003; Liu *et al.*
2004). *FLC* is a strong repressor of flowering (Whittaker and Dean 2017). In the Col-0
background, *ago1-27* plants initiated flowering ~9 days later and have ~2 more leaves, albeit the
traits were less tightly correlated than in wild type (**Figure 3C,** compare blue and green dots). In
stark contrast, in the STAIRS 9472 background, *ago1-27* plants showed no change in the number
of days to flowering; however, these plants showed dramatically fewer leaves at the onset of
flowering, developing on average only ~4 leaves. In fact, the severity of the rosette leaf number
phenotype of STAIRS9472;*ago1-27* was comparable to that observed in loss-of-function early
flowering mutants (Pouteau *et al.* 2004; Undurraga *et al.* 2012). In short, *AGO1* perturbation in
the STAIRS line specifically affected the trait rosette leaf number while not affecting the trait
days to flowering.
196
The close correlation of the traits days to flowering and rosette leaf number traits relies on
***FLC* in the Col-0 background**
The L*er* fragment in STAIRS9472 encompasses several known flowering time genes, including
FLC which delays flowering by repressing the gene *FLOWERING LOCUS T (FT)*. *FLC*
expression is repressed when plants are exposed to cold temperatures for a prolonged period of
time (*i.e.*, vernalization or winter period), allowing *FT* expression and onset of flowering
(Andrés and Coupland 2012; Whittaker and Dean 2017). Genetic variation in *FLC* and in
FRIGIDA (FRI), a positive regulator of *FLC*, accounts for the vast majority of differences in
flowering time across *Arabidopsis* strains (Shindo *et al.* 2005; Kim *et al.* 2009; Bloomer and
Dean 2017). Many *Arabidopsis* strains do not require vernalization to initiate flowering because
they carry *FLC* mutations, as is the case for L*er*, or *FRI* mutations, as is the case for Col-0. The
STAIRS9472 line carries the non-functional L*er FLC* allele.
209
We wondered if the lack of functional *FLC* in STAIRS9472;*ago1-27* contributed to its unusual
phenotype. To test this possibility, we examined the consequences of repressing *FLC* through
vernalization for both flowering time traits in Col-0 wild-type, STAIRS9472, Col-0 *ago1-27* and
STAIRS9472;*ago1-27* plants (**Supplemental Figure 3**). Vernalization did not erase the
difference in rosette leaf number between Col-0 *ago1-27* and STAIRS9472;*ago1-27* plants, with
the latter still showing significantly fewer leaves ($p = 5.704e^{-12}$ Wilcoxon test). However,

vernalization uncoupled both flowering time traits in an *AGO1*-dependent manner in the Col-0
background. Although vernalized *ago1-27* plants initiated flowering ∼5 days later than
vernalized Col-0 wild-type plants, they had ∼4 fewer leaves rather than more leaves. We
conclude that the close association of days to flowering and rosette leaf number in the Col-0
background requires the presence of functional *FLC* and *AGO1*. Perturbation of *AGO1* alone
diminished the close correlation between both traits but did not reverse it. L*er* and other natural
FLC mutants must have rewired flowering time pathways such that the traits days to flowering
and rosette leaf number remain closely correlated in the absence of functional *FLC*.
224
MIR156 polymorphisms are unlikely to cause *AGO1*-dependent phenotype
To identify the causal polymorphism(s) underlying the *AGO1*-dependent STAIRS9472
phenotype, we examined other genes within the L*er* segment with functions in flowering time
(Song *et al.* 2013, 2015; Spanudakis and Jackson 2014) and candidate polymorphisms between
Col-0 and L*er* (Nordborg *et al.* 2005; Borevitz *et al.* 2007; Ossowski *et al.* 2008) (**Figure 3D**).
We measured expression of these candidate genes among the four genotypes Col-0 wild-type,
STAIRS9472, Col-0 *ago1-27* and STAIRS9472;*ago1-27*; for the three *MIR156* genes (e, d, f),
and *MIR172b*, we measured expression of major target genes (Ji *et al.* 2011). As expected, *FLC*
expression was barely detectable in STAIRS9472 and STAIRS9472; *ago1-27* plants (**Figure
3D**), consistent with the known disruption of *FLC* in L*er* (Michaels *et al.* 2003; Liu *et al.* 2004).
FLC expression increased in Col-0 *ago1-27* plants relative to Col-0 wild-type plants, consistent
with the late flowering phenotype of the former genotype. As a general trend, target genes of
MIR156 increased in expression in the STAIRS *ago1-27* background compared to target gene
expression in the Col-0 *ago1-27* background, suggesting that *MIR156* may be less functional in
the STAIRS line. *MIR156* represses the expression of several *SQUAMOSA PROMOTER
BINDING LIKE* (SPL) transcription factors (miR156-SPL module, **Figures 3E, 5D**) that regulate
flowering by activating and repressing other transcription factors and miRNAs (Aukerman and
Sakai 2003; Yamaguchi *et al.* 2009; Wu *et al.* 2009). Overexpression of *MIR156* leads to delayed
onset of flowering with many more rosette leaves (Wu *et al.* 2009; Xu *et al.* 2016), suggesting
that less functional *MIR156* may diminish rosette leaf number.
245
We searched for L*er*-specific polymorphisms in the *MIR156* genes in available genome
assemblies and found a predicted single nucleotide polymorphism (SNP) within the loop of
MIR156f. Resequencing of all three *MIR156* genes confirmed this SNP and identified an
additional deletion of 14 nucleotides near the base of the stem loop. As the *MIR156* genes are
highly conserved in the plant kingdom (Cuperus *et al.* 2011; Luo *et al.* 2013), we examined their
natural variation among other *Arabidopsis* strains, sequencing an additional 55 strains. Of all
sequenced strains, 42 carried the L*er*-specific C-to-T SNP, one carried a C-to-G SNP, and 32
carried the 14-nt deletion (**Supplemental Figure 2**). The presence or absence of the deletion was
highly correlated with the presence or absence of the SNP (R^2 = 0.3506, p = 0.0007, Pearson
correlation test). To address whether either one or both L*er*-specific *MIR156f* polymorphisms
affect rosette leaf number, we tested for association with this trait across these accessions
(phenotypic data from (Lempe *et al.* 2005)). No association was found. Although this result did
not rule out the *MIR156f* polymorphisms as the causative *AGO1*-dependent alleles, it made it less
likely that these polymorphisms would explain the unusual STAIRS9472;*ago1-27* phenotype.
260

Identifying the *AGO-1* dependent L*er*-specific polymorphism with a bulk segregant analysis.

To identify the L*er*-specific variant(s) causing the observed trait uncoupling in STAIRS9472; *ago1-27* plants, we used a classic bulk segregant analysis followed by high-throughput sequencing (Cuperus *et al.* 2010; Sun and Schneeberger 2015). To generate a segregating population for the tested alleles, we crossed STAIRS9472 with *ago1-27*, and allowed for selfing to generate F_2 seeds. F_2 plants were measured for days to flowering and the number of rosette leaves at this point (**Figure 4A**). From this F_2 experiment, we pooled plants based on phenotype, defining the STAIRS9472;*ago1-27* phenotype as plants with 6 or fewer rosette leaves (**Figures 3A-C, 4A**), and isolated their DNA. We combined equal DNA amounts for 100 plants with the *AGO1*-dependent STAIRS9472 phenotype and 100 plants with higher numbers of rosette leaves. Using short-read sequencing, we aligned reads to the relevant chromosome 5 segment using SHOREmap (Sun and Schneeberger 2015), relying on the many known polymorphisms between L*er* and Col-0 to distinguish L*er*- and Col-0-specific reads. If successful, bulk segregant analysis will show increasing enrichment of homozygosity near the causal locus, with the causal locus at the center of a peak region (Salathia *et al.* 2007; Schneeberger *et al.* 2009; Cuperus *et al.* 2010; Sun and Schneeberger 2015). This mapping approach works best if variation at a single locus causes a segregating phenotype, and if phenotypes can be clearly distinguished from each another in order to pool samples with high confidence.

Although our phenotype of interest was quantitative in nature, *i.e.*, a range of leaf numbers rather than an absence or presence of a feature, we observed a skew towards L*er* alleles on chromosome 5 with a SHOREmap peak region at chr5:7,600,000 to chr5:7,800,000 (**Figure 4B**). Of the known flowering time-associated genes, only one fell in this peak region, *HUA2* (AT5G23150). Some L*er* backgrounds, including the STAIRS9742 line, carry a premature stop codon mutation in *HUA2*, likely disrupting function (Chen and Meyerowitz 1999; Doyle *et al.* 2005; Zapata *et al.* 2016). *HUA2* function is less well characterized than that of other flowering time genes; however, *hua2* mutants in a Col-0 background show reduced *FLC* levels and fewer rosette leaves at onset of flowering (Doyle *et al.* 2005) . *MIR156f* did not reside in the peak region, consistent with the previously described lack of genotype-phenotype association (**Figure 4B, C**).

To confirm that loss of functional *HUA2* was responsible for the *AGO1*-dependent phenotype in STAIRS9472, we used a Col-0-derived *hua2* mutant allele, *hua2-4*, and generated a double mutant *hua2-4;ago1-27* in the Col-0 background. We predicted that this homozygous double mutant would exhibit the uncoupling of days to flowering and rosette leaf number traits observed in the STAIRS9472; *ago1-27* line. Using a segregating F_2 population, we simultaneously measured days to flowering and rosette leaf number, and genotyped each plant (**Figure 5A-C**). The *hua2-4* single mutant plants and the *hua2-4;ago1-27* double mutant plants showed no significant difference in days to flowering but rosette leaf number was markedly reduced in double mutant plants, recapitulating our original finding with STAIRS9472;*ago1-27* plants. The observed uncoupling of these traits was independent of *FLC* which is not disrupted in the *hua2-4;ago1-27* double mutant. This result suggests that the L*er*-specific, non-functional *hua2* allele may compensate for the L*er*-specific *FLC* disruption, thereby maintaining the close association of days to flowering and rosette leaf number traits.

***HUA2* effects on gene expression suggest *SPL4* as a likely *HUA2* target.**
To understand in more detail how *HUA2* affects the complex flowering gene network, we
conducted RNA-seq experiments examining wild-type Col-0, single mutants *hua2-4* and *ago1-*
27 and *hua2-4;ago1-27* double mutant seedlings. As expected, *ago1-27* mutants and Col-0 wild-
type showed differential expression of many miRNA target genes (**Supplemental Table 6**). The
expression of the known *HUA2* targets *FLC* and *FLOWERING LOCUS M* (*FLM, MAF1*) was
reduced in both the single *hua2-4* mutant and the *hua2-4;ago1-27* double mutant seedlings,
excluding them as sources of the *AGO1*-dependent phenotype.
315
However, the comparison of *ago1-27* and *hua2-4;ago1-27* plants showed strong upregulation of
SPL4 expression in the latter (**Figure 5D**), consistent with our finding that *SPL4* was strongly
upregulated STAIRS9472;*ago1-27* (**Figure 3E**). Other important flowering time genes were also
differentially expressed in *hua2-4;ago1-27* double mutant plants, including the master regulator
FT, LATE ELONGATED HYPOCOTYL(LHY), SUPPRESSOR OF OVEREXPRESSION OF
CONSTANS 1 (SOC1), AGAMOUS-LIKE 8 (AGL8, FRUITFUL), and *MIR159b* (**Figure 5D**).
These genes interact in complex ways to control the transition to flowering (**Figure 5E**). Because
HUA2 is involved in mRNA processing and splicing (Chen and Meyerowitz 1999; Cheng *et al.*
2003; Janakirama 2013), we speculate that *SPL4* may be one of its targets. *SPL4* has three splice
isoforms, and two of them (*SPL4-2, SPL4-3*) lack a miR156-binding site (Yang *et al.*, 2012).
Overexpression of *SPL4-1*, the splice form with the miR156 binding site, does not affect days to
flowering but decreases rosette leaf number. In contrast, overexpression of *SLP4-2* or *SPL4-3*
decreases days to flowering and reduces rosette leaf number (Yang *et al.* 2012). In a *hua2;ago1*
double mutant background the balance of *SPL4* splice forms may be altered, which together with
the absence of functional AGO1 disrupts the close correlation of days to flowering and rosette
leaf number.
332
Discussion
Here we show that AGO1, the principal player in miRNA-mediated control of gene expression in
plants, buffers micro-environmental variation and maintains developmental stability in isogenic
Arabidopsis seedlings. Compared to wild-type Col-0 plants, *ago1* mutant seedlings showed more
lesions on cotyledons (Mason *et al.* 2016), more rosette symmetry defects and abnormal organs,
and increased variation in hypocotyl length of dark-grown seedlings. Given the crucial role that
miRNAs play in plant development, these results are not altogether surprising. MicroRNAs can
impact developmental stability, *i.e.*, the accuracy with which a given genotype produces a trait in
a particular environment, in various ways (Hornstein and Shomron 2006; Voinnet 2009). For
example, miRNAs can buffer gene expression noise as part of incoherent feedforward loops, in
which a transcription factor will activate both the expression of a target gene X and a miRNA,
with the latter repressing target gene X (Hornstein and Shomron 2006; Voinnet 2009).
MicroRNAs enforce developmental patterning decisions through mutual exclusion and spatial or
temporal restrictions in expression, *e.g.*, by suppressing fate-associated transcription factors in
neighboring cells or at a certain time in development (Hornstein and Shomron 2006; Voinnet
2009).
349
In plants, we previously reported increased variation for the same traits in isogenic seedlings
upon perturbation of the chaperone HSP90 (Queitsch *et al.* 2002; Sangster *et al.* 2008b),
consistent with the reported functional relationship of HSP90 and AGO1 (Iki *et al.* 2010, 2012;

Iwasaki *et al.* 2010, 2015; Naruse *et al.* 2018). An exception was the peculiar environmentally-
responsive lesions found on cotyledons in *ago1-27* seedlings (Mason *et al.* 2016). *HSP90* single
mutants produce far fewer seedlings with lesions than *ago1-27* mutants, and double mutants
show many more lesions than *ago1-27* single mutants, inconsistent with simple epistasis. Thus,
we previously suggested that AGO1 is a major, but largely HSP90-independent, factor in
providing environmental robustness to plants.
359
In addition to maintaining developmental stability, HSP90 buffers genetic variation in plants,
fungi, and animals, including humans (Zabinsky *et al.* 2019). The hypothesized mechanism by
which HSP90 overcomes the presence of genetic variation is the chaperone's well-characterized
function in protein folding and maturation (Sangster *et al.* 2004; Jarosz *et al.* 2010; Zabinsky *et*
al. 2019). This hypothesis is supported by the reported differences among disease-associated
protein variants chaperoned by HSP90 versus those chaperoned by HSP70 (Karras *et al.* 2017).
Moreover, across thousands of humans, kinases that are HSP90 clients tend to carry more amino
acid variants than non-client kinases, and these amino acid variants are predicted to be more
damaging to protein folding (Lachowiec *et al.* 2015).
369
In contrast, it is harder to envision a simple, direct mechanism by which AGO1 overcomes the
presence of genetic variation in either miRNAs or their targets, unless such buffering involves
AGO1's close relationship with HSP90 for the latter. Although we observed several instances in
which AGO1 perturbation revealed and concealed genetic variation in the same traits in which
we previously found HSP90-dependent variation (Sangster *et al.* 2008b), there was no overlap in
the genetic loci buffered by AGO1 and HSP90. While this result was consistent with our study of
the AGO1-dependent lesions (Mason *et al.* 2016), it raised anew the question as to how AGO1
may buffer genetic variation. In flies, proper expression of mir-9a, a miRNA acting on the
transcription factor *Senseless*, buffers genomic variation (Cassidy *et al.* 2013). Reducing mir-9a
regulation of *Senseless* leads to phenotypic variation in sensory cell fate in genetically diverse
flies, with candidate causal variants in genes that belong to the *Senseless*-dependent proneural
network governing sensory organ fate. In other words, in this case, AGO1-dependent buffering
via mir-9a occurs at the network level, consistent with the mechanisms by which miRNAs buffer
development stability and micro-environmental fluctuations.
384
To fully understand an instance of AGO1-dependent genetic variation, we focused on the
uncoupling of the traits days to flowering and rosette leaf number in STAIRS9472. Both traits
involve the mir156-SPL module and the key players *FLC* and *FRI* (**Fig. 5D**). We show that in
the Col-0 background the coupling of these traits requires functional *FLC* and *AGO1*. In
STAIRS9472, *FLC* is non-functional because the gene resides in the L*er*-introgression segment.
Without *FLC*, how are days to flowering and rosette leaf number coupled in L*er*? Using bulk
segregant analysis, we identified the non-functional *hua2* L*er*-allele as the likely causal AGO1-
dependent polymorphism. Indeed, we were able to recapitulate the uncoupling phenotype in the
hua2-4; ago1-27 double mutant in the Col-0 background.
It is noteworthy that this non-functional *HUA2* allele arose only recently and likely in the
laboratory (Zapata *et al.* 2016); there are several L*er* strains without this allele (Chen and
Meyerowitz 1999; Doyle *et al.* 2005; Zapata *et al.* 2016). These strains and other *Arabidopsis*
accessions with nonfunctional *FLC* genes must have acquired different polymorphisms to
maintain the coupling of both traits. The inbreeding nature of *Arabidopsis* and the propagation of

commonly used accessions like L*er* in controlled laboratory conditions readily allows fixation of
such polymorphisms. However, there are prior reports that natural variation in *HUA2* can affect
flowering time and plant morphology. The Sy-0 accession carries a gain-of function *HUA2* allele
that enhances *FLC* expression leading to larger rosettes, in addition to suppressing *AGAMOUS*
leading to indeterminate development of floral meristems (Wang *et al.* 2007). The Ws accession
carries a 12bp-deletion in *HUA2*, possibly weakening *HUA2* function (Doyle *et al.* 2005).
405
Our expression analysis offered some clues as to how *HUA2* may facilitate the close coupling of
days to flowering and rosette leaf number (**Figure 5D, E**). Comparing gene expression in *ago1-*
27 and *ago1-27; hua2-4* plants, we found that the mir156-SPL module gene *SPL4* was highly
upregulated in the double mutant. *SPL4* is expressed in three splice isoforms (*SPL4-1, SPL4-2,*
SPL4-3) with only one, *SPL4-1,* regulated by miR156 (Yang *et al.*, 2012). Overexpression of
SPL4-1 in transgenic plants does not alter days to flowering but reduces rosette leaf number. In
contrast, overexpression of *SLP4-2* or *SPL4-3* decreases both days to flowering and reduces
rosette leaf number (Yang *et al.* 2012). Because *HUA2* functions in mRNA processing and
splicing (Chen and Meyerowitz 1999; Cheng *et al.* 2003; Janakirama 2013), *SPL4* may be one of
its targets. Non-functional *HUA2* may lead to increased presence of the *SPL4-1* splice form,
which is exacerbated when mir156-dependent suppression of *SPL4-1* fails in the *ago1-27; hua2-*
4 double mutant, disrupting the close correlation of days to flowering and rosette leaf number
(**Figure 5E**). Thus, similar to the scenario in flies (Cassidy *et al.* 2013), AGO1 appears to buffer
genetic variation via microRNA-dependent network connections in plants. Disruption of the
miRNA-dependent network path in *ago1*-mutants can reveal genetic variants such as the non-
functional *HUA2* allele in other paths controlling the same trait (**Figure 5E**).
422
Taken together, our study holds several important lessons. First, AGO1 buffers phenotypic
variation in isogenic seedlings and genetic variation in genetically divergent ones. Second,
AGO1 does so independently of the chaperone HSP90 despite their close functional relationship,
suggesting that epistasis is largely a first-order phenomenon, specific to two interacting loci.
Indeed, this observed specificity of epistasis can extend to specific variants in pairwise
interacting loci. We previously showed that HSP90 can buffer genetic variation in Ste12, a
transcription factor that governs mating and invasion in yeast. However, HSP90-dependent
variants in Ste12 are rare; they reside in only two positions that are close to one another and alter
charge and DNA binding (Dorrity *et al.* 2018). This surprising specificity of epistatic interactions
calls into question the utility of current large-scale efforts to understand the phenotypic
contributions of epistasis by combining null mutants in human cells and in other organisms.
Third, our results provide a cautionary tale in interpreting phenocopies. Mutants in AGO1 and
HSP90 show highly similar phenotypes (Bohmert *et al.* 1998; Queitsch *et al.* 2002; Morel 2002;
Vaucheret *et al.* 2004; Sangster *et al.* 2007; Mason *et al.* 2016), and yet the underlying
mechanisms appear to differ, at least in part. Fourth, unlike HSP90, AGO1 suppresses the
phenotypic consequences of genetic variation by enabling miRNA-dependent network paths
rather than acting directly on variant-containing molecules, thereby extending the buffering
concept. Fifth and last, we show that key pathways can involve different molecular players even
in closely related strains of the same species. The uncoupling of highly correlated traits could be
a useful tool for plant breeders who want to improve one trait without compromising another
tightly coupled trait. Our study suggests miRNAs as good candidates for such targeted breeding
and engineering efforts.

Plant Materials and Growth Conditions: The following parental lines were used: Col-0, *ago1-27* in the Col-0 background, and STAIRS N9448, N9456, N9472, N9501(Morel 2002; Koumproglou *et al.* 2002). *ago1-27* plants were crossed into the STAIR lines and F_2's that carried the wild-type and *ago1-27* allele in both Col-0 and the STAIRS backgrounds were isolated. Selected F_2's and their progeny were used to perform the described experiments. For the hypocotyl and root length assays, the plants were grown on MS media containing 0.0005% MES hydrate, 0.004% vitamin solution, 3% phytoagar, and 1% sucrose.

Genotyping of F_2 plants: We used PCR to genotype the F_2's from each STAIRS – *ago1-27* cross. PCR conditions for *ago1-27* genotyping is as follows: 5' at 94 °C, followed by 35 cycles at 30 s at 94 °C, 30 s at 55 °C, 1 min at 72 °C. PCR product was then digested at 37°C with Bsp1286I, which cuts wild-type sequence. PCR conditions for MIR156F genotyping is as follows: 2' at 95°C, followed by 35 cycles at 30 s at 94°C, 50 s at 57°C, 40 s at 72°C.

Hypocotyl and root length assays: Seeds from different genotypes were plated on agar plates (10 seeds/per plate, equally spaced). The plates were stacked in racks to ensure vertical position, wrapped in aluminum foil, and transferred to 4˚C for five days to promote germination. They were then unwrapped, and exposed to light for two hours. After that, the plates were wrapped in aluminum foil again, to prevent further light exposure and were transferred to a 23˚C tissue culture incubator for seven days. The plants were grown vertically. After that, the plates were taken out, and photographed. The photographs were used to measure the seedlings' hypocotyls and roots using the ImageJ software (http://rsbweb.nih.gov/ij/).

Early morphology traits analysis: Seeds from the different genotypes were plated on agar (36 seeds/per plate). The plates were wrapped in aluminum foil and transferred to 4˚C for five days. Plates were unwrapped and transferred to long days (LD) in 23˚C tissue culture incubator for 10 days. The plants were grown horizontally. The plates were rotated every day to prevent biases due to location in the incubator. On the 10th day, the seedlings were scored for their morphological traits.

Flowering time experiments: Seeds from different genotypes were embedded in 1ml of 0.1% agar, and then stratified for five days at 4˚C. They were sown on soil in 36-pot trays. Flowering time was measured by scoring both the number of rosette leaves and days to flowering when the primary inflorescence of the plant had reached a height of 1cm. Flowering time experiments were performed in long days (LD, 16 hours of light, 8 hours of dark), at 23˚C.

Rosette diameter measurements: The diameter of the rosette was measured on the day that the primary inflorescence of the plant reached a height of 1cm.

Vernalization treatment: Seeds were stratified for five days at 4˚C and then sown on soil. They were allowed to grow for five days at 23˚C in LD or short days (SD) conditions and then transferred to 4˚C for forty days, according to recommendations from Sung *et al.*, 2006 (Sung et al. 2006).

Gene expression analysis: To determine the expression levels via qPCR, total RNA was
isolated from the aerial parts of 14-day old plants at ZT16 using the SV Total Isolation System
(Promega). RNA quality was determined using a Nanodrop and only high-quality samples
(A260/A230 > 1.8 and A260/A280> 1.8) were used for subsequent qPCR experiments. To
remove possible DNA contamination, RNA was treated with DNaseI (Ambion) for 60 minutes at
37˚C. We used the Transcriptor First Strand cDNA Synthesis Kit (Roche) for cDNA synthesis.
The qPCR primers were designed using the Universal Probe Library Assay Design Center tool
(Roche), and Primer3 (Untergasser et al. 2012). Specific amplification was confirmed before
conducting the qPCR experiments. The qPCRs experiments were carried out in 96-well plates
with a LightCycler480 (Roche) using SYBR green. The following program was used for the
amplification: pre-denaturation for 5 min at 95˚C, followed by 35 cycles of denaturation for 15 s
at 95˚C, annealing for 20 s at 55˚C, and elongation for 30 s at 72˚C. All qPCR experiments were
carried out with two biological replicates (independent samples harvested on different days), and
with three technical replicates per sample.
504
RNA-seq samples were prepared similarly as for qPCR, and then using the Illumina stranded
Tru-seq kit following the standard protocol. Samples were sequenced using the Nextseq550
platform. We used TopHat (v2.1.2) to align RNA-seq reads to the TAIR10 genome annotation
(Trapnell *et al.* 2009), htseq-count (v0.12.4) to calculate counts per gene (Anders *et al.* 2015),
using a minimum map quality of 10 and Cuffdiff (v2.2.1) to generate FPKMs (Trapnell *et al.*
2013), and DESeq2 to identify differentially expressed genes among genotypes (Love *et al.*
2014).
512
513   **Sequencing of miR156F, D and E in diverse *A. thaliana* strains:** The genes *MIR156F*,
514   *MIR156D* and *MIR156E* were amplified using the primers listed on **Supplemental Table 7**. PCR
515   products were sequenced by the Sanger method. The sequences were aligned using T-coffee.
516
Bulk segregant analysis - library preparation and sequencing: Approximately 400 F_2 plants
were sown, and leaf samples of equal size were collected from 100 plants that resembled the
STAIRS9472;*ago1-27* phenotype (6 leaves or fewer at flowering) and 100 plants with a greater
number of leaves. Individual plants were genotyped. In parallel, leaf samples were collected for
all genotypes. DNA was extracted using CTAB extraction (Weigel and Glazebrook 2002) and
quantified using the Qubit HS dsDNA assays. Libraries were quality checked on the Agilent
2100 bioanalyzer using a DNA 1000 chip (Agilent). Samples were pooled and libraries were
generated using the Nextera sample kit according to the manufacturer's instruction. DNA
concentration of the amplified libraries was measured with the DNA 1000 kit as well as the DNA
high-sensitivity kit for diluted libraries (both Agilent). Samples were sequenced on an Illumina
Nextseq in a 75-bp paired-end run.
528
529   **Bulk segregant analysis – data analysis:** Using the function SHORE import, raw reads were
530   trimmed or discarded based on quality values with a cutoff Phred score of +38. After correcting
531   the paired-end alignments with an expected insert size of 300 bp, we applied SHORE consensus
532   to identify variation among mutants and reference. We applied SHOREmap using the included
533   L*er*/Col-0 SNPs. Plot boost was applied to further define a mapping interval.
534

Data availability
All RNA sequencing reads and genomic sequence used for the SHOREmap analysis are
deposited in the NCBI Sequence Read Archive under the BioProject accession PRJNA836875.

Acknowledgements
This work was supported by the National Human Genome Research Institute Interdisciplinary
Training in Genomic Sciences (T32 training grant HG000035 to G.A.M.) and the National
Institute of Health (New Innovator award DP2OD008371 and NIGMS 1R35GM139532-01 to
C.Q.). We thank the editor and two anonymous reviewers for insightful comments and additional
references that have improved this manuscript.

Figure Legends:

Figure 1. Perturbation of AGO1 increases phenotypic variation among isogenic seedlings.
A) Early seedling trait measures for wild-type (Col-0 WT), *ago1-46*, and *ago1-27* seedlings.
Ten-day-old seedlings were scored for three different morphological traits: Lesions,
asymmetrical rosettes, and organ defects. The data represent two biological replicates (two
replicates, n = 144 for *ago1* mutants, and n = 216 for Col-0 WT, *p<0.05, ttest). (**B**) Hypocotyl
mean length and variance differ between wild-type and *ago1*-mutant seedlings. Hypocotyl length
was measured for seven-day old, dark-grown seedlings. *ago1-27* mutant seedlings showed
greater variance than Col-0 wild-type seedling in hypocotyl length (Levene's test, p< 1.0E-03; n
= 475 for *ago1-27*, n = 486 for Col-0 WT). Inset: boxplots of hypocotyl length means. Y-axis
represents hypocotyl length (mm), **p< 1.0E-15, Mann-Whitney Wilcoxon test.

Figure 2. Perturbation of AGO1 buffers genetic variation. (**A**) Experimental design to
examine the phenotypic consequences of genetic variation within the Stepped Aligned Inbred
Recombinant Strains (STAIRS) in the context of the *ago1-27* mutation. (**B**) Summary of
examined quantitative traits with evidence for *AGO1*-dependent or L*er*-dependent variation in
each tested STAIRS line. AGO1 perturbation reveals a cryptic genetic variant if this variant's
contribution to a quantitative trait can be detected only in an *ago1*-mutant background. AGO1
perturbation conceals a genetic variant if this variant's contribution to a quantitative trait can no
longer be detected in an *ago1*-mutant background. Genetic variation in the respective L*er*
segments can epistatically interact (*i.e.*, mask) the phenotypic differences observed between Col-
0 wild-type seedlings and *ago1-27* mutant seedlings in the Col-0 background. For STAIRS line
N9472, 78 seedlings were measured for hypocotyl length in the dark, for STAIRS lines N9448,
N9459, and N9501 100 seedlings were measured for this trait. At least 32 plants were measured
for all other traits. See Supplemental Tables 2 and 3 for traits values and assessment of
significance. (**C**) Two examples of *AGO1*-dependent and one example of L*er*-dependent genetic
variation are shown for three different traits. Blue, Col-0 WT; Yellow, STAIRS; Red, *ago1-27* in
the Col-0 background; Green, *ago1-27* in a STAIRS background.

**Figure 3. AGO1 perturbation uncouples the traits days to flowering and rosette leaf
number.** Plants for Col-0 WT, STAIRS9472, Col-0 *ago1-27* and STAIRS9472; *ago1-27* were
grown in a random block design in long days, n = 30-36. Days to flowering were recorded and
rosette leaf numbers at the onset of flowering were counted. Blue, Col-0 WT; Yellow,
STAIRS9472; Red, *ago1-27* in the Col-0 background; Green, *ago1-27* in the STAIRS9472

background (**A**) Days to flowering. The *ago-1* mutant flowered ~9 days later than Col-0 WT
(p=5.51E-12, Mann-Whitney Wilcoxon test); no significant difference was found between
STAIRS9472 and the *ago-1* mutant in the STAIRS9472 background (p=0.4714, Mann-Whitney
Wilcoxon test). The L*er* introgression in STAIRS9472 was epistatic to *ago1-27* *p<0.0001,
Mann-Whitney Wilcoxon test. (**B**) Rosette leaf number. Col-0 WT plants showed fewer leaves
than *ago1-27* mutant plants, consistent with the mutant's late flowering phenotype. In the
STAIRS 9472 background, *ago1-27* mutant plants showed no change in the number of days to
flowering; however, these plants developed significantly fewer leaves (p=3.45E-12, Mann-
Whitney Wilcoxon test). (**C**) Scatter plot of the two measured traits days to flowering and rosette
leaf number in the four tested genotypes. The traits were less well correlated in the *ago-1* mutant
in the Col-0 background (compare blue and green dots); however, the normally tight correlation
was fully lost in the STAIRS background (compare red and yellow dots). (**D**) Known flowering
time genes are residing within the L*er* chromosome 5 region of the STAIRS9472 line. (**E**)
Quantitative PCR measurements for candidate gene expression. *TFL1* and *SPL4* were
significantly increased in expression in the STAIRS9472;*ago1-27* background. 14-day old plant
tissue was collected at ZT16 (Zeitgeber 16; 16 hours after dawn). Mean expression data
represent two biological replicates, each with three technical replicates. Standard error is
indicated. (* = p < 0.05, ** = p < 0.005, T-test).
599
Figure 4: Bulk segregant analysis identifies the non-functional L*er hua2* allele as a
candidate *AGO1*-dependent locus. (**A**) F$_2$ plants from *ago1-27* x STAIRS9472 cross were
grown in long days, phenotypes were recorded, and plants were genotyped for the *ago1-27*
allele. For bulk segregant analysis, we selected plants that were homozygous for the *ago1-27*
mutation and flowered with six or fewer leaves (n=100), resembling the *AGO1*-dependent
STAIRS9472 phenotype. Representative F$_2$ *ago1-27* plants at flowering are shown. Scale bar =
1cm. (**B**) Bulk segregant analysis. Red dots represent L*er* allele frequencies on chromosome 5
(bp, x-axis). Allele frequencies (y-axis) were estimated as the fraction of reads supporting a L*er*
allele divided by the number of reads mapping to that locus. Dashed blue line represents sliding
window-based allele frequencies as estimated by SHOREmap. Dashed black line represents
window-based plot boost as estimated by SHOREmap. The L*er hua2-5* allele emerged as the
candidate *AGO1*-dependent locus because L*er* allele frequencies were highest at this locus
compared with other regions on chromosome 5. (**D**) F$_2$ plants homozygous for the *ago1-27*
mutation with six or fewer leaves at flowering were PCR genotyped for alleles at *FLC*, *HUA2*,
and *MIR156f* loci. The near perfect enrichment of L*er hua2-5* allele validates the result of our
bulk segregant analysis.
616
Figure 5: The *ago1-27; hua2-4* double mutant uncouples the traits days to flowering time
and rosette leaf number in the Col-0 background. An F$_2$ population segregating for the *ago1-*
27 and *hua2-4* mutant alleles was grown in long days. Days to flowering were recorded and
rosette leaf numbers at the onset of flowering were counted. Grey, Col-0 WT; white *ago1-27*
parent Red, *ago1-27;HUA2+/+* F2; Blue, *hua2-4* parent; Green, *ago1-27;hua2-4*. See
Supplemental Table 5 for further details (**A**) Plants carrying a homozygous *ago1-27* allele
flowered ~8.6 days later than Col-0 WT with ~16 leaves. Plants carrying a homozygous *hua2-4*
allele initiated flowering ~4.5 days earlier than Col-0 WT. As observed for STAIRS9472;*ago1-*
27, the *hua2-4* mutant allele was epistatic to *ago1-27*. *p<0.0283, **p<1.0E-06, Mann-Whitney
Wilcoxon test. The double mutant *ago1-27;hua2-4* plants showed a similar mean value but

greater trait variance. (**B**) The rosette leaf number phenotype of the double mutant *ago1-*
27;hua2-4 plants resembles that of the STAIRS9472;*ago1-27* line. *ago1-27;hua2-4* plants flower
with 5 leaves on average. (*p<0.0283, **p<1.0E-06, Mann-Whitney Wilcoxon test, for both A
and B) (**C**) Scatter plot with rosette leaf number on the x-axis and days to flowering on the y-
axis. Data are shown for F_2 plants that are homozygous for the *ago1-27* mutant allele and
segregate for the *hua2-4* mutant allele. (*D*) Known flowering time genes with differential
expression in the *ago1-27;hua2-4* double mutant as determined by RNA-seq. (**E**) Suggested
intersection of miRNAs and HUA2 in flowering time pathways. Depicted in red is the proposed
connection between *HUA2* and *SPL4* such that *SPL4* expression is regulated by both mir156 and
HUA2. In bold, genes that are overexpressed in the double mutant *ago1-27,hua-2* relative to the
single mutant *ago1-27*.
638
639 **Supplemental Figure 1**.Five phenotypes were measured for four STAIRS inbred lines, resulting
640 in cases of revealing, concealing and epistatic phenotypic changes. The five phenotypes
641 measured were days to flowering, rosette leaf number, rosette diameter, hypocotyl length and
642 root length.
643
644 **Supplemental Figure 2.** (A) Landsberg, Col-0 and the inbred strain STAIRS9472 we sequenced
645 across the MIR156F gene, supporting that there was both a SNP and a 14-nucleotide deletion in
646 both the Landsberg and STAIRS9472 strains compared to Col-0. (B) Further sequencing across
647 55 A. thaliana strains. Of the sequenced strains, , 42 carried the L*er*-specific C-to-T SNP, one
648 carried a C-to-G SNP, and 32 strains carried the 14-nt deletion.
649
650 **Supplemental Figure 3.** Seeds were stratified for five days at 4˚C and then sown on soil. They
651 were allowed to grow for five days at 23˚C in LD or short days (SD) conditions and then
652 transferred to 4˚C for forty days to vernalize plants. (**A**) Rosette leaf number at flowering with
653 vernalized plants in long day conditions. (**B**) Days to flowering with vernalized plants in long
654 day conditions.
655

References
Anders S., P. T. Pyl, and W. Huber, 2015 HTSeq--a Python framework to work with high-

throughput sequencing data. Bioinformatics 31: 166–169.

Andrés F., and G. Coupland, 2012 The genetic basis of flowering responses to seasonal cues.

Nat. Rev. Genet. 13: 627–639.

Aukerman M. J., and H. Sakai, 2003 Regulation of flowering time and floral organ identity by a

MicroRNA and its APETALA2-like target genes. Plant Cell 15: 2730–2741.

Axtell M. J., 2013 Classification and comparison of small RNAs from plants. Annu. Rev. Plant

Biol. 64: 137–159.

Bloomer R. H., and C. Dean, 2017 Fine-tuning timing: natural variation informs the mechanistic

basis of the switch to flowering in Arabidopsis thaliana. J. Exp. Bot. 68: 5439–5452.

Bohmert K., I. Camus, C. Bellini, D. Bouchez, M. Caboche, *et al.*, 1998 AGO1 defines a novel

locus of Arabidopsis controlling leaf development. EMBO J. 17: 170–180.

Bologna N. G., and O. Voinnet, 2014 The diversity, biogenesis, and activities of endogenous

silencing small RNAs in Arabidopsis. Annu. Rev. Plant Biol. 65: 473–503.

Borevitz J. O., S. P. Hazen, T. P. Michael, G. P. Morris, I. R. Baxter, *et al.*, 2007 Genome-wide

patterns of single-feature polymorphism in Arabidopsis thaliana. Proc. Natl. Acad. Sci.

U. S. A. 104: 12057–12062.

Cassidy J. J., A. R. Jha, D. M. Posadas, R. Giri, K. J. T. Venken, *et al.*, 2013 miR-9a minimizes

the phenotypic impact of genomic diversity by buffering a transcription factor. Cell 155:

1556–1567.

Chen X., and E. M. Meyerowitz, 1999 HUA1 and HUA2 are two members of the floral homeotic

AGAMOUS pathway. Mol. Cell 3: 349–360.

Cheng Y., N. Kato, W. Wang, J. Li, and X. Chen, 2003 Two RNA binding proteins, HEN4 and

HUA1, act in the processing of AGAMOUS pre-mRNA in Arabidopsis thaliana. Dev.

Cell 4: 53–66.

Cuperus J. T., T. A. Montgomery, N. Fahlgren, R. T. Burke, T. Townsend, *et al.*, 2010

Identification of MIR390a precursor processing-defective mutants in Arabidopsis by

direct genome sequencing. Proc. Natl. Acad. Sci. U. S. A. 107: 466–471.

Cuperus J. T., N. Fahlgren, and J. C. Carrington, 2011 Evolution and Functional Diversification

of MIRNA Genes. THE PLANT CELL ONLINE tpc.110.082784-.

Dong Q., B. Hu, and C. Zhang, 2022 MicroRNAs and their roles in plant development. Front.

Plant Sci. 13: 824240.

Dorrity M. W., J. T. Cuperus, J. A. Carlisle, S. Fields, and C. Queitsch, 2018 Preferences in a

trait decision determined by transcription factor variants. Proc. Natl. Acad. Sci. U. S. A.

115: E7997–E8006.

Doyle M. R., C. M. Bizzell, M. R. Keller, S. D. Michaels, J. Song, *et al.*, 2005 HUA2 is required

for the expression of floral repressors in Arabidopsis thaliana. Plant J. 41: 376–385.

Gangaraju V. K., H. Yin, M. M. Weiner, J. Wang, X. A. Huang, *et al.*, 2011 Drosophila Piwi

functions in Hsp90-mediated suppression of phenotypic variation. Nat. Genet. 43: 153–

158.

Hilgers V., N. Bushati, and S. M. Cohen, 2010 Drosophila microRNAs 263a/b confer robustness

during development by protecting nascent sense organs from apoptosis. PLoS Biol. 8:

e1000396.

Hornstein E., and N. Shomron, 2006 Canalization of development by microRNAs. Nat. Genet.

38 Suppl: S20-4.

Iki T., M. Yoshikawa, M. Nishikiori, M. C. Jaudal, E. Matsumoto-Yokoyama, *et al.*, 2010 In

vitro assembly of plant RNA-induced silencing complexes facilitated by molecular

chaperone HSP90. Mol. Cell 39: 282–291.

Iki T., M. Yoshikawa, T. Meshi, and M. Ishikawa, 2012 Cyclophilin 40 facilitates HSP90-

mediated RISC assembly in plants. EMBO J. 31: 267–278.

Iwasaki S., M. Kobayashi, M. Yoda, Y. Sakaguchi, S. Katsuma, *et al.*, 2010 Hsc70/Hsp90

chaperone machinery mediates ATP-dependent RISC loading of small RNA duplexes.

Mol. Cell 39: 292–299.

Iwasaki S., H. M. Sasaki, Y. Sakaguchi, T. Suzuki, H. Tadakuma, *et al.*, 2015 Defining

fundamental steps in the assembly of the Drosophila RNAi enzyme complex. Nature 521:

533–536.

Janakirama P., 2013 Functional characterization of the HUA2 gene family in Arabidopsis

thaliana

Jarosz D. F., M. Taipale, and S. Lindquist, 2010 Protein homeostasis and the phenotypic

manifestation of genetic diversity: principles and mechanisms. Annu. Rev. Genet. 44:

189–216.

Jarosz D. F., and S. Lindquist, 2010 Hsp90 and environmental stress transform the adaptive

value of natural genetic variation. Science 330: 1820–1824.

Ji L., X. Liu, J. Yan, W. Wang, R. E. Yumul, *et al.*, 2011 ARGONAUTE10 and ARGONAUTE1

Regulate the Termination of Floral Stem Cells through Two MicroRNAs in Arabidopsis,

(L.-J. Qu, Ed.). PLoS Genet. 7: e1001358.

Johnston M., M.-C. Geoffroy, A. Sobala, R. Hay, and G. Hutvagner, 2010 HSP90 protein

stabilizes unloaded Argonaute complexes and microscopic P-bodies in human cells. Mol.

Biol. Cell 21: 1462–1469.

Karras G. I., S. Yi, N. Sahni, M. Fischer, J. Xie, *et al.*, 2017 HSP90 shapes the consequences of

human genetic variation. Cell 168: 856-866.e12.

Kim D.-H., M. R. Doyle, S. Sung, and R. M. Amasino, 2009 Vernalization: winter and the

timing of flowering in plants. Annu. Rev. Cell Dev. Biol. 25: 277–299.

Koumproglou R., T. M. Wilkes, P. Townson, X. Y. Wang, J. Beynon, *et al.*, 2002 STAIRS: a

new genetic resource for functional genomic studies of Arabidopsis. Plant J. 31: 355–

364.

Lachowiec J., T. Lemus, E. Borenstein, and C. Queitsch, 2015 Hsp90 promotes kinase evolution.

Mol. Biol. Evol. 32: 91–99.

Lachowiec J., G. A. Mason, K. Schultz, and C. Queitsch, 2018 Redundancy, Feedback, and

Robustness in the Arabidopsis thaliana BZR/BEH Gene Family. Front. Genet. 9: 523.

Lempe J., S. Balasubramanian, S. Sureshkumar, A. Singh, M. Schmid, *et al.*, 2005 Diversity of

flowering responses in wild Arabidopsis thaliana strains. PLoS Genet. 1: 109–118.

Lempe J., J. Lachowiec, A. M. Sullivan, and C. Queitsch, 2013 Molecular mechanisms of

740   robustness in plants. Curr. Opin. Plant Biol. 16: 62–69.

741   Li X., J. J. Cassidy, C. A. Reinke, S. Fischboeck, and R. W. Carthew, 2009 A MicroRNA

742   imparts robustness against environmental fluctuation during development. Cell 137: 273–

743   282.

744   Liu J., Y. He, R. Amasino, and X. Chen, 2004 siRNAs targeting an intronic transposon in the

745   regulation of natural flowering behavior in Arabidopsis. Genes Dev. 18: 2873–2878.

746   Love M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and dispersion

747   for RNA-seq data with DESeq2. Genome Biol. 15: 550.

748   Luo Y., Z. Guo, and L. Li, 2013 Evolutionary conservation of microRNA regulatory programs in

749   plant flower development. Dev. Biol. 380: 133–144.

750   Masel J., and M. L. Siegal, 2009 Robustness: mechanisms and consequences. Trends Genet. 25:

751   395–403.

752   Mason G. A., T. Lemus, and C. Queitsch, 2016 The mechanistic underpinnings of an ago1-

753   mediated, environmentally dependent, and stochastic phenotype. Plant Physiol. 170:

754   2420–2431.

755   Michaels S. D., Y. He, K. C. Scortecci, and R. M. Amasino, 2003 Attenuation of FLOWERING

756   LOCUS C activity as a mechanism for the evolution of summer-annual flowering

757   behavior in Arabidopsis. Proc. Natl. Acad. Sci. U. S. A. 100: 10102–10107.

Miyoshi T., A. Takeuchi, H. Siomi, and M. C. Siomi, 2010 A direct role for Hsp90 in pre-RISC

formation in Drosophila. Nat. Struct. Mol. Biol. 17: 1024–1026.

Morel J.-B., 2002 Fertile Hypomorphic ARGONAUTE (ago1) Mutants Impaired in Post-

Transcriptional Gene Silencing and Virus Resistance. THE PLANT CELL ONLINE 14:

629–639.

Naruse K., E. Matsuura-Suzuki, M. Watanabe, S. Iwasaki, and Y. Tomari, 2018 In vitro

reconstitution of chaperone-mediated human RISC assembly. RNA 24: 6–11.

Nordborg M., T. T. Hu, Y. Ishino, J. Jhaveri, C. Toomajian, *et al.*, 2005 The pattern of

polymorphism in Arabidopsis thaliana. PLoS Biol. 3: e196.

Okazaki K., H. Kato, T. Iida, K. Shinmyozu, J.-I. Nakayama, *et al.*, 2018 RNAi-dependent

heterochromatin assembly in fission yeast Schizosaccharomyces pombe requires heat-

shock molecular chaperones Hsp90 and Mas5. Epigenetics Chromatin 11.

https://doi.org/10.1186/s13072-018-0199-8

Ossowski S., K. Schneeberger, R. M. Clark, C. Lanz, N. Warthmann, *et al.*, 2008 Sequencing of

natural strains of Arabidopsis thaliana with short reads. Genome Res. 18: 2024–2033.

Pouteau S., V. Ferret, V. Gaudin, D. Lefebvre, M. Sabar, *et al.*, 2004 Extensive phenotypic

variation in early flowering mutants of Arabidopsis. Plant Physiol. 135: 201–211.

Queitsch C., T. A. Sangster, and S. Lindquist, 2002 Hsp90 as a capacitor of phenotypic variation.

Nature 417: 618–624.

Rohner N., D. F. Jarosz, J. E. Kowalko, M. Yoshizawa, W. R. Jeffery, *et al.*, 2013 Cryptic

variation in morphological evolution: HSP90 as a capacitor for loss of eyes in cavefish.

Science 342: 1372–1375.

Rutherford S. L., and S. Lindquist, 1998 Hsp90 as a capacitor for morphological evolution.

Nature 396: 336–342.

Salathia N., H. N. Lee, T. A. Sangster, K. Morneau, C. R. Landry, *et al.*, 2007 Indel arrays: an

affordable alternative for genotyping. Plant J. 51: 727–737.

Sangster T. A., S. Lindquist, and C. Queitsch, 2004 Under cover: causes, effects and implications

of Hsp90-mediated genetic capacitance. Bioessays 26: 348–362.

Sangster T. A., A. Bahrami, A. Wilczek, E. Watanabe, K. Schellenberg, *et al.*, 2007 Phenotypic

diversity and altered environmental plasticity in Arabidopsis thaliana with reduced Hsp90

levels. PLoS One 2: e648.

Sangster T. A., N. Salathia, H. N. Lee, E. Watanabe, K. Schellenberg, *et al.*, 2008a HSP90-

buffered genetic variation is common in Arabidopsis thaliana. Proceedings of the

National Academy of Sciences 105: 2969–2974.

Sangster T. A., N. Salathia, S. Undurraga, R. Milo, K. Schellenberg, *et al.*, 2008b HSP90 affects

the expression of genetic variation and developmental stability in quantitative traits.

Proceedings of the National Academy of Sciences 105: 2963–2968.

Schneeberger K., S. Ossowski, C. Lanz, T. Juul, A. H. Petersen, *et al.*, 2009 SHOREmap:

simultaneous mapping and mutation identification by deep sequencing. Nat. Methods 6:

550–551.

Schopf F. H., M. M. Biebl, and J. Buchner, 2017 The HSP90 chaperone machinery. Nat. Rev.

Mol. Cell Biol. 18: 345–360.

Shindo C., M. J. Aranzana, C. Lister, C. Baxter, C. Nicholls, *et al.*, 2005 Role of FRIGIDA and

FLOWERING LOCUS C in determining variation in flowering time of Arabidopsis.

Plant Physiol. 138: 1163–1173.

Smith M. R., M. R. Willmann, G. Wu, T. Z. Berardini, B. Möller, *et al.*, 2009 Cyclophilin 40 is

required for microRNA activity in Arabidopsis. Proc. Natl. Acad. Sci. U. S. A. 106:

5424–5429.

Song Y. H., S. Ito, and T. Imaizumi, 2013 Flowering time regulation: photoperiod- and

temperature-sensing in leaves. Trends Plant Sci. 18: 575–583.

Song Y. H., J. S. Shim, H. A. Kinmonth-Schultz, and T. Imaizumi, 2015 Photoperiodic

flowering: time measurement mechanisms in leaves. Annu. Rev. Plant Biol. 66: 441–464.

Spanudakis E., and S. Jackson, 2014 The role of microRNAs in the control of flowering time. J.

Exp. Bot. 65: 365–380.

Sun H., and K. Schneeberger, 2015 SHOREmap v3.0: fast and accurate identification of causal

mutations from forward genetic screens. Methods Mol. Biol. 1284: 381–395.

Takahashi M., and H. Morikawa, 2014 Nitrogen dioxide accelerates flowering without changing

the number of leaves at flowering in Arabidopsis thaliana. Plant Signal. Behav. 9:

e970433.

Trapnell C., L. Pachter, and S. L. Salzberg, 2009 TopHat: discovering splice junctions with

RNA-Seq. Bioinformatics 25: 1105–1111.

Trapnell C., D. G. Hendrickson, M. Sauvageau, L. Goff, J. L. Rinn, *et al.*, 2013 Differential

analysis of gene regulation at transcript resolution with RNA-seq. Nat. Biotechnol. 31:

46–53.

Undurraga S. F., M. O. Press, M. Legendre, N. Bujdoso, J. Bale, *et al.*, 2012 Background-

dependent effects of polyglutamine variation in the Arabidopsis thaliana gene ELF3.

Proc. Natl. Acad. Sci. U. S. A. 109: 19363–19367.

Vaucheret H., F. Vazquez, P. Crété, and D. P. Bartel, 2004 The action of ARGONAUTE1 in the

miRNA pathway and its regulation by the miRNA pathway are crucial for plant

development. Genes Dev. 18: 1187–1197.

Voinnet O., 2009 Origin, biogenesis, and activity of plant microRNAs. Cell 136: 669–687.

Wang Q., U. Sajja, S. Rosloski, T. Humphrey, M. C. Kim, *et al.*, 2007 HUA2 caused natural

variation in shoot morphology of A. thaliana. Curr. Biol. 17: 1513–1519.

Wang Y., R. Mercier, T. C. Hobman, and P. LaPointe, 2013 Regulation of RNA interference by

Hsp90 is an evolutionarily conserved process. Biochim. Biophys. Acta 1833: 2673–2681.

Weigel D., and J. Glazebrook, 2002 *Arabidopsis: A Laboratory Manual*. CSHL Press.

Whitacre J. M., 2012 Biological robustness: paradigms, mechanisms, and systems principles.

Front. Genet. 3: 67.

Whittaker C., and C. Dean, 2017 The FLC locus: A platform for discoveries in epigenetics and

adaptation. Annu. Rev. Cell Dev. Biol. 33: 555–575.

Woehrer S. L., L. Aronica, J. H. Suhren, C. J.-L. Busch, T. Noto, *et al.*, 2015 A Tetrahymena

Hsp90 co-chaperone promotes siRNA loading by ATP-dependent and ATP-independent

mechanisms. EMBO J. 34: 559–577.

Wu G., M. Y. Park, S. R. Conway, J.-W. Wang, D. Weigel, *et al.*, 2009 The sequential action of

miR156 and miR172 regulates developmental timing in Arabidopsis. Cell 138: 750–759.

Xu M., T. Hu, J. Zhao, M.-Y. Park, K. W. Earley, *et al.*, 2016 Developmental Functions of

miR156-Regulated SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL)

Genes in Arabidopsis thaliana, (M. Tsiantis, Ed.). PLoS Genet. 12: e1006263.

Yamaguchi A., M.-F. Wu, L. Yang, G. Wu, R. S. Poethig, *et al.*, 2009 The microRNA-regulated

SBP-Box transcription factor SPL3 is a direct upstream activator of LEAFY,

FRUITFULL, and APETALA1. Dev. Cell 17: 268–278.

Yang X., H. Zhang, and L. Li, 2012 Alternative mRNA processing increases the complexity of

microRNA-based gene regulation in Arabidopsis. Plant J. 70: 421–431.

Yeyati P. L., R. M. Bancewicz, J. Maule, and V. van Heyningen, 2007 Hsp90 selectively

modulates phenotype in vertebrate development. PLoS Genet. 3: e43.

Zabinsky R. A., G. A. Mason, C. Queitsch, and D. F. Jarosz, 2019 It's not magic – Hsp90 and its

854   effects on genetic and epigenetic variation. Semin. Cell Dev. Biol. 88: 21–35.

855   Zapata L., J. Ding, E.-M. Willing, B. Hartwig, D. Bezdan, *et al.*, 2016 Chromosome-level

856   assembly of *Arabidopsis thaliana* L *er* reveals the extent of translocation and inversion

857   polymorphisms. Proc. Natl. Acad. Sci. U. S. A. 113: E4052–E4060.

858
859
860

Figure 1

Figure 2

Figure 3

A

1cm

4 leaves   5 leaves   6 leaves   8 leaves   9 leaves

B

Known flowering time associated genes

PNY *PRR7* *LM1* *TFL1* *MIR172b* *MYB33* *CHE* *FLC* *MIR156d* *EMF1* *MIR156e* *NFYA1/HAP2A* *FY* *ALG15* *CONSTANS* *FRL1* *TFL2* *HUA2* *PRR5* *FPF1* *MIR156f* *GID1C*
ELF6

Allele frequency (*Ler*)

1.0
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1

SHOREmap 3.0 with "plot boost"
51x coverage, n=100

1   2000001   4000001   6000001   8000001   10000001

Chromosome 5 position

C

■ Col-0 allele
■ *Ler* allele
■ Heterozygous

n=28
FLC genotype frequency

n=48
HUA2 genotype frequency

n=99
MIR156f genotype frequency

Figure 4

Figure 5