<u>Avian Influenza A Virus polymerase can utilise human ANP32 proteins to support</u>

<u>cRNA but not vRNA synthesis</u>

Olivia C. Swann, Amalie B. Rasmussen, Thomas P. Peacock, Carol M. Sheppard*,

Wendy S. Barclay*[1]

Department of Infectious Diseases, Faculty of Medicine, Imperial College London,

UK

*Joint corresponding author

1. Lead author

carol.sheppard08@imperial.ac.uk

w.barclay@imperial.ac.uk

<u>Abstract</u>

Host restriction limits the emergence of novel pandemic strains from the Influenza A

Virus avian reservoir. For efficient replication in mammalian cells, the avian influenza

RNA-dependent RNA polymerase must adapt to use human orthologues of the host

factor ANP32, which lack a 33 amino acid insertion relative to avian ANP32A. Here

we find that influenza polymerase requires ANP32 proteins to support both steps of

replication: cRNA and vRNA synthesis. Nevertheless, avian strains are only restricted

in vRNA synthesis in human cells. Therefore, avian polymerase can use human

ANP32 orthologues to support cRNA synthesis, without acquiring mammalian

adaptations. This implies a fundamental difference in the mechanism by which ANP32

proteins support cRNA vs vRNA synthesis.

Importance

In order to infect humans and cause a pandemic, avian influenza must first learn how

to use human versions of the proteins the virus hijacks for replication – instead of the

avian versions found in bird cells. One such protein is ANP32. Understanding the

details of how host proteins such as ANP32 support viral activity may allow the design

of new antiviral treatments that disrupt these interactions. In this work, we use cells

that lack ANP32 to unambiguously demonstrate ANP32 is needed for both steps

of influenza genome replication. Surprisingly however, we find that avian influenza can

use human ANP32 proteins for the first step of replication without any adaptation, but

only avian ANP32 for the second step of replication. This suggests ANP32 may have

an additional role in supporting the second step of replication, and it is this activity that

is specifically blocked when avian influenza infects human cells.

 <u>Introduction</u>

Influenza A viruses (IAVs) pose a pandemic risk: whilst the natural hosts of IAV are

aquatic birds, the virus is associated with sporadic zoonotic jumps, which may trigger

widespread disease in an immunologically naïve population (1). The enormous

consequences of such events are illustrated by the historical 1918 'Spanish flu'

pandemic, and the ongoing coronavirus pandemic caused by SARS-CoV-2.

Prior to emerging as a pandemic strain, zoonotic IAV must adapt to overcome multiple

host range barriers. One important restriction is adaptation of the RNA-dependent

RNA polymerase (FluPol) for efficient activity within the mammalian cellular

environment (1, 2). FluPol is a heterotrimer consisting of three viral proteins: PB1, PB2

and PA. Each of the eight viral genomic RNAs (vRNAs) of the IAV genome are

encapsidated by nucleoproteins (NP) and associate with a FluPol protomer in a viral

ribonucleoprotein complex (vRNP). During infection, FluPol drives both transcription

and replication of the viral genome (3). Transcription occurs in a primer-dependent

manner to produce capped and poly-adenylated positive sense viral mRNAs. Genome

replication occurs in a two-step process: first, negative sense vRNA is copied into a

full-length positive-sense cRNA intermediate which is packaged into complementary

RNPs (cRNPs) by acquiring an encapsidating polymerase and NP. Nascent vRNPs

are then produced from cRNPs (3, 4). cRNA and vRNA synthesis differ in *de novo*

initiation strategy. Whilst cRNA synthesis uses terminal initiation, vRNA synthesis uses

internal initiation followed by primer realignment that is dependent on an additional

trans-activating FluPol (5–8).

Structural studies suggest the functional flexibility of FluPol is possible due to rearrangements of peripheral domains outside the catalytic core (3). These depend on the presence or absence and nature of the bound RNA promoter, or upon dimerization with additional polymerase molecules (9–11). FluPol also co-opts host factors to co-ordinate its activity, some of which may stabilise different functional states of the polymerase (10, 12).

ANP32 proteins are a family of proviral host factors that are essential for influenza replication (13, 14). Species differences between ANP32 proteins underly host restriction of avian influenza polymerase in mammalian cells (15). Prototypical avian influenza polymerase bearing a glutamic acid at position 627 of the PB2 protein (hereafter referred to as FluPol 627E) cannot be supported by mammalian ANP32 proteins that lack the 33 amino acid insertion present in avian ANP32A orthologues, and accordingly show restricted replication in human cells. In contrast, FluPol bearing a single amino acid substitution at the PB2 627 position from glutamic acid to lysine (the quintessential mammalian-adapting mutation, FluPol 627K), can utilise mammalian ANP32 proteins that lack this insertion, and such viruses replicate to high titres in human cells (2, 15). Interestingly, although chicken ANP32A (chANP32A) shows enhanced binding to FluPol relative to human ANP32A (huANP32A), that is dependent on the 33 amino acid insertion, this pattern occurs independent of FluPol 627 signature, i.e., species-specific functionality is not dictated by differences in binding affinity (16–18).

Previous studies suggest that ANP32 proteins are specifically required to support vRNA synthesis, and accordingly that host restriction occurs at this step of replication.

Sugiyama et al found that purified huANP32A stimulated RNA replication from a short

cRNA but not vRNA template in *in vitro* replication assays (19). Moreover, studies

comparing infection in mammalian cells with strains bearing either FluPol 627E or

627K found that 627E was specifically restricted in vRNA synthesis, but not cRNP

stabilisation (20, 21). Nevertheless, other reports implicate ANP32 in both steps of

replication (11, 22). The recent structures of huANP32A and chANP32A in complex

with FluPol from Influenza C virus demonstrate that the N terminal Leucine Rich

Repeat (LRR) domain of ANP32 bridges a novel asymmetric dimer of two FluPol

enzymes. The unstructured C terminal Low Complexity Acidic Region (LCAR) domain

of ANP32 remains largely unresolved, apart from additional density sandwiched in a

groove formed between the two 627 domains. This dimer was interpreted as a

'replication complex', with the RNA-bound FluPol (FluPol$_R$) adopting a replication-

competent structure, and the additional apo-enzyme (FluPol$_E$) adopting a novel

conformation: poised to encapsidate the nascent RNA into an RNP complex.

Interestingly, this structure was obtained with FluPol$_R$ bound to a 47nt short vRNA,

i.e., primed for cRNA synthesis, which was not previously implemented in requiring

support by ANP32. Moreover, mutations introduced to disrupt the FluPol asymmetric

dimer interface resulted in a significant reduction in cRNA encapsidation. This,

combined with the logic that both cRNA and vRNA require encapsidation into RNPs

during replication, suggests that ANP32 may play a role in cRNA synthesis as well as

vRNA synthesis (11).

Here we use ANP32 knockout cell lines to clarify the role of ANP32 proteins in cRNA

synthesis. We establish an RNA fluorescence in situ (FISH) assay for directly

visualising cRNA and demonstrate that ANP32 proteins are essential for primary

cRNA synthesis in authentic infection, as well as under experimental conditions in

which vRNA synthesis is inhibited. Nevertheless, we find that avian FluPol 627E does

not show restricted cRNA accumulation in mammalian cells. This observation is

consistent regardless of whether $FluPol_R$, $FluPol_E$ or both FluPol molecules in the

replication complex bear the avian-like 627E signature. Moreover, the cRNPs

produced by $FluPol_R$ 627E are functional for onward replication. To conclude, our

study suggests that huANP32 proteins are sufficient to support avian FluPol cRNA

synthesis, and that host restriction acts specifically at the level of vRNA synthesis. This

suggests a fundamental difference in the mechanism by which the host factor ANP32

supports vRNA and cRNA synthesis.

Results

cRNA synthesis is inhibited in human cells lacking ANP32A/B

Human cells express three members of the ANP32 protein family: huANP32A, huANP32B and huANP32E (23). Previous work has demonstrated that huANP32A/B are functionally redundant for proviral activity, whilst huANP32E does not support FluPol activity (13, 14). To investigate the role of ANP32 proteins in cRNA synthesis, wild type human eHAP cells (eHAP WT) and human eHAP cells in which ANP32A/B have been ablated (eHAP dKO) (13) were infected with A/Puerto Rico/8/1934(H1N1) (PR8) and the accumulation of segment 4 (haemagglutinin (HA)) vRNA, cRNA and mRNA quantified using a tagged RT-qPCR approach (24). Following infection, there was no increase in either vRNA or cRNA levels over time in the absence of ANP32A/B, confirming ANP32 proteins are essential for replication (Figure 1A,B). Significantly, no increase in cRNA accumulation was observed over input, suggesting a direct role for ANP32 proteins in the pioneering round of cRNA synthesis. In contrast, HA mRNA transcripts increased 50-fold in dKO cells from 0 hours post infection (hpi) to 2hpi, illustrating that primary transcription does not require ANP32 proteins (Figure 1C). In a separate experiment, we analysed the accumulation of segment 6 (neuraminidase (NA)) vRNA, cRNA and mRNA at an early and late timepoint in eHAP WT and dKO cells, using an analogous tagged RT-qPCR approach. As with segment 4, at late timepoints a significant reduction in all three RNA species was observed (Figure 1D,E,F). At 3hpi, a ~10 fold reduction in the accumulation of cRNA was already apparent in the dKO cells, despite no difference in vRNA accumulation having yet occurred. Again, this suggested a direct role for huANP32A/B in supporting cRNA synthesis.

Human ANP32 proteins play a direct role in primary cRNA synthesis

To eliminate the possibility that ANP32 proteins are only required for vRNA synthesis,

and the lack of cRNA accumulation is an indirect effect, we made use of cRNP

stabilisation assays. cRNP stabilisation assays manipulate cellular conditions to allow

only the primary round of cRNA synthesis from incoming viral genomes (Vreede et al,

2004, Nilsson et al, 2017). vRNA synthesis, and consequently further secondary

rounds of cRNA synthesis, are inhibited. The assay involves pre-expressing FluPol

and NP, and then infecting cells with virus. Importantly, the FluPol that is exogenously

expressed is a catalytically dead mutant (PB1 D446Y) that will allow stabilisation of

nascent cRNA into catalytically inactive cRNPs. Cells are then drug treated with either

actinomycin D (ActD) or cycloheximide (CHX) during infection to inhibit the synthesis

of nascent viral proteins which would otherwise enable normal replication.

Initially, ActD-treated cRNP stabilisation assays were performed in eHAP WT and dKO

cells. ActD inhibits viral transcription, therefore in these assays only primary cRNA

synthesis occurs (Figure 2A). When analysed by RT-qPCR (Figure 2B,C,D) a highly

significant reduction in cRNA was observed in dKO cells as compared to WT cells. As

expected, no accumulation of NA vRNA or mRNA occurred over background levels

(controls lacking transfected PB1, dotted line), in either cell type. Equal transfection

efficiency was confirmed by western blot (Figure S1A).

We next established *in situ* assays for directly visualising IAV replication products to

allow single cell, spatial information to be collected alongside bulk assay readouts. To

achieve this, the RNA FISH assay RNAscope® was used (25). Probes were designed

to target either segment 6 negative sense RNA (NA vRNA probe) or positive sense

RNA (NA +RNA probe). The NA +RNA probe is unable to distinguish NA cRNA/mRNA

due to the minimal sequence differences between these two RNA species. To counter

this, simultaneous cRNP stabilisation and replication assays were performed with

ActD treatment, to inhibit viral transcription and allow +RNA staining to be attributed

to cRNA. Replication assays are performed as for cRNP stabilisation assays, however

active polymerase is pre-expressed, so that multiple rounds of cRNA and vRNA

synthesis take place. Validation of the approach is shown in Figure S1B-F. In an *in*

situ cRNP stabilisation assay, +RNA accumulation could be clearly detected 3hpi in

the nuclei of infected WT cells (Figure 2E). In contrast, whilst incoming vRNPs could

be observed in the nuclei of dKO cells, no +RNA staining was observed. This

corroborates the view that huANP32A/B are required for primary cRNA synthesis.

Our infection data (Figure 1) suggest that ANP32 proteins are not required for primary

transcription. Nevertheless, at later infection timepoints, reduced mRNA accumulation

is observed in dKO cells, compared to WT cells (Figure 1C,F). To confirm this is an

indirect effect due to reduced vRNP template, we chose to undertake a CHX-treated

cRNP stabilisation assay in eHAP wild type, dKO or in eHAP cells lacking expression

of all three huANP32 proteins: A, B and E (eHAP tKO). In this version of the assay,

both cRNA synthesis and primary transcription occur in the absence of vRNA

synthesis (Figure 2F). No accumulation of vRNA over input was observed in any of

the cell types over time (Figure 2G), as expected. In agreement with ActD cRNP

stabilisation assays (Figure 2C), a significant decrease in cRNA accumulation was

seen in both dKO and tKO cells compared to WT by 5hpi (Figure 2H). A small but

significant difference (p=0.0018) was observed between cRNA accumulation in the

dKO and tKO cells, implying huANP32E may support a low level of cRNP stabilisation.

In contrast, no significant difference was observed in mRNA accumulation in the

presence or absence of any ANP32 proteins from 0 to 5hpi (Figure 2I). This confirms

that primary transcription does not require ANP32 proteins. Comparable transfection

efficiency was confirmed via western blot (Figure S2G).

Avian polymerase is not restricted in cRNA synthesis in mammalian cells

Restriction of avian signature FluPol (FluPol 627E) in mammalian cells is attributed to

species differences in ANP32 proteins. Consequently, as we have confirmed that

ANP32 is required to support both cRNA and vRNA synthesis, we would expect FluPol

627E to be restricted in both steps of replication. Nevertheless, previous work maps

restriction to occur specifically at the level of vRNA synthesis (20, 21, 26). To

investigate this apparent contradiction, we used a pair of isogenic viruses based on

the avian strain A/turkey/England/50-92/1991 (H5N1) (5092) that differ only in the

residue at PB2 position 627: either the wild type PB2 627E (hereafter referred to as

5092E) or the humanising mutation PB2 E627K (5092K) (27).

First, we confirmed that 5092 polymerase is also dependent on ANP32 proteins for

cRNA synthesis both during authentic infection (Figure S2A,B,C) and in cRNP

stabilisation assays (Figure S2D,E,F). Next, we undertook simultaneous cRNP

stabilisation and replication assays in human cells, where only the shorter

huANP32A/B/E proteins, that are incompatible with FluPol 627E, are available. We

pre-expressed NP with either catalytically dead 5092 FluPol 627E or 627K (cRNP

stabilisation assay) or WT 5092 FluPol 627E or 627K (replication assay), followed by

infection with either 5092K or 5092E virus in the presence of ActD. Significantly, these

assays allow the effect of PB2 residue 627 in either $FluPol_R$ or $FluPol_E$ to be

differentiated. The resulting FluPol combinations for both cRNA and vRNA synthesis

are outlined in Figure 3A.

During the pioneering round of cRNA synthesis, measured in the cRNP stabilisation

assay, FluPol$_E$ is provided by the pre-expressed polymerase whilst FluPol$_R$ is brought

in on the vRNPs from the infecting virus. No vRNA accumulated in the cRNP

stabilisation assay, as expected (Figure 3B). Interestingly, neither the PB2 signature

of FluPol$_R$ nor FluPol$_E$ impacted cRNP stabilisation, with all four FluPol combinations

accumulating equivalent amounts of cRNA (Figure 3C). This demonstrates that avian

and human signature virus can undergo cRNA synthesis equally well in human cells,

i.e., avian FluPol is compatible with huANP32 proteins for cRNA synthesis and

stabilisation.

In the replication assay, samples with pre-expressed FluPol 627K accumulated

equivalent quantities of vRNA and cRNA, regardless of whether the incoming virus

was 5092K or 5092E (Figure 3Ai,ii,B,C). Similarly, samples with pre-expressed FluPol

627E were equally impaired in vRNA and cRNA synthesis, independent of infecting

virus signature (Figure 3Aiii,iv,B,C). It has previously been described that host

restricted FluPol is impaired in nuclear import, due to incompatibilities with the

importin-alpha isoforms present in human cells (28). However, this cannot explain the

data obtained here, as FluPol 627E was fully functional in supporting the pioneering

round of cRNA synthesis and stabilisation within the nucleus during cRNP stabilisation

assays. As replication assays allow multiple cycles of replication, the reduced cRNA

synthesis observed with pre-expressed FluPol 627E is likely due to the secondary

effect of reduced vRNP template. Exogenous expression of chANP32A was able to

rescue replication for all FluPol combinations (Figure 3D,E).

<u>Discussion</u>

Here we have used cell lines that lack expression of ANP32 proteins to unambiguously

confirm huANP32 is required to support cRNA synthesis. Nevertheless, we did not

observe restriction of avian virus cRNA synthesis in human cells, confirming that host

range restriction imparted by species differences in ANP32 acts at the level of vRNA

synthesis.

In the structure of the ANP32 and FluPol complex described by Carrique et al, the

ANP32 LRR domain bridges the FluPol dimer. The flexible LCAR remains largely

unresolved, although there is additional density present in a groove formed between

the two FluPol 627 domains. As the groove is acidic in FluPol 627E, the acidic

huANP32 LCAR is likely incompatible with this interaction, whereas the mixture of

acidic and basic residues in the 33 amino acid insertion of avian chANP32A

overcomes this block (11). Our data suggests this mismatch between the huANP32

LCAR and FluPol 627E can be tolerated for cRNA synthesis. This conclusion aligns

with the observation that addition of chANP32A to human cells does not stimulate

FluPol 627E cRNA synthesis (26).

The ANP32-stabilised replication complex is thought to be required to load an

encapsidating FluPol onto the promoter of the extruded strand of nascent c/vRNA.

Moreover, it has been proposed that the unstructured ANP32 LCAR domain may act

as a molecular whip, which recruits NP for loading onto the nascent RNA (11, 29).

Previous studies have suggested that host restricted avian FluPol can generate cRNA

in human cells, but that the resulting cRNPs are non-functional for onwards replication

(20, 22, 26). However, here we show that cRNA produced from $FluPol_R$ 627E is fully

functional for vRNA synthesis, provided the encapsidating FluPol bears 627K for

onward vRNA synthesis. Therefore, functional cRNPs are produced in the absence of

a compatible 627-LCAR interaction, suggesting host restriction does not act at the

level of cRNA encapsidation. In combination, our data therefore suggests that the 627-

LCAR interaction is required for a further mechanistic role that is specific to vRNA

synthesis. For example, a compatible 627-LCAR interaction could be of critical

importance to recruit the trans-activating FluPol (York et al., 2013).

In summary, this work has established that huANP32 is required for cRNA synthesis,

as well as vRNA synthesis, in human cells. Nonetheless, host restriction of avian

FluPol 627E acts specifically at the level of vRNA synthesis, as huANP32 paralogues

are sufficient for supporting avian cRNA synthesis and encapsidation. This suggests

a fundamental difference in the mechanism by which ANP32 proteins support cRNA

vs vRNA synthesis. We have also established an RNA FISH assay which allows

visualisation of FluPol replication products. This allows single-cell, spatial information

to be collated, which in future could be used to improve our understanding of the

spatial elements of FluPol regulation.

Materials and methods

Cells and cell culture

Human-engineered Haploid cells (eHAP, Horizon Discovery), eHAP cells with both

huANP32A and huANP32B (eHAP dKO) ablated via CRISPR-Cas9, as previously

described (13), or eHAP cells with huANP32A, huANP32B and huANP32E ablated via

CRISPR-Cas9 (eHAP tKO) (gift from Ecco Staller and Ervin Fodor) were maintained

in Iscove's modified Dulbecco's medium (IMDM, Thermo Fisher) supplemented with

10% fetal bovine serum (FBS, Labtech), 1% penicillin-streptomycin (pen-strep, Gibco)

and 1% non-essential amino acids (NEAA, Gibco). Human embryonic kidney (293Ts,

ATCC) and Madin-Darby canine kidney (MDCK, ATCC) cells were maintained in

Dulbecco's modified Eagle medium (DMEM) supplemented with 10% FBS, 1% pen-

strep and 1% NEAA. When used for infection, 293T cells were cultured on poly-L-

lysine coated plates to aid adherence. All cells were maintained at 37°C and 5% CO_2.

Plasmids

pCAGGS expression plasmids encoding the polymerase subunits (PB2, PB1, PA) and

NP from PR8 were subcloned from pPolI plasmids. pCAGGS expression plasmids

encoding 50-92 PB2, PB1, PA and NP have previously been described (30). The

catalytic mutant PB1 D446Y has previously been described (31, 32). pCAGGS

expression plasmid encoding FLAG-tagged chicken ANP32 has been described

previously (15).

Viral infections

The full strain names of viruses used in this study are A/Puerto Rico/8/1934(H1N1)

(PR8) and A/turkey/England/50-92/1991(H5N1) (50-92). For experiments using 50-

92, all infections were performed with recombinant viruses containing the HA, NA and M segments from PR8, the PB1, PA, NP and NS segments from 50-92 and either the WT 50-92 PB2 segment containing a lysine at position 627 (5092E) or a modified PB2 with a glutamic acid at position 627 (5092K), as previously described (27). For infections, virus was diluted in serum-free media to the correct multiplicity of infection (described in figure legends). For comparative 50-92E/K experiments, viral inputs were normalised based on genome copy number. To synchronise infection, viral inoculation was performed at 4°C. In brief, cells were pre-incubated at 4°C for 15 minutes, before addition of viral inoculum and a further incubation at 4°C for 45 minutes. Viral inoculum was then replaced with pre-warmed full media and infected plates incubated at 37°C, 5% CO_2. At the appropriate time point, cells were processed for RT-qPCR analysis, imaging analysis or immunoblot as described below.

Replication/cRNP stabilisation assays

For replication/cRNP stabilisation assays, cells were transfected using Lipofectamine 3000 (Invitrogen) with pCAGGS expression plasmid mixtures encoding polymerase components in the ratios 2:2:1:4, PB2:PB1:PA:NP, where 1=20ng, 40ng or 80ng (24 well plate, 12 well plate or 6 well plate respectively). For experiments including chANP32A, pCAGGS expression plasmid encoding FLAG-tagged chANP32A was included in the transfection mix at a ratio of 4. For replication assays WT PB1 plasmid was pre-expressed, whilst for cRNP stabilisation assays catalytically dead polymerase (PB1 D446Y) was pre-expressed. 20 hours post transfection, cells were infected as described above, with the addition of actinomycin D (5 μg/mL), cycloheximide (100 μg/mL) or DMSO control as indicated. At the appropriate time point, cells were processed for RT-qPCR analysis, imaging analysis or immunoblot as described below.

Tagged RT-qPCR against vRNA, cRNA and mRNA

For RT-qPCR analysis, 293T or eHAP cells were cultured in 24 well plates, with each condition in triplicate. Following infection/transfection, cells were lysed using buffer RLT or RLT plus (Qiagen), frozen at -80, then total RNA was extracted using either the RNeasy RNA extraction kit (Qiagen) with 30 minute on column DNAseI digest, or the QIAsymphony RNA kit (Qiagen). Quantification for segment 4 and segment 6 vRNA, cRNA and mRNA was based on the tagged primer approach developed by Kawakami et al (24). For each sample, four reverse transcription reactions were set up using 200ng RNA/reaction, RevertAid H Minus Reverse Transcriptase (Thermo Scientific) as per the manufacturer's instructions, plus a tagged primer targeting either vRNA or cRNA, a tagged polydT (for viral mRNA) or an untagged polydT (for GAPDH internal control). For NA vRNA, cRNA and mRNA, primers used were GGCCGTCATGGTGGCGAATGAAACCATAAAAAGTTGGAGGAAG, GCTAGCTTCAGCTAGGCATCAGTAGAAACAAGGAGTTT and CCAGATCGTTCGAGTCGTTTTTTTTTTTTTTTTTTT respectively, tags underlined)) whilst primers used for HA vRNA, cRNA and mRNA were GGCCGTCATGGTGGCGAATGGAGAGTGCCCAAAATACGT, GCTAGCTTCAGCTAGGCATCAGTAGAAACAAGGGTGTT and CCAGATCGTTCGAGTCGTTTTTTTTTTTTTTTTTTT respectively, tags underlined). Tagged cDNA was then diluted 1 in 10 and quantified using real-time quantitative PCR using Fast SYBR green master mix (Thermo Scientific). Primer pairs used were: CCTTCCCCTTTTCGATCTTG/ GGCCGTCATGGTGGCGAAT (NA vRNA), CTTTTTGTGGCGTGAATAGTG/ GCTAGCTTCAGCTAGGCATC (NA cRNA), CTTTTTGTGGCGTGAATAGTG/ CCAGATCGTTCGAGTCGT (NA mRNA),

CATACCATCCATCTATCATTCC/ GGCCGTCATGGTGGCGAAT (HA vRNA),

GGGGGCAATCAGTTTCTG/ GCTAGCTTCAGCTAGGCATC (HA cRNA),

GATTCTGGCGATCTACTCAACTGTC/ CCAGATCGTTCGAGTCG (HA mRNA) and

AATCCCATCACCATCTTCCA/ TGGACTCCACGACGTACTCA (GAPDH). qPCR

analysis was carried out in duplicate or triplicate on a Viia 7 real-time PCR system

(Thermo Fisher). Fold changes in gene expression relative to either input (0hpi) or

mock infected controls (as indicated in figure legends) were calculated using the $2^{-\Delta\Delta CT}$

method with GAPDH expression as internal control.

RNAscope/immunofluorescence co-staining

For imaging analysis, cells were cultured on glass cover slips coated in poly-L-lysine

in 12 well plates. At the appropriate timepoint, infected cells were washed in phosphate

buffered saline (PBS, Gibco) and fixed in 4% paraformaldehyde for 30 minutes, prior

to further washes in PBS and dehydration in an ethanol gradient (50% EtOH, 5 mins,

70% EtOH, 5 mins, 100% EtOH, 5 mins, fresh 100% EtOH, 10 mins). Cover slips were

stored in 100% ethanol at -20°C until further processing. For

RNAscope/immunofluorescence co-staining, RNA was stained first using RNAscope

probes (ACDBio). Probes were designed to target PR8 NA vRNA (channel 1) and PR8

NA cRNA/mRNA (+RNA) (channel 2). Cover slips were rehydrated in an ethanol

gradient (70% EtOH, 2 mins, 50% EtOH, 2 mins, PBS, 10 mins), treated with protease

III diluted 1 in 15 in PBS and staining undertaken using the fluorescent multiplex kit

v1, following the manufacturer's instructions up until and including incubation in the

final fluorophore mixture (Fl-Amp4). At this point, cover slips were blocked in PBS with

2% bovine serum albumin (BSA) and 0.1% tween for 30 minutes, incubated in rabbit

α-PB2 (catalog no. GTX125926, Genetex) antibody for 1 hour at room temperature

(RT), followed by goat α-rabbit AF647 (Invitrogen) plus DAPI for 1 hour, RT. Images

were obtained using a Leica SP5 inverted confocal microscope and processing

undertaken using FIJI software (33, 34).

Immunoblot analysis

To confirm equivalent protein expression during replication/cRNP stabilisation assays,

cells transfected in parallel were lysed with homemade radioimmunoprecipitation

assay (RIPA) buffer (150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1%

SDS, 50 mM Tris, pH 7.4) supplemented with an EDTA-free protease inhibitor cocktail

tablet (Roche). Lysates were clarified, then mixed with 4x Laemmli sample buffer (Bio-

Rad) with 10% β-mercaptoethanol. Membranes were probed with rabbit α-vinculin

(EPR8185, Abcam) or mouse α-tubulin (ab7291, Abcam) and mouse α-NP (C43,

Abcam), followed by near-infrared secondary antibodies (IRDye 680RD goat anti-

rabbit (IgG) secondary antibody (Abcam) and IRDye 800CW goat anti-mouse (IgG)

secondary antibody (Abcam)). Western blots were visualised using an Odyssey

imaging system (Li-Cor Biosciences).

Statistical analysis

Statistics throughout this study were performed using one-way analysis of variance

(ANOVA) or student's T-test as described in figure legends.

We thank Ervin Fodor and Ecco Staller for the kind gift of the ANP32A/B/E triple

knockout eHAP cells. We thank Andreas Bruckbauer and Stephen Rothery at the

Facility for Imaging by Light Microscopy (FILM) at Imperial for support with microscopy.

This work was supported by Wellcome Trust grant 205100. In addition, O.C.S. was

supported by a Wellcome Trust studentship, A.B.R. was supported by a Medical

Research Council (MRC) studentship, and T.P.P. was supported by Biotechnology

and Biological Sciences Research Council (BBSRC) grant BB/R013071/1.

O.C.S., A.M.B., C.M.S and W.S.B. conceived and planned experiments; O.C.S. and

A.M.B. performed experiments and analysed data, O.C.S., T.P.P., C.M.S. and W.S.B.

provided supervision, O.C.S. and W.S.B. wrote the original manuscript, O.C.S.,

A.M.B., T.P.P., C.M.S. and W.S.B. reviewed and edited the manuscript.

The authors declare no competing interests.

References

425   References

426   1.   Long JS, Mistry B, Haslam SM, Barclay WS. 2019. Host and viral determinants
427   of influenza A virus species specificity. Nature Reviews Microbiology. Nature
428   Publishing Group https://doi.org/10.1038/s41579-018-0115-z.

429   2.   Subbarao EK, London W, Murphy BR. 1993. A single amino acid in the PB2
430   gene of influenza A virus is a determinant of host range. J Virol 67:1761–4.

431   3.   Pflug A, Lukarska M, Resa-Infante P, Reich S, Cusack S. 2017. Structural
432   insights into RNA synthesis by the influenza virus transcription-replication
433   machine. Virus Research. Elsevier B.V.
434   https://doi.org/10.1016/j.virusres.2017.01.013.

435   4.   Fodor E, Velthuis AJWT. 2020. Structure and function of the influenza virus
436   transcription and replication machinery. Cold Spring Harbor Perspectives in
437   Medicine 10:1–14.

438   5.   Fan H, Walker AP, Carrique L, Keown JR, Serna Martin I, Karia D, Sharps J,
439   Hengrung N, Pardon E, Steyaert J, Grimes JM, Fodor E. 2019. Structures of
440   influenza A virus RNA polymerase offer insight into viral genome replication.
441   Nature 573:287–290.

442   6.   York A, Hengrung N, Vreede FT, Huiskonen JT, Fodor E. 2013. Isolation and
443   characterization of the positive-sense replicative intermediate of a negative-
444   strand RNA virus. Proc Natl Acad Sci U S A 110.

445   7.   Deng T, Vreede FT, Brownlee GG. 2006. Different De Novo Initiation
446   Strategies Are Used by Influenza Virus RNA Polymerase on Its cRNA and
447   Viral RNA Promoters during Viral RNA Replication. Journal of Virology
448   80:2337–2348.

8.   Jorba N, Coloma R, Ortín J. 2009. Genetic trans-complementation establishes

a new model for influenza virus RNA transcription and replication. PLoS

Pathogens 5.

9.   Hengrung N, el Omari K, Serna Martin I, Vreede FT, Cusack S, Rambo RP,

Vonrhein C, Bricogne G, Stuart DI, Grimes JM, Fodor E. 2015. Crystal

structure of the RNA-dependent RNA polymerase from influenza C virus.

Nature 527:114–117.

10.   Thierry E, Guilligay D, Kosinski J, Bock T, Gaudon S, Round A, Pflug A,

Hengrung N, el Omari K, Baudin F, Hart DJ, Beck M, Cusack S. 2016.

Influenza Polymerase Can Adopt an Alternative Configuration Involving a

Radical Repacking of PB2 Domains. Molecular Cell 61:125–137.

11.   Carrique L, Fan H, Walker AP, Keown JR, Sharps J, Staller E, Barclay WS,

Fodor E, Grimes JM. 2020. Host ANP32A mediates the assembly of the

influenza virus replicase. Nature 587:638–643.

12.   Peacock TP, Sheppard CM, Staller E, Barclay WS. 2019. Host Determinants

of Influenza RNA Synthesis. Annual Reviews in Virology 6:215–233.

13.   Staller E, Sheppard CM, Neasham PJ, Mistry B, Peacock TP, Goldhill DH,

Long JS, Barclay WS. 2019. ANP32 Proteins Are Essential for Influenza Virus

Replication in Human Cells. Journal of Virology 93.

14.   Zhang H, Zhang Z, Wang Y, Wang M, Wang X, Zhang X, Ji S, Du C, Chen H,

Wang X. 2019. Fundamental Contribution and Host Range Determination of

ANP32A and ANP32B in Influenza A Virus Polymerase Activity. Journal of

Virology 93.

15.   Long JS, Giotis ES, Moncorgé O, Frise R, Mistry B, James J, Morisson M,

Iqbal M, Vignal A, Skinner MA, Barclay WS. 2016. Species difference in

ANP32A underlies influenza A virus polymerase host restriction. Nature

529:101–104.

16. Baker SF, Ledwith MP, Mehle A. 2018. Differential Splicing of ANP32A in Birds

Alters Its Ability to Stimulate RNA Synthesis by Restricted Influenza

Polymerase. Cell Reports 24:2581-2588.e4.

17. Domingues P, Hale BG. 2017. Functional Insights into ANP32A-Dependent

Influenza A Virus Polymerase Host Restriction. Cell Reports 20:2538–2546.

18. Mistry B, Long JS, Schreyer J, Staller E, Yusef Sanchez-David R, Barclay WS.

2020. Elucidating the Interactions between Influenza Virus Polymerase and

Host Factor ANP32A. Journal of Virology 94.

19. Sugiyama K, Kawaguchi A, Okuwaki M, Nagata K. 2015. PP32 and APRIL are

host cell-derived regulators of influenza virus RNA synthesis from cRNA. Elife

4:1–19.

20. Mänz B, Brunotte L, Reuther P, Schwemmle M. 2012. Adaptive mutations in

NEP compensate for defective H5N1 RNA replication in cultured human cells.

Nature Communications 3.

21. Nilsson BE, te Velthuis AJW, Fodor E. 2017. Role of the PB2 627 Domain in

Influenza A Virus Polymerase Function. Journal of Virology 91.

22. Nilsson-Payant BE, tenOever BR, te Velthuis AJW. 2021. The host factor

ANP32A is required for influenza A virus vRNA and cRNA synthesis. Journal

of Virology https://doi.org/10.1128/jvi.02092-21.

23. Reilly PT, Yu Y, Hamiche A, Wang L. 2014. Cracking the ANP32 whips:

Important functions, unequal requirement, and hints at disease implications.

BioEssays 36:1062–1071.

24.   Kawakami E, Watanabe T, Fujii K, Goto H, Watanabe S, Noda T, Kawaoka Y.

499   2011. Strand-specific real-time RT-PCR for distinguishing influenza vRNA,

500   cRNA, and mRNA. Journal of Virological Methods 173:1–6.

501   25.   Wang F, Flanagan J, Su N, Wang LC, Bui S, Nielson A, Wu X, Vo HT, Ma XJ,

502   Luo Y. 2012. RNAscope: A novel in situ RNA analysis platform for formalin-

503   fixed, paraffin-embedded tissues. Journal of Molecular Diagnostics 14:22–29.

504   26.   Bi Z, Ye H, Wang X, Fang A, Yu T, Yan L, Zhou J. 2019. Insights into species-

505   specific regulation of ANP32A on the mammalian-restricted influenza virus

506   polymerase activity. Emerging Microbes and Infections 8:1465–1478.

507   27.   Long JS, Howard WA, Núñez A, Moncorgé O, Lycett S, Banks J, Barclay WS.

508   2013. The Effect of the PB2 Mutation 627K on Highly Pathogenic H5N1 Avian

509   Influenza Virus Is Dependent on the Virus Lineage. Journal of Virology

510   87:9983–9996.

511   28.   Gabriel G, Herwig A, Klenk HD. 2008. Interaction of polymerase subunit PB2

512   and NP with importin α1 is a determinant of host range of influenza A virus.

513   PLoS Pathogens 4.

514   29.   Wang F, Sheppard CM, Mistry B, Staller E, Barclay WS, Grimes JM, Fodor E,

515   Fan H. 2022. The C-terminal LCAR of host ANP32 proteins interacts with the

516   influenza A virus nucleoprotein to promote the replication of the viral RNA

517   genome. Nucleic Acids Research https://doi.org/10.1093/nar/gkac410.

518   30.   Moncorgé O, Mura M, Barclay WS. 2010. Evidence for Avian and Human Host

519   Cell Factors That Affect the Activity of Influenza Virus Polymerase. Journal of

520   Virology 84:9978–9986.

31.   Cauldwell A v., Moncorge O, Barclay WS. 2013. Unstable Polymerase-

Nucleoprotein Interaction Is Not Responsible for Avian Influenza Virus

Polymerase Restriction in Human Cells. Journal of Virology 87:1278–1284.

32.   Biswas SK, Nayak DP. 1994. Mutational Analysis of the Conserved Motifs of

Influenza A Virus Polymerase Basic Protein 1. Journal of Virology 68:1819–

1826.

33.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,

Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ,

Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source

platform for biological-image analysis. Nat Methods 9:676–82.

34.   Schneider C a, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25

years of image analysis. Nature Methods 9:671–675.

Figure 1. cRNA synthesis is inhibited in human cells lacking ANP32A/B. (A-C)

Segment 4 (A) vRNA (B) cRNA or (C) mRNA accumulation over time in eHAP WT vs

dKO cells following infection with PR8 (MOI=3). Fold change was calculated over input

(0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed at

2hpi using an unpaired t-test following log transformation. (D-F) Segment 6 (D) vRNA

(E) cRNA or (F) mRNA accumulation over time in eHAP WT vs dKO cells following

infection with PR8 (MOI=3). Fold change was calculated over mock infected cells. n=3

biological replicates, plotted as mean ± s.d.. Significance was assessed using multiple

unpaired t-tests following log transformation, corrected for multiple comparisons using

false discovery rate. hpi=hours post infection. ns=not significant; **=$p<0.01$;

***=$p<0.001$; ****=$p<0.0001$.

Figure 2. Human ANP32 proteins play a direct role in primary cRNA synthesis. (A)

Schematic illustrating influenza polymerase activity under the conditions of a cRNP

stabilisation assay with ActD. (B-E) cRNP stabilisation assay with ActD in eHAP WT

vs dKO cells, following infection with PR8 (MOI=3), 3hpi. Segment 6 (B) vRNA (C)

cRNA or (D) mRNA accumulation. Dotted line indicates background RNA present in

control samples transfected with a plasmid mix lacking PB1. Fold change was

calculated over mock infected cells. n=3 biological replicates, plotted as mean ± s.d..

Significance was assessed using an unpaired t-test following log transformation. (E)

Accumulation of segment 6 vRNA and +RNA in eHAP WT and dKO cells, analysed

using RNAscope. Magenta arrowheads highlight a subset of NA vRNA-stained puncta.

Images are representative maximum intensity projections. (F) Schematic illustrating

influenza polymerase activity under the conditions of a cRNP stabilisation assay with

CHX. (G-I) cRNP stabilisation assay with CHX in eHAP WT, dKO and tKO cells.

Segment 4 (G) vRNA (H) cRNA or (I) mRNA accumulation following infection with PR8 (MOI=3). Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using multiple unpaired t-tests following log transformation, corrected for multiple comparisons using false discovery rate. hpi=hours post infection. ns=not significant; **=$p<0.01$; ***=$p<0.001$; ****=$p<0.0001$.

Figure 3. Avian polymerase is not restricted in cRNA synthesis in mammalian cells. (A) Schematic outlining assay set-up and expected polymerase combinations during replication. As indicated on the left, in the cRNP stabilisation assay, only the first layer of activity (primary cRNA synthesis) will occur. Both layers of activity can occur in replication assays. Unknown polymerase activity is indicated by a dashed black arrow. (B,C) Simultaneous cRNP stabilisation and replication assays with ActD, 6 hpi. Segment 6 (B) vRNA and (C) cRNA accumulation following infection with either 5092E or 5092K as indicated (MOI=0.1). Dotted line indicates levels of vRNA/cRNA present in a control lacking PB2 in the transfection mix. Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using one way ANOVA with Dunnett's multiple comparison test, following log transformation. (D,E) Replication assays with ActD and chANP32A, 6 hpi. Segment 6 (D) vRNA and (E) cRNA accumulation following infection with either 5092E or 5092K as indicated (MOI=0.1). Dotted line indicates levels of vRNA/cRNA present in a control lacking PB2 in the transfection mix. Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using one way ANOVA with Dunnett's multiple comparison test, following log transformation. Pre-expressed polymerase mixes: Stabilisation K=PB2 627K/PB1 D446Y/PA/NP;

Stabilisation E=PB2 627E/PB1 D446Y/PA/NP; Replication K=PB2 627K/PB1/PA/NP;

Replication E=PB2 627E/PB1/PA/NP. hpi=hours post infection. ns=not significant; *=

$p < 0.05$; ****=$p < 0.0001$.

Figure S1. Supporting data for figure 2. (A) Matched western blot for Fig. 2B,C,D. (B-

F) Validation of replication and stabilisation assays with ActD analysed by RNA FISH.

Assays were performed in HEK 293T cells, following infection with PR8 (MOI=3). (B)

Accumulation of NA vRNA and +RNA, 3hpi, analysed by RNAscope with indirect

immunofluorescence against PB2. Pre-transfected PR8 polymerase mixes are

indicated on the left-hand side: FluPol WT=PB2/PB1/PA/NP; FluPol D446Y=PB2/PB1

D446Y/PA/NP; -PB1=PB2/PA/NP. Images are representative maximum intensity

projections. (C) Matched western blot confirming equal transfection of different

polymerase mixtures. (D,E,F) Matched RT-qPCR analysis of segment 6 (D) vRNA (E)

cRNA and (F) mRNA accumulation, 6hpi. Pre-transfected polymerase mixes are

indicated on the x-axis. Dotted line indicates input levels of RNA (0hpi). Fold change

was calculated over mock infected cells. n=3 biological replicates, plotted as mean ±

s.d.. Significance compared to –PB1 was assessed using one-way ANOVA with

Dunnett's multiple comparison test, following log transformation. (G) Matched western

blot for Fig. 2G,H,I. Untransf= Untransfected; hpi=hours post infection. ns=not

significant; ****=$p < 0.0001$.

Figure S2. Human ANP32 proteins play a direct role in primary cRNA synthesis of

H5N1 5092. (A-C) Segment 6 (A) vRNA (B) cRNA or (C) mRNA accumulation over

time in eHAP WT vs tKO cells following infection with 5092K (MOI=3). Fold change

was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d..

Significance at 2hpi was assessed using an unpaired t-test following log

transformation. (D,E,F) cRNP stabilisation assay with CHX in eHAP WT and tKO cells.

Segment 4 (D) vRNA (E) cRNA or (F) mRNA accumulation following infection with

5092K (MOI=3). Fold change was calculated over input (0hpi). Dotted line indicates

background levels of vRNA/cRNA present in WT cells transfected with a control

plasmid mix minus PB1, 5hpi. n=3 biological replicates, plotted as mean ± s.d..

Significance was assessed using multiple unpaired t-tests following log

transformation, corrected for multiple comparisons using false discovery rate.

hpi=hours post infection. ns=not significant; *= p<0.05; **=p<0.01; ***=p<0.001;

****=p<0.0001.

Figure 1. cRNA synthesis is inhibited in human cells lacking ANP32A/B. (A-C) Segment 4 (A) vRNA (B) cRNA or (C) mRNA accumulation over time in eHAP WT vs dKO cells following infection with PR8 (MOI=3). Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed at 2hpi using an unpaired t-test following log transformation. (D-F) Segment 6 (D) vRNA (E) cRNA or (F) mRNA accumulation over time in eHAP WT vs dKO cells following infection with PR8 (MOI=3). Fold change was calculated over mock infected cells. n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using multiple unpaired t-tests following log transformation, corrected for multiple comparisons using false discovery rate. hpi=hours post infection. ns=not significant; **=$p<0.01$; ***=$p<0.001$; ****=$p<0.0001$.

Figure 2. Human ANP32 proteins play a direct role in primary cRNA synthesis. (A) Schematic illustrating influenza polymerase activity under the conditions of a cRNP stabilisation assay with ActD. (B-E) cRNP stabilisation assay with ActD in eHAP WT vs dKO cells, following infection with PR8 (MOI=3), 3hpi. Segment 6 (B) vRNA (C) cRNA or (D) mRNA accumulation. Dotted line indicates background RNA present in control samples transfected with a plasmid mix lacking PB1. Fold change was calculated over mock infected cells. n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using an unpaired t-test following log transformation. (E) Accumulation of segment 6 vRNA and +RNA in eHAP WT and dKO cells, analysed using RNAscope. Magenta arrowheads highlight a subset of NA vRNA-stained puncta. Images are representative maximum intensity projections. (F) Schematic illustrating influenza polymerase activity under the conditions of a cRNP stabilisation assay with CHX. (G-I) cRNP stabilisation assay with CHX in eHAP WT, dKO and tKO cells. Segment 4 (G) vRNA (H) cRNA or (I) mRNA accumulation following infection with PR8 (MOI=3). Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using multiple unpaired t-tests following log transformation, corrected for multiple comparisons using false discovery rate. hpi=hours post infection. ns=not significant; **=p<0.01; ***=p<0.001; ****=p<0.0001.

Figure 3. Avian polymerase is not restricted in cRNA synthesis in mammalian cells. (A) Schematic outlining assay set-up and expected polymerase combinations during replication. As indicated on the left, in the cRNP stabilisation assay, only the first layer of activity (primary cRNA synthesis) will occur. Both layers of activity can occur in replication assays. Unknown polymerase activity is indicated by a dashed black arrow. (B,C) Simultaneous cRNP stabilisation and replication assays with ActD, 6 hpi. Segment 6 (B) vRNA and (C) cRNA accumulation following infection with either 5092E or 5092K as indicated (MOI=0.1). Dotted line indicates levels of vRNA/cRNA present in a control lacking PB2 in the transfection mix. Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using one way ANOVA with Dunnett's multiple comparison test, following log transformation. (D,E) Replication assays with ActD and chANP32A, 6 hpi. Segment 6 (D) vRNA and (E) cRNA accumulation following infection with either 5092E or 5092K as indicated (MOI=0.1). Dotted line indicates levels of vRNA/cRNA present in a control lacking PB2 in the transfection mix. Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using one way ANOVA with Dunnett's multiple comparison test, following log transformation. Pre-expressed polymerase mixes: Stabilisation K=PB2 627K/PB1 D446Y/PA/NP; Stabilisation E=PB2 627E/PB1 D446Y/PA/NP; Replication K=PB2 627K/PB1/PA/NP; Replication E=PB2 627E/PB1/PA/NP. hpi=hours post infection. ns=not significant; *= p<0.05; ****=p<0.0001.

Figure S1. Supporting data for figure 2. (A) Matched western blot for Fig. 2B,C,D. (B-F) Validation of replication and stabilisation assays with ActD analysed by RNA FISH. Assays were performed in HEK 293T cells, following infection with PR8 (MOI=3). (B) Accumulation of NA vRNA and +RNA, 3hpi, analysed by RNAscope with indirect immunofluorescence against PB2. Pre-transfected PR8 polymerase mixes are indicated on the left-hand side: FluPol WT=PB2/PB1/PA/NP; FluPol D446Y=PB2/PB1 D446Y/PA/NP; -PB1=PB2/PA/NP. Images are representative maximum intensity projections. (C) Matched western blot confirming equal transfection of different polymerase mixtures. (D,E,F) Matched RT-qPCR analysis of segment 6 (D) vRNA (E) cRNA and (F) mRNA accumulation, 6hpi. Pre-transfected polymerase mixes are indicated on the x-axis. Dotted line indicates input levels of RNA (0hpi). Fold change was calculated over mock infected cells. n=3 biological replicates, plotted as mean ± s.d.. Significance compared to –PB1 was assessed using one-way ANOVA with Dunnett's multiple comparison test, following log transformation. (G) Matched western blot for Fig. 2G,H,I. Untransf= Untransfected; hpi=hours post infection. ns=not significant; ****=p<0.0001.

Figure S2. Human ANP32 proteins play a direct role in primary cRNA synthesis of H5N1 5092. (A-C) Segment 6 (A) vRNA (B) cRNA or (C) mRNA accumulation over time in eHAP WT vs tKO cells following infection with 5092K (MOI=3). Fold change was calculated over input (0hpi). n=3 biological replicates, plotted as mean ± s.d.. Significance at 2hpi was assessed using an unpaired t-test following log transformation. (D,E,F) cRNP stabilisation assay with CHX in eHAP WT and tKO cells. Segment 4 (D) vRNA (E) cRNA or (F) mRNA accumulation following infection with 5092K (MOI=3). Fold change was calculated over input (0hpi). Dotted line indicates background levels of vRNA/cRNA present in WT cells transfected with a control plasmid mix minus PB1, 5hpi. n=3 biological replicates, plotted as mean ± s.d.. Significance was assessed using multiple unpaired t-tests following log transformation, corrected for multiple comparisons using false discovery rate. hpi=hours post infection. ns=not significant; *= p<0.05; **=p<0.01; ***=p<0.001; ****=p<0.0001.