Presynaptic Rac1 controls synaptic strength through the regulation of synaptic vesicle priming

(Abbreviated title): Presynaptic Rac1 regulates synaptic strength

Christian Keine[1,2,3], Mohammed Al-Yaari[1], Tamara Radulovic[1,2,3], Connon I. Thomas[6], Paula Valino Ramos[1], Debbie Guerrero-Given[6], Mrinalini Ranjan[4,5], Holger Taschenberger[4] Naomi Kamasawa[6], and Samuel M. Young, Jr.[1,7^].

[1]Department of Anatomy and Cell Biology, University of Iowa, Iowa City, IA, USA
[2]Department of Human Medicine, Carl-von-Ossietzky University Oldenburg, Oldenburg, Germany
[3]Research Center Neurosensory Science, Oldenburg, Germany
[4]Department of Molecular Neurobiology, Max Planck Institute for Multidisciplinary Sciences, Göttingen, Germany
[5]Göttingen Graduate School for Neurosciences, Biophysics, and Molecular Biosciences, Göttingen, Germany
[6]Electron Microscopy Core Facility, Max Planck Florida Institute, Jupiter, FL, USA
[7]Department of Otolaryngology, Iowa Neuroscience Institute, University of Iowa, Iowa City, IA, USA.

^Correspondence
Samuel M. Young, Jr. PhD
Department of Anatomy and Cell Biology
University of Iowa
PBDB 5322
169 Newton Road
Iowa City, IA 52242
Email: samuel-m-young@uiowa.edu

ORCIDs:
Christian Keine: 0000-0002-8953-2593
Mohammed Al-Yaari   0000-0003-3196-266X
Tamara Radulovic: 0000-0002-2825-9773
Connon I. Thomas: 0000-0003-0995-9667
Mrinalini Ranjan: 0000-0003-4310-3811
Holger Taschenberger: 0000-0003-3186-3231
Naomi Kamasawa: 0000-0002-8926-5309
Samuel M. Young, Jr.:   0000-0002-7589-7612

Abstract

Synapses contain a limited number of synaptic vesicles (SVs) that are released in response to action potentials (APs). Therefore, sustaining synaptic transmission over a wide range of AP firing rates and timescales depends on SV release and replenishment. Although actin dynamics impact synaptic transmission, how presynaptic regulators of actin signaling cascades control SV release and replenishment remains unresolved. Rac1, a Rho GTPase, regulates actin signaling cascades that control synaptogenesis, neuronal development, and postsynaptic function. However, the presynaptic role of Rac1 in regulating synaptic transmission is unclear. To unravel Rac1's roles in controlling transmitter release, we performed selective presynaptic ablation of Rac1 at the mature mouse calyx of Held synapse. Loss of Rac1 increased synaptic strength, accelerated EPSC recovery after conditioning stimulus trains, and augmented spontaneous SV release with no change in presynaptic morphology or AZ ultrastructure. Analyses with constrained short-term plasticity models revealed faster SV priming kinetics and, depending on model assumptions, elevated SV release probability or higher abundance of tightly docked fusion-competent SVs in Rac1-deficient synapses. We conclude that presynaptic Rac1 is a key regulator of synaptic transmission and plasticity mainly by regulating the dynamics of SV priming and potentially SV release probability.

Introduction

Information encoding in the nervous system requires synaptic transmission to drive and sustain action potentials (APs) over rapidly changing and highly variable AP firing rates (Reinagel and Laughlin, 2001; Theunissen and Elie, 2014; Brette, 2015; Azarfar et al., 2018). However, the number of synaptic vesicles (SVs) available for fusion in response to an AP, the readily releasable pool (RRP), is limited (Alabi and Tsien, 2012). Therefore, tight regulation of SV release and RRP replenishment is required for synaptic reliability and temporally precise information encoding (Neher, 2010; Hallermann and Silver, 2013). Priming is the process that generates fusion-competent SVs. It is a critical step in the SV cycle that primarily regulates RRP size and SV pool replenishment. Priming also controls SV release probability (P_r) by determining SV fusogenicity ('molecular priming') and regulating the spatial coupling between docked SV and presynaptic Ca^{2+} entry ('positional priming') (Klug et al., 2012; Schneggenburger and Rosenmund, 2015; Neher and Brose, 2018). In some synapses, the SV priming kinetics are highly dependent on presynaptic cytosolic Ca^{2+} levels, which are activity-dependent. Importantly, human mutations in molecules that regulate priming are associated with neurological disorders (Waites et al., 2011; Wondolowski and Dickman, 2013; Torres et al., 2017; Bonnycastle et al., 2021). Therefore, elucidating the molecular mechanisms that regulate SV priming is critical to understanding the diversity of neuronal information encoding in health and disease.

Actin is a central component of both the presynaptic and postsynaptic compartments, with diverse roles in regulating synaptic function and neuronal circuit development. Manipulation of actin dynamics or interference with presynaptic AZ proteins implicated in regulating actin dynamics affects transmitter release and SV replenishment, as well as P_r (Morales et al., 2000; Sakaba and Neher, 2003; Cingolani and Goda, 2008; Sun and Bamji, 2011; Waites et al., 2011; Lee et al., 2012; Lee et al., 2013; Montesinos et al., 2015; Rust and Maritzen, 2015). However, due to disparate results, the role of presynaptic actin signaling cascades in regulating transmitter release and SV pool replenishment is controversial. Finally, in contrast to the postsynaptic

compartment, our understanding of how presynaptic regulators of actin signaling cascades

control synaptic transmission and short-term plasticity (STP) are in the early stages.

Rac1, a Rho GTPase, is a critical regulator of actin signaling cascades (Bosco et al.,

2009; Yasuda, 2017), and human mutations in Rac1 are associated with neurological disorders

(Bai et al., 2015; Reijnders et al., 2017; Zamboni et al., 2018). Rac1 is involved in multiple

processes which control synaptogenesis, axon guidance, neuronal development, and

postsynaptic function to regulate neuronal circuit function (Bai et al., 2015). Although Rac1 is

expressed in both the pre- and postsynaptic compartment (Threadgill et al., 1997; Doussau et

al., 2000; Kumanogoh et al., 2001; O'Neil et al., 2021), its presynaptic role in regulating synaptic

transmission is not well understood. Recent work using cultured hippocampal neurons proposed

that presynaptic Rac1 is a negative regulator of SV pool replenishment (O'Neil et al., 2021),

though at which steps in the SV cycle it exerts its regulatory role is unknown. Finally, how Rac1

regulates the temporal dynamics of transmitter release and pool replenishment in a native

neuronal circuit remains elusive.

To unravel the roles of presynaptic Rac1 in regulating transmitter release, we utilized the

calyx of Held, a glutamatergic axosomatic presynaptic terminal in the auditory brainstem, which

is the sole input to drive AP firing in the principal cells of the medial nucleus of the trapezoid

body (MNTB) (Borst and Soria van Hoeve, 2012; Joris and Trussell, 2018). In the calyx of Held,

RRP dynamics are tightly regulated to support a nearly failsafe synaptic transmission with high

temporal precision, but the molecular machinery of transmitter release and SV pool

replenishment and their regulation are similar to conventional presynaptic terminals in the central

nervous system (Iwasaki and Takahashi, 1998; Iwasaki et al., 2000; Iwasaki and Takahashi,

2001; Satzler et al., 2002; Taschenberger et al., 2002; Rollenhagen and Lubke, 2006; Neher and

Sakaba, 2008; Alabi and Tsien, 2012; Eggermann et al., 2012; Hallermann and Silver, 2013;

Schneggenburger and Rosenmund, 2015). In addition, molecular manipulations specific to only

the calyceal terminal can be carried out at different developmental stages (Wimmer et al., 2006;

Young and Neher, 2009; Chen et al., 2013; Lubbert et al., 2019). To elucidate the roles of

presynaptic Rac1 in regulating transmitter release while avoiding interference with its role in

synaptogenesis, axon guidance, and neuronal development, we selectively deleted Rac1 two

days after hearing onset in postnatal day (P) 14 mouse calyx of Held synapses. At this time

point, the synapse is functionally mature, and neuronal properties of brainstem circuits are

considered largely "adult-like". Subsequently, we determined how the loss of Rac1 impacted

calyx of Held/ MNTB principal cell transmission at the adult stage (P28 onwards) (Sonntag et al.,

2009; Crins et al., 2011; Sonntag et al., 2011). Presynaptic Rac1 deletion did neither affect the

calyx of Held morphology nor active zone (AZ) ultrastructure but led to increased synaptic

strength, faster SV pool replenishment, and augmented spontaneous SV release. Additionally,

we found that loss of Rac1 delayed the EPSC onset and potentiated asynchronous release

during high-frequency trains.

Analysis of the experimental data with constrained STP models confirmed faster SV

priming kinetics in Rac1-deficient synapses. Methods of quantal analysis, which assume a single

and homogenous pool of readily releasable SVs (Neher, 2015; Schneggenburger and

Rosenmund, 2015), reported an increased P_r and a tendency towards an increased RRP after

Rac1 loss. Both experimental findings were corroborated in numerical simulation using a single

pool STP model (Weis et al., 1999). In contrast, simulations based on a sequential two-step

priming scheme which assumes two distinct states of docked/primed SVs, a loosely-docked

(LS), immature priming state which is not fusion competent, and a tightly docked (TS), mature

priming state which is fusion competent (Neher and Taschenberger, 2021; Lin et al., 2022),

required only changes in SV priming kinetics but no change in P_r or the number of release sites

to reproduce experimental data. Simulations based on the sequential two-step SV priming and

fusion scheme fully accounted for the increased synaptic strength in $Rac1^{-/-}$ synapses by a

larger abundance of tightly docked SVs. Therefore, we propose that presynaptic Rac1 is a key

molecule that controls synaptic strength and STP primarily by regulating the SV priming

dynamics and potentially P_r. Finally, we conclude that presynaptic Rac1 is a critical regulator of

synaptic transmission and plasticity.

Results

Presynaptic deletion of Rac1 after hearing onset does not impact calyx of Held morphology or AZ ultrastructure

Presynaptic terminals contain Rac1 (Doussau et al., 2000; O'Neil et al., 2021) which regulates synaptogenesis, axon guidance, and neuronal development (Xu et al., 2019; Zhang et al., 2021). In this study, we aimed to elucidate Rac1's presynaptic function in controlling synaptic transmission and plasticity independent of its role in regulating synapse development and maturation at the calyx of Held synapse. To do so, we injected HdAd co-expressing Cre recombinase and EGFP into the cochlear nucleus (CN) of P14 $Rac1^{flox/flox}$ mice when the calyx of Held synapse is considered "adult-like" and commenced synapse analysis at P28 onwards (Fig. 1A) (Sonntag et al., 2009; Crins et al., 2011; Sonntag et al., 2011). Since Rac1 controls synapse development and morphology, it was essential to determine whether the loss of Rac1 after hearing onset altered the calyx of Held morphology or AZ ultrastructure. We analyzed calyx morphology from 3D reconstructions of confocal z-stack images acquired from $Rac1^{+/+}$ and $Rac1^{-/-}$ calyces at P28 and found no difference in calyx surface area or volume (Fig. 1B). To determine if the loss of Rac1 impacted AZ ultrastructure, we performed ultrathin-section TEM and analyzed AZ length, SV distribution, and the number of docked SVs and found no difference between $Rac1^{+/+}$ and $Rac1^{-/-}$ (Fig. 1C). Therefore, we conclude that after hearing onset, Rac1 does not regulate calyx of Held morphology and AZ ultrastructure.

Loss of Rac1 increases synaptic strength and relative synaptic depression.

Perturbations of the presynaptic actin cytoskeleton impact synaptic transmission and plasticity in multiple model systems and synapses (Cole et al., 2000; Sakaba and Neher, 2003; Bleckert et al., 2012; Lee et al., 2012; Rust and Maritzen, 2015; Miki et al., 2016; Gentile et al., 2022; Wu and Chan, 2022). Since Rac1 is an actin cytoskeleton regulator, we examined how the loss of Rac1 impacted AP-evoked synaptic transmission and STP. Afferent fibers of calyx synapses were electrically stimulated with 50 APs at two stimulation frequencies (50 and 500 Hz),

representing typical *in-vivo* firing rates at the calyx. AMPAR-mediated ESPCs were recorded in

MNTB principal cells innervated by transduced ($Rac1^{-/-}$) and non-transduced ($Rac1^{+/+}$) calyces

in 1.2 mM external Ca^{2+} and at 36-37°C to closely mimic *in-vivo* conditions (Fig. 2A, B). By

analyzing the initial response ($EPSC_1$) of the EPSC trains, we found a robust increase in

synaptic strength upon genetic Rac1 deletion ($Rac^{+/+}$ = 1.3 ± 0.4 nA vs. $Rac1^{-/-}$ = 3 ± 1.1 nA,

p<0.001, n = 15/15, Fig. 2C, D1) with no changes in EPSC waveform. Plotting EPSC amplitudes

vs. stimulus number revealed substantial differences in STP between $Rac1^{+/+}$ and $Rac1^{-/-}$

synapses. At 50 Hz stimulation, both $Rac1^{+/+}$ and $Rac1^{-/-}$ showed short-term depression, which

was more pronounced in $Rac1^{-/-}$ showing increased steady-state depression ($EPSC_{ss}$ / $EPSC_1$)

(Fig. 2A). Despite the stronger relative short-term depression, absolute steady-state EPSC

amplitudes were almost two-fold larger in $Rac1^{-/-}$ ($Rac1^{+/+}$ = 0.47 ± 0.14 nA vs.

$Rac1^{-/-}$ = 0.84 ± 0.22 nA, p < 0.001, n = 15/15, Fig. 2D1). At 500 Hz stimulation, $Rac1^{+/+}$ showed

robust short-term facilitation, which was absent in $Rac1^{-/-}$ (PPR: $Rac1^{+/+}$ = 1.2 ± 0.1 vs.

$Rac1^{-/-}$ = 1 ± 0.1, p < 0.001, n = 15/15, Fig. 2B). Similar to 50 Hz stimulation, $Rac1^{-/-}$ showed

more pronounced relative steady-state depression at 500 Hz stimulation. Notably, absolute

steady-state EPSC amplitudes at 500 Hz stimulation frequency were similar for $Rac1^{+/+}$ and

$Rac1^{-/-}$ (Fig. 2D1).

Since the loss of Rac1 increased synaptic strength and altered STP, we aimed to identify the

underlying mechanisms and evaluated RRP size and SV release probability (P_r) using

established quantal analysis methods (Elmqvist and Quastel, 1965; Neher, 2015; Thanawala

and Regehr, 2016). We estimated RRP size using 500 Hz stimulus trains which effectively

depleted the RRP in both $Rac1^{+/+}$ and $Rac1^{-/-}$ synapses by applying three conventional

paradigms based on the common assumption of quantal release originating from a single and

functionally homogenous pool of readily-releasable SVs (Neher, 2015; Schneggenburger and

Rosenmund, 2015): EQ, NpRf, and SMN with correction (Elmqvist and Quastel, 1965; Neher,

2015; Thanawala and Regehr, 2016). All three methods reported a moderate increase in RRP

size in $Rac1^{-/-}$ calyces (Fig. 2D2), however, this was statistically significant only for SMN

analysis. The initial P_r of resting synapses was estimated by dividing the $EPSC_1$ amplitude by

the estimated RRP sizes. All three analysis methods revealed an approximately two-fold

increase in P_r (SMN: $Rac1^{+/+} = 0.09 \pm 0.02$ vs. $Rac1^{-/-} = 0.15 \pm 0.03$, $p < 0.001$; NpRf:

$Rac1^{+/+} = 0.09 \pm 0.02$ vs. $Rac1^{-/-} = 0.17 \pm 0.03$, $p < 0.001$; EQ: $Rac1^{+/+} = 0.09 \pm 0.02$ vs.

$Rac1^{-/-} = 0.17 \pm 0.04$, $p < 0.001$, n = 15/15, Fig. 2D3). Therefore, based on the assumption of a

single and functionally homogenous RRP, this analysis indicates that presynaptic Rac1 deletion

increases synaptic strength and short-term depression primarily by elevating initial P_r with little

increase in RRP size. This suggests that Rac1 controls synaptic strength as a negative regulator

of P_r.

Loss of Rac1 increases mEPSC frequency but not amplitude

The RRP size and the P_r of the fusion-competent SVs is determined by the SV priming process,

which involves the assembly of the molecular fusion apparatus, defined as 'molecular priming'.

In addition, P_r also depends on the spatial coupling between docked SVs and Ca^{2+} entry sites

which may be adjusted by a distinct 'positional priming' step. Thus, 'molecular priming'

encompasses the steps that render SVs fusion competent and regulate their intrinsic

fusogenicity (Basu et al., 2007; Xue et al., 2010; Schneggenburger and Rosenmund, 2015;

Schotten et al., 2015), while positional priming consists of the steps that place molecularly

primed SVs close to voltage-gated calcium channels (VGCCs) (Neher, 2010). The spatial

coupling between SV and VGCCs critically determines the speed and efficacy of AP-evoked

release (Eggermann et al., 2012; Stanley, 2016). Spontaneous SV release is not or only little

dependent on VGCCs (Schneggenburger and Rosenmund, 2015; Kavalali, 2020), and the

frequency of miniature EPSC (mEPSCs) can, thus, be interpreted as a readout of intrinsic SV

fusogenicity at basal Ca^{2+} with increased SV fusogenicity causing higher mEPSC frequencies

(Basu et al., 2007; Schotten et al., 2015; Dong et al., 2018).Therefore, to determine if an

increased intrinsic SV fusogenicity caused the increase in P_r, we recorded mEPSCs from

Rac1$^{+/+}$ and *Rac1$^{-/-}$* calyx synapses (Fig. 3) and found a four-fold increase in mEPSC frequency

(*Rac1$^{+/+}$* = 3.4 ± 4.4 Hz vs. *Rac1$^{-/-}$* = 14.3 ± 6.5 Hz, p < 0.001, n = 15/11) with no change in

mEPSC amplitude or waveform. To rule out that the increased mEPSC frequencies were due to

changes in presynaptic Ca^{2+} currents, we recorded mEPSCs in the presence of 200 µM Cd^{2+}, a

non-specific VGCC blocker. Since Cd^{2+} did not affect mEPSC frequencies we conclude that

Rac1 loss increases intrinsic SV fusogenicity at basal Ca^{2+}.

Loss of Rac1 increases EPSC onset delays and decreases synchronicity of AP-evoked

release

Although we found an increase in SV fusogenicity, this does not rule out an additional role for

Rac1 in regulating spatial coupling distances between molecularly primed SVs and VGCCs. In

the mature calyx of Held, AP-evoked SV release is governed by local Ca^{2+} nanodomains,

ensuring a fast onset and highly synchronous AP-evoked EPSCs to faithfully encode auditory

information (Fedchyshyn and Wang, 2005). In addition to the gating kinetics of presynaptic

VGCCs and postsynaptic AMPARs, the time between presynaptic AP and EPSC onset (EPSC

onset delay) is determined by the coupling distance between SVs and VGCCs. The coupling

distance defines the time for Ca^{2+} to diffuse and bind to the Ca^{2+} sensor and initiate SV fusion

(Fedchyshyn and Wang, 2007; Nakamura et al., 2015). Thus, EPSC onset delays can serve as

a readout of changes in spatial coupling distances, as increased onset delays are consistent

with SVs being more loosely coupled to VGCCs and vice versa. Therefore, we measured EPSC

onset delays during 50 Hz and 500 Hz stimulation (Fig. 4A, B) and found them to become

progressively larger during stimulation for *Rac1$^{+/+}$* and *Rac1$^{-/-}$* calyces. At 50 Hz, the increase in

EPSC onset delays during stimulus trains was comparable between *Rac1$^{+/+}$* and *Rac1$^{-/-}$*,

amounting to about 60 µs between the first and the last ten EPSCs. At 500 Hz stimulation,

however, EPSC onset delays increased more rapidly in *Rac1$^{-/-}$*, with steady-state EPSC onset

delays being significantly larger for *Rac1$^{-/-}$* compared to *Rac1$^{+/+}$* (*Rac1$^{+/+}$* = 94± 32 µs vs.

Rac1$^{-/-}$ = 131 ± 29 µs, p = 0.004, n = 15/15).

In addition to modulating EPSC onset delays, coupling distances between SV and

VGCCs affect the time course of synchronous release and the relative contribution of

synchronous versus asynchronous release during AP trains (Wadel et al., 2007; Chen et al.,

2015; Stanley, 2016; Yang et al., 2021). This is because synchronous release is dominated by

tightly coupled SVs which rapidly fuse in response to high local $[Ca^{2+}]$, while asynchronous

release likely represents a stronger contribution of more loosely coupled SVs (Sakaba, 2006;

Schneggenburger and Rosenmund, 2015). An approximate measure for changes in the time

course of AP-evoked release can be obtained by analyzing the EPSC charge over EPSC

amplitude ratio ('effective EPSC duration') representing the width of a square current pulse with

same amplitude as the EPSC peak and same integral as the EPSC charge. Therefore, we

calculated the effective EPSC duration for both 50 Hz and 500 Hz stimulation (Fig. 4C) and

found its value for $EPSC_1$ comparable between $Rac1^{+/+}$ and $Rac1^{-/-}$. At steady-state during

50 Hz stimulation, however, the effective EPSC duration was slightly longer in $Rac1^{-/-}$

($Rac1^{+/+}$ = 0.32 ± 0.03 ms vs. $Rac1^{-/-}$ = 0.35 ± 0.03 ms, p = 0.006). At steady-state during

500 Hz stimulation, the effective EPSC duration in $Rac1^{-/-}$ calyces was prolonged further and

increased   by   ~25%   compared   to $Rac1^{+/+}$ ($Rac1^{+/+}$ = 0.43 ± 0.04 ms   vs.

$Rac1^{-/-}$ = 0.55 ± 0.12 ms, p < 0.001). These findings are consistent with transmitter release

being less synchronous in $Rac1^{-/-}$ synapses, especially during sustained activity at high AP

firing rates.

In summary, we found that $Rac1^{-/-}$ synapses had longer EPSC onset delays and showed

more strongly increasing effective EPSC durations during stimulus trains, especially at high

stimulation frequencies, implying less synchronous release. Since tighter SV to VGCCs coupling

has the opposite effect, i.e., generates shorter EPSC onset delays and more tightly

synchronized release, we conclude that the increase of synaptic strength in $Rac1^{-/-}$ calyces is

not due to tighter spatial coupling between SVs and VGCCs.

Loss of Rac1 facilitates EPSC recovery and RRP replenishment

The kinetics of molecular priming regulates RRP replenishment and determines steady-state release rates during high-frequency stimulation (Lipstein et al., 2013; Ritzau-Jost et al., 2018; Lipstein et al., 2021). Since *Rac1* deletion increased steady-state release during 50 Hz stimulus trains, we hypothesized that SV pool replenishment proceeds faster in the absence of Rac1. To test how Rac1 loss influences RRP replenishment, we applied afferent fiber stimulation using a paired train protocol consisting of a 500 Hz depleting train (50 APs) followed by a second 500 Hz test train at varying recovery intervals (Fig. 5). Recovery was then measured for both the initial EPSC amplitude and the RRP estimate of the test trains. Recovery of the initial EPSC amplitude was quantified in terms of both its absolute (Fig. 5A2) and its fractional value (Fig. 5A3), with the latter being the ratio $(EPSC_{test} - EPSC_{ss}) / (EPSC_1 - EPSC_{ss})$, where $EPSC_{test}$, $EPSC_1$ and $EPSC_{ss}$ are the initial amplitude of the test train, and the first and the steady-state amplitudes of the 500 Hz conditioning train, respectively. Recovery of absolute $EPSC_1$ amplitude was significantly different between $Rac1^{+/+}$ and $Rac1^{-/-}$ ($Rac1^{+/+}$: A = 1.4, τ_{fast} = 17 ms, τ_{slow} = 2.7 s, f_{slow} = 0.67, τ_w = 1.8 s vs. $Rac1^{-/-}$: A = 3.4, τ_{fast} = 29 ms, τ_{slow} = 2.3 s, f_{slow} = 0.71, τ_w = 1.6 s, p < 0.001, n = 15/15) and fractional EPSC recovery was almost 50% faster in $Rac1^{-/-}$ ($Rac1^{+/+}$: τ_{fast} = 36 ms, τ_{slow} = 2.1 s, f_{slow} = 0.89, τ_w = 1.9 s vs. $Rac1^{-/-}$: τ_{fast} = 46 ms, τ_{slow} = 1.7 s, f_{slow} = 0.76, τ_w = 1.3 s, p < 0.001, Fig. 5A). Next, we analyzed fractional RRP recovery by dividing the RRP estimate of the test train by the RRP estimate of the depleting train and found that RRP recovery rates were about 40% faster in $Rac1^{-/-}$ ($Rac1^{+/+}$: τ_{fast} = 20 ms, τ_{slow} = 1.6 s, f_{slow} = 0.69, τ_w = 1.1 s vs. $Rac1^{-/-}$: τ_{fast} = 39 ms, τ_{slow} = 1.3 s, f_{slow} = 0.52, τ_w = 0.7 s, p < 0.001, Fig. 5B). Finally, we compared paired-pulse ratios ($EPSC_2 / EPSC_1$) at individual recovery intervals. Independent of recovery interval, PPR was consistently lower in $Rac1^{-/-}$ ($Rac1^{+/+}$ = 1.3 ± 0.1 vs. $Rac1^{-/-}$ = 1.1 ± 0.1, p < 0.001, Fig. 5C), consistent with an increase in P_r.

Numerical simulations of STP and EPSC recovery are consistent with altered SV priming after Rac1 loss

Since the loss of presynaptic Rac1 caused three principal changes in synaptic function: (*i*) increased synaptic strength (Fig. 2D1), (*ii*) increased steady-state release during 50 Hz stimulation (Fig. 2A2), and (*iii*) accelerated EPSC recovery following conditioning 500 Hz trains (Fig. 5), we next sought to corroborate our conclusions about underlying synaptic mechanisms by reproducing experimental data in numerical simulations. To do so, we used two distinct STP models: (1) a single pool model with a Ca^{2+}-dependent SV pool replenishment similar to the release-site model of Weis et al. (1999) and (2) a recently established sequential two-step priming scheme (Neher and Taschenberger, 2021; Lin et al., 2022).

The single-pool model assumes a single type of release site to which SVs can reversibly dock, and SV docking and priming is described by a single transition step (Fig. 6A1). The kinetics of the forward (priming) transition (determined by the rate constant k_f) is characterized by a Michaelis-Menten-like dependence on cytosolic $[Ca^{2+}]$ (Fig. 6A2 inset) while the backward (unpriming) transition has a fixed rate constant (k_b). For resting synapses, the equilibrium between empty sites (ES) and sites occupied by a docked and primed SV (DS) is given by the ratio k_f/k_b at basal $[Ca^{2+}]$ in this scheme. The total number of SV docking sites (N_{total}), priming and unpriming rate constants (k_f and k_b), Ca-dependence of the priming step (parameters σ and $K_{0.5}$), release probability P_r, and the time course of $[Ca^{2+}]$ regulating the priming speed ('effective $[Ca^{2+}]$') were free parameters and adjusted by trial and error (Fig. 6 – Figure supplement 1) to reproduce experimentally observed differences in the time course of fractional recovery (Fig. 6A3), initial synaptic strength, and time course of STP during 50 and 500 Hz stimulation (Fig. 6A4). A comparison between simulated and experimental data shows that the single pool model can describe the functional differences between *Rac1*$^{+/+}$ and *Rac1*$^{-/-}$ synapses, reproducing larger initial synaptic strength, larger steady-state release during 50 Hz trains in Rac1-deficient synapses, and similar steady-state release during 500 Hz trains in both genotypes (Fig. 6A5).

To describe both $Rac1^{+/+}$ and $Rac1^{-/-}$ synapses adequately, we had to introduce changes

in N_{total}, priming kinetics and P_r. The best fit was achieved by adjusting the ratio k_f/k_b at basal

[Ca^{2+}] to yield a number of release sites occupied with a docked/primed SV at rest (RRP) of

2150 and 2532 and by setting P_r to 0.08 and 0.165 for $Rac1^{+/+}$ and $Rac1^{-/-}$ synapses,

respectively (Fig. 6 – Figure supplement 1). Larger initial synaptic strength in $Rac1^{-/-}$ synapses

resulted from ~1.2 fold and ~2 fold changes in RRP and P_r, respectively, which is consistent with

the analysis shown in Fig. 2, which rests on similar assumptions as the single-pool STP model.

The higher steady state release in $Rac1^{-/-}$ synapses is primarily a result of their higher priming

rate constant k_1 for effective [Ca^{2+}] up to ~1 µM, while a faster saturation of k_1 with effective

[Ca^{2+}] above ~1 µM in $Rac1^{-/-}$ synapses (Fig. 6A2) accounts for steady-state release during

500 Hz stimulation being similar to $Rac1^{+/+}$ (Fig. 6A5).

Subsequently, we simulated our experimental data using a two-step priming scheme,

which postulates a sequential build-up of the SV fusion apparatus and distinguishes two distinct

priming states - an immature loosely-docked state (LS) and a mature tightly-docked (TS) state

(Neher and Brose, 2018; Neher and Taschenberger, 2021; Lin et al., 2022). The two-step

priming scheme (Fig. 6B1) reproduces functional changes in $Rac1^{-/-}$ synapses, including the

accelerated EPSC recovery after pool depletion (Fig. 6B3), increased initial EPSC amplitudes,

and elevated steady-state release during 50 Hz trains (Fig. 6B4). In contrast to simulations using

the single-pool model (Fig. 6A1), only the model parameters determining the SV priming kinetics

needed to be adjusted to reproduce the changes in $Rac1^{-/-}$ synapses when using the sequential

two-step model (Fig. 6B1). In resting $Rac1^{-/-}$ synapses, the priming equilibrium was shifted

towards a higher fraction of TS SVs in addition to a slight reduction of the fraction of empty sites

(ES) because of the increased priming rate constants k_1 and k_2 at resting cytosolic [Ca^{2+}]. The

higher abundance of fusion-competent SVs in resting $Rac1^{-/-}$ synapses ($Rac1^{+/+}$ = 691 TS SVs

vs. $Rac1^{-/-}$ = 1666 TS SVs) fully accounts for their increased synaptic strength, while the P_r and

the total number of sites N_{total} were constrained to the same values for $Rac1^{+/+}$ and $Rac1^{-/-}$

synapses. Proper adjustment of the Ca^{2+}-dependence of k_1 and k_2 (Fig. 6B2) reproduces the

different steady-state release in $Rac1^{-/-}$ compared to $Rac1^{+/+}$ synapses (Fig. 6B4). The steeper

increase in k_1 and k_2 with increasing $[Ca^{2+}]$ (Fig. 6B2) accounts for the faster recovery early after

cessation of 500 Hz conditioning when $[Ca^{2+}]$ is still elevated (Fig. 6B3). Thus, the sequential

two-step model is capable of replicating the observed changes in $Rac1^{-/-}$ synapses (Fig. 6B3,

6B5) solely by modifying the priming equilibrium between LS and TS SVs in resting synapses

and carefully adjusting the Ca^{2+}-dependence of the two SV priming steps with unaltered model

parameter values for P_r and N_{total}.

357   In summary, the experimental data available do not allow us to unambiguously favor one

model over the other (Fig. 6A1 vs. 6B1). Both models reproduce differences in STP and EPSC

recovery between $Rac1^{-/-}$ and $Rac1^{+/+}$ synapses, and both models require increased SV priming

speed at resting and intermediate $[Ca^{2+}]$ in $Rac1^{-/-}$ synapses to reproduce the data faithfully.

While the single-pool model (Fig. 6A1) requires different values for N_{total} and P_r, to account for

the changes observed between $Rac1^{-/-}$ and $Rac1^{+/+}$ synapses, necessary changes to model

parameters were limited to those determining the SV priming equilibrium (k_1, b_1, k_2, b_2) when

using the sequential two-step model (Fig. 6B1).

Loss of Rac1 impacts delay and temporal precision of AP firing in response to SAM

stimuli

Animal vocalization, including human speech, is characterized by rapid amplitude modulations

(Joris et al., 2004). These amplitude modulations can be mimicked by sinusoidal amplitude-

modulated stimuli (SAM), which will produce periodic fluctuations in the firing rates of the calyx

of Held (Mc Laughlin et al., 2008; Tolnai et al., 2008). Therefore, different SAM frequencies will

differentially stress SV release and recovery mechanisms and can be used to reveal how the

observed changes in synaptic strength and SV pool recovery potentially impact auditory

signaling in $Rac1^{-/-}$ calyces. Using *in vivo* responses to SAM stimuli (Tolnai et al., 2008) which

contained modulation frequencies between 20 and 2000 Hz as templates for afferent fiber

stimulation, we recorded postsynaptic APs in the loose-patch configuration (Fig. 7). First, we analyzed the success probability of presynaptic stimuli triggering a postsynaptic AP to assess the reliability of synaptic transmission and found no difference between $Rac1^{+/+}$ and $Rac1^{-/-}$ for all modulation frequencies tested (Fig. 7B1). Since temporal precision is crucial in auditory signaling, we then analyzed if loss of Rac1 affected the temporal precision of AP generation by calculating the standard deviation of AP delays ('AP jitter'). For all modulation frequencies, temporal precision was unchanged when analyzed for the complete stimulus (Fig. 7B2). However, our previous analysis showed that EPSC onset delay and synchronicity were only affected at high stimulation frequencies. Since the SAM stimuli generate periods of high firing activity interspersed with periods of low activity, we tested how preceding activity influenced AP jitter and AP delay. First, we used the preceding inter-spike interval (ISI) to estimate preceding activity and calculated AP jitter and AP delay for different ISIs (Fig. 7C1). Both AP jitter and AP delay were comparable between $Rac1^{+/+}$ and $Rac1^{-/-}$ for most ISIs, but AP delay in $Rac1^{-/-}$ increased slightly at short ISIs. Previous studies showed that EPSC/EPSP amplitudes, and subsequent AP generation, are influenced by preceding neuronal activity (Haustein et al., 2008; Englitz et al., 2009; Lorteije et al., 2009; Sonntag et al., 2011; Yang and Xu-Friedman, 2015; Ghanbari et al., 2020). Since the analysis shown in Fig. 7C1 only considered the last ISI, it may result in an incomplete estimate of preceding activity. Therefore, we additionally estimated activity levels by calculating the sum of all preceding APs weighted by their distance to the AP under observation (Fig. 7C2). Using this approach, we found that AP jitter was similar in $Rac1^{+/+}$ and $Rac1^{-/-}$ synapses for low to moderate activity levels but diverged at increasing activity levels, with $Rac1^{-/-}$ showing higher AP jitter ($Rac1^{+/+} = 65 \pm 32$ µs vs. $Rac1^{-/-} = 99 \pm 62$ µs, $p < 0.001$, n = 14/18). Likewise, AP delays were similar for low activity levels but increased more strongly in $Rac1^{-/-}$ with increasing activity ($Rac1^{+/+} = 110 \pm 52$ µs vs. $Rac1^{-/-} = 158 \pm 101$ µs, $p = 0.03$). These data suggest that the activity-dependent increase in EPSC onset and reduction

400   in EPSC synchronicity in *Rac1*$^{-/-}$ at high stimulation frequencies translates into a small but

401   consistent increase in AP delays and AP jitter.

Discussion

By genetically ablating Rac1 at the calyx of Held after hearing onset, we identified presynaptic roles of Rac1 in regulating synaptic transmission and plasticity in a native neuronal circuit. Based on our experimental data and numerical simulations, we identify presynaptic Rac1 as a key regulator of synaptic strength and SV pool replenishment by controlling SV priming kinetics and, depending on model assumptions, by either regulating P_r (single-pool model) or by regulating the abundance of tightly docked and fusion-competent SVs (sequential two-step priming model). In conclusion, we propose that presynaptic Rac1 is a critical regulator for encoding information flow in neuronal circuits.

Presynaptic Rac1 regulates initial synaptic strength

Our finding that Rac1 regulates initial synaptic strength contrasts with a recent study in primary cultured hippocampal neurons which concluded that presynaptic Rac1 had no such effect (O'Neil et al., 2021). Multiple reasons may explain these differences. One cause may be due to the specific conditions of a native neuronal circuit which are not fully replicated under *in vitro* conditions in cultured neuronal circuits. We specifically abated Rac1 at an advanced developmental stage, two days after hearing onset, at which neuronal circuit properties are well defined, and the calyx of Held is functionally largely mature (Englitz et al., 2009; Sonntag et al., 2009; Sonntag et al., 2011; Borst and Soria van Hoeve, 2012). Although Rac1 was ablated at 10 days *in vitro* in the study by O'Neil et al. (2021), the corresponding developmental *in-vivo* stage is difficult to estimate. In addition, it is unknown how and to what extent culture conditions determine synaptic transmission and STP characteristics of hippocampal synapses developing *in vitro*. Another critical difference between our study and that of O'Neil et al. (2021) was the recording condition. We used 1.2 mM external $[Ca^{2+}]$ and physiological temperature to mimic *in-vivo* conditions (Lorteije et al., 2009; Borst, 2010; Forsberg et al., 2019), while O'Neil et al. (2021) performed experiments at 2 mM external $[Ca^{2+}]$ and room temperature. As a result, our estimates for initial P_r based on the ratio of $EPSC_1$ over RRP had a mean of ~0.1 in *Rac1*$^{+/+}$

calyces, and values for individual synapses never exceeded 0.15, which allowed us to observe a

two-fold increase following Rac1 ablation. In the cultured neurons, using a similar analysis, a

mean initial P_r of ~0.4 was reported for $Rac1^{+/+}$ with values for individual synapses frequently

exceeding 0.5 in both excitatory and inhibitory neurons. It is therefore possible that in cultured

neurons, P_r was close to saturation, thereby occluding any further increase in initial synaptic

strength in $Rac1^{-/-}$ neurons. In addition, many regulatory steps in the SV cycle are temperature-

dependent (Chanaday and Kavalali, 2020), and in mouse hippocampal synapses, actin-

dependent synaptic release enhancement is restricted to physiological temperatures (Jensen et

al., 2007).

Presynaptic Rac1 regulates SV pool replenishment

The availability of fusion-competent SVs critically determines synaptic strength during ongoing

stimulation. The steady-state occupancy of the RRP varies with stimulation rates (Hallermann

and Silver, 2013; Neher, 2015) and is determined by the SV pool replenishment kinetics, which

are critical for maintaining synaptic transmission. Pool replenishment is typically quantified by

two methods (Hallermann and Silver, 2013; Neher, 2015): (1) the replenishment rate during

ongoing stimulation can be obtained as the slope of line fits to the steady-state portion of

cumulative EPSC trains assuming that during ongoing stimulation quantal release is balanced by

newly replenished SVs. (2) Replenishment rate constants after stimulation can be estimated

from the time constants of exponential fits to the time course of the fractional EPSC or pool

recovery plotted as a function of the inter-stimulus interval following conditioning trains.

We found that $Rac1^{-/-}$ calyces had larger steady-state EPSC amplitudes, and the slope

of the line fits to cumulative EPSC trains revealed a faster replenishment rate at 50 Hz

stimulation, similar to observations in hippocampal cultures at 20 and 40 Hz stimulation (O'Neil

et al., 2021). In contrast, during ongoing 500 Hz stimulation, steady-state EPSC amplitudes and

slope of the line fits were similar in $Rac1^{+/+}$ and $Rac1^{-/-}$ synapses. However, time courses of

EPSC or pool recovery after 500 Hz conditioning showed an almost 50% faster recovery when fit

by bi-exponential functions. This acceleration was largely due to a speed-up of τ_{slow} and a larger

contribution of τ_{fast}, which according to our simulations, reflects a faster recovery occurring at

cytosolic [Ca^{2+}] relatively close to resting values. In addition, we estimated a slightly larger

relative fraction of the fast recovery component in *Rac1^{−/−}* synapses, which, according to our

simulations, reflects the magnitude of Ca-dependent acceleration of SV pool replenishment

immediately after cessation of 500 Hz conditioning while cytosolic [Ca^{2+}] decays back to resting

values. These observations are consistent with the experimental finding of an enhanced steady-

state release in *Rac1^{−/−}* synapses during 50 Hz stimulation, which reflects their faster pool

replenishment at [Ca^{2+}] <1 μM because of the steeper increase of the forward priming rate

constants with increasing cytosolic [Ca^{2+}]. The steady-state release was similar during 500 Hz

trains when the synapses regenerate fusion-competent SVs at similar maximum rates in *Rac1^{+/+}*

and *Rac1^{−/−}*. This was primarily caused by a stronger saturation of the Ca-dependence of the

priming rate constant for [Ca^{2+}] >1 μM in *Rac1^{−/−}* synapses. Because of the similar steady-state

EPSC levels measured at 500 Hz, it is unlikely that Rac1 controls an upstream limit on the

priming process.

Presynaptic Rac1 regulation of spontaneous release

At mature calyx of Held synapse, blocking presynaptic Ca^{2+} influx through VGCCs does

not change spontaneous release rates (Dong et al., 2018). Thus, mEPSC frequencies can be

interpreted as a readout of intrinsic SV fusogenicity at basal [Ca^{2+}] with higher mEPSC

frequencies reflecting increased SV fusogenicity (Basu et al., 2007; Schotten et al., 2015; Dong

et al., 2018), provided that pool size remains unaltered. However, SVs undergoing AP-evoked

and spontaneous release may not necessarily originate from the same SV pools. Nevertheless,

all SVs have to undergo a priming step to acquire fusion competence. Actin is found throughout

the presynaptic terminal and application of latrunculin causes a rapid increase in mEPSCs rates

in the presence of internal Ca^{2+} chelators (BAPTA-AM, EGTA-AM) while also augmenting AP

evoked release (Morales et al., 2000). Thus, although Rac1 loss may differentially affect

mEPSCs rates and rates of AP-evoked release, increased mEPSC frequencies following Rac1

loss are consistent with a higher fusogenicity of those SVs contributing to spontaneous release

presumably due to Rac1's role as actin regulator.

Presynaptic Rac1 does not affect the spatial coupling between docked SV and VGCCs

After docking to release sites, SVs become fusion-competent during the assembly of the

molecular fusion machinery, which defines the intrinsic SV fusogenicity. In addition to the

intrinsic SV fusogenicity, the probability of SVs undergoing fusion upon AP arrival is determined

by their spatial coupling distances to VGCCs, which determines the local $[Ca^{2+}]$ 'seen' by the

vesicular Ca^{2+} sensor for fusion (Neher, 2010; Schneggenburger and Rosenmund, 2015).

Therefore, the proximity of SVs to VGCCs (also termed 'positional priming') is a critical

determinant of transmitter release (Wadel et al., 2007; Chen et al., 2015; Stanley, 2016). At the

calyx of Held after hearing onset, AP-evoked SV release is controlled by local $[Ca^{2+}]$

nanodomains generated around $Ca_V2.1$ channels which are located at ~25 nm distance from

docked SVs (Fedchyshyn and Wang, 2005; Chen et al., 2015; Nakamura et al., 2015), resulting

in fast synchronous transmitter release. Since Rac1 regulates actin dynamics, loss of Rac1 may

result in tighter spatial coupling between SV and VGCCs, thereby increasing synaptic strength. If

SVs were located in closer proximity to VGCCs, we would expect to see decreased initial EPSC

onset delay times or decreased EPSC onset delay times during both 50 and 500 Hz stimulation.

However, we found that EPSCs in $Rac1^{-/-}$ calyx synapses had increased EPSC onset delays

and less synchronous EPSCs, particularly at high firing rates, which is inconsistent with SVs

being more tightly coupled to VGCCs.

502   The EPSC onset delay includes the AP conduction delay, the time between triggering an

AP by afferent fiber stimulation and its arrival at the calyx terminal, and the transmitter release

delay, the time between presynaptic AP and SV fusion which is dependent on the distance

between SVs to VGCCs (Fedchyshyn and Wang, 2007). Therefore, we cannot exclude that loss

of Rac1 may prolong conduction delay, thereby obscuring a role of Rac1 in regulating SV to

VGCC coupling distances. However, EPSC onset delays of the first EPSC or during 50 Hz train

stimuli were similar between $Rac1^{+/+-}$ and $Rac1^{-/-}$ calyces. The EPSC onset delay was only

increased for steady-state EPSCs during 500 Hz stimulation. In addition, if longer conduction

delays were solely responsible for the increase in EPSC onset delays, no changes in release

kinetics would be expected. However, during both 50 and 500 Hz stimulus trains, the effective

EPSC duration increased more strongly in $Rac1^{-/-}$ calyces compared to $Rac1^{+/+}$ consistent with

a less synchronized release time course in the former, possibly because of a larger fraction of

SVs located distally from VGCCs contributing to release in $Rac1^{-/-}$.

Another possibility is that presynaptic APs in $Rac1^{-/-}$ were broader and thereby increased

presynaptic Ca^{2+} influx which contributed to increased synaptic strength and faster EPSC

recovery. Although broader presynaptic APs are expected to widen the release transient and

may delay presynaptic Ca^{2+} influx occurring mainly during AP repolarization (Borst and

Sakmann, 1998; Li et al., 2007; Wang et al., 2008; Kochubey et al., 2009) and thereby increase

synaptic delays, we did not observe an increased effective EPSC duration or longer EPSC onset

delays between $Rac1^{+/+-}$ and $Rac1^{-/-}$ calyces for $EPSC_1$. Even though we cannot exclude a

Rac1-dependent regulation of presynaptic AP waveform, we do not consider such scenario very

likely. Using TEA, a Kv3 channel blocker, to broaden the presynaptic AP impairs high-frequency

firing at the calyx of Held (Wang et al., 1998; Wang and Kaczmarek, 1998; Johnston et al.,

2010). Since $Rac1^{-/-}$ calyces were able to follow high-frequency stimulation and EPSC onset

delays were similar for $EPSC_1$ and throughout the 50 Hz stimulus train it is unlikely loss of Rac1

caused a general broadening of APs. Therefore, in the absence of experimental evidence

supporting Rac1 loss-induced changes in SV to VGCC coupling distances, we propose that

Rac1 regulates synaptic strength and RRP recovery at the level of molecular priming, either by

increasing the intrinsic fusogenicity of SVs and their P_r or by increasing the abundance of fusion-

competent tightly docked SVs. In both scenarios, increased P_r (single pool model) or a shift in

priming equilibrium in favor of TS SVs (two-step model) could account for the increased initial

synaptic strength in $Rac1^{-/-}$ synapses and, at least in part, for their higher mEPSC rates.

Although our experimental data do not allow us to favor one scenario over the other

unambiguously, both models suggest that Rac1 regulates SV priming. While it was not required

to postulate differences in P_r when simulating $Rac1^{+/+-}$ and $Rac1^{-/-}$ synapse with a sequential

two-step model, we cannot rule out that P_r is indeed changed after Rac1 loss. It is well possible

to implement a higher P_r in $Rac1^{-/-}$ calyces within the framework of a sequential two-step model

and future experiments combined with simulations will be needed to explore this possibility

further.

Loss of Rac1 might also impact endocytosis through changes in actin signaling (Wu and

Chan, 2022). However, at the calyx of Held, the endocytic role of F-actin appears to be

negligible as actin depolymerization using latrunculin did not affect endocytosis (Eguchi et al.,

2017; Piriya Ananda Babu et al., 2020). Furthermore, endocytosis acts on a time scale of

seconds and would be too slow to affect SV pool replenishment during short stimulus trains

(Armbruster and Ryan, 2011). Although ultrafast endocytosis occurs on a timescale of tens of

milliseconds it is unlikely to contribute to the $Rac1^{-/-}$ phenotype as SV reformation is too slow

with a time scale of tens of seconds (Watanabe et al., 2013a; Watanabe et al., 2013b). In

addition, endocytosis is postulated to play a role in release site clearance (Neher, 2010; Sakaba

et al., 2013) as the perturbation of endocytosis increases the rate of synaptic depression and

slows RRP recovery (Wen et al., 2013) which we did not observe in the $Rac1^{-/-}$ calyces.

Rac1 regulation of auditory signaling and information processing

The ability to accurately encode sound information requires synaptic transmission in the lower

auditory brainstem to drive and sustain precise AP firing over rapid and large fluctuations of AP

firing rates up to the kilohertz range (Grothe et al., 2010; Borst and Soria van Hoeve, 2012;

Friauf et al., 2015). The calyx of Held - MNTB principal cell synapse is a failsafe and reliable

synaptic relay that transforms afferent AP spike patterns from the CN into precisely timed

inhibition to several mono- and binaural nuclei (Friauf and Ostwald, 1988; Spirou et al., 1990;

Joris et al., 2004; Englitz et al., 2009; Lorteije et al., 2009; Sonntag et al., 2011). The use of SAM

stimuli which mimics the neuronal response to environmental sounds (Joris et al., 2004; Tolnai

et al., 2008), suggests that loss of Rac1 did not deteriorate the faithful auditory signaling

properties over different amplitude modulation frequencies. However, we found that increased

EPSC onset and decreased SV release synchronicity in $Rac1^{-/-}$ translated into larger AP delay

and AP jitter at high activity levels. Although the absolute changes were modest, the need for

high temporal precision in the auditory brainstem might cause a more severe impact in

downstream nuclei, such as the lateral and medial superior olive. Since the calyx of Held

generates suprathreshold EPSPs even under short-term depression (Lorteije et al., 2009;

Lorteije and Borst, 2011), changes in synaptic strength caused by Rac1 ablation may affect

postsynaptic AP timing but should have little effect on postsynaptic AP firing success probability

in MNTB principal neurons. Synaptic transmission at other auditory synapses in the SPON,

MSO, or LSO that operate close to AP threshold or rely on synaptic integration to encode

information is likely more substantially affected by Rac1 ablation.

Ideas and speculation

Rac1 regulation of F-actin dynamics controls SV priming

F-actin regulates several SV cycle steps, which could impact SV replenishment rates (Morales et

al., 2000; Sakaba and Neher, 2003; Cingolani and Goda, 2008; Sun and Bamji, 2011; Waites et

al., 2011; Lee et al., 2012; Lee et al., 2013; Montesinos et al., 2015; Rust and Maritzen, 2015).

Specifically, F-actin may act as a physical barrier within the presynaptic AZ (Cingolani and

Goda, 2008) as the application of latrunculin transiently increases P_r and mEPSC rates (Morales

et al., 2000). Studies in which Septin 5 (Yang et al., 2010) or the actin cytoskeleton was

disrupted (Lee et al., 2012) are in support of the idea of F-actin as a structural barrier potentially

impacting molecular priming or SV to VGCC coupling (Yang et al., 2010). Assuming that SV

docking distance relative to the plasma membrane corresponds to SV priming (Imig et al., 2014;

Jung et al., 2016; Pulido and Marty, 2018), we propose that Rac1's regulation of F-actin may

impact the physical barrier and thus affect the conversion from LS to TS, although the P_r of the

final TS state in Rac1-deficient terminals remains unknown (Fig. 8). Based on our simulations,

we speculate that the loss of Rac1 results in local actin depolymerization in the AZ, thereby

promoting the LS to TS transition and/or increasing P_r of fusion-competent SVs by allowing

vesicle docking closer to the plasma membrane. In support of our model, increased synaptic

strength has been demonstrated to correlate with shortened tethering filaments that resulted in

SVs positioned more closely to the presynaptic membrane (Jung et al., 2021). It is also in line

with the finding that GIT proteins (Montesinos et al., 2015), Arp2/3 (O'Neil et al., 2021), and

Piccolo (Waites et al., 2011), which are all regulators of F-actin dynamics, also control synaptic

strength and suggests that molecules regulating F-actin dynamics at the AZ could be key

regulators of SV replenishment.

Mutations in Rac1 that result in loss or gain of function are associated with intellectual

disability (Reijnders et al., 2017; Zamboni et al., 2018). While much attention has focused on

Rac1 dysregulation in the dendritic compartment or the role of Rac1 in neuronal development,

our work demonstrates that presynaptic loss of Rac1 increases synaptic strength and EPSC

recovery independent of its developmental role. Human mutations of Rac1 may affect synaptic

strength at many different synaptic connections and potentially alter the excitation-inhibition

balance and synaptic information processing in neuronal circuits associated with neurological

disorders and addiction (Dietz et al., 2012; Bai et al., 2015; Wright et al., 2020).

Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (*Mus musculus*)	Rac1^{tm1Djk}/J (Rac1$^{flox/flox}$)	Jackson Laboratory (Glogauer et al., 2003)	RRID:IMSR_JAX:005550	either sex
antibody	anti-GFP (rabbit polyclonal)	Abcam	Cat# ab6556 RRID:AB_305564	EM (0.1 µg/mL)
antibody	6 nm colloidal Gold-AffiniPure anti-rabbit IgG (donkey polyclonal)	Jackson ImmunoResearch	Cat# 711-195-152 RRID:AB_2340609	EM (1:100)
sequence-based-reagent	primer: 5'-TCC AAT CTG TGC TGC CCA TC-3'	(Glogauer et al., 2003)		
sequence-based-reagent	primer: 5'-GAT GCT TCT AGG GGT GAG CC-3'	(Glogauer et al., 2003)		
recombinant DNA reagent	HdAd 28E4 hsyn iCre EGFP (viral vector)	Samuel M. Young, Jr., University of Iowa		
recombinant DNA reagent	HdAd 28E4 hsyn iCre mEGFP (viral vector)	Samuel M. Young, Jr., University of Iowa		
recombinant DNA reagent	HdAd 28E4 hsyn mEGFP (viral vector)	Samuel M. Young, Jr., University of Iowa		
chemical compound, drug	kynurenic acid	Tocris Bioscience	Cat# 0223	
chemical compound, drug	lidocaine N-ethyl bromide (QX-314)	Sigma Aldrich	Cat# L5783	
chemical compound, drug	D-AP5	Tocris Bioscience	Cat# 0106	
chemical compound, drug	(-)-bicuculline methochloride	Tocris Bioscience	Cat# 0131	
chemical compound, drug	strychnine hydrochloride	Tocris Bioscience	Cat# 2785	
chemical	tetraethylammonium	Sigma Aldrich	Cat# T-2265	

compound, drug	chloride			
chemical compound, drug	tetrodotoxin	Alomone labs	Cat# T-550	
chemical compound, drug	Cadmium chloride hemi(pentahydrate)	Sigma Aldrich	Cat# C3141	
software, algorithm	Matlab	The Mathworks	RRID:SCR_001622; v9.10	
software, algorithm	Patchmaster	HEKA; Harvard Bioscience	RRID:SCR_000034; v2x90.2	
software, algorithm	Igor Pro	Wavemetrics	RRID:SCR_000325; v6.37	
software, algorithm	Fiji	https://fiji.sc/	RRID:SCR_002285	
software, algorithm	Patcher's Power Tools	Max Planck Institute for Biophysical Chemistry; Gottingen; Germany	RRID:SCR_001950; v2.19	
software, algorithm	StereoDrive	Neurostar	N/A; v3.1.5	
software, algorithm	Live Acquisition	Thermo Fisher Scientific	N/A; v2.1.0.10	
software, algorithm	Neuromatic	(Rothman and Silver, 2018)	RRID:SCR_004186	

Animals

All experiments were performed following animal welfare laws and approved by the Institutional Committee for Care and Use of Animals at the University of Iowa (PHS Assurance No. D16-00009 (A3021-01) (Animal Protocol 0021952) and complied with accepted ethical best practices. Animals were housed at a 12-hour light/dark cycle and had access to food and water ad libitum. Experiments were performed on Rac1^{tm1Djk} (*Rac1$^{fl/fl}$*) mice (Glogauer et al., 2003) (RRID:IMSR_JAX: 005550, The Jackson Laboratory, Bar Harbor, USA) of either sex. Animals of this line possess a *loxP* site flanking exon 1 of the *Rac1* gene disrupting Rac1 expression after recombination mediated by Cre recombinase. Genotyping was performed using PCR-amplification with the following primers 5'-TCC AAT CTG TGC TGC CCA TC-3' and 5'-GAT GCT TCT AGG GGT GAG CC-3' and amplification products (*Rac1$^{fl/fl}$*: 242 bp, wildtype: 115 bp)

618 were separated gel electrophoresis on a 1.5% agarose gel (Glogauer et al., 2003). Viral vectors

619 were injected at postnatal day 14 (P14) and experiments were performed at P28-30. All

620 available measures were taken to minimize animal pain and suffering.

DNA construct and recombinant viral vector production

Helper-dependent adenoviral vectors (HdAd) expressing a codon-optimized Cre recombinase

(Cre) (Shimshek et al., 2002) were produced as previously described (Montesinos et al., 2016;

Lubbert et al., 2017). These HdAd vectors contain two independent transgene cassettes that

independently express Cre recombinase and EGFP or myristoylated EGFP (mEGFP) under the

control of 470 bp human synapsin promoter (Lubbert et al., 2017). In brief, the expression

cassette with Cre recombinase was cloned into the pdelta28E4 SynEGFP plasmid using the

*Asc*I enzyme digestion site. The final plasmid has been modified to contain a separate EGFP or

myristoylated EGFP (mEGFP) expression cassette. Then, the pHAD plasmid was linearized by

*Pme*I enzyme to expose the ends of the 5' and 3' inverted terminal repeats (ITRs) and

transfected into 116 producer cells (Profection® Mammalian Transfection System, Promega,

Madison, WI, USA). Helper virus (HV) was added the following day for HdAd production. Forty-

eight hours postinfection, after cytopathic effects have taken place, cells were subjected to three

freeze/thaw cycles for lysis and release of the viral particles. HdAd was purified by CsCl

ultracentrifugation. HdAd was stored at -80 °C in storage buffer (10 mM HEPES, 1 mM MgCl$_2$,

250 mM sucrose, pH 7.4).

Virus injections

Virus injections at P14 were performed as previously described (Lubbert et al., 2019). Briefly,

mice were anesthetized with 5% isoflurane inhalation and anesthesia was maintained with 2%

isoflurane throughout the procedure. Subcutaneous injection of physiological saline, lidocaine,

bupivacaine, and meloxicam was used to treat loss of liquid and alleviate pain. The injection site

was determined using the Stereodrive software (Neurostar) and corrected for head orientation

and tilt. A small hole (diameter <1 mm) was drilled into the skull using a foot-pedal controlled drill

MH-170 (Foredom). The virus solution was injected via a glass pipet (Drummond) at a rate of

100 nL/min with a nanoliter injector (NanoW, Neurostar). Following the injection, the glass

needle was left in place for 1 min to dissipate the pressure and then slowly retracted. Animals

were then placed under an infrared heat lamp and allowed to recover before being returned to

their respective cages with their mother.

Confocal imaging and reconstruction of presynaptic terminals

For reconstruction of the calyx of Held terminals, animals were injected with viral vectors

expressing either mEGFP (yielding $Rac1^{+/+}$) or Cre recombinase and mEGFP (yielding $Rac1^{-/-}$).

Mice were anesthetized with an intraperitoneal injection of tribromoethanol (250 mg/kg body

weight) and transcardially perfused with ice-cold 0.1 M phosphate buffer (PB, pH 7.4) followed

by 4% paraformaldehyde (PFA) in 0.1 M PB. Brains were removed and post-fixed overnight in

4% PFA solution at 4°C. The next day, brains were sliced into 40 μm sections on a Leica

VT1200 vibratome. mEGFP positive slices were identified and mounted on cover slips with Aqua

Polymount (Polysciences, Inc., Warrington, PA, USA). Confocal images were acquired with a

Zeiss LSM 700 or 880 confocal scanning microscope using a 63x/1.3 NA Apochromat multi-

immersion objective. Image stacks were collected using 0.44 um plane line scans with line

average of four times. Images were processed using Fiji (Schindelin et al., 2012) (http://fiji.sc.,

RRID:SCR 002285). Calyx reconstructions were performed blind to genotype using Imaris

Measurement Pro (BitPlane) using automatic signal detection manual curation in single planes in

Z-stack confocal images as previously described (Radulovic et al., 2020).

Electron microscopy

Preembedding immuno-electron microscopy was performed as previously described

(Montesinos et al., 2015; Dong et al., 2018). Briefly, $Rac^{fl/fl}$ mice injected at P14 with HdAd co-

expressing Cre-recombinase and EGFP were anesthetized and perfused transcardially at P28

with phosphate-buffered saline (PBS, 150 mM NaCl, 25 mM Sørensen's phosphate buffer,

pH 7.4) followed by fixative solution for 7-9 min containing 4% PFA, 0.5% glutaraldehyde, and

0.2% picric acid in 100 mM Sørensen's phosphate buffer (PB, pH 7.4). Brains were post-fixed

with 4% PFA in PB overnight and 50-µm coronal sections of the brainstem were obtained on a

vibratome (Leica VT1200S). Expression of EGFP at the calyx of Held terminals was visualized

using an epifluorescence inverted microscope (CKX41, Olympus) equipped with an XCite Series

120Q lamp (Excelitas Technologies), and only those samples showing EGFP were further

processed. After washing with PB, sections were cryoprotected with 10%, 20%, and 30%

sucrose in PB, submersed into liquid nitrogen and then thawed. Sections were incubated in a

blocking solution containing 10% normal goat serum (NGS), 1% fish skin gelatin (FSG), in 50

mM Tris-buffered saline (TBS, 150 mM NaCl, 50 mM Tris, pH 7.4) for 1h, and incubated with an

anti-GFP antibody (0.1 µg/mL, ab6556, Abcam, RRID: AB_305564) diluted in TBS containing

1% NGS, 0.1% FSG plus 0.05% NaN_3 at 4°C for 48h. After washing with TBS, sections were

incubated overnight in nanogold conjugated donkey anti-rabbit IgG (1:100, Jackson

Immunoresearch, RRID: AB_2340609) diluted in TBS containing 1% NGS and 0.1% FSG.

Immunogold-labeled sections were washed in PBS, briefly fixed with 1% glutaraldehyde in PBS,

and silver intensified using an HQ silver intensification kit (Nanoprobe). After washing with PB,

sections were treated with 0.5% OsO_4 in 0.1 M PB for 20 min, en-bloc stained with 1% uranyl

acetate, dehydrated and flat embedded in Durcupan resin (Sigma-Aldrich). After trimming out

the MNTB region, ultrathin sections were prepared with 40 nm-thickness using an

ultramicrotome (EM UC7, Leica). Sections were counterstained with uranyl acetate and lead

citrate and examined in a Tecnai G2 Spirit BioTwin transmission electron microscope (Thermo

Fisher Scientific) at 100 kV acceleration voltage. Images were taken with a Veleta CCD camera

(Olympus) operated by TIA software (Thermo Fisher Scientific). Images used for quantification

were taken at 60 000x magnification.

TEM image analysis

Calyces positive for Cre expression ($Rac1^{-/-}$) were identified by immunogold labeling with an

anti-GFP antibody and compared to EGFP-negative terminals ($Rac1^{+/+}$) in the same slice or

calyces in the wildtype sample. All TEM data were analyzed using Fiji imaging analysis software

(Schindelin et al., 2012). Each presynaptic active zone (AZ) was defined as the membrane

directly opposing the postsynaptic density, and the length of each AZ was measured. Vesicles

within 200 nm from each AZ were manually selected and their distances relative to the AZ were

calculated using a 32-bit Euclidean distance map generated from the AZ. Synaptic vesicle

distances were binned every 5 nm and counted (Montesinos et al., 2015; Dong et al., 2018).

Vesicles less than 5 nm from the AZ were considered "docked". Three animals for each

condition and ~40 AZs per animal were analyzed. Three researchers performed analysis blind to

genotype independently and results were averaged.

Electrophysiology

Acute coronal brainstem slices (~200 μm) containing the MNTB were prepared as previously

described (Chen et al., 2013; Thomas et al., 2019). Briefly, after decapitation of the animal, the

brain was immersed in low-calcium artificial cerebrospinal fluid (aCSF) solution containing (in

mM): 125 NaCl, 2.5 KCl, 3 $MgCl_2$, 0.1 $CaCl_2$, 10 glucose, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 0.4 L-

ascorbic acid, 3 myo-inositol, and 2 Na-pyruvate, pH 7.3–7.4). Brain slices were obtained using

a Leica VT 1200S vibratome equipped with zirconia ceramic blades (EF-INZ10, Cadence

Blades) as previously described. The blade was advanced at a speed of 20-50 μm/s. Slices

were immediately transferred to an incubation beaker containing standard extracellular solution

(same as above but using 1 mM $MgCl_2$ and 1.2 mM $CaCl_2$) at ~37°C and continuously bubbled

with 95% O_2–5% CO_2. After approximately 45 min of incubation, slices were transferred to a

recording chamber with the same saline solution at physiological temperature (~37°C).

Electrical stimulation of afferent fibers was performed as previously described (Forsythe and

Barnes-Davies, 1993). Briefly, a bipolar platinum-iridium electrode was positioned medially of the

MNTB to stimulate afferent fibers (FHC, Model MX214EP). Postsynaptic MNTB neurons were

whole-cell voltage-clamped at –60 mV using an EPC10/2 amplifier controlled by Patchmaster

Software (version 2x90.2, HEKA Elektronik, RRID:SCR_000034). Slices were continuously

perfused with standard aCSF solution at a rate of 1 mL/min and visualized by an upright

microscope (BX51WI, Olympus) through a 60x water-immersion objective (LUMPlanFL N,

Olympus) and an EMCCD camera (Andor Luca S, Oxford Instruments). To identify calyces

expressing Cre-recombinase and EGFP, the slice was illuminated at an excitation wavelength of

480 nm using a Polychrome V xenon bulb monochromator (TILL Photonics). For whole-cell

voltage-clamp recordings, the standard extracellular solution was supplemented with 1 mM

kynurenic acid to avoid saturation of postsynaptic AMPA receptors, 50 µM D-AP-5 to block

NMDA receptors, and 20 µM bicuculline and 5 µM strychnine to block inhibitory GABA- and

glycine receptors, respectively (all Tocris Bioscience). Patch pipettes had a resistance of ~3–4

MΩ and were filled with the following (in mM): 130 Cs-gluconate, 20 tetraethylammonium (TEA)-

Cl, 10 HEPES, 5 Na_2-phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP, 6 QX-314, and 5 EGTA, pH 7.2

(315 mOsm. Reported voltages are uncorrected for a calculated liquid junction potential of 13

mV. For loose-patch recordings, the extracellular solution was supplemented with 20 µM

bicuculline and 5 µM strychnine, and the patch pipettes were filled with aCSF. To simulate *in*

vivo activity levels, afferent fibers were stimulated with activity patterns previously recorded *in*

vivo in response to sinusoidal amplitude-modulated (SAM) sound stimulation (Tolnai et al.,

2008).

mEPSCs were recorded with the same aCSF supplemented with 50 mM D-AP5, 20 µM

bicuculline, 5 µM strychnine, 1 µM TTX, and 20 mM TEA. In a subset of $Rac1^{-/-}$ recordings, Cd^{2+}

(cadmium chloride), a non-selective Ca^{2+} channel blocker was flushed in during recordings to

determine the impact of VGCC on spontaneous SV release. The baseline mEPSC frequency

was established by recording each cell for at least one minute before Cd^{2+} was flushed in via the

bath perfusion. MNTB principal neurons were whole-cell voltage-clamped at -80 mV and

recorded until enough mEPSCs (>25) were recorded. Overlapping events were excluded from

the analysis.

Data were acquired at a sampling rate of 100 kHz and lowpass filtered at 6 kHz. Series

resistance (3–8 MΩ) was compensated online to <3 MΩ, except for mEPSC recordings where

series resistance (<9 MΩ) was not compensated. All experiments were performed at near-

physiological temperature (36-37°C), and the temperature was maintained by a heated bath

chamber (HCS, ALA Scientific Instruments) and perfusion system (HPC-2, ALA Scientific

Instruments). The temperature of the bath solution was monitored during the experiment using a

glass-coated micro thermistor.

Electrophysiological data analysis

Electrophysiological data were imported to Matlab (version 9.10; The Mathworks,

RRID:SCR_001622) using a custom-modified version of sigTool (Lidierth, 2009) and Igor Pro

(version 8.0.4.2, Wavemetrics, RRID:SCR_000325) equipped with Patcher's Power Tools

(version 2.19, RRID:SCR_001950) and NeuroMatic (Rothman and Silver, 2018)

(RRID:SCR_004186), and analyzed offline with custom-written functions in Matlab and IgorPro.

The remaining series resistance was compensated offline to 0 MΩ with a time lag of 10 µs

(Traynelis, 1998). EPSC amplitudes were measured as peak amplitude minus baseline

preceding the EPSC. RRP size and P_r were calculated using the EQ method (Elmqvist and

Quastel, 1965), back-extrapolation method (SMN with correction) (Neher, 2015), and NpRf

model (Thanawala and Regehr, 2016) as previously described (Lubbert et al., 2019). Onset time

of EPSCs was determined by fitting a Boltzmann function to the EPSC rising flank and

calculating the time point of maximum curvature, as described previously (Fedchyshyn and

Wang, 2007). The duration between stimulus and EPSC onset was defined as EPSC onset

delay. The synchronicity of SV release for individual EPSCs was estimated from the effective

EPSC duration by dividing EPSC charge by EPSC amplitude. This 'effective EPSC duration'

indicates the width of a square current pulse with amplitude and charge identical to that of the

EPSC (Lopez-Murcia et al., 2019). A shorter 'effective width' indicates a shorter release transient

provided that mEPSC kinetics are unchanged.

Recordings of mEPSCs were analyzed using NeuroMatic in Igor Pro. Potential events were

detected when exceeding an amplitude threshold at 4-6 times the standard deviation of the

baseline and all events were manually curated to exclude false positives. Rise time was

measured between 10% and 90% of the peak amplitude.

Recovery of SVs was estimated by non-linear least-square fits to single EPSC and RRP

recovery time courses. Both types of recovery time courses were best fit with a bi-exponential

function of the form:

$$A\left(1 - \left(f\left(e^{-\frac{t}{\tau_{d1}}}\right) + (1-f)\left(e^{-\frac{t}{\tau_{d2}}}\right)\right)\right)$$

with variable A constraint to 1 when fitting fractional recovery. An F-test was used to determine

the better fit. Fractional recovery of the first EPSC was calculated as $(EPSC^1_{test} - EPSC^{SS}_{cond})/$

$(EPSC^1_{cond} - EPSC^{SS}_{cond})$ where $EPSC^1_{test}$ is the amplitude of the first EPSC of the test train,

$EPSC^{SS}_{cond}$ the steady-state amplitude of the conditioning train and $EPSC^1_{cond}$ the amplitude of the

first EPSC of the conditioning train.

To estimate preceding neuronal activity during SAM stimulation, we calculated the interspike

interval (ISI) as the distance to the preceding AP. To analyze AP responses during SAM

stimulation, we used the preceding ISI to estimate the influence of previous activity during SAM

recordings. Since short-term plasticity can extend well beyond the last ISI, we also estimated

preceding activity by calculating the sum of all preceding events, with each event weighted by its

distance to the AP under observation. This weighting was implemented as a single-exponential

decaying function, emphasizing temporally close events over more distant ones (Sonntag et al.,

2011; Keine et al., 2016). The time constant was set to 30 ms, consistent with previous reports

(Yang and Xu-Friedman, 2015). The calculation was also performed with time constants of

10 ms and 100 ms yielding similar results.

Experimental design and statistical analysis

Individual neurons were considered independent samples for electrophysiological data analysis and morphological reconstruction. For thin-section TEM analysis, AZs were considered independent samples (244 total AZs, 6 animals, ~40 AZs each). Statistical analysis was conducted in MATLAB (version 9.4, RRID:SCR_001622) and GraphPad Prism (RRID:SCR_002798). Data distributions were tested for Gaussianity using the Shapiro-Wilk test. To compare two groups, we used a two-tailed unpaired Student's t-test with Welch's correction (normal distribution) or a two-tailed Mann-Whitney U test (non-normal distribution). An RM ANOVA was performed to compare more than two groups with within-subject factors, and p-values were Bonferroni-adjusted for multiple comparisons. Fits to data were subject to F-tests to determine the better model (mono- or biexponential) and for comparison between groups. A p-value of 0.05 was deemed significant for interpreting all statistical tests. Effect sizes were calculated using the MES toolbox in MATLAB (Hentschke and Stuttgen, 2011) and are reported as Cohen's U_1 for two-sample comparison and eta-squared (η^2) for RM ANOVA. No statistical test was performed to pre-determine sample sizes. Exact p-values, test statistics, and effect sizes for all statistical comparisons are summarized in supplemental tables. Boxplots show median, interquartile, and minimum/maximum within 1.5 times the interquartile range. Average data in the text are reported as mean ± standard deviation.

Numerical simulations of STP and EPSC recovery

Simulations of synaptic STP in response to 50 and 500 Hz stimulus trains and of the recovery of EPSC amplitudes after SV pool depletion were performed using two types of kinetics schemes for SV priming and fusion: (*i*) a simple single pool model with a Ca^{2+}-dependent SV pool replenishment as described below, and (*ii*) the sequential two-step model as recently proposed by Lin et al. (2022).

Single pool model

The simple single pool model consisted of a single type of release site to which SVs can

reversibly dock. The total number of sites (N_{total}) is, therefore, given at any given time point *t* by

the sum of empty release sites (N_e(t)) and the release sites occupied by a primed and fusion-

competent SV (N_o(t)): $N_{tot} = N_e(t) + N_o(t)$.

Transitions between N_e(t) and N_o(t) are described by forward (k_f) and backward (k_b) rate

constants according to:

$$\frac{d}{dt} N_o(t) = k_f \cdot N_e(t) - k_b \cdot N_o(t)$$

$$\frac{d}{dt} N_e(t) = k_b \cdot N_o(t) - k_f \cdot N_e(t)$$

826   While the backward (unpriming) rate constant had a fixed value, k_f was assumed to increase

827   with increasing cytosolic $[Ca^{2+}]$ ('effective $[Ca^{2+}]$'). The Ca-dependence of k_f was described by a

828   Michaelis-Menten like saturation according to:

$$k_f(t) = \left(k_{f,rest} + \sigma \cdot ([Ca^{2+}](t) - [Ca^{2+}]_{rest}) \right) / \left(1 + ([Ca^{2+}](t) - [Ca^{2+}]_{rest}) / K_{0.5)} \right)$$

829   where $k_{f,rest}$ is the value of k_f at resting $[Ca^{2+}]$ ($[Ca^{2+}]_{rest}$; assumed to be 50 nM), σ is a slope

830   factor and $K_{0.5}$ is the Michaelis-Menten K_D value.

831   The effective $[Ca^{2+}]$ ($[Ca^{2+}](t)$) was assumed to increase instantaneously at action potential (AP)

832   arrival and to decay back to its resting value $[Ca^{2+}]_{rest}$ with a rate constant k_{Ca} according to the

833   rate equation:

$$\frac{d}{dt} [Ca^{2+}](t) = -k_{Ca} \cdot ([Ca^{2+}](t) - [Ca^{2+}]_{rest})$$

834   Release probability (P_r) at arrival of the j^{th} AP was modelled according to

$$P_{r,j} = P_{r,1} \cdot y_j^{4.5} \cdot z_j$$

with y≥1 and z≤1. Here, $P_{r,1}$ designates the release probability for the first EPSC in a train, y_j

accounts for changes in local $[Ca^{2+}]$ during repetitive stimulation ($y_j = [Ca^{2+}]_j / [Ca^{2+}]_1$), likely

due to presynaptic Ca^{2+} current facilitation, and/or saturation of local Ca^{2+} buffers, and z

accounts for a small reduction of P_r during repetitive stimulation.

Both variables y_j and z_j were initialized to 1 at the onset of a stimulus train. The variable y was

incremented after each AP by

$$y_{inc} = y_{inc,1} \cdot (y_{max} - y_j)$$

841   and z was decremented by

$$z_{dec} = z_{dec,1} \cdot (z_j - z_{min})$$

842   During inter-stimulus intervals, time courses of $y(t)$ and $z(t)$ were determined by the rate

843   equations:

$$\frac{d}{dt} y(t) = (1 - y(t)) \cdot k_y$$

$$\frac{d}{dt} z(t) = (1 - z(t)) \cdot k_z$$

(for details see Lin et al. (2022)).

For each release event, the quantal content m_j of the EPSC$_j$ triggered by stimulus j was

calculated as the product of $P_{r,j} \cdot N_o(t_j)$ with both quantities evaluated immediately before

stimulus arrival. Between APs, the differential equations were solved numerically using the fifth-

order Runge-Kutta-Fehlberg algorithm implemented in IgorPro (Wavemetrics). All model

parameters on the time course of effective $[Ca^{2+}]$ and P_r during stimulus trains were constrained

to the same values for both genotypes. The remaining model parameters (N_{total}, initial P_r, k_f, k_b,

σ, $K_{0.5}$) were adjusted by trial and error to reproduce experimentally observed differences

between $Rac1^{+/+}$ and $Rac1^{-/-}$ synapses (Figure 6 figure supplement 1). The total number of SV

docking sites (N_{total}), priming and unpriming rate constants, Ca^{2+}-dependence of the priming

step, and P_r were free parameters and adjusted by trial and error to reproduce experimental data

on 1) the time course of fractional recovery after 500 Hz conditioning stimuli (Fig. 6A3), 2) initial

synaptic strength, and 3) the time course of STP during 50 and 500 Hz stimulation (Fig. 6A4).

Sequential two-step model

The sequential two-step model was implemented as previously described (Lin et al., 2022), with

the standard model parameters modified to reproduce experimentally observed STP and EPSC

recovery after conditioning trains in >P28 mouse calyx synapses and at physiological

temperature. All model parameters defining the time course of effective $[Ca^{2+}]$ and P_r during

stimulus trains were constrained to the same values for both genotypes. Except for initial P_r and

N_{total}, the remaining model parameters were adjusted by trial and error to reproduce

experimentally observed differences between *Rac1*$^{+/+}$ and *Rac1*$^{-/-}$ synapses. Thus, only those

parameters determining the SV state transitions (k_1, b_1, σ_1, $K_{0.5}$, k_2, b_2, σ_2) were allowed to differ

between *Rac1*$^{+/+}$ and *Rac1*$^{-/-}$ synapses while initial P_r and N_{total} identical for both genotypes

(Figure 6 - figure supplement 2).

Data and software accessibility

All numerical data used to generate the figures are part of the respective source files.

Experimental raw data and custom-written software central to the conclusion of this study are

available at http://dx.doi.org/10.17632/c4b7gn8bh7 under the Creative Commons Attribution 4.0

License (CC BY 4.0).

Acknowledgments

We would like to thank the members of the Young lab for their comments on the manuscript.

Competing interests

The authors declare no competing interests.

References
Alabi AA, Tsien RW (2012) Synaptic vesicle pools and dynamics. Cold Spring Harbor
perspectives in biology 4:a013680.
Armbruster M, Ryan TA (2011) Synaptic vesicle retrieval time is a cell-wide rather than
individual-synapse property. Nat Neurosci 14:824-826.
Azarfar A, Calcini N, Huang C, Zeldenrust F, Celikel T (2018) Neural coding: A single neuron's
perspective. Neurosci Biobehav Rev 94:238-247.
Bai Y, Xiang X, Liang C, Shi L (2015) Regulating Rac in the nervous system: molecular function
and disease implication of Rac GEFs and GAPs. Biomed Res Int 2015:632450.
Basu J, Betz A, Brose N, Rosenmund C (2007) Munc13-1 C1 domain activation lowers the
energy barrier for synaptic vesicle fusion. J Neurosci 27:1200-1210.
Bleckert A, Photowala H, Alford S (2012) Dual pools of actin at presynaptic terminals. Journal of
neurophysiology 107:3479-3492.
Bonnycastle K, Davenport EC, Cousin MA (2021) Presynaptic dysfunction in
neurodevelopmental disorders: Insights from the synaptic vesicle life cycle. J Neurochem
157:179-207.
Borst JG (2010) The low synaptic release probability in vivo. Trends in neurosciences 33:259-
266.
Borst JG, Sakmann B (1998) Calcium current during a single action potential in a large
presynaptic terminal of the rat brainstem. J Physiol 506:143-157.
Borst JG, Soria van Hoeve J (2012) The calyx of held synapse: from model synapse to auditory
relay. Annual review of physiology 74:199-224.
Bosco EE, Mulloy JC, Zheng Y (2009) Rac1 GTPase: a "Rac" of all trades. Cell Mol Life Sci
66:370-374.
Brette R (2015) Philosophy of the Spike: Rate-Based vs. Spike-Based Theories of the Brain.
Front Syst Neurosci 9:151.
Chanaday NL, Kavalali ET (2020) Is Ca(2+) Essential for Synaptic Vesicle Endocytosis? Trends
in neurosciences 43:77-79.
Chen Z, Cooper B, Kalla S, Varoqueaux F, Young SM, Jr. (2013) The Munc13 proteins
differentially regulate readily releasable pool dynamics and calcium-dependent recovery
at a central synapse. The Journal of neuroscience : the official journal of the Society for
Neuroscience 33:8336-8351.
Chen Z, Das B, Nakamura Y, DiGregorio DA, Young SM, Jr. (2015) Ca2+ channel to synaptic
vesicle distance accounts for the readily releasable pool kinetics at a functionally mature
auditory synapse. J Neurosci 35:2083-2100.
Cingolani LA, Goda Y (2008) Actin in action: the interplay between the actin cytoskeleton and
synaptic efficacy. Nat Rev Neurosci 9:344-356.
Cole JC, Villa BR, Wilkinson RS (2000) Disruption of actin impedes transmitter release in snake
motor terminals. The Journal of physiology 525 Pt 3:579-586.
Crins TT, Rusu SI, Rodriguez-Contreras A, Borst JG (2011) Developmental changes in short-
term plasticity at the rat calyx of Held synapse. The Journal of neuroscience : the official
journal of the Society for Neuroscience 31:11706-11717.
Dietz DM et al. (2012) Rac1 is essential in cocaine-induced structural plasticity of nucleus
accumbens neurons. Nat Neurosci 15:891-896.
Dong W, Radulovic T, Goral RO, Thomas C, Suarez Montesinos M, Guerrero-Given D,
Hagiwara A, Putzke T, Hida Y, Abe M, Sakimura K, Kamasawa N, Ohtsuka T, Young
SM, Jr. (2018) CAST/ELKS Proteins Control Voltage-Gated Ca(2+) Channel Density and
Synaptic Release Probability at a Mammalian Central Synapse. Cell Rep 24:284-293
e286.

Doussau F, Gasman S, Humeau Y, Vitiello F, Popoff M, Boquet P, Bader MF, Poulain B (2000)
A Rho-related GTPase is involved in Ca(2+)-dependent neurotransmitter exocytosis. J
Biol Chem 275:7764-7770.
Eggermann E, Bucurenciu I, Goswami SP, Jonas P (2012) Nanodomain coupling between
Ca(2)(+) channels and sensors of exocytosis at fast mammalian synapses. Nat Rev
Neurosci 13:7-21.
Eguchi K, Taoufiq Z, Thorn-Seshold O, Trauner D, Hasegawa M, Takahashi T (2017) Wild-Type
Monomeric alpha-Synuclein Can Impair Vesicle Endocytosis and Synaptic Fidelity via
Tubulin Polymerization at the Calyx of Held. J Neurosci 37:6043-6052.
Elmqvist D, Quastel DM (1965) A quantitative study of end-plate potentials in isolated human
muscle. The Journal of physiology 178:505-529.
Englitz B, Tolnai S, Typlt M, Jost J, Rubsamen R (2009) Reliability of synaptic transmission at
the synapses of Held in vivo under acoustic stimulation. PLoS One 4:e7014.
Fedchyshyn MJ, Wang LY (2005) Developmental transformation of the release modality at the
calyx of Held synapse. The Journal of neuroscience : the official journal of the Society for
Neuroscience 25:4131-4140.
Fedchyshyn MJ, Wang LY (2007) Activity-dependent changes in temporal components of
neurotransmission at the juvenile mouse calyx of Held synapse. The Journal of
physiology 581:581-602.
Forsberg M, Seth H, Bjorefeldt A, Lyckenvik T, Andersson M, Wasling P, Zetterberg H, Hanse E
(2019) Ionized calcium in human cerebrospinal fluid and its influence on intrinsic and
synaptic excitability of hippocampal pyramidal neurons in the rat. J Neurochem 149:452-
470.
Forsythe ID, Barnes-Davies M (1993) The binaural auditory pathway: excitatory amino acid
receptors mediate dual timecourse excitatory postsynaptic currents in the rat medial
nucleus of the trapezoid body. Proc Biol Sci 251:151-157.
Friauf E, Ostwald J (1988) Divergent projections of physiologically characterized rat ventral
cochlear nucleus neurons as shown by intra-axonal injection of horseradish peroxidase.
Experimental brain research Experimentelle Hirnforschung Experimentation cerebrale
73:263-284.
Friauf E, Fischer AU, Fuhr MF (2015) Synaptic plasticity in the auditory system: a review. Cell
Tissue Res 361:177-213.
Gentile JE, Carrizales MG, Koleske AJ (2022) Control of Synapse Structure and Function by
Actin and Its Regulators. Cells 11.
Ghanbari A, Ren N, Keine C, Stoelzel C, Englitz B, Swadlow HA, Stevenson IH (2020) Modeling
the Short-Term Dynamics of in Vivo Excitatory Spike Transmission. J Neurosci 40:4185-
4202.
Glogauer M, Marchal CC, Zhu F, Worku A, Clausen BE, Foerster I, Marks P, Downey GP,
Dinauer M, Kwiatkowski DJ (2003) Rac1 deletion in mouse neutrophils has selective
effects on neutrophil functions. J Immunol 170:5652-5657.
Grothe B, Pecka M, McAlpine D (2010) Mechanisms of sound localization in mammals.
Physiological reviews 90:983-1012.
Hallermann S, Silver RA (2013) Sustaining rapid vesicular release at active zones: potential
roles for vesicle tethering. Trends in neurosciences 36:185-194.
Haustein MD, Reinert T, Warnatsch A, Englitz B, Dietz B, Robitzki A, Rubsamen R, Milenkovic I
(2008) Synaptic transmission and short-term plasticity at the calyx of Held synapse
revealed by multielectrode array recordings. J Neurosci Methods 174:227-236.
Hentschke H, Stuttgen MC (2011) Computation of measures of effect size for neuroscience data
sets. The European journal of neuroscience 34:1887-1894.

Imig C, Min SW, Krinner S, Arancillo M, Rosenmund C, Sudhof TC, Rhee J, Brose N, Cooper
BH (2014) The morphological and molecular nature of synaptic vesicle priming at
presynaptic active zones. Neuron 84:416-431.
Iwasaki S, Takahashi T (1998) Developmental changes in calcium channel types mediating
synaptic transmission in rat auditory brainstem. The Journal of physiology 509 (Pt
2):419-423.
Iwasaki S, Takahashi T (2001) Developmental regulation of transmitter release at the calyx of
Held in rat auditory brainstem. The Journal of physiology 534:861-871.
Iwasaki S, Momiyama A, Uchitel OD, Takahashi T (2000) Developmental changes in calcium
channel types mediating central synaptic transmission. J Neurosci 20:59-65.
Jensen V, Walaas SI, Hilfiker S, Ruiz A, Hvalby O (2007) A delayed response enhancement
during hippocampal presynaptic plasticity in mice. The Journal of physiology 583:129-
143.
Johnston J, Forsythe ID, Kopp-Scheinpflug C (2010) Going native: voltage-gated potassium
channels controlling neuronal excitability. The Journal of physiology 588:3187-3200.
Joris PX, Trussell LO (2018) The Calyx of Held: A Hypothesis on the Need for Reliable Timing in
an Intensity-Difference Encoder. Neuron 100:534-549.
Joris PX, Schreiner CE, Rees A (2004) Neural processing of amplitude-modulated sounds.
Physiological reviews 84:541-577.
Jung JH, Szule JA, Marshall RM, McMahan UJ (2016) Variable priming of a docked synaptic
vesicle. Proc Natl Acad Sci U S A 113:E1098-1107.
Jung JH, Kirk LM, Bourne JN, Harris KM (2021) Shortened tethering filaments stabilize
presynaptic vesicles in support of elevated release probability during LTP in rat
hippocampus. Proc Natl Acad Sci U S A 118.
Kavalali ET (2020) Neuronal Ca(2+) signalling at rest and during spontaneous
neurotransmission. The Journal of physiology 598:1649-1654.
Keine C, Rubsamen R, Englitz B (2016) Inhibition in the auditory brainstem enhances signal
representation and regulates gain in complex acoustic environments. Elife 5.
Klug A, Borst JG, Carlson BA, Kopp-Scheinpflug C, Klyachko VA, Xu-Friedman MA (2012) How
do short-term changes at synapses fine-tune information processing? J Neurosci
32:14058-14063.
Kochubey O, Han Y, Schneggenburger R (2009) Developmental regulation of the intracellular
Ca^{2+} sensitivity of vesicle fusion and Ca^{2+}-secretion coupling at the rat calyx of Held. J
Physiol 587:3009-3023.
Kumanogoh H, Miyata S, Sokawa Y, Maekawa S (2001) Biochemical and morphological
analysis on the localization of Rac1 in neurons. Neurosci Res 39:189-196.
Lee JS, Ho WK, Lee SH (2012) Actin-dependent rapid recruitment of reluctant synaptic vesicles
into a fast-releasing vesicle pool. Proceedings of the National Academy of Sciences of
the United States of America 109:E765-774.
Lee JS, Ho WK, Neher E, Lee SH (2013) Superpriming of synaptic vesicles after their
recruitment to the readily releasable pool. Proceedings of the National Academy of
Sciences of the United States of America 110:15079-15084.
Li L, Bischofberger J, Jonas P (2007) Differential gating and recruitment of P/Q-, N-, and R-type
Ca^{2+} channels in hippocampal mossy fiber boutons. J Neurosci 27:13420-13429.
Lidierth M (2009) sigTOOL: A MATLAB-based environment for sharing laboratory-developed
software to analyze biological signals. J Neurosci Methods 178:188-196.
Lin KH, Taschenberger H, Neher E (2022) A sequential two-step priming scheme reproduces
diversity in synaptic strength and short-term plasticity. Proc Natl Acad Sci U S A
119:e2207987119.
Lipstein N, Chang S, Lin KH, Lopez-Murcia FJ, Neher E, Taschenberger H, Brose N (2021)
Munc13-1 is a Ca^{2+}-phospholipid-dependent vesicle priming hub that shapes synaptic

short-term plasticity and enables sustained neurotransmission. Neuron 109:3980-4000
e3987.
Lipstein N, Sakaba T, Cooper BH, Lin KH, Strenzke N, Ashery U, Rhee JS, Taschenberger H,
Neher E, Brose N (2013) Dynamic control of synaptic vesicle replenishment and short-
term plasticity by Ca^{2+}-calmodulin-Munc13-1 signaling. Neuron 79:82-96.
Lopez-Murcia FJ, Reim K, Jahn O, Taschenberger H, Brose N (2019) Acute Complexin
Knockout Abates Spontaneous and Evoked Transmitter Release. Cell Rep 26:2521-2530
e2525.
Lorteije JA, Borst JG (2011) Contribution of the mouse calyx of Held synapse to tone adaptation.
The European journal of neuroscience 33:251-258.
Lorteije JA, Rusu SI, Kushmerick C, Borst JG (2009) Reliability and precision of the mouse calyx
of Held synapse. The Journal of neuroscience : the official journal of the Society for
Neuroscience 29:13770-13784.
Lubbert M, Goral RO, Satterfield R, Putzke T, van den Maagdenberg AM, Kamasawa N, Young
SM, Jr. (2017) A novel region in the CaV2.1 alpha1 subunit C-terminus regulates fast
synaptic vesicle fusion and vesicle docking at the mammalian presynaptic active zone.
Elife 6.
Lubbert M, Goral RO, Keine C, Thomas C, Guerrero-Given D, Putzke T, Satterfield R,
Kamasawa N, Young SM, Jr. (2019) CaV2.1 alpha1 Subunit Expression Regulates
Presynaptic CaV2.1 Abundance and Synaptic Strength at a Central Synapse. Neuron
101:260-273 e266.
Mc Laughlin M, van der Heijden M, Joris PX (2008) How secure is in vivo synaptic transmission
at the calyx of Held? The Journal of neuroscience : the official journal of the Society for
Neuroscience 28:10206-10219.
Miki T, Malagon G, Pulido C, Llano I, Neher E, Marty A (2016) Actin- and Myosin-Dependent
Vesicle Loading of Presynaptic Docking Sites Prior to Exocytosis. Neuron 91:808-823.
Montesinos MS, Satterfield R, Young SM, Jr. (2016) Helper-Dependent Adenoviral Vectors and
Their Use for Neuroscience Applications. Methods Mol Biol 1474:73-90.
Montesinos MS, Dong W, Goff K, Das B, Guerrero-Given D, Schmalzigaug R, Premont RT,
Satterfield R, Kamasawa N, Young SM, Jr. (2015) Presynaptic Deletion of GIT Proteins
Results in Increased Synaptic Strength at a Mammalian Central Synapse. Neuron
88:918-925.
Morales M, Colicos MA, Goda Y (2000) Actin-dependent regulation of neurotransmitter release
at central synapses. Neuron 27:539-550.
Nakamura Y, Harada H, Kamasawa N, Matsui K, Rothman JS, Shigemoto R, Silver RA,
DiGregorio DA, Takahashi T (2015) Nanoscale distribution of presynaptic Ca(2+)
channels and its impact on vesicular release during development. Neuron 85:145-158.
Neher E (2010) What is Rate-Limiting during Sustained Synaptic Activity: Vesicle Supply or the
Availability of Release Sites. Frontiers in synaptic neuroscience 2:144.
Neher E (2015) Merits and Limitations of Vesicle Pool Models in View of Heterogeneous
Populations of Synaptic Vesicles. Neuron 87:1131-1142.
Neher E, Sakaba T (2008) Multiple roles of calcium ions in the regulation of neurotransmitter
release. Neuron 59:861-872.
Neher E, Brose N (2018) Dynamically Primed Synaptic Vesicle States: Key to Understand
Synaptic Short-Term Plasticity. Neuron 100:1283-1291.
Neher E, Taschenberger H (2021) Non-negative Matrix Factorization as a Tool to Distinguish
Between Synaptic Vesicles in Different Functional States. Neuroscience 458:182-202.
O'Neil SD, Racz B, Brown WE, Gao Y, Soderblom EJ, Yasuda R, Soderling SH (2021) Action
potential-coupled Rho GTPase signaling drives presynaptic plasticity. Elife 10.

Piriya Ananda Babu L, Wang HY, Eguchi K, Guillaud L, Takahashi T (2020) Microtubule and
1077   Actin Differentially Regulate Synaptic Vesicle Cycling to Maintain High-Frequency
1078   Neurotransmission. J Neurosci 40:131-142.
1079 Pulido C, Marty A (2018) A two-step docking site model predicting different short-term synaptic
1080   plasticity patterns. J Gen Physiol 150:1107-1124.
1081 Radulovic T, Dong W, Goral RO, Thomas CI, Veeraraghavan P, Montesinos MS, Guerrero-
1082   Given D, Goff K, Lubbert M, Kamasawa N, Ohtsuka T, Young SM, Jr. (2020) Presynaptic
1083   development is controlled by the core active zone proteins CAST/ELKS. The Journal of
1084   physiology 598:2431-2452.
1085 Reijnders MRF et al. (2017) RAC1 Missense Mutations in Developmental Disorders with Diverse
1086   Phenotypes. Am J Hum Genet 101:466-477.
1087 Reinagel P, Laughlin S (2001) Natural stimulus statistics. Network 12:237-240.
1088 Ritzau-Jost A, Jablonski L, Viotti J, Lipstein N, Eilers J, Hallermann S (2018) Apparent calcium
1089   dependence of vesicle recruitment. The Journal of physiology 596:4693-4707.
1090 Rollenhagen A, Lubke JH (2006) The morphology of excitatory central synapses: from structure
1091   to function. Cell Tissue Res 326:221-237.
1092 Rothman JS, Silver RA (2018) NeuroMatic: An Integrated Open-Source Software Toolkit for
1093   Acquisition, Analysis and Simulation of Electrophysiological Data. Front Neuroinform
1094   12:14.
1095 Rust MB, Maritzen T (2015) Relevance of presynaptic actin dynamics for synapse function and
1096   mouse behavior. Experimental cell research.
1097 Sakaba T (2006) Roles of the fast-releasing and the slowly releasing vesicles in synaptic
1098   transmission at the calyx of Held. J Neurosci 26:5863-5871.
1099 Sakaba T, Neher E (2003) Involvement of actin polymerization in vesicle recruitment at the calyx
1100   of Held synapse. The Journal of neuroscience : the official journal of the Society for
1101   Neuroscience 23:837-846.
1102 Sakaba T, Kononenko NL, Bacetic J, Pechstein A, Schmoranzer J, Yao L, Barth H, Shupliakov
1103   O, Kobler O, Aktories K, Haucke V (2013) Fast neurotransmitter release regulated by the
1104   endocytic scaffold intersectin. Proc Natl Acad Sci U S A 110:8266-8271.
1105 Satzler K, Sohl LF, Bollmann JH, Borst JG, Frotscher M, Sakmann B, Lubke JH (2002) Three-
1106   dimensional reconstruction of a calyx of Held and its postsynaptic principal neuron in the
1107   medial nucleus of the trapezoid body. The Journal of neuroscience : the official journal of
1108   the Society for Neuroscience 22:10567-10579.
1109 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
1110   C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P,
1111   Cardona A (2012) Fiji: an open-source platform for biological-image analysis. Nature
1112   methods 9:676-682.
1113 Schneggenburger R, Rosenmund C (2015) Molecular mechanisms governing Ca(2+) regulation
1114   of evoked and spontaneous release. Nat Neurosci 18:935-941.
1115 Schotten S, Meijer M, Walter AM, Huson V, Mamer L, Kalogreades L, ter Veer M, Ruiter M,
1116   Brose N, Rosenmund C, Sorensen JB, Verhage M, Cornelisse LN (2015) Additive effects
1117   on the energy barrier for synaptic vesicle fusion cause supralinear effects on the vesicle
1118   fusion rate. Elife 4:e05531.
1119 Shimshek DR, Kim J, Hubner MR, Spergel DJ, Buchholz F, Casanova E, Stewart AF, Seeburg
1120   PH, Sprengel R (2002) Codon-improved Cre recombinase (iCre) expression in the
1121   mouse. Genesis 32:19-26.
1122 Sonntag M, Englitz B, Kopp-Scheinpflug C, Rubsamen R (2009) Early postnatal development of
1123   spontaneous and acoustically evoked discharge activity of principal cells of the medial
1124   nucleus of the trapezoid body: an in vivo study in mice. The Journal of neuroscience : the
1125   official journal of the Society for Neuroscience 29:9510-9520.

Sonntag M, Englitz B, Typlt M, Rubsamen R (2011) The calyx of held develops adult-like
dynamics and reliability by hearing onset in the mouse in vivo. The Journal of
neuroscience : the official journal of the Society for Neuroscience 31:6699-6709.
Spirou GA, Brownell WE, Zidanic M (1990) Recordings from cat trapezoid body and HRP
labeling of globular bushy cell axons. Journal of neurophysiology 63:1169-1190.
Stanley EF (2016) The Nanophysiology of Fast Transmitter Release. Trends in neurosciences
39:183-197.
Sun Y, Bamji SX (2011) beta-Pix modulates actin-mediated recruitment of synaptic vesicles to
synapses. The Journal of neuroscience : the official journal of the Society for
Neuroscience 31:17123-17133.
Taschenberger H, Leao RM, Rowland KC, Spirou GA, von Gersdorff H (2002) Optimizing
synaptic architecture and efficiency for high-frequency transmission. Neuron 36:1127-
1143.
Thanawala MS, Regehr WG (2016) Determining synaptic parameters using high-frequency
activation. J Neurosci Methods 264:136-152.
Theunissen FE, Elie JE (2014) Neural processing of natural sounds. Nat Rev Neurosci 15:355-
366.
Thomas CI, Keine C, Okayama S, Satterfield R, Musgrove M, Guerrero-Given D, Kamasawa N,
Young SM, Jr. (2019) Presynaptic Mitochondria Volume and Abundance Increase during
Development of a High-Fidelity Synapse. J Neurosci 39:7994-8012.
Threadgill R, Bobb K, Ghosh A (1997) Regulation of dendritic growth and remodeling by Rho,
Rac, and Cdc42. Neuron 19:625-634.
Tolnai S, Hernandez O, Englitz B, Rubsamen R, Malmierca MS (2008) The medial nucleus of
the trapezoid body in rat: spectral and temporal properties vary with anatomical location
of the units. The European journal of neuroscience 27:2587-2598.
Torres VI, Vallejo D, Inestrosa NC (2017) Emerging Synaptic Molecules as Candidates in the
Etiology of Neurological Disorders. Neural Plast 2017:8081758.
Traynelis SF (1998) Software-based correction of single compartment series resistance errors. J
Neurosci Methods 86:25-34.
Wadel K, Neher E, Sakaba T (2007) The coupling between synaptic vesicles and Ca2+
channels determines fast neurotransmitter release. Neuron 53:563-575.
Waites CL, Leal-Ortiz SA, Andlauer TF, Sigrist SJ, Garner CC (2011) Piccolo regulates the
dynamic assembly of presynaptic F-actin. The Journal of neuroscience : the official
journal of the Society for Neuroscience 31:14250-14263.
Wang LY, Kaczmarek LK (1998) High-frequency firing helps replenish the readily releasable
pool of synaptic vesicles. Nature 394:384-388.
Wang LY, Neher E, Taschenberger H (2008) Synaptic vesicles in mature calyx of Held synapses
sense higher nanodomain calcium concentrations during action potential-evoked
glutamate release. J Neurosci 28:14450-14458.
Wang LY, Gan L, Forsythe ID, Kaczmarek LK (1998) Contribution of the Kv3.1 potassium
channel to high-frequency firing in mouse auditory neurones. The Journal of physiology
509 (Pt 1):183-194.
Watanabe S, Liu Q, Davis MW, Hollopeter G, Thomas N, Jorgensen NB, Jorgensen EM (2013a)
Ultrafast endocytosis at Caenorhabditis elegans neuromuscular junctions. Elife
2:e00723.
Watanabe S, Rost BR, Camacho-Perez M, Davis MW, Sohl-Kielczynski B, Rosenmund C,
Jorgensen EM (2013b) Ultrafast endocytosis at mouse hippocampal synapses. Nature
504:242-247.
Weis S, Schneggenburger R, Neher E (1999) Properties of a model of Ca^{++}-dependent vesicle
pool dynamics and short term synaptic depression. Biophys J 77:2418-2429.

Wen H, Hubbard JM, Rakela B, Linhoff MW, Mandel G, Brehm P (2013) Synchronous and
asynchronous modes of synaptic transmission utilize different calcium sources. Elife
2:e01206.
Wimmer VC, Horstmann H, Groh A, Kuner T (2006) Donut-like topology of synaptic vesicles with
a central cluster of mitochondria wrapped into membrane protrusions: a novel structure-
function module of the adult calyx of Held. The Journal of neuroscience : the official
journal of the Society for Neuroscience 26:109-116.
Wondolowski J, Dickman D (2013) Emerging links between homeostatic synaptic plasticity and
neurological disease. Front Cell Neurosci 7:223.
Wright WJ, Graziane NM, Neumann PA, Hamilton PJ, Cates HM, Fuerst L, Spenceley A,
MacKinnon-Booth N, Iyer K, Huang YH, Shaham Y, Schluter OM, Nestler EJ, Dong Y
(2020) Silent synapses dictate cocaine memory destabilization and reconsolidation. Nat
Neurosci 23:32-46.
Wu LG, Chan CY (2022) Multiple Roles of Actin in Exo- and Endocytosis. Frontiers in synaptic
neuroscience 14:841704.
Xu Z, Chen Y, Chen Y (2019) Spatiotemporal Regulation of Rho GTPases in Neuronal
Migration. Cells 8.
Xue M, Craig TK, Xu J, Chao HT, Rizo J, Rosenmund C (2010) Binding of the complexin N
terminus to the SNARE complex potentiates synaptic-vesicle fusogenicity. Nat Struct Mol
Biol 17:568-575.
Yang CH, Ho WK, Lee SH (2021) Postnatal maturation of glutamate clearance and release
kinetics at the rat and mouse calyx of Held synapses. Synapse 75:e22215.
Yang H, Xu-Friedman MA (2015) Skipped-stimulus approach reveals that short-term plasticity
dominates synaptic strength during ongoing activity. J Neurosci 35:8297-8307.
Yang YM, Fedchyshyn MJ, Grande G, Aitoubah J, Tsang CW, Xie H, Ackerley CA, Trimble WS,
Wang LY (2010) Septins regulate developmental switching from microdomain to
nanodomain coupling of Ca(2+) influx to neurotransmitter release at a central synapse.
Neuron 67:100-115.
Yasuda R (2017) Biophysics of Biochemical Signaling in Dendritic Spines: Implications in
Synaptic Plasticity. Biophys J 113:2152-2159.
Young SM, Jr., Neher E (2009) Synaptotagmin has an essential function in synaptic vesicle
positioning for synchronous release in addition to its role as a calcium sensor. Neuron
63:482-496.
Zamboni V, Jones R, Umbach A, Ammoni A, Passafaro M, Hirsch E, Merlo GR (2018) Rho
GTPases in Intellectual Disability: From Genetics to Therapeutic Opportunities. Int J Mol
Sci 19.
Zhang H, Ben Zablah Y, Zhang H, Jia Z (2021) Rho Signaling in Synaptic Plasticity, Memory,
and Brain Disorders. Front Cell Dev Biol 9:729076.

1214

1215

 Figures legends

 Fig. 1: Loss of presynaptic Rac1 after hearing onset does not affect calyx of Held gross

 morphology or ultrastructure. (**A**) Cre recombinase-expressing HdAds were injected into the

 cochlear nucleus of $Rac1^{fl/fl}$ mice at P14, yielding $Rac1^{-/-}$ calyces of Held. All experiments were

 performed at around four weeks of age. Cre-recombinase-expressing calyces could be visually

 identified by simultaneous expression of EGFP. (**B1**) Representative reconstruction of calyx

 terminals of $Rac1^{+/+}$ (left) and $Rac1^{-/-}$ (right) animals. (**B2**) Calyx morphology assessed by

 surface area (left) and volume (right) was not affected by the loss of Rac1. (**C1**) Representative

 EM images of the active zone (yellow) and docked SV (blue) to assess synaptic ultrastructure.

 (**C2**) AZ length and number of docked SV were comparable between $Rac1^{+/+}$ and $Rac1^{-/-}$. (**C3**)

 SV distribution as a function of distance to AZ was not different between $Rac1^{+/+}$ and $Rac1^{-/-}$.

 Box plot whiskers extend to the minimum/maximum within the 1.5 interquartile range; open

 markers indicate individual data points. For EM data, the results of three independent

 investigators were averaged. All data shown in the figure and the detailed results of statistical

 tests are part of the supplementary file.

 Figure 1 – source data 1: Excel file containing the data shown in Figure 1 and the results of

 statistical analysis.

 Fig. 2: Presynaptic Rac1 regulates synaptic vesicles release probability and synaptic

 strength. Synaptic transmission at the calyx of Held – MNTB synapse was studied at P28 after

 deletion of Rac1 at P14 at different stimulation frequencies. (**A1, B1**) Representative evoked

 EPSCs for $Rac1^{+/+}$ (black) and $Rac1^{-/-}$ (orange) at 50 Hz and 500 Hz stimulation frequency.

 Stimulus artifacts were blanked for clarity. (**C**) Magnification of the first EPSC. Ablation of

 presynaptic Rac1 resulted in increased EPSC amplitude with no change in EPSC dynamics.

 (**A2-A4**) At 50 Hz stimulation frequency, $Rac1^{-/-}$ showed stronger short-term depression and

 larger steady-state EPSC amplitudes. (**B2-B4**) At 500 Hz stimulation frequency, loss of Rac1

resulted in a lack of short-term facilitation and increased synaptic depression with no change in

steady-state EPSC amplitude. (**D1**) Population data showing an increase in first EPSC amplitude

in $Rac1^{-/-}$. Steady-state EPSC amplitudes were increased in $Rac1^{-/-}$ at 50 Hz but not at 500 Hz

stimulation frequency. (**D2**) Population data of RRP using three different estimation methods,

suggesting little to no change in RRP size (**D3**) Population data indicating that release probability

in $Rac1^{-/-}$ was elevated independent of estimation method. All data shown in the figure and the

detailed results of statistical tests are part of the supplementary file.

Figure 2 – source data 1: Excel file containing the data shown in Figure 2 and the results of

statistical analysis.

1251

Fig. 3: Presynaptic loss of Rac1 increase calcium-independent neurotransmitter release.

(**A**) Representative recordings of mEPSCs for $Rac1^{+/+}$ (left, black) and $Rac1^{-/-}$ (right, orange).

(**B1-B4**) Rac1 deletion increased mEPSC frequency but did not affect mEPSC amplitude, rise

time, or full width at half-maximal amplitude (FWHM). (**C**) The increased mEPSC rates at

$Rac1^{-/-}$ were independent of presynaptic voltage-gated calcium channels (VGCC), as blocking

VGCC with cadmium (Cd^{2+}) did not affect mEPSC frequency. All data shown in the figure and

the detailed results of statistical tests are part of the supplementary file.

Figure 3 – source data 1: Excel file containing the data shown in Figure 3 and the results of

statistical analysis.

1261

Fig. 4: Presynaptic loss of Rac1 decreases SV synchronicity and prolongs EPSC onset at

high-frequency stimulation. (**A**) Experiments were performed at low (50 Hz, A1) and high

(500 Hz, A2) stimulation frequencies. Representative recordings of first ($EPSC_1$) and last

($EPSC_{50}$) EPSC in the stimulus train. Traces are aligned at the EPSC onset of the first EPSC.

Stimulus artifacts are partially blanked for better visibility. Note the shift in the onset of $EPSC_{50}$ in

$Rac1^{-/-}$ compared to $Rac1^{+/+}$ at 500 Hz but not 50 Hz. (**B1**) Absolute EPSC onset delay for

50 Hz (gray and light orange) and 500 Hz (black and orange) stimulation. (**B2**) EPSC onset

delay relative to $EPSC_1$ for 50 Hz (gray and light orange) and 500 Hz (black and orange). At

50 Hz, the EPSC onset delay was similar between $Rac1^{+/+}$ and $Rac1^{-/-}$. At 500 Hz, the EPSC

onset delay was substantially larger at $Rac1^{-/-}$. For better visualization, only every second data

point is shown. (**B3**) EPSC onset delay of the last ten EPSCs relative to $EPSC_1$ for 50 Hz and

500 Hz stimulation. EPSC delay of the last ten EPSC was not different between $Rac1^{+/+}$ and

$Rac1^{-/-}$ at 50 Hz but increased for $Rac1^{-/-}$ at 500 Hz stimulation frequency. (**C1**) Analysis of

'effective EPSC duration' to estimate SV release synchronicity during 50 Hz and 500 Hz

stimulation. Synchronicity was estimated from 'effective EPSC duration' by dividing the EPSC

charge by the EPSC amplitude. Note the increase in effective EPSC duration for $Rac1^{-/-}$ at

500 Hz stimulation. For better visualization, only every second data point is shown (**C2**) EPSC

duration was not different for $EPSC_1$ but slightly longer for late EPSCs in $Rac1^{-/-}$ at 50 Hz and

substantially longer at 500 Hz stimulation frequency. (**C3**) EPSC duration of the last ten EPSCs

normalized to $EPSC_1$. Note the progressive increase in EPSC duration in $Rac1^{-/-}$ with increasing

stimulation frequency. All data shown in the figure and the detailed results of statistical tests are

part of the supplementary file.

Figure 4 – source data 1: Excel file containing the data shown in Figure 4 and the results of

statistical analysis.

1286

Fig. 5: Loss of presynaptic Rac1 facilitates synaptic vesicle recovery. Recovery of single

EPSC and RRP recovery was measured by two consecutive train stimuli (conditioning stimulus

and recovery stimulus) at 500 Hz at varying recovery intervals. (**A**) Single EPSC recovery. (**A1**)

Representative traces for $Rac1^{+/+}$ (black) and $Rac1^{-/-}$ (orange) for recovery intervals ranging

from 20 ms to 16 s. (**A2**) Recovery of absolute EPSC amplitudes as a function of recovery

interval with a magnification of short intervals (right). (**A3**) Fractional EPSC recovery as a

function of recovery interval. (**B**) Recovery of the RRP (**B1**) Representative recovery traces

following a 100 ms and 2 s recovery interval. The depleting stimulus train is the same as in A1.

(**B2**) Fractional RRP recovery was faster in $Rac1^{-/-}$ compared to $Rac1^{+/+}$. (**C**) Recovery of the

paired-pulse ratio (PPR) of the first two EPSCs of the recovery train. PPR was consistently lower

in $Rac1^{-/-}$, but the difference was emphasized at longer recovery intervals. All data shown in the

figure and the detailed results of statistical tests are part of the supplementary file.

Figure 5 – source data 1: Excel file containing the data shown in Figure 5 and the results of

statistical analysis.

1301

Fig. 6: Numerical simulations of 50 and 500 Hz STP and EPSC recovery after conditioning

500 Hz trains are consistent with *Rac1*-loss induced changes in SV priming.

Experimental observations were equally well reproduced by either of two kinetic schemes of SV

priming and fusion: a single pool model (**A**) or a recently proposed (Lin et al., 2022) sequential

two-step SV priming scheme (**B**). (**A1**) Diagram of vesicle states for the single pool model. SVs

reversibly dock at empty release sites (ES). SVs in the docked and primed state (DS) undergo

fusion with the probability P_r upon AP arrival. Vacated sites become immediately available for

SV docking and priming. Transitions represented by dashed lines occur instantaneously, while

those represented by solid lines occur with rate constants as indicated. Forward transition rate

constant is Ca^{2+}-dependent. For the single pool model, P_r, and total number of sites (N_{total}) were

free parameters for both genotypes. The model predicts an increased P_r from 0.08 ($Rac1^{+/+}$) to

0.165 ($Rac1^{-/-}$) and an increase in the number of docked SVs (RRP) from 2150 SVs ($Rac1^{+/+}$) to

2532 SVs ($Rac1^{-/-}$). (**A2**) Dependence of k_f on cytosolic $[Ca^{2+}]$ (effective $[Ca^{2+}]$) for $Rac1^{+/+}$

(black) and $Rac1^{-/-}$ (orange) synapses. The inset illustrates the time course of the effective

$[Ca^{2+}]$ during a 500 Hz train consisting of 40 stimuli. (**A3**) Predictions of the single pool model

(lines) for the time course of the fractional recovery of $EPSC_{test}$ after 500 Hz conditioning trains

superimposed onto experimental data for $Rac1^{+/+}$ (black circles) and $Rac1^{-/-}$ (orange triangles)

synapses (data from Fig. 5A3). (**A4**) Predictions of the single pool model (lines) for the time

course of STP during 50 Hz and 500 Hz trains superimposed onto experimental data (circles) for

Rac1$^{+/+}$ (black, top panel) and *Rac1*$^{-/-}$ (orange, bottom panel) synapses. (**A5**) Model predictions

for the time course of STP during 50 Hz and 500 Hz trains for *Rac1*$^{+/+}$ (gray and black) and

Rac1$^{-/-}$ (light and dark orange) synapses shown superimposed to facilitate comparison. (**B1**)

Diagram of vesicle states for the sequential two-step priming scheme. SVs reversibly dock at

empty release sites (ES) and become fusion-competent by undergoing a sequence of two

priming steps. After initial docking, SVs in the loosely docked state (LS) reversibly transition to

the tightly docked state (TS) from which they undergo fusion upon AP arrival with the probability

P_r. Vacated sites become immediately available for SV docking and priming. Transitions

represented by dashed lines occur instantaneously, while those represented by solid lines occur

with rate constants as indicated. Forward transition rate constants are Ca^{2+}-dependent. For the

sequential two-step priming scheme, P_r, and total number of sites (N_{total}) were constrained to the

same values for *Rac1*$^{+/+}$ and *Rac1*$^{-/-}$ synapses and only parameters determining the kinetics of

the two priming steps were allowed to differ between genotypes. (**B2**) Dependence of k_1 (top

panel) and k_2 (bottom panel) on cytosolic $[Ca^{2+}]$ for *Rac1*$^{+/+}$ (black) and *Rac1*$^{-/-}$ (orange)

synapses. The inset illustrates the time course of the effective $[Ca^{2+}]$ during a 500 Hz train

consisting of 40 stimuli. (**B3**) Predictions of the sequential two-step model (lines) for the time

course of the fractional recovery of EPSC$_{test}$ after 500 Hz conditioning trains superimposed onto

experimental data for *Rac1*$^{+/+}$ (black circles) and *Rac1*$^{-/-}$ (orange triangles) synapses (data from

Fig. 5A3). (**B4**) Predictions of the sequential two-step model (lines) for the time course of STP

during 50 Hz and 500 Hz trains superimposed onto experimental data (circles) for *Rac1*$^{+/+}$

(black, top panel) and *Rac1*$^{-/-}$ (orange, bottom panel) synapses. (**B5**) Model predictions for the

time course of STP during 50 Hz and 500 Hz trains for *Rac1*$^{+/+}$ (gray and black) and *Rac1*$^{-/-}$

(light and dark orange) synapses shown superimposed to facilitate comparison.

1344

Fig. 6 – Figure supplement 1: Model predictions and parameters of simple single pool

models fitted to *Rac1*$^{+/+}$ and *Rac1*$^{-/-}$ STP and recovery data sets.

(**A**) Model predictions for synaptic STP during regular stimulus trains consisting of 40 APs and

delivered at frequencies from 0.5 to 500 Hz for $Rac1^{+/+}$ (**A1**, black) and $Rac1^{-/-}$ (**A2**, orange)

synapses. Note the strongly enhanced initial synaptic strength but similar steady-state release at

the highest stimulus frequency. (**B**) Model predictions for the paired-pulse ratio

(PPR = $EPSC_2$/$EPSC_1$) for $Rac1^{+/+}$ (black circles) and $Rac1^{-/-}$ (orange triangles) synapses.

While $Rac1^{+/+}$ synapses show substantial net facilitation at ISIs of 2 and 5 ms, net facilitation is

absent from for $Rac1^{-/-}$ synapses. (**D**) Model parameters for $Rac1^{+/+}$ (**D1**, black) and $Rac1^{-/-}$

(**D2**, orange) synapses. Parameter values that differ between the models for $Rac1^{+/+}$ and $Rac1^{-/-}$

synapses are listed in bold. For details, please refer to Lin et al. (2022). For converting EPSC

amplitudes measured in the presence of 1 mM kynurenic acid into quantal content, a quantal

size of $q = 7.48$ pA was assumed.

1358

Fig. 6 – Figure supplement 2: Model predictions and parameters of sequential two-step

models fitted to $Rac1^{+/+}$ and $Rac1^{-/-}$ STP and recovery data sets.

(**A**) Model predictions for synaptic STP during regular stimulus trains consisting of 40 APs and

delivered at frequencies from 0.5 to 500 Hz for $Rac1^{+/+}$ (**A1**, black) and $Rac1^{-/-}$ (**A2**, orange)

synapses. Note the strongly enhanced initial synaptic strength but similar steady-state release at

the highest stimulus frequency. (**B**) Model predictions for the paired-pulse ratio

(PPR = $EPSC_2$/$EPSC_1$) for $Rac1^{+/+}$ (black circles) and $Rac1^{-/-}$ (orange triangles) synapses.

While $Rac1^{+/+}$ synapses show substantial net facilitation at ISIs of 2 and 5 ms, net facilitation is

absent from for $Rac1^{-/-}$ synapses. (**D**) Model parameters for $Rac1^{+/+}$ (**D1**, black) and $Rac1^{-/-}$

(**D2**, orange) synapses. Parameter values that differ between the models for $Rac1^{+/+}$ and $Rac1^{-/-}$

synapses are listed in bold. For details, please refer to Lin et al. (2022). For converting EPSC

amplitudes measured in the presence of 1 mM kynurenic acid into quantal content, a quantal

size of $q = 7.48$ pA was assumed.

1372

Fig. 7: Alterations in presynaptic release probability did not impair the reliability of action

potential generation during *in vivo*-like activity. (**A1**) AP firing was recorded in response to *in*

vivo-like stimulation patterns derived from responses to sinusoidal amplitude-modulated sounds

at different modulation frequencies. Raster plot shows three representative stimulation patterns.

(**A2**) Representative traces of loose-patch recordings during afferent fiber stimulation with in

vivo-like activity for $Rac1^{+/+}$ (black) and $Rac1^{-/-}$ (orange). Triangles indicate the stimulus time

points. Stimulus artifacts were blanked for clarity. (**B1**) Fraction of successful AP generation was

not different between $Rac1^{+/+}$ and $Rac1^{-/-}$ independent of modulation frequency (**B2**) AP jitter

defined as the standard deviation of AP latencies was not changed in $Rac1^{-/-}$ neither for the

complete stimulus nor as a function of amplitude modulation frequency. (**C1**) AP jitter and AP

delay were largely independent of preceding inter-spike interval. (**C2**) AP jitter and AP delay as a

function of preceding activity level. The preceding activity was calculated as the sum of all

preceding APs (green, S_{-1}, S_{-2}, …, S_{-n}) weighted by their temporal distance to the AP under

observation (purple, W_{-1}, W_{-2}, …, W_{-n}). The weighting was implemented as an exponentially

decaying kernel (blue shaded area). Note the increased AP jitter and AP delay in $Rac1^{-/-}$ at

higher activity levels. All data shown in the figure and the detailed results of statistical tests are

part of the supplementary file.

Figure 7 – source data 1: Excel file containing the data shown in Figure 7 and the results of

statistical analysis.

1392

1393 **Fig. 8: Proposed model of Rac1's presynaptic role in regulating synaptic transmission.** In

1394 the proposed model, loss of Rac1 results in changes in F-actin at the active zone, thereby

1395 reducing the physical barrier between SVs and the plasma membrane resulting in increased

1396 synaptic strength through faster priming and potentially higher P_r.

1397

1398

A1

Cre

GBC

cochlear
nucleus

calyx of Held

MNTB

A2

Ψ | EGFP | hSyn | Cre-recombinase | hSyn

A3

hearing
onset

P12 **P14** P30

HdAd
injection

experiments

B1 *3D reconstruction*

Rac1^{+/+}   *Rac1^{−/−}*

B2

Calyx surface area (μm²)   p=0.22

(50)   (46)

Rac1^{+/+}   *Rac1^{−/−}*

Calyx volume (μm³)   p=0.42

Rac1^{+/+}   *Rac1^{−/−}*

C1 *Electron microscopy*

Rac1^{+/+}   *Rac1^{−/−}*

100 nm   100 nm

C2

AZ length (nm)   p=0.42

(121)   (123)

Rac1^{+/+}   *Rac1^{−/−}*

docked SV/AZ   p=0.27

Rac1^{+/+}   *Rac1^{−/−}*

C3

SV/bin

Distance from AZ (nm)

Rac1^{+/+}   *Rac1^{−/−}*

A1 *first EPSC recovery*

Rac1⁺/⁺

1 nA
20 ms
20 ms
16 s
0.5 ms

Rac1⁻/⁻

20 ms
16 s

B1 *RRP recovery*

100 ms   2 s

A2

Rac1⁺/⁺ (τ_w=1.8 s)   Rac1⁻/⁻ (τ_w=1.6 s)

Recovery EPSC_test (nA)
Recovery interval (s)
p=1.0e-12

B2

Fractional recovery RRP
Rac1⁺/⁺ (τ_w=1.1 s)   Rac1⁻/⁻ (τ_w=0.68 s)
Recovery interval (s)
p=8.2e-09

A3

Fractional recovery EPSC_test
Rac1⁺/⁺ (τ_w=1.9 s)   Rac1⁻/⁻ (τ_w=1.3 s)
Recovery interval (s)
p=2.6e-07

C *Paired-pulse ratio*

Rac1⁺/⁺   Rac1⁻/⁻
Paired-pulse ratio
50 ms   100 ms   1 s   4 s   8 s
Recovery interval

Rac1$^{+/+}$ *Rac1*$^{-/-}$

pre

post

pre

post

🔶 Ca$_v$2.1   🕸 F-actin   ✦ SNARE complex