Redox-based sensing controls bacterial chemotaxis to copper

Gwennaëlle Louis[1], Pauline Cherry[1], Catherine Michaux[2], Sophie Rahuel-Clermont[3], Marc Dieu[4], Françoise Tilquin[1], Laurens Maertens[1,5], Rob Van Houdt[5], Patricia Renard[4], Eric Perpete[2] and Jean-Yves Matroule[1*]

[1]Research Unit in Biology of Microorganisms (URBM), Department of Biology, Namur Research Institute for Life Sciences (NARILIS), University of Namur, Namur, Belgium.

[2] Laboratoire de Chimie Physique des Biomolécules, Namur Research Institute for Life Sciences (NARILIS) and Namur Institute of Structured Matter (NISM), University of Namur, Rue de Bruxelles, 61, 5000 Namur, Belgium.

[3] Université de Lorraine, CNRS, IMoPA, F-54000 Nancy, France.

[4]MaSUN, Mass Spectrometry Facility, University of Namur, Namur, Belgium.

[5]Microbiology Unit, Interdisciplinary Biosciences, Belgian Nuclear Research Centre (SCK CEN), 2400 Mol, Belgium.

*Corresponding author. Mailing address: Research Unit in Biology of Microorganisms (URBM), 61 Rue de Bruxelles 5000 Namur, Belgium. Phone: (32)81-72-42-45. E-mail: jean-yves.matroule@unamur.be

Abstract

Chemotaxis is a widespread strategy used by unicellular and multicellular living organisms to maintain their fitness in stressful environments. In bacteria, the control of chemotaxis relies on the sensing of environmental changes by chemoreceptors, which modulate flagellar rotation via the well-characterized CheA-CheY two-component system. However, the molecular mechanisms underlying signal recognition and activation of the chemoreceptors remain largely unclear.

Here we address this gap in the frame of a chemotactic response sustaining bacterial flight from toxic copper (Cu). We found that a significant Cu accumulation in the cytosol triggers an oxidative stress, which can be abrogated by the overexpression of the superoxide dismutase SodB or of the catalase KatG. Interestingly, the inhibition of ROS production not only improves cell growth but also impedes Cu-chemotaxis, suggesting that ROS derived from cytosolic Cu mediate the control of Cu chemotaxis. We also provide evidence that cellular Cu accumulation favors Cu binding to the sensor domain of the newly identified chemoreceptor McpR. Finally, we discovered that the redox cycling of the McpR-bound Cu modulates McpR conformation. Overall our study provides mechanistic insights on a redox-based sensing by a chemoreceptor, indicating that the cellular redox status can play a key role in bacterial chemotaxis.

Introduction

Cell motility is widely distributed across the three domains of life. For instance, in animals it is involved in fertilization and in neutrophils recruitment to an inflammatory area. It also allows archaea and bacteria to seek nutrients in scarce environments and to flee from toxic compounds while it supports to some extent host tropism in the frame of symbiotic relationships.

Most of the time, cell motility is driven by chemical gradients and is therefore named chemotaxis. In flagellated bacteria, positive (attraction) and negative (repulsion) chemotactic responses rely on the well characterized CheA-CheY two-component system that modulates flagellar rotation, thereby inducing stochastic direction changes [1]. The activation state of CheA-CheY is controlled by finely tuned cytoplasmic or inner membrane-spanning methyl-accepting chemotaxis proteins (Mcp) monitoring chemical variations in the cytosol or in the periplasm [2]. However, how the ligand is recognized and bound to the MCP and how this event controls MCP activity is still largely unexplored.

Here we addressed this gap in chemotaxis regulation by deciphering a sensing mechanism underlying the bacterial chemotactic response to a toxic Cu-rich environment. We provide evidence that a high intracellular Cu concentration triggers the production of reactive oxygen species (ROS), which is a prerequisite to Cu chemotaxis. We also demonstrate that the redox cycling of a Cu ion bound to the sensor domain of the newly-identified McpR modulates McpR conformation. This original redox sensing mechanism may ensure a continuous monitoring of the intracellular Cu status, providing a real-time control of the chemotactic response.

Results

In a previous study, we identified an original bimodal strategy to cope with Cu stress where the flagellated morphotype (swarmer cells) of *C. crescentus* triggers a prompt negative chemotaxis from the Cu source [3]. In the same study, we also showed that a significant intracellular Cu accumulation is a prerequisite to sustain this chemotactic response, suggesting that the sensing of Cu variations when the swarmer (SW) cell swims along the Cu gradient is performed intracellularly.

In order to add a subcellular resolution to this Cu accumulation, we measured the Cu content by ICP-OES in SW cell fractionates. In accordance with our previous study [3], Cu exposure led to an important cellular Cu accumulation albeit at a much lesser extent in the cytoplasm than in the periplasm (**Fig. 1A**).

The transient maintenance of a high intracellular Cu content in the SW cells may potentially lead to an oxidative stress via the production of superoxide anion ($O_2^{\bullet-}$), hydrogen peroxide (H_2O_2) and hydroxyl radical ($OH^{\bullet}$) (**Fig. S1A**). In line with this idea, the relative fluorescence intensity of the ROS sensitive turn-on CellROX Deep Red probe is significantly increased in SW cells exposed to Cu or H_2O_2, the latter being used as a positive control (**Fig. 1B**). Accordingly, the fluorescence of the cytoplasmic turn-off rxYFP redox biosensor is partially quenched under the same conditions (**Fig. 1C**), mirroring an imbalance of the cytoplasmic GSH-GSSG pool upon oxidative stress (**Fig. S1B**). These findings are reinforced by the differential expression of genes involved in oxidative stress response upon Cu stress, including those encoding the catalase KatG (CCNA_03138) and the superoxide dismutases SodA (CCNA_03672) and SodB (CCNA_01855) [4]. Together these data argue for the rapid generation of an oxidative stress in SW cells exposed to Cu.

In order to determine whether the ROS derived from cellular Cu accumulation play a role

in Cu chemotaxis, we envisioned that boosting the aforementioned antioxidant defense system

would limit Cu-induced ROS production in SW cells. We ectopically expressed the *sodB* or the

katG genes in the WT background from the low copy pMR10 plasmid under the control of the

strong and constitutive lac promoter. The resulting *sodB*$^+$ and *katG*$^+$ strains prevent the increase

of the CellROX Deep Red fluorescence upon Cu exposure, reflecting a potentiation of

antioxidant defenses (**Fig. 1D**). In support of this data, the growth of these two optimized *sodB*$^+$

and *katG*$^+$ strains is significantly improved under moderate Cu stress (**Fig. 1E**), arguing for a

role of $O_2^{\cdot-}$, H_2O_2 in Cu toxicity.

In order to assess the potential role of these ROS in Cu chemotaxis, we took advantage of

the aforementioned *sodB*$^+$ and *katG*$^+$ strains, where ROS formation is largely hindered. By

using the previously designed Live Chemotaxis Imaging (LCI) [3], we showed that the *sodB*$^+$

and *katG*$^+$ strains are impaired in Cu chemotaxis, at a similar extent to the Δ*cheA1* mutant (**Fig.**

1F, **SI Appendix**). Consistent with a role of KatG in H_2O_2 detoxification, the *katG*$^+$ strain is

also impeded in H_2O_2 chemotaxis (**Fig. S2**). These data support a major role of the ROS

resulting from Cu accumulation in Cu chemotaxis.

To further dissect the detailed mechanisms of Cu sensing, we sought to identify the

chemoreceptor involved in Cu chemotaxis. The genome of *C. crescentus* is predicted to encode

19 Mcp (named McpA to McpS) [5]. No canonical Cu binding site, including the CXXC motif

or the βαββαβ fold [6], could be found by running manual and JPred4-based analyses of the

19 Mcp sequences [7]. Therefore, we proceeded to single in-frame deletions of the 19 *mcp*

genes, and their respective chemotactic behavior to Cu was assessed by LCI (**Fig. 2A**). Most of

the Δ*mcp* mutants displayed a WT-like behavior with a flight percentage after 25 min varying

between 15% and 51%. Only Δ*mcpA*, Δ*mcpB* and Δ*mcpR* mutants did not significantly cross

the arbitrary threshold of 10% flight, suggesting that Cu chemotaxis might be severely affected

in these genetic backgrounds. The Δ*mcpA*, Δ*mcpB* and Δ*mcpR* mutants could be complemented

by the expression of an ectopic version of *mcpA*, *mcpB* and *mcpR* on the low copy pMR10

plasmid, respectively, suggesting that they may potentially control Cu chemotaxis (**Fig. S3**).

However, McpA and McpB are likely involved in chemotaxis at a more global extent

considering (i) the position of the *mcpA* and *mcpB* genes within and in the vicinity of the major

chemotaxis cluster, respectively [8], and (ii) the reduced motility halo of the Δ*mcpA* mutant

observed in a Cu-free swarming assay (**Fig. S4**). In addition, McpB displays the terminal

sequence of the CheB/CheR docking site that may be necessary for the adaptive response [9],

which will not be considered in this study. Therefore, we aimed at further characterizing the

role of McpR in Cu chemotaxis.

Owing to the multimeric organization of Mcp clusters in bacteria [10], one may argue that

the loss of Cu chemotaxis in the Δ*mcpR* mutant results from a loss of motility and/or a disruption

of the general chemotactic machinery. The ability of the Δ*mcpR* mutant to form a motility halo

in a Cu-free swarming assay (**Fig. S5A**) and to sustain negative chemotaxis to other chemicals,

such as zinc (Zn) (**Fig. S5B**), refutes this hypothesis. It is also worth mentioning that the loss

of Cu chemotaxis in the Δ*mcpR* mutant is not due to a loss of ROS production under Cu stress

since the Δ*mcpR* mutant exhibits a similar decrease of the rxYFP fluorescence to the WT strain

upon Cu exposure (**Fig. S6**).

Mcp are able to sense attractants or repellents in a direct or indirect manner [11]. Therefore,

we decided to assess by intrinsic fluorescence spectroscopy the direct binding of Cu to purified

McpR. When mixed with Cu(I) or Cu(II) at 1:1, 1:10 and 1:100 ratios, the intrinsic McpR

fluorescence at 346 nm is quenched in a dose-dependent manner, reflecting a modification of

the McpR aromatic residues environment (**Fig. 2B**, see **Fig. S7** for full spectra). The

dissociation constants (Kd) of 48 μM +/- 10 μM and 37 μM +/- 9 μM for Cu(I) and Cu(II),

respectively, attest to a similar affinity of both ionic forms for McpR. However, the micromolar

range of Cu(I) and Cu(II) Kds exceeds from several logs the Kd of well-known cuproteins

[12, 13], arguing for a rather low affinity of McpR for Cu. Therefore, one may argue that the

high Cu cellular content measured in the Cu-exposed SW cells is necessary to promote a rather

labile interaction between McpR and Cu. Interestingly, McpR seems to preferentially bind Cu

since Zn, cadmium (Cd), manganese (Mn) and nickel (Ni) had no effect on McpR intrinsic

fluorescence (**Fig. S8**).

The complete loss of Cu-chemotaxis in the *sodB+* and *katG+* genetic backgrounds (**Fig.**

1F) indicates that Cu binding to McpR is solely not sufficient to sustain Cu chemotaxis.

Consistent with this idea, we found that H_2O_2 chemotaxis is fully abolished in the Δ*mcpR*

mutant, indicative of a role of McpR as a sensor of the oxidative stress (**Fig. 2C**). Owing to the

oxidative conditions generated by Cu accumulation in the cytosol (**Fig. 1B-C**), we sought to

assess by LC/MS the oxidation status of McpR with a particular focus on the sensor domain

(residues 1-161) after incubating purified McpR with Cu(II) or Cu(I) +/- H_2O_2 at a 1:100 ratio.

Only the Cu(I) + H_2O_2 condition, and to a lesser extent Cu(I), led to a significant oxidation of

the McpR sensor domain, and more specifically of the H99 residue. In this particular condition,

about 30% of the H99 residues acquired an additional oxygen ion, most likely yielding 2-

oxoHis (**Fig. 3A**). In accordance with this observation, the expression of an ectopic *mcpR$_{H99A}$*

allele in the Δ*mcpR* mutant does not restore its ability to sustain Cu chemotaxis, indicating that

H99 is a key residue in McpR Cu chemotaxis (**Fig. 3B**). Noteworthy, H99A mutation does not

impede Cd chemotaxis for which McpR is also essential (**Fig. S9**), confirming the functionality

of McpR$_{H99A}$ and the proper assembly of the chemotactic machinery.

Metal catalyzed oxidation (MCO) of histidines via the generation of OH$^{\bullet}$ has been

previously described in the *Bacillus subtilis* PerR repressor [14], in yeast SOD1 [15], and in

amyloid beta (Aβ) peptide [16]. The exact role of this modification remains unknown. In

addition, no redox cycling involving 2-oxoHis has been described so far, questioning the

relevance of H99 oxidation in Cu/ROS sensing. MCO mostly targets the metal-binding residues

in a caged process [17], suggesting that H99 may coordinate Cu. To test this hypothesis, we

calculated the Kd values of the McpR$_{H99A}$/Cu(I) and McpR$_{H99A}$/Cu(II) complexes (**Fig. 3C**).

H99A mutation triggers a 9.4- and 5.8-fold increase of Cu(I) and Cu(II) Kd, respectively,

relative to McpR, suggesting a key role of H99 in Cu(I) and Cu(II) coordination. The important

variability associated with the McpR$_{H99A}$ Kd values likely reflects a potential instability of this

mutant in solution upon Cu(I) and Cu(II) exposure, resulting in the presence of several

subpopulations of McpR$_{H99A}$.

McpR sequence analysis by the TMHMM 2.0 [18] and SignalIP 6.0 [19] tools did not

reveal any McpR transmembrane domains and signal peptide, respectively, arguing for a

cytoplasmic McpR localization and therefore a Cu sensing. The 3D structure prediction of

McpR by a fold recognition method (Phyre2 [20]) highlights a cysteine (C75) residue in the

vicinity of H99 (**Fig. 3D**). Cysteine residues often contribute to Cu coordination and the

measured 19.8 Å distance in the predicted model is compatible with Cu coordination by C75

and H99 residues [21]. In line with this idea, a Δ*mcpR* mutant expressing an ectopic *mcpR$_{C75S}$*

allele was unable to trigger Cu chemotaxis (**Fig. 3B**), indicating that C75 also plays a key role

in Cu-sensing. As a control, the McpR$_{C75S}$ mutant was still able to flee from H$_2$O$_2$, for which

McpR is also required (**Fig. S10**), attesting the functionality of the McpR$_{C75S}$ mutant within the

chemotaxis machinery. In order to determine whether the loss-of-function of the McpR$_{C75S}$

mutant in Cu chemotaxis resulted from a loss of Cu binding, we calculated the Kd values of the

McpR$_{C75S}$/Cu(I) and McpR$_{C75S}$/Cu(II) complexes. Similar to the H99A mutation, the C75S

mutation also impeded Cu(I) binding as attested by a 3.5-fold Kd increase. Surprisingly, Cu(II)

binding to the McpR$_{C75S}$ mutant remained unchanged (**Fig. 3C**), suggesting that C75 is required

to coordinate Cu(I) but not Cu(II). Consistent with this observation, H99 oxidation by Cu(I) +/-

H_2O_2 is fully abolished in the McpR$_{C75S}$ mutant (**Fig. 3E**). The Kd values calculated for the

McpR$_{C75S}$ and McpR$_{H99A}$ mutants are in agreement with the known selective binding of Cu(I)

and Cu(II) to cysteine and histidine residues, respectively [21, 22].

The aforementioned observations led us to the idea that Cu(I) coordination by H99 and

C75 is decisive for McpR-mediated Cu chemotaxis. Given the structural changes elicited by

the binding of a ligand to a cognate chemoreceptor [23], we envisioned that Cu(I) binding to

McpR modulates its conformation. To test this idea, Far-UV circular dichroism (CD)

spectroscopy was used to monitor the impact of Cu(I) and Cu(II) on McpR secondary structure.

Accordingly, Cu(I) has a more significant impact than Cu(II) on McpR secondary structure

(**Fig. 3F**), increasing the α-helix percentage from 23% to 44%.

Based on our findings, we propose that Cu excess within the cytoplasm catalyzes the

production of ROS including $O_2^{\cdot-}$ and H_2O_2 that are required for Cu chemotaxis. Cu excess also

favors a rather labile Cu binding to the McpR sensor domain. Cu(I) coordination by McpR C75

and H99 residues triggers McpR conformational change. Interestingly, Cu(II) only binds McpR

H99 residue and does not modify the apo form-like conformation. These observations suggest

that a redox cycling of the McpR-bound Cu atom could modulate McpR conformation (**Fig. 4**).

Discussion

Although the signaling pathway controlling flagellar rotation in bacterial chemotaxis has

been extensively studied, the sensing mechanisms by the upstream chemoreceptors (MCP) are

still largely unknown. We sought to address this gap in the frame of negative chemotaxis to

toxic copper (Cu) in the bacterial model *Caulobacter crescentus*.

Our study identified the first Cu sensing MCP hereafter named McpR. McpR is devoid of

any transmembrane domains or signal peptide and is therefore predicted to localize in the

cytoplasm. Cytoplasmic MCPs have been shown to monitor the cellular metabolic state [24],

whereas inner membrane MCPs usually sense the periplasmic environment, as a proxy for

extracellular changes. The cytoplasmic Cu content is significantly increased albeit to a lesser

extent than in the periplasm when the SW cells are exposed to a toxic Cu concentration. To

explain the rationale of using a cytoplasmic MCP to monitor extracellular changes, one could

hypothesize that the sensing of cytoplasmic Cu provides a more sensitive monitoring of subtle

variations of Cu concentrations when the SW cell swims down a Cu gradient.

In a previous study, we showed that the maintenance of a high cellular Cu concentration is

critical for Cu chemotaxis [3]. Here, we provide evidence that McpR binds Cu with low affinity,

suggesting that this binding is only possible when the Cu concentration within the cytoplasm is

high. Considering that Cu binding to McpR is a prequisite for Cu chemotaxis, one could propose

that when the SW cell swim away from the Cu source, the cytoplasmic Cu concentration

decreases and ultimately impede Cu binding to McpR and the resulting chemotactic signaling

pathway.

We also demonstrate that the cellular Cu accumulation triggers by an unknown mechanism

the production of ROS that can be impeded by the overexpression of SodB and KatG catalase,

suggesting that at least O_2^{-} and H_2O_2 are generated. The contribution of the ROS generated by

cellular Cu in Cu toxicity is still largely debated [25, 26]. Here, we demonstrate that ROS

derived from cellular Cu not only impede cell growth to some extent, but also play a key role

in Cu chemotaxis. The exact target of ROS in Cu chemotaxis was not identified in this study

but one could hypothesize that O_2^{-} is rapidly converted into H_2O_2. The latter would feed a local

Fenton-like reaction catalyzed by the Cu-(I) atom coordinated by McpR C75 and McpR H99

residues. This would lead to a local OH· production and to the oxidation of McpR-bound Cu(I)

into McpR-bound Cu(II). Owing to the reducing cytoplasmic environment, McpR-bound Cu(II)

could in turn be reduced by cytoplasmic electron donors such as GSH, NADH and thioredoxin

to restore the Cu(I) ionic form. Thereby, the McpR-bound Cu would undergo a redox cycling

involving the Cu-(I) and Cu-(II) ionic forms, which would be maintained as long as H_2O_2 is

produced by cytoplasmic Cu. Cu-generated ROS would therefore act as second messengers in

a bacterial stress response.

The modulation of the CheA-CheY two-component system results from MCP

conformational changes occurring upon ligand binding [27]. We demonstrated here that McpR

conformation is modulated by the ionic status of the Cu atom bound to McpR. It should be

emphasized that MCPs are usually organized in trimers of dimers and assembled in large arrays

in cells [28]. Therefore, it is possible that Cu coordination by McpR C75 and McpR H99

residues within the McpR array implies two McpR monomers from one or two distinct dimers.

Nevertheless, McpR conformational change likely regulates the downstream effectors and the

resulting chemotactic response.

Materials and Methods

Strains, plasmids and growth conditions

Caulobacter crescentus (NA1000 strain) was grown at 30°C in Peptone Yeast Extract (PYE)

medium [29] with 5 μg/ml kanamycin, 1 μg/ml chloramphenicol and/or an appropriate

concentration of $CuSO_4.5H_2O$ when required. Cultures in exponential phase were used for all

experiments. Strains and plasmids are listed in Table S1, and the strategies for their construction

are available upon request. SW cells isolation was achieved according to [30].

Growth curve measurements

C. crescentus exponential phase cultures were diluted in PYE medium to a final O.D.$_{660\ nm}$ of 0.05 and inoculated in 96-well plates with appropriate CuSO$_4$ concentration when required. Bacteria were then grown for 24 h at 30°C under continuous shaking in an Epoch 2 Microplate Spectrophotometer from BioTek (Santa Clara, CA, USA). The absorbance at O.D.$_{660\ nm}$ was measured every 10 min.

Motility assay

C. crescentus cultures grown in PYE were diluted to an O.D.$_{660\ nm}$ of 0.4 and inoculated with sterile toothpicks on a PYE swarming plates (PYE + 0.25% agar). The plates were incubated at 30°C for 3 days, then scanned.

Cellular fractionation and ICP-OES measurements

C. crescentus cells were fixed for 20 min in 2% paraformaldehyde at 4°C and then washed three times with an ice-cold wash buffer (10 mM Tris-HCl pH 6.8, 100 μM EDTA). The cells were then incubated with 0.125% zwitterionic detergent. A final centrifugation at 14,500 x g for 15 min separated the cytoplasm (and inner membrane fraction) from the periplasm (and outer membrane fraction). The two fractions were incubated overnight with a final concentration of 1 M HNO$_3$ and then analyzed by inductively coupled plasma - optical emission spectrometry (ICP-OES) with an Optima 8000 ICP-OES from PerkinElmer. Cellular Cu concentration were calculated as described in [3].

Oxidative stress measurement with the CellROX Deep Red probe

An exponential phase culture of *C. crescentus* was synchronized to isolate the SW cells. The isolated SW cells were diluted to an OD$_{660nm}$ of 0.2 and incubated for 30 min at 30°C under moderate shaking with 5 μM of CellROX Deep Red probe (ThermoFisher) and appropriate concentrations of CuSO$_4$ or H$_2$O$_2$. The bacteria were then washed three times with 20 mM

276 Phosphate Buffer to remove the non-internalized probes. The bacteria were finally transferred

277 into a black bottom 96-wells plate and the fluorescence was monitored at 640/690 nm with a

278 SpectraMax iD3 (Molecular Devices, San Jose, CA, USA).

Oxidative stress measurement with the rxYFP biosensor

280 The SW cells were isolated from an exponential phase culture overexpressing the rxYFP

281 biosensor [31]. The SW cells were diluted to an OD_{660nm} of 0.4 in a black bottom 96-well plate

282 and mixed with different concentrations of $CuSO_4$ and H_2O_2. The rxYFP fluorescence was then

283 monitored every minute for 20 min at 510/556 nm with a SpectraMax iD3 (Molecular Devices,

284 San Jose, CA, USA).

Live Chemotaxis Imaging

The live chemotaxis imaging (LCI) assay was adapted from[3]. Eight microliters of melted

1.5% agarose H_2O with $CuSO_4$, H_2O_2, $ZnSO_4$ or $CdSO_4$ were loaded into the chamber through

the external inlet channels to generate the plug. One hundred and fifty microliters of isolated

SW cells were in turn injected into the bacterial chamber through the central inlet channel.

Images were collected every min for 25 min in one focal plan (near the liquid surface) in the

vicinity of the plug with a Nikon Eclipse Ti2-E inverted microscope equipped with a Nikon

20X/0.5 Ph2 objective and with a Hamamatsu C13440 digital camera. Quantitative analysis of

the time-lapse images was performed with MicrobeJ [32]. The number of bacteria in a same

focal plane was determined at each time point.

Protein expression and purification

A BL21 pET28a-McpR$_{His}$ exponential phase culture was incubated for 3 h under shacking at

37°C with 1 mM Isopropyl-β-D-thiogalactopyranoside (IPTG) and then harvested by

centrifugation for 30 min at 4,000 x g at 4°C. McpR$_{His}$ induction was verified by 12 % Tricine–

sodium dodecyl sulfate–polyacrylamide gel electrophoresis (Tricine–SDS–PAGE), followed

by Coomassie Blue staining and by WB using an anti-His antibody (1/5000). In order to purify

McpR$_{His}$, the IPTG-induced culture was washed with 0.1 M phosphate buffer (pH 8) and

centrifuged for 15 min at 9,000 x g at 4°C. Bacterial cells were lysed at 2.4 kbar using a Cell

disruptor (Cell Disruption System – one shot model, Constant). As McpR$_{His}$ was mainly

forming inclusion bodies, the pellet fraction was resuspended for 1 h under shaking at 4°C in a

0.1 M phosphate buffer (pH 8) containing 6 M guanidine. The solubilized inclusion bodies were

centrifuged for 30 min at 9,000 x g at 4 °C. The collected supernatant was dialyzed overnight

in a 0.1 M phosphate buffer (pH 8) to remove guanidine.

Fluorescence quenching by Cu and other metal ions

309   McpR protein concentration was determined (i) by measuring the absorbance of the sample at

310   280 nm with the SpectraMax iD3 or (ii) with a Bradford protein assay. 3.58 µM of McpR was

311   incubated with different ratios of CuSO$_4$, Tetrakis(acetonitrile)copper(I) hexafluorophosphate,

312   ZnSO$_4$, CdSO$_4$, MnSO$_4$ and NiSO$_4$ in black bottom 96-wells plates. The fluorescence emission

313   spectra were recorded upon 280 nm excitation with the SpectraMax iD3.

Kd determination

315   The titration of McpR by Cu(I) or Cu(II) monitored by the protein intrinsic fluorescence

316   intensity quenching was fitted to a single-site hyperbolic model by non-linear regression using

317   the Scidavis 2.7 software to determine the affinity constants. Dissociation constants are reported

318   as the mean value ± standard deviation obtained from a minimum of three independent

319   experiments performed on distinct protein productions.

Identification of McpR oxidation by mass spectrometry

Protein digestion

McpR samples were treated using Filter-aided sample preparation (FASP) using the following

protocol. To first wash the filter, 100 µl of formic acid 1 % were placed in each Millipore

Microcon 30 MRCFOR030 Ultracel PL-30 before a 15 min centrifugation at 14,100 x g. For

each sample, 40 µg of protein adjusted in 150 µl of 8 M urea buffer (8 M urea in 0.1 M Tris

buffer at pH 8.5) were placed individually in a column and centrifuged for 15 min at 14,100 x

g. The filtrate was discarded and the columns were washed three times by adding 200 µl of urea

buffer followed by a 15 min centrifugation at 14,100 x g. For the reduction step, 100 µl of

dithiothreitol (DTT) were added and mixed for 1 min at 400 rpm with a thermomixer before a

15 min incubation at 24 °C. The samples were then centrifugated for 15 min at 14,100 x g, the

filtrate was discarded and the filter was washed by adding 100 µl of urea buffer before another

15 min centrifugation at 14,100 x g. An alkylation step was performed by adding 100 µl of

iodoacetamide ((IAA), in urea buffer) in the column and mixing for 1 min at 400 rpm in the

dark followed by a 20 min incubation in the dark and a 10 min centrifugation at 14,100 x g. To

remove the excess of IAA, 100 µl of urea buffer were added and the samples were centrifuged

for 15 min at 14,100 x g. To quench the rest of IAA, 100 µl of DTT were placed on the column,

mixed for 1 min at 400 rpm and incubated for 15 min at 24 °C before a 10 min centrifugation

at 14,100 x g. To remove the excess of DTT, 100 µl of urea buffer were placed on the column

and centrifuged for 15 min at 14,100 x g. The filtrate was discarded and the column was washed

three times by adding 100 µl of 50 mM sodium bicarbonate buffer ((ABC), in ultrapure water)

followed by a 10 min centrifugation at 14,100 x g. The remaining 100 µl were kept at the bottom

of the column to avoid any evaporation in the column. The digestion process was performed by

adding 80 µl of mass spectrometry grade trypsin (1/50 in ABC buffer) in the column and mixed

for 1 min at 400 rpm followed by an overnight incubation at 24°C in a water saturated

environment. The Microcon columns were placed in a 1.5 ml LoBind tube and centrifuged for

10 min at 14,100 x g. Fourty µl of ABC buffer were placed on the column before a 10 min

centrifugation at 14,100 x g. Ten % Trifluoroacetic acid (TFA) in ultrapure water were added

into the LoBind Tube to obtain 0.2 % TFA. The samples were dried in a SpeedVac up to 20 µl

and transferred into an injection vial.

Mass spectrometry

The digest was analyzed using nano-LC-ESI-MS/MS tims TOF Pro (Bruker, Billerica, MA,

USA) coupled with an UHPLC nanoElute (Bruker).

Peptides were separated by nanoUHPLC (nanoElute, Bruker) on a 75 µm ID, 25 cm C18

column with integrated CaptiveSpray insert (Aurora, ionopticks, Melbourne) at a flow rate of

400 nl/min, at 50°C. LC mobile phases A was water with 0.1% formic acid (v/v) and B ACN

with 0.1% formic acid (v/v). Samples were loaded directly on the analytical column at a

constant pressure of 800 bar. The digest (1 µl) was injected, and the organic content of the

mobile phase was increased linearly from 2% B to 15 % in 36 min, from 15 % B to 25% in 19

min, from 25% B to 37 % in 5 min and from 37% B to 95% in 5 min. Data acquisition on the

tims TOF Pro was performed using Hystar 5.1 and timsControl 2.0. tims TOF Pro data were

acquired using 100 ms TIMS accumulation time, mobility (1/K0) range from 0.6 to 1.6 Vs/cm².

Mass-spectrometric analysis were carried out using the parallel accumulation serial

fragmentation (PASEF) acquisition method. One MS spectra followed by ten PASEF MSMS

spectra per total cycle of 1.1 s.

Data analysis was performed using PEAKS Studio X Pro with ion mobility module

(Bioinformatics Solutions Inc., Waterloo, ON). Protein identifications were conducted using

PEAKS search engine with 15 ppm as parent mass error tolerance and 0.05 Da as fragment

mass error tolerance. Carbamidomethylation was allowed as fixed modification, oxidation of

methionine, histidine, tryptophan and acetylation (N-term) as variable modification. Enzyme

specificity was set to trypsin, and the maximum number of missed cleavages per peptide was

set at three. The peak lists were searched against the *E. coli*-K12 (11002 sequences) and

Caulobacter NA1000 (3863 sequences) from UNIREF 100. Peptide spectrum matches and

protein identifications were normalized to less than 1.0% false discovery rate.

The identification results were processed with the PTM profile tool from PEAKS, which

provides quantitative information of modified peptides compared with unmodified peptides for

the modification sites of the protein across all MS samples [33, 34].

Circular dichroism

8.95 µM of McpR was incubated with different $CuSO_4$ and Tetrakis(acetonitrile)copper(I)

hexafluorophosphate concentrations. Measurements were performed with a MOS-500

spectropolarimeter in the far-UV region (190-260 nm), using a 1 mm pathlength cell. Four scans

were averaged and base lines were subtracted. Secondary structure analyses were performed on

the circular dichroism data with the BeStSel server.

383
384

Acknowledgements

We would like to thank Jakob R. Winther for providing the rxYFP construct. We are also grateful to Valérie Charles and Carmela Aprile (CMI laboratory, NISM, UNamur) for the ICP measurements and to Virgile Neyman for technical support during the protein purification. Finally, we would like to acknowledge Jean-François Collet and Xavier De Bolle for critical reading of this manuscript and the URBM members and Johan Wouters (Chemistry Department, UNamur) for fruitful discussions. This work was supported by the University of Namur. G. L. was supported by the Belgian Fund for Industrial and Agricultural Research Associate (FRIA). Catherine Michaux and Eric Perpète thank the Belgian National Fund for Scientific Research for their respective associate and senior research associate positions.

Bibliography

1. Wadhams, G.H., and Armitage, J.P. (2004). Making sense of it all: bacterial chemotaxis. Nat Rev Mol Cell Biol *5*, 1024-1037.

2. Yang, W., and Briegel, A. (2020). Diversity of Bacterial Chemosensory Arrays. Trends Microbiol *28*, 68-80.

3. Lawaree, E., Gillet, S., Louis, G., Tilquin, F., Le Blastier, S., Cambier, P., and Matroule, J.Y. (2016). Caulobacter crescentus intrinsic dimorphism provides a prompt bimodal response to copper stress. Nat Microbiol *1*, 16098.

4. Maertens, L., Cherry, P., Tilquin, F., Van Houdt, R., and Matroule, J.Y. (2021). Environmental Conditions Modulate the Transcriptomic Response of Both Caulobacter crescentus Morphotypes to Cu Stress. Microorganisms *9*.

5. Nierman, W.C., Feldblyum, T.V., Laub, M.T., Paulsen, I.T., Nelson, K.E., Eisen, J.A., Heidelberg, J.F., Alley, M.R., Ohta, N., Maddock, J.R., et al. (2001). Complete genome sequence of Caulobacter crescentus. Proc Natl Acad Sci U S A *98*, 4136-4141.

6. Lamb, A.L., Wernimont, A.K., Pufahl, R.A., Culotta, V.C., O'Halloran, T.V., and Rosenzweig, A.C. (1999). Crystal structure of the copper chaperone for superoxide dismutase. Nat Struct Biol *6*, 724-729.

7. Drozdetskiy, A., Cole, C., Procter, J., and Barton, G.J. (2015). JPred4: a protein secondary structure prediction server. Nucleic Acids Res *43*, W389-394.

8. Berne, C., and Brun, Y.V. (2019). The Two Chemotaxis Clusters in Caulobacter crescentus Play Different Roles in Chemotaxis and Biofilm Regulation. J Bacteriol *201*.

9. Barnakov, A.N., Barnakova, L.A., and Hazelbauer, G.L. (1998). Comparison in vitro of a high- and a low-abundance chemoreceptor of Escherichia coli: similar kinase activation but different methyl-accepting activities. J Bacteriol *180*, 6713-6718.

10. Briegel, A., Wong, M.L., Hodges, H.L., Oikonomou, C.M., Piasta, K.N., Harris, M.J., Fowler, D.J., Thompson, L.K., Falke, J.J., Kiessling, L.L., et al. (2014). New insights into bacterial chemoreceptor array structure and assembly from electron cryotomography. Biochemistry *53*, 1575-1585.

11. Neumann, S., Hansen, C.H., Wingreen, N.S., and Sourjik, V. (2010). Differences in signalling by directly and indirectly binding ligands in bacterial chemotaxis. EMBO J *29*, 3484-3495.

12. Xiao, Z., Brose, J., Schimo, S., Ackland, S.M., La Fontaine, S., and Wedd, A.G. (2011). Unification of the copper(I) binding affinities of the metallo-chaperones Atx1, Atox1, and related proteins: detection probes and affinity standards. J Biol Chem *286*, 11047-11055.

13. Kahra, D., Kovermann, M., and Wittung-Stafshede, P. (2016). The C-Terminus of Human Copper Importer Ctr1 Acts as a Binding Site and Transfers Copper to Atox1. Biophys J *110*, 95-102.

14. Lee, J.W., and Helmann, J.D. (2006). The PerR transcription factor senses H2O2 by metal-catalysed histidine oxidation. Nature *440*, 363-367.

15. Tiwari, M.K., Hagglund, P.M., Moller, I.M., Davies, M.J., and Bjerrum, M.J. (2019). Copper ion / H2O2 oxidation of Cu/Zn-Superoxide dismutase: Implications for enzymatic activity and antioxidant action. Redox Biol *26*, 101262.

16. Ramteke, S.N., Walke, G.R., Joshi, B.N., Rapole, S., and Kulkarni, P.P. (2014). Effects of oxidation on redox and cytotoxic properties of copper complex of Abeta1-16 peptide. Free Radic Res *48*, 1417-1425.

17. Stadtman, E.R. (1993). Oxidation of free amino acids and amino acid residues in proteins by radiolysis and by metal-catalyzed reactions. Annu Rev Biochem *62*, 797-821.

18. Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E.L. (2001). Predicting
transmembrane protein topology with a hidden Markov model: application to complete
genomes. J Mol Biol *305*, 567-580.
19. Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gislason, M.H., Pihl, S.I.,
Tsirigos, K.D., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022). SignalP
6.0 predicts all five types of signal peptides using protein language models. Nat
Biotechnol.
20. Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J. (2015). The
Phyre2 web portal for protein modeling, prediction and analysis. Nat Protoc *10*, 845-
858.
21. Rubino, J.T., and Franz, K.J. (2012). Coordination chemistry of copper proteins: how
nature handles a toxic cargo for essential function. J Inorg Biochem *107*, 129-143.
22. Rubino, J.T., Chenkin, M.P., Keller, M., Riggs-Gelasco, P., and Franz, K.J. (2011). A
comparison of methionine, histidine and cysteine in copper(I)-binding peptides reveals
differences relevant to copper uptake by organisms in diverse environments.
Metallomics *3*, 61-73.
23. Chervitz, S.A., and Falke, J.J. (1996). Molecular mechanism of transmembrane
signaling by the aspartate receptor: a model. Proc Natl Acad Sci U S A *93*, 2545-2550.
24. Salah Ud-Din, A.I.M., and Roujeinikova, A. (2017). Methyl-accepting chemotaxis
proteins: a core sensing element in prokaryotes and archaea. Cell Mol Life Sci *74*, 3293-
3303.
25. Dupont, C.L., Grass, G., and Rensing, C. (2011). Copper toxicity and the origin of
bacterial resistance--new insights and applications. Metallomics *3*, 1109-1118.
26. Macomber, L., Rensing, C., and Imlay, J.A. (2007). Intracellular copper does not
catalyze the formation of oxidative DNA damage in Escherichia coli. J Bacteriol *189*,
1616-1626.
27. Parkinson, J.S., Hazelbauer, G.L., and Falke, J.J. (2015). Signaling and sensory
adaptation in Escherichia coli chemoreceptors: 2015 update. Trends Microbiol *23*, 257-
266.
28. Khursigara, C.M., Wu, X., and Subramaniam, S. (2008). Chemoreceptors in
Caulobacter crescentus: trimers of receptor dimers in a partially ordered hexagonally
packed array. J Bacteriol *190*, 6805-6810.
29. Poindexter, J.S. (1981). The caulobacters: ubiquitous unusual bacteria. Microbiol Rev
45, 123-179.
30. Evinger, M., and Agabian, N. (1977). Envelope-associated nucleoid from Caulobacter
crescentus stalked and swarmer cells. J Bacteriol *132*, 294-301.
31. Ostergaard, H., Tachibana, C., and Winther, J.R. (2004). Monitoring disulfide bond
formation in the eukaryotic cytosol. J Cell Biol *166*, 337-345.
32. Ducret, A., Quardokus, E.M., and Brun, Y.V. (2016). MicrobeJ, a tool for high
throughput bacterial cell detection and quantitative analysis. Nat Microbiol *1*, 16077.
33. Meier, F., Brunner, A.D., Koch, S., Koch, H., Lubeck, M., Krause, M., Goedecke, N.,
Decker, J., Kosinski, T., Park, M.A., et al. (2018). Online Parallel Accumulation-Serial
Fragmentation (PASEF) with a Novel Trapped Ion Mobility Mass Spectrometer. Mol
Cell Proteomics *17*, 2534-2545.
34. Zhang, J., Xin, L., Shan, B., Chen, W., Xie, M., Yuen, D., Zhang, W., Zhang, Z., Lajoie,
G.A., and Ma, B. (2012). PEAKS DB: de novo sequencing assisted database search for
sensitive and accurate peptide identification. Mol Cell Proteomics *11*, M111 010587.

Supporting information

Caulobacter crescentus genome is predicted to encode two chemotaxis clusters including genes coding for CheA histidines kinases, CheY response regulators, CheW adaptors, CheB methylesterases, CheR methytransferases, McpA and McpK and other uncharacterized Che proteins. Most of the studies conducted in *C. crescentus* so far focused on the major chemotaxis cluster (cluster 1) (1-4) while the second alternate chemotaxis cluster (cluster 2) was only recently investigated (1). These studies concluded that the cluster 1 is essential for chemotaxis, while cluster 1 and cluster 2 are involved in biofilm formation and holdfast production (1, 2). Consistent with these observations, CheAI (cluster 1 CheA) turns out to be the main histidine kinase involved in chemotaxis while the exact function of CheAII (cluster 2 CheA) remains unknown[1]. In order to determine which CheA is involved in Cu-chemotaxis, we constructed the Δ*cheAI*, Δ*cheAII* and Δ*cheAI-II* mutants and assessed their chemotactic behavior to Cu by LCI (**Fig. 1**). The Δ*cheAI* mutant completely lost its chemotactic response to Cu, while the Δ*cheAII* mutant displayed a WT phenotype. Accordingly, the Δ*cheAI-II* double mutant showed a similar behavior to the Δ*cheAI* mutant. These data confirm that the CheAI histidine kinase is the main histidine kinase involved in Cu chemotaxis.

Figure 1: CheAI is the main histidine kinase involved in Cu chemotaxis. Percentage of WT, Δ*cheAI*, Δ*cheAII* and Δ*cheAI-II* SW cells remaining in the vicinity of the Cu plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Cu plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

References

1   Alley, M. R., Maddock, J. R. & Shapiro, L. Polar localization of a bacterial chemoreceptor. *Genes Dev* **6**, 825-836, doi:10.1101/gad.6.5.825 (1992).
2   Alley, M. R., Maddock, J. R. & Shapiro, L. Requirement of the carboxyl terminus of a bacterial chemoreceptor for its targeted proteolysis. *Science* **259**, 1754-1757, doi:10.1126/science.8456303 (1993).
3   Tsai, J. W. & Alley, M. R. Proteolysis of the McpA chemoreceptor does not require the *Caulobacter* major chemotaxis operon. *J Bacteriol* **182**, 504-507, doi:10.1128/JB.182.2.504-507.2000 (2000).
4   Tsai, J. W. & Alley, M. R. Proteolysis of the *Caulobacter* McpA chemoreceptor is cell cycle regulated by a ClpX-dependent pathway. *J Bacteriol* **183**, 5001-5007, doi:10.1128/JB.183.17.5001-5007.2001 (2001).
5   Berne, C. & Brun, Y. V. The Two Chemotaxis Clusters in *Caulobacter crescentus* Play Different Roles in Chemotaxis and Biofilm Regulation. *J Bacteriol* **201**, doi:10.1128/JB.00071-19 (2019).

Figure 1

A.

B.

C.

D.

E.

F.

Figure 1: Cu-induced ROS sustain Cu chemotaxis. A. Intracellular Cu concentration in the cytoplasm (and inner membrane fraction) and in the periplasm (and outer membrane fraction) in SW cells exposed to 175 µM Cu for 5 min. **B.** Relative fluorescence intensity of SW cells incubated with 5 µM CellROX Deep Red probe and exposed to 175 µM Cu or 400 µM H_2O_2. **C.** Relative fluorescence intensity of SW cells expressing the rxYFP biosensor and exposed to 175 µM Cu or 400 µM H_2O_2 for 20 min. **D.** Relative fluorescence intensity of WT SW cells and SW cells overexpressing SodB ($sodB^+$) and KatG ($katG^+$) incubated with 5 µM CellROX Deep Red probe and exposed to 175 µM Cu. **E.** Growth profiles at $OD_{660\ nm}$ of WT, $sodB^+$ and $katG^+$ strains in PYE (orange) and exposed to 175 µM Cu (blue). **F.** Percentage of WT, $\Delta cheAI$, $sodB^+$ and $katG^+$ SW cells in the vicinity of the Cu plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Cu plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p- values* were calculated using an ANOVA when appropriate (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 2

A.

B.

C.

Figure 2: McpR is involved in Cu chemotaxis and Cu binding. A. Percentage of WT and Δ*mcp (mcpA* to *mcpS)* SW cells remaining in the vicinity of the Cu plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Cu plug) is indicated in red. **B.** Relative intrinsic fluorescence intensity of purified McpR (3.58 µM) incubated with Cu(II) and Cu(I) at 1:1, 1:10 and 1:100 ratios. **C.** Percentage of WT and Δ*mcpR* SW cells remaining in the vicinity of the H_2O_2 plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the H_2O_2 plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA when appropriate (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 3

Figure 3: Cu cycling modulates its coordination by the H99 and C75 residues. A. Percentage of oxidation of the H99 residue from purified McpR incubated with Cu(II), Cu(I), H_2O_2 and Cu(I)+H_2O_2 at a 1:100 ratio. **B.** Percentage of SW cells with a WT, $\Delta mcpR$ and $\Delta mcpR$ overexpressing McpR, McpR$_{H99A}$ or McpR$_{C75S}$ background remaining in the vicinity of the Cu plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Cu plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). **C.** Measure, by intrinsic fluorescence, of the Kd values of purified McpR, McpR$_{H99A}$ and McpR$_{C75S}$ (1.79 µM) incubated with 0 to 500 µM Cu(I) and Cu(II). Mean +/- s.d. **D.** 3D structure prediction of the sensor domain of McpR by Phyre 2. H99 and C75 residues are highlighted. The distance was measured with PyMOL viewer **E.** Percentage of oxidation of the H99 residue from purified McpR$_{C75S}$ incubated Cu(II), Cu(I), H_2O_2 and Cu(I)+H_2O_2 at a 1:100 ratio. **F.** Circular Dichroism spectra of purified McpR (8.95 µM) incubated with Cu(I) and Cu(II) at a 1:100 ratio.

Figure 4

Figure 4: Hypothetical model of Cu sensing by McpR. SW cell exposure to Cu increases the cytoplasmic Cu concentration, leading to the generation of H_2O_2 (left). In addition, high cytoplasmic Cu content favors Cu(I) coordination by the C75 and H99 residues within the cytoplasmic McpR, triggering a conformational change of the chemoreceptor. Cu(II) binding to H99 residue has no impact on McpR apo-like conformation, suggesting that a redox-cycling of the McpR-bound Cu atom could modulate McpR conformation and very likely its activity. This redox Cu cycling would be maintained along the extracellular Cu gradient until the SW cell reaches a Cu-poor environment (right). Under these conditions, the cytoplasmic Cu concentration together with the ROS level would return to their basal level and the labile Cu binding to McpR would be hindered, bringing McpR to its apo form conformation.

Figure S1

Figure S1: Potential pathways leading to ROS production by Cu. A. *In vitro* generation of Cu-induced ROS by Fenton-like reaction and Haber-Weiss cycles. The Cu-GSH complexes may also trigger ROS production (27, 28). The SODs and KatG are involved in $O_2^{\bullet -}$ and H_2O_2 detoxification, respectively **B.** The rxYFP biosensor is in equilibrium with the GSH pool. GSH is involved in different ways in ROS buffering. As a result, ROS production triggers an increase of the GS-SG pool, which will be partially reduced to GSH by using rxYFP as an electron donor. The resulting oxidized rxYFP will lose its fluorescence properties. GSH (glutathione), PRX (Peroxiredoxins), TRX (thioredoxin) and GRX (glutaredoxin).

Figure S2

Figure S2: Overexpression of KatG prevents H$_2$O$_2$ chemotaxis. Percentage of WT and *katG*$^+$ SW

cells remaining in the vicinity of the H$_2$O$_2$ plug after 25 min. An arbitrary threshold of 10% flight

(corresponding to 90% of cells remaining in the vicinity of the H$_2$O$_2$ plug) is indicated in red. Mean +/-

s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$,

*** $p < 0.001$).

Figure S3

Figure S3: McpA, McpB and McpR are potentially involved in Cu chemotaxis. Percentage of WT, Δ*mcpA*, Δ*mcpA* overexpressing McpA, Δ*mcpB*, Δ*mcpB* overexpressing McpB, Δ*mcpR* and Δ*mcpR* overexpressing McpR SW cells remaining in the vicinity of the Cu plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Cu plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure S4

Figure S4: The Δ*mcpA* mutant is defective in chemotaxis. Chemotaxis-driven swarming pattern of the WT and Δ*mcpA* strains.

Figure S5

A.

B.

Figure S5: The chemotactic machinery is functional in the Δ*mcpR* mutant. A. Chemotaxis-driven swarming pattern of the WT and Δ*mcpR* strains. **B.** Percentage of WT and Δ*mcpR* SW cells remaining in the vicinity of the Zn plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Zn plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using a *t-test* (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure S6

Figure S6: The Cu-induced oxidative stress is not affected in the Δ*mcpR* mutant. Relative intensity of fluorescence of the WT and Δ*mcpR* SW cells overexpressing the rxYFP biosensor and exposed to 175 μM Cu for 20 min. Mean +/- s.d., at least 3 biological replicates. *p- values* were calculated using a *t-test* (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure S7

Figure S7: Representative fluorescence spectra of purified McpR. Representative intrinsic fluorescence spectra upon 280 nm excitation of purified McpR (3.58 µM) incubated with Cu(II) (top) and Cu(I) (bottom) at 1:1, 1:10 and 1:100 ratios.

Figure S8

Figure S8: Zn, Cd, Mn and Ni do not directly bind to McpR. Relative intrinsic fluorescence of purified McpR (3.58 µM) incubated with $ZnSO_4$, $CdSO_4$, $MnSO_4$ and $NiSO_4$ at 1:10 and 1:100 ratios. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA and were not significant ($p > 0.05$).

Figure S9

Figure S9: The McpR$_{H99A}$ mutant is still proficient for Cd chemotaxis. Percentage of WT, Δ*mcpR* and Δ*mcpR* overexpressing McpR$_{H99A}$ SW cells remaining in the vicinity of the Cd plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the Cd plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure S10

Figure S10: The McpR$_{C75S}$ mutant is still proficient for H$_2$O$_2$ chemotaxis. Percentage of WT, Δ*mcpR* and Δ*mcpR* overexpressing McpR$_{C75S}$ SW cells remaining in the vicinity of the H$_2$O$_2$ plug after 25 min. An arbitrary threshold of 10% flight (corresponding to 90% of cells remaining in the vicinity of the H$_2$O$_2$ plug) is indicated in red. Mean +/- s.d., at least 3 biological replicates, except for Δ*mcpR* overexpressing McpR$_{C75S}$ (2 biological replicates). *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Table S1

Strains	Relevant genotype or description	References
C. crescentus		
WT	CB15N (or NA1000), synchronizable variant strain of CB15	(Evinger and Agabian, 1977)
WTrxYFP	WT strain carrying the *rxYFP* gene on the pJS14 under the control of the lac promoter; CmR	This study
Δ*cheAI*	Knockout strain for *cheAI* gene (CCNA_XXX)	This study
Δ*A*	Knockout strain for *mcpA* gene	This study
Δ*A+A*	Knockout strain for *mcpA* gene carrying a copy of *mcpA* on the pMR10 under the control of the lac promoter; KanR	This study
Δ*B*	Knockout strain for *mcpB* gene	This study
Δ*B+B*	Knockout strain for *mcpB* gene carrying a copy of *mcpB* on the pMR10 under the control of the lac promoter; KanR	This study
Δ*C*	Knockout strain for *mcpC* gene	This study
Δ*D*	Knockout strain for *mcpD* gene	This study
Δ*E*	Knockout strain for *mcpE* gene	This study
Δ*F*	Knockout strain for *mcpF* gene	This study
Δ*G*	Knockout strain for *mcpG* gene	This study
Δ*H*	Knockout strain for *mcpH* gene	This study
Δ*I*	Knockout strain for *mcpI* gene	This study
Δ*J*	Knockout strain for *mcpJ* gene	This study
Δ*K*	Knockout strain for *mcpK* gene	This study
Δ*L*	Knockout strain for *mcpL* gene	This study
Δ*M*	Knockout strain for *mcpM* gene	This study
Δ*N*	Knockout strain for *mcpN* gene	This study
Δ*O*	Knockout strain for *mcpO* gene	This study
Δ*P*	Knockout strain for *mcpP* gene	This study
Δ*Q*	Knockout strain for *mcpQ* gene	This study
Δ*R*	Knockout strain for *mcpR* gene	This study
Δ*RrxYFP*	Knockout strain for *mcpR* gene carrying the *rxYFP* gene on the pJS14 under the control of the lac promoter; CmR	This study
Δ*R+R*	Knockout strain for *mcpR* gene carrying a copy of *mcpR* on the pMR10 under the control of the lac promoter; KanR	This study
Δ*R+RH99A*	Knockout strain for *mcpR* gene carrying a copy of *mcpR* with the mutation H99A on the pMR10 under the control of the lac promoter; KanR	This study
Δ*R+RC75S*	Knockout strain for *mcpR* gene carrying a copy of *mcpR* with the mutation C75S on the pMR10 under the control of the lac promoter; KanR	This study
Δ*S*	Knockout strain for *mcpS* gene	This study
sodB+	WT strain carrying a copy of *sodB* on the pMR10 under the control of the lac promoter; KanR	This study
katG+	WT strain carrying a copy of *katG* on the pMR10 under the control of the lac promoter; KanR	This study
WTrxYFP	WT strain carrying the *rxYFP* gene on the pJS14 under the control of the lac promoter; CmR	This study
E. coli		
S17-1	RP4-2, Tc::Mu, KM-Tn7, for plasmid mobilization	
BL21	BL21 *E. coli* strain used for overexpression of proteins	
Plasmids	**Relevant genotype or description**	
pNPTS138	mobRP4+ ori-R6K sacB; integrative vector in *C. crescentus* for in-frame deletions; KanR	
pMR10	DH10B strain carrying the pMR10 low copy number replicative cloning vector in *E. coli* and *C. crescentus*; KanR	
pJS14	DH10B strain carrying the pJS14 medium copy number replicative cloning vector in *E. coli* and *C. crescentus*; CmR	
pET28a	BL21 strain carrying the pET28a high copy number replicative cloning vector in *E. coli*; KanR	

Table S1: Strains and plasmids