A cytoplasmic chemoreceptor and Reactive Oxygen Species mediate bacterial chemotaxis to copper

Gwennaëlle Louis[1], Pauline Cherry[1], Catherine Michaux[2], Sophie Rahuel-Clermont[3], Marc Dieu[4], Françoise Tilquin[1], Laurens Maertens[1,5], Rob Van Houdt[5], Patricia Renard[4], Eric Perpete[2] and Jean-Yves Matroule[1*]

[1]Research Unit in Biology of Microorganisms (URBM), Department of Biology, Namur Research Institute for Life Sciences (NARILIS), University of Namur, Namur, Belgium.

[2] Laboratoire de Chimie Physique des Biomolécules, Namur Research Institute for Life Sciences (NARILIS) and Namur Institute of Structured Matter (NISM), University of Namur, Rue de Bruxelles, 61, 5000 Namur, Belgium.

[3] Université de Lorraine, CNRS, IMoPA, F-54000 Nancy, France.

[4]MaSUN, Mass Spectrometry Facility, University of Namur, Namur, Belgium.

[5]Microbiology Unit, Interdisciplinary Biosciences, Belgian Nuclear Research Centre (SCK CEN), 2400 Mol, Belgium.

*Corresponding author. Mailing address: Research Unit in Biology of Microorganisms (URBM), 61 Rue de Bruxelles 5000 Namur, Belgium. Phone: (32)81-72-42-45. E-mail: jean-yves.matroule@unamur.be

Running title

Bacterial chemotaxis to copper and oxidative stress

Keywords

Chemotaxis, bacteria, copper, reactive oxygen species (ROS), stress response

Abstract

Chemotaxis is a widespread strategy used by unicellular and multicellular living organisms to maintain their fitness in stressful environments. We previously showed that bacteria can trigger a negative chemotactic response to a copper (Cu)-rich environment. Cu ions toxicity on bacterial cell physiology has been mainly linked to mismetallation events and ROS production, although the precise role of Cu-generated ROS remains largely debated.

Here, we found that the cytoplasmic Cu ions content mirrors variations of the extracellular Cu ions concentration and triggers a dose-dependent oxidative stress, which can be abrogated by superoxide dismutase and catalase overexpression. The inhibition of ROS production in the cytoplasm not only improves bacterial growth but also impedes Cu-chemotaxis, indicating that ROS derived from cytoplasmic Cu ions mediate the control of bacterial chemotaxis to Cu.

We also identified the Cu chemoreceptor McpR, which binds Cu ions with low affinity, suggesting a labile interaction. In addition, we demonstrate that the cysteine 75 and histidine 99 within the McpR sensor domain are key residues in Cu chemotaxis and Cu coordination. Finally, we

discovered that *in vitro* both Cu(I) and Cu(II) ions modulate McpR conformation in a distinct manner. Overall, our study provides mechanistic insights on a redox-based control of Cu chemotaxis, indicating that the cellular redox status can play a key role in bacterial chemotaxis.

Introduction

Cell motility is widely distributed across the three domains of life. For instance, in animals it is involved in fertilization, and in neutrophils recruitment to an inflammatory area. It also allows archaea and bacteria to seek nutrients in scarce environments, and to flee from toxic compounds while it supports to some extent host tropism in the frame of symbiotic relationships.

Most of the time, cell motility is driven by chemical gradients and is therefore named chemotaxis. In flagellated bacteria, positive (attraction) and negative (repulsion) chemotactic responses imply stochastic direction changes triggered by the inversion of flagellar rotation (1,2). This switch is caused by the binding of the phosphorylated response regulator, CheY~P, to the flagellar motor. Spontaneous or phosphatase-mediated CheY~P dephosphorylation by the CheZ phosphatase or by a member of the CheC/CheX/FliY phosphatase family, triggers CheY release from the flagellar motor (3-5). The release of CheY promotes a straight bacterial swimming pattern until CheY is phosphorylated again and triggers a new direction change. CheY phosphorylation relies on the histidine kinase CheA, which is recruited and autophosphorylated within a chemotaxis cluster composed of cytoplasmic or inner membrane-spanning methyl-accepting chemotaxis proteins (Mcp), also named chemoreceptors (6). Chemoreceptors are classically composed of a N-terminal sensing domain, trans-membrane (TM) domains (which are absent in cytoplasmic chemoreceptors), a HAMP domain and the C-terminal (Mcp) signaling domain. Depending on their cellular topology, chemoreceptors monitor chemical variations in the cytosol or in the periplasm *via* their N-terminal sensing domain. Upon substrate binding to their sensing domain, chemoreceptors undergo a conformational change leading to the increase (repellent response) or the decrease (attractant response) of CheA histidine kinase activity (7,8). In *E. coli* Tar and Tsr

chemoreceptors, this conformational change consists in a 2 Å shift of one of the TM helices (TM2), causing a "piston-like" movement in the direction of the membrane that impacts the HAMP domain (9). The methylation of the Mcps by the CheR methyltransferase results in an increase of CheA activity and their demethylation by the CheA-activated CheB methylesterase leads to the decrease of CheA activity (9). This adaptation response sets back the Mcps in a pre-stimulus state.

Copper (Cu) has been used for centuries as an antimicrobial and antifouling strategy in medicine and in various anthropogenic activities including agriculture and industry. Cu toxicity in solution mainly results from its high redox potential causing the displacement of native metallic ions from metalloproteins, and the disruption of Fe-S clusters (10). Cu ability to generate reactive oxygen species (ROS) has been shown *in vitro* (11) but the role of these ROS *in vivo* yet remains unclear. In a previous study, we identified an original bimodal strategy to cope with Cu stress where the flagellated morphotype (swarmer cell) of *Caulobacter crescentus* triggers a prompt negative chemotaxis from the Cu source (12). In addition to the negative Cu-chemotaxis, *C. crescentus* exhibits a positive chemotactic behavior toward xylose (13) and O_2 (14) but no dedicated Mcp could be identified so far.

C. crescentus genome harbors two chemotaxis operons. The major chemotaxis operon encodes McpA and McpB, CheYI and CheYII, CheAI, one CheW, CheRI, CheBI and the CheY-like CleA. This operon also contains the *cheD*, *cheU* and *cheE* genes, which have been hardly described. CheYII and CheBI are likely the major CheY and CheB in *C. crescentus* chemotaxis (15,16). The alternate chemotaxis operon encodes McpG and McpK, two CheY, CheAII, one CheW, CheBII, and CheRII. The other *cheY* genes and the third *cheW* are dispersed within the genome. Only the first chemotaxis operon seems to be essential for chemotaxis while both operons are involved in

biofilm formation and holdfast production, by regulating the expression of the *hfiA* holdfast inhibitor (16,17).

No CheZ homolog has been found in *C. crescentus* genome but the multiple CheY are proposed to act as phosphate sinks to allow the termination process, such as in *Rhizobium meliloti (18)*. Five of the CheY have been redefined as Cle proteins (CheY-like c-di-GMP effectors) and renamed CleA to CleE. CleA and CleD could compete with CheYII to prevent the rotational switch of the flagellum and to promote straight run (15). The *C. crescentus* MCPs array exhibits a higher-ordered hexagonal structure constituted of approximatively one to two thousand MCPs per array (19). Among the *C. crescentus* MCPs, only the inner membrane McpA and the cytoplasmic McpB encoded by the major chemotaxis operon were described as polar chemoreceptors, although their exact function remains unknown (20).

In the present study, we aimed to decipher the sensing mechanism underlying *C. crescentus* chemotaxis to a toxic Cu-rich environment. We provide evidence that an increased intracellular Cu concentration triggers a dose-dependent production of reactive oxygen species (ROS), which is a prerequisite to Cu chemotaxis. We also demonstrate that the redox status of the Cu ions bound to the sensor domain of the newly identified McpR modulates McpR conformation *in vitro*. This original redox sensing mechanism may ensure a continuous monitoring of the intracellular Cu status, providing a real-time control of the chemotactic response.

Results

To further dissect the molecular mechanisms underlying the negative Cu-chemotaxis of *Caulobacter crescentus* swarmer (SW) cells (12), we first sought to identify the chemoreceptor involved in Cu chemotaxis. The genome of *C. crescentus* is predicted to encode 19 chemoreceptors (named McpA to McpS) based on the presence of an Mcp domain binding CheA/CheW (21). No canonical Cu ions-binding site, including the CXXC motif or the βαββαβ fold (22), could be found by running manual and JPred4-based analyses of the 19 chemoreceptor protein sequences (23). Therefore, we proceeded to single in-frame deletions of the 19 *mcp* genes, and their respective chemotactic behavior to Cu ions was assessed by using the previously designed Live Chemotaxis Imaging (LCI) (12). Most of the Δ*mcp* mutants displayed a WT-like behavior with a flight percentage after 25 min varying between 15% and 51%. Only Δ*mcpA*, Δ*mcpB* and Δ*mcpR* mutants showed a significant decrease of Cu-chemotaxis relative to the WT strain (**Fig. 1A**), suggesting that these chemoreceptors might be involved in Cu-chemotaxis. The Δ*mcpA*, Δ*mcpB* and Δ*mcpR* mutants could be complemented by the expression of an ectopic version of *mcpA*, *mcpB* and *mcpR* on a low copy plasmid, respectively, ruling out any polar effect of the deletions (**Fig. S1**). Owing to the position of the *mcpA* gene within the major chemotaxis cluster (16), McpA is likely involved in chemotaxis to a less specific extent. Consistent with this hypothesis, the Δ*mcpA* mutant displays a reduced motility halo in a Cu ions-free swarming assay, while the Δ*mcpB* and Δ*mcpR* mutants exhibit a WT-like chemotaxis profile in the same condition (**Fig. S2A**). The presence of the terminal sequence of the CheB/CheR docking site in McpB (absent from McpR) led us to the idea that McpB might be necessary for the adaptive response as in *E. coli* (24), which

will not be considered in this study. Therefore, we decided to determine how McpR modulates Cu-chemotaxis.

The ability of the Δ*mcpR* mutant to form a motility halo in a Cu ions-free swarming assay (**Fig. S2A**) and to sustain negative chemotaxis to other chemicals, such as zinc (Zn) ions (**Fig. S2B**) indicate that the loss of Cu-chemotaxis in the Δ*mcpR* mutant does not result from a loss of motility and/or a disruption of the general chemotactic machinery.

McpR harbors the canonical domain organization found in chemoreceptors including a N-terminal sensing domain followed a HAMP domain and a C-terminal Mcp domain. McpR sequence analysis using the TMHMM 2.0 (25) and SignalIP 6.0 (26) tools did not reveal any McpR transmembrane domain and signal peptide respectively, arguing for a cytoplasmic localization for McpR, and therefore a sensing of the cytoplasmic Cu content.

Chemoreceptors can sense attractants or repellents in a direct or indirect manner *via* their sensing domain (27). The 3D structure prediction of McpR by Alphafold2 (28,29) highlights a cysteine (C75) in the vicinity of a histidine (H99) in the McpR sensing domain (**Fig. 1B**). Cysteine and histidine residues have often been described as key residues in Cu coordination (30), suggesting that McpR may bind Cu ions directly.

Therefore, we used intrinsic fluorescence spectroscopy to assess the direct binding of Cu to the purified McpR. Upon the addition of either Cu(I) or Cu(II) ions, the intrinsic McpR fluorescence at 346 nm is quenched in a dose-dependent manner, reflecting a modification of the McpR aromatic residues environment (**Fig. 1C**). The dissociation constants (*K*d) (see Materials and Methods for calculation method) of 48 µM +/- 10 µM for Cu(I) ions and 37 µM +/- 9 µM for Cu(II) ions attest

to a similar affinity of both ionic forms for McpR. However, the micromolar range of Cu(I) ions and Cu(II) ions Kds exceeds from several logs the Kd of well-known cuproteins (31,32), arguing for a rather labile interaction between McpR and Cu. Nevertheless, McpR seems to preferentially bind Cu ions since Zn, cadmium (Cd), manganese (Mn) and nickel (Ni) ions had no effect on purified McpR intrinsic fluorescence (**Fig. S3**).

To determine whether the cytoplasmic Cu ions content could reflect variations of the extracellular Cu ions concentration when the SW cell swims away from a Cu ions source, we measured by ICP-OES the Cu ions content in cytoplasmic fractions from SW cells subjected to increasing Cu ions concentrations. Although the measurement cannot discriminate between free and complexed Cu ions, Cu(I) and Cu(II), we measured a dose-dependent accumulation of Cu ions in the cytoplasm (**Fig. 1D**). This observation is in accordance with previous data from our lab, showing that an artificial decrease of Cu ions content *via* the overexpression of the PcoB efflux pump impeded Cu-chemotaxis (12). Together, these data suggest that the sensing of cytoplasmic Cu ions by McpR could be used as a proxy by the SW cells to mirror variations of the concentration of Cu ions in *C. crescentus* environment to control Cu-chemotaxis.

The transient maintenance of a high intracellular Cu ions content in the SW cells may potentially lead to an oxidative stress *via* the production of superoxide anion ($O_2^{\bullet-}$), hydrogen peroxide (H_2O_2) and hydroxyl radical ($OH^{\bullet}$) (**Fig. S4A**). In line with this idea, the relative fluorescence intensity of the ROS sensitive turn-on CellROX Deep Red probe is significantly increased in SW cells exposed to Cu ions, and H_2O_2, the latter being used as a positive control (**Fig. 2A**). Accordingly, the fluorescence of the cytoplasmic turn-off rxYFP redox biosensor is partially quenched under the same conditions (**Fig. 2B**), mirroring an imbalance of the cytoplasmic GSH-GSSG pool upon

oxidative stress (**Fig. S4B**). These findings are reinforced by the differential expression of genes involved in oxidative stress response upon Cu stress, including those encoding the catalase KatG and the superoxide dismutases SodA and SodB (33). As observed for the cytoplasmic Cu ions content (**Fig. 1D**), the amount of ROS is increased when the SW cells are exposed to increasing extracellular Cu ions concentrations, indicating that the cytoplasmic ROS level mirrors the cytoplasmic Cu ions content (**Fig. 2C**).

We envisioned that boosting the antioxidant defense system would limit ROS production by Cu ions. Therefore, we ectopically expressed the *sodB* or the *katG* genes in the WT background from a low copy plasmid under the control of the strong and constitutive *lac* promoter. The resulting *sodB*$^+$ and *katG*$^+$ strains prevent the increase of the CellROX Deep Red fluorescence upon SW cells exposure to Cu ions, reflecting a potentiation of antioxidant defenses (**Fig. 2D**). In support of this data, the growth of these two optimized *sodB*$^+$ and *katG*$^+$ strains is improved under moderate Cu stress (**Fig. 2E**), arguing for a role of $O_2^{\bullet-}$ and H_2O_2 in Cu toxicity.

Interestingly, the *sodB*$^+$ and *katG*$^+$ strains are impeded in Cu-chemotaxis similarly to the Δ*cheAI* mutant (**Fig. 2F**, **SI Appendix and Fig. S5**), indicating that the ROS resulting from the accumulation of cellular Cu ions play a role in Cu-chemotaxis. It is worth mentioning that the loss of Cu-chemotaxis in the Δ*mcpR* mutant is not due to a loss of ROS production under Cu stress since the Δ*mcpR* mutant exhibits a similar decrease of the rxYFP fluorescence to the WT strain upon Cu ions exposure (**Fig. S6**). These data support a major role of the ROS resulting from Cu ions accumulation in Cu-chemotaxis and indicate that Cu ions binding to McpR is solely not sufficient to sustain Cu-chemotaxis.

Owing to the oxidative conditions resulting from Cu ions accumulation in the cytosol (**Fig. 2A-C**), we sought to assess by LC/MS the oxidation status of McpR *in vitro* after incubating the purified McpR with Cu(II) ions or Cu(I) ions +/- H_2O_2. Only the Cu(I) ions + H_2O_2 condition, and to a lesser extent Cu(I) ions, led to a significant oxidation of the McpR sensor domain (residues 1-161), and more specifically of the H99 residue (**Fig. 3A, left panel**). In this condition, about 30% of the H99 residues acquired an additional oxygen atom, most likely yielding 2-oxoHis. Metal catalyzed oxidation (MCO) of histidines via the generation of $OH^\bullet$ has been previously described in the *Bacillus subtilis* PerR repressor (34), in yeast SOD1 (35), and in amyloid beta (Aβ) peptide (36). The exact role of this modification remains unknown and no redox cycling involving 2-oxoHis has been described so far, questioning the relevance of H99 oxidation in Cu/ROS sensing. MCO mostly targets the metal-binding residues in a caged process (37), suggesting that H99 may coordinate Cu. To test this hypothesis, we purified a McpR$_{H99A}$ mutant protein and calculated the Kd values of the McpR$_{H99A}$/Cu(I) ions and McpR$_{H99A}$/Cu(II) ions complexes (**Fig. 3B, Fig. S7A for the fitting curves**). H99A mutation triggers a 9.4- and 5.8-fold increase of Cu(I) ions and Cu(II) ions Kd, respectively, relative to McpR, suggesting a key role of H99 in Cu(I) ions and Cu(II) ions coordination. The important variability associated with the McpR$_{H99A}$ Kd values likely reflects a potential instability of this mutant in solution upon Cu(I) ions and Cu(II) ions exposure, resulting in the presence of several subpopulations of McpR$_{H99A.}$

To assess the potential implication of the neighboring C75 residue in Cu ions coordination, we purified the McpR$_{C75S}$ mutant protein and calculated the Kd values of the McpR$_{C75S}$/Cu(I) ions and McpR$_{C75S}$/Cu(II) ions complexes. As for the H99A mutation, the C75S mutation also impeded Cu(I) ions binding as attested by a 3.5-fold Kd increase. Cu(II) ions binding to the McpR$_{C75S}$

mutant remained unchanged (**Fig. 3B, Fig. S7B for the fitting curves**), suggesting that McpR C75 is required to coordinate Cu(I) ions but not Cu(II) ions. Consistent with this observation, *in vitro* H99 oxidation by Cu(I) ions +/- H_2O_2 is fully abolished in the McpR$_{C75S}$ variant (**Fig. 3A, right panel**). The *K*d values calculated for the McpR$_{C75S}$ and McpR$_{H99A}$ variants are in agreement with the known selective binding of Cu(I) and Cu(II) ions to cysteine and histidine residues (30,38).

To determine whether Cu coordination by McpR H99 and C75 residues is essential to Cu-chemotaxis, we decided to express an ectopic *mcpR$_{H99A}$* or *mcpR$_{C75S}$* allele in the Δ*mcpR* mutant and to monitor the chemotactic behaviour of both strains to Cu ions. None of the mutant allele was able to complement the Cu-chemotaxis defect of the Δ*mcpR* mutant (**Fig. 3C**). One could argue that the McpR$_{H99A}$ and McpR$_{C75S}$ variants are unstable *in vivo*, explaining the Δ*mcpR* mutant phenotype. In accordance with the pleiotropic sensing ability of some Mcps, we discovered that McpR is also playing a key role in cadmium (Cd)-chemotaxis (**Fig. S8**). Interestingly, the *mcpR$_{H99A}$* and *mcpR$_{C75S}$* mutant strains were not impeded in Cd-chemotaxis (**Fig. S8**). This latter observation attests the integrity of the McpR$_{H99A}$ and McpR$_{C75S}$ variants *in vivo* and suggests that Cd sensing by McpR does not rely on the H99 and C75 residues.

These data led us to the idea that Cu(I) ions coordination by McpR H99 and C75 residues is decisive for McpR-mediated Cu chemotaxis. Given the structural changes elicited by the binding of a ligand to a cognate chemoreceptor (39), we envisioned that Cu(I) ions binding to McpR could modulate its conformation. To test this idea, Far-UV circular dichroism (CD) spectroscopy was used to monitor the impact of Cu(I) ions and Cu(II) ions on McpR secondary structure *in vitro*. Accordingly, Cu(I) ions has a more significant impact than Cu(II) ions on McpR secondary structure (**Fig. 3D**), increasing the α-helix percentage from 23% to 44%.

Based on our findings, we propose that Cu exposure triggers a dose-dependent Cu accumulation within the cytoplasm triggering a dose-dependent production of ROS that will sustain Cu chemotaxis. The increase in cytoplasmic Cu ions concentration will likely favor a rather labile Cu ions binding to the McpR sensor domain. Cu(I) ions coordination by McpR C75 and H99 residues triggers McpR conformational change. Interestingly, Cu(II) ions seem to preferentially bind McpR H99 residue and does not modify the apo form-like conformation. These observations suggest that a redox cycling of the McpR-bound Cu atom could modulate McpR conformation (**Fig. 4**).

Discussion

The role of ROS in Cu toxicity and Cu stress response remains largely unknown in bacteria. We sought to address this gap in the frame of negative chemotaxis to toxic Cu in the dimorphic bacterial model *C. crescentus*.

Our study identified the first Cu sensing chemoreceptor hereafter named McpR. McpR is devoid of any transmembrane domains or signal peptide and is therefore predicted to localize in the cytoplasm. Cytoplasmic chemoreceptors have been shown to monitor the cellular metabolic state (40), whereas inner membrane chemoreceptors usually sense the periplasmic environment, as a proxy for extracellular changes. The cytoplasmic Cu content varies in a dose-dependent manner when the SW cells are exposed to Cu, suggesting that the sensing of cytoplasmic Cu could provide a sensitive monitoring of subtle variations of Cu concentrations when the SW cell swims down a Cu gradient.

In a previous study, we showed that the maintenance of a high cellular Cu concentration is critical for Cu chemotaxis (12). Consistent with this observation, we provide evidence that McpR binds Cu with low affinity. Although aspecific Cu binding to non-cognate residues cannot be ruled out, the

significant difference in *Kd* values measured between the WT McpR and the McpR$_{H99A}$ and McpR$_{C75S}$ variants argue for a Cu binding to these residues. Considering that Cu binding to McpR is a prequisite for Cu chemotaxis, one could propose that when the SW cell is sufficiently far from the Cu source, the resulting low cytoplasmic Cu content impedes Cu binding to McpR and in turn the chemotactic response.

We also demonstrate that the cellular Cu accumulation triggers a dose-dependent production of ROS. This dose dependency in ROS production argues that the increase of extracellular Cu ions content triggers an increase of the cytoplasmic Cu ions labile pool, which is likely involved in the labile Cu binding to McpR. ROS production can be prevented by the overexpression of SodB superoxide dismutase and KatG catalase, suggesting that at least $O_2^{\cdot-}$ and H_2O_2 are generated under Cu stress. The contribution of the ROS derived from cellular Cu ions in Cu toxicity is still largely debated (41,42). Here, we demonstrate that ROS derived from cellular Cu ions not only impede cell growth to some extent, but also play a key role in Cu-chemotaxis. The exact target of ROS in Cu-chemotaxis was not identified in this study, but one could hypothesize that $O_2^{\cdot-}$ is rapidly converted into H_2O_2. The latter would feed a local Fenton-like reaction catalyzed by the Cu(I) atom coordinated by McpR C75 and McpR H99 residues. This would lead to a local $OH^{\cdot}$ production and to the oxidation of McpR-bound Cu(I) into McpR-bound Cu(II). Owing to the reducing cytoplasmic environment, McpR-bound Cu(II) could in turn be reduced by cytoplasmic electron donors such as GSH, NADH and thioredoxin to restore the Cu(I) ionic form. Thereby, the McpR-bound Cu ion would undergo a redox cycling involving the Cu(I) and Cu(II) ionic forms, which would be maintained as long as H_2O_2 is produced by cytoplasmic Cu ions. Cu-generated ROS would therefore act as second messengers in a bacterial stress response.

The modulation of the CheA-CheY two-component system results from a conformational change of the chemoreceptor occurring upon ligand binding (9). We demonstrated here that McpR conformation is modulated by the ionic status of the Cu atom bound to McpR. It should be emphasized that chemoreceptors are usually organized in trimers of dimers and assembled in large arrays in cells (43). Therefore, it is possible that Cu coordination by the McpR C75 and H99 residues within the McpR array implies two McpR monomers from one or two distinct dimers. Nevertheless, McpR conformational change likely regulates the downstream effectors such as CheAI and the resulting chemotactic response.

The diversity of chemoreceptors produced by a single bacterial species varies from 5 to 60 chemoreceptors and is often correlated to its lifestyle (40). Nevertheless, chemoreceptors are not always specific to one single substrate and can also perform indirect sensing *via* a substrate transporter (27). For instance, the *E. coli* Tar chemoreceptor directly senses aspartate and performs an indirect sensing of maltose through a periplasmic maltose binding protein (44).

The extrapolation of our findings to other motile bacterial species based on sequences analyses is complicated by (i) the large diversity of Cu coordination mechanisms and thereby the low predictability of Cu-binding sites by *in silico* analyses and (ii) the low conservation of the Mcp sensor domains. Nevertheless, it is very likely that the use of ROS as second messengers is widely conserved in bacterial response to Cu.

Experimental procedures

Bacterial strains, plasmids and growth conditions

Caulobacter crescentus (NA1000 strain) was grown at 30°C in Peptone Yeast Extract (PYE) medium (45) with 5 μg/ml kanamycin, 1 µg/ml chloramphenicol and/or an appropriate concentration of $CuSO_4.5H_2O$ when required. Cultures in exponential phase were used for all experiments. Strains, plasmids and primers used in this study are listed in Table S2, and the strategies for their construction are available upon request. SW cells isolation was achieved according to (46). Briefly, the bacteria were collected at an $O.D._{660\ nm}$ of 0.4 and centrifuged in Ludox LS colloidal silica gradient. After the centrifugation, SW are separated from the other cellular forms (ST and PD) based on a density difference. The SW cells were collected and washed several times with PBS prior the experiments.

Growth curve measurements

C. crescentus exponential phase cultures were diluted in PYE medium to a final $O.D._{660\ nm}$ of 0.05 and inoculated in 96-well plates with appropriate $CuSO_4$ concentration when required. Bacteria were then grown for 24 h at 30°C under continuous shaking in an Epoch 2 Microplate Spectrophotometer from BioTek (Santa Clara, CA, USA). The absorbance at $O.D._{660\ nm}$ was measured every 10 min.

Motility assay

C. crescentus cultures grown in PYE were diluted to an $O.D._{660\ nm}$ of 0.4 and inoculated with sterile toothpicks on a PYE swarming plate (PYE + 0.25% agar). The plates were incubated at 30°C for 3 days, then imaged.

Determination of cytoplasmic Cu content

C. crescentus cells were fixed for 20 min in 2% paraformaldehyde at 4°C and then washed three times with an ice-cold wash buffer (10 mM Tris-HCl pH 6.8, 100 μM EDTA). The cells were then incubated for 10 min with zwitterionic detergent (0.0125 % Zwittergent 3-14 detergent, 200 mM Tris-HCl pH 7.6), to disrupt the outer membrane but not the inner membrane. A final centrifugation at 14,500 x g for 15 min separated the spheroplasts (cytoplasm and inner membrane fraction) from the periplasm (and outer membrane fraction). The spheroplasts were lysed under 2.4 kbar by using a cell disrupter (Cell Disruption System, one shot model, Constant). Cells debris were removed by centrifugation at 10,000 x g for 10 min. and cell lysates were diluted in HNO_3 1 %. Samples were finally analyzed by inductively coupled plasma - optical emission spectrometry (ICP-OES) with an Optima 8000 ICP-OES from PerkinElmer. Cellular Cu concentration were calculated as described in (12). Briefly, 10 μl of the samples after fixation were diluted 1:100 in PBS and counted with a BD FACSVerseTM flow cytometer. Data were analyzed with the BD FACSuite V1.0.5 software. A ratio between the number of bacteria and the OD_{660nm} was determined. The combination of both flow cytometry and ICP-OES analyses allows to determine the number of Cu atoms in one spheroplast. Cu concentration was calculated using the following formula;

$$\frac{\dfrac{Amount\ of\ Cu\ (mg)}{Molecular\ weight\ of\ Cu\ (63,546\ \frac{mg}{L})} \times Avogadro\ constant\ (6.022\ \times 10^{23})}{Number\ of\ bacteria}$$

Oxidative stress measurement with the CellROX Deep Red probe

An exponential phase culture of *C. crescentus* was synchronized to isolate the SW cells. The isolated SW cells were diluted to a final OD_{660nm} of 0.2 and incubated for 30 min at 30°C under moderate shaking with 5 μM of CellROX Deep Red probe (ThermoFisher) and appropriate

concentrations of $CuSO_4$ or H_2O_2. The bacteria were then washed three times with 20 mM Phosphate Buffer (12.5 mM Na_2HPO_4, 8 mM KH_2PO_4, pH 7) to remove the non-internalized probe. The bacteria were finally transferred into a black bottom 96-wells plate and the fluorescence was monitored at Em/Ex of 640/690 nm with a SpectraMax iD3 (Molecular Devices, San Jose, CA, USA). The relative fluorescence intensity was determined by defining 100% as the fluorescence from control condition.

Oxidative stress measurement with the rxYFP biosensor

The SW cells were synchronized from an exponential phase culture overexpressing the rxYFP biosensor (47). The SW cells were diluted to a final OD_{660nm} of 0.4 in a black bottom 96-well plate and mixed with different concentrations of $CuSO_4$ and H_2O_2. The rxYFP fluorescence was then monitored every minute for 20 min at Em/Ex of 510/556 nm with a SpectraMax iD3 (Molecular Devices, San Jose, CA, USA). The relative fluorescence intensity was determined by defining 100% as the initial measured fluorescence at time 0.

Live Chemotaxis Imaging

The live chemotaxis imaging (LCI) assay was adapted from (12). Briefly, chemotaxis devices were made by casting solubilized 10:1 polydimethylsiloxane (PDMS Slygard 184, Dow Corning) in a small glass pot (d = 50 mm; h = 30 mm; glassware from Lenz laborglass instrument) in which coverslips were initially installed to mold the future bacterial chamber. The PDMS was left overnight, allowing degassing and curing. The different channels were drilled into the PDMS devices. A microscope slide and the PDMS cube were washed successively with acetone (only the slide), isopropanol, methanol and rinsed with milli-Q H_2O. The materials were blown dry between each wash. The PDMS cube was sealed to the slide by pressure. Eight microliters of melted 1.5% agarose H_2O with appropriate

concentration of $CuSO_4$, $ZnSO_4$ or $CdSO_4$ were loaded into the chamber through the external inlet channels to generate the plug. One hundred and fifty microliters of isolated SW cells were in turn injected into the bacterial chamber through the central inlet channel. Images were collected every minute for 25 min. in one focal plan (near the liquid surface) in the vicinity of the plug with a Nikon Eclipse Ti2-E inverted microscope equipped with a Nikon 20X/0.5 Ph2 objective and with a Hamamatsu C13440 digital camera. Quantitative analysis of the time-lapse images was performed with MicrobeJ (48). The number of bacteria in a same focal plane was determined at each time point; the number of bacteria at the beginning of the experiment is considered as 100%.

Protein expression and purification

A BL21 pET28a-McpR$_{His}$ exponential phase culture was incubated for 3 h under shacking at 37°C with 1 mM Isopropyl-β-D-thiogalactopyranoside (IPTG) and then harvested by centrifugation for 30 min at 4,000 x g at 4°C. McpR$_{His}$ induction was verified by 12 % Tricine–sodium dodecyl sulfate–polyacrylamide gel electrophoresis (Tricine–SDS–PAGE), followed by Coomassie Blue staining and by Western-Blot using an anti-His antibody (MBL, PM032). To purify McpR$_{His}$, the IPTG-induced culture was washed with binding buffer (20 mM Tris-HCl (pH 8.0), 500 mM NaCl, 10% glycerol, 10 mM $MgCl_2$) supplemented with 400 mg lysozyme (Sigma), complete EDTA-free protease cocktail inhibitor (Roche), 5 mg DNase I (Roche) and 0.4 g SDS and centrifuged for 15 min at 9,000 x g at 4°C. Bacterial cells were lysed at 2.4 kbar using a Cell disruptor (Cell Disruption System – one shot model, Constant). As McpR$_{His}$ was mainly forming inclusion bodies, the pellet fraction was resuspended for 1 h under shaking at 4°C in the binding buffer containing 6 M guanidine. The solubilized inclusion bodies were centrifuged for 30 min at 9,000 x g at 4 °C. The collected supernatant was dialyzed overnight in a 0.1 M phosphate buffer

(12.5 mM Na_2HPO_4, 8 mM KH_2PO_4, pH 8)) to remove guanidine, allowing the refolding of McpR. The same methodology was applied for the C75S and H99A mutants.

McpR protein concentration was determined (i) by measuring the absorbance of the sample at 280 nm with the SpectraMax iD3 (Molecular Devices, San Jose, CA, USA) or (ii) with a Bradford protein assay.

Fluorescence quenching by Cu and other metal ions

Purified McpR at a concentration of 3.58 μM was incubated with different ratios of $CuSO_4$, Tetrakis(acetonitrile)Cu(I) hexafluorophosphate, $ZnSO_4$, $CdSO_4$, $MnSO_4$ and $NiSO_4$ in black bottom 96-wells plates. The fluorescence emission spectra were recorded upon 280 nm excitation with the SpectraMax iD3.

***K*d determination**

The titration of McpR by Cu(I) or Cu(II) monitored by the protein intrinsic fluorescence intensity quenching was fitted to a single-site hyperbolic model by non-linear regression using the Scidavis 2.7 software to determine the affinity constants, according to the following equation:

Fluorescence intensity = F0 + A*[Cu]/(Kd + [Cu]),

with F0 corresponding to the fluorescence intensity of the apo McpR, A to the total fluorescence amplitude change upon Cu binding and *K*d to the affinity dissociation constant. Dissociation constants are reported as the mean value ± standard deviation obtained from a minimum of three independent experiments performed on distinct protein productions.

Identification of McpR oxidation by mass spectrometry

Protein digestion

McpR samples were treated using Filter-aided sample preparation (FASP) using the following protocol. To first wash the filter, 100 µl of formic acid 1 % were placed in each Millipore Microcon 30 MRCFOR030 Ultracel PL-30 before a 15 min centrifugation at 14,100 x g. For each sample, 40 µg of protein adjusted in 150 µl of 8 M urea buffer (8 M urea in 0.1 M Tris buffer at pH 8.5) were placed individually in a column and centrifuged for 15 min at 14,100 x g. The filtrate was discarded, and the columns were washed three times by adding 200 µl of urea buffer followed by a 15 min centrifugation at 14,100 x g. For the reduction step, 100 µl of dithiothreitol (DTT) were added and mixed for 1 min at 400 rpm with a thermomixer before a 15 min incubation at 24 °C. The samples were then centrifugated for 15 min at 14,100 x g, the filtrate was discarded, and the filter was washed by adding 100 µl of urea buffer before another 15 min centrifugation at 14,100 x g. An alkylation step was performed by adding 100 µl of iodoacetamide ((IAA), in urea buffer) in the column and mixing for 1 min at 400 rpm in the dark followed by a 20 min incubation in the dark and a 10 min centrifugation at 14,100 x g. To remove the excess of IAA, 100 µl of urea buffer were added and the samples were centrifuged for 15 min at 14,100 x g. To quench the rest of IAA, 100 µl of DTT were placed on the column, mixed for 1 min at 400 rpm and incubated for 15 min at 24 °C before a 10 min centrifugation at 14,100 x g. To remove the excess of DTT, 100 µl of urea buffer were placed on the column and centrifuged for 15 min at 14,100 x g. The filtrate was discarded, and the column was washed three times by adding 100 µl of 50 mM sodium bicarbonate buffer ((ABC), in ultrapure water) followed by a 10 min centrifugation at 14,100 x g. The remaining 100 µl were kept at the bottom of the column to avoid any evaporation in the column. The digestion process was performed by adding 80 µl of mass spectrometry grade trypsin (1/50 in ABC buffer) in the column and mixed for 1 min at 400 rpm followed by an overnight incubation at 24°C in a water saturated environment. The Microcon columns were placed in a 1.5

ml LoBind tube and centrifuged for 10 min at 14,100 x g. Fourty µl of ABC buffer were placed on the column before a 10 min centrifugation at 14,100 x g. Ten % Trifluoroacetic acid (TFA) in ultrapure water were added into the LoBind Tube to obtain 0.2 % TFA. The samples were dried in a SpeedVac up to 20 µl and transferred into an injection vial.

Mass spectrometry

The digest was analyzed using nano-LC-ESI-MS/MS tims TOF Pro (Bruker, Billerica, MA, USA) coupled with an UHPLC nanoElute (Bruker).

Peptides were separated by nanoUHPLC (nanoElute, Bruker) on a 75 µm ID, 25 cm C18 column with integrated CaptiveSpray insert (Aurora, ionopticks, Melbourne) at a flow rate of 400 nl/min, at 50°C. LC mobile phases A was water with 0.1% formic acid (v/v) and B ACN with 0.1% formic acid (v/v). Samples were loaded directly on the analytical column at a constant pressure of 800 bar. The digest (1 µl) was injected, and the organic content of the mobile phase was increased linearly from 2% B to 15 % in 36 min, from 15 % B to 25% in 19 min, from 25% B to 37 % in 5 min and from 37% B to 95% in 5 min. Data acquisition on the tims TOF Pro was performed using Hystar 5.1 and timsControl 2.0. tims TOF Pro data were acquired using 100 ms TIMS accumulation time, mobility (1/K0) range from 0.6 to 1.6 Vs/cm². Mass-spectrometric analysis were carried out using the parallel accumulation serial fragmentation (PASEF) acquisition method. One MS spectra followed by ten PASEF MSMS spectra per total cycle of 1.1 s.

Data analysis was performed using PEAKS Studio X Pro with ion mobility module (Bioinformatics Solutions Inc., Waterloo, ON). Protein identifications were conducted using PEAKS search engine with 15 ppm as parent mass error tolerance and 0.05 Da as fragment mass

error tolerance. Carbamidomethylation was allowed as fixed modification, oxidation of methionine, histidine, tryptophan, and acetylation (N-term) as variable modification. Enzyme specificity was set to trypsin, and the maximum number of missed cleavages per peptide was set at three. The peak lists were searched against the *E. coli*-K12 (11002 sequences) and *Caulobacter* NA1000 (3863 sequences) from UNIREF 100. Peptide spectrum matches and protein identifications were normalized to less than 1.0% false discovery rate.

The identification results were processed with the PTM profile tool from PEAKS, which provides quantitative information of modified peptides compared with unmodified peptides for the modification sites of the protein across all MS samples (49,50).

Circular dichroism

Purified McpR at a concentration of 8.95 µM was incubated with different $CuSO_4$ and Tetrakis(acetonitrile)Cu(I) hexafluorophosphate concentrations. Measurements were performed with a MOS-500 spectropolarimeter in the far-UV region (190-260 nm), using a 1 mm pathlength cell. Four scans were averaged, and base lines were subtracted. Secondary structure analyses were performed on the circular dichroism data with the BeStSel server.

Supporting information

This article contains supporting information.

Acknowledgements

We would like to thank Jakob R. Winther for providing the rxYFP construct. We are also grateful to Valérie Charles and Carmela Aprile (CMI laboratory, NISM, UNamur) for the ICP measurements and to Virgile Neyman for technical support during the protein purification. Finally, we would like to acknowledge Jean-François Collet and Xavier De Bolle for critical reading of this manuscript and the URBM members and Johan Wouters (Chemistry Department, UNamur) for fruitful discussions.

Funding and additional information

This work was supported by the University of Namur. G. L. was supported by the Belgian Fund for Industrial and Agricultural Research Associate (FRIA). Catherine Michaux and Eric Perpète thank the Belgian National Fund for Scientific Research for their respective associate and senior research associate positions.

Conflict of interest

The authors declare no competing interests

Data Availability

All data are contained within the manuscript and Supporting Information section. The raw data can be shared upon request to jean-yves.matroule@unamur.be

References

1. Wadhams, G. H., and Armitage, J. P. (2004) Making sense of it all: bacterial chemotaxis. *Nat Rev Mol Cell Biol* **5**, 1024-1037

2. Thomas, M. A., Kleist, A. B., and Volkman, B. F. (2018) Decoding the chemotactic signal. *J Leukoc Biol* **104**, 359-374

3. Silversmith, R. E. (2010) Auxiliary phosphatases in two-component signal transduction. *Curr Opin Microbiol* **13**, 177-183

4. Sircar, R., Greenswag, A. R., Bilwes, A. M., Gonzalez-Bonet, G., and Crane, B. R. (2013) Structure and activity of the flagellar rotor protein FliY: a member of the CheC phosphatase family. *J Biol Chem* **288**, 13493-13502

5. Lertsethtakarn, P., Howitt, M. R., Castellon, J., Amieva, M. R., and Ottemann, K. M. (2015) Helicobacter pylori CheZ(HP) and ChePep form a novel chemotaxis-regulatory complex distinct from the core chemotaxis signaling proteins and the flagellar motor. *Mol Microbiol* **97**, 1063-1078

6. Briegel, A., Wong, M. L., Hodges, H. L., Oikonomou, C. M., Piasta, K. N., Harris, M. J., Fowler, D. J., Thompson, L. K., Falke, J. J., Kiessling, L. L., and Jensen, G. J. (2014) New insights into bacterial chemoreceptor array structure and assembly from electron cryotomography. *Biochemistry* **53**, 1575-1585

7. Samanta, D., Borbat, P. P., Dzikovski, B., Freed, J. H., and Crane, B. R. (2015) Bacterial chemoreceptor dynamics correlate with activity state and are coupled over long distances. *Proc Natl Acad Sci U S A* **112**, 2455-2460

8. Flack, C. E., and Parkinson, J. S. (2022) Structural signatures of Escherichia coli chemoreceptor signaling states revealed by cellular crosslinking. *Proc Natl Acad Sci U S A* **119**, e2204161119

9. Parkinson, J. S., Hazelbauer, G. L., and Falke, J. J. (2015) Signaling and sensory adaptation in Escherichia coli chemoreceptors: 2015 update. *Trends Microbiol* **23**, 257-266

10. Macomber, L., and Imlay, J. A. (2009) The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc Natl Acad Sci U S A* **106**, 8344-8349

11. Giachino, A., and Waldron, K. J. (2020) Copper tolerance in bacteria requires the activation of multiple accessory pathways. *Mol Microbiol* **114**, 377-390

12. Lawaree, E., Gillet, S., Louis, G., Tilquin, F., Le Blastier, S., Cambier, P., and Matroule, J. Y. (2016) Caulobacter crescentus intrinsic dimorphism provides a prompt bimodal response to copper stress. *Nat Microbiol* **1**, 16098

13. Kovarik, M. L., Brown, P. J., Kysela, D. T., Berne, C., Kinsella, A. C., Brun, Y. V., and Jacobson, S. C. (2010) Microchannel-nanopore device for bacterial chemotaxis assays. *Anal Chem* **82**, 9357-9364

14. Morse, M., Colin, R., Wilson, L. G., and Tang, J. X. (2016) The Aerotactic Response of Caulobacter crescentus. *Biophys J* **110**, 2076-2084

15. Nesper, J., Hug, I., Kato, S., Hee, C. S., Habazettl, J. M., Manfredi, P., Grzesiek, S., Schirmer, T., Emonet, T., and Jenal, U. (2017) Cyclic di-GMP differentially tunes a bacterial flagellar motor through a novel class of CheY-like regulators. *Elife* **6**

16. Berne, C., and Brun, Y. V. (2019) The Two Chemotaxis Clusters in Caulobacter crescentus Play Different Roles in Chemotaxis and Biofilm Regulation. *J Bacteriol* **201**

17. Tsai, J. W., and Alley, M. R. (2000) Proteolysis of the McpA chemoreceptor does not require the Caulobacter major chemotaxis operon. *J Bacteriol* **182**, 504-507

18. Sourjik, V., and Schmitt, R. (1998) Phosphotransfer between CheA, CheY1, and CheY2 in the chemotaxis signal transduction chain of Rhizobium meliloti. *Biochemistry* **37**, 2327-2335

19. Briegel, A., Ding, H. J., Li, Z., Werner, J., Gitai, Z., Dias, D. P., Jensen, R. B., and Jensen, G. J. (2008) Location and architecture of the Caulobacter crescentus chemoreceptor array. *Mol Microbiol* **69**, 30-41

20. Potocka, I., Thein, M., M, O. S., Jenal, U., and Alley, M. R. (2002) Degradation of a Caulobacter soluble cytoplasmic chemoreceptor is ClpX dependent. *J Bacteriol* **184**, 6635-6641

21. Nierman, W. C., Feldblyum, T. V., Laub, M. T., Paulsen, I. T., Nelson, K. E., Eisen, J. A., Heidelberg, J. F., Alley, M. R., Ohta, N., Maddock, J. R., Potocka, I., Nelson, W. C., Newton, A., Stephens, C., Phadke, N. D., Ely, B., DeBoy, R. T., Dodson, R. J., Durkin, A. S., Gwinn, M. L., Haft, D. H., Kolonay, J. F., Smit, J., Craven, M. B., Khouri, H., Shetty,

J., Berry, K., Utterback, T., Tran, K., Wolf, A., Vamathevan, J., Ermolaeva, M., White, O., Salzberg, S. L., Venter, J. C., Shapiro, L., and Fraser, C. M. (2001) Complete genome sequence of Caulobacter crescentus. *Proc Natl Acad Sci U S A* **98**, 4136-4141

22. Lamb, A. L., Wernimont, A. K., Pufahl, R. A., Culotta, V. C., O'Halloran, T. V., and Rosenzweig, A. C. (1999) Crystal structure of the copper chaperone for superoxide dismutase. *Nat Struct Biol* **6**, 724-729

23. Drozdetskiy, A., Cole, C., Procter, J., and Barton, G. J. (2015) JPred4: a protein secondary structure prediction server. *Nucleic Acids Res* **43**, W389-394

24. Barnakov, A. N., Barnakova, L. A., and Hazelbauer, G. L. (1998) Comparison in vitro of a high- and a low-abundance chemoreceptor of Escherichia coli: similar kinase activation but different methyl-accepting activities. *J Bacteriol* **180**, 6713-6718

25. Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E. L. (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**, 567-580

26. Teufel, F., Almagro Armenteros, J. J., Johansen, A. R., Gislason, M. H., Pihl, S. I., Tsirigos, K. D., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022) SignalP 6.0 predicts all five types of signal peptides using protein language models. *Nat Biotechnol*

27. Neumann, S., Hansen, C. H., Wingreen, N. S., and Sourjik, V. (2010) Differences in signalling by directly and indirectly binding ligands in bacterial chemotaxis. *EMBO J* **29**, 3484-3495

28. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli, P., and Hassabis, D. (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589

29. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D., Stroe, O., Wood, G., Laydon, A., Zidek, A., Green, T., Tunyasuvunakool, K., Petersen, S., Jumper, J., Clancy, E., Green, R., Vora, A., Lutfi, M., Figurnov, M., Cowie, A., Hobbs, N.,

Kohli, P., Kleywegt, G., Birney, E., Hassabis, D., and Velankar, S. (2022) AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439-D444

30. Rubino, J. T., and Franz, K. J. (2012) Coordination chemistry of copper proteins: how nature handles a toxic cargo for essential function. *J Inorg Biochem* **107**, 129-143

31. Xiao, Z., Brose, J., Schimo, S., Ackland, S. M., La Fontaine, S., and Wedd, A. G. (2011) Unification of the copper(I) binding affinities of the metallo-chaperones Atx1, Atox1, and related proteins: detection probes and affinity standards. *J Biol Chem* **286**, 11047-11055

32. Kahra, D., Kovermann, M., and Wittung-Stafshede, P. (2016) The C-Terminus of Human Copper Importer Ctr1 Acts as a Binding Site and Transfers Copper to Atox1. *Biophys J* **110**, 95-102

33. Maertens, L., Cherry, P., Tilquin, F., Van Houdt, R., and Matroule, J. Y. (2021) Environmental Conditions Modulate the Transcriptomic Response of Both Caulobacter crescentus Morphotypes to Cu Stress. *Microorganisms* **9**

34. Lee, J. W., and Helmann, J. D. (2006) The PerR transcription factor senses H2O2 by metal-catalysed histidine oxidation. *Nature* **440**, 363-367

35. Tiwari, M. K., Hagglund, P. M., Moller, I. M., Davies, M. J., and Bjerrum, M. J. (2019) Copper ion / H2O2 oxidation of Cu/Zn-Superoxide dismutase: Implications for enzymatic activity and antioxidant action. *Redox Biol* **26**, 101262

36. Ramteke, S. N., Walke, G. R., Joshi, B. N., Rapole, S., and Kulkarni, P. P. (2014) Effects of oxidation on redox and cytotoxic properties of copper complex of Abeta1-16 peptide. *Free Radic Res* **48**, 1417-1425

37. Stadtman, E. R. (1993) Oxidation of free amino acids and amino acid residues in proteins by radiolysis and by metal-catalyzed reactions. *Annu Rev Biochem* **62**, 797-821

38. Rubino, J. T., Chenkin, M. P., Keller, M., Riggs-Gelasco, P., and Franz, K. J. (2011) A comparison of methionine, histidine and cysteine in copper(I)-binding peptides reveals differences relevant to copper uptake by organisms in diverse environments. *Metallomics* **3**, 61-73

39. Chervitz, S. A., and Falke, J. J. (1996) Molecular mechanism of transmembrane signaling by the aspartate receptor: a model. *Proc Natl Acad Sci U S A* **93**, 2545-2550

40. Salah Ud-Din, A. I. M., and Roujeinikova, A. (2017) Methyl-accepting chemotaxis proteins: a core sensing element in prokaryotes and archaea. *Cell Mol Life Sci* **74**, 3293-3303

41. Dupont, C. L., Grass, G., and Rensing, C. (2011) Copper toxicity and the origin of bacterial resistance--new insights and applications. *Metallomics* **3**, 1109-1118

42. Macomber, L., Rensing, C., and Imlay, J. A. (2007) Intracellular copper does not catalyze the formation of oxidative DNA damage in Escherichia coli. *J Bacteriol* **189**, 1616-1626

43. Khursigara, C. M., Wu, X., and Subramaniam, S. (2008) Chemoreceptors in Caulobacter crescentus: trimers of receptor dimers in a partially ordered hexagonally packed array. *J Bacteriol* **190**, 6805-6810

44. Webre, D. J., Wolanin, P. M., and Stock, J. B. (2003) Bacterial chemotaxis. *Curr Biol* **13**, R47-49

45. Poindexter, J. S. (1981) The caulobacters: ubiquitous unusual bacteria. *Microbiol Rev* **45**, 123-179

46. Evinger, M., and Agabian, N. (1977) Envelope-associated nucleoid from Caulobacter crescentus stalked and swarmer cells. *J Bacteriol* **132**, 294-301

47. Ostergaard, H., Tachibana, C., and Winther, J. R. (2004) Monitoring disulfide bond formation in the eukaryotic cytosol. *J Cell Biol* **166**, 337-345

48. Ducret, A., Quardokus, E. M., and Brun, Y. V. (2016) MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. *Nat Microbiol* **1**, 16077

49. Meier, F., Brunner, A. D., Koch, S., Koch, H., Lubeck, M., Krause, M., Goedecke, N., Decker, J., Kosinski, T., Park, M. A., Bache, N., Hoerning, O., Cox, J., Rather, O., and Mann, M. (2018) Online Parallel Accumulation-Serial Fragmentation (PASEF) with a Novel Trapped Ion Mobility Mass Spectrometer. *Mol Cell Proteomics* **17**, 2534-2545

50. Zhang, J., Xin, L., Shan, B., Chen, W., Xie, M., Yuen, D., Zhang, W., Zhang, Z., Lajoie, G. A., and Ma, B. (2012) PEAKS DB: de novo sequencing assisted database search for sensitive and accurate peptide identification. *Mol Cell Proteomics* **11**, M111 010587

Figure legends

Figure 1: McpR is involved in Cu chemotaxis and Cu binding. A. Percentage of WT and Δ*mcp* *(mcpA* to *mcpS)* SW cells remaining in the vicinity of the Cu plug after 25 min. **B.** 3D structure prediction of McpR of McpR by AlphaFold. H99 and C75 residues are highlighted. **C.** Representative McpR titration by Cu(I) (squares) or Cu(II) (circles) monitored by protein intrinsic fluorescence intensity quenching, fitted to a single-site hyperbolic model by non-linear regression analysis (solid line). **D.** Number of Cu atoms per spheroplast from SW cells exposed to various Cu concentrations for 5 min. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Figure 2: Cu-induced ROS sustain Cu chemotaxis. A. Raw fluorescence of SW cells incubated with 5 µM CellROX Deep Red probe and exposed to 175 µM Cu or 400 µM H_2O_2. **B.** Relative fluorescence intensity of SW cells expressing the rxYFP biosensor and exposed to 175 µM Cu or 400 µM H_2O_2 for 20 min. **C.** Relative fluorescence intensity of SW cells expressing the rxYFP biosensor and exposed to various Cu concentrations for 20 min. **D.** Relative fluorescence intensity of WT SW cells and SW cells overexpressing SodB (*sodB+*) and KatG (*katG+*) incubated with 5 µM CellROX Deep Red probe and exposed to 175 µM Cu. **E.** Growth profiles at $OD_{660\ nm}$ of WT, *sodB+* and *katG+* strains in PYE (orange) and exposed to 175 µM Cu (blue). **F.** Percentage of WT, Δ*cheAI*, *sodB+* and *katG+* SW cells in the vicinity of the Cu plug after 25 min. Mean +/- s.d., at least 3 biological replicates, except for 300 µM Cu (2 biological replicates). *p- values* were calculated using an ANOVA when appropriate (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Figure 3: Cu cycling modulates its coordination by McpR. A. Percentage of oxidation of the H99 residue from purified McpR (left panel) and McpR$_{C75S}$ (right panel) incubated with Cu(II), Cu(I), H$_2$O$_2$ and Cu(I)+H$_2$O$_2$ at a 1:100 ratio. **B.** Measure, by intrinsic fluorescence, of the Kd values of purified McpR, McpR$_{H99A}$ and McpR$_{C75S}$ (1.79 µM) incubated with 0 to 500 µM Cu(I) and Cu(II). Mean +/- s.d. **C.** Percentage of WT, Δ*mcpR* and Δ*mcpR* overexpressing McpR, McpR$_{H99A}$ or McpR$_{C75S}$ SW cells remaining in the vicinity of the Cu plug after 25 min. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**). **D.** Circular Dichroism spectra of purified McpR (8.95 µM) incubated with Cu(I) and Cu(II) at a 1:100 ratio.

Figure 4: Hypothetical model of Cu sensing by McpR. SW cell exposure to Cu increases the cytoplasmic Cu concentration, leading to the generation of ROS (left). In addition, high cytoplasmic Cu content favors Cu(I) coordination by the C75 and H99 residues within the cytoplasmic McpR, triggering a conformational change of the chemoreceptor. Cu(II) binding to H99 residue has no impact on McpR apo-like conformation, suggesting a redox-dependent modulation of McpR conformation. This process would be maintained along the extracellular Cu gradient until the SW cell reaches a Cu-poor environment (right). Under these conditions, the cytoplasmic Cu concentration together with the ROS level would return to their basal level and the labile Cu binding to McpR would be hindered, bringing McpR to its apo form conformation.

Figure 1

A.

B.

C.

D.

Figure 2

A.

B.

C.

D.

E.

F.

Figure 3

A.

McpR / C75S

Ctl / Cu(II) / Cu(I) / H₂O₂ / Cu(I) + H₂O₂

Oxidation (%) / Oxidation (%)

Oxidized / Non-oxidized

B.

	Cu(I)	Cu(II)
	Kd (μM)	Kd (μM)
McpR	48 ± 10	37 ± 9
McpR$_{H99A}$	451 ± 205	213 ± 140
McpR$_{C75S}$	168 ± 83	46 ± 23

C.

*** / ** / **

Bacteria remaining near the Cu plug (%)

WT / +R / +R$_{H99A}$ / +R$_{C75S}$

ΔR

D.

Ellipticity

Wavelength (nm)

Ctl / Cu(I) / Cu(II)

Figure 4

Figure S1

Figure S1: McpA, McpB and McpR are potentially involved in Cu chemotaxis. Percentage of WT, Δ*mcpA*, Δ*mcpA* overexpressing McpA, Δ*mcpB*, Δ*mcpB* overexpressing McpB, Δ*mcpR* and Δ*mcpR* overexpressing McpR SW cells remaining in the vicinity of the Cu plug after 25 min. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Figure S2

A.

B.

Figure S2: The chemotactic machinery is functional in the Δ*mcpR* mutant. A. Chemotaxis-driven swarming pattern of the WT, Δ*mcpA*, Δ*mcpB* and Δ*mcpR* strains in a Cu ions-free medium. **B.** Percentage of WT and Δ*mcpR* SW cells remaining in the vicinity of the Zn plug after 25 min. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using a *t-test* (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Figure S3

Figure S3: Zn, Cd, Mn and Ni do not directly bind to McpR. Relative intrinsic fluorescence of purified McpR (3.58 µM) incubated with $ZnSO_4$, $CdSO_4$, $MnSO_4$ and $NiSO_4$ at 1:10 and 1:100 ratios. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using a t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Figure S4

A.

B.

Figure S4: Potential pathways leading to ROS production by Cu. A. *In vitro* generation of Cu-induced ROS by Fenton-like reaction and Haber-Weiss cycles. The Cu-GSH complexes may also trigger ROS production (27, 28). The SODs and KatG are involved in $O_2\bullet-$ and H_2O_2 detoxification, respectively **B.** The rxYFP biosensor is in equilibrium with the GSH pool. GSH is involved in different ways in ROS buffering. As a result, ROS production triggers an increase of the GS-SG pool, which will be partially reduced to GSH by using rxYFP as an electron donor. The resulting oxidized rxYFP will lose its fluorescence properties. GSH (glutathione), PRX (Peroxiredoxins), TRX (thioredoxin) and GRX (glutaredoxin).

Figure S5

Figure S5: CheAI is the main histidine kinase involved in Cu chemotaxis. Percentage of WT, ΔcheAI, ΔcheAII and ΔcheAI-II SW cells remaining in the vicinity of the Cu plug after 25 min. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$): WT vs. ΔcheAI = 0.0019; WT vs. ΔcheAII = >0.9999; WT vs. ΔcheAI-II = 0.0406.

Figure S6

Figure S6: The Cu-induced oxidative stress is not affected in the Δ*mcpR* mutant. Relative intensity of fluorescence of the WT and Δ*mcpR* SW cells expressing the rxYFP biosensor and exposed to 175 µM Cu for 20 min. Mean +/- s.d., at least 3 biological replicates. *p- values* were calculated using a *t-test* (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Figure S7

A.

H99A
- ■ Cu(I)
- ○ Cu(II)

Fluorescene intensity units

[Cu(I), Cu(II)], µM

B.

C75S
- ■ Cu(I)
- ○ Cu(II)

Fluorescence intensity units

[Cu(I), Cu(II)], µM

Figure S7: Fitting curves of the McpR$_{H99A}$ and McpR$_{C75S}$ variants. Representative H99A (**A.**) and C75S (**B.**) McpR titration by Cu(I) (squares) or Cu(II) (circles) monitored by protein intrinsic fluorescence intensity quenching, fitted to a single-site hyperbolic model by non-linear regression analysis (solid line). The fluorescence intensity axis scales correspond to the same amplitude for B and C.

Figure S8

Figure S7: The McpR$_{C75S}$ and McpR$_{H99A}$ mutants are still functional. Percentage of WT, Δ$mcpR$ and Δ$mcpR$ overexpressing McpR, McpR$_{H99A}$ or McpR$_{C75S}$ SW cells remaining in the vicinity of the Cd plug after 25 min. Mean +/- s.d., at least 3 biological replicates. *p-values* were calculated using an ANOVA (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) (**Table S1**).

Table S1: *pvalues.* One-way ANOVA and t-test were performed using GraphPad Prism.

Figure 1A (ANOVA)	pvalue	summary
WT vs. ΔA	<0.0001	****
WT vs. ΔB	0.0338	*
WT vs. ΔC	0.9993	ns
WT vs. ΔD	0.9992	ns
WT vs. ΔE	>0.9999	ns
WT vs. ΔF	0.6094	ns
WT vs. ΔG	0.9991	ns
WT vs. ΔH	0.8952	ns
WT vs. ΔI	0.9689	ns
WT vs. ΔJ	0.999	ns
WT vs. ΔK	0.9991	ns
WT vs. ΔL	>0.9999	ns
WT vs. ΔM	0.9796	ns
WT vs. ΔN	0.9993	ns
WT vs. ΔO	0.9996	ns
WT vs. ΔP	0.9996	ns
WT vs. ΔQ	0.9988	ns
WT vs. ΔR	0.0001	***
WT vs. ΔS	0.9995	ns

Figure 1D (ANOVA)	pvalue	summary
50 µM Cu vs. 100 µM Cu	0.0128	*
50 µM Cu vs. 175 µM Cu	0.0011	**
100 µM Cu vs. 175 µM Cu	0.2469	ns

Figure 2A (ANOVA)	pvalue	summary
Ctl vs. Cu	<0.0001	****
Ctl vs. H2O2	0.0007	***

Figure 2B (ANOVA)	pvalue	summary
Ctl vs. Cu	<0.0001	****
Ctl vs. H2O2	<0.0001	****

Figure 2C (ANOVA)	pvalue	summary
Ctl vs. 20 µM	<0.0001	****
Ctl vs. 175 µM	<0.0001	****
Ctl vs. 300 µM	0.0004	***
Ctl vs. 1.16 mM	<0.0001	****

Figure 2D (ANOVA)	pvalue	summary
WT vs. sodB+	0.0005	***
WT vs. katG+	0.001	***

Figure 2F (ANOVA)	pvalue	summary
WT vs. cheAI	0.0007	***
WT vs. sodB+	0.0156	*
WT vs. katG+	0.0045	**

Figure 3C (ANOVA)	pvalue	summary
WT vs. ΔR	0.0001	***
WT vs. ΔR+R	>0.9999	ns
WT vs. H99A	0.2356	ns
WT vs. C75S	0.2503	ns
ΔR+R vs. C75S	0.0056	**
ΔR+R vs. H99A	0.0053	**

Figure S1 (ANOVA)	pvalue	summary
WT vs. ΔA	<0.0001	****
WT vs. ΔA+A	>0.9999	ns
WT vs. ΔB	0.0338	*
WT vs. ΔB+B	>0.9999	ns
WT vs. ΔR	0.0001	***
WT vs. ΔR+R	0.9954	ns

Figure S2 (t-test)	pvalue	summary
WT vs. ΔR	0.4878	ns

Figure S3 (t-test)	pvalue	summary
Cu(II) 1:10 vs. Cu(II) 1:100	1.0372E-05	***
Cu(I) 1:10 vs. Cu(I) 1:100	0.00784501	**
Zn 1:10 vs. Zn 1:100	0.9479843	ns
Cd 1:10 vs. Cd 1:100	0.59277248	ns
Mn 1:10 vs. Mn 1:100	0.70207278	ns
Ni 1:10 vs. Ni 1:100	0.58457574	ns

Figure S5 (t-test)	pvalue	summary
WT Ctl vs. R Ctl	0.9651	ns
WT Cu vs. R Cu	0.6783	ns

Figure S7 (ANOVA)	pvalue	summary
WT vs. ΔR	0.0171	*
WT vs. H99A	0.8499	ns
WT vs. C75S	0.9949	ns

Table S2: Strains, primers and plasmids (1/4)

Strains	Relevant genotype or description	References
C. crescentus		
WT	CB15N (or NA1000), synchronizable variant strain of CB15	(Evinger and Agabian, 1977)
WTrxYFP	WT strain carrying the *rxYFP* gene on the pJS14 under the control of the lac promoter; CmR	This study
ΔcheAI	Knockout strain for *cheAI* gene (CCNA_00442)	This study
ΔA	Knockout strain for *mcpA* gene (CCNA_00439)	This study
ΔA+A	Knockout strain for *mcpA* gene carrying a copy of *mcpA* on the pMR10 under the control of the lac promoter; KanR	This study
ΔB	Knockout strain for *mcpB* gene (CCNA_00437)	This study
ΔB+B	Knockout strain for *mcpB* gene carrying a copy of *mcpB* on the pMR10 under the control of the lac promoter; KanR	This study
ΔC	Knockout strain for *mcpC* gene (CCNA_00348)	This study
ΔD	Knockout strain for *mcpD* gene (CCNA_01727)	This study
ΔE	Knockout strain for *mcpE* gene (CCNA_02364)	This study
ΔF	Knockout strain for *mcpF* gene (CCNA_02773)	This study
ΔG	Knockout strain for *mcpG* gene (CCNA_00626)	This study
ΔH	Knockout strain for *mcpH* gene (CCNA_03459)	This study
ΔI	Knockout strain for *mcpI* gene (CCNA_02940)	This study
ΔJ	Knockout strain for *mcpJ* gene (CCNA_03247)	This study
ΔK	Knockout strain for *mcpK* gene (CCNA_00629)	This study
ΔL	Knockout strain for *mcpL* gene (CCNA_03468)	This study
ΔM	Knockout strain for *mcpM* gene (CCNA_02402)	This study
ΔN	Knockout strain for *mcpN* gene (CCNA_00538)	This study
ΔO	Knockout strain for *mcpO* gene (CCNA_02935)	This study
ΔP	Knockout strain for *mcpP* gene (CCNA_01465)	This study
ΔQ	Knockout strain for *mcpQ* gene (CCNA_00064)	This study
ΔR	Knockout strain for *mcpR* gene (CCNA_02901)	This study
ΔRrxYFP	Knockout strain for *mcpR* gene carrying the *rxYFP* gene on the pJS14 under the control of the lac promoter; CmR	This study
ΔR+R	Knockout strain for *mcpR* gene carrying a copy of *mcpR* on the pMR10 under the control of the lac promoter; KanR	This study
ΔR+RH99A	Knockout strain for *mcpR* gene carrying a copy of *mcpR* with the mutation H99A on the pMR10 under the control of the lac promoter; KanR	This study
ΔR+RC75S	Knockout strain for *mcpR* gene carrying a copy of *mcpR* with the mutation C75S on the pMR10 under the control of the lac promoter; KanR	This study
ΔS	Knockout strain for *mcpS* gene (CCNA_00160)	This study
sodB+	WT strain carrying a copy of *sodB (CCNA_01855)* on the pMR10 under the control of the lac promoter; KanR	This study
katG+	WT strain carrying a copy of *katG (CCNA_03138)* on the pMR10 under the control of the lac promoter; KanR	This study
E. coli		
S17-1	RP4-2, Tc::Mu, KM-Tn7, for plasmid mobilization	
BL21	BL21 *E. coli* strain used for overexpression of proteins	
Plasmids	Relevant genotype or description	
pNPTS138	mobRP4+ ori-R6K sacB; integrative vector in *C. crescentus* for in-frame deletions; KanR	
pMR10	DH10B strain carrying the pMR10 low copy number replicative cloning vector in *E. coli* and *C. crescentus*; KanR	
pJS14	DH10B strain carrying the pJS14 medium copy number replicative cloning vector in *E. coli* and *C. crescentus*; CmR	
pET28a	BL21 strain carrying the pET28a high copy number replicative cloning vector in *E. coli*; KanR	

Table S2: Strains, primers and plasmids (2/4)

Target	Sequence
Gblock used for the KO of *cheAI*	TGG CTTCATCGAGG CGGTG CG CG TCGACGACGACTACCG CG CGATCCCGATCCTGGTG CTGACCACCGAAAGCGACCCTGCCAAG AAGCAG CGGG CTCG CGAGG CGGG CGC CACGG GCTGGATCGTCAAG CCG TTCAAC CCGGAAAAG CTCGTCGACG CCATCCG CCG CGTCG CCGCCTGAT CCCGGTCACGGACCTCAGTACCCCG TTATTGAATACCCGTTG CTAGACGCGAATTAGGACTTACGGGACTGACG CTGGG CTATCG CGAGGAA AGTCCG CATCCG ACGGATG G CGTGGTG CTG CTGGTCGAAGG CGAAGACGG CTCG CGGG CCG CCCTGGTCG CCGACG CCATCCATGG CCAG CGTCAGGTGGTCATCAAGTCTCTGGAG CAGAACTATCAGCAGGTCGAAGG CGTCG CCG CCG CGACG ATCCTCGG TGACGGCCG TGTGG CTC TGATCCTCGACGTCGACG CCACGATCAACCTCCG CCGTCG CGAAG
Gblock used for the KO of *mcpA*	ACTGGACG CAGGAG CAAATTCGCAGG TG CAAG TTAAGATATTCTGAGTCG CACGG TAAGATTGCGG GTTTACTTTCAGGTAAC CACGATCTG CCG CTTTGGTGACTCTCATTG TG CGG CGAGGG TTTCG CAGATTCCATG CGG CGGAG CCTGCGACCAAACGAC CACATGCTCCAAAATGGGACGAAAATCTCTGACTTCCTTTAACCAAGTCAGAGTCTGTTAGGAATCTATCGGGATTATCACCCTTCTGATGG CCCGCTTCCAGGTGGGCTCCG GTTCGTCGTCCTACGCCCGTCCGGCCGTCGCCGACGCGGGTCATCACGCCCCCGCGCGAAATCCGGTGGCCGAGGCAACAGGCCCGTCTGAA CACCTTCGCCCGTCCGGGTCGGAGCAGCGGTTCGGCGGGCCCTCGCCCAGGCCCCCGCCTCGGATGG TTGGGAGGAATTCtaaTGGTTGCCGC GATCTCGCTGCCCGAAAACCTGGATCTCAAGACCGCAGCC
Gblock used for the KO of *mcpB*	CCAGGTCCAGCAGCTGATCATTGCGCGTGAAACGCTGAAGCGCGGCGGGTGAGTGCGACACCCTGTCGCCCCCTCCGCCCGTCGGCCGTATGCAATCGATAAACCATCCCCTCGTTCCCACGATATTGACCGTTCAGTCTGTGCAAACGGTTGCAACCTGAGCGGGGGTGCGGCGGGTGCGGCGGAGCGCCGCACTCGCTGCGAGAATATTTAAGCGCTCATTAGGGATAAGGGTCCGAGGGATTTCCCCAAGCGAAACACGCGCGGACCCTCGAGCGCCCTGGGTTCAGGCATGTTCGCTTAGACCGTACGAACGCGCTCAGCAGATTTGGTTCGGGAGCTTCTCGACGCGGGGCCGTGTCGTCTTGGCGCGGTCTCTCGTCGTTCCGGCGCCAACATCGCCCCAGACAGCAGAACGCCCCCGCCGACCAGGTCGACGAGGGCGTTTTGTTGTTCCGATCGGAAAGACGCTTACGCGCCGCCCGGGAAGCGGAACGGAACCTTCACGGT
Gblock used for the KO of *mcpC*	CGACCACGCAACGTTACACCCAGGTGGACGCCGCCGGCCTGTTGGCCGCCTATCAGGCCGCCCACCCTAAGGGATGACCCCGAGCAAGCACGCCGGGAGGTTCATCGCGCCCGTCCGTCGCGATGCGACACACTGTCGCGAAGGGCGACCGCCGAAACGCCATTAACCGCGATTTTACCGCATCAGGGCGTTGTGGGGGCTTGAGAAATTCCCTCTGCGTCCCGCGTCTGATCGCTGACGAGGCCCCCTACCGGTTTAGTAAGCGTAAGCCCCGCCGCCCCCCTCTGGCGGGGTTCCCCCCCTCGAAAAACAGACTGGCCCCGCCTCCTCAAGAGGCGGGGCCTTTTCTTGGCTGTATTCGCGTTGAGCGTGGGCGCTCACGCGGATTGGAAATCCTCGTCCTTCGACAAGCTCAGGATGAGGGATTTTGTGCGAACCGAGGCCAACAGTCGTCCTCACCCTGAGCTTGTCGAAGGGCGAGGGCGGCCGCCACGACGCA
Gblock used for the KO of *mcpD*	TCGCGCCGTTCATGGTCGATCTGCTGCGCACTGATCGCGAGGCGGCCAAGGGTGTGGTTCGCTTCAAGGAACGCGGCCAGGAATCAGCGTCGATCGACGCCTGACCGATTCGGCGGCGCCGCCTTCATGGACGCAAGGGTTGTCTGACGCTGTCAGACTTCCACTCTGGATGGCGATTCCTAGACCGCGTTCGGCGATAGGTCGCCCCAAACGGGAGCATTAGCGGCTCGTTAACCATGCACGGCCAGTTTCCCGTGGGGGGACGAACCCGCAGTAACGGGTTCCAGATGGCCGCCGCTGACAAAGCGGCGGCCATTTCGGCTCTGCTTAGCGCCCTTTAGGGCGACCGGGCCCGGTGTTCCCACGCACTGTGCGCGGAACCCGCAGCCGGACGGCGCGGGCGGCGGGCGGCCTTGCCGCCGGCCCGTCTTGGTCCCGACCTTGTAGCTGGGCGTCGTGGCGACCTTCTTCTGGGCGCGCGGTCCCTTGGCCTTCACGGACTTG
Gblock used for the KO of *mcpE*	CGAGGGGACCGATTCGGCCACGCCCCTGGCCTGGAGCCACGCGGAGTACGTCAAGCTGCTGCGATCGGTCAGCGACGGGCAGGTCTGGGATCACTATGCGCCCGTAAAGGCGCGGTACGCGCGCTAGAGGGGCCAAAAAGTGCGCGACAATGTCGCGGTCGCGCATTCCTGAAACGTGGTTAACGGGGCTTTAGGCATTAGGCCGTGAACTTGAGCGTAATCGCCTAGCTCACGGCCGGAGGCCTTTGTTCGCTCCGACGCGCGAGAGCTTCCGGCAGCGCTACGTGCAGGGCGCCAACGCCCTGAAGGTCGCCCCCAGCTCCCGTCCCGGGGAATGGGAAGAGTTCtagGCCTGACCGCTCTCAGCCGTGCGCAGGGCGAAACGGAGCTCTGACAGAGCGATACCGTTCTGGGCCTTTTCGGCTTCGCGAACGCTGACCAGGAGCTCATTCATGCCATTGGCGACGGCCCAGTCCCAATGGGCGTGACCTGGGGTCGCC
Gblock used for the KO of *mcpF*	CCTGTTCTTCGGCCTCGATCACGCCCGCCAGAAGCTCGCTGCCTGGGCCGAAGACTACAACACACGCCGGCCGCACTCGTCGATCGGCTACCAGACCCCCGCGGCCTATGCCGCGCACCTGAACGCAACAGGCCGACCCGCTGCGCAATGTGGGAGCTCCACGGCTCGCCGCTGTTGCTACAACCGCGCCATCGGGCGTAACCAATGCCGAGACTCTGGTCCGCGCTGGATGAAAGATCAGAGGCAGGTCAGTCATGCTTTGGCCATGGAGGCTGACAACCTGACCGTACTGATGGGGCAGTTTCGGATCGGAGAAACTCCAGAAGCGCAGTCCAATCGCAGGGACTTAAGCAAAGGCGCCGGCGTCCCGTCTAATTTGCGACCGATAGCGCAAAGCCACGCTCGTGCGGGATCCGCACCGCAAAGATTGATGGCTGGGACGAGTTCtaaCCCAGAACACAATGGTCTCTGCGCGCGCCAAAGGCGATCAAAAGGCCCGT
Gblock used for the KO of *mcpG*	TCACCGCGCTTCCGGAGCCAGAGTCTCCCTCCGAGGCCCGTCAACGGGAGCTCTGGCTCAACGGTCCCGCGCGTCCCGCCGAAGGACTCTCGCAGGCCGACGCTGACCGTCTTTTCGAATAGCCATCGGCCCAGCGTGGCCGTGTCGGCTTGTTAACCGTTTCCGCCAGATAGTCGAACCTGGGTTCAAGGTCAGCGATCatgTCGCCCCATTCATTCTCGCCGAAAGGCGCTAACAGGTTCGATCAGCCGGCTACGGCGCGCGCTCGCCCCTGCAGCGTTCACACGTCGCGCGGCGCCGCAGCCTAGGACGCCGAACGCACCATCCGCTCGAGGCGACGCAGCAGCACCCGCGCCTCGAACGGCTTACCGATAAAGTCGGCGGCCCCAAGGCTCATCGACCGGCGAATATCCTCGGGCGAGGTCTGGGCGGTCAGCATCAAGGCGGGGATCTCCGCGAACGCGGGATCGCGACGCACCGCGTGGAGCAAGGCCAGGCCAT
Gblock used for the KO of *mcpH*	ACGCCCCCAATCCGAGCTGAGCGACAAAAGGCCACACTATTCGAAAGGTTatgTTTTACCCGAAATTCACCGAGACAAATCAGCGTTATATGGACTTGGGCGGTGTTCCGCCTCGCAAGACGCACGGATCGCGGCGAATGATCAAACTTCTGAACGGCTCCGCGGGCGACGCCAGCCAGAAGATCGACGCACTCGACAAGTCGCTCGCGATGATCGAGTTCGACGTCAAGGGAACGATCCTCGCGGCCAAGATCATCAACCGCCCCCCGCGCGCCCCCCGCGCCGGCTGATGGGCCACATCAGATCGGGTGAAGAGAGCTGTCCTTGGCCGCCGACAAGGGGGCCGGCGGCGCGATCGGCGCTCGAACCGTCTTCCGCGCCACATAGCGGCGCACACGCTTGCGGGTCTTCTTCAGCAGGCGCCAGCGGGCCTTGCGCAGGACCAGCTTCTCGGTGCGCAGCATCTGTTGCCAGACCACCAGCAGCACCTCCGGCTCGCG
Gblock used for the KO of *mcpI*	GGCTTCGGAGCCTGTATCGTCatgAACGTCACCTTCTCCAACCTCGACGCCCAACGCCAGATCGGCGGCAAGCTGATCATCGCCCTCGCCTGGCTGCTGATCCCGATCGTGCTGGGCGCGCGCCTGGCCGTGCAGGCGCCCATCCTGGGGCTCTCGATCGCCGCGGTGCTGGTCGCCACGACCCTCAACACCGACGCCACCCAGCTGGCCGATCTGGTCGCCCAGTTCGACCTACCGCGAGGCCATATCGAGCAGTCCCAGCGGCGGGTCGCCtagGACGCGGCGCGGCGCCCTTTCGCGCCGCCGCCGTTGCAAGCTCAAGCTTGGCCCCCTTGCGTTACCGGAACCGGGCGCCTATCCTCCCCAGCCCCCGGTTGTGCGCGCCCGCGACTGGAAGATCTTCCTCCCCAAGGCGTCCGAGCTAAAATCCGCAATGGAAAAA
Gblock used for the KO of *mcpJ*	CCTCGGAATTCTTTTAGTCTTGCGACTTTGGACACCTAACGCCGACGCCTCTTAAGACCGTATAAATTACGACCAGTCTGTCTGGTGACACTATGTCGCATGGTTAGGAAGGCGCCACCGAAACGTCGTTAAGAGGCGAGCAATCACTCTTCTATAGGTCTGGGGACGGCCGCGTGGCGCTAGGTCCGTCGCATAGGCGATTTCGCCTAGAGGACGGATCAGCGCCGCGGCCGCGGGAGGACCGCCCACCCCGGGCGGCGGGCGCCGGGCCCCTCCATCAATCCATCCATCCATAGCTATAGCGTCGAGCGCCCCCACCGGAGCGCTCGCGCCGGCACGCTTTCGCCTCGCGAGCCAGAGCTCGCGCTGCAGATCATGGCGATTGTAACCACGACATAATGTCGCACCACAAAGTCGCTGCTGATAAGTTCGGGTTTACCTTGTTGGTATAGTAGAAAAATCAACAAACACCCAAAAGTATATATAACTAAGAAA
Gblock used for the KO of *mcpK*	GCCTAGTAGGTTCAGCTCGACGCTAAAAGGCTTCAGTATTCCTCTAAACTCGCCACACTCAAAATGAGTATAGATCGCGCCTCTTAGAGTTATTCGGGGCCGGGGCTCGGGGCGGGTAACGCGCGGTTAATCGCGCGGATCAGAGATTGACGACCCATGCGATTTGCCCAGGCCGCCTCCATTCCTCGGATGAAGGCGACGGGTCGGAGCGCGATGAAGGCGGGTTTTAAGACTCTGCGGCGTTGGGCCCGCGAGATCGCGCGCAACACCTCCGAAGCCGCCAACGGCGCTCAAGACGTGTCGCGGAACATTTCCGACGTCTTGCATGGCGCCCAACAGACCGGCTCCGCCTCGCACCAGGTGGTCAGCGCCTCGCAGGAGCTGGGGCGTCAAGCCGCAGCGCTGCGTGGTCACCTTCCTGAAGACGGTGCGGGCCGCCTAGCGCGCTCCGGCGCGGTTAGGGCCTTCGGAGATAGCGATGGATGATCTGAT

Gblock used for the KO of *mcpL*	CAATCTGTCG CGGGTATGG TTACGG CCGCGTTTA CCATCGG CCGAG CATCG TTG CAG CG CAGTTCAAG CTTGATCATTCAAC CCCAAG CCCC GGGTTCAGatgCTG CG CCTCAACG CCTTCCTG AAACTG CTGACCGTCG CG TTGATGG TCGG CACACAATCCTTG CCATCG CCG CAG CGACGTTCG GCCTTTTG CATCTG CGGGTCGG CGGACCGATCGCCAACCGTCAAG CCG CCG CCTTCGATCTCGTCAGCCCTCTAGCTTTGGATGGGAGCATCC AAAG CGTTGGAAAGAGATCTAAATATCTGATTTTAAAG CCTTACATCCG CTCTGG CG CTTCGATGCCCAGCAGGTCCAACG CCAGCTCCAGCT GGCGCAGGGTGGTCTCGGCCAGGGTCAGGCGCGAGGCGCGCAGCGTCTCGGTGTCAGCGCTCATGATCGGGCAGGCCGCGTAGAACTTCG AGAACGACTGGGCCAGCTTGTAGGCGTGCTCGGCCACGA
Gblock used for the KO of *mcpM*	GTCGCAGCGGTTGACGCCGGTAAGGTCAAAGCGCAGGTCGCGACGGTCTTCGAGCGCCTCGGTGAGGCGAGGCCCGGCGTCAAACAGGCC CCGCGCCGTCCAGTCCCCGGCGAGTACGGCGACCAGGCCTCCGTCCTGTTCGGTAAGGGTGAAGTCCGCCGGAAGCGCCATGAAGTCCTTG GGGTCTGGTCCCGGTTATATCCAAGCCCACGCAAGGGCCTGACAAGCCCAAAACGCGCCAAAGCGCAGCCACGGCCCTCGGCCAAGTGAAC GCCGCCGTGACCCAGCTCGACCAGGTGACCCAGCGCAACGCCGCCATGGTTTCGCAATCGACCGACGCCACCCACGCGCTCCGTCGTCGAGG CGGCCGATCTGTCGAACCGCGTCGGAGCTTTCCGCCTGGGGGTCGGTCACAGCCGACCCATCAGGAACAGCCCGCCGAGAATCCCGTGCA CGCCGCCCGCGCGCGCGTCGCCGCTTTCGCCCGTCCCGGGCGCTGA
Gblock used for the KO of *mcpN*	CGGCCGAGATCGCGCTTTCGCGACGCCGAATAGGCGCCCGAAACAGTCGAAGACCCGGACGCCAAAAGTGTCCGGGGCGACGCCGAGCAAG GTTTTCGCCAATTGATTGAGGAAAGGCCCGGGAGAGATCCCGGCGCCTTTCTTTTGGGTTGAGTGATCGCCCACAGATTGGGTAACCGGTG GAACAACGTTTTCCTTGCGTTAAGAATCGACGCGCATCTTTCCTTAAGGGGAGTATGCAAACGATGTCAACAAGCGCCGAGACCGTCCGAAA GAGCGTCACGGCGCCGGGAGGAATGTCACAATCACGCGCCGGTACCGGGTCAGGGCCGGCGCGCCCTTGTCGTCGATCGCCACGATGAAG TCCCCGCCTCAGGATAGTTGAACCAGAGAAACGACAGGCTGTCGCC
Gblock used for the KO of *mcpO*	TGCGCCCCTGAACGATGGCCCGACGTTCAAGGTTCAGCAAGATTTACCatgCCTTCTCTCGCCACAGTAGAGTTTGGGGGCGTGAAGATGGCC GGTGCGTTCAAGCGGTTATCTCTGGCGGGGAAACTGATGGCGGCCGGGGCGGGGCCCTTGGCGCGCTGTTGATCGCAGCCTCGCTGCTG ATCGCCTGGAGCAGCAATGACGCTGTTCGAGACCTGGCCGGCCGAACGCTACGCCGCCAGCATGACCGGAGCCCTTTAGCTTTAGATGGAATCATC TAAAGCGTTTGAAAGGGCTCTAAGTATCTGATTTTAGAGCCTGTTTATCTTGTTTAGATGGTTCCATCTAAACCAGACGGGCTCTAAGGCGCG AACGCCGCCTCAGGCCGAGGCGGTTTCGCATCTCGCTCATCACCGCCGCGCGCCAGCCTCTTTCGAGCGTCCAGGCGGCGATGGCCAGCCAG AGCAGGCCCAGGAACGCGTGCCCGCGCTGCCAGTAG
Gblock used for the KO of *mcpP*	ACGTAGACACGCGTATGCGACAAAATGCGCGCGATCGCACTGAAGTCACGGACGGTTAATCCCTAGATCCAATGCTTGGCCGGAATCTCCA GGGGCGGCCatgATCGAGTTCGACAGCATTGAGGCCCAGCGGCGCACGGGCGGGAAGCTGGTCTTGGCCTGCAATGCGGCGATGGTGGCG TTGACGCCACTGGCCAGTCTTGTCGCGGGCAACGCGGTCGCGAAACTGGCCGGAATCGCCACGGTCCTGTCGGCTTCTGGCATCTGCGCTGA ACCGCCCTTGCGGCGGCGGGTCCAGAACAGCGAACGCGCTGGAGCGCTTTCCGATCGTATGAAGCGGTGATCGAATGAAGAGCCGTCTACC GTAAAGGTCAGCGAGCGGCGCAGCGCACAGCGGCCCTTTCGAGGCTCCAGTAGCGCCGGATGACCAGGGCGAGGGTCGCCAGCGTGAAG CCGGCGGCGACGTAGCACGCTGGGCAATGGCCCGCGAACGCCCAGCT
Gblock used for the KO of *mcpQ*	AGTTCCACGCCGCCTTCGGCCGCTATTTCGAGATCGCCGCCGTTTACCGTCAGGTGGCGCTGGTGAAGACCTCCGAAATCCCGCGCTAGCGA CGGTTCTCTAGAAATTCATTTAAATACAATATTTTAGGTATTTCGCGACATCATGCCACCTCGCGACATCATGCCACCCATGCGAGAATGTCATGGT TAAGGCCCGCTAGACTAGTTGGGCCGCATGCTCCGCCCATTCGGATCTGCGGAGAGGTCGCGATCACCGCCGCCAGCCATACCCTGGCGCA GGAAGCCACACGACTGGCGCAGCGCATGGGCGAACTGCGGATCATGCGCGCCGGAATGCAGGCGGCGGCCTAGAGCGTTTCCGATCTGT TTGGATCGGAGAAACGCTCTAGGCCTTTGTATTAACGCATTTTCTGACGACGAACCGGATCCACTTCGTCGGAAAATGCTCTAGATCACACGC CGCCGGCGGTCGAGGAACGGGCCGGTCGCGGGGTCTAGAG
Gblock used for the KO of *mcpR*	GACGTTAATGCTATGTTAGCCGTTTTGCTCGACCTCATTAACCTAATACGttgCGGGCGCGGCGTGCGAGCGGGGGGCTCGGGGATATGTGG TTCAAAGAGATCGCCCGACGGCGTCGCGCCGAGACGCGCCGCCTGAAGGAACTGGAGAGCCTGGCCAAGGCCATGGAGCGGTCGCAGGCC GGCTGGAACTGCGGCCCAATGCGACGGTCGTCCGCGCCAACGGCGCGTTCCAGCGCCTGACCGGCTAGGGATGCTGTGTCATCCCGGAAA GCGTGTAGCGCTTGTCCGGGACCCAGGGGCGGCAAGCGCGGCGCTCCCGCAGTCCCTGGGTCCCGGCTCTACCCCCGGCTTTCGCCGGGGT TCGGCCGGGATGACACAGGATATTTCTGTCAGGGGTGGCGGAGGTGGCGAGCATGTGCGTCGATCCTGCGACTCACGAAACCCGAGACCCT CCATGAAGTCCCTGTTCGCCGCCTCGGCTCTCGCCCTGCTGATCG
Gblock used for the KO of *mcpS*	ACCACGCCGACGACGCCGACGACATGTTCAGCCGCCTTATGGGTGACTTGGTCGAGCCGCGCCGAGAGTTCATCCAGGAGAACGCCCTCGA CGCCGAGGTCGACGTCTAAAGCTCATCCTTGCAGCTAAGTGTTGTTGCAAGTGTTTGTTCCATAAGCGCCGGGCGCCCAATCCAGGGGCGGC CCGGCGTTTTCTTTTCCAGAAACGCCCGTCTCGCCTTGGCCGCGGGCCCCAGTCGCTTCGCGTCGCCGCGGGAATGTTGAACGCTCGCGCAG AGAAAGAGTACGCATTTGTTATCGTCCGCGTCCAACCCAATACGGGATGGCATGCTTCCGGGCGCCTTCCTGCGGAATTTTTGAACTAGTTTT TCGCCCTGAAAGCGAATTCGTTCAGAACTCTGGCGCCGTCGTTGGATTCGATATGACATTTGTTGTCTTGCCAGCGTCATCATGTTTGTATAGC GCATTGCCGGCTTGCTGTTTCAGTCGCTGGACACAACC

Table S2: Strains, primers and plasmids (4/4)

Target	Plasmid	Forward Primer (5'-3')	Reverse Primer (5'-3')
Primers used to control the deletion of *cheAI*		GCCCGAAAACCTGGATCT	CGAAGCTGCAGTACTCGTTG
Primers used to control the deletion of *mcpA*		TGTTCCGATCGGAAAGACGC	CCCGTAAGTCCTAATTCGCG
Primers used to overexpress *mcpA*	pMR10 (with HindIII and XbaI)	CCAAGCTTatgTTGGCGATCCGTGGG	GCTCTAGAttaGAATTCCTCCCAACC
Primers used to control the deletion of *mcpB*		CGAAGGCTTCAAGGTCGC	CCGCAATCTTACCGTGCG
Primers used to overexpress *mcpB*	pMR10 (with HindIII and XbaI)	CCCAAGCTTatgGGGACCGCCATGAAC	GCTCTAGAGCctaGAATTCTTCCCACTCTTCC
Primers used to control the deletion of *mcpC*		CCCTCTCTTTGCGGATGAGC	ACAAGGTCGGCAAGTTCCCG
Primers used to control the deletion of *mcpD*		ATGGCCTGGATCACCTCGGC	TCAGTGGTCGTCGCGTGAAGC
Primers used to control the deletion of *mcpE*		TCTACCGCGTTCGCTACAG	CCTGCCCAGAAATAGCTGTC
Primers used to control the deletion of *mcpF*		TTCGTGCACGACCAGTTCGC	GTCGCGCTTGCTCAAGTTCG
Primers used to control the deletion of *mcpG*		ATCGTCAAGCCTTTCAGCGC	CGGTGGGACAGATCAAGGC
Primers used to control the deletion of *mcpH*		AGAGGTTCTGGTTCCAGCGC	CGATACTCGTTCCAGGCCCG
Primers used to control the deletion of *mcpI*		ACATCGGGCTCTCCGAACGC	AATCTAGCGGCCTCGCTCCG
Primers used to control the deletion of *mcpJ*		GATAGCGCATCCAGTCAGC	GTTCGCGTTGCCTTCACCG
Primers used to control the deletion of *mcpK*		CTCATCCAGGTTCGTGAACCCGAGC	ATGGTCTGAGGACCCAGGCTCTCGC
Primers used to control the deletion of *mcpL*		GGCGATGATCTGAGTGTCGC	GCTTTCCGAGGAGCAGTTCG
Primers used to control the deletion of *mcpM*		TTCGCGCATCACTGAGATGC	AGACGGTGTGATCCTCACCCCGC
Primers used to control the deletion of *mcpN*		GGGTGACCACCTTGATAAGC	TTCTGAACCTGATCCCCACG
Primers used to control the deletion of *mcpO*		GTGATCACCGGCGACTTGC	GGCCTGTTGCTGGTCTTCG
Primers used to control the deletion of *mcpP*		GAGGGTCATCAAGCGAAGC	GGATCGTGTGTGAGACCAGC
Primers used to control the deletion of *mcpQ*		AGGCGATCCGCTACATGGAGATCGGC	ACAGCAGGGCCACAATCGCCATGC
Primers used to control the deletion of *mcpR*		GGATGGTCAAGCGCTACGGC	CCATGTGGATGTCGTGGCCG
Primers used to overexpress *mcpR*	pMR10 (with XhoI and BamHI)	CCCTCGAGttgCGGGCGCGGCGTGC	CGGGATCCCGGACAAGCGCTACACGCTT
Primers used to overexpress *mcpR*, *mcpRH99A* and *mcpRC75S* (for purification)	pET28a (with NdeI and BamHI)	GGAATTCcatatgttgCGGGCGCGGCG	CGggatccCGGACAAGCGCTACACGCTT
Primers used for the H99A mutation (with Q5 Site-Directed Mutagenesis Kit)		GCGCGCGCTGgctCGCGGCGAGC	CAGAATTCGCGGTACTCATCGCTG
Primers used for the H99A mutation (with Q5 Site-Directed Mutagenesis Kit)		ATTGTCGGGTctCCCCACAGCCTG	CTCCGGAGACGCGTAGCC
Primers used to control the deletion of *mcps*		ATCTCTACGCCGAGGAAGGC	GGCGAAAGAGCCGCAAAGC
Primers used to overexpress *sodB*	pMR10 (with XbaI and EcoRI)	GCTCTAGAatgCAACGCTTGCTGACCG	CGGAATTCctaGCGCGCGGCGGCG
Primers used to overexpress *katG*	pMR10 (with XbaI and XhoI)	GCTCTAGAgtgATGGAAACACCTGCGCG	CCCTCGAGctaAGCCGCCAGATCCAGGC