1   **Sofosbuvir and its tri-phosphate metabolite inhibit the RNA-dependent RNA polymerase activity of non-Structural protein 5 from the Kyasanur forest disease virus**

4   Mansi Malik, Parvathy Vijayan, Deepak K Jagannath, Rakesh K Mishra, Anirudha Lakshminarasimhan*

6   **Correspondence**

7   Tata Institute for Genetics and Society, NCBS campus, GKVK, Bellary Road, Bengaluru - 560065, KA, India

9   *Email: anirudha.lakshminarasimhan@gmail.com

Abstract

Kyasanur forest disease is a neglected zoonotic disease caused by a single-stranded RNA-based flavivirus, the incidence of which was first recorded in 1957 in the Southern part of India. Kyasanur forest disease virus is transmitted to monkeys and humans through the infected tick bite of *Haemophysalis spinigera*. Kyasanur forest disease is a febrile illness, which in severe cases, results in neurological complications leading to mortality. The current treatment regimens are symptomatic and supportive, and no targeted therapies are available for this disease. In this study, we evaluated the ability of FDA-approved drugs sofosbuvir (and its active metabolite) and Dasabuvir to inhibit the RNA-dependent RNA polymerase activity of NS5 protein from the Kyasanur forest disease virus. NS5 protein containing the N-terminal methyl transferase domain and C-terminal RNA-dependent RNA polymerase domain was expressed in *Escherichia coli*, and RNA-dependent RNA polymerase activity was demonstrated with the purified protein. The RNA-dependent RNA polymerase assay conditions were optimized, followed by the determination of apparent $K_{m,ATP}$ to validate the enzyme preparation. Half maximal-inhibitory concentrations against RNA-dependent RNA polymerase activity were determined for Sofosbuvir and its active metabolite. Dasabuvir did not show detectable inhibition with the tested conditions. This is the first demonstration of the inhibition of RNA-dependent RNA polymerase activity of NS5 protein from the Kyasanur forest disease virus with small molecule inhibitors. These initial findings can potentially facilitate the discovery and development of targeted therapies for treating Kyasanur forest disease.

Keywords

Kyasanur forest disease, Kyasanur forest disease virus, Non-structural protein 5, RNA-dependent RNA polymerase, Sofosbuvir, Flavivirus

Abbreviations

KFD – Kyasanur forest disease

KFDV – Kyasanur forest disease virus

NS5 – Non-structural protein 5

RdRp – RNA-dependent RNA polymerase

MTase – Methyl transferase

HCV – Hepatitis C virus

Introduction

Kyasanur forest disease (KFD) is an infectious viral hemorrhagic disease primarily infecting human and non-human primates [1]. Mortality in monkeys, especially among the black-faced langurs (*Presbytis entellus*) and red-faced bonnet monkeys (*Macaca radiata*), has been reported during multiple outbreaks of this disease [2]. In addition, birds, squirrels, gerbils, mice, rats, and bats also function as hosts for the virus [3]. The hard-bodied tick, *Haemaphysalis spinigera*, is the primary vector for the Kyasanur forest disease virus (KFDV) [4]. KFDV was also isolated from various genera of ticks, such as *Ixodes, Argas, Ornithodoros, Hyalomma, Dermacentor, and Rhipicephalus* [3]. Clinical symptoms of KFD include

high-grade fever, chills, headache, Conjunctival inflammation, vomiting, abdominal pain, diarrhea, maculopapular eruptions, muscle tenderness, nausea, and persistent cough [5]. In extreme cases, neurological and hemorrhagic manifestations are observed, with an estimated fatality of ~ 3-5% [6]. A microchip-based point-of-care test (Truenat KFD), developed by ICMR-NIV and Molbio Diagnostics, Goa, is currently the gold standard for diagnosis in India [7]. Formalin inactivated tissue-culture vaccine is available for KFD, administered in two doses to individuals aged 7–65 years at one-month intervals [8]. However, the immunity conferred by this vaccine is short-lived because of which booster doses are recommended after 6–9 months of primary vaccination for five consecutive years [8]. Current treatment regimens for KFD are limited to providing symptomatic relief and management [6]. There are no direct-acting antiviral drugs available against KFD. Failure of optimal coverage provided by the vaccine, with the absence of antiviral interventions against KFDV, makes it a public health challenge, necessitating attention toward developing second-generation vaccines and targeted therapies.

KFDV is a spherical (~40-65 nm in size) virus containing an icosahedral nucleocapsid enclosing a positive-sense RNA of 10,774 bases [9]. Kyasanur forest disease virus genome is closely related to Alkhurma Virus from Saudi Arabia, Omsk hemorrhagic fever virus from Siberia, and Nanjianyin Virus from China [10-12]. The RNA strand codes for a polyprotein comprising three structural (C, E & M) and seven non-structural proteins (NS1, NS2a, 2b, NS3, NS4a, 4b, and NS5 [13]. NS5 belonging to the Flaviviridae family are bi-functional proteins comprising of an N-terminal methyl transferase

domain (MTase) and a C-terminal RNA-dependent RNA polymerase domain

(RdRp) separated by a linker [14, 15](Fig. 1A). MTase domain

methylates N-7 of guanine and 2'-OH ribose of viral RNA cap, in the

presence of S-adenosyl methionine as a co-factor. N-7 methylation is

critical for enhancing the efficiency of viral translation, and 2'-O

methylation is essential for the subversion of the innate immune

response during viral infection [15]. RdRp domain of flaviviruses

catalyzes the de novo template-dependent formation of phosphodiester

bonds in the presence of nucleoside triphosphate (NTP) and divalent

metal ions. The canonical structure of the Flaviviridae RdRp domain

consists of an encircled right hand containing three subdomains -

thumb, palm, and fingers folded, surrounding the active site. RdRps

consist of channels, the entry of which is lined with positively

charged residues, facilitating NTPs and template RNA entry into the

active site [14]. The absence of proofreading exonuclease activity

in RdRps may lead to the incorporation of mutations and,

consequently, the selection of variants of viruses [16]. Hence, RdRp

of NS5 plays a crucial role in viral genome replication and can

serve as a potential antiviral drug target [17]. Nucleoside and non-

nucleoside inhibitors against NS5 RdRp from different viruses

belonging to the Flaviviridae family have been identified [18-25].

Sofosbuvir is an FDA-approved drug currently in use for the

treatment of Hepatitis C virus (HCV) infection [26, 27]. In this

study, we recombinantly expressed KFDV NS5 protein and characterized

the purified protein to identify a potential inhibitor of RdRp

activity. Sofosbuvir (Pro-drug) and its metabolite were demonstrated

to inhibit the RdRp activity of the KFDV NS5 protein.

Materials and Methods

Expression and purification of NS5 from KFDV

Codon-optimized gene corresponding to 5 to 897 amino acids of KFDV

NS5 with an N-terminal Hexa-histidine tag was synthesized (Genscript

USA) and sub-cloned into pET-28a (+) vector between NdeI and HindIII

restriction sites. Transformants of Rosetta2(DE3) pLysS containing

pET28a – KFDV NS5 were grown to an OD of 2.0 and induced with 0.5 mM

IPTG for 16 hours at 18°C. Induced cells were centrifuged at 5000 g

for 10 minutes at 4°C, and the pellet was resuspended in a lysis

Buffer containing 20 mM Na_2HPO_4, 0.5 M NaCl, 50 mM L-Arginine, 10 mM

$MgSO_4$, 5 mM β-mercaptoethanol, 5 mM Imidazole, 1 mM PMSF, 10%

Glycerol, pH 7 [28]. The cell suspension was sonicated, and the

lysate was centrifuged at 20000 g for 60 minutes at 4°C. The

clarified lysate was filtered with Durapore PVDF 0.22 μm membrane

and loaded onto a Ni-NTA affinity column pre-equilibrated with 20 mM

Na_2HPO_4, 0.5 M NaCl, 10 % Glycerol pH 7.0. The column was washed with

equilibration buffer containing 50 mM Imidazole and bound protein

eluted with 20 mM Na_2HPO_4, 0.5 M NaCl, 400 mM Imidazole, 10% Glycerol

pH 7.0. Eluates containing a prominent band corresponding to the

expected molecular weight of ~104 kDa were injected into Superdex

200 Increase 10/300 GL column pre-equilibrated with 50 mM Tris-HCl

pH 8.0, 150 mM NaCl, 5 mM $MgCl_2$, 1 mM DTT, 5% Glycerol, 0.05% Tween

20. The purified protein was quantified using Bradford assay and

densitometry analysis. Purified fractions were transferred onto the

PVDF membrane and probed with mouse anti-His primary antibody

(1:3000) and anti-mouse IgG HRP—conjugate (1:6000). Gel peptic

digest and LC-MS/MS analysis were performed for the gel filtration

fractions to ascertain the identity of the NS5 protein.

RNA-dependent RNA polymerase activity of purified NS5

RNA-dependent RNA polymerase (RdRp) activity of KFDV NS5 protein was

determined by employing a fluorometric endpoint assay based on the

intercalating property of a fluorescent dye, SYTO 9, which binds to

double-stranded (ds) RNA and not to single-stranded (ss) RNA [29].

250 nM of purified NS5 protein was added to a standard reaction mix

comprising 50 mM HEPES pH 7.0, 2.5 mM $MnCl_2$, 500 μM ATP, 20 μg/mL

poly(U), and 0.1mg/mL BSA, and the reaction mix was incubated at

30°C for 60 minutes in 96 well flat-bottom black plates. The

reaction was quenched by adding 25 mM EDTA, following which 0.25 μM

SYTO 9 was added and further incubated at room temperature for 7

minutes. Fluorescence was recorded in a plate reader (Infinite 200

Pro M Plex, TECAN, Switzerland) using excitation and emission

filters at 485 nm and 520 nm, respectively. The background

fluorescence levels were subtracted from the samples using a protein

blank control reaction. The effect of pH, temperature, NaCl, and

glycerol concentrations on RdRp activity was determined to identify

the optimal conditions using the purified KFDV NS5 protein [30].

Buffers with pH ranging from 5.0 to 9.0 (Citrate pH 5.0, Tricine pH

6.0, HEPES pH 7.0, Tris-HCl PH 7.5, Tris-HCl pH 8.0, and Bicine pH

9.0) were assayed. NS5 protein was incubated at different

temperatures of 20°C, 25°C, 30°C, 37°C, and 42°C, different NaCl

concentrations (5 mM, 25 mM, 50 mM, 75 mM, 100 mM, 300 mM, and 500

mM) and varying percentages of glycerol (5% v/v, 10% v/v, 20% v/v,

30% v/v, 40% v/v, 50% v/v). The mean values were computed from the

experiments with standard deviation.

NS5 protein was titrated against a fixed concentration of 500 μM ATP

to arrive at the protein concentration of 250 nM to determine

apparent $K_{m,ATP}$. At concentrations ranging from 50 μM to 500 μM, ATP

was titrated with 250 nM of purified NS5 protein and 20 μg/mL of

poly(U). Mean values were computed with standard deviations (n = 3),

and K_m was calculated using GraphPad Prism, 9.3.0.

Determination of inhibitory concentrations for sofosbuvir, its

metabolite, and dasabuvir

Sofosbuvir, Sofosbuvir triphosphate, and Dasabuvir (MedChemExpress,

Monmouth Junction, NJ) were dissolved in DMSO and diluted freshly

for the experiment. The inhibitory effect of Sofosbuvir, sofosbuvir

triphosphate, and Dasabuvir was determined at different

concentrations ranging from 5 μM to 20 μM. The assay was performed

in presence of 250 nM of NS5 protein, 20 μg/mL poly(U) and 500 μM

ATP in the reaction buffer containing 50 mM HEPES pH 7.0, 5 mM NaCl,

10% glycerol, 0.1 mg/ml BSA. IC_{50} values were determined using a non-

linear regression model with GraphPad Prism, 9.3.0. The mean values

were computed with standard deviation (n = 3).

Results

KFDV NS5 was expressed in bacteria and purified with affinity column

and size exclusion chromatography

Multiple sequence alignment of KFDV polyprotein (Uniprot id: D7RF80)

with NS5 proteins from tick-borne encephalitis virus (TBEV), Yellow

fever virus (YEFV), Zika virus (ZKV), Japanese encephalitis virus (JAEV), West Nile virus (WNV) and Dengue virus serotype 2 (DENV2), led to the identification of KFDV NS5 protein, i.e., between amino acids, 2514 to 3416 in the polyprotein sequence. NS5 protein is widely conserved across the Flaviviridae family (50 – 70% identity/similarity). Residues 5 to 897 of NS5 protein containing the MTase and RdRp domain (Fig. 1A) were expressed with an N-terminal Hexa histidine tag in bacteria and purified with Ni-NTA affinity chromatography (Fig. 1B). A lower molecular weight contaminant band (Fig. 1) was observed along with the band corresponding to the expected molecular weight of NS5 (~104 kDa), on the Coomassie blue stained SDS-PAGE gel. Anti-His antibody also recognized the lower molecular weight contaminant band with western blotting, suggesting it to be an N-terminal degradation product. It was confirmed with amino acid sequencing of tryptic digests with LC-MS/MS. The sequence corresponded to 5 to 282 amino acids of the NS5 protein (Data not shown). This histidine-tagged N-terminal fragment co-eluting with NS5 protein in affinity chromatography was removed with superdex 200 increase 10/300 GL size exclusion column chromatography (Peak P2 from Fig. 1C) to the extent that it was not observed in a Coomassie-stained SDS-PAGE gel. The identity of the NS5 protein - Peak, P1 corresponding to the expected molecular weight of 104 kDa, was confirmed with LC-MS/MS analysis and western blot analysis using an anti-His antibody (Fig. 1D). The yield of NS5 protein after size exclusion chromatography (P1 fraction) was 1 mg/liter of bacterial culture.

Apparent $K_{m,ATP}$ for NS5 RDRP from KFDV is comparable to that determined for other viruses

KFDV NS5 construct containing both the MTase and RdRp domains was

used in this study to rule out any effect of the MTase domain on

RdRp activity. The interplay of both the RdRp and MT domains was

earlier demonstrated for Dengue and Zika NS5 [31-33]. Compared to 5%

glycerol, KFDV NS5 showed an increase in RdRp activity at 10%

glycerol, which significantly reduced at 20% (Fig 2). Inhibition of

RdRp activity of KFDV NS5 was observed at lower concentration ranges

of NaCl consistent with an earlier report from Usutu virus NS5 [30].

RdRp from HCV tolerated NaCl concentrations until 100 mM without

significant RdRp activity loss [34]. However, it is unclear how

Dengue and KFDV/Usutu NS5 proteins differ in their sensitivity to

different concentrations of NaCl. Mechanistically, the binding of

Dengue NS5 to ssRNA has been shown to release six Na^+ ions [35].

Based on this report, the loss of activity can be speculated to be

due to the dissociation of RNA at higher concentrations of NaCl and

not due to the denaturation of protein. At a pH of 7.0, maximal

activity was observed, tapered at pHs 5.0 and 9.0. The optimal

activity was observed at a temperature of 30°C at pH 7, without

significant activity loss at 42°C (Fig 2). Apparent $K_{m,ATP}$ was

determined to be 42.15 $\pm$ 5.81 µM for KFDV NS5, under the tested

conditions, as described in methods (Fig. 3). The reported values of

$k_{m,GTP}$ for glutathionylated Dengue and Zika NS5 were 189.8 $\pm$ 52.5 µM,

and 137.2 $\pm$ 41.3 µM, respectively [36]. Motif F from the ring domain

of RdRp with highly conserved Lys, Glu, Arg residues, and motif G

from pinky finger domain, with conserved ala, binds to NTPs, and

+1/+2 junction of the template strand, respectively (Fig. 4) [37].

The comparable values of apparent $K_{m,ATP}$ to $K_{m,GTP}$ validates the use of

the purified protein for inhibition studies.

Sofosbuvir and sofosbuvir triphosphate inhibit the RDRP activity of KFDV NS5.

With HCV NS5, the half-maximal inhibitory concentrations of sofosbuvir triphosphate and sofosbuvir were reported to be 0.12 μM and 2.06 ug/mL (corresponding to 3.8905 μM), respectively (24, 57). X-ray crystal structure of HCV NS5B with sofosbuvir (PDB: 4WTG) showed a critical interaction of Asn291 with the 2'-F group of Sofosbuvir, leading to its incorporation into the growing RNA chain and, subsequently, chain termination (Fig. 4) [39]. As indicated earlier, active site residues in the Flaviviridae family are highly conserved, including Asn291 increasing the likelihood of Sofosbuvir inhibiting KFDV NS5. Sofosbuvir and sofosbuvir triphosphate exhibited IC_{50} of 3.45 ± 0.012 and 3.73 ± 0.033 μM, respectively against KFDV NS5 (Fig. 5). Interestingly, Dasabuvir, a nonnucleoside analog, did not show inhibition of RdRp activity, even at 100 μM, under the tested conditions. IC_{50} of Dasabuvir varied between nanomolar and micromolar concentrations when tested against different genotypes of HCV RdRps [25]. Also, Dasabuvir is not predicted to bind to conserved residues in RdRps across other viruses [40].

Discussion

Kyasanur forest disease virus is a high-risk category pathogen endemic to southern India. The current line of treatment for KFD includes maintaining hydration, blood work, and management of neurological and hemorrhagic conditions [5,6]. Sofosbuvir, also known as L-Alanine, N-[[P(S),2'R]-2'-deoxy-2'-fluoro-2'-methyl-P-

phenyl-5'-uridylyl]-,1-methyl ethyl ester ($C_{22}H_{29}FN_3O_9P$), is an orally

available nucleotide analog, approved for the treatment of HCV viral

infections [26]. In preclinical studies, Sofosbuvir has exhibited

inhibitory activity against all genotypes of HCV [41]. Sofosbuvir

has been indicated for chronic HCV infection (genotype 1, 2, 3, or

4) in adults as part of combination therapy and in children older

than 3 years with genotype for chronic HCV infections with

ribavirin. Many Clinical trials are ongoing for evaluating

sofosbuvir in different populations and dosing regimens in

combination with other drugs against HCV [42 – 46]. Sofosbuvir also

exhibited inhibitory activity against Dengue, Chikungunya, Yellow

fever virus, Hepatitis E virus, and SARS-CoV2 [47 -51]. Sofosbuvir

(PSI-7977) is metabolized to its triphosphate metabolite, β-D-2'-

deoxy-2'-fluoro-2'-C-methyluridine 5'-triphosphate (PSI-7409), by

hydrolysis of the carboxyl ester moiety, spontaneous release of the

Phenyl group, deamidation and phosphorylation [52, 53]. KFDV NS5

expression construct used in this study also included the MTase

domain, as the MTase was demonstrated to influence the activity RdRp

domain by interaction with the template RNA [38]. KFDV NS5 protein

was recombinantly expressed in bacteria, and so were NS5 proteins

from Usutu, Zika, Japanese encephalitis, Dengue, HCV, and West Nile

virus [30, 33, 37, 54, 55, 56]. Recombinantly purified NS5 RDRP

protein from the Zika virus exhibited an IC_{50} of 0.38 $\pm$ 0.03 μM

against sofosbuvir triphosphate [56, 47]. The inhibitory

concentration (IC_{50}) of sofosbuvir triphosphate and sofosbuvir were

in the same range for KFDV NS5. This could be due to differences in

the architecture of sofosbuvir binding sites across viruses.

This is the first report on the expression and purification of the

NS5 protein of KFDV, its assay establishment, and the demonstration of inhibition of RdRp activity by sofosbuvir. Further, the established biochemical assay can screen potential inhibitors of KFDV RdRp. Drug repurposing approaches against KFD, a neglected disease, can pave the way for identifying novel inhibitors for developing targeted therapeutic interventions against the disease.

Acknowledgments

The authors thank Tata Trusts for funding this study. We would like to acknowledge the NCBS-inStem-CCAMP mass spectrometry facility for the technical assistance provided for this study.

Conflict of interest

The authors declare no conflict of interest.

Availability of data and materials

All the relevant data used to support the findings of this study are included in the article.

References

[1]   T.H. Work, H. Trapido, D.P. Narasimha Murthy, R. Laxmana Rao, P.N. Bhatt, K.G. Kul-karni, Kyasanur Forest Disease III. A Preliminary Report on the Nature of the Infection and Clinical

Manifestations in Human Beings., Indian J. Med. Sci. 11 (1957) 619–
45.

[2] K. V. Shah, C.N. Dandawate, P.N. Bhatt, Kyasanur forest
disease virus: viremia and challenge studies in monkeys with
evidence of cross-protection by Langat virus infection,
F1000Research. 1 (2012). https://doi.org/10.12688/F1000RESEARCH.1-
61.V1.

[3] K. Muraleedharan, Wildlife Arthropods of Karnataka with
Special Reference to KFD Endemic Area of Shivamogga District: Those
Parasitic on Small and Large Mammals, (2016).
www.jakraya.com/journal/vri.

[4] M.G.R. Varma, H.E. Webb, K.M. Pavri, Studies on the
transmission of Kyasanur Forest disease virus by Haemaphysalis
spinigera newman, Trans. R. Soc. Trop. Med. Hyg. 54 (1960) 509–516.
https://doi.org/10.1016/0035-9203(60)90024-9.

[5] A. Munivenkatappa, R.R. Sahay, P.D. Yadav, R. Viswanathan,
D.T. Mourya, Clinical & epidemiological significance of Kyasanur
forest disease, Indian J. Med. Res. 148 (2018) 145.
https://doi.org/10.4103/IJMR.IJMR_688_17.

[6] Kyasanur forest disease, Centers for Disease Control and
Prevention, National Center for Emerging and Zoonotic Infectious
Diseases (NCEZID), Division of High-Consequence Pathogens and
Pathology (DHCPP), Viral Special Pathogens Branch (VSPB).
https://www.cdc.gov/vhf/kyasanur/index.html

[7] T. Majumdar, A. Shete, P. Yadav, S. Patil, D. Mali, A.
Waghmare, P. Gawande, Point of care real-time polymerase chain

reaction-based diagnostic for Kyasanur forest disease, Int. J. Infect. Dis. 108 (2021) 226–230. https://doi.org/10.1016/J.IJID.2021.05.036.

[8] G.S. Kasabi, M. V. Murhekar, V.K. Sandhya, R. Raghunandan, S.K. Kiran, G.H. Channabasappa, S.M. Mehendale, Coverage and Effectiveness of Kyasanur Forest Disease (KFD) Vaccine in Karnataka, South India, 2005-10, PLoS Negl. Trop. Dis. 7 (2013) e2025. https://doi.org/10.1371/JOURNAL.PNTD.0002025.

[9] K.A. Dodd, B.H. Bird, M.L. Khristova, C.G. Albariño, S.A. Carroll, J.A. Comer, B.R. Erickson, P.E. Rollin, S.T. Nichol, Ancient Ancestry of KFDV and AHFV Revealed by Complete Genome Analyses of Viruses Isolated from Ticks and Mammalian Hosts, PLoS Negl. Trop. Dis. 5 (2011) e1352. https://doi.org/10.1371/JOURNAL.PNTD.0001352.

[10] B. Bhatia, H. Feldmann, A. Marzi, Kyasanur Forest Disease and Alkhurma Hemorrhagic Fever Virus—Two Neglected Zoonotic Pathogens, Microorg. 2020, Vol. 8, Page 1406. 8 (2020) 1406. https://doi.org/10.3390/MICROORGANISMS8091406.

[11] D. Lin, L. Li, D. Dick, R.E. Shope, H. Feldmann, A.D.T. Barrett, M.R. Holbrook, Analysis of the complete genome of the tick-borne flavivirus Omsk hemorrhagic fever virus, Virology. 313 (2003) 81–90. https://doi.org/10.1016/S0042-6822(03)00246-0.

[12] J. Wang, H. Zhang, S. Fu, H. Wang, D. Ni, R. Nasci, Q. Tang, G. Liang, Isolation of Kyasanur Forest Disease Virus from Febrile Patient, Yunnan, China, Emerg. Infect. Dis. 15 (2009) 326. https://doi.org/10.3201/EID1502.080979.

[13] K. Venugopal, T. Gritsun, V.A. Lashkevich, E.A. Gould,
Analysis of the structural protein gene sequence shows Kyasanur
Forest disease virus as a distinct member in the tick-borne
encephalitis virus serocomplex, J. Gen. Virol. 75 (1994) 227–232.
https://doi.org/10.1099/0022-1317-75-1-227.

[14] J. Wu, W. Liu, P. Gong, P. Gong, A Structural Overview of RNA-
Dependent RNA Polymerases from the Flaviviridae Family, Int. J. Mol.
Sci. 2015, Vol. 16, Pages 12943-12957. 16 (2015) 12943–12957.
https://doi.org/10.3390/IJMS160612943.

[15] H. Dong, K. Fink, R. Züst, S.P. Lim, C.F. Qin, P.Y. Shi,
Flavivirus RNA methylation, J. Gen. Virol. 95 (2014) 763–778.
https://doi.org/10.1099/VIR.0.062208-0/CITE/REFWORKS.

[16] S.F. Elena, R. Sanjuán, Adaptive Value of High Mutation Rates
of RNA Viruses: Separating Causes from Consequences, J. Virol. 79
(2005) 11555–11558. https://doi.org/10.1128/JVI.79.18.11555-
11558.2005.

[17] P.O. Fernandes, M.A. Chagas, W.R. Rocha, A.H. Moraes, Non-
structural protein 5 (NS5) as a target for antiviral development
against established and emergent flaviviruses, Curr. Opin. Virol. 50
(2021) 30–39. https://doi.org/10.1016/J.COVIRO.2021.07.001.

[18] X. Yao, S. Guo, W. Wu, J. Wang, S. Wu, S. He, Y. Wan, K.S.
Nandakumar, X. Chen, N. Sun, Q. Zhu, S. Liu, Q63, a novel DENV2 RdRp
non-nucleoside inhibitor, inhibited DENV2 replication and infection,
J. Pharmacol. Sci. 138 (2018) 247–256.
https://doi.org/10.1016/J.JPHS.2018.06.012.

[19] R. Arora, C.W. Liew, T.S. Soh, D.A. Otoo, C.C. Seh, K. Yue, S.

Nilar, G. Wang, F. Yokokawa, C.G. Noble, Y.L. Chen, P.-Y. Shi, J.

Lescar, T.M. Smith, T.E. Benson, S.P. Lim, Two RNA Tunnel Inhibitors

Bind in Highly Conserved Sites in Dengue Virus NS5 Polymerase:

Structural and Functional Studies, J. Virol. 94 (2020).

https://doi.org/10.1128/JVI.01130-20

[20] Y. Chen, Z. Li, P. Pan, Z. Lao, J. Xu, Z. Li, S. Zhan, X. Liu,

Y. Wu, W. Wang, G. Li, Cinnamic acid inhibits Zika virus by

inhibiting RdRp activity, Antiviral Res. 192 (2021) 105117.

https://doi.org/10.1016/J.ANTIVIRAL.2021.105117.

[21] Y. Lin, H. Zhang, W. Song, S. Si, Y. Han, J. Jiang,

Identification and characterization of Zika virus NS5 RNA-dependent

RNA polymerase inhibitors, Int. J. Antimicrob. Agents. 54 (2019)

502–506. https://doi.org/10.1016/J.IJANTIMICAG.2019.07.010.

[22] P. Coulerie, M. Nour, A. Maciuk, C. Eydoux, J.C. Guillemot, N.

Lebouvier, E. Hnawia, K. Leblanc, G. Lewin, B. Canard, B. Figadère,

Structure-Activity Relationship Study of Biflavonoids on the Dengue

Virus Polymerase DENV-NS5 RdRp, Planta Med. 79 (2013) 1313–1318.

https://doi.org/10.1055/S-0033-1350672.

[23] M.A. Abu-Zaied, S.F. Hammad, F.T. Halaweish, G.H. Elgemeie,

Sofosbuvir Thio-analogues: Synthesis and Antiviral Evaluation of the

First Novel Pyridine- And Pyrimidine-Based Thioglycoside

Phosphoramidates, ACS Omega. 5 (2020) 14645–14655.

https://doi.org/10.1021/ACSOMEGA.0C01364

[24] Fenaux, M., Lin, X., Yokokawa, F., Sweeney, Z., Saunders, O.,

Xie, L., ... & Zhong, W. (2016). Antiviral nucleotide incorporation

by recombinant human mitochondrial RNA polymerase is predictive of

increased in vivo mitochondrial toxicity risk. Antimicrobial agents

and chemotherapy, 60(12), 7077-7085.

[25] W. Kati, G. Koev, M. Irvin, J. Beyer, Y. Liu, P. Krishnan, T.

Reisch, R. Mondal, R. Wagner, A. Molla, C. Maring, C. Collins, In

vitro activity and resistance profile of dasabuvir, a nonnucleoside

hepatitis c virus polymerase inhibitor, Antimicrob. Agents

Chemother. 59 (2015) 1505–1511. https://doi.org/10.1128/AAC.04619-

14.

[26] I. Gentile, F. Borgia, A. R. Buonomo, G. Castaldo, G. Borgia,

A Novel Promising Therapeutic Option Against Hepatitis C Virus: An

Oral Nucleotide NS5B Polymerase Inhibitor Sofosbuvir, Curr Med Chem

20(30) 3733-3742. https://doi.org/10.2174/09298673113209990178.

[27] E. Lawitz, I.M. Jacobson, D.R. Nelson, S. Zeuzem, M.S.

Sulkowski, R. Esteban, D. Brainard, J. Mcnally, W.T. Symonds, J.G.

Mchutchison, D. Dieterich, E. Gane, Development of sofosbuvir for

the treatment of hepatitis C virus infection, Ann. N. Y. Acad. Sci.

1358 (2015) 56–67. https://doi.org/10.1111/NYAS.12832.

[28] R.P. Dos Santos Alves, L.R. Pereira, D.L.N. Fabris, F.S.

Salvador, R.A. Santos, P.M. De Andrade Zanotto, C.M. Romano, J.H.

Amorim, L.C. De Souza Ferreira, Production of a recombinant dengue

virus 2 NS5 protein and potential use as a vaccine antigen, Clin.

Vaccine Immunol. 23 (2016) 460–469.

https://doi.org/10.1128/CVI.00081-16.

[29] Y. Sáez-Álvarez, A. Arias, C. del Águila, R. Agudo,
Development of a fluorescence-based method for the rapid
determination of Zika virus polymerase activity and the screening of
antiviral drugs, Sci. Reports 2019 91. 9 (2019) 1–11.
https://doi.org/10.1038/s41598-019-41998-1.

[30] L. Albentosa-González, P. Clemente-Casares, R. Sabariegos, A.
Mas, Polymerase activity, protein-protein interaction, and cellular
localization of the Usutu virus NS5 protein, Antimicrob. Agents
Chemother. 64 (2020). https://doi.org/10.1128/AAC.01573-19.

[31] Y. Zhao, T.S. Soh, J. Zheng, K.W.K. Chan, W.W. Phoo, C.C. Lee,
M.Y.F. Tay, K. Swaminathan, T.C. Cornvik, S.P. Lim, P.Y. Shi, J.
Lescar, S.G. Vasudevan, D. Luo, A Crystal Structure of the Dengue
Virus NS5 Protein Reveals a Novel Inter-domain Interface Essential
for Protein Flexibility and Virus Replication, PLOS Pathog. 11
(2015) e1004682. https://doi.org/10.1371/JOURNAL.PPAT.1004682.

[32] S.P. Lim, J.H.K. Koh, C.C. Seh, C.W. Liew, A.D. Davidson, L.S.
Chua, R. Chandrasekaran, T.C. Cornvik, P.Y. Shi, J. Lescar, A
crystal structure of the dengue virus non-structural protein 5 (NS5)
polymerase delineates interdomain amino acid residues that enhance
its thermostability and de novo initiation activities, J. Biol.
Chem.   288   (2013)   31105–31114.
https://doi.org/10.1074/jbc.M113.508606.

[33] B. Zhao, G. Yi, F. Du, Y. C. Chuang, R. C. Vaughan, B.
Sankaran, et al., Structure and function of the Zika virus full-
length NS5 protein, Nat. Commun. 8 (2017) 14762.
https://doi:10.1038/ncomms14762.

[34] V. Lohmann, A. Roos, F. Körner, J.O. Koch, R. Bartenschlager,

Biochemical and Kinetic Analyses of NS5B RNA-Dependent RNA

Polymerase of the Hepatitis C Virus, Virology. 249 (1998) 108–118.

https://doi.org/10.1006/VIRO.1998.9311.

[35] M.R. Szymanski, M.J. Jezewska, P.J. Bujalowski, C. Bussetta,

M. Ye, K.H. Choi, W. Bujalowski, Full-length Dengue Virus RNA-

dependent RNA Polymerase-RNA/DNA Complexes, J. Biol. Chem. 286

(2011) 33095–33108. https://doi.org/10.1074/jbc.m111.255034.

[36] C. Saisawang, A. Kuadkitkan, P. Auewarakul, D.R. Smith, A.J.

Ketterman, Glutathionylation of dengue and Zika NS5 proteins affects

guanylyltransferase and RNA dependent RNA polymerase activities,

PLoS   One.   13   (2018)   e0193133.

https://doi.org/10.1371/JOURNAL.PONE.0193133.

[37]   G. Lu, P. Gong, Crystal Structure of the Full-Length

Japanese   Encephalitis   Virus   NS5   Reveals   a   Conserved

Methyltransferase-Polymerase   Interface,   PLOS   Pathog. 9 (2013)

e1003549. https://doi.org/10.1371/JOURNAL.PPAT.1003549.

[38] T. Rusanov, T. Kent, M. Saeed, T.M. Hoang, C. Thomas, C.M.

Rice, R.T. Pomerantz, Identification of a Small Interface between

the Methyltransferase and RNA Polymerase of NS5 that is Essential

for Zika Virus Replication, Sci. Reports 2018 81. 8 (2018) 1–11.

https://doi.org/10.1038/s41598-018-35511-3.

[39] T.C. Appleby, J.K. Perry, E. Murakami, O. Barauskas, J. Feng,

A. Cho, D. Fox, D.R. Wetmore, M.E. McGrath, A.S. Ray, M.J. Sofia, S.

Swaminathan, T.E. Edwards, Structural basis for RNA replication by

the hepatitis C virus polymerase, Science (80-.). 347 (2015) 771–

775. https://doi.org/10.1126/SCIENCE.1259210.

[40] P.P.L.F. de Albuquerque, L.H.S. Santos, D. Antunes, E.R.

Caffarena, A.S. Figueiredo, Structural insights into NS5B protein of

novel equine hepaciviruses and pegiviruses complexed with polymerase

inhibitors, Virus Res. 278 (2020) 197867.

https://doi.org/10.1016/J.VIRUSRES.2020.197867.

[41] B. Han, R. Martin, S. Xu, A. Parvangada, E.S. Svarovskaia, H.

Mo, H. Dvory-Sobol, Sofosbuvir susceptibility of genotype 1 to 6 HCV

from DAA-naïve subjects, Antiviral Res. 170 (2019) 104574.

https://doi.org/10.1016/J.ANTIVIRAL.2019.104574.

[42] A.S. Price, A.K. Nelson, A. Ghosh, S. Kottilil, J. V. Chua, A

Phase 2 open label study of ledipasvir/sofosbuvir for 12 weeks in

subjects with hepatitis B virus infection, J. Med. Virol. (2022).

https://doi.org/10.1002/JMV.28105.

[43] V. Virlogeux, L. Choupeaux, P. Pradat, M. Maynard, F. Bailly,

C. Scholtès, M.C. Gagnieu, F. Zoulim, Sofosbuvir plus daclatasvir

with or without ribavirin for chronic hepatitis C infection: Impact

of drug concentration on viral load decay, Dig. Liver Dis. 48 (2016)

1351–1356. https://doi.org/10.1016/J.DLD.2016.07.014.

[44] M. H. El-Sayed, Clinical Pharmacokinetics of

Daclatasvir/Sofosbuvir in Adolescents with Hepatitis C Virus.

ClinicalTrials.gov. https://clinicaltrials.gov/ct2/show/NCT03540212.

[45] A. A. A. Elmoety, Evaluation of Sofosbuvir and Daclatasvir

Combo in COIVD-19 Patients in Egypt. ClinicalTrials.gov.

https://www.clinicaltrials.gov/ct2/show/NCT04773756.

[46]   C. Chappell, Safety, Tolerability, and Outcomes of

Velpatasvir/SofosbuviR in Treatment of Chronic Hepatitis C Virus

During   Pregnancy   (STORC).   Clinicaltrials.gov.

https://clinicaltrials.gov/ct2/show/NCT05140941.

[47] H.T. Xu, S.P. Colby-Germinario, S.A. Hassounah, C. Fogarty, N.

Osman, N. Palanisamy, Y. Han, M. Oliveira, Y. Quan, M.A. Wainberg,

Evaluation   of   Sofosbuvir   (β-D-2'-deoxy-2'-α-fluoro-2'-β-C-

methyluridine) as an inhibitor of Dengue virus replication, Sci.

Reports 2017 71. 7 (2017) 1–11. https://doi.org/10.1038/s41598-017-

06612-2.

[48] A.C. Ferreira, P.A. Reis, C.S. de Freitas, C.Q. Sacramento,

L.V.B. Hoelz, M.M. Bastos, M. Mattos, N. Rocha, I.G. de Azevedo

Quintanilha, C. da Silva Gouveia Pedrosa, L.R.Q. Souza, E.C. Loiola,

P. Trindade, Y.R. Vieira, G. Barbosa-Lima, H.C. de Castro Faria

Neto, N. Boechat, S.K. Rehen, K. Brüning, F.A. Bozza, P.T. Bozza,

T.M.L. Souza, Beyond members of the Flaviviridae family, sofosbuvir

also inhibits chikungunya virus replication, Antimicrob. Agents

Chemother. 63 (2019). https://doi.org/10.1128/AAC.01389-18.

[49] C.S. de Freitas, L.M. Higa, C.Q. Sacramento, A.C. Ferreira,

P.A. Reis, et al, Yellow fever virus is susceptible to sofosbuvir

both in vitro and in vivo, PLoS Negl. Trop. Dis. 13 (2019) e0007072.

https://doi.org/10.1371/JOURNAL.PNTD.0007072.

[50] Z. Liang, J. Shu, Q. He, F. Zhang, L. Dai, L. Wang, F. Lu, L.

Wang, High dose sofosbuvir and sofosbuvir-plus-ribavirin therapy

inhibit Hepatitis E Virus (HEV) replication in a rabbit model for

acute HEV infection, Antiviral Res. 199 (2022) 105274.

https://doi.org/10.1016/J.ANTIVIRAL.2022.105274.

[51] C.Q. Sacramento, N. Fintelman-Rodrigues, J.R. Temerozo, 579 A.D.P.D. Da Silva, S.D.S.G. Dias, et al, In vitro antiviral activity 580 of the anti-HCV drugs daclatasvir and sofosbuvir against SARS-CoV-2, 581 the aetiological agent of COVID-19, J. Antimicrob. Chemother. 76 582 (2021) 1874–1885. https://doi.org/10.1093/JAC/DKAB072.

583 [52] E. Murakami, T. Tolstykh, H. Bao, C. Niu, H.M. Micolochick 584 Steuer, D. Bao, W. Chang, C. Espiritu, S. Bansal, A.M. Lam, M.J. 585 Otto, M.J. Sofia, P.A. Furman, Mechanism of activation of PSI-7851 586 and its diastereoisomer PSI-7977, J. Biol. Chem. 285 (2010) 34337– 587 34347. https://doi.org/10.1074/jbc.M110.161802.

588 [53] S. Chen, J. Wu, X. Yang, Q. Sun, S. Liu, F. Rashid, E.E. 589 Dzakah, H. Wang, J. Wang, P. Gong, S. Tang, RNA-Dependent RNA 590 Polymerase of the Second Human Pegivirus Exhibits a High-Fidelity 591 Feature, Microbiol. Spectr. (2022). 592 https://doi.org/10.1128/SPECTRUM.02729-22.

593 [54] T. Rusanov, T. Kent, M. Saeed, T.M. Hoang, C. Thomas, C.M. 594 Rice, R.T. Pomerantz, Identification of a Small Interface between 595 the Methyltransferase and RNA Polymerase of NS5 that is Essential 596 for Zika Virus Replication, Sci. Reports 2018 81. 8 (2018) 1–11. 597 https://doi.org/10.1038/s41598-018-35511-3

[55] J.-W. Oh, T. Ito, M.M.C. Lai, A Recombinant Hepatitis C 599 Virus RNA-Dependent RNA Polymerase Capable of Copying the Full- 600 Length Viral RNA, J. Virol. 73 (1999) 7694–7702. 601 https://doi.org/10.1128/JVI.73.9.7694-7702.1999.

[56] Sacramento, C. Q., de Melo, G. R., de Freitas, C. S., Rocha, 603 N., Hoelz, L. V. B., Miranda, M., ... & Souza, T. M. L. (2017). The

clinically approved antiviral drug sofosbuvir inhibits Zika virus

replication. Scientific reports, 7(1), 1-12.

57. Abu-Zaied, M. A., Hammad, S. F., Halaweish, F. T., & Elgemeie,

G. H. (2020). Sofosbuvir Thio-analogues: Synthesis and Antiviral

Evaluation of the First Novel Pyridine-and Pyrimidine-Based

Thioglycoside Phosphoramidates. ACS omega, 5(24), 14645-14655.

Figure Legends

Fig. 1. NS5 protein of Kyasanur forest disease virus. A. Schematic

representation of NS5 protein domains - N-terminal methyl

transferase (MTase – 1 to 263) and C-terminal RNA dependent RNA

polymerase (RdRp – 274 to 903). Linker (L – 264 to 273 amino acids)

residues connecting the two domains are not conserved among

flaviviruses (35). B.   Coomassie-stained SDS-PAGE gel (12%)

containing Ni-NTA affinity column elutes. Lanes containing a

prominent band of the expected molecular weight of 104 kDa,

corresponding to 5 to 897 amino acids of NS5, are seen in the elutes

(indicated by an arrow). Also seen is a lower molecular weight band

between the molecular weight markers - 35 kDa and 25 kDa (*). C.

Superdex increase 200 10/300 GL fractionation of affinity purified

fraction of NS5 protein. Peaks P1 and P2 correspond to NS5 and the

lower molecular weight protein (*). Coomassie-stained SDS-PAGE gel

is on the left panel. D. anti- Histidine tagged antibody probed

western blot of P1 and P2 fractions. The prominent bands in P1 (NS5

protein indicated with arrowhead) and P2 (lower molecular weight

contaminant band marked with asterisks) were recognized by an anti-

His antibody.

Fig. 2. Optimization of reaction conditions for RdRp activity of

KFDV NS5. Effect of pH (A), NaCl (B), glycerol (C), and temperature

(D) on RdRp activity. Activity is represented as arbitrary

fluorescence units (A.U.) on the Y-axis. For every sample well,

fluorescence units were subtracted from a blank control devoid of

NS5 protein under the same conditions. The standard deviation was

computed and represented as error bars for every data point.

Fig. 3. Determination of apparent $K_{m,ATP}$ for RdRp domain of KFDV NS5.

Apparent K_m for ATP was determined by Michaelis-Menten equation

using graph pad prism 9.3.0. The representative graph contains

values averaged from at least three independent experiments (n=3).

Activity is represented as arbitrary fluorescence units (A.U.) on

the Y-axis. The standard deviation error bars are shown for every

data point.

Fig. 4. Amino acid sequence alignment of motifs F, G, and B of NS5

from the Flaviviridae family. Kyasanur forest disease virus (KFDV),

Alkhurma hemorrhagic fever (ALKV), Tick-borne encephalitis virus

(TBEV), Yellow fever virus (YEFV), Zika virus (ZKV), Japanese

encephalitis virus (JAEV), West Nile virus (WNV) Dengue virus

serotype 2 (DENV2), Hepatitis C virus (HCV). Conserved residues in

Motif F and motif G, from ring and pinky finger domains involved in

interacting with NTPs and the template, respectively, are

highlighted. Asn291 from the NS5 RdRp domain of HCV makes a critical

interaction with Sofosbuvir [33]. Asn291 is highly conserved across

the sequences shown in the figure (marked as *). There is a higher

level of conservation among the flavivirus genus than flaviviruses

and HCV (which belong to the genus Hepacivirus).

Fig. 5. Determination of IC_{50} for sofosbuvir (A) and sofosbuvir

triphosphate (B). Increasing concentrations of Sofosbuvir and

sofosbuvir triphosphate were titrated using the optimized assay

conditions described in the methods. The log of compound

concentration is plotted against percentage activity and $IC_{50,}$

computed from the profiles using Graph pad prism 9.3.0 (n=3). The

structures of compounds are shown above the IC_{50} graphs.

Motif F

KFDV	452	VYNMMG**K**R**E**KKLG----------EFGVAKGS**R**AIWYM 478
ALKV	452	VYNMMG**K**R**E**KKLG----------EFGVAKGS**R**AIWYM 478
TBEV	452	VYNMMG**K**R**E**KKLG----------EFGVAKGS**R**AIWYM 478
YEFV	452	EYNMMG**K**R**E**KKLS----------EFGKAKGS**R**AIWYM 478
ZKV	452	VYNMMG**K**R**E**KKQG----------EFGKAKGS**R**AIWYM 478
JAEV	453	IYNMMG**K**R**E**KKPG----------EFGKAKGS**R**AIWFM 479
WNV	453	IYNMMG**K**R**E**KKPG----------EFGKAKGS**R**AIWFM 479
DEN_V2	450	VYNMMG**K**R**E**KKLG----------EFGKAKGS**R**AIWYM 476
HCV	135	DTTIMA**K**S**E**VFCVQPEKGGEKDPAKGVEKRP**R**VIQYP 163

Motif G   Motif B

KFDV	403	KVRSNA**A**LGA 412	601	QRG**SG**QVV**T**YAL**N**TL	615
ALKV	403	KVRSNA**A**LGA 412	601	QRG**SG**QVV**T**YAL**N**TL	615
TBEV	403	KV**K**SNA**A**LGA 412	601	QRG**SG**QVV**T**YAL**N**TL	615
YEFV	403	KVRSHA**A**VGA 412	601	QRG**SG**QVV**T**YAL**N**TI	615
ZKV	403	KVRSNA**A**LGA 412	601	QRG**SG**QVV**T**YAL**N**TF	615
JAEV	404	KVNSNA**A**LGA 413	602	QRG**SG**QVV**T**YAL**N**TF	616
WNV	404	KVNSNA**A**LGA 413	602	QRG**SG**QVV**T**YAL**N**TF	616
DEN_V2	401	KVRSNA**A**LGA 411	599	QRG**SG**QVG**T**YGL**N**TF	613
HCV	90	LTPPHS**A**KSK 99	279	CRA**SG**VLT**T**SCG**N**TL	293