Fibroblastic cells are a site of mouse cytomegalovirus *in vivo* lytic replication and latent

persistence oppositely regulated by *Stat1*

Katarzyna M. Sitnik[1,2 §*], Fran Krstanović[3], Natascha Gödecke[1], Ulfert Rand[1], Tobias Kubsch[1], Henrike

Maaß[1], Yeonsu Kim[1], Ilija Brizić[3], Luka Čičin-Šain[1,4,5 §*]

[1] Department of Viral Immunology, Helmholtz Centre for Infection Research, 38124 Braunschweig,

Germany

[2] Institute of Animal Breeding and Genetics, University of Veterinary Medicine Vienna, 1210 Vienna,

Austria

[3] Center for Proteomics, Faculty of Medicine, University of Rijeka, 51000 Rijeka, Croatia

[4] Centre for Individualized Infection Medicine, a joint venture of HZI and MHH, 30625 Hannover,

Germany

[5] German Centre for Infection Research (DZIF), Hannover-Braunschweig site, 38124 Braunschweig,

Germany

§ Senior author

* Corresponding author

Correspondence: Luka Čičin-Šain, Luka.Cicin-Sain@helmholtz.hzi.de (Lead contact) and Katarzyna

Sitnik, Katarzyna.Sitnik@vetmeduni.ac.at

Abstract

To date, no herpesvirus has been shown to latently persist in fibroblastic cells. Here, we demonstrate

that mouse CMV (MCMV), a β-herpesvirus, persists for the long term and across organs in PDGFRα$^+$

fibroblastic cells, with similar or higher genome loads than in the previously known sites of MCMV

latency. Whereas MCMV gene transcription in PDGFRα$^+$ fibroblastic cells was almost completely

silenced at 5 months post-infection, these cells gave rise to reactivated virus *ex vivo*, arguing that they

supported latent MCMV infection. Notably, PDGFRα$^+$ fibroblastic cells also supported productive virus

replication during primary MCMV infection. Mechanistically, *Stat1*-deficiency resulted in increased lytic

but abolished latent infection of fibroblastic cells *in vivo*. In sum, fibroblastic cells have a dual role as a

site of lytic MCMV replication and a reservoir of latent MCMV *in vivo* and *Stat1* is critically involved in

the regulation of MCMV latency.

Introduction

Mouse CMV (MCMV) belongs to the genus *Muromegalovirus* in the subfamily

Betaherpesvirinae. It is most closely related to the primate-infecting viruses of the genus

Cytomegalovirus, comprising human CMV (HCMV). Together, CMVs are characterized by large DNA

genomes (>200 kb), induction of cytomegaly in infected cells and strict species tropism [1]. As for all

herpesviruses, the life cycle of CMVs comprises both lytic (productive) and latent (non-productive)

infection, the latter underpinning their ability to persist for life in their respective hosts. During lytic

infection the CMV genome is transcribed to express high levels of immediate–early (IE), early (E) and

late (L) viral gene products, added in a temporal sequence, that carry the virus through the phases of

genome replication and virion assembly, culminating in the release of infectious virions and cell lysis.

By contrast, during latent infection CMV genomic DNA persists in the cell nucleus but the viral progeny

is not produced. Viral gene transcription in latency is considerably silenced and the major CMV IE

promoter is associated with histones bearing repressive post-translational marks [2, 3]. Under

appropriate conditions, the virus productive cycle is reactivated, leading to the genesis of new

infectious progeny. A formal criterium of latency is the appearance of infectious virus upon, but not

prior to, co-cultivation of the population of interest with cells permissive to lytic infection. *In vivo*, lytic

CMV infection prevails for some time following virus inoculation, enabling viral amplification and

spread, but in immunocompetent hosts it is eventually extinguished, after which a life-long phase of *in*

vivo latency ensues when only latent virus can be detected [4].

Herpesviruses are recognized for using distinct cell types as sites of lytic replication and latent

persistence [5]. In line with this, myeloid progenitors and monocytes are well-characterised sites of

HCMV latency [6]. Latent infection of these cell types by MCMV is less clear, due to conflicting results

in studies [7, 8] and remains a controversial topic. On the other hand, and somewhat curiously for a

herpesvirus, latent MCMV was found to reside during *in vivo* latency in liver endothelial cells [8] and

peritoneal macrophages [9, 10], which represent cell types that are permissive to lytic MCMV infection

[9, 11]. Fibroblasts are also well-known to support lytic MCMV replication and are commonly used to

grow virus stocks [12]. Interestingly, experiments of Mercer *et al.* [13] and Pomeroy *et al.* [14]

performed more than 30 years ago, demonstrated enrichment for MCMV genomes and virus

reactivation from stroma-enriched spleen fragments, while essentially excluding the contribution of

hematopoietic cells. The authors proposed that splenic fibroblasts were involved but could not prove it

due to the lack of available methods for splenic fibroblast purification. Notably, the notion about

latency in fibroblastic cells has since not been investigated for any CMV or other herpesvirus, and

remains unconfirmed.

Like other viral infections, MCMV is crucially controlled by the host's response to interferons

(IFNs) [15], which signal through signal transducer and activator of transcription 1 (STAT1) [16]. Mice

deficient for *Stat1* or otherwise rendered unresponsive to IFNs succumb within a few days to infection

with only a few hundred plaque forming units (PFU) of MCMV [17, 18]. Besides orchestrating antiviral

immunity through effects on the functions of the cells of the immune system, IFNs directly inhibit

productive MCMV growth by elevating cell-intrinsic antiviral activity [15, 19, 20]. Notably, IFN treatment

inhibits the activity of MCMV major IE promoter, which controls the initiation of the lytic cycle [20]. We

have shown that MCMV infection in permissive cells infected at low multiplicity of infection (MOI) in the

presence of IFN is transiently repressed until IFN is removed. This reversible state satisfied the formal

definition of latency, where MCMV genomes are maintained in absence of replicating virus, but the

virus has the capacity to reactivate [20]. However, it remained unclear if similar phenomena may be

observed in a long-term *in vivo* model of CMV latency.

In this study, using state-of-the-art methods for the purification of tissue-resident subsets from

mouse organs, we comprehensively mapped the cellular sites of MCMV genome persistence during

latency *in vivo*. Our study unravels that MCMV commonly persists in PDGFRα$^+$ fibroblastic cells for the

long term, in a state that fulfils all the formal criteria of latent infection. We further show that PDGFRα$^+$

fibroblastic cells are simultaneously permissive to lytic MCMV infection and have a role in viral

reproduction during the acute phase of *in vivo* infection. Finally, we identify a physiological role of

Stat1 in promoting latent persistence of MCMV in fibroblastic cells *in vivo* whilst concurrently inhibiting

their lytic infection with the virus.

Results

PDGFRα^+ fibroblastic cells carry MCMV genomes during latent MCMV infection *in vivo*.

To address whether fibroblastic cells might be a site of MCMV latent persistence *in vivo*, we

initially performed a comprehensive characterisation of the distribution of MCMV genomes in tissue-

resident cells across mouse organs during latency *in vivo*. Specifically, we determined the number of

viral genome copies per 100,000 cells (henceforth referred to as MCMV genome load) in FACS-sorted

fibroblastic-, endothelial-, epithelial- and tissue-resident macrophage populations from the liver,

salivary glands (SG), spleen, visceral adipose tissue (VAT) and lungs of mice latently infected via

intraperitoneal route (Fig. 1a). Intraperitoneal administration of MCMV is the most widely used MCMV

infection model that results in infection of multiple organs due to virus systemic spread in the acute

phase of infection [21, 22]. Populations of tissue-resident macrophages (Mϕ), such as F4/80$^+$CD11c$^+$

SG Mϕ, lung alveolar Mϕ, liver Kupfer cells, spleen red pulp Mϕ and VAT Mϕ were identified using

organ-specific gating strategies [23-25]. Endothelial cells (EC) were identified as CD45$^-$ PDGFRα^-

CD31$^+$ cells and were shown to co-express CD146 in the liver (Supplementary Fig. 1b). Of note, EC

from VAT were not analysed as they could not be isolated in sufficient numbers. PDGFRα^+ fibroblastic

cells (PDGFRα^+ FC) were identified as CD45$^-$CD31$^-$PDGFRα^+ cells in VAT [26], CD45$^-$CD31$^-$EpCAM$^-$

PDGFRα^+ cells in the lungs [27] and SG [28]; and as CD45$^-$CD31$^-$ITGB1$^+$PDGFRα^+ cells in the spleen

[29]. In the liver we focused on a defined CD45$^-$CD31$^-$EpCAM$^-$PDGFRα^+ FC subset, vitamin A$^+$ liver

stellate cells [30, 31]. We chose to focus on FC that express PDGFRα since they are relatively well-

defined across mouse organs [32]. For detailed phenotypes and gating schemes for all cell

populations mentioned above see Fig. 1 and Supplementary Fig. 1a. To quantify MCMV genome load,

genomic DNA isolated from sorted cell subsets was used as input in a quantitative PCR assay with

primers specific to viral gB gene and to mouse *Pthrp* gene. Absolute quantification was performed

using serial dilutions of a plasmid carrying both gB and *Pthrp* sequences, according to a previously

described method [33], with a quantitation limit of 10 copies of the plasmid standard in a single qPCR

reaction (Supplementary Fig. 1c). Strikingly, PDGFRα^+ FC carried the highest viral genome load

across multiple tissues, in the liver (ca 2-fold), in SG (ca 10-fold) and most substantially, in the spleen

(ca 100-fold) (Fig. 1b-i). Only in the lungs were MCMV genomes absent from PDGFRα^+ FC (Fig. 1j-k).

EC carried MCMV genomes in liver and lungs, which is consistent with previous reports [8, 34], but

their MCMV genome content appeared lower in the spleen and was barely detectable in the SG (Fig.

1b-k). MCMV genomes were detected in VAT Mϕ at levels similar to PDGFRα^+ FC from the same site,

but were absent or barely detectable in Mϕ from other organs (Fig. 1b-k). Next, to compare the viral

genome load in PDGFRα^+ FC to that of circulating myeloid cells, we analysed monocytes isolated from

bone marrow (BM) or the spleen of latently infected mice. We detected less than 10 copies of MCMV

genomes, which is 10-100-fold lower than carried by PDGFRα^+ FC in all organs tested except the

lungs. Importantly, MCMV genomes persisted in PDGFRα^+ FC even 1 year after infection, as shown

for PDGFRα^+ FC from the spleen, lymph nodes (LNs) [35] and VAT (Fig. 1m-o). Together, these

results reveal that MCMV genomes are commonly present in PDGFRα^+ FC of multiple organs during

MCMV latency *in vivo*.

Figure 1. PDGFRα⁺ fibroblastic cells carry MCMV genomes during latent MCMV infection *in vivo.* Analysis of MCMV genome load in indicated cell subsets purified from mice latently infected with MCMV administered intraperitoneally **a-l**, 5 months or **m-o**, 1 year prior. **a**, Experimental set up for b-l. SG, submandibular salivary glands; Li, liver; Spl, spleen; VAT, visceral adipose tissue; Lu, lungs. BM, bone marrow. **m**, Experimental set up for n-o. Lymph nodes (LNs) represent pooled inguinal, brachial, axillary, cervical and mesenteric LNs. **b**, **d**, **f**, **h**, **j**, **n** Gating strategy for sorting of the indicated cell subsets. Numbers are percentage of cells in the indicated

gates. **c**, **e**, **g**, **i**, **k**, **l**, **o** Number of virus genomes per 100,000 cells. Horizontal lines depict median from n = 3-6
biological replicates (depicted as symbols). Biological replicates represent cells sorted from pooled preparations
from **c**, **k**, 1-2 mice; **e**, **i**, **l**, 2 mice; or **g**, **o**, 4 mice and are from 2 independent experiments per organ. Sorting
was done either from 1 organ (VAT) or simultaneously from 2 organs (SG and Lu; Li and BM; SPL and LNs). q.l.,
quantification limit; n.d., not detected; n.a., not analysed

The route of infection determines the load of MCMV genomes in PDGFRα⁺ FC of latently

infected lungs.

Our finding that following intraperitoneal infection MCMV genomes persist in PDGFRα⁺ FC in all

organs except the lungs was intriguing. We previously compared mucosal routes of infection, such as

intranasal or intragastric, with intraperitoneal infection and observed that intranasal infection is the

more efficient route of MCMV infection for the lungs [36]. Nevertheless, intranasal infection results in a

more restricted pattern of virus spread than the intraperitoneal infection route, involving only the lungs

and SG, which was in line with a previous report [37]. Notably, the cellular sites of MCMV latency after

intranasal infection, which is thought to better model the natural MCMV infection [38], are yet to be

investigated. Therefore, we next determined MCMV genome load in PDGFRα⁺ FC, EC as well as

epithelial cells and Mφ from the lungs and SG of mice latently infected via intranasal route (Fig. 2a).

146
147
148 **Figure 2. The route of infection determines the MCMV genome content of PDGFRα⁺ FC in latently**
149 **infected lungs.** Analysis of MCMV genome load in indicated cell subsets purified from mice latently infected
150 with MCMV administered intranasally 5 months prior. **a** Experimental set up. **b**, **c** Number of virus genomes per
151 100,000 cells. Horizontal lines depict median from n = 3-4 biological replicates (depicted as symbols). SG and Lu
152 subsets were simultaneously obtained from 1-2 mice per replicate. q.l., quantification limit; n.d., not detected.
153

Notably, PDGFRα$^+$ FC carried the highest MCMV genome load of all tested subsets in both lungs and

SG of mice infected intranasally (Fig. 2b-c). Thus, distinct routes of infection result in differences in

MCMV genome distribution in PDGFRα$^+$ FC in the lungs. Whether this is due to putative differences in

the exposure/accessibility of PDGFRα$^+$ FC to the virus during acute infection and/or putative

differential effects on virus persistence in PDGFRα$^+$ FC remains to be clarified in future studies.

Differences in MCMV genome load between PDGFRα$^+$ and PDGFRα$^-$ FC during latency *in vivo*.

Next, we asked whether other populations of FC, which do not express PDGFRα, carried similar

MCMV genome load in latently infected mice as PDGFRα$^+$ FC. To this end, we determined MCMV

genome load in distinct subsets of FC from the lungs of mice latently infected either intraperitoneally or

intranasally (Fig. 3a). The CD45$^-$CD31$^-$EpCAM$^-$ cell fraction from the lungs consisted of two discrete

compartments, a major subset of PDGFRα$^+$ FC, which is known to comprise alveolar, peribronchial

and adventitial fibroblasts [27], and a minor FC population that lacked PDGFRα but expressed

PDGFRβ, hereafter termed PDGFRα$^-$PDGFRβ$^+$ FC (Fig. 3b). The majority of PDGFRα$^-$PDGFRβ$^+$ FC

expressed CD146 (Supplementary Fig. 1d), fitting with the universal phenotype of mural cells [39].

Following intraperitoneal infection, which results in the preferential distribution of MCMV genomes to

EC, both PDGFRα$^+$ FC and PDGFRα$^-$PDGFRβ$^+$ FC were a minor source of MCMV genomes with the

load of less than 10 viral genome copies per 100,000 cells (Fig. 3c). Following intranasal infection,

which results in the preferential distribution of MCMV genomes to PDGFRα$^+$ FC, viral genomes were

robustly detected in both FC subsets but the genome load in PDGFRα$^-$PDGFRβ$^+$ FC was ca 3-fold

lower compared to PDGFRα$^+$ FC (Fig. 3d). We also mapped the distribution of MCMV genomes

during latency in the subpopulations of FC from the spleens of mice infected intraperitoneally (Fig. 3e).

We fractionated CD45$^-$CD31$^-$ITGB1$^+$ splenic FC [29] into Ly6C$^+$ red pulp FC, BST-1$^+$ white pulp FC,

and BST-1$^-$Ly6C$^-$ FC (hereafter termed, other FC), which are mainly composed of CD146$^+$ mural cells

[29]. As shown in Fig. 3f, Ly6C$^+$ red pulp FC uniformly expressed high levels of PDGFRα, BST-1$^+$

white pulp FC contained a minor fraction of PDGFRα$^-$ cells while other FC consisted mainly of

PDGFRα$^-$ cells. Notably, MCMV genome load was nearly 10-fold higher in Ly6C$^+$ red pulp FC

compared to the other two FC subpopulations (Fig. 3g). In sum, MCMV genome load is higher in the

PDGFRα-positive fraction of FC in the lungs and spleen than in other FC subsets.

Figure 3. Differences in MCMV genome load between PDGFRα⁺ and PDGFRα⁻ FC during latency *in vivo*.
Analysis of MCMV genome load in indicated FC subsets purified from the lungs of mice latently infected with
MCMV administered intraperitoneally or intranasally (**a-d**), or the spleen of mice latently infected with MCMV
administered intraperitoneally (**e-g**). **a** Experimental set up for b-d. **e** Experimental set up for f-g. **b, f** Gating
strategy for sorting of the indicated cell subsets. **c**, **d**, **g** Number of virus genomes per 100,000 cells. **c**, **d**
Horizontal line depicts the arithmetic mean ± SD from n = 4 biological replicates (depicted as symbols) pooled
from 2 independent experiments. Biological replicates represent cells sorted from pooled preparations from 2
mice. ***, p-value < 0.001, one-way ANOVA with Dunnett's multiple comparisons test **g** Horizontal line depicts
the arithmetic mean ± SD from n = 2-4 biological replicates (depicted as symbols) pooled from 2 independent
experiments. Biological replicates represent cells sorted from pooled preparations from 4 mice. ****, p-value <
0.0001, unpaired t test. q.l., quantification limit; n.d., not detected.
194

Silenced and stochastic viral gene transcription in PDGFRα⁺ FC from latently infected mice

Next, we investigated if PDGFRα⁺ FC that carried viral genomes during MCMV latency *in vivo* fulfilled

the formal criteria of latent infection. We first asked whether viral gene expression in PDGFRα⁺ FC

from latently infected mice was consistent with a putative ongoing lytic viral cycle. To this aim, we

measured the expression of viral transcripts from the immediate-early (*ie1*/*ie3*), early (*M38*) or late

(*M48/SCP*) phase of the lytic infection in RNA isolated from PDGFRα⁺ FC from latently infected mice

by RT-qPCR (Table 1). We analysed PDGFRα⁺ FC from various organs and conditions in which they

were shown to carry MCMV genomes (see Fig. 1 and 2), such as the spleen, VAT, SG and liver of

mice infected intraperitoneally, and from PDGFRα⁺ FC from the lungs of mice infected intranasally.

PDGFRα⁺ FC from the lungs of mice infected intraperitoneally, which are devoid of MCMV genomes

(see Fig. 2), were used as a negative control. Liver EC and VAT Mϕ from mice infected via

intraperitoneal route were also included. Following reverse transcription, copies of cDNA representing

each transcript were quantified using serial dilutions of MCMV-BAC [33], with a quantitation limit of 10

copies of the BAC standard in a single qPCR reaction (Supplementary Fig. 2a). Although we used

primer pairs that did not distinguish between viral DNA and RNA, our assay specifically detected viral

RNA, since no positive signals were obtained when the same RNA samples were processed with

omission of reverse-transcriptase (negative data not shown). To ascertain that comparable amounts of

cDNA were used, we standardized our readouts on mouse *Gapdh* expression in all samples. The

number of cells corresponding to the cDNA input used per qPCR reaction was estimated for each

sample and was in the range of 5,000-15,000 cells (Table 1). Overall, expression of viral mRNAs was

detected only sporadically and as no more than a few hundred of copies (Table 1). Specifically, *ie1/ie3*

expression was detected in 8, *M38* in 2 and *M48/SCP* in 1 out of 18 tested PDGFRα⁺ FC samples, but

none of the samples expressed all three viral genes (Table 1). This transcriptional profile is

inconsistent with lytic infection while it fits with the pattern of viral transcription in latently infected lungs

described recently by Griessl *et al.*, which was found to be intermittent and stochastic [34]. *ie1/ie3*

expression was detected in 1 out 3 tested liver EC samples and in 1 out 3 tested VAT Mϕ samples,

indicating that viral transcription is also silenced in EC and Mϕ carrying MCMV genomes during

latency *in vivo.* In sum, the profile of viral gene expression in PDGFRα⁺ FC was at odds with

persistent lytic infection.

224
225
226
227

228 **Table 1**

		Est. cell no.	MCMV transcript copies detected per reaction					
			ie1/3		M38		M48(SCP)	
5 months after intraperitoneal infection	Spleen PDGFRα⁺ FC	8,911	0	0	0	0	0	0
		9,153	0	0	0	0	0	0
		7,630	0	0	0	0	0	0
	AT PDGFRα⁺ FC	8,359	0	0	0	0	0	0
		7,659	21	8	0	0	0	0
		11,799	0	0	0	0	0	0
	SG PDGFRα⁺ FC	5,468	0	0	0	0	0	0
		6,242	123	140	0	0	0	0
		6,242	0	0	0	9	58	60
	Liver PDGFRα⁺ FC	6,224	0	0	0	0	0	0
		6,735	0	0	0	0	0	0
		6,715	0	0	0	0	0	0
	Liver EC	9,333	0	0	0	0	0	0
		6,618	0	0	0	0	0	0
		9,488	8	7	0	0	0	0
	Lung PDGFRα⁺ FC (negative control)	14,431	0	0	0	0	0	0
		15,234	0	0	0	0	0	0
		16,971	0	0	0	0	0	0
5 months after intranasal infection	Lung PDGFRα⁺ FC	16,971	200	128	0	0	0	0
		14,756	405	303	7	16	0	0
		17,261	317	433	0	0	0	0
3 months after intraperitoneal infection	VAT PDGFRα⁺ FC	10,626	10	14	0	0	0	0
		11,751	11	29	0	0	0	0
		9,498	40	64	0	0	0	0
	VAT Mφ	10,551	0	0	0	0	0	0
		12,228	0	0	0	0	0	0
		10,975	5	5	0	0	0	0

229
230
231 **Table 1.** RT-qPCR analysis of the expression of MCMV genes in the indicated subsets obtained from mice
232 latently infected with MCMV via the indicated route and for the indicated amount of time. Each row represents
233 cDNA from a different biological replicate analysed in 2 reactions (technical duplicates) per MCMV gene.
234 Biological replicates represent cells sorted from pooled preparations from 1-2 (Lu, Li, SG, VAT) or 4 (Spl) mice
235 per replicate.
236

PDGFRα⁺ FC carry latent MCMV *in vivo*.

To test for functional criteria of latency, we next determined if MCMV can reactivate from PDGFRα⁺

FC purified from latently infected mice. To this aim, we devised an *ex vivo* reactivation assay, in which

PDGFRα⁺ FC isolated from the lungs of mice latently infected via intranasal route were seeded in half-

area 96-well plates at the density of 100,000 cells per well (resulting in the deposition of ~100 MCMV

genomes per well) (Fig. 4a). Upon 7 days of *ex vivo* incubation, viral reactivation was assessed in

individual wells using three criteria: i) visual inspection of plaque formation, ii) IE1

immunofluorescence as well as iii) detection of infectious virus in culture supernatant by plaque assay

on monolayer of mouse embryonic fibroblasts (MEFs). Notably, infectious virus emerged in 30-50 % of

wells from 4 out of 4 samples of lung PDGFRα⁺ FC (Fig. 4b-c). As expected, virus did not grow out

when lung PDGFRα^+ FC isolated from mice infected intraperitoneally were used, as they were shown

to be devoid of MCMV genomes (Fig. 4d). Further importantly, no virus outgrowth was detected when

lysates prepared from lung PDGFRα^+ FC isolated from mice infected intranasally were incubated on a

monolayer of MEFs for the same amount of time (Fig. 4e-f). The results showing that virus grew out

only from cells but not from cell lysates (even though the assay was able to detect a single PFU of

MCMV), argues that PDGFRα^+ FC carried latent MCMV which reactivated upon *ex vivo* cultivation. To

confirm this using PDGFRα^+ FC from another tissue, but also to study the kinetic of reactivation

events, virus reactivation was also monitored in PDGFRα^+ FC isolated from the VAT of mice latently

infected via intraperitoneal route. Mice were administered with reporter MCMV expressing major IE

promoter-driven GFP (MCMV^{IE-GFP}) [29], which resulted in the same latent viral load in VAT PDGFRα^+

FC (Supplementary Fig. 2b) while it enabled monitoring of the kinetic of re-initiation of lytic gene

transcription by life cell imaging (Fig. 4g). By 7 days of *in vitro* incubation, the virus grew out in ~10%

of wells from 9 out of 10 samples (Fig. 4h-i). Virus outgrowth was specifically associated with the

presence of PDGFRα^+ FC, since cultivation of VAT cell preparations that had been depleted of

PDGFRα^+ FC did not yield any virus (Fig. 4j). It should be noted that this result does not provide

evidence for the lack of reactivation from VAT Mϕ or EC since the culture conditions did not support

their survival (data not shown). As shown for lung PDGFRα^+ FC, virus was recovered only from live

PDGFRα^+ FC from VAT and not from cell lysates (Fig. 4k-l). Thus, the virus could reactivate from

PDGFRα^+ FC from both lungs and VAT, with frequency correlating with the level of IE gene

transcription in these cells, which was lower in VAT than in the lungs (see Table 1). Based on the

emergence of initial GFP$^+$ cells, re-initiation of lytic gene transcription occurred either 1.5 - 2 days after

seeding (50% of reactivations) or no sooner than 4 - 5 days after seeding (40% of reactivations) (Fig.

4i). Thus, reactivation events appeared to follow a biphasic kinetic. Together, these results indicate

that MCMV latently persists in PDGFRα^+ FC *in vivo*.

Figure 4. MCMV establishes latent infection in PDGFRα⁺ FC *in vivo*. *Ex vivo* virus reactivation assay from
PDGFRα⁺ FC purified from **a-f**, the lungs of mice latently infected with MCMV administered intranasally or **g-m**,
from the VAT of mice latently infected with MCMV administered intraperitoneally. **a**, Experimental set up for b-d.
e, Experimental set up for f. **g**, Experimental set up for h-j, m. **k**, Experimental set up for l. **b**, **d**, **f**, **i**, **j**, **l**,
Frequency of viral outgrowth from wells with indicated cells. Biological replicates are from 1-2 mice per replicate.
Numbers on the right side are the proportion of wells with virus outgrowth for the indicated biological replicate. **c**
Representative immunofluorescence image showing detection of viral IE1 protein. **h** Representative
fluorescence microscopy image of a GFP⁺ plaque that arose in the *ex vivo* reactivation assay. **m** Kinetic of re-
initiation of viral lytic cycle. Shown is the proportion of wells in which GFP signal had first emerged at the
indicated time-point.

PDGFRα⁺ FC are a site of lytic virus replication during primary MCMV infection

The apparent latent persistence of MCMV in PDGFRα⁺ FC *in vivo* implied the question whether or not

they could also support productive virus cycle during primary MCMV infection. To directly assess the

permissiveness of PDGFRα⁺ FC for lytic MCMV infection, we purified PDGFRα⁺ FC from the lungs of

naive mice and infected them *ex vivo* with recombinant MCMV[tFP-GRB], which expresses major IE

promoter-driven green fluorescent protein (GFP) and late gene promoter-driven red fluorescent protein

(mCherry) (Fig. 5a). Analysis of viral replication kinetics during *in vitro* culture demonstrated that

PDGFRα⁺ FC are permissive for lytic MCMV infection (Fig. 5b-c). To address this *in vivo*, we first

investigated if lytic gene expression was induced in PDGFRα⁺ FC during the acute phase of infection

with MCMV. Given that spleen becomes infected with MCMV within hours following intraperitoneal

inoculation [22], we isolated PDGFRα⁺ FC from the spleens of mice 12 hours post intraperitoneal

infection, and analysed the expression of viral genes *ie1/3*, *M38* and *M48(SCP)* by RT-qPCR. At 12

hours post infection, splenic PDGFRα⁺ FC expressed high levels of all three viral transcripts at levels

at least 1,000-fold higher than in any tested subset of PDGFRα⁺ FC from latently infected mice (Table

2, see also Table 1). Therefore, splenic PDGFRα⁺ FC supported lytic viral transcription shortly after

the inoculation of mice with MCMV.

Table 2

			MCMV transcript copies detected per reaction					
		Est. cell no.	*ie1/3*		*M38*		*M48(SCP)*	
12 hours after intraperitoneal infection	**Spleen PDGFRα⁺ FC**	7,944	219,224	211,938	411,485	411,485	61,694	61,289
uninfected	**Spleen PDGFRα⁺ FC**	6,113	0	0	0	0	0	0

300
301   **Table 2.** RT-qPCR analysis of the expression of MCMV genes in PDGFRα⁺ FC at 12 hours post infection. cDNA
302   isolated from pooled cell preparations from 4 mice was analysed in 2 technical duplicates per MCMV gene.
303
304
To confirm that PDGFRα⁺ FC also supported productive replication of the virus *in vivo*, we used mice

heterozygous for *Pdgfra*-CreER[T2] knock-in allele [40] (hereafter termed *Pdgfra*-CreER[T2] (+/-), that

express tamoxifen-activatable Cre recombinase from one endogenous *Pdgfra* locus while the other

locus is preserved, and infected them with a reporter virus MCMV[floxSTOP-GFP] [41]. MCMV[floxSTOP-GFP]

undergoes deletion of the stop cassette preventing GFP expression in cells that express active Cre

recombinase and was used previously with Tie2-Cre or Alb-Cre mice to define virus replication in

endothelial cells or in hepatocytes [11]. Following the infection of *Pdgfra*-CreER[T2] (+/-) mice with

MCMV[floxSTOP-GFP] it is expected that the viruses that replicate in PDGFRα⁺ FC become genetically

recombined so their progeny gives rise to GFP⁺ plaques on monolayers of MEFs. *Pdgfra*-CreER^{T2} (+/-)

mice were infected with MCMV$^{floxSTOP-GFP}$ via the intraperitoneal route on day 0, fed tamoxifen from -1

to 3 days post infection to activate the Cre recombinase, and sacrificed on day 4 post infection (Fig.

5d). Organ lysates from the spleen, VAT, liver and lungs were used to determine the frequency of

GFP⁺ plaques. On day 4 post infection, GFP⁺ virus was detected as ~10% of plaques in the liver, ~20

% in the VAT and ~50 % in the spleen and lungs, although in the latter organ the percentage of GFP⁺

plaques could only be reliably measured in 5 out of 10 samples due to generally low infectious virus

titer in this organ (Fig. 5e-f). The emergence of GFP⁺ virus was entirely dependent on the presence of

Cre recombinase since similar virus titer but no GFP⁺ plaques were found in organs of infected Cre-

negative littermate controls (data not shown).

Figure 5

Figure 5. PDGFRα⁺ FC are a site of lytic virus replication during primary MCMV infection. a-c, Analysis of
virus replication kinetics in lung-derived PDGFRα⁺ FC following their *ex vivo* infection of with MCMV$^{tFP-GRB}$, which
expresses IE promoter driven-green fluorescent protein (GFP) and late gene promoter-driven red fluorescent
protein (mCherry). **a** Experimental set up for b-c. **b** Data are presented as arithmetic mean ± SD of n = 3
technical replicates and are from one representative experiment of 2 using cells sorted from pooled lung
preparations from 3 mice. **c** Representative fluorescence microscopy images of GFP and mCherry expression 5
days after *ex vivo* infection. **d-f**, Contribution of PDGFRα⁺ FC to MCMV propagation *in vivo* determined after

intraperitoneal infection of *Pdgfra*-CreER[T2] (+/-) mice with a Cre-inducible reporter virus, MCMV[floxSTOP-GFP]. **d**

Experimental set up for e-f. **e** Median virus titer in indicated organs. Circles represent samples from individual

mice. Grey-filled circles are samples for which percentage of GFP[+] plaques could be determined. White-filled

circles are samples for which percentage of GFP[+] plaques could not be determined. **f** Median proportion of GFP[+]

plaques in indicated organs. Circles represent samples from individual mice.

Thus, multiple organs contained the progeny of viruses that had replicated in PDGFRα[+] cells,

indicating that PDGFRα[+] FC play a role in MCMV *in vivo* replication. Of note, given that MCMV may

have spread from the sites of initial replication within 4 days, it is not possible to distinguish whether

the virus had replicated in PDGFRα[+] FC present in the given organ or elsewhere. In sum, PDGFRα[+]

FC supported both productive infection and latent MCMV persistence *in vivo*.

Stat1* is required for MCMV genome persistence in fibroblastic cells *in vivo

Next, we sought to identify host pathway/-s that affect MCMV's ability to latently persist in PDGFRα[+]

FC *in vivo*. We hypothesized that the same signal/-s that inhibit lytic cycle entry by MCMV may

concomitantly enable the virus to establish and/or maintain latent infection. Since IFNβ was sufficient

to silence IE transcription [20], we chose to assess the putative involvement of STAT1, which crucially

mediates responses to IFNs [16] by testing lytic and latent MCMV infection in *Stat1*[-/-] mice. To

circumvent the caveats linked to the increased sensitivity of *Stat1*[-/-] mice to MCMV, we generated and

used a single-cycle GFP-reporter MCMV mutant, which we name MCMV[ΔgLGFP]. To this effect we

disrupted the gene for the essential virion glycoprotein L (gL/M115) in our GFP reporter virus,

MCMV[IE1/3-GFP] [29] on a Δm157 background. MCMVs lacking gL can be propagated in gL-

complementing cells, to generate virus stocks consisting of pseudotyped gL[+] virions that are able to

infect and replicate for one cycle in gL deficient cells, whereupon their progeny virions will lack the gL

and thus cannot spread further to other cells [42-44]. As shown in Fig. 6a, WT and *Stat1*[-/-] mice were

infected with MCMV[ΔgLGFP], followed by analysis of lytic infection and latent persistence of the virus

respectively at 22 hours and 60 days post infection in fibroblastic cells in the spleen, which are known

to be directly infected by MCMV virions injected intraperitoneally or intravenously [22]. We focused

specifically on red pulp FC, firstly because their high MCMV genome content during *in vivo* latency

offered a broad dynamic range and secondly, to eliminate confounding effects due to a potential

change in the composition of spleen stromal cells in *Stat1*⁻ᐟ⁻ mice. Since Ly6C is an IFN-regulated

gene, Ly6C⁺ red pulp FC were identified as PDGFRα⁺BST-1⁻CD146⁻ FC, as previously reported [29].

The abundance of red pulp FC was comparable between WT and *Stat1*⁻ᐟ⁻ mice (Supplementary Fig.

2c). At 22 hours after intraperitoneal or intravenous injection of MCMV^ΔgLGFP, *Stat1*⁻ᐟ⁻ mice showed a

significantly higher percentage of GFP⁺ red pulp FC (Fig. 6b-d). Consistent with being lytically

infected, GFP⁺ red pulp FC from WT and *Stat1*⁻ᐟ⁻ mice expressed high levels of *ie1/ie3*, *M38* and

M48(SCP) transcripts (Table 3). Therefore, more red pulp FC became lytically infected in the absence

of STAT1. At 60 days post infection, red pulp FC were devoid of GFP⁺ cells (Fig. 6f) and viral gene

expression was silenced (Table 3). Strikingly, whereas MCMV genomes persisted in red pulp FC from

WT mice at ~ 300 copies per 100,000 cells, these cells were profoundly depleted of MCMV genomes

and contained only ~ 10 copies per 100,000 cells in *Stat1*⁻ᐟ⁻ mice (Fig. 6g). Together, the above data

implicate a dual role of STAT1 as a negative regulator of lytic MCMV infection and a positive regulator

of latent MCMV persistence in fibroblastic cells *in vivo*.

Figure 6. *Stat1* is required for MCMV genome persistence in fibroblastic cells *in vivo*. a-d, Analysis of the
lytic infection of red pulp FC in WT and *Stat1*⁻ᐟ⁻ mice 22 hours after the administration of MCMV^ΔgL-GFP
intraperitoneally or intravenously. **a** Experimental set up for b-d. **b-d** Percentage of GFP⁺ cells in red pulp FC. **b**
Representative flow cytometry plot. Numbers indicate the percentage of gated cells. **c, d** Data are presented as
the arithmetic mean ± SD of n = 3 biological replicates (individual mice) and are from 1 experiment performed for
each infection route. *, p-value < 0.05 (exact p value indicated in the plot), unpaired t test. **e-g**, Analysis of latent
MCMV persistence in red pulp FC from WT and *Stat1*⁻ᐟ⁻ mice 60 days after intraperitoneal administration of
MCMV^ΔgL-GFP. **e** Experimental set up for f-g. **f** Plots show the lack of GFP⁺ cells in red pulp FC. **g** Number of virus
genomes per 100,000 of red pulp FC. Horizontal line depicts arithmetic mean ± SD from n = 4 biological
replicates (depicted as symbols) and are pooled from 2 independent experiments. Biological replicates represent
cells sorted from pooled preparations from 2 mice. **, p-value < 0.01, Welch's t test. q.l., quantification limit; n.d.,
not detected.

386
387
388 **Table 3**

| | | Est. cell no. | MCMV transcript copies detected per reaction | | | | | |
			ie1/3		M38		M48(SCP)	
22 hours after intravenous infection	GFP⁺ WT red pulp FC	310	72,214	73,673	677,316	713,744	65,231	60,544
		470	151,387	139,745	1,253,481	1,278,349	100,668	114,507
		201	35,075	31,316	369,285	355,057	42,951	42,372
	GFP⁺ Stat1⁻ᐟ⁻ red pulp FC	242	82,357	79,127	1,208,111	1,281,450	102,291	98,214
		311	41,163	36,264	710,806	724,908	44,131	44,734
		518	59,401	65,214	905,681	999,157	66,734	65,391
60 days after intraperitoneal infection	WT red pulp FC	21,864	ND	ND	ND	4	ND	ND
		18,811	ND	ND	ND	ND	ND	ND
		17,837	ND	ND	ND	ND	ND	ND
		16,063	ND	ND	ND	ND	ND	ND
		18,503	ND	ND	ND	ND	15	ND
	Stat1⁻ᐟ⁻ red pulp FC	23,573	ND	ND	ND	ND	ND	ND
		21,565	ND	ND	ND	ND	ND	ND
		17,029	ND	ND	ND	ND	ND	ND
		25,591	ND	ND	ND	ND	ND	ND

389
390 **Table 3.** RT-qPCR analysis of the expression of MCMV genes in the indicated subsets purified from WT or
391 *Stat1*⁻ᐟ⁻ mice that had been infected with MCMV$^{\Delta gL\text{-}GFP}$ for 22 hours or 60 days. Each row represents cDNA from
392 a different biological replicate analysed in 2 reactions (technical duplicates) per MCMV gene. Each biological
393 replicate represents cells sorted from a single mouse in a single experiment (22 hours post infection) or cells
394 sorted from pooled preparations of 2 mice and are from 2 independent experiments (60 days post infection).
395

Discussion

This work provides to our knowledge the first comprehensive characterisation of the

distribution of MCMV genomes in multiple tissue-resident cell types across numerous organs during

MCMV latency *in vivo*, and the first demonstration of latent MCMV persistence in fibroblastic cells.

Latent MCMV persistence in PDGFRα⁺ FC was not restricted to a single organ but represented a

wide-spread phenomenon engaging many organs, suggesting that fibroblasts are a physiologically

relevant, and in some organs may be a major site of MCMV *in vivo* latency. Our findings are not in

opposition with previous identification of endothelial cells and macrophages as cellular sites of MCMV

latency *in vivo* but rather confirm and extend previous knowledge by broadening the scope of tissues

in which these cell types were tested for MCMV genome presence. In our model, which involved

infection of adult C57BL/6 mice with tissue culture derived MCMV, we found the content of MCMV

genomes in monocytes to be below 10 copies per 100,000 cells at 5 months post infection. This is

similar to a recent study by Liu *et al.*, in which 3 week-old BalbC mice were infected with MCMV Smith

strain and analysed 3-6 months later, where the MCMV genome load in monocytes was more robust

but still not higher than 10 copies per 100,000 cells [7]. It has been suggested that the presence of

latent HCMV in endothelial cells may be a consequence of transdifferentiating latently infected myeloid

progenitor cells [45], which might in theory explain our observations. However, the scarcity of viral

genomes in monocytes in relation to tissue-resident cells, argues against this explanation in MCMV

latency. Furthermore, given modern knowledge about the origin of fibroblastic cells in the mouse we

deem this scenario very unlikely for MCMV. It is currently well established that $CD45^-PDGFR\alpha^+$ or

$PDGFR\beta^+$ fibroblastic cells found in mouse organs under homeostatic conditions are mesenchymal-

derived and not a progeny of circulating progenitors originating from hematopoietic or mesenchymal

BM-derived stem cells [27]. Although differentiation of monocyte/Mϕ-like cells into fibroblasts has been

documented in the context of organ injury or fibrosis [46, 47], the low MCMV content of monocytes

and Mϕ in most studied organs is at odds with the high enrichment of $CD45^-PDGFR\alpha^+$ for MCMV

genomes observed in our study. An alternative hematopoietic source of fibroblast-like cells in a

perturbed state can be $CD34^+Col1a1^+$ fibrocytes, but they and their progeny were excluded from our

analysis based on their positivity for CD45 [48]. Thus, available indirect evidence strongly suggests

that MCMV not only latently persist in but also latently infects $CD45^-PDGFR\alpha^+$ FC.

Notably, $CD45^-PDGFR\alpha^+$ FC also became lytically infected by the virus and contributed to its *in*

vivo replication during the acute stage following primary infection. These findings shed new light on

the role of fibroblasts in MCMV's life cycle *in vivo* by showing that not only does this cell type support

viral lytic cycle, as previously understood, but it nevertheless serves as the site of latent MCMV

persistence. It is important to note that the above conclusion can only be inferred for the entire

population of $PDGFR\alpha^+$ FC from a given organ, but not on a single-cell basis. $PDGFR\alpha^+$ FC are known

to be heterogenous [32] and it is entirely possible that lytic and latent infection are allocated to distinct

subsets or maturation stages. Additional studies involving single cell transcriptomics will be required to

resolve this matter in the future.

While our study was focused on demonstrating bone-fide latent virus persistence in fibroblastic

cells and identification of a cellular signalling pathway on which it is critically dependent, some

important follow up questions remain that lie outside of the scope of the current project. For instance, although our mechanistic studies in Stat1$^{-/-}$ mice unravelled that endogenous Stat1 was required to enable latent persistence of MCMV in fibroblastic cells, we did not determine whether or not the requirement for Stat1 was cell intrinsic. To decipher this, mice bearing conditional deletion of Stat1 in fibroblasts will need to be implemented. Should Stat1 be specifically required in fibroblasts, it will further need to be clarified whether or not Stat1 affects the ability of the virus to establish latent infection or rather persist in a latent form, for example by preventing spontaneous reactivation. Addressing these questions will likely require careful probing of single-cell transcriptomes of fibroblastic cells at various time points after infection in vitro or in vivo. Regardless of the results of the future investigations addressing these outstanding questions, it is important to observe that the data generated in the course of this study unravel that one of the most prominent antiviral host factors, which is crucial for the resolution of virus infections and host survival, concomitantly enables a herpesvirus to latently persist in its host, at least in certain cell types. In sum, the findings of this study significantly broaden our understanding of the functions of murine Stat1 during in vivo infection with a β-herpesvirus.

Methods

Mice

Experiments were performed with female mice on C57BL/6 background. Mice were infected at 8-10 weeks of age, except in Fig. 6 where the infection was done at 14 weeks of age. C57BL/6JrJ mice (purchased from Janvier Labs), Pdgfra-CreER^{T2} mice (IMSR_JAX:032770) [40] and Stat1$^{-/-}$ mice (generated by mating Stat1^{flox} mice with CMV-Cre deleter strain) [49] were bred and maintained under specific pathogen free (SPF) conditions according to Federation of European Laboratory Animal Science Associations (FELASA) guidelines respectively at central animal facility of HZI Braunschweig, Germany; Faculty of Medicine, University of Rijeka, Croatia, and University of Veterinary Medicine Vienna, Austria. Animal procedures were approved as due by The Lower Saxony State Office of Consumer Protection and Food Safety; by The Ethics Committee at the Faculty of Medicine, Rijeka and Ethics Committee of the Veterinary Department of the Ministry of Agriculture, Croatia, and by the

Ethics and Animal Welfare Committee of the University of Veterinary Medicine Vienna and the

Austrian Federal Ministry of Science and Research. Virus was administered in 200 μl PBS

intraperitoneally or in 20 μl in PBS intranasally as described previously [50]. *Pdgfra*-CreER^{T2} mice

received tamoxifen (0.1mg/g; Sigma) diluted in corn oil by oral gavage daily from day -1 to day 3 post

infection.

Viruses

Viral stocks were generated in tissue culture by transfecting MCMV genome-carrying BACs into

NIH3T3 cells (ATCC CRL-1658) using FuGene HD (Promega). Reconstituted viral particles were

passaged five times on M2-10B4 cells (ATCC CRL-1972) before generating sucrose cushion-purified

viral stocks. All MCMVs used in this study were based on the Mck2-repaired BAC-encoded MCMV

strain, clone pSM3fr-MCK-2fl 3.3 [51]. MCMV^{IE-GFP} [29] and MCMV$^{floxSTOP-GFP}$ [41] were described

before. MCMV$^{tFP-GRB}$ expressing three different fluorescent proteins under the control of endogenous

viral promoters was generated by *en passant* mutagenesis [52] using expression cassettes described

earlier [53]. In short, constructs encoding fluorescent reporters (EGFP, mTagBFP-NLS and mCherry)

plus a distal P2A-encoding sequence were inserted into the start codon of IE1/3 (EGFP), E2

(mTagBFP-NLS), and SCP (mCherry). MCMV^{ΔgLGFP} was generated by disrupting m157 and gL/M115

genes in MCMV^{IE-GFP}. Viral genes were disrupted by frameshift mutations induced by replacement of

38 bp of each gene's open reading frame (ORF) with two STOP codons targeted to the ORF's start

(gL) or after codon 22 (m157). MCMV^{ΔgLGFP} was propagated in gL-expressing NIH-3T3 cell line [42] (a

kind gift from Ian Humphreys, Cardiff University).

Cell Isolation

Stromal-vascular fractions from VAT were prepared as previously described [35] except that liberase

TM in the digestion medium was replaced with collagenase P (0.4 mg/ml). Spleen, LNs and SGs were

processed according to a previously established protocol [29]. In short, enzymatic treatment with

collagenase P (0.4 mg/ml), dispase II (2 mg/ml) and DNase I (50 μg/ml) was performed for 30 min at

37 °C and then repeated for additional 20 min, followed by incubation with 5 mM EDTA for 5 min.

CD45$^-$ cell fractions from the spleen and LNs were enriched by depletion of CD45$^+$ cells while splenic

Mϕ were enriched by positive selection of VCAM-1$^+$ cells using MACS (Miltenyi Biotech). Lungs were

injected with 2.5 ml digestion solution containing collagenase P (0.4 mg/ml), dispase II (2 mg/ml) and

DNase I (50 µg/ml), minced with scissors and then processed in gentleMACS Dissociator (Miltenyi

Biotech) using program "37C_m_LDK_1". Liver digestion was performed using the retrograde

perfusion technique via the inferior vena cava described by Mederacke *et al.* [30] with modifications.

Briefly, organs were *in situ* perfused with 10 ml of pre-warmed Liver Perfusion Medium (Gibco)

followed by injection of 20 ml of pre-warmed digestion medium (1 mg/ml collagenase P, 1 mg/ml

dispase II, 50 µg/ml DNase I in Gibco Hank's Balanced Salt Solution (HBSS) with 10 mM HEPES and

1.5 mM calcium chloride) over 5-7 min. Afterwards, organs were dissected, minced with scissors and

in vitro digested for 45 min at 37°C. Non-parenchymal cells were enriched by two rounds of

centrifugation at 50 × g for 5 min whereupon only supernatant was collected.

Flow Cytometry and Cell Sorting

Antibodies used in this study are listed in Supplementary Table 1. Dead cells (identified using 7-AAD

or Zombie Fixable Viability Kits; both from BioLegend) and cell aggregates (identified on FSC-A

versus FSC-H scatter plots) were excluded from all analyses. Sorting was performed on ARIA-Fusion,

Aria-II SORP or Aria-II (BD Biosciences). Analysis was done using FlowJo software (version 10.8.1,

BD Biosciences).

Immunofluorescence

Cells were fixed in 4 % (wt/vol) paraformaldehyde, permeabilized in 0.1% Triton-X and blocked with

2% BSA. Staining with anti-IE1 antibody (IE 1.01, CapRi) was done overnight at 4°C and was followed

by incubation with anti-mouse IgG (H+L) F(ab')$_2$ conjugated to Alexa Fluor 647 (Cell Signaling

Technology). Images were acquired with ZEISS LSM 980 laser-scanning confocal microscope and

analysed using ZEN software (Carl Zeiss MicroImaging).

Quantification of MCMV Genome Load by qPCR

Quantification of MCMV genome load was performed according to the method described by Simon *et*

al. [33] with modifications. Briefly, DNA was extracted using AllPrep DNA/RNA Micro Kit (QIAGEN),

and after elution precipitated with ethanol, washed, and resuspended in 40 µl dH$_2$O. Samples were

assayed for viral gB and mouse Pthrp DNA copies in technical duplicates using 9 µl of sample DNA

per reaction. Serial dilutions of 10^6-10^1 copies per reaction of the pDrive_gB_PTHrP_Tdy plasmid [33]

were used to generate standard curves for both qPCR reactions. Dilutions of 10^4-10^1 plasmid copies

contained 1 ng carrier RNA (QIAGEN). QPCR was performed in a LightCycler 480 Instrument II

(Roche) using Fast Plus EvaGreen qPCR master mix (Biotium). Cycling conditions were as follows:

enzyme activation, 2 min 95°C, followed by 50 cycles of denaturation for 10 sec at 95°C, annealing for

20 sec at 56°C and extension for 30 sec at 72°C. Specificity of amplified sequences was validated by

inspection of melting curves. Primer sequences: *Pthrp* 5'-ggtatctgccctcatcgtctg-3' and 5'-

cgtttcttcctccaccatctg-3'; gB 5'-gcagtctagtcgctttctgc-3' and 5'-aaggcgtggactagcgataa-3'.

Analysis of MCMV Transcripts by RT-qPCR

Total RNA was extracted using AllPrep DNA/RNA Micro Kit with on-column DNA removal (QIAGEN).

cDNA was synthesized with SuperScript IV Reverse Transcriptase and a 1:1 mixture of oligo-dT and

random oligonucleotide hexamers (Invitrogen). QPCR was performed in a LightCycler 480 Instrument

II (Roche) using Fast Plus EvaGreen qPCR master mix (Biotium). Serial dilutions of 10^6-10^1 copies per

reaction of MCMV genome containing BAC (clone pSM3fr-MCK-2fl 3.3 [51]) were used to generate

standard curves. Cycling conditions were as follows: enzyme activation, 2 min 95°C, followed by 50

cycles of denaturation for 10 sec at 95°C, annealing for 20 sec at 57°C and extension for 30 sec at

72°C. Specificity of amplified sequences was validated by inspection of melting curves. RT-qPCR

reactions prepared from equal amount of sample RNA but with the omission of reverse transcriptase

served as controls for the absence of carry-over viral DNA. Primer sequences: *Gapdh* 5'-

tgtgtccgtcgtggatctga-3' and 5'-cctgcttcaccaccttcttgat-3'; ie1/3 5'-gggcatgaagtgtgggtaca-3' and 5'-

cgcatcgaaagacaacgcaa-3'; M38 5'-cccatgtccacgtttggtg-3' and 5'-tctcaagtgcaatcgcctca-3'; M48/SCP

5'-aggacgtgttgaacttcttctcc-3' and 5'-tcgatgtccgtctccttcac-3'.

***Ex vivo* Reactivation Assay**

FC were seeded into fibronectin-coated 96 Half Area Well Flat Bottom Microplates (Corning) (lung FC)

or gelatin- or fibronectin-coated 96 Well Flat Bottom Microplates (Greiner Bio-One) (VAT FC) at a

density of 100,000 cells per well in culture medium (high-glucose DMEM supplemented with 1 %

(vol/vol) GlutaMAX, 10 mM HEPES, 50 μM 2-mercaptoethanol, 200 U/ml penicillin, 200 U/ml

streptomycin, 50 μg/ml gentamicin and 5 % FCS) and incubated in a humidified incubator under 5 %

CO_2 and at 37°C for 7 days without replacing the medium. Cell lysates prepared in gentleMACS M

Tubes for tissue homogenisation using gentleMACS Dissociator (Miltenyi Biotech, program

"mlung_02") were assayed in parallel by incubation with MEFs. To track the emergence of GFP$^+$ cells,

plates were imaged with IncuCyte S3 automated fluorescence microscope (Sartorius). After 7 days,

plates were directly inspected for the presence of plaques and additionally, the culture supernatant

from each well was tested for the presence of infectious virus by plaque assay on MEFs. To ensure

detection of even minimal amounts of virus [54], following addition of the supernatant, MEFs were

spun down at 800 g for 30 min and then incubated at 37°C for another 30 min. Afterwards, the

inoculum was removed and the cells overlaid with medium containing 0.75 % (wt/vol) methylcellulose

(Sigma). After 4 days at 37 °C, the plates were examined for the presence of plaques under an

inverted microscope.

Ex vivo **Infection of Cells**

Lung-derived FC at 80 % confluency were infected with MCMV$^{tFP-GRB}$ at an MOI of 0.1 for an hour at

37°C. Afterwards, the medium was changed to cultivation medium (high-glucose DMEM (Gibco)

supplemented with 1 % (vol/vol) GlutaMAX, 10 mM HEPES, 50 μM 2-mercaptoethanol, 200 U/ml

penicillin, 200 U/ml streptomycin, 50 μg/ml gentamicin and 5 % FCS). Supernatants were collected

from a different set of wells every 24 hours until 5 days post infection and stored at -80°C. Titration of

the virus was done by infecting MEF cell monolayers grown in 48-well plates with serially diluted

supernatants for an hour at 37 °C. Thereupon, the inoculum was removed, and the cells were overlaid

with culture medium containing 0.75 % (wt/vol) methylcellulose (Sigma). Plaques were counted

following incubation of cells for 4 days at 37 °C.

Statistics and Reproducibility

Unpaired t test, Welch's t test and one-way ANOVA with Dunnett's multiple comparisons test were

performed with GraphPad Prism 9. No values were excluded from statistical analysis as outliers. P-

values < 0.05 were considered significant. * $p<0.05$; ** $p<0.01$; *** $p<0.001$; **** $p<0.0001$. Error bars

denote mean ± SD. Sample size is reported in figure captions. Results were reproduced by two or

more independent experiments performed.

Data Availability

Source data are provided with this paper.

Acknowledgements

This study was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) – Projektnummer 158989968 - SFB 900; the RESIST Excellence cluster (EXC 2155,

project B6); the German Centre for Infection Research (DZIF) of the German Federal Ministry of

Science and Education (BMBF) through TTU 07.834 (to L.C-S.) and the European Union's Horizon

2020 research and innovation program under Grant Agreement 793858 (to K.M.S.). Ilija Brizić is

supported by Research Cooperability program of the Croatian Science Foundation funded by the

European Union from the European Social Fund under the Operational Programme Efficient Human

Resources 2014–2020 (PZS-2019-02-7879). We thank Inge Hollatz-Rangosch and Ayse Barut for

technical assistance. We acknowledge Lothar Gröbe and Maria Höxter from HZI Core Flow Cytometry

Facility as well as Marina Pils, Katrin Schlarman and Petra Beyer from HZI Animal Facility. We thank

Mathias Müller and Birgit Strobl (University of Veterinary Medicine Vienna) for providing $Stat1^{-/-}$ mice.

We acknowledge Caroline Lassnig for providing help with *in vivo* experiments. We thank Hansjörg

Hauser for useful discussions.

Author contributions

Concept and study design, K.M.S. and L.C-S.; Investigation, K.M.S., F.K., N.G., H.M., Y.K.;

Methodology, K.M.S. and L.C-S., Materials: U.R. and T.K.; Resources, L.C-S. and I.B.; Funding

acquisition, L.C-S., K.M.S. and I.B.; Writing of the manuscript, K.M.S. and L.C-S.; Supervision, K.M.S.

and L.C-S.

Competing interests

L.C-S. has submitted a patent for the use of CMV as a vaccine vector. The authors declare no further

competing interests.

References

1. Gatherer, D., et al., *ICTV Virus Taxonomy Profile: Herpesviridae 2021.* J Gen Virol, 2021. **102**(10).

2. Liu, X.F., et al., *Establishment of murine cytomegalovirus latency in vivo is associated with changes in*
histone modifications and recruitment of transcriptional repressors to the major immediate-early
promoter. J Virol, 2008. **82**(21): p. 10922-31.

3. Sinclair, J., *Chromatin structure regulates human cytomegalovirus gene expression during latency,*
reactivation and lytic infection. Biochim Biophys Acta, 2010. **1799**(3-4): p. 286-95.

4. *Cytomegaloviruses.* 2006, Poole: Caister Academic Press.

5. Cohen, J.I., *Herpesvirus latency.* J Clin Invest, 2020. **130**(7): p. 3361-3369.

6. Griffiths, P. and M. Reeves, *Pathogenesis of human cytomegalovirus in the immunocompromised host.*
Nat Rev Microbiol, 2021. **19**(12): p. 759-773.

7. Liu, X.F., et al., *A novel murine model of differentiation-mediated cytomegalovirus reactivation from*
latently infected bone marrow haematopoietic cells. J Gen Virol, 2019. **100**(12): p. 1680-1694.

8. Seckert, C.K., et al., *Liver sinusoidal endothelial cells are a site of murine cytomegalovirus latency and*
reactivation. J Virol, 2009. **83**(17): p. 8869-84.

9. Brautigam, A.R., et al., *Pathogenesis of murine cytomegalovirus infection: the macrophage as a*
permissive cell for cytomegalovirus infection, replication and latency. J Gen Virol, 1979. **44**(2): p. 349-
59.

10. Pollock, J.L., et al., *Latent murine cytomegalovirus infection in macrophages.* Virology, 1997. **227**(1): p.
168-79.

11. Sacher, T., et al., *The major virus-producing cell type during murine cytomegalovirus infection, the*
hepatocyte, is not the source of virus dissemination in the host. Cell Host Microbe, 2008. **3**(4): p. 263-
72.

12. Brune, W., H. Hengel, and U.H. Koszinowski, *A mouse model for cytomegalovirus infection.* Curr Protoc
Immunol, 2001. **Chapter 19**: p. Unit 19.7.

13. Mercer, J.A., C.A. Wiley, and D.H. Spector, *Pathogenesis of murine cytomegalovirus infection:*
identification of infected cells in the spleen during acute and latent infections. J Virol, 1988. **62**(3): p.
987-97.

14. Pomeroy, C., P.J. Hilleren, and M.C. Jordan, *Latent murine cytomegalovirus DNA in splenic stromal cells*
of mice. J Virol, 1991. **65**(6): p. 3330-4.

15. McNab, F., et al., *Type I interferons in infectious disease.* Nat Rev Immunol, 2015. **15**(2): p. 87-103.

16. Najjar, I. and R. Fagard, *STAT1 and pathogens, not a friendly relationship.* Biochimie, 2010. **92**(5): p.
425-44.

17. Presti, R.M., et al., *Interferon gamma regulates acute and latent murine cytomegalovirus infection and*
chronic disease of the great vessels. J Exp Med, 1998. **188**(3): p. 577-88.

18. Gil, M.P., et al., *Biologic consequences of Stat1-independent IFN signaling.* Proc Natl Acad Sci U S A,
2001. **98**(12): p. 6680-5.

19. Kropp, K.A., et al., *Reversible inhibition of murine cytomegalovirus replication by gamma interferon*
(IFN-γ) in primary macrophages involves a primed type I IFN-signaling subnetwork for full establishment
of an immediate-early antiviral state. J Virol, 2011. **85**(19): p. 10286-99.

20. Dağ, F., et al., *Reversible silencing of cytomegalovirus genomes by type I interferon governs virus*
latency. PLoS Pathog, 2014. **10**(2): p. e1003962.

21. Farrell, H.E., et al., *Murine Cytomegalovirus Spreads by Dendritic Cell Recirculation.* mBio, 2017. **8**(5).

22. Hsu, K.M., et al., *Murine cytomegalovirus displays selective infection of cells within hours after systemic*
administration. J Gen Virol, 2009. **90**(Pt 1): p. 33-43.

23. Thom, J.T., et al., *Salivary gland resident APCs are Flt3L- and CCR2-independent macrophage-like cells*
incapable of cross-presentation. Eur J Immunol, 2014. **44**(3): p. 706-14.

24. Scott, C.L., et al., *The Transcription Factor ZEB2 Is Required to Maintain the Tissue-Specific Identities of*
Macrophages. Immunity, 2018. **49**(2): p. 312-325.e5.

25. Bellomo, A., et al., *Reticular Fibroblasts Expressing the Transcription Factor WT1 Define a Stromal Niche*
that Maintains and Replenishes Splenic Red Pulp Macrophages. Immunity, 2020. **53**(1): p. 127-142.e7.

26.   Spallanzani, R.G., et al., *Distinct immunocyte-promoting and adipocyte-generating stromal components*
coordinate adipose tissue immune and metabolic tenors. Sci Immunol, 2019. **4**(35).
27.   Plikus, M.V., et al., *Fibroblasts: Origins, definitions, and functions in health and disease.* Cell, 2021.
184(15): p. 3852-3872.
28.   Yamamoto, S., et al., *Platelet-derived growth factor receptor regulates salivary gland morphogenesis*
via fibroblast growth factor expression. J Biol Chem, 2008. **283**(34): p. 23139-49.
29.   Pezoldt, J., et al., *Single-cell transcriptional profiling of splenic fibroblasts reveals subset-specific innate*
immune signatures in homeostasis and during viral infection. Commun Biol, 2021. **4**(1): p. 1355.
30.   Mederacke, I., et al., *High-yield and high-purity isolation of hepatic stellate cells from normal and*
fibrotic mouse livers. Nat Protoc, 2015. **10**(2): p. 305-15.
31.   Wilhelm, A., et al., *CD248/endosialin critically regulates hepatic stellate cell proliferation during chronic*
liver injury via a PDGF-regulated mechanism. Gut, 2016. **65**(7): p. 1175-85.
32.   Buechler, M.B., et al., *Cross-tissue organization of the fibroblast lineage.* Nature, 2021. **593**(7860): p.
575-579.
33.   Simon, C.O., et al., *Role for tumor necrosis factor alpha in murine cytomegalovirus transcriptional*
reactivation in latently infected lungs. J Virol, 2005. **79**(1): p. 326-40.
34.   Griessl, M., et al., *Stochastic Episodes of Latent Cytomegalovirus Transcription Drive CD8 T-Cell*
"Memory Inflation" and Avoid Immune Evasion. Front Immunol, 2021. **12**: p. 668885.
35.   Sitnik, K.M., et al., *Context-Dependent Development of Lymphoid Stroma from Adult CD34(+)*
Adventitial Progenitors. Cell Rep, 2016. **14**(10): p. 2375-88.
36.   Oduro, J.D., et al., *Murine cytomegalovirus (CMV) infection via the intranasal route offers a robust*
model of immunity upon mucosal CMV infection. J Gen Virol, 2016. **97**(1): p. 185-95.
37.   Cardin, R.D., et al., *The M33 chemokine receptor homolog of murine cytomegalovirus exhibits a*
differential tissue-specific role during in vivo replication and latency. J Virol, 2009. **83**(15): p. 7590-601.
38.   Farrell, H.E. and P.G. Stevenson, *Cytomegalovirus host entry and spread.* J Gen Virol, 2019. **100**(4): p.
545-553.
39.   Muhl, L., et al., *Single-cell analysis uncovers fibroblast heterogeneity and criteria for fibroblast and*
mural cell identification and discrimination. Nat Commun, 2020. **11**(1): p. 3953.
40.   Chung, M.I., et al., *Niche-mediated BMP/SMAD signaling regulates lung alveolar stem cell proliferation*
and differentiation. Development, 2018. **145**(9).
41.   Tegtmeyer, P.K., et al., *STING induces early IFN-β in the liver and constrains myeloid cell-mediated*
dissemination of murine cytomegalovirus. Nat Commun, 2019. **10**(1): p. 2830.
42.   Snyder, C.M., et al., *Cross-presentation of a spread-defective MCMV is sufficient to prime the majority*
of virus-specific CD8+ T cells. PLoS One, 2010. **5**(3): p. e9681.
43.   Snyder, C.M., et al., *Sustained CD8+ T cell memory inflation after infection with a single-cycle*
cytomegalovirus. PLoS Pathog, 2011. **7**(10): p. e1002295.
44.   Farrell, H.E., et al., *Murine Cytomegalovirus Spread Depends on the Infected Myeloid Cell Type.* J Virol,
2019. **93**(15).
45.   Dupont, L. and M.B. Reeves, *Cytomegalovirus latency and reactivation: recent insights into an age old*
problem. Rev Med Virol, 2016. **26**(2): p. 75-89.
46.   Guerrero-Juarez, C.F., et al., *Single-cell analysis reveals fibroblast heterogeneity and myeloid-derived*
adipocyte progenitors in murine skin wounds. Nat Commun, 2019. **10**(1): p. 650.
47.   Sinha, M., et al., *Direct conversion of injury-site myeloid cells to fibroblast-like cells of granulation*
tissue. Nat Commun, 2018. **9**(1): p. 936.
48.   Reinhardt, J.W. and C.K. Breuer, *Fibrocytes: A Critical Review and Practical Guide.* Front Immunol, 2021.
12: p. 784401.
49.   Wallner, B., et al., *Generation of mice with a conditional Stat1 null allele.* Transgenic Res, 2012. **21**(1): p.
217-24.
50.   Oduro, J.D., et al., *Murine cytomegalovirus (CMV) infection via the intranasal route offers a robust*
model of immunity upon mucosal CMV infection. J Gen Virol, 2016. **97**(1): p. 185-195.

51.   Jordan, S., et al., *Virus progeny of murine cytomegalovirus bacterial artificial chromosome pSM3fr show*
699   *reduced growth in salivary Glands due to a fixed mutation of MCK-2.* J Virol, 2011. **85**(19): p. 10346-53.
700   52.   Tischer, B.K., G.A. Smith, and N. Osterrieder, *En passant mutagenesis: a two step markerless red*
701   *recombination system.* Methods Mol Biol, 2010. **634**: p. 421-30.
702   53.   Rand, U., et al., *A Novel Triple-Fluorescent HCMV Strain Reveals Gene Expression Dynamics and Anti-*
703   *Herpesviral Drug Mechanisms.* Front Cell Infect Microbiol, 2021. **10**: p. 536150.
704   54.   Kurz, S., et al., *Latency versus persistence or intermittent recurrences: evidence for a latent state of*
705   *murine cytomegalovirus in the lungs.* J Virol, 1997. **71**(4): p. 2980-7.
706