High-throughput platform of reconstituted depolarization-induced Ca^{2+} release for
skeletal muscle disease mutation validation and drug discovery

Takashi Murayama[1][*], Nagomi Kurebayashi[1], Takuro Numaga-Tomita[2], Takuya Kobayashi[1],
Satoru Okazaki[1], Kyosuke Yamashiro[1], Tsutomu Nakada[2], Shuichi Mori[3], Ryosuke Ishida[3],
Hiroyuki Kagechika[3], Mitsuhiko Yamada[2], Takashi Sakurai[1]

[1]Department of Pharmacology, Juntendo University School of Medicine, Tokyo, Japan.

[2]Department of Molecular Pharmacology, Shinshu University School of Medicine, Matsumoto,
Japan.

[3]Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, Tokyo,
Japan.

Short running title: High-throughput platform for reconstituted DICR
15

[*]Corresponding author: Takashi Murayama, Ph.D., Department of Pharmacology, Juntendo
University School of Medicine, Tokyo 113-8421, Japan. E-mail: takashim@juntendo.ac.jp

Summary

Mutations in essential components for depolarization-induced Ca^{2+} release (DICR) are

implicated into various skeletal muscle diseases. Murayama et al. establish high-throughput

platform for the reconstituted DICR in nonmuscle cells for evaluation of disease-causing

mutations and drug discovery.

Abstract

In skeletal muscle excitation-contraction (E-C) coupling, depolarization of the plasma

membrane triggers Ca^{2+} release from the sarcoplasmic reticulum (SR), referred to as

depolarization-induced Ca^{2+} release (DICR). DICR occurs via the type 1 ryanodine receptor

(RyR1), which physically interacts with the dihydropyridine receptor Cav1.1 subunit in specific

machinery formed with additional essential components including β1a, Stac3 adaptor protein

and junctophilins. Exome sequencing has accelerated the discovery of many novel mutations

in DICR machinery genes in various skeletal muscle diseases. However, functional validation

is time-consuming because it must be performed in a skeletal muscle environment. In this study,

we established a high-throughput platform of the reconstituted DICR in HEK293 cells. The

essential components were effectively transduced into HEK293 cells expressing RyR1 using

baculovirus vectors, and Ca^{2+} release was quantitatively measured with R-CEPIA1er, a

fluorescent ER Ca^{2+} indicator, without contaminant of extracellular Ca^{2+} influx. In these cells,

$[K^+]$-dependent Ca^{2+} release was triggered by chemical depolarization with the aid of inward

rectifying potassium channel, indicating a successful reconstitution of DICR. Using the

platform, we evaluated several Cav1.1 mutations that are implicated in malignant hyperthermia

and myopathy. We also tested several RyR1 inhibitors; whereas dantrolene and Cpd1 inhibited

DICR, procaine had no effect. Furthermore, twitch potentiators such as perchlorate and

thiocyanate accelerated DICR without affecting Ca^{2+}-induced Ca^{2+} release. These results well

reproduced the findings with the muscle fibers and the cultured myotubes. This high-throughput

DICR platform will be highly useful for validation of mutations and drug discovery for skeletal

muscle diseases.

Introduction

In vertebrate skeletal muscle, depolarization of the plasma membrane triggers Ca^{2+} release from the sarcoplasmic reticulum (SR), which is commonly referred to as excitation-contraction (E-C) coupling. This process occurs at triad junctions in which a transverse (T) tubule is closely apposed to terminal cisterna of the SR(Franzini-Armstrong, 2018). In excitation-contraction coupling, the type 1 ryanodine receptor (RyR1), a Ca^{2+} release channel in the SR, is activated by a conformational change of the voltage sensor in the Cav1.1 subunit of the dihydropyridine receptor (DHPR) upon depolarization of the T-tubule membrane. This is referred to as depolarization-induced Ca^{2+} release (DICR)(Rios and Pizarro, 1991; Schneider, 1994). Studies with knockout animals have revealed that functional and structural interactions between Cav1.1 and RyR1 in muscle require at least three additional components, a β1a auxiliary subunit of DHPR(Flucher and Campiglio, 2019; Shishmarev, 2020), the Stac3 adaptor protein(Horstick et al., 2013; Nelson et al., 2013; Reinholt et al., 2013), and junctophilins(Takeshima et al., 2000), which may form specific machinery for DICR(Flucher and Campiglio, 2019; Shishmarev, 2020; Rufenach and Van Petegem, 2021).

Mutations in genes that encode DICR machinery components have been implicated in various types of muscle diseases, including congenital myopathies, periodic paralysis, and malignant hyperthermia (MH)(Agrawal et al., 2018; Flucher, 2020; Lawal et al., 2020). Rapid exome sequencing has accelerated the discovery of novel mutations; however, validation of these mutations by functional testing is critically important for the diagnosis and treatment of these diseases. This has been performed using myotubes from mice that lack DICR machinery components, for example knockout mice of Cav1.1(Tanabe et al., 1988), β1a(Gregg et al., 1996) and Stac3(Polster et al., 2016). While this approach has evaluated several disease-causing mutations in the DICR components (Weiss et al., 2004; Pirone et al., 2010; Eltit et al., 2012; Polster et al., 2016; Perez et al., 2018), it has certain limitations. For example, it is expensive and labor intensive to maintain the mouse lines. Differentiation of cells into myotubes is time consuming and is affected by various factors, such as the cell line used and experimental conditions. Furthermore, transduction of foreign genes into myotubes is often inefficient.

There are no specific treatments for most diseases related to mutations in the DICR machinery. A major cause for the slow pace of novel drug development is the lack of an appropriate drug screening platform for these diseases. We have recently established an efficient high-throughput platform for screening RyR1-targeted drugs using endoplasmic reticulum (ER)

Ca^{2+} measurements in HEK293 cells(Murayama et al., 2018; Murayama and Kurebayashi,
2019) and using this platform we developed Compound 1 (Cpd1), a novel RyR1 inhibitor for
treatment of MH(Mori et al., 2019; Yamazawa et al., 2021). Therefore, functional screening
using HEK293 cells is a promising platform for drug discovery.

Recently, Perni et al. successfully reconstituted DICR by co-expressing five essential
components, RyR1, Cav1.1, β1a, junctophilin-2 (JP2) and Stac3, in non-muscle tsA201
cells(Perni et al., 2017). They demonstrated formation of ER-plasma membrane junctions in
which Cav1.1 is arranged into tetrads, which is indicative of physical links to RyR1(Block et
al., 1988). Depolarization of the plasma membrane using the patch-clamp technique induced a
rapid Ca^{2+} transient without influx of extracellular Ca^{2+}. This provides a basis for developing
reconstituted DICR.

In this study, we established a high-throughput platform for DICR that was
reconstituted in HEK293 cells. Baculovirus infection of essential components greatly improved
transduction efficiency. Chemical depolarization by high concentration potassium [K$^+$]
solutions with the aid of an inward-rectified potassium channel (Kir2.1) enabled the
simultaneous stimulation of many cells. Furthermore, a genetically-encoded ER Ca^{2+} indicator
quantitatively measured DICR without influence of extracellular Ca^{2+} influx. Using this
platform, we successfully evaluated disease-causing mutations in Cav1.1 and tested several
RyR1 inhibitors and DICR modulators. Our high-throughput platform will accelerate the
validation of mutations and drug discovery for skeletal muscle diseases.

Materials and Methods

Generation of stable inducible HEK293 cell lines

HEK293 cells stably expressing R-CEPIA1er and RyR1 were generated as described
previously(Murayama et al., 2018; Murayama and Kurebayashi, 2019). Briefly, a full-length
rabbit RyR1 cDNA was cloned in a tetracycline-induced expression vector (pcDNA5/FRT/TO;
Life Technologies, CA, USA)(Tong et al., 1997). Flp-In T-REx HEK293 cells were co-
transfected with this expression vector and pOG44 in accordance with the manufacturer's
instructions. Clones with suitable doxycycline-induced expression of RyR1 were selected and
used for experiments. The cells were then transfected with an R-CEPIA1er cDNA (a generous
gift from Masamitsu Iino, The University of Tokyo) for stable expression of R-CEPIA1er. Cells
were cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum,

2 mM L-glutamine, 15 µg/ml blasticidin, and 100 µg/ml hygromycin.

Production of baculovirus

Baculovirus that expresses vesicular stomatitis virus G-protein (VSVG) on the virus
envelope(Tani et al., 2001) was produced using the Bac-to-Bac system (LifeTechnologies) as
described previously(Uehara et al., 2017). cDNAs for Cav1.1, β1a, JP2, Stac3, and Kir2.1 were
cloned from mouse skeletal muscle using reverse transcription-PCR (primers pair are listed in
Table 1) and cloned into the modified donor vector (pFastBac1-VSVG-CMV-WPRE).
Baculovirus was produced in Sf9 cells according to the manufacturer's instructions
(LifeTechnologies). Baculovirus titers were immunologically determined using an anti-gp64
antibody (AcV5, Santa Cruz Biotechnology)(Kitts and Green, 1999). Titers were $1–2×10^8$
pfu/mL for all viruses. The baculovirus solutions were stored in the dark at 4°C and used within
one month after production.

Immunocytochemistry

RyR1-HEK293 cells infected with baculovirus for the essential components (5 BV cells) were
fixed with ice-cold methanol and rinsed with phosphate-buffered saline (PBS). The cells were
blocked with 5% BSA/PBS for 30 min and incubated overnight with primary antibodies against
Cav1.1 (sc-514685, Santa Cruz Biotechnology), β1a (MA5-27714, Invitrogen), JP2 (sc-377086,
Santa Cruz Biotechnology), Stac3 (1G8, Abnova), or Kir2.1 (S112B-14, Invitrogen) in 1%
BSA/PBS. After washing three times with PBS, the cells were incubated for 2 h with Alexa594-
labeled goat anti-mouse IgG (Invitrogen) in 1% BSA/PBS. After washing with PBS, the cells
were observed using an all-in-one fluorescence microscope (BZ-X710, Keyence, Osaka, Japan).

Western blotting

RyR1-HEK293 cells with or without baculovirus infection for the essential components were
lysed with Pro-Prep protein extraction solution (iNtRON Biotechnology). The extracted
proteins were separated on 3–15% polyacrylamide gels and transferred to PVDF membranes.
Membranes were probed with the primary antibodies described above, followed by HRP-
labeled anti-mouse IgG (04-18-18, KPL). Positive bands were detected by chemiluminescence
using ImmunoStar LD (Fujifilm Wako Chemicals) as a substrate.

Single-cell Ca^{2+} imaging

RyR1/R-CEPPIA1er HEK293 cells grown on a glass-bottomed dish were treated with

doxycycline to induce the expression of RyR1 and infected with 5 BV 24 h prior to

measurement. Single-cell Ca^{2+} imaging was performed as described previously(Murayama et

al., 2015; Uehara et al., 2017). Measurements were performed at 26°C by perfusing solutions

using an in-line solution heater/cooler (Warner Instruments, Holliston, MA, USA). ER $[Ca^{2+}]$

signals were determined by R-CEPIA1er fluorescence excited at 561 nm using a 40× objective

lens. Cells were perfused with normal Krebs solution (140 mM NaCl, 5 mM KCl, 2 mM $CaCl_2$,

1 mM $MgCl_2$, 11 mM glucose, 5 mM HEPES, pH 7.4) for 30 s, high $[K^+]$ Krebs solution

containing 50 mM $[K^+]$ for 50 s, and normal Krebs solution for 1.5 min. Fluorescence intensities

in individual cells were determined by region of interest (ROI) analysis using AquaCosmos

software (Hamamatsu Photonics, Hamamatsu, Japan).

Time lapse ER $[Ca^{2+}]$ Measurements using a fluorometer

Time lapse ER $[Ca^{2+}]$ measurements were performed using the FlexStation3 fluorometer

(Molecular Devices, San Jose, CA)(Murayama et al., 2018; Murayama and Kurebayashi, 2019).

Stable RyR1/R-CEPIA1er HEK293 cells were seeded on 96-well flat-, clear-bottom black

microplates (#3603, Corning, New York, NY) at a density of $3×10^4$ cells/well in 100 µL culture

medium. One day after seeding, 100 µL culture medium containing 2 µg/mL doxycycline (for

induction of RyR1) and baculovirus solutions for Cav1.1, β1a, JP2, Stac3, and Kir2.1 (2 µL

each) were added to each well. After 24 h, the culture medium were replaced with 81 µL normal

Krebs solution, and the microplate was placed in a FlexStation3 fluorometer preincubated at

37°C. Signals from R-CEPIA1er, which was excited at 560 nm and emitted at 610 nm, were

captured every 5 seconds for 150 seconds. Thirty seconds after starting, 54 µL of the high $[K^+]$

or caffeine solution was applied to the cells. The fluorescence change was expressed as F/F_0 in

which averaged fluorescence intensity of the last 25 seconds (F) was normalized to that of the

initial 25 seconds (F_0). For drug testing, compounds were added to normal Krebs solution at

the indicated concentrations.

For determination of resting ER $[Ca^{2+}]$, cells were incubated with 81 µL normal Krebs

solution, and 54 µL Krebs solution containing 50 mM Ca^{2+} and 50 µM ionomycin was added

30 sec after starting (final concentrations of Ca^{2+} and ionomycin were 20 mM and 20 µM,

respectively). R-CEPIA1er fluorescence was captured every 10 seconds for 6 min.

Measurement of resting membrane potential

The resting membrane potential of baculovirus-infected RyR1-HEK293 cells with or without

Kir2.1 expression was measured under current-clamp conditions in the whole cell configuration

of the patch-clamp method with an Axopatch 200B amplifier (Molecular Devices, San Jose,

CA). Pipettes were fabricated from borosilicate glass capillaries (Kimble) with a vertical pipette

puller (PP-83, Narishige, Tokyo, Japan). Pipettes had series resistance of 3–4 MΩ when filled

with the pipette solution containing (in mM) 120 D-glutamate, 20 KCl, 10 NaCl, 0.1 EGTA, 2

$MgCl_2$, 3 MgATP, and 5 Hepes (pH 7.4 adjusted with KOH). The series resistance was routinely

compensated by ~75% with the amplifier. Bath solution contained (in mM): 140 NaCl, 5 KCl,

11 glucose, 2 $CaCl_2$, 1 $MgCl_2$, and 10 Hepes (pH 7.4 adjusted with NaOH). High [K^+] bath

solution contained (in mM): 95 NaCl, 50 KCl, 11 glucose, 2 $CaCl_2$, 1 $MgCl_2$, and 10 Hepes (pH

7.4 adjusted with NaOH). The resting membrane potential was measured immediately after

breaking into cells.

Statistics

Data are presented as means ± SD. Unpaired Student's t test was used for comparisons between

two groups. One-way analysis of variance (ANOVA), followed by Dunnett's test, was

performed to compare three or more groups. Two-tailed tests were used for all analyses.

Statistical analysis was performed using Prism v9 (GraphPad Software, Inc., La Jolla, USA).

Results

Establishment of a high-throughput platform for DICR using a fluorescence microplate
reader

To reconstitute DICR, we used HEK293 cells stably expressing RyR1 and R-

CEPIA1er, a genetically-encoded fluorescent ER Ca^{2+} indicator(Suzuki et al., 2014), that were

generated for high-throughput screening of novel RyR1 inhibitors(Murayama et al., 2018;

Murayama and Kurebayashi, 2019) (**Fig. 1 A**, left). The other components for DICR, Cav1.1,

β1a, Stac3 and JP2, which were used by Perni et al.(Perni et al., 2017), were transduced using

VSV-G pseudotyped baculovirus (BV), which efficiently infects mammalian cells(Tani et al.,

2001; Uehara et al., 2017) (**Fig. 1 A**, right). In these cells, DICR can be detected by a decrease

in ER Ca^{2+}. To prevent extracellular Ca^{2+} influx through Cav1.1 by depolarization, a N617D

mutation was introduced that abolishes Ca^{2+} conductance(Schredelseker et al., 2010). In

addition, we expressed Kir2.1, an inward-rectified potassium channel in skeletal

muscle(DiFranco et al., 2015), to hyperpolarize the membrane potential because the resting

membrane potential in HEK293 cells is much less (−20 mV) than in skeletal muscle (−90

mV)(Kirkton and Bursac, 2011). Imunocytochemistry revealed that the five proteins (Cav1.1,

β1a, Stac3, JP2, and Kir2.1) were successfully expressed in almost all cells, which we term

hereafter 5 BV cells (**Fig. S1 A**). BV-dependent protein expression was also confirmed by

western blotting of cell lysate (**Fig. S1 B**). Therefore, BV infection effectively transduced the

essential components of DICR in RyR1/R-CEPIA1er-HEK293 cells.

To detect DICR, we used a fluorescence microplate reader (FlexStation3), which

measures time-lapse R-CEPIA1er fluorescence in 96-well plates. Fluorescence signals were

collected in a 1.5 mm diameter area; therefore, an averaged signal from thousands of cells was

detected in each well. We adopted chemical depolarization using high $[K^+]$ solutions, which

can simultaneously depolarize all the cells to trigger DICR. The cells were initially incubated

with normal Krebs solution containing 5 mM $[K^+]$. Then, solutions containing various $[K^+]$

concentrations were added to give $[K^+]$ concentrations of 5–61 mM. In cells without BV

infection, no apparent changes in fluorescence were observed with any $[K^+]$ concentration (**Fig.**

1 B, left). Caffeine (10 mM) reduced the R-CEPIA1er fluorescence, confirming the functional

expression of RyR1. In 5 BV cells, high $[K^+]$ solution rapidly reduced the R-CEPIA1er

fluorescence in a $[K^+]$-dependent manner (**Fig. 1 B**, right). To monitor ER Ca^{2+} changes in

individual cells, we also observed R-CEPIA1er fluorescence using laser scanning confocal

microscopy. In 5 BV cells, high $[K^+]$ solution caused a large and transient reduction in the R-

CEPIA1er fluorescence in 90% of cells (**Movie S1** and **Fig. S2**).

For quantitative analysis, we normalized the averaged fluorescence intensity of the last

25 seconds (F) to that of the initial 25 seconds (F_0) to correct variations in the initial

fluorescence intensity between wells. The F/F_0 plot as a function of $[K^+]$ concentrations

demonstrated $[K^+]$ dependence with an EC_{50} of 9.8 $\pm$ 0.1 mM $[K^+]$ (**Fig. 1 C**). The initial

fluorescence change after stimulation (i.e., 35 seconds), which may reflect the rate of Ca^{2+}

release, provided larger EC_{50} (17.0 ± 0.4 mM). However, variations were larger, especially near

EC_{50} value because of rapid changes in fluorescence intensity (**Fig. 1 C**). The fluorescence

change by 61 mM $[K^+]$, which is expressed as $1-F/F_0$ (**Fig. 1 C**), reached ~80% of that induced

by caffeine in 5 BV cells, indicating a high degree of DICR reconstitution (**Fig. 1 D**). Taken

together, these results indicate successful establishment of a high-throughput platform for DICR.

The original method for DICR reconstitution monitors cytoplasmic $[Ca^{2+}]$(Perni et al.,

2017); therefore, we also assessed DICR by measuring cytoplasmic Ca^{2+} using G-GECO1.1, a

genetically-encoded Ca^{2+} indicator(Zhao et al., 2011). In cells without BV infection, a

substantial increase in $[Ca^{2+}]$ was observed by high $[K^+]$, indicating the existence of endogenous

Ca^{2+} influx pathways by depolarization (**Fig. S3 A**). In 5 BV cells, DICR was detected by rapid

and large increases in $[Ca^{2+}]$, which declined with time (**Fig. S3 B**). The initial fluorescence

change after stimuli (i.e., 35 seconds) provided dose-dependent curve with EC_{50} of 16.4 ± 0.5

mM, which is similar to the value determined by initial fluorescence change in ER Ca^{2+}

measurement (see **Fig. 1 C**).

Validation of the reconstituted DICR

We next determined whether the reconstituted platform reproduces DICR in skeletal

muscle. Initially, we examined the role of the essential components. Removal of Cav1.1 resulted

in complete loss of high $[K^+]$-induced Ca^{2+} release (**Fig. 2A-C**). Ca^{2+} release was also lost in

cells lacking β1a or Stac3, and in cells expressing RyR2 instead of RyR1 (**Fig. 2 C**). Removal

of JP2 significantly reduced the fluorescence change by 61 mM $[K^+]$ with a large rightward

shift in $[K^+]$ dependence (**Fig. 2 B, C**). These results are consistent with previous findings of

mice lacking each component(Tanabe et al., 1988; Takeshima et al., 1994; Gregg et al., 1996;

Takeshima et al., 2000; Polster et al., 2016).

Skeletal muscle Cav1.1 forms a complex with three auxiliary subunits, β1a, $α_2$-δ and

$γ_1$(Catterall, 2000). $α_2$-δ and $γ_1$ may play modulatory roles(Flucher et al., 2005). Additional

expression of $α_2$-δ had no effect on high $[K^+]$-induced Ca^{2+} release (**Fig. 2 D, E**). Expression of

γ1 did not significantly affect high $[K^+]$-induced Ca^{2+} release at 20 mM or lower $[K^+]$ but it

reduced the fluorescence change at higher $[K^+]$ concentrations (**Fig. 2 D, E**). In $γ_1$ cells, R-

CEPIA1er fluorescence decreased but increased again with time, suggesting inactivation of the

Ca^{2+} release (**Fig. 2 A**). These findings are consistent with previous reports of the effects of $α_2$-

δ(Obermair et al., 2005) and $γ_1$(Ursu et al., 2004) on DICR. Therefore, our reconstituted

platform can reproduce DICR in skeletal muscle and is valid for evaluating disease-causing

mutations or drugs.

We also tested the importance of Kir2.1 on the reconstituted DICR platform. We

measured membrane potential of cells under current-clamp conditions using pipette solution

containing ~145 mM $[K^+]$ (**Fig. 3 A**). The membrane potential of 5 BV-infected cells was -71

± 3 mV in normal Krebs solution and became depolarized to -14 ± 3 mV by high $[K^+]$ solution

containing 50 mM $[K^+]$. In contrast, membrane potential was already depolarized in the cells

without Kir2.1 (-18 ± 2 mV) in normal Krebs solution, which is consistent with the previous

report(Kirkton and Bursac, 2011). Without Kir2.1, R-CEPIA1er fluorescence intensity was

substantially reduced and high [K$^+$]-induced Ca^{2+} release was completely abolished (**Fig. 3 B**).

We quantified resting ER [Ca^{2+}] by determining the maximum fluorescence intensity of R-

CEPIA1er using ionomycin/Ca^{2+} (**Fig. 3 C**). The resting ER [Ca^{2+}] was severely reduced in

cells without Kir2.1 (**Fig. 3 D**). These results suggest that persistently activated DICR activity

under the depolarized conditions causes ER Ca^{2+} depletion. In support of this, further removal

of β1a or Stac3, which completely abolishes DICR activity, restored the ER [Ca^{2+}] (**Fig. 3 D**).

Therefore, Kir2.1 is indispensable for our reconstituted DICR platform.

Evaluation of disease-causing mutations in Cav1.1

Using the high-throughput DICR platform, we evaluated mutations in Cav1.1 which

are implicated in various muscle diseases(Beam et al., 2017; Flucher, 2020). We initially tested

known MH mutations, R174W(Carpenter et al., 2009), R1086H(Stewart et al., 2001), and

T1354S(Pirone et al., 2010) (**Fig. 4 A**). We found that ER [Ca^{2+}] was severely reduced for

R1086H (**Fig. 4 B**) and that R1086H exhibited a smaller fluorescence change in response to

high [K$^+$] (**Fig. 4 C, D**) compared with the wild type (WT) or the other mutants. This was partly

caused by low ER [Ca^{2+}], which reduces maximum fluorescence changes. In addition, R1086H

showed an enhanced [K$^+$]-dependence, i.e., lower EC$_{50}$ value for [K$^+$] (**Fig. 4 C, E**). In contrast,

no or only minor changes in ER [Ca^{2+}] and [K$^+$] dependence were observed for R174W and

T1354S cells (**Fig. 4 B–E**). We anticipated that the reduction in ER [Ca^{2+}] in R1086H cells was

caused by activation of Cav1.1 at resting membrane potential because of its hypersensitivity.

To test the hypothesis, we hyperpolarized the membrane potential which reduced the activation

of Cav1.1. Reduction of [K$^+$] in the Krebs solution from 5 mM to 2.5 mM successfully

hyperpolarized the membrane potential by ~15 mV (**Fig. S4 A**). This substantially increased

ER [Ca^{2+}] in R1086H cells to a level comparable to that in WT cells (**Fig. S4 B**). R1086H cells

exhibited a clear greater sensitivity to [K$^+$] than WT cells (**Fig. S4 C–E**), supporting our

hypothesis of R1086H hyperactivation.

Previous functional characterization of these Cav1.1 mutants consistently showed

enhanced caffeine sensitivity, i.e., acceleration of Ca^{2+}-induced Ca^{2+} release (CICR)(Weiss et

al., 2004; Pirone et al., 2010; Eltit et al., 2012). We tested caffeine dependence of these mutants

in our platform. Caffeine released Ca^{2+} from WT cells in a dose-dependent manner (**Fig. 4 F**).

R1086H cells also exhibited a caffeine dependence but with reduced fluorescence change by

61 mM [K$^+$] (**Fig. 4 G**) and a significantly smaller EC$_{50}$ (**Fig. 4 H**), indicating an enhanced

caffeine sensitivity. In contrast, caffeine dependence was not changed in R174W or T1354S

cells (**Fig. 4 F–H**).

We next examined the effect of myopathy-related mutations in Cav1.1 on DICR
activity. We chose four mutations, E100K, F275L, P742Q, and L1367V(Schartner et al., 2017),
none of which have been functionally characterized (**Fig. 5 A**). ER [Ca^{2+}] was not changed by
any of these mutations (**Fig. 5 B**). DICR activity was completely lost in the F275L mutant (**Fig.**
5 C, D) and there was a substantial rightward shift in [K$^+$] dependence for P742Q (**Fig. 5 C**);
The EC$_{50}$ value was threefold higher in P742Q (33.2 ± 1.0 mM) compared with WT (10.3 ± 0.9
mM) (**Fig. 5 E**). The other two mutations (E100K and L1367V) had no significant effects on
[K$^+$] dependence (**Fig. 5 C-E**). We also tested the effects of these mutations on caffeine-induced
Ca^{2+} release. No substantial changes were observed with any of the mutations (**Fig. 5 F–H**).

Effect of drugs on DICR activity

The high-throughput DICR platform is expected to be useful for screening drugs for
muscle diseases. To test this possibility, we examined the effects of known DICR modulators.
We initially tested three RyR1 inhibitors (**Fig. 6 A**). Dantrolene is a well-known RyR1 inhibitor
that is clinically used for MH(Riazi et al., 2018). Cpd1 is a novel potent RyR1-selective
inhibitor that we recently developed(Mori et al., 2019; Yamazawa et al., 2021). Procaine
inhibits CICR but not DICR in frog skeletal muscle(Thorens and Endo, 1975; Kashiyama et al.,
2010). Dantrolene (10 μM) and Cpd1 (3 μM) caused rightward shifts in [K$^+$] dependence
compared with the control, indicating that they inhibit DICR (**Fig. 6 B, C**). This is consistent
with the suppression of twitch and tetanic tension in isolated muscles and with reduced *in vivo*
muscle weakness by dantrolene(Meyler et al., 1979; Leslie and Part, 1981) and
Cpd1(Yamazawa et al., 2021). Procaine (5 mM), in contrast, did not inhibit DICR (**Fig. 6 B,**
C). All three compounds shifted the caffeine dependence rightward, indicating inhibition of
CICR (**Fig. 6 D, E**). These results indicate that dantrolene and Cpd1 inhibit both DICR and
CICR, whereas procaine is a CICR-selective inhibitor.

Lyotropic anions, such as perchlorate and thiocyanate, potentiate E-C coupling in
skeletal muscle(Luttgau et al., 1983; Huang, 1986; Csernoch et al., 1987; Delay et al., 1990;
Gonzalez and Rios, 1993). Perchlorate (10 mM) and thiocyanate (10 mM) significantly shifted
the [K$^+$] dependence leftward (**Fig. 7 A, B**). In contrast, they did not affect caffeine dependence
(**Fig. 7 C, D**). These results indicate that lyotropic anions potentiate DICR but not CICR. This
is consistent with previous findings showing that the potentiating effects of lyotropic anions are
primarily caused by shifting voltage dependence of the charge movement toward more negative

potentials(Luttgau et al., 1983; Huang, 1986; Csernoch et al., 1987; Delay et al., 1990; Gonzalez and Rios, 1993). Taken together, our high-throughput platform reproduces the effects of known DICR modulators, indicating that it can be used to screen for and test drugs that affect DICR.

Discussion

In this study, we established a high-through platform for DICR that was reconstituted in HEK293 cells expressing RyR1 and R-CEPIA1er. We made three key improvements to the original method by Perni et al. (Perni et al., 2017). First, the essential components were transduced using VSV-G pseudotyped baculovirus, which can effectively infect a wide variety of mammalian cells without toxicity(Tani et al., 2001). This greatly increased the transduction efficiency; almost all cells expressed the essential components (**Fig. S1**). Second, depolarization of the plasma membrane was induced by high $[K^+]$ solution. This enabled us to simultaneously stimulate many cells (**Fig. 1, Fig. S2** and **movie S1**). For this purpose, we expressed Kir2.1, an inward-rectifying potassium channel, which effectively hyperpolarized the membrane potential (Kirkton and Bursac, 2011) (**Fig. 3**). Third, ER $[Ca^{2+}]$, instead of cytoplasmic $[Ca^{2+}]$, was measured to detect DICR. This is critically important to avoid contamination of signals from extracellular Ca^{2+} influx. Indeed, ER $[Ca^{2+}]$ was not changed by high $[K^+]$ depolarization in cells without essential components (**Fig. 1 A**), in contrast to substantial increases in cytoplasmic $[Ca^{2+}]$ (**Fig. S4 A**). These improvements allow quantitative measurements of DICR using a microplate reader.

In our reconstitution, five components, RyR1, Cav1.1, β1a, JP2 and Stac3, are essential for DICR; removal of each component abolished or reduced the DICR activity (**Fig. 2 A, B**). This is consistent with previous findings from mice lacking each component(Tanabe et al., 1988; Takeshima et al., 1994; Gregg et al., 1996; Takeshima et al., 2000; Polster et al., 2016). In addition, the effects of α2-δ and γ1 auxiliary subunits on DICR activity corresponded with findings of previous reports (Ursu et al., 2004; Obermair et al., 2005) (**Fig. 2 C, D**). Therefore, our reconstituted platform successfully reproduces DICR in skeletal muscle. We found that a substantial DICR still occurred without JP2 (**Fig. 2 A, B**). This is in contrast to reports by Perni et al.(Perni et al., 2017; Perni and Beam, 2022), in which no voltage-gated Ca^{2+} release was observed in reconstituted cells without junctophilins. A possible reason for this is differences in expression methods. Our VSV-G baculovirus is more effective at transduction than standard lipofection; therefore, sufficient proteins were expressed that might spontaneously interact with each other without junctophilins.

We quantitatively evaluated the effect of disease-casing mutations or drugs using the EC_{50}

value for $[K^+]$. In our experimental conditions, the EC_{50} value for $[K^+]$ was about 10 mM (**Fig.**

1 C). The value is lower than that for skeletal muscle(Caputo and Bolanos, 1994) or

myotubes(Yang et al., 2006) in high $[K^+]$-induced Ca^{2+} transients. This may be partly explained

by measurement differences. We calculated the EC_{50} from the fluorescence intensity for 125–

150 seconds after stimulation (**Fig. 1 C**). This reduced the EC_{50} compared with the value

obtained just after stimulation in ER $[Ca^{2+}]$ (**Fig. 1 C**) or cytoplasmic $[Ca^{2+}]$ measurements (**Fig.**

S3). Nevertheless, our results reproduced the effects of disease-causing mutations (**Fig. 3–5**) or

drugs (**Fig. 6, 7**), strongly indicating that the EC_{50} value for $[K^+]$ is a valid parameter for

quantitative evaluation.

Formation of DICR machinery can cause peripheral localization of the components, e.g.,

RyR1 and Cav1.1, in the reconstituted cells(Perni et al., 2017; Perni and Beam, 2022). In our

reconstituted cells, however, no such peripheral-specific localization was observed; the

components were localized throughout cells except for in nuclei (**Fig. S1**). This may be because

of high levels of the components expressed by baculovirus, which not only accelerates

formation of the DICR machinery (as discussed above), but also increases the number of

uncoupled components.

Mutations in Cav1.1 are implicated in various muscle diseases, including MH and

myopathy(Beam et al., 2017; Flucher, 2020). Among three MH mutations (R174W, R1086H

and T1354S), we demonstrated that R1086H, but not R174W orT1354S, significantly

accelerated DICR (**Fig. 4 C–E**). This is consistent with previous findings using Cav1.1-

deficient (dysgenic) myotubes (Weiss et al., 2004; Pirone et al., 2010; Eltit et al., 2012). All

three mutations show increased sensitivity to caffeine-induced Ca^{2+} release compared with WT,

indicating enhanced CICR activity (Weiss et al., 2004; Pirone et al., 2010; Eltit et al., 2012).

However, we showed that caffeine sensitivity was increased only for R1086H (**Fig. 4 F–H**). A

possible reason for this difference is the cells used. We used stable RyR1-overexpressing cells.

This may cause an excess amount of 'uncoupled' RyR1, which might mask the enhanced

sensitivity of caffeine dependence of the 'coupled' RyR1 by Cav1.1 mutations.

Most myopathy-related mutations identified in Cav1.1 have not been functionally

characterized. We evaluated four mutations (E100K, F275L, P742Q and L1367V) found in

patients with congenital myopathy(Schartner et al., 2017) (**Fig. 5**). We demonstrated that P742Q

substantially inhibited DICR with a large rightward shifted $[K^+]$ dependence. This proline

resides in the II-III loop which interacts with RyR1 or Stac3(Wong King Yuen et al., 2017). Substitution of the proline to threonine, the residue found in the corresponding position of CaV1.2, diminished DICR in dysgenic myotubes(Kugler et al., 2004). Therefore, P742Q may cause myopathy by reducing DICR activity. E100K and L1367V had no or only minor impacts on DICR activity. These mutations might cause muscle weakness by reasons other than reduced DICR. F275L is recessive because it was found in two compound heterozygous sisters carrying F275L and the nonsense mutation, p.L791Cfs*37(Schartner et al., 2017). F275L completely abolished DICR without any changes in ER $[Ca^{2+}]$ or caffeine sensitivity. This does not fit with a recessive mutation; if this is the case, patients will have no DICR activity. Further investigations are required for this mutation.

There are no specific treatments for most DICR-related diseases; therefore, there is an urgent need to develop novel drugs for these diseases(Agrawal et al., 2018; Flucher, 2020; Lawal et al., 2020). We here evaluated DICR modulators. Among three known RyR1 inhibitors, dantrolene and Cpd1 suppressed DICR (**Fig. 6**). This is consistent with the suppression of twitch and tetanic tensions of isolated muscles and of reduced *in vivo* muscle weakness by dantrolene(Meyler et al., 1979; Leslie and Part, 1981) and Cpd1(Yamazawa et al., 2021). Interestingly, procaine did not suppress DICR (**Fig. 6**). Procaine inhibits CICR but not DICR in frog skeletal muscle(Thorens and Endo, 1975) and reconstituted frog α-RyR and β-RyR in RyR1-deficient myotubes(Kashiyama et al., 2010). Our data show similar procaine action in a mammalian system. We also demonstrated that lyotropic anions (perchlorate and thiocyanate) accelerate DICR (**Fig. 7**). This reproduces previous findings using skeletal muscle fibers(Luttgau et al., 1983; Huang, 1986; Csernoch et al., 1987; Delay et al., 1990; Gonzalez and Rios, 1993). Interestingly, lyotropic anions did not affect caffeine dependence, suggesting no effect on CICR (**Fig. 7**). Therefore, our high-throughput platform will also be useful for drug testing and screening for novel drugs that affect DICR.

In summary, we established a high-throughput platform for skeletal muscle DICR. This platform is highly efficient and quantitative thus will be useful for both evaluation of disease-causing mutations and the development of novel drugs for DICR-related diseases.

Acknowledgments

We thank Ikue Hiraga and the Laboratory of Proteomics and Biomolecular Science, Biomedical Research Core Facilities, Juntendo University Graduate School of Medicine, for technical

assistance. We thank Jeremy Allen, PhD, from Edanz (https://jp.edanz.com/ac) for editing a
draft of this manuscript. This work was supported by JSPS KAKENHI (19H03404 and
22H02805 to T.M., 19K07105 and 22K06652 to N.K., 20K11368 to T.K.), the Platform Project
for Supporting Drug Discovery and Life Science Research (Basis for Supporting Innovative
Drug Discovery and Life Science Research (BINDS)) from the Japan Agency for Medical
Research and Development (AMED) (JP20am0101080 to T.M. and N.K. and JP20am0101098
to H.K.), an Intramural Research Grant for Neurological and Psychiatric Disorders from the
National Center of Neurology and Psychiatry (2-5 to T.M.), the Vehicle Racing
Commemorative Foundation (6237 and 6303 to T.M.), and the Cooperative Research Project
of Research Center for Biomedical Engineering (to H.K.). The authors declare no competing
financial interests.

Author contributions

T. Murayama, N. Kurebayashi, S. Okazaki, K. Yamashiro carried out the cell biological
experiments; T. Numaga-Tomita and M. Yamada performed patch-clamp experiments; T.
Nakada, S. Mori, R. Ishida, and H. Kagechika provided experimental tools; T. Murayama, N.
Kurebayashi, T. Numaga-Tomita, T. Kobayashi, T. Nakada, M. Yamada, and T. Sakurai
analyzed the experimental data; T. Murayama, N. Kurebayashi, T. Numaga-Tomita and M.
Yamada wrote the manuscript. All authors reviewed and approved the manuscript.

References

Agrawal, A., G. Suryakumar, and R. Rathor. 2018. Role of defective Ca(2+) signaling in
skeletal muscle weakness: Pharmacological implications. *J Cell Commun Signal*.
12:645-659.

Beam, T.A., E.F. Loudermilk, and D.F. Kisor. 2017. Pharmacogenetics and pathophysiology of
CACNA1S mutations in malignant hyperthermia. *Physiol Genomics*. 49:81-87.

Block, B.A., T. Imagawa, K.P. Campbell, and C. Franzini-Armstrong. 1988. Structural evidence
for direct interaction between the molecular components of the transverse
tubule/sarcoplasmic reticulum junction in skeletal muscle. *J Cell Biol*. 107:2587-2600.

Caputo, C., and P. Bolanos. 1994. Fluo-3 signals associated with potassium contractures in
single amphibian muscle fibres. *J Physiol*. 481 (Pt 1):119-128.

Carpenter, D., C. Ringrose, V. Leo, A. Morris, R.L. Robinson, P.J. Halsall, P.M. Hopkins, and
M.A. Shaw. 2009. The role of CACNA1S in predisposition to malignant hyperthermia.
BMC Med Genet. 10:104.

Catterall, W.A. 2000. Structure and regulation of voltage-gated Ca2+ channels. *Annu Rev Cell*
Dev Biol. 16:521-555.

Csernoch, L., L. Kovacs, and G. Szucs. 1987. Perchlorate and the relationship between charge
movement and contractile activation in frog skeletal muscle fibres. *J Physiol*. 390:213-
227.

Delay, M., D.E. Garcia, and J.A. Sanchez. 1990. The effects of lyotropic anions on charge
movement, calcium currents and calcium signals in frog skeletal muscle fibres. *J*
Physiol. 425:449-469.

DiFranco, M., C. Yu, M. Quinonez, and J.L. Vergara. 2015. Inward rectifier potassium currents
in mammalian skeletal muscle fibres. *J Physiol*. 593:1213-1238.

Eltit, J.M., R.A. Bannister, O. Moua, F. Altamirano, P.M. Hopkins, I.N. Pessah, T.F. Molinski,
J.R. Lopez, K.G. Beam, and P.D. Allen. 2012. Malignant hyperthermia susceptibility
arising from altered resting coupling between the skeletal muscle L-type Ca2+ channel
and the type 1 ryanodine receptor. *Proc Natl Acad Sci U S A*. 109:7923-7928.

Flucher, B.E. 2020. Skeletal muscle CaV1.1 channelopathies. *Pflugers Arch*. 472:739-754.

Flucher, B.E., and M. Campiglio. 2019. STAC proteins: The missing link in skeletal muscle EC
coupling and new regulators of calcium channel function. *Biochim Biophys Acta Mol*
Cell Res. 1866:1101-1110.

Flucher, B.E., G.J. Obermair, P. Tuluc, J. Schredelseker, G. Kern, and M. Grabner. 2005. The
role of auxiliary dihydropyridine receptor subunits in muscle. *J Muscle Res Cell Motil*.
26:1-6.

Franzini-Armstrong, C. 2018. The relationship between form and function throughout the
history of excitation-contraction coupling. *J Gen Physiol*. 150:189-210.

Gonzalez, A., and E. Rios. 1993. Perchlorate enhances transmission in skeletal muscle
excitation-contraction coupling. *J Gen Physiol*. 102:373-421.

Gregg, R.G., A. Messing, C. Strube, M. Beurg, R. Moss, M. Behan, M. Sukhareva, S. Haynes,
J.A. Powell, R. Coronado, and P.A. Powers. 1996. Absence of the beta subunit (cchb1)
of the skeletal muscle dihydropyridine receptor alters expression of the alpha 1 subunit
and eliminates excitation-contraction coupling. *Proc Natl Acad Sci U S A*. 93:13961-
13966.

Horstick, E.J., J.W. Linsley, J.J. Dowling, M.A. Hauser, K.K. McDonald, A. Ashley-Koch, L.
Saint-Amant, A. Satish, W.W. Cui, W. Zhou, S.M. Sprague, D.S. Stamm, C.M. Powell,
M.C. Speer, C. Franzini-Armstrong, H. Hirata, and J.Y. Kuwada. 2013. Stac3 is a
component of the excitation-contraction coupling machinery and mutated in Native
American myopathy. *Nat Commun*. 4:1952.

Huang, C.L. 1986. The differential effects of twitch potentiators on charge movements in frog
skeletal muscle. *J Physiol*. 380:17-33.

Kashiyama, T., T. Murayama, E. Suzuki, P.D. Allen, and Y. Ogawa. 2010. Frog alpha- and beta-
ryanodine receptors provide distinct intracellular Ca2+ signals in a myogenic cell line.
PLoS One. 5:e11526.

Kirkton, R.D., and N. Bursac. 2011. Engineering biosynthetic excitable tissues from
unexcitable cells for electrophysiological and cell therapy studies. *Nat Commun*. 2:300.

Kitts, P.A., and G. Green. 1999. An immunological assay for determination of baculovirus titers
in 48 hours. *Anal Biochem*. 268:173-178.

Kugler, G., R.G. Weiss, B.E. Flucher, and M. Grabner. 2004. Structural requirements of the
dihydropyridine receptor alpha1S II-III loop for skeletal-type excitation-contraction
coupling. *J Biol Chem*. 279:4721-4728.

Lawal, T.A., J.J. Todd, J.W. Witherspoon, C.G. Bonnemann, J.J. Dowling, S.L. Hamilton, K.G.
Meilleur, and R.T. Dirksen. 2020. Ryanodine receptor 1-related disorders: an historical
perspective and proposal for a unified nomenclature. *Skelet Muscle*. 10:32.

Leslie, G.C., and N.J. Part. 1981. The action of dantrolene sodium on rat fast and slow muscle
in vivo. *Br J Pharmacol*. 72:665-672.

Luttgau, H.C., G. Gottschalk, L. Kovacs, and M. Fuxreiter. 1983. How perchlorate improves
excitation-contraction coupling in skeletal muscle fibers. *Biophys J*. 43:247-249.

Meyler, W.J., I. Mols-Thurkow, A.H. Scaf, S. Sargo, and H. Wesseling. 1979. The effect of
dantrolene sodium on rat skeletal muscle in relation to the plasma concentration. *Eur J*
Pharmacol. 53:335-342.

Mori, S., H. Iinuma, N. Manaka, M. Ishigami-Yuasa, T. Murayama, Y. Nishijima, A. Sakurai,
R. Arai, N. Kurebayashi, T. Sakurai, and H. Kagechika. 2019. Structural development
of a type-1 ryanodine receptor (RyR1) Ca(2+)-release channel inhibitor guided by
endoplasmic reticulum Ca(2+) assay. *Eur J Med Chem*. 179:837-848.

Murayama, T., and N. Kurebayashi. 2019. Assays for Modulators of Ryanodine Receptor
(RyR)/Ca(2+) Release Channel Activity for Drug Discovery for Skeletal Muscle and
Heart Diseases. *Curr Protoc Pharmacol*. 87:e71.

Murayama, T., N. Kurebayashi, M. Ishigami-Yuasa, S. Mori, Y. Suzuki, R. Akima, H. Ogawa,
J. Suzuki, K. Kanemaru, H. Oyamada, Y. Kiuchi, M. Iino, H. Kagechika, and T. Sakurai.
2018. Efficient High-Throughput Screening by Endoplasmic Reticulum Ca^{2+}
Measurement to Identify Inhibitors of Ryanodine Receptor Ca^{2+}-Release Channels. *Mol*
Pharmacol. 94:722-730.

Murayama, T., N. Kurebayashi, T. Yamazawa, H. Oyamada, J. Suzuki, K. Kanemaru, K. Oguchi,
M. Iino, and T. Sakurai. 2015. Divergent Activity Profiles of Type 1 Ryanodine Receptor
Channels Carrying Malignant Hyperthermia and Central Core Disease Mutations in the
Amino-Terminal Region. *PLoS One*. 10:e0130606.

Nelson, B.R., F. Wu, Y. Liu, D.M. Anderson, J. McAnally, W. Lin, S.C. Cannon, R. Bassel-
Duby, and E.N. Olson. 2013. Skeletal muscle-specific T-tubule protein STAC3 mediates
voltage-induced Ca2+ release and contractility. *Proc Natl Acad Sci U S A*. 110:11881-
11886.

Obermair, G.J., G. Kugler, S. Baumgartner, P. Tuluc, M. Grabner, and B.E. Flucher. 2005. The
Ca2+ channel alpha2delta-1 subunit determines Ca2+ current kinetics in skeletal muscle
but not targeting of alpha1S or excitation-contraction coupling. *J Biol Chem*. 280:2229-
2237.

Perez, C.F., J.M. Eltit, J.R. Lopez, D. Bodnar, A.F. Dulhunty, S. Aditya, and M.G. Casarotto.
2018. Functional and structural characterization of a novel malignant hyperthermia-

susceptible variant of DHPR-beta1a subunit (CACNB1). *Am J Physiol Cell Physiol.* 314:C323-C333.

Perni, S., and K. Beam. 2022. Junctophilins 1, 2, and 3 all support voltage-induced Ca2+ release despite considerable divergence. *J Gen Physiol.* 154.

Perni, S., M. Lavorato, and K.G. Beam. 2017. De novo reconstitution reveals the proteins required for skeletal muscle voltage-induced Ca(2+) release. *Proc Natl Acad Sci U S A.* 114:13822-13827.

Pirone, A., J. Schredelseker, P. Tuluc, E. Gravino, G. Fortunato, B.E. Flucher, A. Carsana, F. Salvatore, and M. Grabner. 2010. Identification and functional characterization of malignant hyperthermia mutation T1354S in the outer pore of the Cavalpha1S-subunit. *Am J Physiol Cell Physiol.* 299:C1345-1354.

Polster, A., B.R. Nelson, E.N. Olson, and K.G. Beam. 2016. Stac3 has a direct role in skeletal muscle-type excitation-contraction coupling that is disrupted by a myopathy-causing mutation. *Proc Natl Acad Sci U S A.* 113:10986-10991.

Reinholt, B.M., X. Ge, X. Cong, D.E. Gerrard, and H. Jiang. 2013. Stac3 is a novel regulator of skeletal muscle development in mice. *PLoS One.* 8:e62760.

Riazi, S., N. Kraeva, and P.M. Hopkins. 2018. Updated guide for the management of malignant hyperthermia. *Can J Anaesth.* 65:709-721.

Rios, E., and G. Pizarro. 1991. Voltage sensor of excitation-contraction coupling in skeletal muscle. *Physiol Rev.* 71:849-908.

Rufenach, B., and F. Van Petegem. 2021. Structure and function of STAC proteins: Calcium channel modulators and critical components of muscle excitation-contraction coupling. *J Biol Chem.* 297:100874.

Schartner, V., N.B. Romero, S. Donkervoort, S. Treves, P. Munot, T.M. Pierson, I. Dabaj, E. Malfatti, I.T. Zaharieva, F. Zorzato, O. Abath Neto, G. Brochier, X. Lornage, B. Eymard, A.L. Taratuto, J. Bohm, H. Gonorazky, L. Ramos-Platt, L. Feng, R. Phadke, D.X. Bharucha-Goebel, C.J. Sumner, M.T. Bui, E. Lacene, M. Beuvin, C. Labasse, N. Dondaine, R. Schneider, J. Thompson, A. Boland, J.F. Deleuze, E. Matthews, A.N. Pakleza, C.A. Sewry, V. Biancalana, S. Quijano-Roy, F. Muntoni, M. Fardeau, C.G. Bonnemann, and J. Laporte. 2017. Dihydropyridine receptor (DHPR, CACNA1S) congenital myopathy. *Acta Neuropathol.* 133:517-533.

Schneider, M.F. 1994. Control of calcium release in functioning skeletal muscle fibers. *Annu*

577   *Rev Physiol*. 56:463-484.

578   Schredelseker, J., M. Shrivastav, A. Dayal, and M. Grabner. 2010. Non-Ca2+-conducting Ca2+
579   channels in fish skeletal muscle excitation-contraction coupling. *Proc Natl Acad Sci U*
580   *S A*. 107:5658-5663.

581   Shishmarev, D. 2020. Excitation-contraction coupling in skeletal muscle: recent progress and
582   unanswered questions. *Biophys Rev*. 12:143-153.

583   Stewart, S.L., K. Hogan, H. Rosenberg, and J.E. Fletcher. 2001. Identification of the
584   Arg1086His mutation in the alpha subunit of the voltage-dependent calcium channel
585   (CACNA1S) in a North American family with malignant hyperthermia. *Clin Genet*.
586   59:178-184.

587   Suzuki, J., K. Kanemaru, K. Ishii, M. Ohkura, Y. Okubo, and M. Iino. 2014. Imaging
588   intraorganellar Ca2+ at subcellular resolution using CEPIA. *Nat Commun*. 5:4153.

589   Takeshima, H., M. Iino, H. Takekura, M. Nishi, J. Kuno, O. Minowa, H. Takano, and T. Noda.
590   1994. Excitation-contraction uncoupling and muscular degeneration in mice lacking
591   functional skeletal muscle ryanodine-receptor gene. *Nature*. 369:556-559.

592   Takeshima, H., S. Komazaki, M. Nishi, M. Iino, and K. Kangawa. 2000. Junctophilins: a novel
593   family of junctional membrane complex proteins. *Mol Cell*. 6:11-22.

594   Tanabe, T., K.G. Beam, J.A. Powell, and S. Numa. 1988. Restoration of excitation-contraction
595   coupling and slow calcium current in dysgenic muscle by dihydropyridine receptor
596   complementary DNA. *Nature*. 336:134-139.

597   Tani, H., M. Nishijima, H. Ushijima, T. Miyamura, and Y. Matsuura. 2001. Characterization of
598   cell-surface determinants important for baculovirus infection. *Virology*. 279:343-353.

599   Thorens, S., and M. Endo. 1975. Calcium-induced calcium release and "depolarization"-
600   induced calcium release: their physiological significance. *Proc Jpn Acad*. 51:473-478.

601   Tong, J., H. Oyamada, N. Demaurex, S. Grinstein, T.V. McCarthy, and D.H. MacLennan. 1997.
602   Caffeine and halothane sensitivity of intracellular Ca2+ release is altered by 15 calcium
603   release channel (ryanodine receptor) mutations associated with malignant hyperthermia
604   and/or central core disease. *J Biol Chem*. 272:26332-26339.

605   Uehara, A., T. Murayama, M. Yasukochi, M. Fill, M. Horie, T. Okamoto, Y. Matsuura, K.
606   Uehara, T. Fujimoto, T. Sakurai, and N. Kurebayashi. 2017. Extensive Ca2+ leak
607   through K4750Q cardiac ryanodine receptors caused by cytosolic and luminal Ca2+
608   hypersensitivity. *J Gen Physiol*. 149:199-218.

Ursu, D., R.P. Schuhmeier, M. Freichel, V. Flockerzi, and W. Melzer. 2004. Altered inactivation
of Ca2+ current and Ca2+ release in mouse muscle fibers deficient in the DHP receptor
gamma1 subunit. *J Gen Physiol*. 124:605-618.

Weiss, R.G., K.M. O'Connell, B.E. Flucher, P.D. Allen, M. Grabner, and R.T. Dirksen. 2004.
Functional analysis of the R1086H malignant hyperthermia mutation in the DHPR
reveals an unexpected influence of the III-IV loop on skeletal muscle EC coupling. *Am
J Physiol Cell Physiol*. 287:C1094-1102.

Wong King Yuen, S.M., M. Campiglio, C.C. Tung, B.E. Flucher, and F. Van Petegem. 2017.
Structural insights into binding of STAC proteins to voltage-gated calcium channels.
Proc Natl Acad Sci U S A. 114:E9520-E9528.

Yamazawa, T., T. Kobayashi, N. Kurebayashi, M. Konishi, S. Noguchi, T. Inoue, Y.U. Inoue, I.
Nishino, S. Mori, H. Iinuma, N. Manaka, H. Kagechika, A. Uryash, J. Adams, J.R.
Lopez, X. Liu, C. Diggle, P.D. Allen, S. Kakizawa, K. Ikeda, B. Lin, Y. Ikemi, K.
Nunomura, S. Nakagawa, T. Sakurai, and T. Murayama. 2021. A novel RyR1-selective
inhibitor prevents and rescues sudden death in mouse models of malignant hyperthermia
and heat stroke. *Nat Commun*. 12:4293.

Yang, T., J. Riehl, E. Esteve, K.I. Matthaei, S. Goth, P.D. Allen, I.N. Pessah, and J.R. Lopez.
2006. Pharmacologic and functional characterization of malignant hyperthermia in the
R163C RyR1 knock-in mouse. *Anesthesiology*. 105:1164-1175.

Zhao, Y., S. Araki, J. Wu, T. Teramoto, Y.F. Chang, M. Nakano, A.S. Abdelfattah, M. Fujiwara,
T. Ishihara, T. Nagai, and R.E. Campbell. 2011. An expanded palette of genetically
encoded Ca(2)(+) indicators. *Science*. 333:1888-1891.

Figure legends

Figure 1 **DICR measurements in 96-well plates using a fluorescence microplate reader. (A)**

Schematic drawing of RyR1/R-CEPIA1er HEK293 cells without (No BV, left) and with (5 BV,

right) baculovirus infection of essential components for DICR (Cav1.1, β1a, JP2, Stac3) and

Kir2.1. **(B)** Typical results of time-lapse R-CEPIA1er fluorescence measurement using a

FlexStation3 fluorescence microplate reader for RyR1/R-CEPIA1er HEK293 cells without (left,

No BV) or with (right, 5 BV) BV infection. High [K$^+$] solution ranging from 5–61 mM (symbols

shown in right) or 10 mM caffeine (light blue) was applied at 30 seconds after starting (blue

arrowheads). **(C)** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in No BV (black) and 5

BV cells (red). F/F$_0$ was obtained by normalizing the fluorescence (F) at 35 sec (open circles)

or the averaged fluorescence of 125-150 sec to that of the first 25 seconds (F$_0$). Data are means

$\pm$ SD (n = 6). Note that [K$^+$] dependent plot with data at 35 sec (open red) was shifted

rightward compared with that from data at 125–150 sec (filled red) with larger variations,

especially around EC$_{50}$ values. **(D)** Fluorescence change by 61 mM [K$^+$] or 10 mM caffeine.

Fluorescence change is expressed as 1-F/F$_0$ (see panel **C**). Data are means $\pm$ SD (n = 6) and

were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test. Note that 5

BV cells show large fluorescence change with high [K$^+$] which is nearly 80% of that with

caffeine.

Figure 2 **Validation of the reconstituted DICR platform. (A)** Typical results of R-CEPIA1er

fluorescence in cells infected with 5 BV (left), BV without Cav1.1 (center), and 5 BV plus γ1

(right). **(B)** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in 5 BV (black), -Cav1.1 (red),

and -JP2 (dark blue). Data are means ± SD (n = 9 for 5 BV, 6 for -Cav1.1 and -JP2). **(C)**

Fluorescence change by 61 mM [K$^+$] in the BV-infected cells lacking one of essential

components (-Cav1.1, -β1a, -Stac3, or -JP2) or with RyR2 cells instead of RyR1. Data are

means ± SD (n = 6) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test. **(D)** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells infected

with 5 BV (black), 5 BV plus α$_2$-δ (purple) and 5 BV plus γ1 (orange). Data are means ± SD (n

= 6). **(E)** Fluorescence change by 61 mM [K$^+$] in 5 BV, +α$_2$-δ, and +γ1 cells. Data are means ±

SD (n = 6) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 3 **Importance of Kir2.1 on DICR platform. (A)** Resting membrane potential of cells

infected with 5 BV (left) or BV without Kir2.1 (-Kir2.1) (right). Data are means ± SD (n = 10

and 16 for 5 mM and 50 mM [K$^+$], respectively, for 5 BV and 10 and 14 for 5 mM and 50 mM

[K$^+$], respectively, for -Kir2.1) and were analyzed by two-way ANOVA with Bonferroni's

multiple comparisons test. **(B)** Typical results of R-CEPIA1er fluorescence in cells infected

with 5 BV (left) and BV without Kir2.1 (right). High [K$^+$] solution ranging from 5–61 mM

(symbols shown in left) was applied at 30 seconds after starting (blue arrowheads). **(C)** Typical

results of ER [Ca^{2+}] measurement in cells with 5 BV (left), without Kir2.1, without Kir2.1 and

β1a, and without Kir2.1 and Stac3. **(D)** Resting ER [Ca^{2+}] in cells infected with 5 BV, BV

without Kir2.1 (-Kir2.1), Kir2.1 and β1a (-Kir2.1, -β1a), or Kir2.1 and Stac3 (-Kir2.1, -Stac3).

Data are means ± SD (n = 8) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test.

Figure 4 **Effect of MH mutations in Cav1.1 on DICR activity. (A)** Schematic drawing of the

location of MH mutations (R174W, R1086H and T1354S) in Cav1.1. **(B)** Resting ER [Ca^{2+}] in

cells expressing WT and mutant Cav1.1. Data are means ± SD (n = 12) and were analyzed by

one-way ANOVA with Dunnett's multiple comparisons test. **(C)** [K$^+$] dependence of R-

CEPIA1er fluorescence (F/F$_0$) in WT (black), R174W (red), R1086H (blue) and T1354S

(green) Cav1.1 cells. Data are means ± SD (n = 9). **(D, E)** Fluorescence change by 61 mM [K$^+$]

(D) and EC$_{50}$ values for [K$^+$] **(E)**. Data are means ± SD (n = 9) and were analyzed by one-way

ANOVA with Dunnett's multiple comparisons test. **(F)** Caffeine dependence of v(F/F$_0$) in WT

(black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means ± SD

(n = 9). **(G, H)** Fluorescence change by 20 mM caffeine **(G)** and EC$_{50}$ values for caffeine **(H)**.

Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test.

Figure 5 **Effect of myopathy mutations in Cav1.1 on DICR activity. (A)** Schematic drawing

of the location of myopathy mutations (E100K, F275L, P742Q and L1367V) in Cav1.1. **(B)**

Resting ER [Ca^{2+}] in cells expressing WT and mutant Cav1.1. Data are means ± SD (n = 12)

and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** [K$^+$]

dependence of R-CEPIA1er fluorescence (F/F$_0$) in WT (black), E100K (red), F275L (blue),

P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means ± SD (n = 9). **(D, E)**

Fluorescence change by 61 mM [K$^+$] **(D)** and EC$_{50}$ values for [K$^+$] **(E)**. Data are means ± SD

(n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(F)**

Caffeine dependence of R-CEPIA1er fluorescence (F/F$_0$) in WT (black), E100K (red), F275L

(blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means ± SD (n = 9). **(G,**

H) Fluorescence change by 20 mM caffeine (**G**) and EC_{50} values for caffeine (**H**). Data are

means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test.

Figure 6 **Effect of RyR1 inhibitors on DICR activity. (A)** Chemical structures of dantrolene,

Cpd1 and procaine. **(B)** $[K^+]$ dependence of fluorescence change (F/F_0) in cells without (Control,

black) or with dantrolene (10 μM, red), Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are

means ± SD (n = 9). **(C)** EC_{50} values for $[K^+]$. Data are means ± SD (n = 9) and were analyzed

by one-way ANOVA with Dunnett's multiple comparisons test. **(D)** Caffeine dependence of R-

CEPIA1er fluorescence (F/F_0) in cells without (Control, black) or with dantrolene (10 μM, red),

Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are means ± SD (n = 9). **(E)** EC_{50} values

for caffeine. Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with

Dunnett's multiple comparisons test.

Figure 7 **Effect of lyotropic anions on DICR activity. (A)** $[K^+]$ dependence of fluorescence

change (F/F_0) in cells in the absence (Control, black) or presence of perchlorate (10 mM, red)

or thiocyanate (10 mM, blue). Data are means ± SD (n = 9). **(B)** EC_{50} values for $[K^+]$. Data are

means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test. **(C)** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in cells in the

absence (Control, black) or presence of perchlorate (10 mM, red) or thiocyanate (10 mM, blue).

Data are means ± SD (n = 9). **(D)** EC_{50} values for caffeine. Data are means ± SD (n = 9) and

were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Table 1. List of PCR primers.

mCav1.1-F, AAAAAGCTTGCCACCATGGAGCCGCCCTCACCCCAG

mCav1.1-R, TTTACGCGTTCAAGGCTCCCAGGGCACGGC

mβ1a-F, AAAGCTAGCCACCATGGTCCAGAAGAGCGGCATG

mβ1a-R, TTTGATATCCTACATGGCGTGCTCCTGAGG

mJP2-F, AAAGCTAGCCACCATGAGCGGGGGCCGCTTTGAC

mJP2-R, TTTGATATCTCAAGTCAGGAGGTGAACAAA

mStac3-F, AAAGCTAGCCACCATGACAGAAAAGGAAGTGGTG

mStac3-R, TTTACGCGTTTAAATCTCCTCCAGGAAGTC

mKir2.1-F, AAAGCTAGCCACCATGGGCAGTGTGAGAACCAAC

mKir2.1-R, TTTACGCGTTCATATCTCCGATTCTCGCCT

rα2δ-F, AAAGCTAGCCACCATGGCTGCTGGCTGCCTGCTG

rα2δ-R, TTTACGCGTTCACCATAGATAGTGTCTGCT

mγ1-F, AAAGCTAGCCACCATGTCACAGACCAAAACAGCG

mγ1-R, TTTACGCGTCTAGTGCTCTGGCTCAGCGTC

Figure 1

Figure 1 **DICR measurements in 96-well plates using a fluorescence microplate reader. (A)** Schematic drawing of RyR1/R-CEPIA1er HEK293 cells without (No BV, left) and with (5 BV, right) baculovirus infection of essential components for DICR (Cav1.1, β1a, JP2, Stac3) and Kir2.1. **(B)** Typical results of time-lapse R-CEPIA1er fluorescence measurement using a FlexStation3 fluorescence microplate reader for RyR1/R-CEPIA1er HEK293 cells without (left, No BV) or with (right, 5 BV) BV infection. High [K$^+$] solution ranging from 5–61 mM (symbols shown in right) or 10 mM caffeine (light blue) was applied at 30 seconds after starting (blue arrowheads). **(C)** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in No BV (black) and 5 BV cells (red). F/F$_0$ was obtained by normalizing the fluorescence (F) at 35 sec (open circles) or the averaged fluorescence of 125-150 sec to that of the first 25 seconds (F$_0$). Data are means $\pm$ SD (n = 6). Note that [K$^+$] dependent plot with data at 35 sec (open red) was shifted rightward compared with that from data at 125–150 sec (filled red) with larger variations, especially around EC$_{50}$ values. **(D)** Fluorescence change by 61 mM [K$^+$] or 10 mM caffeine. Fluorescence change is expressed as 1-F/F$_0$ (see panel **C**). Data are means $\pm$ SD (n = 6) and were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test. Note that 5 BV cells show large fluorescence change with high [K$^+$] which is nearly 80% of that with caffeine.

Figure 2

Figure 2 **Validation of the reconstituted DICR platform. (A)** Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV (left), BV without Cav1.1 (center), and 5 BV plus γ1 (right). **(B)** [K^+] dependence of R-CEPIA1er fluorescence (F/F_0) in 5 BV (black), -Cav1.1 (red), and -JP2 (dark blue). Data are means $\pm$ SD (n = 9 for 5 BV, 6 for -Cav1.1 and -JP2). **(C)** Fluorescence change by 61 mM [K^+] in the BV-infected cells lacking one of essential components (-Cav1.1, -β1a, -Stac3, or -JP2) or with RyR2 cells instead of RyR1. Data are means $\pm$ SD (n = 6) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(D)** [K^+] dependence of R-CEPIA1er fluorescence (F/F_0) in cells infected with 5 BV (black), 5 BV plus α2-δ (+α2-δ, purple) and 5 BV plus γ1 (+γ1, orange). Data are means $\pm$ SD (n = 6). **(E)** Fluorescence change by 61 mM [K^+] in 5 BV, +α2-δ, and +γ1 cells. Data are means $\pm$ SD (n = 6) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 3

Figure 3 **Importance of Kir2.1 on DICR platform. (A)** Membrane potential of cells infected with 5 BV (left) or BV without Kir2.1 (-Kir2.1) (right). Data are means $\pm$ SD (n = 10 and 16 for 5 mM and 50 mM K$^+$, respectively, for 5 BV and 10 and 14 for 5 mM and 50 mM K$^+$, respectively, for -Kir2.1) and were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test. **(B)** Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV (left) and BV without Kir2.1 (right). High [K$^+$] solution ranging from 5–61 mM (symbols shown in left) was applied at 30 seconds after starting (blue arrowheads). **(C)** Typical results of ER [Ca^{2+}] measurement in cells with 5 BV, without Kir2.1, without Kir2.1 and β1a, and without Kir2.1 and Stac3. **(D)** Resting ER [Ca^{2+}] in cells infected with 5 BV, BV without Kir2.1 (-Kir2.1), Kir2.1 and β1a (-Kir2.1, -β1a), or Kir2.1 and Stac3 (-Kir2.1, -Stac3). Data are means $\pm$ SD (n = 8) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 4

Figure 4 **Effect of MH mutations in Cav1.1 on DICR activity. (A)** Schematic drawing of the location of MH mutations (R174W, R1086H and T1354S) in Cav1.1. **(B)** Resting ER $[Ca^{2+}]$ in cells expressing WT and mutant Cav1.1. Data are means $\pm$ SD (n = 12) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** $[K^+]$ dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means $\pm$ SD (n = 9). **(D, E)** Fluorescence change by 61 mM K^+ **(D)** and EC_{50} values for K^+ **(E)**. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(F)** Caffeine dependence of v(F/F_0) in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means $\pm$ SD (n = 9). **(G, H)** Fluorescence change by 20 mM caffeine **(G)** and EC_{50} values for caffeine **(H)**. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 5

Figure 5 **Effect of myopathy mutations in Cav1.1 on DICR activity. (A)** Schematic drawing of the location of myopathy mutations (E100K, F275L, P742Q and L1367V) in Cav1.1. **(B)** Resting ER [Ca^{2+}] in cells expressing WT and mutant Cav1.1. Data are means $\pm$ SD (n = 12) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** [K^+] dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means $\pm$ SD (n = 9). **(D, E)** Fluorescence change by 61 mM K^+ **(D)** and EC_{50} values for K^+ **(E)**. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(F)** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means $\pm$ SD (n = 9). **(G, H)** Fluorescence change by 20 mM caffeine **(G)** and EC_{50} values for caffeine **(H)**. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 6

Figure 6 **Effect of RyR1 inhibitors on DICR activity. (A)** Chemical structures of dantrolene, Cpd1 and procaine. **(B)** [K$^+$] dependence of fluorescence change (F/F$_0$) in cells without (Control, black) or with dantrolene (10 μM, red), Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are means $\pm$ SD (n = 9). **(C)** EC$_{50}$ values for [K$^+$]. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(D)** Caffeine dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells without (Control, black) or with dantrolene (10 μM, red), Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are means $\pm$ SD (n = 9). **(E)** EC$_{50}$ values for caffeine. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 7

Figure 7 **Effect of lyotropic anions on DICR activity. (A)** [K$^+$] dependence of fluorescence change (F/F$_0$) in cells in the absence (Control, black) or presence of perchlorate (10 mM, red) or thiocyanate (10 mM, blue). Data are means $\pm$ SD (n = 9). **(B)** EC$_{50}$ values for K$^+$. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** Caffeine dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells in the absence (Control, black) or presence of perchlorate (10 mM, red) or thiocyanate (10 mM, blue). Data are means $\pm$ SD (n = 9). **(D)** EC$_{50}$ values for caffeine. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.