Reconstituted depolarization-induced Ca^{2+} release platform for skeletal muscle disease

mutation validation and drug discovery

Takashi Murayama[1*], Nagomi Kurebayashi[1], Takuro Numaga-Tomita[2], Takuya Kobayashi[1],

Satoru Okazaki[1], Kyosuke Yamashiro[1], Tsutomu Nakada[2], Shuichi Mori[3], Ryosuke Ishida[3],

Hiroyuki Kagechika[3], Mitsuhiko Yamada[2], Takashi Sakurai[1]

[1]Department of Pharmacology, Juntendo University School of Medicine, Tokyo, Japan.

[2]Department of Molecular Pharmacology, Shinshu University School of Medicine,

Matsumoto, Japan.

[3]Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, Tokyo,

Japan.

Short running title: Development of reconstituted DICR platform

[*]Corresponding author: Takashi Murayama, Ph.D., Department of Pharmacology, Juntendo

University School of Medicine, Tokyo 113-8421, Japan. E-mail: takashim@juntendo.ac.jp

Summary

Mutations in essential components for depolarization-induced Ca^{2+} release (DICR) are

implicated into various skeletal muscle diseases. Murayama et al. establish a reconstituted

DICR platform in nonmuscle cells for evaluation of disease-causing mutations and drug

discovery.

Abstract

In skeletal muscle excitation-contraction (E-C) coupling, depolarization of the plasma

membrane triggers Ca^{2+} release from the sarcoplasmic reticulum (SR), referred to as

depolarization-induced Ca^{2+} release (DICR). DICR occurs via the type 1 ryanodine receptor

(RyR1), which physically interacts with the dihydropyridine receptor Cav1.1 subunit in

specific machinery formed with additional essential components including β1a, Stac3 adaptor

protein and junctophilins. Exome sequencing has accelerated the discovery of many novel

mutations in genes encoding DICR machinery in various skeletal muscle diseases. However,

functional validation is time-consuming because it must be performed in a skeletal muscle

environment. In this study, we established a platform of the reconstituted DICR in HEK293

cells. The essential components were effectively transduced into HEK293 cells expressing

RyR1 using baculovirus vectors, and Ca^{2+} release was quantitatively measured with

R-CEPIA1er, a fluorescent ER Ca^{2+} indicator, without contaminant of extracellular Ca^{2+}

influx. In these cells, [K^+]-dependent Ca^{2+} release was triggered by chemical depolarization

with the aid of inward rectifying potassium channel, indicating a successful reconstitution of

DICR. Using the platform, we evaluated several Cav1.1 mutations that are implicated in

malignant hyperthermia and myopathy. We also tested several RyR1 inhibitors; whereas

dantrolene and Cpd1 inhibited DICR, procaine had no effect. Furthermore, twitch potentiators

such as perchlorate and thiocyanate shifted the voltage dependence of DICR to more negative

potentials without affecting Ca^{2+}-induced Ca^{2+} release. These results well reproduced the

findings with the muscle fibers and the cultured myotubes. Since the procedure is simple and

reproducible, the reconstituted DICR platform will be highly useful for validation of

mutations and drug discovery for skeletal muscle diseases.

Introduction

In vertebrate skeletal muscle, depolarization of the plasma membrane triggers Ca^{2+} release from the sarcoplasmic reticulum (SR), which is commonly referred to as excitation-contraction (E-C) coupling. This process occurs at triad junctions in which a transverse (T) tubule is closely apposed to terminal cisterna of the SR (Franzini-Armstrong, 2018). In excitation-contraction coupling, the type 1 ryanodine receptor (RyR1), a Ca^{2+} release channel in the SR, is activated by a conformational change of the voltage sensor in the Cav1.1 subunit of the dihydropyridine receptor (DHPR) upon depolarization of the T-tubule membrane. This is referred to as depolarization-induced Ca^{2+} release (DICR) (Rios and Pizarro, 1991; Schneider, 1994). Studies with knockout animals have revealed that functional and structural interactions between Cav1.1 and RyR1 in muscle require at least three additional components, a β1a auxiliary subunit of DHPR (Gregg et al., 1996; Schredelseker et al., 2005), the Stac3 adaptor protein (Horstick et al., 2013; Nelson et al., 2013; Reinholt et al., 2013), and junctophilins (Takeshima et al., 2000; Yang et al., 2022), which may form specific machinery for DICR (Flucher and Campiglio, 2019; Shishmarev, 2020; Rufenach and Van Petegem, 2021).

Mutations in genes that encode DICR machinery components have been implicated in various types of muscle diseases, including congenital myopathies, periodic paralysis, and malignant hyperthermia (MH) (Agrawal et al., 2018; Flucher, 2020; Lawal et al., 2020). Rapid exome sequencing has accelerated the discovery of novel mutations; however, validation of these mutations by functional testing is critically important for the diagnosis and treatment of these diseases. This has been performed using myotubes from mice that lack DICR machinery components, for example knockout mice of Cav1.1 (Tanabe et al., 1988), β1a (Gregg et al., 1996) and Stac3 (Nelson et al., 2013). While this approach has evaluated several disease-causing mutations in the DICR components (Weiss et al., 2004; Pirone et al., 2010; Eltit et al., 2012; Polster et al., 2016; Perez et al., 2018), it has certain limitations. For example, it is expensive and labor intensive to maintain the mouse lines. Differentiation of cells into myotubes is time consuming and is affected by various factors, such as the cell line used and experimental conditions. Furthermore, transduction of foreign genes into myotubes is often inefficient.

There are no specific treatments for most diseases related to mutations in the DICR machinery. A major cause for the slow pace of novel drug development is the lack of an

appropriate drug screening platform for these diseases. We have recently established an

efficient high-throughput platform for screening RyR1-targeted drugs using endoplasmic

reticulum (ER) Ca^{2+} measurements in HEK293 cells (Murayama et al., 2018; Murayama and

Kurebayashi, 2019) and using this platform we developed Compound 1 (Cpd1), a novel RyR1

inhibitor for treatment of MH (Mori et al., 2019; Yamazawa et al., 2021). Therefore,

functional screening using HEK293 cells is a promising platform for drug discovery.

Recently, Perni et al. successfully reconstituted DICR by co-expressing five essential

components, RyR1, Cav1.1, β1a, junctophilin-2 (JP2) and Stac3, in non-muscle tsA201 cells

(Perni et al., 2017). They demonstrated formation of ER-plasma membrane junctions in which

Cav1.1 is arranged into tetrads, which is indicative of physical links to RyR1 (Block et al.,

1988). Depolarization of the plasma membrane using the patch-clamp technique induced a

rapid Ca^{2+} transient without influx of extracellular Ca^{2+}. This provides a basis for developing

reconstituted DICR.

In this study, we established a platform for DICR that was reconstituted in HEK293

cells. Baculovirus (BV) infection of essential components greatly improved transduction

efficiency. Chemical depolarization by high concentration potassium $[K^+]$ solutions with the

aid of an inward-rectified potassium channel (Kir2.1) enabled simultaneous stimulation of

many cells. Furthermore, a genetically-encoded ER Ca^{2+} indicator quantitatively measured

DICR without influence of extracellular Ca^{2+} influx. Using this platform, we successfully

evaluated disease-causing mutations in Cav1.1 and tested several RyR1 inhibitors and DICR

modulators. Our reconstituted DICR platform will accelerate the validation of mutations and

drug discovery for skeletal muscle diseases.

Materials and Methods

Generation of stable inducible HEK293 cell lines

HEK293 cells stably expressing R-CEPIA1er and RyR1 were generated as described

previously (Murayama et al., 2018; Murayama and Kurebayashi, 2019). Briefly, a full-length

rabbit RyR1 cDNA was cloned in a tetracycline-induced expression vector

(pcDNA5/FRT/TO; Life Technologies, CA, USA) (Tong et al., 1997). Flp-In T-REx HEK293

cells were co-transfected with this expression vector and pOG44 in accordance with the

manufacturer's instructions. Clones with suitable doxycycline-induced expression of RyR1

were selected and used for experiments. The cells were then transfected with an R-CEPIA1er

cDNA (a generous gift from Masamitsu Iino, The University of Tokyo) for stable expression

of R-CEPIA1er. Cells were cultured in Dulbecco's modified Eagle's medium supplemented

with 10% fetal calf serum, 2 mM L-glutamine, 15 µg/ml blasticidin, and 100 µg/ml

hygromycin.

BV Production

Baculovirus (BV) that expresses vesicular stomatitis virus G-protein (VSVG) on the virus

envelope (Tani et al., 2001) was produced using the Bac-to-Bac system (LifeTechnologies) as

described previously (Uehara et al., 2017). cDNAs for Cav1.1, β1a, JP2, Stac3, and Kir2.1

were cloned from mouse skeletal muscle using reverse transcription-PCR (primer pairs are

listed in **Table 1**). Ca^{2+} non-conducting mutation (N617D) (Schredelseker et al., 2010) and

disease-causing mutations were introduced in Cav1.1 cDNA by inverse PCR (primer pairs are

listed in **Table 1**). Each cDNA was cloned into the modified donor vector

(pFastBac1-VSVG-CMV-WPRE). BV was produced in Sf9 cells according to the

manufacturer's instructions (LifeTechnologies). BV titers were immunologically determined

using an anti-gp64 antibody (AcV5, Santa Cruz Biotechnology)(Kitts and Green, 1999).

Titers were $1–2 \times 10^8$ pfu/mL for all viruses. The BV solutions were stored in the dark at 4°C

and used within one month after production.

Immunocytochemistry

RyR1-HEK293 cells were seeded on a 35 mm glass bottom dish that had been coated with

collagen solution at a density of 2×10^5 cells/well in 1.5 mL culture media. One day after

seeding, culture media were exchanged to those containing 2 µg/mL doxycycline (for

induction of RyR1) and BV solutions for Cav1.1 (N617D), β1a, JP2, Stac3, and Kir2.1 (30 µL

each in 1.5 mL). Multiplicity of infection (MOI) was 7.5-15 for each BV (depending on the

virus titer). After 24 h, cells were fixed with ice-cold methanol and rinsed with

phosphate-buffered saline (PBS). The cells were blocked with 5% BSA/PBS for 30 min and

incubated overnight with primary antibodies against Cav1.1 (sc-514685, Santa Cruz

Biotechnology), β1a (MA5-27714, Invitrogen), JP2 (sc-377086, Santa Cruz Biotechnology),

Stac3 (1G8, Abnova), or Kir2.1 (S112B-14, Invitrogen) in 1% BSA/PBS. After washing three

times with PBS, the cells were incubated for 2 h with Alexa594-labeled goat anti-mouse IgG

(Invitrogen) in 1% BSA/PBS. After washing with PBS, the cells were observed using an

all-in-one fluorescence microscope (BZ-X710, Keyence, Osaka, Japan).

Single-cell Ca^{2+} imaging

RyR1/R-CEPPIA1er HEK293 cells were seeded on a 35 mm glass bottom dish and culture media were exchanged next day as described above. 24 h after infection, single-cell Ca^{2+} imaging was performed as described previously (Murayama et al., 2015; Uehara et al., 2017). Solutions were perfused using an in-line solution heater/cooler (Warner Instruments, Holliston, MA, USA). Measurements were performed at 26°C instead of 37°C, since cells tended to detach from the dish during perfusion at 37°C. ER [Ca^{2+}] signals were determined by R-CEPIA1er fluorescence excited at 561 nm using a 20× objective lens. Cells were perfused with normal Krebs solution (140 mM NaCl, 5 mM KCl, 2 mM CaCl$_2$, 1 mM MgCl$_2$, 11 mM glucose, 10 mM HEPES, pH 7.4) for 30 s, high [K$^+$] Krebs solution containing 50 mM [K$^+$] for 50 s, and normal Krebs solution for 1.5 min. Fluorescence intensities in individual cells were determined by region of interest (ROI) analysis using AquaCosmos software (Hamamatsu Photonics, Hamamatsu, Japan).

Western blotting

RyR1-HEK293 cells were seeded on 12-well plate at a density of 3×10^5 cells/well in 1 mL culture media. One day after seeding, 1 mL culture media containing 2 µg/mL doxycycline (for induction of RyR1) and BV solutions for Cav1.1 (N617D), β1a, JP2, Stac3, and Kir2.1 (20 µL each) were added to each well. After 24 h, cells were harvested, rinsed with PBS and lysed with 50 µL of Pro-Prep protein extraction solution (iNtRON Biotechnology). After centrifugation at 15,000 rpm for 5 min, the extracted proteins were separated on 3–15% polyacrylamide gels and transferred to PVDF membranes. Membranes were probed with the primary antibodies described above and calnexin (C4731, Sigma-Aldrich, MO, USA) as a loading control, followed by HRP-labeled anti-mouse IgG (04-18-18, KPL) or anti-rabbit IgG (074-1516, KPL). Positive bands were detected by chemiluminescence using ImmunoStar LD (Fujifilm Wako Chemicals) as a substrate.

Time lapse ER [Ca^{2+}] Measurements using a fluorometer

Time lapse ER [Ca^{2+}] measurements were performed using the FlexStation3 fluorometer (Molecular Devices, San Jose, CA) (Murayama et al., 2018; Murayama and Kurebayashi, 2019). Stable RyR1/R-CEPIA1er HEK293 cells were seeded on 96-well flat-, clear-bottom black microplates (#3603, Corning, New York, NY) at a density of 3×10^4 cells/well in 100 µL culture media. One day after seeding, 100 µL culture media containing 2 µg/mL doxycycline

(for induction of RyR1) and BV solutions for Cav1.1 (N617D), β1a, JP2, Stac3, and Kir2.1 (2

μL each) were added to each well. Multiplicity of infection (MOI) was 7.5-15 for each BV

(depending on the virus titer). After 24 h, the culture media were replaced with 81 μL normal

Krebs solution, and the microplate was placed in a FlexStation3 fluorometer preincubated at

37°C. Signals from R-CEPIA1er, which was excited at 560 nm and emitted at 610 nm, were

captured every 5 sec for 150 sec. Thirty sec after starting, 54 μL of the high $[K^+]$ or caffeine

solution was applied to the cells. The fluorescence change was expressed as F/F_0 in which

averaged fluorescence intensity of the last 25 sec (F) was normalized to that of the initial 25

sec (F_0). For drug testing, compounds were added to normal Krebs solution at the indicated

concentrations.

For determination of resting ER $[Ca^{2+}]$, cells were incubated with 81 μL normal

Krebs solution, and 54 μL Krebs solution containing 50 mM Ca^{2+} and 50 μM ionomycin was

added 30 sec after starting (final concentrations of Ca^{2+} and ionomycin were 20 mM and 20

μM, respectively). R-CEPIA1er fluorescence was captured every 10 sec for 6 min.

Measurement of resting membrane potential

The resting membrane potential of BV-infected RyR1-HEK293 cells with or without Kir2.1

expression was measured under current-clamp conditions in the whole cell configuration of

the patch-clamp method with an Axopatch 200B amplifier (Molecular Devices, San Jose, CA).

Pipettes were fabricated from borosilicate glass capillaries (Kimble) with a vertical pipette

puller (PP-83, Narishige, Tokyo, Japan). Pipettes had series resistance of 3–4 MΩ when filled

with the pipette solution containing (in mM) 120 D-glutamate, 20 KCl, 10 NaCl, 0.1 EGTA, 2

$MgCl_2$, 3 MgATP, and 5 HEPES (pH 7.4 adjusted with KOH). The series resistance was

routinely compensated by ~75% with the amplifier. Bath solution contained (in mM): 140

NaCl, 5 KCl, 11 glucose, 2 $CaCl_2$, 1 $MgCl_2$, and 10 HEPES (pH 7.4 adjusted with NaOH).

High $[K^+]$ bath solution contained (in mM): 95 NaCl, 50 KCl, 11 glucose, 2 $CaCl_2$, 1 $MgCl_2$,

and 10 HEPES (pH 7.4 adjusted with NaOH). The resting membrane potential was measured

immediately after breaking into cells.

Statistics

Data are presented as means ± SD. Unpaired Student's t test was used for comparisons

between two groups. One-way analysis of variance (ANOVA), followed by Dunnett's test,

was performed to compare three or more groups. Two-tailed tests were used for all analyses.

Statistical analysis was performed using Prism v9 (GraphPad Software, Inc., La Jolla, USA).

Results

Establishment of a reconstituted DICR platform using a fluorescence microplate reader

To reconstitute DICR, we used HEK293 cells stably expressing RyR1 and

R-CEPIA1er, a genetically-encoded fluorescent ER Ca^{2+} indicator (Suzuki et al., 2014), that

were generated for high-throughput screening of novel RyR1 inhibitors (Murayama et al.,

2018; Murayama and Kurebayashi, 2019) (**Fig. 1 A**, left). The other components for DICR,

Cav1.1, β1a, Stac3 and JP2, which were used by Perni et al. (Perni et al., 2017), were

transduced using VSV-G pseudotyped baculovirus (BV), which efficiently infects mammalian

cells (Tani et al., 2001; Uehara et al., 2017) (**Fig. 1 A**, right). In these cells, DICR can be

detected by a decrease in ER Ca^{2+}. To prevent extracellular Ca^{2+} influx through Cav1.1 by

depolarization, a N617D mutation was introduced that abolishes Ca^{2+} conductance

(Schredelseker et al., 2010). In addition, we expressed Kir2.1, an inward-rectified potassium

channel in skeletal muscle (DiFranco et al., 2015), to hyperpolarize the membrane potential

because the resting membrane potential in HEK293 cells is much less (−20 mV) than in

skeletal muscle (−90 mV) (Kirkton and Bursac, 2011). Imunocytochemistry revealed that the

five proteins (Cav1.1, β1a, Stac3, JP2, and Kir2.1) were successfully expressed in almost all

cells with infection of 5 BV, which we term hereafter 5 BV cells, but not in non-infected cells

(No BV cells) (**Fig. S1 A**). BV-dependent protein expression was also confirmed by Western

blotting of cell lysate (**Fig. S1 B**). Therefore, BV infection effectively transduced the essential

components of DICR in RyR1/R-CEPIA1er-HEK293 cells.

To detect DICR, we used a fluorescence microplate reader (FlexStation3), which

measures time-lapse R-CEPIA1er fluorescence in 96-well plates. Fluorescence signals were

collected in a 1.5 mm diameter area; therefore, an averaged signal from thousands of cells

was detected in each well. We adopted chemical depolarization using high $[K^+]$ solutions,

which can simultaneously depolarize all the cells to trigger DICR. The cells were initially

incubated with normal Krebs solution containing 5 mM $[K^+]$. Then, solutions containing

various $[K^+]$ concentrations were added to give $[K^+]$ concentrations of 5–61 mM. In cells

without BV infection, no apparent changes in fluorescence were observed with any $[K^+]$

concentration (**Fig. 1 B**, left). Caffeine (10 mM) reduced the R-CEPIA1er fluorescence,

confirming the functional expression of RyR1. In 5 BV cells, high $[K^+]$ solution rapidly

reduced the R-CEPIA1er fluorescence in a $[K^+]$-dependent manner (**Fig. 1 B**, right). To

monitor ER Ca^{2+} changes in individual cells, we also observed R-CEPIA1er fluorescence

using laser scanning confocal microscopy. In 5 BV cells, high $[K^+]$ solution caused a large

and transient reduction in the R-CEPIA1er fluorescence in >90% of cells (**Movie S1** and **Fig.**

S2).

For quantitative analysis, we normalized the averaged fluorescence intensity to that

of the initial 25 sec (F_0) to correct variations in the initial fluorescence intensity between wells.

We first used rapid fluorescence change 5 sec after stimulation (i.e., 35 sec), which may

reflect the rate of Ca^{2+} release. The F/F_0 plot as a function of $[K^+]$ concentrations

demonstrated $[K^+]$ dependence with an EC_{50} of 17.0 ± 0.4 mM. However, variations were

relatively large, especially near EC_{50} value because of rapid changes in fluorescence intensity

(**Fig. 1 C**). We instead adopted the average fluorescence intensity for the last 25 sec (125-150

sec). This gave an EC_{50} of 9.8 ± 0.1 mM $[K^+]$, which was lower than that obtained from 35

sec, but with smaller variations (**Fig. 1 C**). The fluorescence change by 61 mM $[K^+]$, which is

expressed as $1-F/F_0$ (**Fig. 1 C**), reached ~80% of that induced by caffeine in 5 BV cells,

indicating a high degree of DICR reconstitution (**Fig. 1 D**). Taken together, these results

indicate successful establishment of a reconstituted DICR platform.

The original method for DICR reconstitution monitors cytoplasmic $[Ca^{2+}]$ (Perni et

al., 2017); therefore, we also assessed DICR by measuring cytoplasmic $[Ca^{2+}]$ using

G-GECO1.1, a genetically-encoded Ca^{2+} indicator (Zhao et al., 2011). In cells without BV

infection, a substantial increase in $[Ca^{2+}]$ was observed by high $[K^+]$, indicating the existence

of endogenous Ca^{2+} influx pathways by depolarization (**Fig. S3 A**). In 5 BV cells, DICR was

detected by rapid and large increases in $[Ca^{2+}]$, which declined with time (**Fig. S3 B**). The

rapid fluorescence change 5 sec after stimuli (i.e., 35 sec) provided dose-dependent curve

with EC_{50} of 16.4 ± 0.5 mM, which is similar to the value determined by rapid fluorescence

change in ER $[Ca^{2+}]$ measurement (17 mM, see **Fig. 1 C**) and those obtained from mouse

skeletal muscle myotubes (~21 mM) in high $[K^+]$-induced Ca^{2+} transients (Yang et al., 2006;

Lopez et al., 2018).

Properties of the reconstituted DICR

We next examined properties of the reconstituted DICR in HEK293 cells. Initially,

we examined the role of the essential components. Removal of Cav1.1 resulted in complete

loss of high $[K^+]$-induced Ca^{2+} release (**Fig. 2 A-C**). Ca^{2+} release was also lost in cells lacking

β1a or Stac3, and in cells expressing RyR2 instead of RyR1 (**Fig. 2 C**). These results are

consistent with previous findings of mice lacking each component that they are essential for

DICR (Tanabe et al., 1988; Takeshima et al., 1994; Gregg et al., 1996; Nelson et al., 2013).

Interestingly, DICR still occurred without JP2, although the fluorescence change by 61 mM

$[K^+]$ was significantly reduced and $[K^+]$ dependence was largely shifted rightward (**Fig. 2 B,**

C).

Skeletal muscle Cav1.1 forms a complex with three auxiliary subunits, β1a, $α_2$-δ and $γ_1$

(Catterall, 2000). $α_2$-δ and $γ_1$ may play modulatory roles (Flucher et al., 2005). Additional

expression of $α_2$-δ had no effect on high $[K^+]$-induced Ca^{2+} release (**Fig. 2 D, E**). Expression

of γ1 did not significantly affect high $[K^+]$-induced Ca^{2+} release at 20 mM or lower $[K^+]$ but it

reduced the fluorescence change at higher $[K^+]$ concentrations (**Fig. 2 D, E**). In $γ_1$ cells,

R-CEPIA1er fluorescence decreased but increased again with time, suggesting inactivation of

the Ca^{2+} release (**Fig. 2 A**). These findings are consistent with previous reports of the effects

of $α_2$-δ (Obermair et al., 2005) and $γ_1$ (Ursu et al., 2004) on DICR. Therefore, our

reconstituted platform can reproduce DICR in HEK293 cells and is valid for evaluating

disease-causing mutations or drugs.

We also tested the importance of Kir2.1 on the reconstituted DICR platform. We

measured membrane potential of cells under current-clamp conditions using pipette solution

containing ~145 mM $[K^+]$ (**Fig. 3 A**). The membrane potential of 5 BV-infected cells was -71

± 3 mV in normal Krebs solution and became depolarized to -14 ± 3 mV by high $[K^+]$

solution containing 50 mM $[K^+]$. In contrast, membrane potential was already depolarized in

the cells without Kir2.1 (-18 ± 2 mV) in normal Krebs solution, which is consistent with the

previous report(Kirkton and Bursac, 2011). Without Kir2.1, R-CEPIA1er fluorescence

intensity was substantially reduced and high $[K^+]$-induced Ca^{2+} release was completely

abolished (**Fig. 3 B**). We quantified resting ER $[Ca^{2+}]$ by determining the maximum

fluorescence intensity of R-CEPIA1er using ionomycin/Ca^{2+} (**Fig. 3 C**). The resting ER

$[Ca^{2+}]$ was severely reduced in cells without Kir2.1 (**Fig. 3 D**). These results suggest that

persistently activated DICR activity under the depolarized conditions causes ER Ca^{2+}

depletion. In support of this, further removal of β1a or Stac3, which completely abolishes

DICR activity, restored the ER $[Ca^{2+}]$ (**Fig. 3 D**). Therefore, Kir2.1 is indispensable for our

reconstituted DICR platform.

Evaluation of disease-causing mutations in Cav1.1

Using the reconstituted DICR platform, we evaluated mutations in Cav1.1 which are

implicated in various muscle diseases (Beam et al., 2017; Flucher, 2020). We initially tested

known autosomal dominant MH mutations, R174W (Carpenter et al., 2009), R1086H

(Stewart et al., 2001), and T1354S (Pirone et al., 2010) (**Fig. 4 A**). Ca^{2+} non-conducting

N617D mutation was also introduced in wild type (WT) and the mutant Cav1.1s. To observe

the effect of mutations clearly, we tested them in homozygous state without WT. The

expression of mutant Cav1.1s was confirmed by Western blot (**Fig. S1 C**). We found that ER

$[Ca^{2+}]$ was severely reduced for R1086H (**Fig. 4 B**). In support of this, R1086H exhibited

lower initial R-CEPIA1er fluorescence intensity and smaller fluorescence change in response

to high $[K^+]$ than those in WT or the other mutants (**Fig. 4 C**). We plotted $[K^+]$ dependence of

DICR using the F/F_0 corrected by ER $[Ca^{2+}]$ (**Fig. 4 D**). This clearly indicates changes in ER

$[Ca^{2+}]$ in each WT or mutant cell. The change in the corrected F/F_0 was substantially smaller

in R1086H (**Fig. 4 E**), being consistent with initial depletion of ER $[Ca^{2+}]$. Interestingly,

R1086H exhibited significantly lower EC_{50} value for $[K^+]$ (8.3 ± 0.3 mM) compared to WT

(10.8 ± 0.7 mM), indicating an enhanced $[K^+]$-dependence (**Fig. 4 F**). No or only minor

changes in ER $[Ca^{2+}]$ and $[K^+]$ dependence were observed for R174W or T1354S cells (**Fig. 4**

B–F). We anticipated that the reduction in ER $[Ca^{2+}]$ in R1086H cells was caused by

activation of Cav1.1 at resting membrane potential because of its hypersensitivity. To test the

hypothesis, we hyperpolarized the membrane potential which reduced the activation of

Cav1.1. Reduction of $[K^+]$ in the Krebs solution from 5 mM to 2.5 mM successfully

hyperpolarized the membrane potential by ~15 mV (**Fig. S4 A**). This substantially increased

ER $[Ca^{2+}]$ in R1086H cells to a level comparable to that in WT cells (**Fig. S4 B**). R1086H

cells exhibited a clear greater sensitivity to $[K^+]$ (5.9 ± 0.2 mM) than WT cells (8.3 ± 0.5 mM)

(**Fig. S4 C–E**), supporting our hypothesis of R1086H hyperactivation.

323   Previous functional characterization of these Cav1.1 mutants consistently showed

enhanced caffeine sensitivity, i.e., acceleration of Ca^{2+}-induced Ca^{2+} release (CICR) (Weiss et

al., 2004; Pirone et al., 2010; Eltit et al., 2012). We tested caffeine dependence of these

mutants in our platform. Caffeine released Ca^{2+} from WT cells in a dose-dependent manner

(**Fig. 4 G**). R1086H cells exhibited a reduced fluorescence change by 20 mM caffeine (**Fig. 4**

H) and a significantly smaller EC_{50} (**Fig. 4 I**), indicating an enhanced caffeine sensitivity.

Caffeine dependence was not changed in R174W or T1354S cells (**Fig. 4 G–I**).

330   We next examined the effect of myopathy-related mutations in Cav1.1 on DICR

activity. We chose four mutations, E100K, F275L, P742Q, and L1367V (Schartner et al.,

2017), none of which have been functionally characterized (**Fig. 5 A**). Ca^{2+} non-conducting

N617D mutation was also introduced in the mutant Cav1.1s. and we tested these mutations in

homozygous states without WT. The expression of mutant Cav1.1s was confirmed by Western

blot (**Fig. S1 C**). ER $[Ca^{2+}]$ was not changed by any of these mutations (**Fig. 5 B**). DICR

activity was completely lost in the F275L mutant and there was a substantial rightward shift in

$[K^+]$ dependence for P742Q (**Fig. 5 C, D**). Whereas fluorescence change by 61 mM $[K^+]$ was

not changed (**Fig. 5 E**), the EC_{50} value was threefold higher in P742Q (33.2 ± 1.0 mM)

compared with WT (10.3 ± 0.9 mM) (**Fig. 5 F**). The other two mutations (E100K and

L1367V) had no significant effects on $[K^+]$ dependence (**Fig. 5 D-F**). We also tested the

effects of these mutations on caffeine-induced Ca^{2+} release. No substantial changes were

observed with any of the mutations (**Fig. 5 G–I**).

Effect of drugs on DICR activity

The reconstituted DICR platform is expected to be useful for screening drugs for

muscle diseases. To test this possibility, we examined the effects of known DICR modulators.

We initially tested three RyR1 inhibitors (**Fig. 6 A**). Dantrolene is a well-known RyR1

inhibitor that is clinically used for MH (Riazi et al., 2018). Cpd1 is a novel potent

RyR1-selective inhibitor that we recently developed (Mori et al., 2019; Yamazawa et al.,

2021). Procaine inhibits CICR but not DICR in frog skeletal muscle (Thorens and Endo,

1975; Kashiyama et al., 2010). Dantrolene (10 µM) and Cpd1 (3 µM) caused rightward shifts

in $[K^+]$ dependence compared with the control (**Fig. 6 B, C**). The maximum response by 61

mM $[K^+]$ was not changed (**Fig. 6 D**), but EC_{50} value was significantly higher in dantrolene

(13.1 ± 0.5 mM) or Cpd1 (13.5 ± 0.5 mM) compared to control (10.4 ± 0.3 mM) (**Fig. 6 E**).

These findings indicate that dantrolene and Cpd1 inhibit DICR. This is consistent with the

suppression of twitch and tetanic tension in isolated muscles and with reduced *in vivo* muscle

weakness by dantrolene (Meyler et al., 1979; Leslie and Part, 1981) or Cpd1 (Yamazawa et al.,

2021). Procaine (5 mM), in contrast, did not inhibit DICR but slightly reduced EC_{50} value

(9.0 ± 0.6 mM) (**Fig. 6 B-E**). All three compounds significantly shifted the caffeine

dependence rightward (**Fig. 6 F**). The maximum response by 20 mM caffeine was not

changed (**Fig. 6 G**), but EC_{50} values were increased by the drugs (1.9 ± 0.2, 3.0 ± 0.2, $3.1 \pm$

0.1, and 3.3 ± 0.4 mM for control, dantrolene, Cpd1, and procaine, respectively) (**Fig. 6 H**).

These results indicate that dantrolene and Cpd1 inhibit both DICR and CICR, whereas

procaine selectively inhibits CICR.

Lyotropic anions, such as perchlorate and thiocyanate, potentiate E-C coupling in

skeletal muscle (Luttgau et al., 1983; Huang, 1986; Csernoch et al., 1987; Delay et al., 1990;

Gonzalez and Rios, 1993). Perchlorate (10 mM) and thiocyanate (10 mM) significantly

shifted the [K^+] dependence leftward (**Fig. 7 A, B**). The EC_{50} values were significantly

reduced (8.6 ± 0.3 and 9.2 ± 0.3 mM for perchlorate and thiocyanate, respectively) compared

to control (10.6 ± 0.3 mM) without changing the maximum response (**Fig. 7 C, D**). In contrast,

they did not affect caffeine dependence (**Fig. 7 E**); no significant changes were observed with

the maximum response (**Fig. 7 F**) nor EC_{50} for caffeine (**Fig. 7 G**). These results indicate that

lyotropic anions potentiate DICR but not CICR. This is consistent with previous findings

showing that the potentiating effects of lyotropic anions are primarily caused by shifting

voltage dependence of the charge movement toward more negative potentials (Luttgau et al.,

1983; Huang, 1986; Csernoch et al., 1987; Delay et al., 1990; Gonzalez and Rios, 1993).

Taken together, our reconstituted DICR platform reproduces the effects of known DICR

modulators, indicating that it can be used to screen drugs that affect DICR.

Discussion

In this study, we established a platform for DICR that was reconstituted in HEK293 cells

expressing RyR1 and R-CEPIA1er. We made three key improvements to the original method

by Perni et al. (Perni et al., 2017). First, the essential components were transduced using

VSV-G pseudotyped baculovirus (BV), which can effectively infect a wide variety of

mammalian cells without toxicity (Tani et al., 2001). This greatly increased the transduction

efficiency; almost all cells expressed the essential components (**Fig. S1**). Second,

depolarization of the plasma membrane was induced by high [K^+] solution. This enabled us to

simultaneously stimulate many cells (**Fig. 1, Fig. S2** and **movie S1**). For this purpose, we

expressed Kir2.1, an inward-rectifying potassium channel, which effectively hyperpolarized

the membrane potential (Kirkton and Bursac, 2011) (**Fig. 3**). Third, ER [Ca^{2+}], instead of

cytoplasmic [Ca^{2+}], was measured to detect DICR. This is critically important to avoid

contamination of signals from extracellular Ca^{2+} influx. Indeed, ER [Ca^{2+}] was not changed

by high [K^+] depolarization in cells without essential components (**Fig. 1 A**), in contrast to

substantial increases in cytoplasmic [Ca^{2+}] (**Fig. S4 A**). These improvements allow

quantitative measurements of DICR using a microplate reader.

In our reconstitution, four components, RyR1, Cav1.1, β1a, and Stac3, are essential for

DICR; removal of each component totally abolished the DICR activity (**Fig. 2 A, B**). This is

consistent with previous findings from mice lacking each component (Tanabe et al., 1988;

Takeshima et al., 1994; Gregg et al., 1996; Nelson et al., 2013). In addition, the effects of α_2-δ

and γ_1 auxiliary subunits on DICR activity corresponded with findings of previous reports

(Ursu et al., 2004; Obermair et al., 2005) (**Fig. 2 C, D**). Therefore, our reconstituted platform

successfully reproduces DICR in HEK293 cells. We found that a substantial DICR still

occurred without JP2 (**Fig. 2 A, B**). This is in contrast to reports by Perni et al. (Perni et al.,

2017; Perni and Beam, 2022), in which no voltage-gated Ca^{2+} release was observed in

reconstituted cells without junctophilins. A possible reason for this is differences in

expression methods. Our VSV-G BV is more effective at transduction than standard

lipofection; therefore, sufficient proteins were expressed that might spontaneously interact

with each other without junctophilins. Thus, our data suggest that junctophilins are not

essential for DICR but may support formation of T-tubule-SR junction (Takeshima et al.,

2000). It would be of interest whether the tetrads (structural signs for functional coupling

between DHPR and RyR1) (Franzini-Armstrong, 2018) are formed in the cells expressing

essential components without junctophilins.

We quantitatively evaluated the effect of disease-causing mutations or drugs using $[K^+]$

dependence. The EC_{50} value for $[K^+]$ in ER $[Ca^{2+}]$ measurement was ~17 mM by calculation

from the rapid fluorescence change (5 sec after stimulation) (**Fig. 1 C**). This is consistent with

the value obtained from cytoplasmic $[Ca^{2+}]$ measurement (~16 mM, **Fig. S3**) or those

obtained from mouse skeletal muscle myotubes (~21 mM) in high $[K^+]$-induced Ca^{2+}

transients (Yang et al., 2006; Lopez et al., 2018). However, since variations were large under

the condition, we instead adopted the averaged fluorescence intensity for 125–150 sec after

stimulation, which was more stable and accurate (**Fig. 1 C**). Although the EC_{50} value was

reduced to ~10 mM, the results reproduced the effects of disease-causing mutations (**Fig. 3–5**)

or drugs (**Fig. 6, 7**), strongly indicating that the EC_{50} value for $[K^+]$ is a valid parameter for

quantitative evaluation of DICR.

Mutations in Cav1.1 are implicated in various muscle diseases, including MH and

myopathy (Beam et al., 2017; Flucher, 2020). Among three MH mutations (R174W, R1086H

and T1354S), we demonstrated that R1086H, but not R174W or T1354S, significantly shifted

the voltage dependence of DICR to more negative potentials (**Fig. 4 C–F**). This is consistent

with previous findings using Cav1.1-deficient (dysgenic) myotubes (Weiss et al., 2004; Pirone

et al., 2010; Eltit et al., 2012). All three mutations are reported to an increased sensitivity to

caffeine-induced Ca^{2+} release compared with WT, indicating enhanced CICR activity (Weiss

et al., 2004; Pirone et al., 2010; Eltit et al., 2012). However, we showed that caffeine

sensitivity was increased only in R1086H (**Fig. 4 G–I**). A possible reason for this difference is
the cells used. We used stable RyR1-overexpressing cells. This may cause an excess amount
of 'uncoupled' RyR1, which might mask the enhanced sensitivity of caffeine dependence of
the 'coupled' RyR1 by Cav1.1 mutations.

Most myopathy-related mutations identified in Cav1.1 have not been functionally
characterized. We evaluated four mutations (E100K, F275L, P742Q and L1367V) found in
patients with congenital myopathy (Schartner et al., 2017) (**Fig. 5**). To observe the effect of
mutations clearly, we tested the mutations in homozygous states. We demonstrated that
P742Q substantially inhibited DICR with a large rightward shifted $[K^+]$ dependence (**Fig. 5**
C-F). This proline reside in the II-III loop interacts with Stac3 (Wong King Yuen et al., 2017).
Substitution of the proline to threonine, the residue found in the corresponding position of
Cav1.2, diminished DICR in dysgenic myotubes (Kugler et al., 2004). Thus, the reduced
voltage dependence of DICR might partly be involved in pathophysiology of the patients.
E100K and L1367V had no or only minor impacts on the DICR activity (**Fig. 5 D-F**).
However, our platform cannot detect rapid kinetics of the channel. Measurements of ionic
currents and charge movements by patch-clamp method would further characterize these
mutants. F275L is recessive because it was found in two compound heterozygous sisters
carrying F275L and the nonsense mutation, p.L791Cfs*37 (Schartner et al., 2017). F275L
completely abolished DICR without any changes in ER $[Ca^{2+}]$ or caffeine sensitivity (**Fig. 5**
C-E). F275L was expressed at a size similar to WT (**Fig. 1 C**) and ER $[Ca^{2+}]$ was preserved
(**Fig. 5 B**). Thus, loss of DICR activity may not be due to loss of protein expression nor ER
$[Ca^{2+}]$ depletion. Loss of DICR in F275L does not fit with a recessive mutation; if this is the
case, patients will have no DICR activity. Further investigations are required for this
mutation.

There are no specific treatments for most DICR-related diseases; therefore, there is an
urgent need to develop novel drugs for these diseases (Agrawal et al., 2018; Flucher, 2020;
Lawal et al., 2020). We here evaluated DICR modulators. Among three known RyR1
inhibitors, dantrolene and Cpd1 suppressed DICR (**Fig. 6**). This is consistent with the
suppression of twitch and tetanic tensions of isolated muscles and of reduced *in vivo* muscle
weakness by dantrolene (Meyler et al., 1979; Leslie and Part, 1981) and Cpd1 (Yamazawa et
al., 2021). Interestingly, procaine did not suppress DICR (**Fig. 6**). Procaine inhibits CICR but
not DICR in frog skeletal muscle (Thorens and Endo, 1975) and reconstituted frog α-RyR and
β-RyR in RyR1-deficient myotubes (Kashiyama et al., 2010). Our data show similar procaine

action in a mammalian system. We also demonstrated that lyotropic anions (perchlorate and

thiocyanate) shifted the voltage-dependence of DICR to more negative potentials (**Fig. 7**).

This reproduces previous findings using skeletal muscle fibers (Luttgau et al., 1983; Huang,

1986; Csernoch et al., 1987; Delay et al., 1990; Gonzalez and Rios, 1993). Interestingly,

lyotropic anions did not affect caffeine dependence, suggesting no effect on CICR (**Fig. 7**).

Therefore, our reconstituted DICR platform will also be useful for drug testing and screening

for novel drugs that affect DICR.

Limitations and future perspective

The present reconstituted DICR platform has several limitations to be considered.

First, we overexpressed the essential components into HEK293 cells. Although this achieved

efficient formation of the DICR machinery, proteins were not only localized to plasma

membrane but also within cells except for in nuclei (Fig. S1 A). This is far from skeletal

muscle triads. The excess amount of uncoupled proteins might affect the results. Peripheral

localization of the components, e.g., RyR1 and Cav1.1, has been shown in the reconstituted

cells by Perni et al. (Perni et al., 2017; Perni and Beam, 2022). In our platform, protein

expression level can be easily controlled by amount of BV. Finding of the best amount and

combination ratio of each component would improve the situation.

Second, we measured DICR using long-lasting (~150 sec) high K^+ depolarization.

This provided efficient stimulation of the reconstituted cells but is markedly different from

the rapid Ca^{2+} release kinetics (~few ms) that occur during an action potential in muscle.

Parallel experiments using patch-clamp method is necessary to characterize the mutants. Our

reconstituted cells are applicable to single cell Ca^{2+} measurements combined with

patch-clamp method. This would measure the rapid kinetics of Ca^{2+} release by direct

membrane depolarization.

Third, we expressed the minimum essential components (RyR1, Cav1.1, β1a, Stac3,

JP2 and Kir2.1) for functional reconstitution of DICR. This is simple but different from the

skeletal muscle environment, in which a number of proteins (e.g., triadin, junctin,

calsequestrin, FKBP12, JP45 etc) associates and regulates the DICR machinery. Drug

screening with reconstitution by the minimum essential components might miss the

compounds that act on the associated proteins. Since VSV-G BV system easily expresses

additional proteins (e.g., α2-δ or γ1 subunit, see Fig. 2 D), it is possible to express the

associated proteins alone or in combination in our DICR platform. It would also be interesting

to express voltage-gated sodium and potassium channels to support electrically-evoked Ca^{2+}
release near future.

Conclusions

498   In summary, we established a reconstituted DICR platform in HEK293 cells. The
499 platform is highly efficient and quantitative thus will be useful for both evaluation of
500 disease-causing mutations and the development of novel drugs for DICR -related diseases.
501 Since the procedure is simple and reproducible, the reconstituted DICR platform will be a
502 powerful tool for accelerating diagnosis and treatment of these diseases.

Acknowledgments

We thank Ikue Hiraga and the Laboratory of Proteomics and Biomolecular Science,
Biomedical Research Core Facilities, Juntendo University Graduate School of Medicine, for
technical assistance. We thank Jeremy Allen, PhD, from Edanz (https://jp.edanz.com/ac) for
editing a draft of this manuscript. This work was supported by JSPS KAKENHI (19H03404
and 22H02805 to T.M., 19K07105 and 22K06652 to N.K., 20K11368 to T.K.), the Platform
Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting
Innovative Drug Discovery and Life Science Research (BINDS)) from the Japan Agency for
Medical Research and Development (AMED) (JP20am0101080 to T.M. and N.K. and
JP20am0101098 to H.K.), an Intramural Research Grant for Neurological and Psychiatric
Disorders from the National Center of Neurology and Psychiatry (2-5 to T.M.), the Vehicle
Racing Commemorative Foundation (6237 and 6303 to T.M.), and the Cooperative Research
Project of Research Center for Biomedical Engineering (to H.K.). The authors declare no
competing financial interests.

Author contributions

T. Murayama, N. Kurebayashi, S. Okazaki, K. Yamashiro carried out the cell biological
experiments; T. Numaga-Tomita and M. Yamada performed patch-clamp experiments; T.
Nakada, S. Mori, R. Ishida, and H. Kagechika provided experimental tools; T. Murayama, N.
Kurebayashi, T. Numaga-Tomita, T. Kobayashi, T. Nakada, M. Yamada, and T. Sakurai
analyzed the experimental data; T. Murayama, N. Kurebayashi, T. Numaga-Tomita and M.
Yamada wrote the manuscript. All authors reviewed and approved the manuscript.

References

Agrawal, A., G. Suryakumar, and R. Rathor. 2018. Role of defective Ca(2+) signaling in
529   skeletal muscle weakness: Pharmacological implications. *J Cell Commun Signal*.
530   12:645-659.
531   Bannister, R.A., and K.G. Beam. 2013. Impaired gating of an L-Type Ca(2+) channel carrying
532   a mutation linked to malignant hyperthermia. *Biophys J*. 104:1917-1922.
533   Beam, T.A., E.F. Loudermilk, and D.F. Kisor. 2017. Pharmacogenetics and pathophysiology
534   of CACNA1S mutations in malignant hyperthermia. *Physiol Genomics*. 49:81-87.
535   Block, B.A., T. Imagawa, K.P. Campbell, and C. Franzini-Armstrong. 1988. Structural
536   evidence for direct interaction between the molecular components of the transverse
537   tubule/sarcoplasmic reticulum junction in skeletal muscle. *J Cell Biol*. 107:2587-2600.
538   Carpenter, D., C. Ringrose, V. Leo, A. Morris, R.L. Robinson, P.J. Halsall, P.M. Hopkins, and
539   M.A. Shaw. 2009. The role of CACNA1S in predisposition to malignant hyperthermia.
540   *BMC Med Genet*. 10:104.
541   Catterall, W.A. 2000. Structure and regulation of voltage-gated Ca2+ channels. *Annu Rev Cell*
542   *Dev Biol*. 16:521-555.
543   Csernoch, L., L. Kovacs, and G. Szucs. 1987. Perchlorate and the relationship between charge
544   movement and contractile activation in frog skeletal muscle fibres. *J Physiol*.
545   390:213-227.
546   Delay, M., D.E. Garcia, and J.A. Sanchez. 1990. The effects of lyotropic anions on charge
547   movement, calcium currents and calcium signals in frog skeletal muscle fibres. *J*
548   *Physiol*. 425:449-469.
549   DiFranco, M., C. Yu, M. Quinonez, and J.L. Vergara. 2015. Inward rectifier potassium
550   currents in mammalian skeletal muscle fibres. *J Physiol*. 593:1213-1238.
551   Eltit, J.M., R.A. Bannister, O. Moua, F. Altamirano, P.M. Hopkins, I.N. Pessah, T.F. Molinski,
552   J.R. Lopez, K.G. Beam, and P.D. Allen. 2012. Malignant hyperthermia susceptibility
553   arising from altered resting coupling between the skeletal muscle L-type Ca2+ channel
554   and the type 1 ryanodine receptor. *Proc Natl Acad Sci U S A*. 109:7923-7928.
555   Flucher, B.E. 2020. Skeletal muscle CaV1.1 channelopathies. *Pflugers Arch*. 472:739-754.
556   Flucher, B.E., and M. Campiglio. 2019. STAC proteins: The missing link in skeletal muscle
557   EC coupling and new regulators of calcium channel function. *Biochim Biophys Acta*
558   *Mol Cell Res*. 1866:1101-1110.
559   Flucher, B.E., G.J. Obermair, P. Tuluc, J. Schredelseker, G. Kern, and M. Grabner. 2005. The
560   role of auxiliary dihydropyridine receptor subunits in muscle. *J Muscle Res Cell Motil*.

26:1-6.

Franzini-Armstrong, C. 2018. The relationship between form and function throughout the
history of excitation-contraction coupling. *J Gen Physiol*. 150:189-210.

Gonzalez, A., and E. Rios. 1993. Perchlorate enhances transmission in skeletal muscle
excitation-contraction coupling. *J Gen Physiol*. 102:373-421.

Gregg, R.G., A. Messing, C. Strube, M. Beurg, R. Moss, M. Behan, M. Sukhareva, S. Haynes,
J.A. Powell, R. Coronado, and P.A. Powers. 1996. Absence of the beta subunit (cchb1)
of the skeletal muscle dihydropyridine receptor alters expression of the alpha 1
subunit and eliminates excitation-contraction coupling. *Proc Natl Acad Sci U S A*.
93:13961-13966.

Horstick, E.J., J.W. Linsley, J.J. Dowling, M.A. Hauser, K.K. McDonald, A. Ashley-Koch, L.
Saint-Amant, A. Satish, W.W. Cui, W. Zhou, S.M. Sprague, D.S. Stamm, C.M. Powell,
M.C. Speer, C. Franzini-Armstrong, H. Hirata, and J.Y. Kuwada. 2013. Stac3 is a
component of the excitation-contraction coupling machinery and mutated in Native
American myopathy. *Nat Commun*. 4:1952.

Huang, C.L. 1986. The differential effects of twitch potentiators on charge movements in frog
skeletal muscle. *J Physiol*. 380:17-33.

Kashiyama, T., T. Murayama, E. Suzuki, P.D. Allen, and Y. Ogawa. 2010. Frog alpha- and
beta-ryanodine receptors provide distinct intracellular Ca2+ signals in a myogenic cell
line. *PLoS One*. 5:e11526.

Kirkton, R.D., and N. Bursac. 2011. Engineering biosynthetic excitable tissues from
unexcitable cells for electrophysiological and cell therapy studies. *Nat Commun*.
2:300.

Kitts, P.A., and G. Green. 1999. An immunological assay for determination of baculovirus
titers in 48 hours. *Anal Biochem*. 268:173-178.

Kugler, G., R.G. Weiss, B.E. Flucher, and M. Grabner. 2004. Structural requirements of the
dihydropyridine receptor alpha1S II-III loop for skeletal-type excitation-contraction
coupling. *J Biol Chem*. 279:4721-4728.

Lawal, T.A., J.J. Todd, J.W. Witherspoon, C.G. Bonnemann, J.J. Dowling, S.L. Hamilton,
K.G. Meilleur, and R.T. Dirksen. 2020. Ryanodine receptor 1-related disorders: an
historical perspective and proposal for a unified nomenclature. *Skelet Muscle*. 10:32.

Leslie, G.C., and N.J. Part. 1981. The action of dantrolene sodium on rat fast and slow muscle
in vivo. *Br J Pharmacol*. 72:665-672.

Lopez, J.R., V. Kaura, C.P. Diggle, P.M. Hopkins, and P.D. Allen. 2018. Malignant
hyperthermia, environmental heat stress, and intracellular calcium dysregulation in a

mouse model expressing the p.G2435R variant of RYR1. *Br J Anaesth*. 121:953-961.

Luttgau, H.C., G. Gottschalk, L. Kovacs, and M. Fuxreiter. 1983. How perchlorate improves

excitation-contraction coupling in skeletal muscle fibers. *Biophys J*. 43:247-249.

Meyler, W.J., I. Mols-Thurkow, A.H. Scaf, S. Sargo, and H. Wesseling. 1979. The effect of

dantrolene sodium on rat skeletal muscle in relation to the plasma concentration. *Eur J*

Pharmacol. 53:335-342.

Mori, S., H. Iinuma, N. Manaka, M. Ishigami-Yuasa, T. Murayama, Y. Nishijima, A. Sakurai,

R. Arai, N. Kurebayashi, T. Sakurai, and H. Kagechika. 2019. Structural development

of a type-1 ryanodine receptor (RyR1) Ca(2+)-release channel inhibitor guided by

endoplasmic reticulum Ca(2+) assay. *Eur J Med Chem*. 179:837-848.

Murayama, T., and N. Kurebayashi. 2019. Assays for Modulators of Ryanodine Receptor

(RyR)/Ca(2+) Release Channel Activity for Drug Discovery for Skeletal Muscle and

Heart Diseases. *Curr Protoc Pharmacol*. 87:e71.

Murayama, T., N. Kurebayashi, M. Ishigami-Yuasa, S. Mori, Y. Suzuki, R. Akima, H. Ogawa,

J. Suzuki, K. Kanemaru, H. Oyamada, Y. Kiuchi, M. Iino, H. Kagechika, and T.

Sakurai. 2018. Efficient High-Throughput Screening by Endoplasmic Reticulum Ca^{2+}

Measurement to Identify Inhibitors of Ryanodine Receptor Ca^{2+}-Release Channels.

Mol Pharmacol. 94:722-730.

Murayama, T., N. Kurebayashi, T. Yamazawa, H. Oyamada, J. Suzuki, K. Kanemaru, K.

Oguchi, M. Iino, and T. Sakurai. 2015. Divergent Activity Profiles of Type 1

Ryanodine Receptor Channels Carrying Malignant Hyperthermia and Central Core

Disease Mutations in the Amino-Terminal Region. *PLoS One*. 10:e0130606.

Nelson, B.R., F. Wu, Y. Liu, D.M. Anderson, J. McAnally, W. Lin, S.C. Cannon, R.

Bassel-Duby, and E.N. Olson. 2013. Skeletal muscle-specific T-tubule protein STAC3

mediates voltage-induced Ca2+ release and contractility. *Proc Natl Acad Sci U S A*.

110:11881-11886.

Obermair, G.J., G. Kugler, S. Baumgartner, P. Tuluc, M. Grabner, and B.E. Flucher. 2005. The

Ca2+ channel alpha2delta-1 subunit determines Ca2+ current kinetics in skeletal

muscle but not targeting of alpha1S or excitation-contraction coupling. *J Biol Chem*.

280:2229-2237.

Perez, C.F., J.M. Eltit, J.R. Lopez, D. Bodnar, A.F. Dulhunty, S. Aditya, and M.G. Casarotto.

2018. Functional and structural characterization of a novel malignant

hyperthermia-susceptible variant of DHPR-beta1a subunit (CACNB1). *Am J Physiol*

Cell Physiol. 314:C323-C333.

Perni, S., and K. Beam. 2022. Junctophilins 1, 2, and 3 all support voltage-induced Ca2+

631   release despite considerable divergence. *J Gen Physiol*. 154.

632 Perni, S., M. Lavorato, and K.G. Beam. 2017. De novo reconstitution reveals the proteins
633   required for skeletal muscle voltage-induced Ca(2+) release. *Proc Natl Acad Sci U S A*.
634   114:13822-13827.

635 Pirone, A., J. Schredelseker, P. Tuluc, E. Gravino, G. Fortunato, B.E. Flucher, A. Carsana, F.
636   Salvatore, and M. Grabner. 2010. Identification and functional characterization of
637   malignant hyperthermia mutation T1354S in the outer pore of the Cavalpha1S-subunit.
638   *Am J Physiol Cell Physiol*. 299:C1345-1354.

639 Polster, A., B.R. Nelson, E.N. Olson, and K.G. Beam. 2016. Stac3 has a direct role in skeletal
640   muscle-type excitation-contraction coupling that is disrupted by a myopathy-causing
641   mutation. *Proc Natl Acad Sci U S A*. 113:10986-10991.

642 Reinholt, B.M., X. Ge, X. Cong, D.E. Gerrard, and H. Jiang. 2013. Stac3 is a novel regulator
643   of skeletal muscle development in mice. *PLoS One*. 8:e62760.

644 Riazi, S., N. Kraeva, and P.M. Hopkins. 2018. Updated guide for the management of
645   malignant hyperthermia. *Can J Anaesth*. 65:709-721.

646 Rios, E., and G. Pizarro. 1991. Voltage sensor of excitation-contraction coupling in skeletal
647   muscle. *Physiol Rev*. 71:849-908.

648 Rufenach, B., and F. Van Petegem. 2021. Structure and function of STAC proteins: Calcium
649   channel modulators and critical components of muscle excitation-contraction coupling.
650   *J Biol Chem*. 297:100874.

651 Schartner, V., N.B. Romero, S. Donkervoort, S. Treves, P. Munot, T.M. Pierson, I. Dabaj, E.
652   Malfatti, I.T. Zaharieva, F. Zorzato, O. Abath Neto, G. Brochier, X. Lornage, B.
653   Eymard, A.L. Taratuto, J. Bohm, H. Gonorazky, L. Ramos-Platt, L. Feng, R. Phadke,
654   D.X. Bharucha-Goebel, C.J. Sumner, M.T. Bui, E. Lacene, M. Beuvin, C. Labasse, N.
655   Dondaine, R. Schneider, J. Thompson, A. Boland, J.F. Deleuze, E. Matthews, A.N.
656   Pakleza, C.A. Sewry, V. Biancalana, S. Quijano-Roy, F. Muntoni, M. Fardeau, C.G.
657   Bonnemann, and J. Laporte. 2017. Dihydropyridine receptor (DHPR, CACNA1S)
658   congenital myopathy. *Acta Neuropathol*. 133:517-533.

659 Schneider, M.F. 1994. Control of calcium release in functioning skeletal muscle fibers. *Annu*
660   *Rev Physiol*. 56:463-484.

661 Schredelseker, J., V. Di Biase, G.J. Obermair, E.T. Felder, B.E. Flucher, C.
662   Franzini-Armstrong, and M. Grabner. 2005. The beta 1a subunit is essential for the
663   assembly of dihydropyridine-receptor arrays in skeletal muscle. *Proc Natl Acad Sci U*
664   *S A*. 102:17219-17224.

Schredelseker, J., M. Shrivastav, A. Dayal, and M. Grabner. 2010. Non-Ca2+-conducting

Ca2+ channels in fish skeletal muscle excitation-contraction coupling. *Proc Natl Acad*
Sci U S A. 107:5658-5663.

Shishmarev, D. 2020. Excitation-contraction coupling in skeletal muscle: recent progress and
unanswered questions. *Biophys Rev*. 12:143-153.

Stewart, S.L., K. Hogan, H. Rosenberg, and J.E. Fletcher. 2001. Identification of the
Arg1086His mutation in the alpha subunit of the voltage-dependent calcium channel
(CACNA1S) in a North American family with malignant hyperthermia. *Clin Genet*.
59:178-184.

Suzuki, J., K. Kanemaru, K. Ishii, M. Ohkura, Y. Okubo, and M. Iino. 2014. Imaging
intraorganellar Ca2+ at subcellular resolution using CEPIA. *Nat Commun*. 5:4153.

Takeshima, H., M. Iino, H. Takekura, M. Nishi, J. Kuno, O. Minowa, H. Takano, and T. Noda.
1994. Excitation-contraction uncoupling and muscular degeneration in mice lacking
functional skeletal muscle ryanodine-receptor gene. *Nature*. 369:556-559.

Takeshima, H., S. Komazaki, M. Nishi, M. Iino, and K. Kangawa. 2000. Junctophilins: a
novel family of junctional membrane complex proteins. *Mol Cell*. 6:11-22.

Tanabe, T., K.G. Beam, J.A. Powell, and S. Numa. 1988. Restoration of excitation-contraction
coupling and slow calcium current in dysgenic muscle by dihydropyridine receptor
complementary DNA. *Nature*. 336:134-139.

Tani, H., M. Nishijima, H. Ushijima, T. Miyamura, and Y. Matsuura. 2001. Characterization
of cell-surface determinants important for baculovirus infection. *Virology*.
279:343-353.

Thorens, S., and M. Endo. 1975. Calcium-induced calcium release and
"depolarization"-induced calcium release: their physiological significance. *Proc Jpn*
Acad. 51:473-478.

Tong, J., H. Oyamada, N. Demaurex, S. Grinstein, T.V. McCarthy, and D.H. MacLennan.
1997. Caffeine and halothane sensitivity of intracellular Ca2+ release is altered by 15
calcium release channel (ryanodine receptor) mutations associated with malignant
hyperthermia and/or central core disease. *J Biol Chem*. 272:26332-26339.

Uehara, A., T. Murayama, M. Yasukochi, M. Fill, M. Horie, T. Okamoto, Y. Matsuura, K.
Uehara, T. Fujimoto, T. Sakurai, and N. Kurebayashi. 2017. Extensive Ca2+ leak
through K4750Q cardiac ryanodine receptors caused by cytosolic and luminal Ca2+
hypersensitivity. *J Gen Physiol*. 149:199-218.

Ursu, D., R.P. Schuhmeier, M. Freichel, V. Flockerzi, and W. Melzer. 2004. Altered
inactivation of Ca2+ current and Ca2+ release in mouse muscle fibers deficient in the
DHP receptor gamma1 subunit. *J Gen Physiol*. 124:605-618.

Weiss, R.G., K.M. O'Connell, B.E. Flucher, P.D. Allen, M. Grabner, and R.T. Dirksen. 2004.
Functional analysis of the R1086H malignant hyperthermia mutation in the DHPR
reveals an unexpected influence of the III-IV loop on skeletal muscle EC coupling. *Am*
J Physiol Cell Physiol. 287:C1094-1102.
Wong King Yuen, S.M., M. Campiglio, C.C. Tung, B.E. Flucher, and F. Van Petegem. 2017.
Structural insights into binding of STAC proteins to voltage-gated calcium channels.
Proc Natl Acad Sci U S A. 114:E9520-E9528.
Yamazawa, T., T. Kobayashi, N. Kurebayashi, M. Konishi, S. Noguchi, T. Inoue, Y.U. Inoue, I.
Nishino, S. Mori, H. Iinuma, N. Manaka, H. Kagechika, A. Uryash, J. Adams, J.R.
Lopez, X. Liu, C. Diggle, P.D. Allen, S. Kakizawa, K. Ikeda, B. Lin, Y. Ikemi, K.
Nunomura, S. Nakagawa, T. Sakurai, and T. Murayama. 2021. A novel RyR1-selective
inhibitor prevents and rescues sudden death in mouse models of malignant
hyperthermia and heat stroke. *Nat Commun*. 12:4293.
Yang, T., J. Riehl, E. Esteve, K.I. Matthaei, S. Goth, P.D. Allen, I.N. Pessah, and J.R. Lopez.
2006. Pharmacologic and functional characterization of malignant hyperthermia in the
R163C RyR1 knock-in mouse. *Anesthesiology*. 105:1164-1175.
Yang, Z.F., P. Panwar, C.R. McFarlane, W.E. Tuinte, M. Campiglio, and F. Van Petegem. 2022.
Structures of the junctophilin/voltage-gated calcium channel interface reveal hot spot
for cardiomyopathy mutations. *Proc Natl Acad Sci U S A*. 119:e2120416119.
Zhao, Y., S. Araki, J. Wu, T. Teramoto, Y.F. Chang, M. Nakano, A.S. Abdelfattah, M. Fujiwara,
T. Ishihara, T. Nagai, and R.E. Campbell. 2011. An expanded palette of genetically
encoded Ca(2)(+) indicators. *Science*. 333:1888-1891.
723
724

725   **Table 1. List of PCR primers.**
726

Protein/mutation	Primer	Nucleotide sequence
For cDNA cloning		
Cav1.1	F	AAA<u>AAGCTT</u>GCCACC ATG GAGCCGCCCTCACCCCAG
	R	TTT<u>ACGCGT</u>TCA AGGCTCCCAGGGCACGGC
β1a	F	AAA<u>GCTAGC</u>CACC ATG GTCCAGAAGAGCGGCATG
	R	TTT<u>GATATC</u>CTA CATGGCGTGCTCCTGAGG
JP2	F	AAA<u>GCTAGC</u>CACC ATG AGCGGGGGCCGCTTTGAC
	R	TTT<u>GATATC</u>TCA AGTCAGGAGGTGAACAAA
Stac3	F	AAA<u>GCTAGC</u>CACC ATG ACAGAAAGGAAGTGGTG
	R	TTT<u>ACGCGT</u>TTA AATCTCCTCCAGGAAGTC
Kir2.1	F	AAA<u>GCTAGC</u>CACC ATG GGCAGTGTGAGAACCAAC
	R	TTT<u>ACGCGT</u>TCA TATCTCCGATTCTCGCCT
For mutagenesis		
N617D	F	**G**ACTCTGTGATGTACAATGGG
	R	CCAGTCTTCACCCGTCAGGAC
R174W	F	**T**GGCTGGTGTCGGGGGTGCCC
	R	CAGGGGTCTGAGCACGCGGAA
R1086H	F	**A**CCAATGCGTACAGTATGCCC
	R	GCTGGTTCTTGTCGAGCTCAC
T1354S	F	**G**CAACTTCGCCTACTACTACT
	R	TGCCGCATGTATGCTCCTCCC
E100K	F	**A**AAGCCGCCATGAAGATCATA
	R	AATGGAGAAGACAATGAGGAA
F275L	F	**C**TCGACAACTTTGGCTTCTCT
	R	GTGGGTGATGCCGTGGTTGGG
P742Q	F	**A**AGGGGACGATGAGGAAGATG
	R	GGAAGTCAGCCGAAGGGTAGG
L1367V	F	**G**TCTGCGCCTTCCTGATCATC
	R	CATGTAGAAACTGATGAAGTA

727
728   F and R represent forward and reverse primers, respectively. Underlines indicate restriction
729   enzyme recognition sequence. Box indicates initiation or termination codon. Bold represents
730   mutated nucleotides.

 Figure legends

 Figure 1 **DICR measurements in 96-well plates using a fluorescence microplate reader.**

 (A) Schematic drawing of RyR1/R-CEPIA1er HEK293 cells without (No BV, left) and with

 (5 BV, right) BV infection of essential components for DICR (Cav1.1 carrying N617D Ca^{2+}

 non-conducting mutation, β1a, JP2, Stac3, and Kir2.1). **(B)** Typical results of time-lapse

 R-CEPIA1er fluorescence measurement using a FlexStation3 fluorescence microplate reader

 for RyR1/R-CEPIA1er HEK293 cells without (left, No BV) or with (right, 5 BV) BV

 infection. High $[K^+]$ solution ranging from 5–61 mM (symbols shown in right) or 10 mM

 caffeine (light blue) was applied at 30 sec after starting (blue arrowheads). **(C)** $[K^+]$

 dependence of R-CEPIA1er fluorescence (F/F_0) in No BV (black) and 5 BV cells (red). F/F_0

 was obtained by normalizing the fluorescence (F) at 35 sec (open circles) or the averaged

 fluorescence of 125-150 sec to that of the first 25 sec (F_0). Data are means $\pm$ SD (n = 9, N =

 3 for 5 BV and n = 6, N = 3 for No BV). Note that $[K^+]$ dependent plot with data at 35 sec

 (open red) was shifted rightward compared with that from data at 125–150 sec (filled red)

 with larger variations, especially around EC_{50} values. **(D)** Fluorescence change by 61 mM $[K^+]$

 or 10 mM caffeine. Fluorescence change is expressed as $1-F/F_0$ (see panel **C**). Data are means

 $\pm$ SD (n = 6, N = 3) and were analyzed by two-way ANOVA with Bonferroni's multiple

 comparisons test. Note that 5 BV cells show large fluorescence change with high $[K^+]$ which is

 nearly 80% of that with caffeine. "n" is the number of wells and "N" is the number of

 independent experiments.

 Figure 2 **Properties of the reconstituted DICR platform. (A)** Typical results of

 R-CEPIA1er fluorescence in cells infected with 5 BV (left), BV without Cav1.1 (center), and

 5 BV plus γ1 (right). **(B)** $[K^+]$ dependence of R-CEPIA1er fluorescence (F/F_0) in 5 BV

 (black), -Cav1.1 (red), and -JP2 (dark blue). Data are means ± SD (n = 9, N = 3 for 5 BV; n =

 6, N = 3 for -Cav1.1 and -JP2). **(C)** Fluorescence change by 61 mM $[K^+]$ in the BV-infected

 cells lacking one of essential components (-Cav1.1, -β1a, -Stac3, or -JP2) or with RyR2 cells

 instead of RyR1. Data are means ± SD (n = 9, N = 3 for 5 BV and n = 6, N = 3 for the others)

 and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. [***], p <

 0.0001 from 5 BV. **(D)** $[K^+]$ dependence of R-CEPIA1er fluorescence (F/F_0) in cells infected

 with 5 BV (black), 5 BV plus α_2-δ (purple) and 5 BV plus γ1 (orange). Data are means ± SD

 (n = 9, N = 3 for 5 BV; n = 6, N = 3 for + α_2-δ and + γ1). **(E)** Fluorescence change by 61 mM

 $[K^+]$ in 5 BV, +α_2-δ, and +γ1 cells. Data are means ± SD (n = 9, N = 3 for 5 BV; n = 6, N = 3

for + α_2-δ and + γ1) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test. "n" is the number of wells and "N" is the number of independent experiments.

Figure 3 **Importance of Kir2.1 on DICR platform. (A)** Resting membrane potential of cells

infected with 5 BV (left) or BV without Kir2.1 (-Kir2.1) (right). Data are means ± SD (n = 10,

N = 3 for 5 mM [K$^+$] with 5 BV; n = 16, N = 3 for 50 mM [K$^+$] with 5 BV; n = 10, N = 3 for 5

mM [K$^+$] with -Kir2.1 and n = 14, N = 3 for 50 mM [K$^+$] with -Kir2.1) and were analyzed by

two-way ANOVA with Bonferroni's multiple comparisons test. **(B)** Typical results of

R-CEPIA1er fluorescence in cells infected with 5 BV (left) and BV without Kir2.1 (right).

High [K$^+$] solution ranging from 5–61 mM (symbols shown in left) was applied at 30 sec after

starting (blue arrowheads). **(C)** Typical results of ER [Ca^{2+}] measurement in cells with 5 BV

(left), without Kir2.1, without Kir2.1 and β1a, and without Kir2.1 and Stac3. **(D)** Resting ER

[Ca^{2+}] in cells infected with 5 BV, BV without Kir2.1 (-Kir2.1), Kir2.1 and β1a (-Kir2.1,

-β1a), or Kir2.1 and Stac3 (-Kir2.1, -Stac3). Data are means ± SD (n = 8, N = 3) and were

analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of

cells (A) or wells (D) and "N" is the number of independent experiments.

Figure 4 **Effect of MH mutations in Cav1.1 on DICR activity. (A)** Schematic drawing of

the location of MH mutations (R174W, R1086H and T1354S) in Cav1.1. Note that WT and

mutant Cav1.1s also carry N617D Ca^{2+} non-conducting mutation. **(B)** Resting ER [Ca^{2+}] in

cells expressing WT and mutant Cav1.1. Data are means ± SD (n = 12; N = 3) and were

analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** Typical results

of R-CEPIA1er fluorescence in cells infected with 5 BV carrying WT (left), R174W (center)

and R1086H (right) Cav1.1. High [K$^+$] solution ranging from 5–61 mM (symbols shown in

left) was applied at 30 sec after starting (blue arrowheads). **(D)** [K$^+$] dependence of

R-CEPIA1er fluorescence corrected by ER [Ca^{2+}] in WT (black), R174W (red), R1086H

(blue) and T1354S (green) Cav1.1 cells. Data are means ± SD (n = 9, N = 3). **(E, F)**

Fluorescence change by 61 mM [K$^+$] corrected by ER Ca^{2+} **(E)** and EC$_{50}$ values for [K$^+$] **(F)**.

Data are means ± SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's

multiple comparisons test. **(G)** Caffeine dependence of R-CEPIA1er fluorescence corrected

by ER Ca^{2+} in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells.

Data are means ± SD (n = 9, N = 3). **(H, I)** Fluorescence change by 20 mM caffeine corrected

by ER Ca^{2+} **(H)** and EC$_{50}$ values for caffeine **(I)**. Data are means ± SD (n = 9, N = 3) and were

analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of

wells and "N" is the number of independent experiments.

Figure 5 **Effect of myopathy mutations in Cav1.1 on DICR activity.** **(A)** Schematic

drawing of the location of myopathy mutations (E100K, F275L, P742Q and L1367V) in

Cav1.1. Note that WT and mutant Cav1.1s also carry N617D Ca^{2+} non-conducting mutation.

(B) Resting ER $[Ca^{2+}]$ in cells expressing WT and mutant Cav1.1. Data are means ± SD (n =

12, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

(C) Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV carrying WT

(left), F275L (center) and P742Q (right) Cav1.1. High $[K^+]$ solution ranging from 5–61 mM

(symbols shown in left) was applied at 30 sec after starting (blue arrowheads). **(D)** $[K^+]$

dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue),

P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means ± SD (n = 9, N = 3). **(E,**

F) Fluorescence change by 61 mM $[K^+]$ **(E)** and EC_{50} values for $[K^+]$ **(F)**. Data are means ±

SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test. **(G)** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black),

E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are

means ± SD (n = 9, N = 3). **(H, I)** Fluorescence change by 20 mM caffeine **(H)** and EC_{50}

values for caffeine **(I)**. Data are means ± SD (n = 9, N = 3) and were analyzed by one-way

ANOVA with Dunnett's multiple comparisons test. "n" is the number of wells and "N" is the

number of independent experiments.

Figure 6 **Effect of RyR1 inhibitors on DICR activity.** **(A)** Chemical structures of dantrolene,

Cpd1 and procaine. **(B)** Typical results of R-CEPIA1er fluorescence in cells without (Control,

left) or with 10 μM dantrolene (center) or 5 mM procaine (right). High $[K^+]$ solution ranging

from 5–61 mM (symbols shown in left) was applied at 30 sec after starting (blue arrowheads).

(C) $[K^+]$ dependence of fluorescence change (F/F_0) in cells without (Cont, black) or with 10

μM dantrolene (Dan, red), 3 μM Cpd1 (Cpd1, blue) or 5 mM procaine (Pro, green). Data are

means ± SD (n = 9, N = 3). **(D, E)** Fluorescence change by 61 mM K^+ **(D)** and EC_{50} values

for $[K^+]$ **(E)**. Data are means ± SD (n = 9 , N = 3) and were analyzed by one-way ANOVA

with Dunnett's multiple comparisons test. **(F)** Caffeine dependence of R-CEPIA1er

fluorescence (F/F_0) in cells without (Control, black) or with 10 μM dantrolene (Dan, red), 3

μM Cpd1 (Cpd1, blue) or 5 mM procaine (Pro, green). Data are means ± SD (n = 9, N = 3).

(G, H) Fluorescence change by 20 mM caffeine (**G**) and EC_{50} values for caffeine (**H**). Data

are means ± SD (n = 9 , N = 3) and were analyzed by one-way ANOVA with Dunnett's

multiple comparisons test. "n" is the number of wells and "N" is the number of independent

experiments.

Figure 7 **Effect of lyotropic anions on DICR activity.** (**A**) Typical results of R-CEPIA1er

fluorescence in cells in the absence (Control, left) and presence of 10 mM perchlorate (ClO_4^-,

center) or 10 mM thiocyanate (SCN^-, right). High [K^+] solution ranging from 5–61 mM

(symbols shown in left) was applied at 30 seconds after starting (blue arrowheads). (**B**) [K^+]

dependence of fluorescence change (F/F$_0$) in cells in the absence (Cont, black) and presence

of 10 mM perchlorate (ClO_4^-, red) or 10 mM thiocyanate (SCN^-, blue). Data are means ± SD

(n = 9, N = 3). (**C, D**) Fluorescence change by 61 mM K^+ (**C**) and EC_{50} values for K^+ (**D**).

Data are means ± SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's

multiple comparisons test. EC_{50} values were significantly reduced with lyotropic anions. (**E**)

Caffeine dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells in the absence (Cont, black)

and presence of 10 mM perchlorate (ClO_4^-, red) or 10 mM thiocyanate (SCN^-, blue). Data are

means ± SD (n = 9, N = 3). (**F, G**) Fluorescence change by 20 mM caffeine (**F**) and EC_{50}

values for caffeine (**G**). Data are means ± SD (n = 9, N = 3) and were analyzed by one-way

ANOVA with Dunnett's multiple comparisons test. No significant changes were observed by

lyotropic anions. "n" is the number of wells and "N" is the number of independent

experiments.

Figure S1 **Expression of essential components for DICR machinery using VSV-G**

pseudotyped BV. (**A**) Immunofluorescent detection of Cav1.1, β1a, Stac3, JP2 and Kir2.1 in

5 BV (left) and No BV (right) cells. Cells were labeled with antibodies to each component,

followed by Alexa594-labeled anti-mouse IgG. Scale bar, 20 µm. Note that red fluorescence

was specifically observed with 5 BV cells. (**B**) Western blots of essential components. Lysates

from No BV and 5 BV cells were separated by SDS-PAGE and probed with antibodies to

each component. Calnexin (CLNX) was used as a loading control. (**C**) Western blot of Cav1.1

mutants. Lysates from 5 BV cells carrying WT, R174W, R1086H, T1354S, E100K, F275L,

P742Q and L1367V Cav1.1 were separated by SDS-PAGE and probed with anti-Cav1.1

antibody. Calnexin (CLNX) was used as loading control.

Figure S2 **Visualization of DICR by laser scanning confocal microscopy. (A)** Typical
results of time-lapse R-CEPIA1er fluorescence measurement of individual 5 BV-infected
RyR1/R-CEPIA1er HEK293 cells using a laser scanning confocal microscope (see also
Movie S1). Cells were incubated with normal Krebs solution and high $[K^+]$ (50 mM) solution
was perfused during measurements (arrow). Note that R-CEPIA1er fluorescence was
transiently decreased by high $[K^+]$ solution and recovered after replacement with normal
Krebs solution. **(B)** Percent responding cells to high $[K^+]$ (50 mM) or caffeine (10 mM)
solution. 100 cells were randomly picked from each of No BV or 5 BV cells. Whereas 95% of
5 BV cells responded to high $[K^+]$, less than 1% of No BV cells responded. All the cells
responded to caffeine. Data are means ± SD (n = 3, N = 3). "n" is the number of cells and "N"
is the number of independent experiments.

Figure S3 **Measurement of DICR with cytoplasmic $[Ca^{2+}]$. (A)** Typical results of
time-lapse G-GECO1.1 fluorescence measurement in No BV (left) and 5 BV (right) cells.
High $[K^+]$ solution ranging from 5–61 mM (shown at right) was applied at 30 sec after
starting (blue arrows). Note that a substantial Ca^{2+} transient was observed in No BV,
indicating the existence of endogenous Ca^{2+} influx pathways by depolarization. **(B)** $[K^+]$
dependence of fluorescence change (F/F_0) in No BV (black) or 5 BV cells (red). F/F_0 was
obtained by normalizing the fluorescence signal at 35 sec (F, grey lines in **A**) to the averaged
signals for the first 25 sec (F_0). Data are means±SD (n = 6, N = 3). "n" is the number of wells
and "N" is the number of independent experiments.

Figure S4 **Effect of the Cav1.1 R1086H mutation on DICR activity at 2.5 mM $[K^+]$. (A)**
Resting membrane potential of 5 BV cells in 5 mM or 2.5 mM $[K^+]$ Krebs solution. Data are
the median with individual data points (n = 10, N = 3 for 5 mM $[K^+]$ and n = 20, N = 3 for 2.5
mM $[K^+]$) and were analyzed by the unpaired two-tailed t test with the Mann-Whitney test.
(B) Resting ER $[Ca^{2+}]$ in WT (black) and R1086H (blue) Cav1.1 cells. Data are means ± SD
(n = 12, N = 3) and were analyzed by the two-tailed T-test with the Mann Whitney test. **(C)**
$[K^+]$ dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black) and R1086H (blue)
Cav1.1 cells. Data are means ± SD (n = 9, N = 3). **(D, E)** Fluorescence change by 61 mM $[K^+]$
(D) and EC_{50} values for $[K^+]$ **(E)**. Data are means ± SD (n = 9, N = 3) and were analyzed by
the two-tailed T-test with the Mann Whitney test. "n" is the number of cells (A) or wells (B-E)
and "N" is the number of independent experiments.

Figure 1 **DICR measurements in 96-well plates using a fluorescence microplate reader. (A)** Schematic drawing of RyR1/R-CEPIA1er HEK293 cells without (No BV, left) and with (5 BV, right) BV infection of essential components for DICR (Cav1.1 carrying N617D Ca^{2+} non-conducting mutation, β1a, JP2, Stac3, and Kir2.1). **(B)** Typical results of time-lapse R-CEPIA1er fluorescence measurement using a FlexStation3 fluorescence microplate reader for RyR1/R-CEPIA1er HEK293 cells without (left, No BV) or with (right, 5 BV) BV infection. High $[K^+]$ solution ranging from 5–61 mM (symbols shown in right) or 10 mM caffeine (light blue) was applied at 30 sec after starting (blue arrowheads). **(C)** $[K^+]$ dependence of R-CEPIA1er fluorescence (F/F_0) in No BV (black) and 5 BV cells (red). F/F_0 was obtained by normalizing the fluorescence (F) at 35 sec (open circles) or the averaged fluorescence of 125-150 sec to that of the first 25 sec (F_0). Data are means ± SD (n = 9, N = 3 for 5 BV and n = 6, N = 3 for No BV). Note that $[K^+]$ dependent plot with data at 35 sec (open red) was shifted rightward compared with that from data at 125–150 sec (filled red) with larger variations, especially around EC_{50} values. **(D)** Fluorescence change by 61 mM $[K^+]$ or 10 mM caffeine. Fluorescence change is expressed as 1-F/F_0 (see panel **C**). Data are means ± SD (n = 6, N = 3) and were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test. Note that 5 BV cells show large fluorescence change with high $[K^+]$ which is nearly 80% of that with caffeine. "n" is the number of wells and "N" is the number of independent experiments.

Figure 2 **Properties of the reconstituted DICR platform. (A)** Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV (left), BV without Cav1.1 (center), and 5 BV plus γ1 (right). **(B)** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in 5 BV (black), -Cav1.1 (red), and -JP2 (dark blue). Data are means $\pm$ SD (n = 9, N = 3 for 5 BV; n = 6, N = 3 for -Cav1.1 and -JP2). **(C)** Fluorescence change by 61 mM [K$^+$] in the BV-infected cells lacking one of essential components (-Cav1.1, -β1a, -Stac3, or -JP2) or with RyR2 cells instead of RyR1. Data are means $\pm$ SD (n = 9, N = 3 for 5 BV and n = 6, N = 3 for the others) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. ***, p < 0.0001 from 5 BV. **(D)** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells infected with 5 BV (black), 5 BV plus α$_2$-δ (purple) and 5 BV plus γ1 (orange). Data are means $\pm$ SD (n = 9, N = 3 for 5 BV; n = 6, N = 3 for + α$_2$-δ and + γ1). **(E)** Fluorescence change by 61 mM [K$^+$] in 5 BV, +α$_2$-δ, and +γ1 cells. Data are means $\pm$ SD (n = 9, N = 3 for 5 BV; n = 6, N = 3 for + α$_2$-δ and + γ1) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of wells and "N" is the number of independent experiments.

Figure 3 **Importance of Kir2.1 on DICR platform. (A)** Resting membrane potential of cells infected with 5 BV (left) or BV without Kir2.1 (-Kir2.1) (right). Data are means $\pm$ SD (n = 10, N = 3 for 5 mM [K⁺] with 5 BV; n = 16, N = 3 for 50 mM [K⁺] with 5 BV; n = 10, N = 3 for 5 mM [K⁺] with -Kir2.1 and n = 14, N = 3 for 50 mM [K⁺] with -Kir2.1) and were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test. **(B)** Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV (left) and BV without Kir2.1 (right). High [K⁺] solution ranging from 5–61 mM (symbols shown in left) was applied at 30 sec after starting (blue arrowheads). **(C)** Typical results of ER [Ca²⁺] measurement in cells with 5 BV (left), without Kir2.1, without Kir2.1 and β1a, and without Kir2.1 and Stac3. **(D)** Resting ER [Ca²⁺] in cells infected with 5 BV, BV without Kir2.1 (-Kir2.1), Kir2.1 and β1a (-Kir2.1, -β1a), or Kir2.1 and Stac3 (-Kir2.1, -Stac3). Data are means $\pm$ SD (n = 8, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of cells (A) or wells (D) and "N" is the number of independent experiments.

Figure 4 **Effect of MH mutations in Cav1.1 on DICR activity. (A)** Schematic drawing of the location of MH mutations (R174W, R1086H and T1354S) in Cav1.1. Note that WT and mutant Cav1.1s also carry N617D Ca^{2+} non-conducting mutation. **(B)** Resting ER $[Ca^{2+}]$ in cells expressing WT and mutant Cav1.1. Data are means $\pm$ SD (n = 12; N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV carrying WT (left), R174W (center) and R1086H (right) Cav1.1. High $[K^+]$ solution ranging from 5–61 mM (symbols shown in left) was applied at 30 sec after starting (blue arrowheads). **(D)** $[K^+]$ dependence of R-CEPIA1er fluorescence corrected by ER $[Ca^{2+}]$ in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means $\pm$ SD (n = 9, N = 3). **(E, F)** Fluorescence change by 61 mM $[K^+]$ corrected by ER Ca^{2+} **(E)** and EC_{50} values for $[K^+]$ **(F)**. Data are means $\pm$ SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(G)** Caffeine dependence of R-CEPIA1er fluorescence corrected by ER Ca^{2+} in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means $\pm$ SD (n = 9, N = 3). **(H, I)** Fluorescence change by 20 mM caffeine corrected by ER Ca^{2+} **(H)** and EC_{50} values for caffeine **(I)**. Data are means $\pm$ SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of wells and "N" is the number of independent experiments.

Figure 5 **Effect of myopathy mutations in Cav1.1 on DICR activity. (A)** Schematic drawing of the location of myopathy mutations (E100K, F275L, P742Q and L1367V) in Cav1.1. Note that WT and mutant Cav1.1s also carry N617D Ca^{2+} non-conducting mutation. **(B)** Resting ER $[Ca^{2+}]$ in cells expressing WT and mutant Cav1.1. Data are means $\pm$ SD (n = 12, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(C)** Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV carrying WT (left), F275L (center) and P742Q (right) Cav1.1. High $[K^+]$ solution ranging from 5–61 mM (symbols shown in left) was applied at 30 sec after starting (blue arrowheads). **(D)** $[K^+]$ dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means $\pm$ SD (n = 9, N = 3). **(E, F)** Fluorescence change by 61 mM $[K^+]$ **(E)** and EC_{50} values for $[K^+]$ **(F)**. Data are means $\pm$ SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(G)** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means $\pm$ SD (n = 9, N = 3). **(H, I)** Fluorescence change by 20 mM caffeine **(H)** and EC_{50} values for caffeine **(I)**. Data are means $\pm$ SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of wells and "N" is the number of independent experiments.

Figure 6 **Effect of RyR1 inhibitors on DICR activity. (A)** Chemical structures of dantrolene, Cpd1 and procaine. **(B)** Typical results of R-CEPIA1er fluorescence in cells without (Control, left) or with 10 μM dantrolene (center) or 5 mM procaine (right). High [K+] solution ranging from 5–61 mM (symbols shown in left) was applied at 30 sec after starting (blue arrowheads). **(C)** [K+] dependence of fluorescence change (F/F_0) in cells without (Cont, black) or with 10 μM dantrolene (Dan, red), 3 μM Cpd1 (Cpd1, blue) or 5 mM procaine (Pro, green). Data are means ± SD (n = 9, N = 3). **(D, E)** Fluorescence change by 61 mM K+ **(D)** and EC_{50} values for [K+] **(E)**. Data are means ± SD (n = 9 , N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **(F)** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in cells without (Control, black) or with 10 μM dantrolene (Dan, red), 3 μM Cpd1 (Cpd1, blue) or 5 mM procaine (Pro, green). Data are means ± SD (n = 9, N = 3). **(G, H)** Fluorescence change by 20 mM caffeine **(G)** and EC_{50} values for caffeine **(H)**. Data are means ± SD (n = 9 , N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. "n" is the number of wells and "N" is the number of independent experiments.

Figure 7 **Effect of lyotropic anions on DICR activity.** (**A**) Typical results of R-CEPIA1er fluorescence in cells in the absence (Control, left) and presence of 10 mM perchlorate (ClO_4^-, center) or 10 mM thiocyanate (SCN^-, right). High $[K^+]$ solution ranging from 5–61 mM (symbols shown in left) was applied at 30 seconds after starting (blue arrowheads). (**B**) $[K^+]$ dependence of fluorescence change (F/F_0) in cells in the absence (Cont, black) and presence of 10 mM perchlorate (ClO_4^-, red) or 10 mM thiocyanate (SCN^-, blue). Data are means $\pm$ SD (n = 9, N = 3). (**C, D**) Fluorescence change by 61 mM K^+ (**C**) and EC_{50} values for K^+ (**D**). Data are means $\pm$ SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. EC_{50} values were significantly reduced with lyotropic anions. (**E**) Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in cells in the absence (Cont, black) and presence of 10 mM perchlorate (ClO_4^-, red) or 10 mM thiocyanate (SCN^-, blue). Data are means $\pm$ SD (n = 9, N = 3). (**F, G**) Fluorescence change by 20 mM caffeine (**F**) and EC_{50} values for caffeine (**G**). Data are means $\pm$ SD (n = 9, N = 3) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. No significant changes were observed by lyotropic anions. "n" is the number of wells and "N" is the number of independent experiments.