High-throughput platform for depolarization-induced Ca^{2+} release for skeletal muscle

disease mutation validation and drug discovery

Takashi Murayama[1]*, Nagomi Kurebayashi[1], Takuro Numaga-Tomita[2], Takuya Kobayashi[1],

Satoru Okazaki[1], Kyosuke Yamashiro[1], Tsutomu Nakada[2], Shuichi Mori[3], Ryosuke Ishida[3],

Hiroyuki Kagechika[3], Mitsuhiko Yamada[2], Takashi Sakurai[1]

[1]Department of Pharmacology, Juntendo University School of Medicine, Tokyo, Japan.

[2]Department of Molecular Pharmacology, Shinshu University School of Medicine, Matsumoto,

Japan.

[3]Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, Tokyo,

Japan.

*Corresponding author: Takashi Murayama, Ph.D., Department of Pharmacology, Juntendo

University School of Medicine, Tokyo 113-8421, Japan. E-mail: takashim@juntendo.ac.jp

Abstract

In skeletal muscle excitation-contraction (E-C) coupling, depolarization of the plasma

membrane triggers Ca^{2+} release from the sarcoplasmic reticulum (SR), referred to as

depolarization-induced Ca^{2+} release (DICR). DICR occurs via the type 1 ryanodine receptor

(RyR1), which physically interacts with the dihydropyridine receptor Cav1.1 subunit in specific

machinery formed with additional components. Exome sequencing has accelerated the

discovery of many novel mutations in DICR machinery genes in various skeletal muscle

diseases. However, functional validation is time-consuming because it must be performed in a

skeletal muscle environment. Here, we describe a high-throughput platform for DICR that is

reconstituted in non-muscle HEK293 cells. We demonstrate that Cav1.1 mutations implicated

in malignant hyperthermia and myopathy exhibit divergent effects on DICR. We also tested

several RyR1 inhibitors and DICR modulators. This high-throughput DICR platform will

accelerate validation of mutations and drug discovery for skeletal muscle diseases.

Introduction

In vertebrate skeletal muscle, depolarization of the plasma membrane triggers Ca^{2+} release from the sarcoplasmic reticulum (SR), which is commonly referred to as excitation-contraction (E-C) coupling. This process occurs at triad junctions in which a transverse (T) tubule is closely apposed to terminal cisterna of the SR[1]. In excitation-contraction coupling, the type 1 ryanodine receptor (RyR1), a Ca^{2+} release channel in the SR, is activated by a conformational change of the voltage sensor in the Cav1.1 subunit of the dihydropyridine receptor (DHPR) upon depolarization of the T-tubule membrane. This is referred to as depolarization-induced Ca^{2+} release (DICR)[2,3]. Studies with knockout animals have revealed that functional and structural interactions between Cav1.1 and RyR1 in muscle require at least three additional components, a β1a auxiliary subunit of DHPR[4,5], the Stac3 adaptor protein[6,7,8], and junctophilins[9], which may form specific machinery for DICR[4,5,10].

Mutations in genes that encode DICR machinery components have been implicated in various types of muscle diseases, including congenital myopathies, periodic paralysis, and malignant hyperthermia (MH)[11,12,13]. Rapid exome sequencing has accelerated the discovery of novel mutations; however, validation of these mutations by functional testing is critically important for the diagnosis and treatment of these diseases. This has been performed using myotubes from mice that lack DICR machinery components, for example knockout mice of Cav1.1[14], β1a[15] and Stac3[16]. While this approach has evaluated several disease-causing mutations in the DICR components [16–20], it has certain limitations. For example, it is expensive and labor intensive to maintain the mouse lines. Differentiation of cells into myotubes is time consuming and is affected by various factors, such as the cell line used and experimental conditions. Furthermore, transduction of foreign genes into myotubes is often inefficient.

There are no specific treatments for most diseases related to mutations in the DICR machinery. A major cause for the slow pace of novel drug development is the lack of an appropriate drug screening platform for these diseases. We have recently established an efficient high-throughput platform for screening RyR1-targeted drugs using endoplasmic reticulum (ER) Ca^{2+} measurements in HEK293 cells[21,22] and using this platform we developed Compound 1 (Cpd1), a novel RyR1 inhibitor for treatment of MH[23,24]. Therefore, functional screening using HEK293 cells is a promising platform for drug discovery.

Recently, Perni et al.[25] successfully reconstituted DICR by co-expressing five essential components, RyR1, Cav1.1, β1a, junctophilin-2 (JP2) and Stac3, in non-muscle tsA201 cells.

They demonstrated formation of ER-plasma membrane junctions in which Cav1.1 is arranged
into tetrads, which is indicative of physical links to RyR1[26]. Depolarization of the plasma
membrane using the patch-clamp technique induced a rapid Ca^{2+} transient without influx of
extracellular Ca^{2+}. This provides a basis for developing reconstituted DICR.

In this study, we established a high-throughput platform for DICR that was
reconstituted in HEK293 cells. Baculovirus infection of essential components greatly improved
transduction efficiency. Chemical depolarization by expression of an inward-rectified
potassium channel (Kir2.1) and high concentration potassium $[K^+]$ solutions enabled the
simultaneous stimulation of many cells. Furthermore, a genetically-encoded ER Ca^{2+} indicator
quantitatively measured DICR without influence of extracellular Ca^{2+} influx. Using this
platform, we successfully evaluated disease-causing mutations in Cav1.1 and tested several
RyR1 inhibitors and DICR modulators. Our high-throughput platform will accelerate the
validation of mutations and drug discovery for skeletal muscle diseases.

Results

Establishment of a high-throughput platform for DICR using a fluorescence microplate
reader

To reconstitute DICR, we used HEK293 cells stably expressing RyR1 and R-
CEPIA1er, a genetically-encoded fluorescent ER Ca^{2+} indicator[27], that were generated for high-
throughput screening of novel RyR1 inhibitors[21,22] (**Fig. 1a**, left). The other components for
DICR, Cav1.1, β1a, Stac3 and JP2, which were used by Perni et al.[25], were transduced using
VSV-G pseudotyped baculovirus (BV), which efficiently infects mammalian cells[28,29] (**Fig. 1a**,
right). In these cells, DICR can be detected by a decrease in ER Ca^{2+}. To prevent extracellular
Ca^{2+} influx through Cav1.1 by depolarization, a N617D mutation was introduced that abolishes
Ca^{2+} conductance[30]. In addition, we expressed Kir2.1, an inward-rectified potassium channel in
skeletal muscle[31], to hyperpolarize the membrane potential because the resting membrane
potential in HEK293 cells is much less (−20 mV) than in skeletal muscle (−90 mV)[32].
Imunocytochemistry revealed that the five proteins (Cav1.1, β1a, Stac3, JP2, and Kir2.1) were
successfully expressed in almost all cells, which we term hereafter 5 BV cells (**Supplementary**
Fig. 1a). BV-dependent protein expression was also confirmed by western blotting of cell lysate
(**Supplementary Fig. 1b**). Therefore, BV infection effectively transduced the essential
components of DICR in RyR1/R-CEPIA1er-HEK293 cells.

To detect DICR, we used a fluorescence microplate reader (FlexStation3), which

measures time-lapse R-CEPIA1er fluorescence in 96-well plates. Fluorescence signals were

collected in a 1.5 mm diameter area; therefore, an averaged signal from thousands of cells was

detected in each well. We adopted chemical depolarization using high $[K^+]$ solutions, which

can simultaneously depolarize all the cells to trigger DICR. The cells were initially incubated

with normal Krebs solution containing 5 mM K^+. Then, solutions containing various K^+

concentrations were added to give K^+ concentrations of 5–61 mM. In cells without BV infection,

no apparent changes in fluorescence were observed with any change in $[K^+]$ (**Fig. 1b**). Caffeine

(10 mM) reduced the R-CEPIA1er fluorescence, confirming the functional expression of RyR1.

In 5 BV cells, high $[K^+]$ solution rapidly reduced the R-CEPIA1er fluorescence in a $[K^+]$-

dependent manner (**Fig. 1c**). To monitor ER Ca^{2+} changes in individual cells, we also observed

R-CEPIA1er fluorescence using laser scanning confocal microscopy. In 5 BV cells, high $[K^+]$

solution caused a large and transient reduction in the R-CEPIA1er fluorescence in 90% of cells

(**Movie S1** and **Supplementary Fig. 2**).

For quantitative analysis, we normalized the averaged fluorescence intensity of the last

25 seconds (F) to that of the initial 25 seconds (F_0) to correct variations in the initial

fluorescence intensity between wells. The F/F_0 plot as a function of K^+ concentrations

demonstrated K^+ dependence with an EC_{50} of ~10 mM K^+ (**Fig. 1d**). The initial fluorescence

change after stimulation (i.e., 35 seconds) may reflect the rate of Ca^{2+} release. However,

variations were too large to accurately determine the EC_{50} values because of rapid changes in

fluorescence intensity (**Supplementary Fig. 3**). The fluorescence change by 61 mM K^+, which

is expressed as $1-F/F_0$ (**Fig. 1d**), reached ~80% of that induced by caffeine in 5 BV cells,

indicating a high degree of DICR reconstitution (**Fig. 1e**). Taken together, these results indicate

successful establishment of a high-throughput platform for DICR.

The original method for DICR reconstitution monitors cytoplasmic $[Ca^{2+}]$[25]; therefore,

we also assessed DICR by measuring cytoplasmic Ca^{2+} using G-GECO1.1, a genetically-

encoded Ca^{2+} indicator[33]. In cells without BV infection, a substantial increase in $[Ca^{2+}]$ was

observed by high $[K^+]$, indicating the existence of endogenous Ca^{2+} influx pathways by

depolarization (**Supplementary Fig. 4a**). In 5 BV cells, DICR was detected by rapid and large

increases in $[Ca^{2+}]$ (**Supplementary Fig. 4b**). However, $[Ca^{2+}]$ rapidly declined with time. This

may be caused by Ca^{2+} extrusion pathways in HEK293 cells. The transient nature of the $[Ca^{2+}]$

rise and substantial background make it difficult to quantitatively measure DICR by

cytoplasmic $[Ca^{2+}]$ in our platform.

Validation of the reconstituted DICR

We next determined whether the reconstituted platform reproduces DICR in skeletal muscle. Initially, we examined the role of the essential components. Removal of Cav1.1 resulted in complete loss of high [K^+]-induced Ca^{2+} release (**Fig. 2a, b**). Ca^{2+} release was also lost in cells lacking β1a or Stac3, and in cells expressing RyR2 instead of RyR1 (**Fig. 2b**). Removal of JP2 significantly reduced the fluorescence change by 61 mM K^+ with a large rightward shift in K^+ dependence (**Fig. 2a, b**). These results are consistent with previous findings of mice lacking each component[9,14–16,34].

Skeletal muscle Cav1.1 forms a complex with three auxiliary subunits, β1a, α_2-δ and γ_1[35]. β1a is essential for DICR, while α_2-δ and γ_1 may play modulatory roles[36]. Additional expression of α_2-δ had no effect on high [K^+]-induced Ca^{2+} release (**Fig. 2c, d**). Expression of γ1 did not significantly affect high [K^+]-induced Ca^{2+} release at 20 mM or lower [K^+] but it reduced the fluorescence change at higher K^+ concentrations (**Fig. 2c, d**). In γ_1 cells, R-CEPIA1er fluorescence decreased but increased again with time, suggesting inactivation of the Ca^{2+} release (**Supplementary Fig. 5**). These findings are consistent with previous reports of the effects of α_2-δ[37] and γ_1[38] on DICR. Therefore, our reconstituted platform can reproduce DICR in skeletal muscle and is valid for evaluating disease-causing mutations or drugs.

We also tested the importance of Kir2.1. Removal of Kir2.1 completely abolished high [K^+]-induced Ca^{2+} release with substantial reduction in the R-CEPIA1er fluorescence intensity (**Fig. 3a–c**). We quantified resting ER [Ca^{2+}] by determining the maximum fluorescence intensity of R-CEPIA1er using ionomycin/Ca^{2+} (**Supplementary Fig. 6**). The resting ER [Ca^{2+}] was severely reduced in cells without Kir2.1 (**Fig. 3d**). The resting membrane potential of 5 BV cells was polarized in normal Krebs solution and became depolarized by high [K^+] solution containing 50 mM K^+ in the presence of ~145 mM intracellular K^+ (**Fig. 3e**). In contrast, resting membrane potential was already depolarized in the cells without Kir2.1 in normal Krebs solution. We hypothesized that spontaneous DICR activity under the depolarized conditions causes ER Ca^{2+} depletion. In support of this idea, further removal of β1a or Stac3, which completely abolishes DICR activity, restored the ER [Ca^{2+}] (**Fig. 3d**). Therefore, Kir2.1 is indispensable for our reconstituted DICR platform.

Evaluation of disease-causing mutations in Cav1.1

Using the high-throughput DICR platform, we evaluated mutations in Cav1.1 which

are implicated in various muscle diseases[12,39]. We initially tested known MH mutations,

R174W[40], R1086H[41], and T1354S[19] (**Fig. 4a**). We found that ER $[Ca^{2+}]$ was severely reduced

for R1086H (**Fig. 4b**) and that R1086H exhibited a smaller fluorescence change in response to

high $[K^+]$ (**Fig. 4c, d**) compared with the wild type (WT) or the other mutants. This was partly

caused by low ER $[Ca^{2+}]$, which reduces maximum fluorescence changes. In addition, R1086H

showed an enhanced K^+-dependence, i.e., lower EC_{50} value for K^+ (**Fig. 4c, e**). In contrast, no

or only minor changes in ER Ca^{2+} and K^+ dependence were observed for R174W and T1354S

cells (**Fig. 4b–e**). We anticipated that the reduction in ER $[Ca^{2+}]$ in R1086H cells was caused

by spontaneous activation of Cav1.1 at resting membrane potential because of its

hypersensitivity. To test the hypothesis, we hyperpolarized the membrane potential which

reduced the spontaneous activation of Cav1.1. Reduction of $[K^+]$ in the Krebs solution from 5

mM to 2.5 mM successfully hyperpolarized the membrane potential by ~15 mV

(**Supplementary Fig. 7a**). This substantially increased ER $[Ca^{2+}]$ in R1086H cells to a level

comparable to that in WT cells (**Supplementary Fig. 7b**). R1086H cells exhibited a clear

greater sensitivity to K^+ than WT cells (**Supplementary Fig. 7c–e**), supporting our hypothesis

of R1086H hyperactivation.

Previous functional characterization of these Cav1.1 mutants consistently showed

enhanced caffeine sensitivity, i.e., acceleration of Ca^{2+}-induced Ca^{2+} release $(CICR)^{[17–19]}$. We

tested caffeine dependence of these mutants in our platform. Caffeine released Ca^{2+} from WT

cells in a dose-dependent manner (**Fig. 4f**). R1086H cells also exhibited a caffeine dependence

but with reduced fluorescence change by 61 mM K^+ (**Fig. 4g**) and a significantly smaller EC_{50}

(**Fig. 4h**), indicating an enhanced caffeine sensitivity. In contrast, caffeine dependence was not

changed in R174W or T1354S cells (**Fig. 4f–h**).

We next examined the effect of myopathy-related mutations in Cav1.1 on DICR

activity. We chose four mutations, E100K, F275L, P742Q, and L1367V[42], none of which have

been functionally characterized (**Fig. 5a**). ER $[Ca^{2+}]$ was not changed by any of these mutations

(**Fig. 5b**). DICR activity was completely lost in the F275L mutant (**Fig. 5c, d**) and there was a

substantial rightward shift in K^+ dependence for P742Q (**Fig. 5c**); The EC_{50} value was threefold

higher in P742Q (33.2 ± 1.0 mM) compared with WT (10.3 ± 0.9 mM) (**Fig. 5e**). The other two

mutations (E100K and L1367V) had no significant effects on K^+ dependence (**Fig. 5c-e**). We

also tested the effects of these mutations on caffeine-induced Ca^{2+} release. No substantial

changes were observed with any of the mutations (**Fig. 5f–h**).

Effect of drugs on DICR activity

The high-throughput DICR platform is expected to be useful for screening drugs for muscle diseases. To test this possibility, we examined the effects of known DICR modulators. We initially tested three RyR1 inhibitors (**Fig. 6a**). Dantrolene is a well-known RyR1 inhibitor that is clinically used for MH[43]. Cpd1 is a novel potent RyR1-selective inhibitor that we recently developed[23,24]. Procaine inhibits CICR but not DICR in frog skeletal muscle[44,45]. Dantrolene (10 µM) and Cpd1 (3 µM) caused rightward shifts in K^+ dependence compared with the control, indicating that they inhibit DICR (**Fig. 6b, c**). This is consistent with the suppression of twitch and tetanic tension in isolated muscles and of reduced *in vivo* muscle weakness by dantrolene[46,47] and Cpd1[24]. Procaine (5 mM), in contrast, did not inhibit DICR (**Fig. 6b, c**). All three compounds shifted the caffeine dependence rightward, indicating inhibition of CICR (**Fig. 6d, e**). These results indicate that dantrolene and Cpd1 inhibit both DICR and CICR, whereas procaine is a CICR-selective inhibitor.

Lyotropic anions, such as perchlorate and thiocyanate, potentiate E-C coupling in skeletal muscle[48–52]. Perchlorate (10 mM) and thiocyanate (10 mM) significantly shifted the K^+ dependence leftward (**Fig. 7a, b**). In contrast, they did not affect caffeine dependence (**Fig. 7c, d**). These results indicate that lyotropic anions potentiate DICR but not CICR. This is consistent with previous findings showing that the potentiating effects of lyotropic anions are primarily caused by shifting voltage dependence of the charge movement toward more negative potentials[48–52]. Taken together, our high-throughput platform reproduces the effects of known DICR modulators, indicating that it can be used to screen for and test drugs that affect DICR.

Discussion

In this study, we established a high-through platform for DICR that was reconstituted in HEK293 cells expressing RyR1 and R-CEPIA1er. We made three key improvements to the original method by Perni et al. [25]. First, the essential components were transduced using VSV-G pseudotyped baculovirus, which can effectively infect a wide variety of mammalian cells without toxicity[28]. This greatly increased the transduction efficiency; almost all cells expressed the essential components (**Supplementary Fig. 1**). Second, depolarization of the plasma membrane was induced by high $[K^+]$ solution. This enabled us to simultaneously stimulate many cells (**Fig. 1, Supplementary Fig. 2** and **movie S1**). For this purpose, we expressed Kir2.1, an inward-rectifying potassium channel, which effectively hyperpolarized the membrane potential [32] (**Fig. 3**). Third, ER $[Ca^{2+}]$, instead of cytoplasmic $[Ca^{2+}]$, was measured

to detect DICR. This is critically important to avoid contamination of signals from extracellular

Ca^{2+} influx. Indeed, ER $[Ca^{2+}]$ was not changed by high $[K^+]$ depolarization in cells without

essential components (**Fig. 1a**). In contrast substantial increases in cytoplasmic $[Ca^{2+}]$ were

observed (**Supplementary Fig. 4a**). These improvements allow quantitative measurements of

DICR using a microplate reader.

In our reconstitution, five components, RyR1, Cav1.1, β1a, JP2 and Stac3, are essential for

DICR; removal of each component abolished or reduced the DICR activity (**Fig. 2a, b**). This is

consistent with previous findings from mice lacking each component[9,14–16,34]. In addition, the

effects of α_2-δ and γ_1 auxiliary subunits on DICR activity corresponded with findings of

previous reports [37,38] (**Fig. 2c, d**). Therefore, our reconstituted platform successfully reproduces

DICR in skeletal muscle. We found that a substantial DICR still occurred without JP2 (**Fig. 2a,**

b). This is in contrast to reports by Perni et al.[25,53], in which no voltage-gated Ca^{2+} release was

observed in reconstituted cells without junctophilins. A possible reason for this is differences

in expression methods. Our VSV-G baculovirus is more effective at transduction than standard

lipofection; therefore, sufficient proteins were expressed that might spontaneously interact with

each other without junctophilins.

We quantitatively evaluated the effect of disease-casing mutations or drugs using the EC_{50}

value for $[K^+]$. In our experimental conditions, the EC_{50} value for K^+ was about 10 mM (**Fig.**

1d). The value is lower than that for skeletal muscle[54] or myotubes[55] in high $[K^+]$-induced Ca^{2+}

transients. This may be partly explained by measurement differences. We calculated the EC_{50}

from the fluorescence intensity for 125–150 seconds after stimulation (**Fig. 1c**). This reduced

the EC_{50} by ~7 mM compared with the value obtained just after stimulation (**Supplementary**

Fig. 3). Nevertheless, our results reproduced the effects of disease-causing mutations (**Fig. 3–**

5) or drugs (**Fig. 6, 7**), strongly indicating that the EC_{50} value for K^+ is a valid parameter for

quantitative evaluation.

Formation of DICR machinery can cause peripheral localization of the components, e.g.,

RyR1 and Cav1.1, in the reconstituted cells[25,53]. In our reconstituted cells, however, no such

peripheral-specific localization was observed; the components were localized throughout cells

except for in nuclei (**Supplementary Fig. 1**). This may be because of high levels of the

components expressed by baculovirus, which not only accelerates formation of the DICR

machinery (as discussed above), but also increases the number of uncoupled components.

Mutations in Cav1.1 are implicated in various muscle diseases, including MH and

myopathy[12,39]. Among three MH mutations (R174W, R1086H and T1354S), we demonstrated

that R1086H, but not R174W orT1354S, significantly accelerated DICR (**Fig. 4c–e**). This is

consistent with previous findings using Cav1.1-deficient (dysgenic) myotubes [17–19]. All three

mutations show increased sensitivity to caffeine-induced Ca^{2+} release compared with WT,

indicating enhanced CICR activity [17–19]. However, we showed that caffeine sensitivity was

increased only for R1086H (**Fig. 4f–h**). A possible reason for this difference is the cells used.

We used stable RyR1-overexpressing cells. This may cause an excess amount of 'uncoupled'

RyR1, which might mask the enhanced sensitivity of caffeine dependence of the 'coupled' RyR1

by Cav1.1 mutations.

Most myopathy-related mutations identified in Cav1.1 have not been functionally

characterized. We evaluated four mutations (E100K, F275L, P742Q and L1367V) found in

patients with congenital myopathy[42] (**Fig. 5**). We demonstrated that P742Q substantially

inhibited DICR with a large rightward shifted K^+ dependence. This proline resides in the II-III

loop which interacts with RyR1 or Stac3[56]. Substitution of the proline to threonine, the residue

found in the corresponding position of CaV1.2, diminished DICR in dysgenic myotubes[57].

Therefore, P742Q may cause myopathy by reducing DICR activity. E100K and L1367V had

no or only minor impacts on DICR activity. These mutations might cause muscle weakness by

reasons other than reduced DICR. F275L is recessive because it was found in two compound

heterozygous sisters carrying F275L and the nonsense mutation, p.L791Cfs*37[42]. F275L

completely abolished DICR without any changes in ER $[Ca^{2+}]$ or caffeine sensitivity. This does

not fit with a recessive mutation; if this is the case, patients will have no DICR activity. Further

investigations are required for this mutation.

There are no specific treatments for most DICR-related diseases; therefore, there is an

urgent need to develop novel drugs for these diseases[11–13]. We here evaluated DICR modulators.

Among three known RyR1 inhibitors, dantrolene and Cpd1 suppressed DICR (**Fig. 6**). This is

consistent with the suppression of twitch and tetanic tensions of isolated muscles and of reduced

in vivo muscle weakness by dantrolene[46,47] and Cpd1[24]. Interestingly, procaine did not suppress

DICR (**Fig. 6**). Procaine inhibits CICR but not DICR in frog skeletal muscle[44] and reconstituted

frog α-RyR and β-RyR in RyR1-deficient myotubes[45]. Our data show similar procaine action in

a mammalian system. We also demonstrated that lyotropic anions (perchlorate and thiocyanate)

accelerate DICR (**Fig. 7**). This reproduces previous findings using skeletal muscle fibers[48–52].

Interestingly, lyotropic anions did not affect caffeine dependence, suggesting no effect on CICR

(**Fig. 7**). Therefore, our high-throughput platform will also be useful for drug testing and

screening for novel drugs that affect DICR.

In summary, we established a high-throughput platform for skeletal muscle DICR. This
platform will be useful for both evaluation of disease-causing mutations and the development
of novel drugs for DICR-related diseases.

Methods

Generation of stable inducible HEK293 cell lines

HEK293 cells stably expressing R-CEPIA1er and RyR1 were generated as described
previously[21,22]. Briefly, a full-length rabbit RyR1 cDNA was cloned in a tetracycline-induced
expression vector (pcDNA5/FRT/TO; Life Technologies, CA, USA)[58]. Flp-In T-REx HEK293
cells were co-transfected with this expression vector and pOG44 in accordance with the
manufacturer's instructions. Clones with suitable doxycycline-induced expression of RyR1
were selected and used for experiments. The cells were then transfected with an R-CEPIA1er
cDNA (a generous gift from Masamitsu Iino, The University of Tokyo) for stable expression of
R-CEPIA1er. Cells were cultured in Dulbecco's modified Eagle's medium supplemented with
10% fetal calf serum, 2 mM L-glutamine, 15 μg/ml blasticidin, and 100 μg/ml hygromycin.

Production of baculovirus

Baculovirus that expresses vesicular stomatitis virus G-protein (VSVG) on the virus envelope[28]
was produced using the Bac-to-Bac system (LifeTechnologies) as described previously[29].
cDNAs for Cav1.1, β1a, JP2, Stac3, and Kir2.1 were cloned from mouse skeletal muscle using
reverse transcription-PCR (primers pair are listed in **Supplementary Table 1**) and cloned into
the modified donor vector (pFastBac1-VSVG-CMV-WPRE). Baculovirus was produced in Sf9
cells according to the manufacturer's instructions (LifeTechnologies). Baculovirus titers were
immunologically determined using an anti-gp64 antibody (AcV5, Santa Cruz Biotechnology)[59].
Titers were $1–2×10^8$ pfu/mL for all viruses. The baculovirus solutions were stored in the dark
at 4°C and used within one month after production.

Immunocytochemistry

RyR1-HEK293 cells infected with baculovirus for the essential components (5 BV cells) were
fixed with ice-cold methanol and rinsed with phosphate-buffered saline (PBS). The cells were

blocked with 5% BSA/PBS for 30 min and incubated overnight with primary antibodies against

Cav1.1 (sc-514685, Santa Cruz Biotechnology), β1a (MA5-27714, Invitrogen), JP2 (sc-377086,

Santa Cruz Biotechnology), Stac3 (1G8, Abnova), or Kir2.1 (S112B-14, Invitrogen) in 1%

BSA/PBS. After washing three times with PBS, the cells were incubated for 2 h with Alexa594-

labeled goat anti-mouse IgG (Invitrogen) in 1% BSA/PBS. After washing with PBS, the cells

were observed using an all-in-one fluorescence microscope (BZ-X710, Keyence, Osaka, Japan).

Western blotting

RyR1-HEK293 cells with or without baculovirus infection for the essential components were

lysed with Pro-Prep protein extraction solution (iNtRON Biotechnology). The extracted

proteins were separated on 3–15% polyacrylamide gels and transferred to PVDF membranes.

Membranes were probed with the primary antibodies described above, followed by HRP-

labeled anti-mouse IgG (04-18-18, KPL). Positive bands were detected by chemiluminescence

using ImmunoStar LD (Fujifilm Wako Chemicals) as a substrate.

Single-cell Ca^{2+} imaging

RyR1/R-CEPPIA1er HEK293 cells grown on a glass-bottomed dish were treated with

doxycycline to induce the expression of RyR1 and infected with 5 BV 24 h prior to

measurement. Single-cell Ca^{2+} imaging was performed as described previously[29,60].

Measurements were performed at 26°C by perfusing solutions using an in-line solution

heater/cooler (Warner Instruments, Holliston, MA, USA). ER Ca^{2+} signals were determined by

R-CEPIA1er fluorescence excited at 561 nm using a 40× objective lens. Cells were perfused

with normal Krebs solution (140 mM NaCl, 5 mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 11 mM

glucose, 5 mM HEPES, pH 7.4) for 30 s, high [K^+] Krebs solution containing 50 mM K^+ for 50

s, and normal Krebs solution for 1.5 min. Fluorescence intensities in individual cells were

determined by region of interest (ROI) analysis using AquaCosmos software (Hamamatsu

Photonics, Hamamatsu, Japan).

Time lapse [Ca^{2+}]$_{ER}$ Measurements using a fluorometer

Time lapse [Ca^{2+}]$_{ER}$ measurements were performed using the FlexStation3 fluorometer

(Molecular Devices, San Jose, CA)[21,22]. Stable RyR1/R-CEPIA1er HEK293 cells were seeded

on 96-well flat-, clear-bottom black microplates (#3603, Corning, New York, NY) at a density

of $3×10^4$ cells/well in 100 μL culture medium. One day after seeding, 100 μL culture medium

containing 2 µg/mL doxycycline (for induction of RyR1) and baculovirus solutions for Cav1.1,

β1a, JP2, Stac3, and Kir2.1 (2 µL each) were added to each well. After 24 h, the culture medium

were replaced with 81 µL normal Krebs solution, and the microplate was placed in a

FlexStation3 fluorometer preincubated at 37°C. Signals from R-CEPIA1er, which was excited

at 560 nm and emitted at 610 nm, were captured every 5 seconds for 150 seconds. Thirty

seconds after starting, 54 µL of the high $[K^+]$ or caffeine solution was applied to the cells. The

fluorescence change was expressed as F/F_0 in which averaged fluorescence intensity of the last

25 seconds (F) was normalized to that of the initial 25 seconds (F_0).

For drug testing, compounds were added to normal Krebs solution at the indicated

concentrations. For determination of $[Ca^{2+}]_{ER}$, cells were incubated with 81 µL normal Krebs

solution, and 54 µL Krebs solution containing 50 mM Ca^{2+} and 50 µM ionomycin was added

30 sec after starting (final concentrations of Ca^{2+} and ionomycin were 20 mM and 20 µM,

respectively). R-CEPIA1er fluorescence was captured every 10 seconds for 6 min.

Measurement of resting membrane potential

The resting membrane potential of baculovirus-infected RyR1-HEK293 cells with or without

Kir2.1 expression was measured under current-clamp conditions in the whole cell configuration

of the patch-clamp method with an Axopatch 200B amplifier (Molecular Devices, San Jose,

CA). Pipettes were fabricated from borosilicate glass capillaries (Kimble) with a vertical pipette

puller (PP-83, Narishige, Tokyo, Japan). Pipettes had series resistance of 3–4 MΩ when filled

with the pipette solution containing (in mM) 120 D-glutamate, 20 KCl, 10 NaCl, 0.1 EGTA, 2

$MgCl_2$, 3 MgATP, and 5 Hepes (pH 7.4 adjusted with KOH). The series resistance was routinely

compensated by ~75% with the amplifier. Bath solution contained (in mM): 140 NaCl, 5 KCl,

11 glucose, 2 $CaCl_2$, 1 $MgCl_2$, and 10 Hepes (pH 7.4 adjusted with NaOH). High $[K^+]$ bath

solution contained (in mM): 95 NaCl, 50 KCl, 11 glucose, 2 $CaCl_2$, 1 $MgCl_2$, and 10 Hepes (pH

7.4 adjusted with NaOH). The resting membrane potential was measured immediately after

breaking into cells.

Statistics

Data are presented as means ± SD. Unpaired Student's t test was used for comparisons between

two groups. One-way analysis of variance (ANOVA), followed by Dunnett's test, was

performed to compare three or more groups. Two-tailed tests were used for all analyses.

Statistical analysis was performed using Prism v9 (GraphPad Software, Inc., La Jolla, USA).

Data availability

All data generated or analyzed in this study are available within the article and its Supplementary Information. All raw data supporting the findings from this study are available from the corresponding author upon request. Source data are provided with this paper.

References

1.   Franzini-Armstrong C. The relationship between form and function throughout the history of excitation-contraction coupling. J Gen Physiol 150, 189-210 (2018).

2.   Rios E, Pizarro G. Voltage sensor of excitation-contraction coupling in skeletal muscle. Physiol Rev 71, 849-908 (1991).

3.   Schneider MF. Control of calcium release in functioning skeletal muscle fibers. Annu Rev Physiol 56, 463-484 (1994).

4.   Shishmarev D. Excitation-contraction coupling in skeletal muscle: recent progress and unanswered questions. Biophys Rev 12, 143-153 (2020).

5.   Flucher BE, Campiglio M. STAC proteins: The missing link in skeletal muscle EC coupling and new regulators of calcium channel function. Biochim Biophys Acta Mol Cell Res 1866, 1101-1110 (2019).

6.   Horstick EJ, et al. Stac3 is a component of the excitation-contraction coupling machinery and mutated in Native American myopathy. Nat Commun 4, 1952 (2013).

7.   Nelson BR, et al. Skeletal muscle-specific T-tubule protein STAC3 mediates voltage-induced Ca2+ release and contractility. Proc Natl Acad Sci U S A 110, 11881-11886 (2013).

8.   Reinholt BM, Ge X, Cong X, Gerrard DE, Jiang H. Stac3 is a novel regulator of skeletal muscle development in mice. PLoS One 8, e62760 (2013).

9.   Takeshima H, Komazaki S, Nishi M, Iino M, Kangawa K. Junctophilins: a novel family of junctional membrane complex proteins. Mol Cell 6, 11-22 (2000).

10.   Rufenach B, Van Petegem F. Structure and function of STAC proteins: Calcium channel modulators and critical components of muscle excitation-contraction coupling. J Biol Chem 297, 100874 (2021).

11.   Agrawal A, Suryakumar G, Rathor R. Role of defective Ca(2+) signaling in skeletal muscle weakness: Pharmacological implications. J Cell Commun Signal 12, 645-659

(2018).

12. Flucher BE. Skeletal muscle CaV1.1 channelopathies. Pflugers Arch 472, 739-754

(2020).

13. Lawal TA*, et al.* Ryanodine receptor 1-related disorders: an historical perspective and

proposal for a unified nomenclature. Skelet Muscle 10, 32 (2020).

14. Tanabe T, Beam KG, Powell JA, Numa S. Restoration of excitation-contraction

coupling and slow calcium current in dysgenic muscle by dihydropyridine receptor

complementary DNA. Nature 336, 134-139 (1988).

15. Gregg RG*, et al.* Absence of the beta subunit (cchb1) of the skeletal muscle

dihydropyridine receptor alters expression of the alpha 1 subunit and eliminates

excitation-contraction coupling. Proc Natl Acad Sci U S A 93, 13961-13966 (1996).

16. Polster A, Nelson BR, Olson EN, Beam KG. Stac3 has a direct role in skeletal muscle-

type excitation-contraction coupling that is disrupted by a myopathy-causing mutation.

Proc Natl Acad Sci U S A 113, 10986-10991 (2016).

17. Weiss RG, O'Connell KM, Flucher BE, Allen PD, Grabner M, Dirksen RT. Functional

analysis of the R1086H malignant hyperthermia mutation in the DHPR reveals an

unexpected influence of the III-IV loop on skeletal muscle EC coupling. Am J Physiol

Cell Physiol 287, C1094-1102 (2004).

18. Eltit JM*, et al.* Malignant hyperthermia susceptibility arising from altered resting

coupling between the skeletal muscle L-type Ca2+ channel and the type 1 ryanodine

receptor. Proc Natl Acad Sci U S A 109, 7923-7928 (2012).

19. Pirone A*, et al.* Identification and functional characterization of malignant hyperthermia

mutation T1354S in the outer pore of the Cavalpha1S-subunit. Am J Physiol Cell

Physiol 299, C1345-1354 (2010).

20. Perez CF*, et al.* Functional and structural characterization of a novel malignant

hyperthermia-susceptible variant of DHPR-beta1a subunit (CACNB1). Am J Physiol

Cell Physiol 314, C323-C333 (2018).

21. Murayama T*, et al.* Efficient High-Throughput Screening by Endoplasmic Reticulum

Ca^{2+} Measurement to Identify Inhibitors of Ryanodine Receptor Ca^{2+}-Release Channels.

Mol Pharmacol 94, 722-730 (2018).

22. Murayama T, Kurebayashi N. Assays for Modulators of Ryanodine Receptor

(RyR)/Ca(2+) Release Channel Activity for Drug Discovery for Skeletal Muscle and

Heart Diseases. Curr Protoc Pharmacol 87, e71 (2019).

23. Mori S*, et al.* Structural development of a type-1 ryanodine receptor (RyR1) Ca(2+)-

release channel inhibitor guided by endoplasmic reticulum Ca(2+) assay. Eur J Med

Chem 179, 837-848 (2019).

24. Yamazawa T, *et al.* A novel RyR1-selective inhibitor prevents and rescues sudden death

in mouse models of malignant hyperthermia and heat stroke. Nat Commun 12, 4293

(2021).

25. Perni S, Lavorato M, Beam KG. De novo reconstitution reveals the proteins required

for skeletal muscle voltage-induced Ca(2+) release. Proc Natl Acad Sci U S A 114,

13822-13827 (2017).

26. Block BA, Imagawa T, Campbell KP, Franzini-Armstrong C. Structural evidence for

direct interaction between the molecular components of the transverse

tubule/sarcoplasmic reticulum junction in skeletal muscle. J Cell Biol 107, 2587-2600

(1988).

27. Suzuki J, Kanemaru K, Ishii K, Ohkura M, Okubo Y, Iino M. Imaging intraorganellar

Ca2+ at subcellular resolution using CEPIA. Nat Commun 5, 4153 (2014).

28. Tani H, Nishijima M, Ushijima H, Miyamura T, Matsuura Y. Characterization of cell-

surface determinants important for baculovirus infection. Virology 279, 343-353 (2001).

29. Uehara A, *et al.* Extensive Ca2+ leak through K4750Q cardiac ryanodine receptors

caused by cytosolic and luminal Ca2+ hypersensitivity. J Gen Physiol 149, 199-218

(2017).

30. Schredelseker J, Shrivastav M, Dayal A, Grabner M. Non-Ca2+-conducting Ca2+

channels in fish skeletal muscle excitation-contraction coupling. Proc Natl Acad Sci U

S A 107, 5658-5663 (2010).

31. DiFranco M, Yu C, Quinonez M, Vergara JL. Inward rectifier potassium currents in

mammalian skeletal muscle fibres. J Physiol 593, 1213-1238 (2015).

32. Kirkton RD, Bursac N. Engineering biosynthetic excitable tissues from unexcitable

cells for electrophysiological and cell therapy studies. Nat Commun 2, 300 (2011).

33. Zhao Y, *et al.* An expanded palette of genetically encoded Ca(2)(+) indicators. Science

333, 1888-1891 (2011).

34. Takeshima H, *et al.* Excitation-contraction uncoupling and muscular degeneration in

mice lacking functional skeletal muscle ryanodine-receptor gene. Nature 369, 556-559

(1994).

35. Catterall WA. Structure and regulation of voltage-gated Ca2+ channels. Annu Rev Cell

Dev Biol 16, 521-555 (2000).

36. Flucher BE, Obermair GJ, Tuluc P, Schredelseker J, Kern G, Grabner M. The role of

auxiliary dihydropyridine receptor subunits in muscle. J Muscle Res Cell Motil 26, 1-6

(2005).

37.   Obermair GJ, Kugler G, Baumgartner S, Tuluc P, Grabner M, Flucher BE. The Ca2+
channel alpha2delta-1 subunit determines Ca2+ current kinetics in skeletal muscle but
not targeting of alpha1S or excitation-contraction coupling. J Biol Chem 280, 2229-
2237 (2005).

38.   Ursu D, Schuhmeier RP, Freichel M, Flockerzi V, Melzer W. Altered inactivation of
Ca2+ current and Ca2+ release in mouse muscle fibers deficient in the DHP receptor
gamma1 subunit. J Gen Physiol 124, 605-618 (2004).

39.   Beam TA, Loudermilk EF, Kisor DF. Pharmacogenetics and pathophysiology of
CACNA1S mutations in malignant hyperthermia. Physiol Genomics 49, 81-87 (2017).

40.   Carpenter D, et al. The role of CACNA1S in predisposition to malignant hyperthermia.
BMC Med Genet 10, 104 (2009).

41.   Stewart SL, Hogan K, Rosenberg H, Fletcher JE. Identification of the Arg1086His
mutation in the alpha subunit of the voltage-dependent calcium channel (CACNA1S) in
a North American family with malignant hyperthermia. Clin Genet 59, 178-184 (2001).

42.   Schartner V, et al. Dihydropyridine receptor (DHPR, CACNA1S) congenital myopathy.
Acta Neuropathol 133, 517-533 (2017).

43.   Riazi S, Kraeva N, Hopkins PM. Updated guide for the management of malignant
hyperthermia. Can J Anaesth 65, 709-721 (2018).

44.   Thorens S, Endo M. Calcium-induced calcium release and "depolarization"-induced
calcium release: their physiological significance. Proc Jpn Acad 51, 473-478 (1975).

45.   Kashiyama T, Murayama T, Suzuki E, Allen PD, Ogawa Y. Frog alpha- and beta-
ryanodine receptors provide distinct intracellular Ca2+ signals in a myogenic cell line.
PLoS One 5, e11526 (2010).

46.   Leslie GC, Part NJ. The action of dantrolene sodium on rat fast and slow muscle in vivo.
Br J Pharmacol 72, 665-672 (1981).

47.   Meyler WJ, Mols-Thurkow I, Scaf AH, Sargo S, Wesseling H. The effect of dantrolene
sodium on rat skeletal muscle in relation to the plasma concentration. Eur J Pharmacol
53, 335-342 (1979).

48.   Luttgau HC, Gottschalk G, Kovacs L, Fuxreiter M. How perchlorate improves
excitation-contraction coupling in skeletal muscle fibers. Biophys J 43, 247-249 (1983).

49.   Delay M, Garcia DE, Sanchez JA. The effects of lyotropic anions on charge movement,
calcium currents and calcium signals in frog skeletal muscle fibres. J Physiol 425, 449-
469 (1990).

50.   Huang CL. The differential effects of twitch potentiators on charge movements in frog
skeletal muscle. J Physiol 380, 17-33 (1986).

51.   Csernoch L, Kovacs L, Szucs G. Perchlorate and the relationship between charge
movement and contractile activation in frog skeletal muscle fibres. J Physiol 390, 213-
227 (1987).

52.   Gonzalez A, Rios E. Perchlorate enhances transmission in skeletal muscle excitation-
contraction coupling. J Gen Physiol 102, 373-421 (1993).

53.   Perni S, Beam K. Junctophilins 1, 2, and 3 all support voltage-induced Ca2+ release
despite considerable divergence. J Gen Physiol 154,   (2022).

54.   Caputo C, Bolanos P. Fluo-3 signals associated with potassium contractures in single
amphibian muscle fibres. J Physiol 481 (Pt 1), 119-128 (1994).

55.   Yang T*, et al.* Pharmacologic and functional characterization of malignant hyperthermia
in the R163C RyR1 knock-in mouse. Anesthesiology 105, 1164-1175 (2006).

56.   Wong King Yuen SM, Campiglio M, Tung CC, Flucher BE, Van Petegem F. Structural
insights into binding of STAC proteins to voltage-gated calcium channels. Proc Natl
Acad Sci U S A 114, E9520-E9528 (2017).

57.   Kugler G, Weiss RG, Flucher BE, Grabner M. Structural requirements of the
dihydropyridine receptor alpha1S II-III loop for skeletal-type excitation-contraction
coupling. J Biol Chem 279, 4721-4728 (2004).

58.   Tong J, Oyamada H, Demaurex N, Grinstein S, McCarthy TV, MacLennan DH.
Caffeine and halothane sensitivity of intracellular Ca2+ release is altered by 15 calcium
release channel (ryanodine receptor) mutations associated with malignant hyperthermia
and/or central core disease. J Biol Chem 272, 26332-26339 (1997).

59.   Kitts PA, Green G. An immunological assay for determination of baculovirus titers in
48 hours. Anal Biochem 268, 173-178 (1999).

60.   Murayama T*, et al.* Divergent Activity Profiles of Type 1 Ryanodine Receptor Channels
Carrying Malignant Hyperthermia and Central Core Disease Mutations in the Amino-
Terminal Region. PLoS One 10, e0130606 (2015).

Acknowledgements

We thank Ikue Hiraga and the Laboratory of Proteomics and Biomolecular Science, Biomedical
Research Core Facilities, Juntendo University Graduate School of Medicine, for technical
assistance. We thank Jeremy Allen, PhD, from Edanz (https://jp.edanz.com/ac) for editing a
draft of this manuscript.

Author information

Affiliations

Department of Pharmacology, Juntendo University School of Medicine, Tokyo, Japan

Takashi Murayama, Nagomi Kurebayashi, Takuya Kobayashi, Satoru Okazaki, Kyosuke Yamashiro and Takashi Sakurai

Department of Molecular Pharmacology, Shinshu University School of Medicine, Matsumoto, Japan.

Takuro Numaga-Tomita, Tsutomu Nakada, and Mitsuhiko Yamada

Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, Tokyo, Japan.

Shuichi Mori, Ryosuke Ishida, and Hiroyuki Kagechika

Contributions

T.M. and N.K. conceived and designed the work. T.M., N.K., T.N.-T., S.O., K.Y., S.M., and R.I. acquired and analyzed data. T.M., N.K., T.N.-T., T.K., S.O., K.Y., T.N., S.M., R.I., K.H., M.Y., T.S. interpreted data. T.M. and N.K. wrote the manuscript. All authors reviewed and approved the manuscript.

Corresponding authors

Correspondence to Takashi Murayama (takashim@juntendo.ac.jp).

Ethics declarations

Competing interests

The authors declare no competing interests.

Figure legends

Fig. 1. DICR measurements in 96-well plates using a fluorescence microplate reader. a
Schematic drawing of RyR1/R-CEPIA1er HEK293 cells without (No BV, left) and with (5 BV,
right) baculovirus infection of essential components for DICR (Cav1.1, β1a, JP2, Stac3) and
Kir2.1. **b**, **c** Typical results of time-lapse R-CEPIA1er fluorescence measurement using a
FlexStation3 fluorescence microplate reader for RyR1/R-CEPIA1er HEK293 cells without (**b**,
No BV) or with (**c**, 5 BV) BV infection. High [K$^+$] solution ranging from 5–61 mM (symbols
shown in **c**) or 10 mM caffeine (light blue) was applied at 30 seconds after starting (blue
arrowheads). **d** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in No BV (black) and 5
BV cells (red). F/F$_0$ was obtained by normalizing the averaged fluorescence of the last 25
seconds (F) to that of the first 25 seconds (F$_0$). Data are means $\pm$ SD (n = 6). **e** Fluorescence
change by 61 mM K$^+$ or 10 mM caffeine. Fluorescence change is expressed as 1-F/F$_0$ (see panel
d). Data are means $\pm$ SD (n = 6) and were analyzed by two-way ANOVA with Bonferroni's
multiple comparisons test. Note that 5 BV cells show large fluorescence change with high [K$^+$]
which is nearly 80% of that with caffeine.

Fig. 2. Validation of the reconstituted DICR platform. a K$^+$ dependence of R-CEPIA1er
fluorescence (F/F$_0$) in 5 BV (black), -Cav1.1 (red), and -JP2 (dark blue). Data are means $\pm$ SD
(n = 9 for 5 BV, 6 for -Cav1.1 and -JP2). **b** Fluorescence change by 61 mM K$^+$ in the BV-
infected cells lacking one of essential components (-Cav1.1, -β1a, -Stac3, or -JP2) or with RyR2
cells instead of RyR1. Data are means $\pm$ SD (n = 6) and were analyzed by one-way ANOVA
with Dunnett's multiple comparisons test. **c** [K$^+$] dependence of R-CEPIA1er fluorescence
(F/F$_0$) in cells infected with 5 BV (black), 5 BV plus α$_2$-δ (+α$_2$-δ, purple) and 5 BV plus γ1 (+γ1,
orange). Data are means $\pm$ SD (n = 6). **d** Fluorescence change by 61 mM K$^+$ in 5 BV, +α$_2$-δ,
and +γ1 cells. Data are means $\pm$ SD (n = 6) and were analyzed by one-way ANOVA with
Dunnett's multiple comparisons test.

Fig. 3. Effects of Kir2.1 on DICR activity. a, b Typical results of R-CEPIA1er fluorescence
in cells infected with 5 BV (**a**) and BV without Kir2.1 (-Kir2.1) (**b**). High [K$^+$] solution ranging
from 5–61 mM (symbols shown in **a**) was applied at 30 seconds after starting (blue arrowheads).
c [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells infected 5 BV (black) and BV

without Kir2.1 (-Kir2.1) (pink). Data are means ± SD (n = 6). **d** Resting ER $[Ca^{2+}]$ in cells

infected with 5 BV (5 BV), BV without Kir2.1 (-Kir2.1), Kir2.1 and β1a (-Kir2.1, -β1a), or

Kir2.1 and Stac3 (-Kir2.1, -Stac3). Data are means ± SD (n = 8) and were analyzed by one-way

ANOVA with Dunnett's multiple comparisons test. **e** Resting membrane potential of cells

infected with 5 BV (left) or BV without Kir2.1 (-Kir2.1) (right). Data are means ± SD (n = 10

and 16 for 5 mM and 50 mM K^+, respectively, for 5 BV and 10 and 14 for 5 mM and 50 mM

K^+, respectively, for -Kir2.1) and were analyzed by two-way ANOVA with Bonferroni's

multiple comparisons test.

Fig. 4. Effect of MH mutations in Cav1.1 on DICR activity. a Schematic drawing of the

location of MH mutations (R174W, R1086H and T1354S) in Cav1.1. **b** Resting ER $[Ca^{2+}]$ in

cells expressing WT and mutant Cav1.1. Data are means ± SD (n = 12) and were analyzed by

one-way ANOVA with Dunnett's multiple comparisons test. **c** $[K^+]$ dependence of R-CEPIA1er

fluorescence (F/F_0) in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1

cells. Data are means ± SD (n = 9). **d, e** Fluorescence change by 61 mM K^+ (**d**) and EC_{50} values

for K^+ (**e**). Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's

multiple comparisons test. **f** Caffeine dependence of v(F/F_0) in WT (black), R174W (red),

R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means ± SD (n = 9). **g, h**

Fluorescence change by 20 mM caffeine (**g**) and EC_{50} values for caffeine (**h**). Data are means

± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 5. Effect of myopathy mutations in Cav1.1 on DICR activity. a Schematic drawing of

the location of myopathy mutations (E100K, F275L, P742Q and L1367V) in Cav1.1. **b** Resting

ER $[Ca^{2+}]$ in cells expressing WT and mutant Cav1.1. Data are means ± SD (n = 12) and were

analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **c** $[K^+]$ dependence of

R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green)

and L1367V (magenta) Cav1.1 cells. Data are means ± SD (n = 9). **d, e** Fluorescence change

by 61 mM K^+ (**d**) and EC_{50} values for K^+ (**e**). Data are means ± SD (n = 9) and were analyzed

by one-way ANOVA with Dunnett's multiple comparisons test. **f** Caffeine dependence of R-

CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and

L1367V (magenta) Cav1.1 cells. Data are means ± SD (n = 9). **g, h** Fluorescence change by 20

mM caffeine (**g**) and EC_{50} values for caffeine (**h**). Data are means ± SD (n = 9) and were

analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 6. Effect of RyR1 inhibitors on DICR activity. a Chemical structures of dantrolene, Cpd1

and procaine. **b** [K$^+$] dependence of fluorescence change (F/F$_0$) in cells without (Control, black)

or with dantrolene (10 μM, red), Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are means

± SD (n = 9). **c** EC$_{50}$ values for K$^+$. Data are means ± SD (n = 9) and were analyzed by one-

way ANOVA with Dunnett's multiple comparisons test. **d** Caffeine dependence of R-CEPIA1er

fluorescence (F/F$_0$) in cells without (Control, black) or with dantrolene (10 μM, red), Cpd1 (3

μM, blue) or procaine (5 mM, green). Data are means ± SD (n = 9). **e** EC$_{50}$ values for caffeine.

Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple

comparisons test.

Fig. 7. Effect of lyotropic anions on DICR activity. a [K$^+$] dependence of fluorescence change

(F/F$_0$) in cells in the absence (Control, black) or presence of perchlorate (10 mM, red) or

thiocyanate (10 mM, blue). Data are means ± SD (n = 9). **b** EC$_{50}$ values for K$^+$. Data are means

± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

c Caffeine dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells in the absence (Control,

black) or presence of perchlorate (10 mM, red) or thiocyanate (10 mM, blue). Data are means

± SD (n = 9). **d** EC$_{50}$ values for caffeine. Data are means ± SD (n = 9) and were analyzed by

one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 1

Fig. 1. DICR measurements in 96-well plates using a fluorescence microplate reader. a Schematic drawing of RyR1/R-CEPIA1er HEK293 cells without (No BV, left) and with (5 BV, right) baculovirus infection of essential components for DICR (Cav1.1, β1a, JP2, Stac3) and Kir2.1. **b**, **c** Typical results of time-lapse R-CEPIA1er fluorescence measurement using a FlexStation3 fluorescence microplate reader for RyR1/R-CEPIA1er HEK293 cells without (**b**, No BV) or with (**c**, 5 BV) BV infection. High [K$^+$] solution ranging from 5–61 mM (symbols shown in **c**) or 10 mM caffeine (light blue) was applied at 30 seconds after starting (blue arrowheads). **d** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in No BV (black) and 5 BV cells (red). F/F$_0$ was obtained by normalizing the averaged fluorescence of the last 25 seconds (F) to that of the first 25 seconds (F$_0$). Data are means $\pm$ SD (n = 6). **e** Fluorescence change by 61 mM K$^+$ or 10 mM caffeine. Fluorescence change is expressed as 1-F/F$_0$ (see panel **d**). Data are means $\pm$ SD (n = 6) and were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test. Note that 5 BV cells show large fluorescence change with high [K$^+$] which is nearly 80% of that with caffeine.

Fig. 2

Fig. 2. Validation of the reconstituted DICR platform. a K$^+$ dependence of R-CEPIA1er fluorescence (F/F$_0$) in 5 BV (black), -Cav1.1 (red), and -JP2 (dark blue). Data are means $\pm$ SD (n = 9 for 5 BV, 6 for -Cav1.1 and -JP2). **b** Fluorescence change by 61 mM K$^+$ in the BV-infected cells lacking one of essential components (-Cav1.1, -β1a, -Stac3, or -JP2) or with RyR2 cells instead of RyR1. Data are means $\pm$ SD (n = 6) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **c** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells infected with 5 BV (black), 5 BV plus α_2-δ (+α_2-δ, purple) and 5 BV plus γ1 (+γ1, orange). Data are means $\pm$ SD (n = 6). **d** Fluorescence change by 61 mM K$^+$ in 5 BV, +α_2-δ, and +γ1 cells. Data are means $\pm$ SD (n = 6) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 3

Fig. 3. Effects of Kir2.1 on DICR activity. a, b Typical results of R-CEPIA1er fluorescence in cells infected with 5 BV (**a**) and BV without Kir2.1 (-Kir2.1) (**b**). High [K$^+$] solution ranging from 5–61 mM (symbols shown in **a**) was applied at 30 seconds after starting (blue arrowheads). **c** [K$^+$] dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells infected 5 BV (black) and BV without Kir2.1 (-Kir2.1) (pink). Data are means $\pm$ SD (n = 6). **d** Resting ER [Ca^{2+}] in cells infected with 5 BV (5 BV), BV without Kir2.1 (-Kir2.1), Kir2.1 and β1a (-Kir2.1, -β1a), or Kir2.1 and Stac3 (-Kir2.1, -Stac3). Data are means $\pm$ SD (n = 8) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **e** Resting membrane potential of cells infected with 5 BV (left) or BV without Kir2.1 (-Kir2.1) (right). Data are means $\pm$ SD (n = 10 and 16 for 5 mM and 50 mM K$^+$, respectively, for 5 BV and 10 and 14 for 5 mM and 50 mM K$^+$, respectively, for -Kir2.1) and were analyzed by two-way ANOVA with Bonferroni's multiple comparisons test.

Fig. 4

Fig. 4. **Effect of MH mutations in Cav1.1 on DICR activity. a** Schematic drawing of the location of MH mutations (R174W, R1086H and T1354S) in Cav1.1. **b** Resting ER [Ca^{2+}] in cells expressing WT and mutant Cav1.1. Data are means ± SD (n = 12) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **c** [K^+] dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means ± SD (n = 9). **d, e** Fluorescence change by 61 mM K^+ (**d**) and EC_{50} values for K^+ (**e**). Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **f** Caffeine dependence of v(F/F_0) in WT (black), R174W (red), R1086H (blue) and T1354S (green) Cav1.1 cells. Data are means ± SD (n = 9). **g, h** Fluorescence change by 20 mM caffeine (**g**) and EC_{50} values for caffeine (**h**). Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 5

Fig. 5. Effect of myopathy mutations in Cav1.1 on DICR activity. a Schematic drawing of the location of myopathy mutations (E100K, F275L, P742Q and L1367V) in Cav1.1. **b** Resting ER [Ca^{2+}] in cells expressing WT and mutant Cav1.1. Data are means $\pm$ SD (n = 12) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **c** [K^+] dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means $\pm$ SD (n = 9). **d, e** Fluorescence change by 61 mM K^+ (**d**) and EC_{50} values for K^+ (**e**). Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **f** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in WT (black), E100K (red), F275L (blue), P742Q (green) and L1367V (magenta) Cav1.1 cells. Data are means $\pm$ SD (n = 9). **g, h** Fluorescence change by 20 mM caffeine (**g**) and EC_{50} values for caffeine (**h**). Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 6

Fig. 6. Effect of RyR1 inhibitors on DICR activity. a Chemical structures of dantrolene, Cpd1 and procaine. **b** [K$^+$] dependence of fluorescence change (F/F$_0$) in cells without (Control, black) or with dantrolene (10 μM, red), Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are means ± SD (n = 9). **c** EC$_{50}$ values for K$^+$. Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **d** Caffeine dependence of R-CEPIA1er fluorescence (F/F$_0$) in cells without (Control, black) or with dantrolene (10 μM, red), Cpd1 (3 μM, blue) or procaine (5 mM, green). Data are means ± SD (n = 9). **e** EC$_{50}$ values for caffeine. Data are means ± SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.

Fig. 7

Fig. 7. Effect of lyotropic anions on DICR activity. a [K^+] dependence of fluorescence change (F/F_0) in cells in the absence (Control, black) or presence of perchlorate (10 mM, red) or thiocyanate (10 mM, blue). Data are means $\pm$ SD (n = 9). **b** EC_{50} values for K^+. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. **c** Caffeine dependence of R-CEPIA1er fluorescence (F/F_0) in cells in the absence (Control, black) or presence of perchlorate (10 mM, red) or thiocyanate (10 mM, blue). Data are means $\pm$ SD (n = 9). **d** EC_{50} values for caffeine. Data are means $\pm$ SD (n = 9) and were analyzed by one-way ANOVA with Dunnett's multiple comparisons test.