1 **Decision-making processes in perceptual learning depend on effectors**

2 Vladyslav Ivanov[1]*, Giorgio Manenti[1,2]*, Sandrin S. Sudmann[1,2,3], Igor Kagan[2,4], Caspar M.

3 Schwiedrzik[1,2,3†]

[1]Neural Circuits and Cognition Lab, European Neuroscience Institute Göttingen – A Joint Initiative of the University Medical Center Göttingen and the Max Planck Society, Grisebachstraße 5, 37077 Göttingen, Germany; [2]Leibniz ScienceCampus Primate Cognition, Göttingen, Germany; [3]Perception and Plasticity Group, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg 4, 37077 Göttingen, Germany; [4]Decision and Awareness Group, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg 4, 37077 Göttingen, Germany

*These authors contributed equally to this work

†To whom correspondence should be addressed: c.schwiedrzik@eni-g.de

Abstract

Visual perceptual learning is traditionally thought to arise in visual cortex. However, typical perceptual

learning tasks also involve systematic mapping of visual information onto motor actions. Because the

motor system contains both effector-specific and effector-unspecific representations, the question

arises whether visual perceptual learning is effector-specific itself, or not. Here, we study this question

in an orientation discrimination task. Subjects learn to indicate their choices either with joystick

movements or with manual reaches. After training, we challenge them to perform the same task with

eye movements. We dissect the decision-making process using the drift diffusion model. We find that

learning effects on the rate of evidence accumulation and the decision criteria depend on effectors,

albeit not fully. This suggests that during perceptual learning, visual information is mapped onto

effector-specific integrators. Overlap of the populations of neurons encoding motor plans for these

effectors may explain partial generalization. Taken together, visual perceptual learning is not limited

to visual cortex, but also affects sensorimotor mapping at the interface of visual processing and

decision making.

Keywords: visual perceptual learning, effector specificity, orientation, sensorimotor mapping

Introduction

A core characteristic of the brain is its ability to learn. Learning is not limited to acquiring new motor skills or a foreign language – visual perception is also highly plastic. Training enables us to improve our visual abilities well into the hyperacuity range (Poggio, Fahle, & Edelman, 1992) and to learn to see what is initially invisible (Schwiedrzik, Singer, & Melloni, 2011). Such training effects are known as visual perceptual learning (VPL). VPL occurs not only during critical periods of development, but also in adult brains. This has profound implications for how we think about brain organization and function, yet little is known about the principles and neural mechanisms underlying adult VPL.

Classical theories suggest that VPL occurs in visual cortex (Fahle, 2004; Sagi, 2011). This idea rests upon the striking correspondence between the known selectivities of neurons in early visual areas and the selectivity of behavioral learning effects that are often found in VPL studies. For example, VPL has been found to be specific to the orientation, location (Schoups, Vogels, & Orban, 1995), polarity (Grieco, Casco, & Roncato, 2006), and even eye of origin (Karni & Sagi, 1991) of the training material, resembling the tuning properties of V1 neurons. Accordingly, classical theories argue that VPL increases the selectivity of neurons in early sensory areas.

While VPL-induced increased selectivity of neurons in early visual cortex has indeed been observed through single cell recordings in monkeys (reviewed in Vogels, 2010), results are somewhat inconsistent across studies (Ghose, Yang, & Maunsell, 2002), the observed changes in selectivity are too small to explain the behavioral benefits (Petrov, Dosher, & Lu, 2005), and modelling points to limitations on how much information can be gained by sharpening tuning curves (Series, Latham, & Pouget, 2004). These results suggest that sharpening of neural tuning in visual cortex is but one mechanism of VPL.

More recent studies have started to investigate a previously neglected aspect of VPL, sensorimotor mapping. In a typical VPL task, two stimulus alternatives, e.g., clockwise or counterclockwise orientation of lines, have to be mapped onto two specific motor actions, e.g., a leftward or rightward saccade. This implies that visual information needs to be brought together with motor plans that correspond to the trainee's choice. This suggests selective strengthening of connections between visual and motor neurons as another candidate mechanism of VPL. Indeed, a groundbreaking study found that VPL changes the activity of neurons in the lateral intraparietal area (LIP), an area in posterior parietal cortex that is involved in oculomotor decision-making and where sensorimotor mappings can take place (Law & Gold, 2008). This resonates with theories that frame VPL as a change in the readout of visual information by decision-making areas (Dosher & Lu, 2017). A role of motor areas in VPL is further suggested by several behavioral findings, e.g., that motor tasks improve VPL consolidation

(Klorfeld-Auslender & Censor, 2019), and, reversely, that VPL can improve oculomotor performance

(Szpiro, Spering, & Carrasco, 2014).

If VPL affects sensory-motor mapping between visual cortex and the motor system, this would suggest

another form of specificity in VPL, namely effector specificity. This is because the motor system

contains neurons that code motor plans for specific effectors. These are highly prevalent in the primary

motor cortex, but also in parietal cortex: many neurons in the so-called parietal reach region (PRR)

code for arm movements, while many neurons in LIP are saccade specific (Christopoulos, Bonaiuto,

Kagan, & Andersen, 2015; Christopoulos, Kagan, & Andersen, 2018; de Lafuente, Jazayeri, & Shadlen,

2015; Dickinson, Calton, & Snyder, 2003). During VPL, connections between stimulus-specific visual

neurons and effector-specific motor neurons could thus be strengthened, resulting in stimulus- and

effector-specific behavioral learning effects. However, areas like LIP also encode components that are

effector-independent (Chang, Calton, Lawrence, Dickinson, & Snyder, 2016; Christopoulos et al., 2018;

Colby & Goldberg, 1999; Gottlieb & Goldberg, 1999; Snyder, Batista, & Andersen, 1997). This raises the

possibility that VPL is effector-independent, too, even if sensorimotor mapping is involved.

Figure 1: Possible models for effector specificity in visual perceptual learning. *Classical model: learning occurs in visual cortex and results in stimulus- but not effector-specific learning effects. Changing the effector after training results in full transfer of learning effects to the new effector. Specific stimulus-response mapping model: visual information is mapped onto effector-specific motor neurons. This would result in stimulus- and effector-specific learning effects. Decision rule model: stimulus-response mapping takes place between visual neurons and effector-independent neurons involved in decision making and movement planning. This results in stimulus-specific learning effects that generalize across effectors. Effector-specific decision rule model: Sensorimotor mapping occurs between visual neurons and effector-specific integrators. Overlap between effectors at the integration stage results in partial transfer between the trained and untrained effectors.*

A number of studies have started to investigate effector-specificity of VPL, but results are mixed,

leaving open several possible models on how sensory and/or motor circuits are affected by training

(Fig. 1): in the above-mentioned *classical model* of VPL, learning occurs in visual cortex and results in

stimulus- but not effector-specific learning effects. Hence, changing the effector after training would

result in full transfer of learning effects to the new effector. Studies that report effector-independent

VPL effects can be interpreted this way (Awada, Bakhtiari, & Pack, 2021). In the *specific stimulus-*

response mapping model, visual information is mapped onto effector-specific motor neurons in motor

cortex (Kazennikov et al., 1999), parietal cortex (Christopoulos et al., 2018; Dickinson et al., 2003), or

subcortical structures (B. H. Liu, Huberman, & Scanziani, 2016). This would result in stimulus- and

effector-specific learning effects. Evidence for high effector-specificity VPL has been reported in

several studies (C. S. Green, Kattner, Siegel, Kersten, & Schrater, 2015; Grzeczkowski, Cretenoud,

Herzog, & Mast, 2017). However, these experiments often also involved a change in the task (e.g.,

binary categorization versus continuous adjustment), which possibly requires different readout

weights. Hence, these results could also be interpreted to reflect the well-known task-specificity of VPL

(Fahle & Morgan, 1996) instead of effector-specificity. Most recently, a *decision rule model* has been

put forward (Awada et al., 2021). Here, stimulus-response mapping takes place between visual

neurons and effector-independent neurons involved in abstract movement planning and decision

making in parietal cortex or subcortical structures like the pulvinar (Wilke, Turchi, Smith, Mishkin, &

Leopold, 2010). This would result in stimulus-specific learning effects that generalize well across

effectors. This has indeed been reported (Awada et al., 2021).

The studies investigating effector-specificity in VPL have so far exclusively focused on accuracy as a

measure of learning and transfer. However, this only considers the end result of a decision-making

process and disregards subcomponents involved in the full sensorimotor processing arc. Here, we use

a model-based approach, the well-established drift diffusion model (DDM) of decision making (Ratcliff,

1978), to dissociate different aspects of this arc and learning effects upon them. The DDM relates

accuracy to reaction times and allows investigation of the rate of evidence accumulation (the drift rate

parameter), the decision bound (i.e., when a decision is reached), non-decision time (related to

encoding of sensory information and execution of motor actions), and decision bias for one or the

other stimulus alternative. This approach has previously been shown to reveal subcomponents of VPL

(Dutilh, Krypotos, & Wagenmakers, 2011; C. C. Liu & Watanabe, 2012; Petrov, Van Horn, & Ratcliff,

2011; Zhang & Rowe, 2014), but has so far not been applied to the question of effector-specificity in

VPL.

We train subjects in an orientation discrimination task. We focus on orientation VPL since stimulus-

specific VPL effects are well documented for this feature (Schoups et al., 1995). We train subjects for

several days to discriminate orientations and to indicate their choice with either a joystick movement

(Experiment 1) or a reach (Experiment 2). After the last training session, we require them to carry out

the same task but to indicate their choice with an eye movement. Our transfer task thus differs from

training only in the effectors, allowing us to dissociate the above-mentioned models.

To preview, across two experiments, we find that VPL depends on the effector. The DDM taking

reaction times into account shows that the specificity arises in the rate of sensory evidence

accumulation (i.e., the drift rate) and the decision bound, which reflects the stopping criterion for

choice, two integral components of decision-making. This is most compatible with a new model, where

visual information is mapped onto effector-specific integrators.

Methods

Participants

A total of 30 healthy human volunteers (18 female, 1 left-handed, mean age 24.97 yrs, *SD* 6.23 yrs) participated in this study. All subjects had normal or corrected-to-normal vision, reported no history of neurological or psychiatric disease, and gave written informed consent before participation. No sample size estimate was performed, but sample size was selected based on previous studies. Because all comparisons were within subjects, we used convenience sampling. 22 subjects (12 female, 1 left-handed, mean age 23.5 yrs, *SD* 2.32 yrs) participated in Experiment 1 involving joystick and eye movements. 3 subjects had to be excluded from data analysis because lack of significant learning (final *n*=19, 12 female, 1 left-handed, mean age 23.68 yrs, *SD* 2.4 yrs). 8 subjects participated in Experiment 2 involving reaches on a touchscreen and eye movements (*n*=8, 6 female, mean age 29 yrs, *SD* 10.92 yrs). 2 subjects were excluded from data analysis because lack of significant learning (final *n*=6, 5 female, mean age 25.67 yrs, *SD* 3.33 yrs). All procedures were in accordance with the Declaration of Helsinki and approved by the Ethics Committee of the University Medical Center Göttingen (protocol number 29/8/17).

Stimuli and procedure

In all experiments, subjects were trained on an orientation discrimination task. Training took place over four to six consecutive days with one training session per day. On the day after the last training session, subjects were instructed to perform the same task, but with a different effector (see below). Stimuli were presented on an LCD monitor (ViewPixx EEG, refresh rate 120 Hz, resolution 1920 × 1080 pixel, viewing distance 65 cm) in a darkened, sound-attenuating booth (Desone Modular Acoustics). Stimulus delivery and response collection were controlled using Psychtoolbox (Brainard, 1997) running in Matlab (The Mathworks, Inc.) on Windows 10. During all experiments, we continuously acquired pupil and gaze measurements using a high-speed, video-based eye tracker (SR Research Eyelink 1000+). Data were sampled at 1000 Hz from both eyes. Subjects were paid €8 per hour. To assure constant motivation over the training sessions, subjects received a bonus of €2 if they improved by 10% from the previous training session.

Experiment 1

In Experiment 1, subjects were trained in orientation discrimination and responded with a joystick movement during training, while they had to respond with a saccade in the transfer session. Subjects had to decide whether a Gabor grating (size 2.4 dva, spatial frequency 3.3 cpd, luminance 43.4 cd/m^2)

was tilted clockwise or counterclockwise with respect to the reference stimulus. The reference stimulus was a Gabor grating with identical parameters tilted 20° counterclockwise from the vertical meridian. The task stimuli were presented in six logarithmically spaced difficulty levels (±Δ = [0°, 0.31°, 0.56°, 1°, 1.77°, 3.16°]) with 20 trials for each of the orientations, respectively. The stimuli were presented in pseudo-random order at 8.65° eccentricity against a grey background (39.5 cd/m^2).

Each trial started with a fixation phase of 2-3 s duration. Then, placeholders for the saccade version of the task appeared for 1.5 s, followed by the presentation of the Gabor grating for 250 ms. Subjects then had 1 s to respond. The start of the response period was indicated by the disappearance of the Gabor grating. Feedback on accuracy was presented by changing the color of the fixation dot to red (incorrect) or green (correct) for 0.5 s. The next trial started 1.5 s later. If subjects did not respond within 1 s, a message "too late" appeared on the screen and the trial was repeated later during the block. Similarly, if subjects broke fixation throughout the trial, the trial was aborted and repeated at the end of the block.

Each training session included 220 trials evenly distributed among four blocks. Subjects were free to take short breaks between blocks. There were additional breaks in the middle of each block to display feedback about performance and reaction times. The reference was shown for 2 s in the center of the screen before and in the middle of each block (in total 8 times per session).

Subjects were instructed to respond as accurately as possible, however, they also had to consider the time limitation. To respond, subjects had to tilt a joystick (Logitech Extreme 3D Pro, either in standard configuration or in a customized configuration for left-handed subjects) with their dominant hand to the left if a stimulus was counterclockwise rotated and to the right if clockwise (while maintaining fixation). Hand dominance was assessed at the beginning of the experiment with the Edinburgh Handedness Inventory (Oldfield, 1971). In the transfer session, subjects were instructed to instead direct their gaze to the left or right placeholder, respectively. Previous research has shown that effectors do not affect sensitivity pre training (Jaskowski & Sobieralska, 2004; Szul, Bompas, Sumner, & Zhang, 2020; Szumska, van der Lubbe, Grzeczkowski, & Herzog, 2016). We focus on effector-specificity since stimulus-specificity does not differentiate the different models for sensorimotor mapping in VPL.

Participant had to reach at least 75% correct responses before we transitioned to the transfer session. All but two subjects reached this performance level within four days; two subjects required five and six days of training, respectively.

<u>Experiment 2</u>

In Experiment 2, subjects were trained on orientation discrimination and responded with a finger

movement during training, while they had to respond with a saccade in the transfer session. We

recorded finger movements by means of a transparent touch screen (Magic Touch, Tyco Touch Inc.),

modelled after Pilacinsky and Lindner (2019), which was mounted in front of the screen. Subjects were

provided with an arm support (ErgoRest) and a finger stylus (TrueTip) to ensure ease of use.

As in Experiment 1, subjects had to decide whether a Gabor grating (size 2.4 dva, spatial frequency 3.3

cpd, luminance 10.9 cd/m^2) was tilted clockwise or counterclockwise with respect to the reference

stimulus. The reference stimulus was a Gabor grating with identical parameters tilted 20°

counterclockwise from the vertical meridian. The task stimuli were presented in six logarithmically

spaced difficulty levels (±Δ = [0°, 0.31°, 0.56°, 1°, 1.77°, 3.16°]) with 20 trials for each of the

orientations, respectively. The stimuli were presented in a pseudo-random order at 8.65° eccentricity

against a dark grey background (6.4 cd/m^2).

Each trial started with a finger and eye fixation phase of 1 to 2 s duration. Then, black placeholders

appeared for 1 or 2 s, followed by the presentation of the Gabor grating for 350 ms, a delay of 2 to 3 s

during which the placeholders' color switched from to blue and yellow, respectively. Subjects had 1 s

to respond. The start of the response period was indicated by change of fixation dot. The next trial

started 0.5 to 2 s later. If subjects did not respond within 1 s, the trial was repeated later during the

block. Similarly, if subjects broke fixation throughout the trial, the trial was aborted and repeated at

the end of the block. Feedback about correct and incorrect responses was provided with a blue or

white dot, respectively, at fixation.

Each training session included 220 trials evenly distributed among four blocks. Subjects were free to

take short breaks between blocks. The reference was shown for 2 s in the center of the screen before

and in the middle of each block (in total 8 times per session).

Subjects were instructed to respond as accurately as possible. To respond, subjects had to move their

finger from the fixation spot to the blue target if the stimulus was tilted clockwise or to the yellow

target if the stimulus was tilted counterclockwise (while maintaining fixation). Hand dominance was

assessed at the beginning of the experiment with the Edinburgh Handedness Inventory (Oldfield,

1971). In the transfer session, subjects were instructed to instead direct their gaze to the left or right

placeholder, respectively. Participant had to reach at least 75% correct responses before they

transitioned to the transfer session. All subjects reached this performance level within four days.

Analyses

The Signal Detection Theory parameter *d'* (D. M. Green & Swets, 1966) was computed per subject and

session. To quantify transfer performance, a Specificity Index (Petrov et al., 2011) was calculated, as

follows:

$$SI = \frac{d'_{last\ training} - d'_{transfer}}{d'_{last\ training} - d'_{first\ training}}$$

The index reflects the training session used to compute the average *d'*. *d'* and reaction times were

entered into repeated measures analyses of variance (rmANOVA) with factors session (4 levels for

training session effects, or 2 levels for last training session to transfer effects), difficulty (5 levels), and

their interaction. Two subjects required more than four days of training (5 and 6) in experiment 1;

therefore, we restrict our analyses to the last four training sessions. Due to technical issues, we lost

the data from the third training session for one subject in Experiment 1. Thus, to be able to compute

rmANOVA across training sessions (1-4), we excluded the subject (total *n*=18) from these analyses, but

we included the subject in the transfer test (last training versus transfer, total *n*=19). For *t*-tests and

rmANOVAs, we compute Hedges' *g* and partial η^2, respectively, as effect sizes, using the Measures of

Effect Size Toolbox (Hentschke & Stüttgen, 2011). All behavioral data are publicly available on Figshare

(private link to be released upon acceptance or reasonable request).

Drift Diffusion Model

Sequential sampling models like the DDM assume that decision making is a stochastic process of

evidence accumulation about a stimulus in time (Ratcliff, 1978). Decisions are made when the

accumulation process reaches an evidence threshold (decision criterion) for one of the alternatives.

The rate of accumulation and the decision criteria determine the reaction time (RT), and fluctuations

in the process are the reason for errors and the distribution of RTs. The DDM can account for effects

of different experimental conditions from accuracy, mean RT for correct and error responses, and the

shape of RT distributions for correct and error responses. The full DDM includes seven parameters:

drift rate (*v*) and its trial-by-trial variability (η), non-decision time (T_{er}) and its trial by-trial variability

(s_t), boundary separation (*a*), the starting point of the decision process (*z*) and its trial-by-trial variability

(s_z). *v* is the key component of the model, as it describes the rate of evidence accumulation and

positively correlates with the accuracy level of subjects. *a* is the distance between decision criteria for

both alternatives, and the bigger the distance is, the longer it takes to make a decision. The evidence

accumulation starts from *z*, which lies between 0 and *a*. *z* can be positioned closer to one of the

alternatives, which biases the decision process toward it. T_{er} accounts for the non-decision phases:

stimulus encoding and response output.

Here, we use the Hierarchical Drift Diffusion Model (HDDM 0.8.0) (Wiecki, Sofer, & Frank, 2013),

available as a toolbox in Python, to fit the model parameters to our data of Experiment 1. The HDDM

employs Bayesian inference to recover the model parameters. First, the HDDM estimates the

parameter values for a population (p_{group}), and then it uses the population fit to constrain the

parameter values for individual subjects (p_{subj}). The hierarchical Bayesian approach performs

significantly better than using χ^2 as the optimization criterion when data per subject is limited (Ratcliff

& Childers, 2015).

We estimate parameters for all sessions jointly while allowing v and a to vary across the sessions and

the difficulty levels (see code below). The stimuli are encoded by their orientation

(clockwise/counterclockwise), and v is estimated for both orientations together. We allow T_{er} and z to

vary only between sessions because a priori, encoding, muscle response time, and the orientation bias

should be the same across difficulty levels. We exclude the variability parameters from the analysis;

for proper estimation of these parameters, the HDDM requires noticeably more samples per condition

than we have available. We run the fitting algorithm four times with 30000 Markow-Chain-Monte-

Carlo samples, 10000 samples burn-in, and discarding every 5th sample.

```
260    model=hddm.HDDMStimCoding(data, split_param='v', include='z',
261    stim_col='orientation', p_outlier=0.05, informative=False, std_depends=False,
262    trace_subjs=True, is_group_model=True, depends_on={'v': ['difficulty', 'session'], 'z':
263    'session', 't': 'session', 'a': ['difficulty', 'session']})

264    model.find_starting_values()

265    model.sample(30000, burn=10000, thin=5, dbname=traces_name, db='pickle')
```

We utilize two formal diagnostics to assess the convergence of the model: the Gelman-Rubin and

Geweke method, respectively. The Gelman-Rubin method compares between-chain and within-chain

variances for each model parameter (Brooks & Gelman, 1998). Significant differences (diagnostics

value >1.2 or 1.1 under strict conditions) between these variances indicate non-convergence (Gelman

& Rubin, 1992). We compute the diagnostic with the help of the kabuki (0.6.3) Python library. The

Geweke method evaluates only within-chain fluctuation (Geweke, 1991). It calculates z-scores of the

difference between the initial and subsequent segments of the chain. If the chain has converged, most

points should be within two standard deviations of zero (z-score ∈ [-2, 2]). We chose the first 10% of

the chain as the initial segment, and the last 50% divided into 20 segments. To compute the Geweke

diagnostic, we use the PyMC (2.3.8) Python library.

To investigate the relationship between drift rate and boundary separation, we use a 2D minimum

energy test (Aslan & Zech, 2005; Székely & Rizzo, 2013), as implemented by B. Lau

(https://github.com/brian-lau/multdist). This test assesses via permutations whether two

multidimensional samples are drawn from the same distribution.

Results

Learning and transfer effects on accuracy and reaction times

We find that across all five difficulty levels, subjects significantly improve their performance (*d'*) as

function of training, with increases ranging between 0.60 and 1.93 (Fig. 2A, mean increase between

first and last training session 1.25, $t(18)$=-11.78, $p<0.001$, g=3.04; rmANOVA main effect of session,

$F(3, 51)$=138.94, $p<0.001$, η_p^2 =0.519). Subjects learn more on easy than on difficult orientation

differences (main effect of difficulty, $F(4, 68)$=96.18, $p<0.001$, η_p^2 =0.647, interaction session ×

difficulty $F(12, 204)$=13.18, $p<0.001$, η_p^2 =0.145). When the effector is changed from a manual joystick

response to a saccadic response in the transfer session, we find a significant decrease in performance

relative to the last session across all difficulty levels (mean decrease 0.32, $t(18)$=6.41, $p<0.001$, g=0.98;

rmANOVA main effect of session, $F(1,18)$= 21.48, $p<0.001$, η_p^2 =0.54 main effect of difficulty, $F(4,72)$=

290.68, $p<0.001$, η_p^2 =0.94, interaction session × difficulty $F(4,72)$=0.487, p=0.745, η_p^2 =0.02). This is

*Figure 2: Accuracy and reaction times for Experiment 1. A. Mean accuracy (in units of d') increases with learning from the first to the last training session (mean difference 1.25, t(18)=-11.78, p<0.001, g=3.04, two-sided) and drops from the level reached in the last training session when the effector is changed from a manual joystick response to a saccadic response (mean difference 0.32, t(18)=6.41, p<0.001, g=0.98, two-sided). B. Mean reaction times for correct and incorrect responses do not differ from each other throughout the sessions and do not change significantly as a function of learning (all p>0.29, all η_p^2 < 0.069). When the effector is changed, reaction times become uniformly shorter (main effect of session, F(1,18)=91.56, p<0.001, $\eta_p^2 = 0.83$, no main effect of accuracy or interaction, all p>0.09, all η_p^2 < 0.14). C. Learning depends on the effector, as evidenced by a mean Specificity Index of 0.25, which is significantly bigger than 0 (t(18)=6.65, p<0.001, g=2.11, one-sided). In all panels, error bars reflect ±2 SDs. *** stand for p < 0.001, ** for p < 0.01, and * for p<0.05.*

also evident when we calculate the SI, which has a mean of 0.25 (Fig. 2C, *t*(18)=6.65, *p*<0.001, *g*=2.11,

one-sided). Mean reaction times for correct and incorrect responses do not differ from each other

throughout the sessions and do not change significantly as a function of learning (all *p*>0.29, all η_p^2 <

0.069). When the effector is changed, reaction times become uniformly shorter (Fig. 2B, main effect

of session, *F*(1,18)=91.56, *p*<0.001, $\eta_p^2 = 0.83$, no main effect of accuracy or interaction, all *p*>0.09, all

η_p^2 < 0.14). This can be attributed to shorter latencies of saccades as compared to hand movements

(Jaskowski & Sobieralska, 2004).

We find very similar results in Experiment 2, where subjects perform the same task with a precise reach

on a touch screen. In this smaller sample (*n*=6), *d'* also improves with training over sessions (Fig. 3A,

mean increase between first and last training session 0.78, *t*(5)=-5.936, *p*=0.002, *g*=0.84; rmANOVA

main effect of session, *F*(3, 15)=35.23, *p*=0.002, $\eta_p^2 = 0.28$), with larger improvements for easier

orientation differences than for difficult orientation differences (main effect of difficulty,

F(4,20)=22.19, *p*<0.001, η_p^2 =0.60, interaction session × difficulty *F*(12,60)=4.81, *p*=0.007, η_p^2 =0.11).

When we change the motor component of the task from reaches to saccades in the transfer session,

we again find a drop in discrimination across difficulty levels, albeit not statistically significant (mean

decrease 0.21, *t*(5)=-1.23 *p*=0.273, *g*=-0.28; rmANOVA main effect of session, *F*(1,5)=1.51, *p*=0.273,

η_p^2 =0.03 main effect of difficulty, $F(4,20)$=41.74, p<0.001, η_p^2 =0.68, interaction session × difficulty

$F(4,20)$=1.362, p=0.282, η_p^2 =0.04). Expressed in mean SI, this is 0.17 ($t(5)$=0.85, p=0.217, g=0.45, one-

sided). However, four out of six subjects do show positive SIs, and after exclusion of a single outlier

with an extremely large negative SI, the mean SI is 0.34 and significant (Fig. 3C, $t(4)$=2.33, p=0.0398,

g=1.34, one-sided).

Average reaction times decrease over sessions (Fig. 3B, mean decrease 0.11 s, $t(5)$=2.82, p=0.037, g=-

0.99; rmANOVA main effect of session, $F(1,5)$=7.934, p=0.037, η_p^2 =0.254, main effect of accuracy,

$F(1,5)$=0.668, p=0.451, η_p^2 =0.001, interaction session × accuracy $F(1,5)$=0.07, p=0.8, η_p^2<0.001) and

further drop significantly when the effector is changed (mean decrease 0.19 s, $t(5)$=-4.97, p=0.004;

g=2.25; rmANOVA main effect of session, $F(1,5)$=0.668, p=0.037, $\eta_p^2 = 0.6356$, main effect of accuracy,

$F(1,5)$=0.012, p=0.909, η_p^2<0.001, interaction session × accuracy $F(1,5)$=1.067, p=0.349, η_p^2 =0.001).

Together, the results from Experiments 1 and 2 show that learning effects in orientation discrimination

depend on the effector without any change in the stimulus material or the task.

*Figure 3: Accuracy and reaction times for Experiment 2. **A.** Mean accuracy (in units of d') increases with learning from the first to the last training session (mean difference 0.78, t(5)=-5.94, p<0.05, g=0.84, two-sided). There is a numerical, but no statistically significant drop in accuracy when the effector is changed from a manual joystick response to a saccadic response (mean difference 0.21, t(5)=-1.23, p=0.273, g=-0.28, two-sided). **B.** Mean reaction times decrease over sessions (mean decrease 0.11 s, t(5)=2.82, p=0.037, g=-0.99) and further drop significantly when the effector is changed (mean decrease 0.19 s, t(5)=-4.97, p=0.004, g=2.25). **C.** Training is specific to the effector in 4 out of 6 subjects, as evidenced by a mean Specificity Index of 0.34, and statistically significantly bigger than 0 after exclusion of one outlier (t(4)=2.33, p=0.0398, g=1.34, one-sided). The dot reflects the outlier with a highly negative Specificity Index. In all panels, Error bars reflect ±2 SDs. *** stand for p < 0.001, ** for p < 0.01, and * for p<0.05.*

Learning and transfer effects on components of the decision-making process

To dissociate different aspects of the decision-making process and learning effects upon them, we use

Hierarchical Drift Diffusion Models (HDDM) to relate accuracy to reaction times. To this end, we

consider the data from Experiment 1 (joystick – saccade) because the task was performed under time

pressure (by means of a fixed response time window of 1 s), which is a requirement for these kinds of

analyses. We consider the parameters drift rate, decision boundary, bias, and non-decision time.

According to Gelman-Rubin method, all parameters converged with a value < 1.01 (Fig. S1). The

Geweke method indicates that all group parameters (n=64) have an average z-score $\in$ [-1.5, 1], and at

worst, 70% of the values are within two standard deviations of zero (Fig. S2). Also, all subject

parameters (n=1138) have an average z-score = ±2, which is within acceptable bounds. Only few

parameters have 50-70% of the values within two standard deviations of zero. Together, this shows

that our HDDMs converged well. Additionally, we verify the model through a posterior predictive

comparison (Table S1), which demonstrates that the predicted accuracy and reaction time

distributions match the empirical data.

*Figure 4: Hierarchical Drift Diffusion Model fit of drift rates and decision boundaries in Experiment 1. **A.** Drift rate (v) increases between the first and the last training session across orientations. When the effector changes in the transfer session, the learning-induced gains in drift rate drop. **B.** The separation between decision boundaries (a) increases between the first and last training session for the easier orientations and decreases or stays the same for the most challenging conditions. Learning effects drop when the effector changes in the transfer session. Squares signify the mean individual fits, errors bars ±2 standard deviations; crosses indicate the group fits. The discrepancy along the abscissa between sessions is intentional to increase visibility of data points.*

<u>Drift rate</u>

Both when assessed on the group and on the subject level, we find that drift rate increases between

the first and the last training session across difficulty levels (Fig. 4A and Table S2, $p_{subj}(v_{last\_training} >$

$v_{first\_training}$) = [0.91, 0.97, 0.99, 0.99, 1.0], $p_{group}(v_{last\_training} > v_{first\_training})$ = [0.99, 1.0, 1.0, 1.0, 1.0] for

±$\Delta_{orientation}$ = [0.31°, 0.56°, 1°, 1.77°, 3.16°], respectively). This is in keeping with previous reports that

show that VPL increases the drift rate in various tasks (e.g., C. C. Liu & Watanabe, 2012; Petrov et al.,

2011). When we change the effector in the transfer session, drift rates drop, which is evident in the

single subject and, especially, group fit of the HDDM ($p_{subj}(v_{last\_training} > v_{transfer})$ = [0.73, 0.61, 0.84, 0.78,

0.91], $p_{group}(v_{last\_training} > v_{transfer})$ = [0.95, 0.78, 0.99, 0.97, 0.99]). This suggests that the rate of evidence

accumulation depends on the effector.

<u>Decision boundaries</u>

We find that the separation between decision boundaries corresponding to clockwise and

counterclockwise choices increases between the first and the last training session, most clearly and

consistently in the easier conditions, but decreases or stays the same for the two most challenging

conditions (Fig. 4B and Table S2, $p_{subj}(a_{last\_training} > a_{first\_training})$ = [0.25, 0.46, 0.72, 0.89, 0.99],

$p_{group}(a_{last\_training} > a_{first\_training})$ = [0.05, 0.44, 0.93, 0.99, 1.0]] for ±$\Delta_{orientation}$ = [0.31°, 0.56°, 1°, 1.77°, 3.16°],

respectively). The separation is reduced when we change the effector in the transfer session

Figure 5: Hierarchical Drift Diffusion Model fit of non-decision times and biases in Experiment 1. A. Non-decision time (t) increases between the first and the last training session, and then decreases in the transfer session B. Subjects had a slight bias (z) in the clockwise direction during the first training session, which disappeared with the last training and remained low for the transfer session. Squares signify the mean individual fits, errors bars ±2 standard deviations; crosses indicate the group fits.

358 ($p_\text{subj}(a_\text{last\_training} > a_\text{transfer})$ = [0.25, 0.46, 0.72, 0.89, 0.99], $p_\text{group}(a_\text{last\_training} > a_\text{transfer})$ = [0.8, 0.87, 0.96, 0.97,

359 0.99]). Hence, like the drift rate, the decision boundary depends on the effector.

Non-decision time

Non-decision time increases between the first training session and the last training session, as is clearly

evident on the group level (Fig. 5A, $p_\text{group}(t_\text{last\_training} > t_\text{first\_training})$ = 0.995), but less consistently on the

subject level ($p_\text{subj}(t_\text{last\_training} > t_\text{first\_training})$ = 0.769). It then decreases massively in the transfer session

($p_\text{subj}(t_\text{last\_training} > t_\text{first\_training})$ = 1.0, $p_\text{group}(t_\text{last\_training} > t_\text{first\_training})$ = 1.0). The latter effect can likely be

attributed to shorter latencies of saccades as compared to hand movements (Jaskowski & Sobieralska,

2004).

Bias

368 We find a slight bias in clockwise orientation (Fig. 5B, z_subj = 0.471, z_group = 0.471) during the first training

369 session, which disappears with the last training (z_subj = 0.505, z_group = 0.505; $p_\text{subj}(z_\text{last\_training} > z_\text{first\_training})$

370 = 0.796, $p_\text{group}(z_\text{last\_training} > z_\text{first\_training})$ = 0.998). Hence, subjects seem to optimally adjust their response

371 bias in accordance with the task demands. During the transfer session, the bias is almost eliminated

372 (z_subj = 0.493, z_group = 0.493; $p_\text{subj}(z_\text{last\_training} > z_\text{first\_training})$ = 0.633, $p_\text{group}(z_\text{last\_training} > z_\text{first\_training})$ = 0.851).

Correlation between drift rate and decision boundary

We find that in our task, higher drift rates (v) are systematically accompanied by larger decision boundaries (a) within a session. The relationship between v and a is firmly linear for all sessions but the first one (Pearson correlation $r_{first\_training}$ = 0.622±0.338SD (p = 0.25±0.22SD), $r_{last\_training}$ = 0.967±0.028SD (p = 0.008±0.012SD), $r_{transfer}$ = 0.929±0.054SD (p = 0.026±0.028SD), see Table S3). This suggests that subjects strategically use the additional time afforded by an increase in drift rate to sample additional information towards higher accuracy instead of saving time on task by decreasing their reaction times through tighter decision bounds. We fit a first-degree polynomial to the data per subject and session, where a depends on v over orientations (difficulty) levels. Thus, the slope of the linear fit reflects how a changes with growing v, and the offset is a when v equals zero. Throughout training, the slope and

Figure 6: Relationship between drift rate and decision boundary. Slope and offset of a linear fit of drift rates against decision boundaries across orientations. Throughout the training sessions, the slope and the offset stabilize at higher and lower values, respectively. This is evidenced by a significant difference in the a-v relationship between the first and last training session (2D minimum energy test, e=2.391, p=0.001). During the transfer session, only the slope changes (t(18)=4.28, p=0.0004), while the offset remains the same as in the last training session (t(18)=-0.77, p=0.44). Colored error bars display the estimates based on the per-subject parameters (mean ±2 SDs), the black crosses show the estimates based on group parameters. Numbers indicate training sessions (1-4), transfer session is red and marked with T.

offset converge to higher and lower values, respectively (Fig. 6). This is evidenced by a significant difference in the *v-a* relationship between the first and last training session (2D minimum energy test, *e*=2.391, *p*=0.001). During the transfer session, only the slope changes (*t*(19)=4.28, *p*=0.0004), while the offset is not different from the last training session (*t*(18)=-0.77, *p*=0.44). Accordingly, the difference between the final training session and the transfer session is statistically significant in 2D (*e*=0.165, *p*=0.04); however, not in comparison with the penultimate training session (*e*=0.040, *p*=0.65). Considering the faster reaction times (by ~150ms) during the transfer session, subjects could compensate to a certain degree for the reduced *v* by raising *a*, which would be reflected in a higher offset value. Overall, this indicates that first, subjects acquire a task strategy that prioritizes accuracy over time; and second, that while task learning is specific to the trained effector, the learned strategy is transferred to another effector.

Discussion

We find that VPL of orientation discrimination depends on the effector with which the task is trained. Analyses with the HDDM reveal that this dependency arises from effector specificity of two subcomponents of the decision-making process, namely the rate of evidence accumulation and the decision bound. Our results thus confirm that sensorimotor mapping is affected by VPL but suggests a new model on how and where this mapping may take place.

Previous models of VPL suggest either full or no transfer between effectors. Full transfer would take place if learning took place entirely in visual cortex (Fahle, 2004; Sagi, 2011), as originally conceived, or if sensorimotor mapping occurs between visual and decision making neurons, e.g., in parietal cortex, in the form of learning an abstract "rule" (Awada et al., 2021). No transfer between effectors would be predicted if sensorimotor mapping would take place between visual and effector-specific neurons in motor or parietal cortex (Grzeczkowski et al., 2017). In theory, learning abstract rules leads to excellent generalization capacity, which has for example been suggested to be the basis of visual training from action video games (C. S. Green, Pouget, & Bavelier, 2010). On the other hand, decomposing tasks into effector-specific components greatly simplifies reinforcement learning and effector-specific reward prediction errors supporting such learning have been found in humans using fMRI (Gershman, Pesaran, & Daw, 2009). Effector specificity has been found in several other forms of learning involving sensorimotor mapping (e.g., Feulner, Postin, Schwiedrzik, & Pooresmaeili, 2021), but generalization across effectors is also not uncommon (reviewed in Halsband & Lange, 2006). Our results can be interpreted as an *effector-specific decision rule*. Here, sensorimotor mapping would occur between visual neurons and effector-specific integrators, e.g., in parietal cortex, similar to the *specific stimulus-response mapping model*.

However, the degree of specificity we find falls between the two previously reported extremes, as transfer is neither full nor none. Given the co-localization of saccade-specific, reach-specific, and effector-invariant motor neurons in parietal cortex (Calton, Dickinson, & Snyder, 2002; Diomedi, Vaccari, Filippini, Fattori, & Galletti, 2020), it is possible that there is overlap of the populations of neurons encoding motor plans for these two effectors. The degree of overlap would then determine the degree of transfer between the trained and untrained effectors in the transfer session. A possible reason for the colocalization of saccade- and reach-selective neurons in parietal cortex is the need for eye-hand coordination in many real-world tasks. Saccades may always be planned along limb movements for visual guidance, but not executed if fixation is required by the task (Levy, Schluppeck, Heeger, & Glimcher, 2007). Such consistent coactivation could also account for partial transfer results, as saccades would be trained along hand movements in an implicit fashion, perhaps through Hebbian plasticity.

Classical VPL models in which learning takes place only in visual cortex predict full transfer and cannot easily explain partial specificity of learning effects to effectors. However, visual cortex activity has been shown to be differentially modulated by saccades versus reaches in humans using fMRI (Gallivan, Chapman, Gale, Flanagan, & Culham, 2019; Levy et al., 2007). In rodents, there is evidence that motor effects in visual cortex increase information about the orientation of visual stimuli (reviewed in Pakan, Francioni, & Rochefort, 2018). This suggests that interactions between visual and motor components of VPL could hypothetically already arise in early visual areas but still points to sensorimotor interactions as a relevant contributor to VPL.

Because neurons in parietal cortex have been shown to integrate visual information towards a decision bound, the parameters of the DDM are often interpreted to arise in parietal cortex as well (Smith & Ratcliff, 2004). This would fit with the known existence of effector-specific integrators and their colocalization in areas like LIP that from the basis of the newly suggested effector-specific but overlapping integrators model. An increased drift rate is usually interpreted as better sensory input to decision-making process (Petrov et al., 2011; Ratcliff, 2014). However, this interpretation cannot fully account for the effector specificity of the drift rate parameter that we found, because the sensory input does not change when the effector changes. Rather, our results suggest that the drift rate (also) reflects the integration process itself, i.e., the action of the integrator. For example, it is possible that integrators in parietal cortex increase their sensitivity to sensory information, which could further contribute to a fast integration of information towards the decision bound.

We find that VPL also has an effect on the decision bound, and that this effect is also (partially) effector-specific. Trainees seem to adjust their decision bound such that faster integration is in a way counteracted by a more conservative decision bound. This policy results in higher accuracy because more information can be sampled but forgoes the opportunity to save time on task by reaching a decision earlier on the basis of a higher drift rate alone. Previous studies have reported a decreased decision bound (C. C. Liu & Watanabe, 2012) and less variability in the decision bound parameter (Petrov et al., 2011) with VPL. However, in both these studies speed was emphasized over accuracy, whereas our task design favored accuracy, which may explain this discrepancy.

Neurally, narrower decision bounds are correlated with an increase in activity at the outset of the integration process in area LIP (Hanks, Kiani, & Shadlen, 2014). Since many neurons in LIP are effector specific, it is possible that the same neurons that may account for the learning effects and the specificity of the drift rate also account for the effects on the decision bound. Our model would thus predict a reduction in spontaneous choice-predictive activity in areas like LIP (Shadlen & Newsome, 2001) with VPL. Alternatively, a central decision bound circuit, e.g., in the basal ganglia (Zhang, 2012),

where correlates of VPL have also been found (Guggenmos, Wilbertz, Hebart, & Sterzer, 2016), could

account for our results.

Our results are in partial conflict with previous reports on full transfer (Awada et al., 2021) and no

transfer between effectors (C. S. Green et al., 2015; Grzeczkowski et al., 2017). As mentioned in the

introduction, the latter results can conceivably be explained by task-specificity of VPL. Several

possibilities could account for the discrepancy with the study by Awada and colleagues: the study by

Awada and colleagues tested a much smaller sample of subjects (6 and 4 subjects in their Experiment

1 and 2, respectively), which may have been insufficient to reveal partial transfer effects. In fact, the

results from Awada and colleagues' Experiment 1 show a small trend towards partial transfer that did

not reach statistical significance. However, it is also possible that differences in the stimuli and task

account for the divergent results. In contrast to our study on orientation discrimination, Awada and

colleagues investigated motion direction discrimination VPL. Motion direction is processed in dorsal

stream areas, which are anatomically closer to parietal decision-making areas than neurons encoding

orientation, which primarily reside in the ventral stream. Furthermore, parietal areas such as LIP

contain intrinsically motion direction sensitive (i.e., sensory) neurons (Fanini & Assad, 2009). Both

these factors could ease sensorimotor mapping to multiple colocalized effectors. Further studies with

high statistical power explicitly comparing motion direction and orientation discrimination tasks will

be able to resolve these alternative interpretations.

Taken together, we find that VPL is not as specific, but also not as generalizable as previously thought.

Taking into account different components of the sensorimotor arc revealed additional aspects of VPL

beyond the traditional focus on early visual areas. Our results also suggest an expanded interpretation

of the drift rate as reflecting not only the amount of accumulable information, but also the sensitivity

of the integrator itself. Finally, in terms of application of VPL (Deveau & Seitz, 2014), where

generalization is key, it may be beneficial to counteract overtraining by varying effectors. Given the

new effector-specific learning rule model we propose, more effector variability during training may

enable more generalization of training outcomes (also see Grzeczkowski, Cretenoud, Mast, & Herzog,

2019).

Acknowledgements

This project was supported by a seed fund from the Leibniz ScienceCampus 'Primate Cognition',

Göttingen, Germany (to CMS and IK), and has received funding from the European Research Council

(ERC) under the European Union's Horizon 2020 research and innovation programme (Grant

agreement No. 802482, to CMS). CMS is supported by the German Research Foundation Emmy

References

Aslan, B., & Zech, G. (2005). Statistical energy as a tool for binning-free, multivariate goodness-of-fit tests, two-sample comparison and unfolding. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, 537*(3), 626-636. doi:10.1016/j.nima.2004.08.071

Awada, A., Bakhtiari, S., & Pack, C. C. (2021). Visual perceptual learning generalizes to untrained effectors. *J Vis, 21*(3), 10. doi:10.1167/jov.21.3.10

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis, 10*(4), 433-436.

Brooks, S. P., & Gelman, A. (1998). General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics, 7*(4), 434-455. doi:10.1080/10618600.1998.10474787

Calton, J. L., Dickinson, A. R., & Snyder, L. H. (2002). Non-spatial, motor-specific activation in posterior parietal cortex. *Nat Neurosci, 5*(6), 580-588. doi:10.1038/nn0602-862

Chang, S. W., Calton, J. L., Lawrence, B. M., Dickinson, A. R., & Snyder, L. H. (2016). Region-Specific Summation Patterns Inform the Role of Cortical Areas in Selecting Motor Plans. *Cereb Cortex, 26*(5), 2154-2166. doi:10.1093/cercor/bhv047

Christopoulos, V. N., Bonaiuto, J., Kagan, I., & Andersen, R. A. (2015). Inactivation of Parietal Reach Region Affects Reaching But Not Saccade Choices in Internally Guided Decisions. *J Neurosci, 35*(33), 11719-11728. doi:10.1523/JNEUROSCI.1068-15.2015

Christopoulos, V. N., Kagan, I., & Andersen, R. A. (2018). Lateral intraparietal area (LIP) is largely effector-specific in free-choice decisions. *Sci Rep, 8*(1), 8611. doi:10.1038/s41598-018-26366-9

Colby, C. L., & Goldberg, M. E. (1999). Space and attention in parietal cortex. *Annu Rev Neurosci, 22*, 319-349. doi:10.1146/annurev.neuro.22.1.319

de Lafuente, V., Jazayeri, M., & Shadlen, M. N. (2015). Representation of accumulating evidence for a decision in two parietal areas. *J Neurosci, 35*(10), 4306-4318. doi:10.1523/JNEUROSCI.2451-14.2015

Deveau, J., & Seitz, A. R. (2014). Applying perceptual learning to achieve practical changes in vision. *Front Psychol, 5*, 1166. doi:10.3389/fpsyg.2014.01166

Dickinson, A. R., Calton, J. L., & Snyder, L. H. (2003). Nonspatial saccade-specific activation in area LIP of monkey parietal cortex. *J Neurophysiol, 90*(4), 2460-2464. doi:10.1152/jn.00788.2002

Diomedi, S., Vaccari, F. E., Filippini, M., Fattori, P., & Galletti, C. (2020). Mixed Selectivity in Macaque Medial Parietal Cortex during Eye-Hand Reaching. *iScience, 23*(10), 101616. doi:10.1016/j.isci.2020.101616

Dosher, B., & Lu, Z. L. (2017). Visual Perceptual Learning and Models. *Annu Rev Vis Sci, 3*, 343-363. doi:10.1146/annurev-vision-102016-061249

Dutilh, G., Krypotos, A. M., & Wagenmakers, E. J. (2011). Task-related versus stimulus-specific practice. *Exp Psychol, 58*(6), 434-442. doi:10.1027/1618-3169/a000111

Fahle, M. (2004). Perceptual learning: a case for early selection. *J Vis, 4*(10), 879-890. doi:10.1167/4.10.4

Fahle, M., & Morgan, M. (1996). No transfer of perceptual learning between similar stimuli in the same retinal position. *Curr Biol, 6*(3), 292-297. doi:10.1016/s0960-9822(02)00479-7

Fanini, A., & Assad, J. A. (2009). Direction selectivity of neurons in the macaque lateral intraparietal area. *J Neurophysiol, 101*(1), 289-305. doi:10.1152/jn.00400.2007

Feulner, B., Postin, D., Schwiedrzik, C. M., & Pooresmaeili, A. (2021). Previous Motor Actions Outweigh Sensory Information in Sensorimotor Statistical Learning. *eNeuro, 8*(5). doi:10.1523/ENEURO.0032-21.2021

Gallivan, J. P., Chapman, C. S., Gale, D. J., Flanagan, J. R., & Culham, J. C. (2019). Selective Modulation of Early Visual Cortical Activity by Movement Intention. *Cereb Cortex, 29*(11), 4662-4678. doi:10.1093/cercor/bhy345

Gelman, A., & Rubin, D. B. (1992). Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science, 7*(4), 457-472.

Gershman, S. J., Pesaran, B., & Daw, N. D. (2009). Human reinforcement learning subdivides structured action spaces by learning effector-specific values. *J Neurosci, 29*(43), 13524-13531. doi:10.1523/JNEUROSCI.2469-09.2009

Geweke, J. F. (1991). *Evaluating the accuracy of sampling-based approaches to the calculation of posterior moments*. Retrieved from https://ideas.repec.org/p/fip/fedmsr/148.html

Ghose, G. M., Yang, T., & Maunsell, J. H. (2002). Physiological correlates of perceptual learning in monkey V1 and V2. *J Neurophysiol, 87*(4), 1867-1888. doi:10.1152/jn.00690.2001

Gottlieb, J., & Goldberg, M. E. (1999). Activity of neurons in the lateral intraparietal area of the monkey during an antisaccade task. *Nat Neurosci, 2*(10), 906-912. doi:10.1038/13209

Green, C. S., Kattner, F., Siegel, M. H., Kersten, D., & Schrater, P. R. (2015). Differences in perceptual learning transfer as a function of training task. *J Vis, 15*(10), 5. doi:10.1167/15.10.5

Green, C. S., Pouget, A., & Bavelier, D. (2010). Improved probabilistic inference as a general learning
mechanism with action video games. *Curr Biol, 20*(17), 1573-1579.
doi:10.1016/j.cub.2010.07.040

Green, D. M., & Swets, J. A. (1966). *Signal detection theory and psychophysics*. New York: Wiley.

Grieco, A., Casco, C., & Roncato, S. (2006). Texture segregation on the basis of contrast polarity of odd-
symmetric filters. *Vision Res, 46*(20), 3526-3536. doi:10.1016/j.visres.2006.05.002

Grzeczkowski, L., Cretenoud, A., Herzog, M. H., & Mast, F. W. (2017). Perceptual learning is specific
beyond vision and decision making. *J Vis, 17*(6), 6. doi:10.1167/17.6.6

Grzeczkowski, L., Cretenoud, A. F., Mast, F. W., & Herzog, M. H. (2019). Motor response specificity in
perceptual learning and its release by double training. *J Vis, 19*(6), 4. doi:10.1167/19.6.4

Guggenmos, M., Wilbertz, G., Hebart, M. N., & Sterzer, P. (2016). Mesolimbic confidence signals guide
perceptual learning in the absence of external feedback. *Elife, 5*. doi:10.7554/eLife.13388

Halsband, U., & Lange, R. K. (2006). Motor learning in man: a review of functional and clinical studies.
J Physiol Paris, 99(4-6), 414-424. doi:10.1016/j.jphysparis.2006.03.007

Hanks, T., Kiani, R., & Shadlen, M. N. (2014). A neural mechanism of speed-accuracy tradeoff in
macaque area LIP. *Elife, 3*. doi:10.7554/eLife.02260

Hentschke, H., & Stüttgen, M. C. (2011). Computation of measures of effect size for neuroscience data
sets. *Eur J Neurosci, 34*(12), 1887-1894. doi:10.1111/j.1460-9568.2011.07902.x

Jaskowski, P., & Sobieralska, K. (2004). Effect of stimulus intensity on manual and saccadic reaction
time. *Percept Psychophys, 66*(4), 535-544. doi:10.3758/bf03194899

Karni, A., & Sagi, D. (1991). Where practice makes perfect in texture discrimination: evidence for
primary visual cortex plasticity. *Proc Natl Acad Sci U S A, 88*(11), 4966-4970.
doi:10.1073/pnas.88.11.4966

Kazennikov, O., Hyland, B., Corboz, M., Babalian, A., Rouiller, E. M., & Wiesendanger, M. (1999). Neural
activity of supplementary and primary motor areas in monkeys and its relation to bimanual
and unimanual movement sequences. *Neuroscience, 89*(3), 661-674. doi:10.1016/s0306-
4522(98)00348-0

Klorfeld-Auslender, S., & Censor, N. (2019). Visual-oculomotor interactions facilitate consolidation of
perceptual learning. *J Vis, 19*(6), 11. doi:10.1167/19.6.11

Law, C. T., & Gold, J. I. (2008). Neural correlates of perceptual learning in a sensory-motor, but not a
sensory, cortical area. *Nat Neurosci, 11*(4), 505-513. doi:10.1038/nn2070

Levy, I., Schluppeck, D., Heeger, D. J., & Glimcher, P. W. (2007). Specificity of human cortical areas for
reaches and saccades. *J Neurosci, 27*(17), 4687-4696. doi:10.1523/JNEUROSCI.0459-07.2007

Liu, B. H., Huberman, A. D., & Scanziani, M. (2016). Cortico-fugal output from visual cortex promotes
plasticity of innate motor behaviour. *Nature, 538*(7625), 383-387. doi:10.1038/nature19818

Liu, C. C., & Watanabe, T. (2012). Accounting for speed-accuracy tradeoff in perceptual learning. *Vision*
Res, 61, 107-114. doi:10.1016/j.visres.2011.09.007

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory.
Neuropsychologia, 9(1), 97-113. doi:10.1016/0028-3932(71)90067-4

Pakan, J. M., Francioni, V., & Rochefort, N. L. (2018). Action and learning shape the activity of neuronal
circuits in the visual cortex. *Curr Opin Neurobiol, 52*, 88-97. doi:10.1016/j.conb.2018.04.020

Petrov, A. A., Dosher, B. A., & Lu, Z. L. (2005). The dynamics of perceptual learning: an incremental
reweighting model. *Psychol Rev, 112*(4), 715-743. doi:10.1037/0033-295X.112.4.715

Petrov, A. A., Van Horn, N. M., & Ratcliff, R. (2011). Dissociable perceptual-learning mechanisms
revealed by diffusion-model analysis. *Psychon Bull Rev, 18*(3), 490-497. doi:10.3758/s13423-
011-0079-8

Pilacinski, A., & Lindner, A. (2019). Distinct contributions of human posterior parietal and dorsal
premotor cortex to reach trajectory planning. *Sci Rep, 9*(1), 1962. doi:10.1038/s41598-019-
39188-0

Poggio, T., Fahle, M., & Edelman, S. (1992). Fast perceptual learning in visual hyperacuity. *Science,*
256(5059), 1018-1021. doi:10.1126/science.1589770

Ratcliff, R. (1978). A theory of memory retrieval. *Psychol Rev, 85*(2), 59-108. doi:10.1037/0033-
295X.85.2.59

Ratcliff, R. (2014). Measuring psychometric functions with the diffusion model. *J Exp Psychol Hum*
Percept Perform, 40(2), 870-888. doi:10.1037/a0034954

Ratcliff, R., & Childers, R. (2015). Individual Differences and Fitting Methods for the Two-Choice
Diffusion Model of Decision Making. *Decision (Wash DC), 2*(4), 237-279.
doi:10.1037/dec0000030

Sagi, D. (2011). Perceptual learning in vision research. *Vision Res, 51*(13), 1552-1566.
doi:10.1016/j.visres.2010.10.019

Schoups, A. A., Vogels, R., & Orban, G. A. (1995). Human perceptual learning in identifying the oblique
orientation: retinotopy, orientation specificity and monocularity. *J Physiol, 483 (Pt 3)*, 797-
810. doi:10.1113/jphysiol.1995.sp020623

Schwiedrzik, C. M., Singer, W., & Melloni, L. (2011). Subjective and objective learning effects dissociate
in space and in time. *Proc Natl Acad Sci U S A, 108*(11), 4506-4511.
doi:10.1073/pnas.1009147108

Series, P., Latham, P. E., & Pouget, A. (2004). Tuning curve sharpening for orientation selectivity: coding
efficiency and the impact of correlations. *Nat Neurosci, 7*(10), 1129-1135. doi:10.1038/nn1321

Shadlen, M. N., & Newsome, W. T. (2001). Neural basis of a perceptual decision in the parietal cortex

(area LIP) of the rhesus monkey. *J Neurophysiol, 86*(4), 1916-1936.

doi:10.1152/jn.2001.86.4.1916

Smith, P. L., & Ratcliff, R. (2004). Psychology and neurobiology of simple decisions. *Trends Neurosci, 27*(3), 161-168. doi:10.1016/j.tins.2004.01.006

Snyder, L. H., Batista, A. P., & Andersen, R. A. (1997). Coding of intention in the posterior parietal

cortex. *Nature, 386*(6621), 167-170. doi:10.1038/386167a0

Székely, G. J., & Rizzo, M. L. (2013). Energy statistics: A class of statistics based on distances. *Journal of*

Statistical Planning and Inference, 143(8), 1249-1272. doi:10.1016/j.jspi.2013.03.018

Szpiro, S. F., Spering, M., & Carrasco, M. (2014). Perceptual learning modifies untrained pursuit eye

movements. *J Vis, 14*(8), 8. doi:10.1167/14.8.8

Szul, M. J., Bompas, A., Sumner, P., & Zhang, J. (2020). The validity and consistency of continuous

joystick response in perceptual decision-making. *Behav Res Methods, 52*(2), 681-693.

doi:10.3758/s13428-019-01269-3

Szumska, I., van der Lubbe, R. H., Grzeczkowski, L., & Herzog, M. H. (2016). Does sensitivity in binary

choice tasks depend on response modality? *Conscious Cogn, 43*, 57-65.

doi:10.1016/j.concog.2016.05.005

Vogels, R. (2010). Mechanisms of visual perceptual learning in macaque visual cortex. *Top Cogn Sci,*

2(2), 239-250. doi:10.1111/j.1756-8765.2009.01051.x

Wiecki, T. V., Sofer, I., & Frank, M. J. (2013). HDDM: Hierarchical Bayesian estimation of the Drift-

Diffusion Model in Python. *Front Neuroinform, 7*, 14. doi:10.3389/fninf.2013.00014

Wilke, M., Turchi, J., Smith, K., Mishkin, M., & Leopold, D. A. (2010). Pulvinar inactivation disrupts

selection of movement plans. *J Neurosci, 30*(25), 8650-8659. doi:10.1523/JNEUROSCI.0953-

10.2010

Zhang, J. (2012). The effects of evidence bounds on decision-making: theoretical and empirical

developments. *Front Psychol, 3*, 263. doi:10.3389/fpsyg.2012.00263

Zhang, J., & Rowe, J. B. (2014). Dissociable mechanisms of speed-accuracy tradeoff during visual

perceptual learning are revealed by a hierarchical drift-diffusion model. *Front Neurosci, 8*, 69.

doi:10.3389/fnins.2014.00069