Cell type- and state- resolved immune transcriptomic profiling identifies glucocorticoid-responsive molecular defects in multiple sclerosis T cells

Tina Roostaei[1], Afsana Sabrin[1], Pia Kivisakk[2], Cristin McCabe[3], Parham Nejad[3], Daniel Felsky[4], Hanane Touil[1], Ioannis Vlachos[3], Daniel Hui[3], Jennifer Fransson[5], Nikolaos A. Patsopoulos[3], Vijay K. Kuchroo[6], Violetta Zujovic[5], Howard L. Weiner[3], Hans-Ulrich Klein[1], Philip L. De Jager[1,7]*

[1] Center for Translational & Computational Neuroimmunology, Department of Neurology, Columbia University Irving Medical Center, New York, USA
[2] Alzheimer's Clinical and Translational Research Unit, Department of Neurology, Massachusetts General Hospital, Boston, USA
[3] Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, USA
[4] Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, University of Toronto, Toronto, Canada
[5] Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS, APHP, Hôpital Pitié Salpétrière Univ. Hosp., DMU Neuroscience 6, Paris, France
[6] Evergrande Center for Immunologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, USA
[7] Columbia Multiple Sclerosis Center, Department of Neurology, Columbia University Irving Medical Center, New York, USA

* Correspondence:
Philip L. De Jager, MD PhD
pld2115@cumc.columbia.edu

Abstract

The polygenic and multi-cellular nature of multiple sclerosis (MS) pathology necessitates cell-type-specific molecular studies in order to improve our understanding of the mechanisms underlying immune cell dysfunction in MS. Here, using a large dataset of 1075 transcriptomes from 209 MS and healthy individuals, we assessed MS-associated transcriptional changes in 6 cell-type-states implicated in MS: naïve and memory CD4$^+$ T cells and CD14$^+$ monocytes purified from peripheral blood, each in primary (unstimulated) and *in vitro* stimulated states. Our analyses identified shared and distinct changes in individual genes, biological pathways, and co-expressed gene modules in MS T cells and monocytes. We found evidence of replication for the association between MS and three (two T cell and one monocyte) co-expressed gene modules in independent data from peripheral blood mononuclear cells (PBMC) and monocyte-derived macrophages. Our computational drug screen identified glucocorticoid receptor agonists as the top therapeutic compounds for reversing the replicated MS-associated T cell modules. This finding was further supported by the significant enrichment of these modules for literature-driven glucocorticoid response gene sets, in addition to results from our *in vitro* experiment on gene expression effects of the glucocorticoid receptor agonist dexamethasone in MS PBMC. In summary, our study provides cell-type-specific information on genes and pathways perturbed in MS T cells and monocytes, and suggests a novel explanation for the therapeutic effects of glucocorticoids in relation to MS T cell pathology.

Introduction

Multiple sclerosis (MS) is a chronic inflammatory disease of the central nervous system (CNS). The earlier phase of the disease is characterized by episodes of focal blood-brain-barrier breakdown and recruitment of peripheral lymphoid and myeloid cells to active demyelinating areas within the white matter, in addition to slow accumulation of lymphocytes in the meninges leading to inflammation and demyelination in the underlying gray matter[1]. The enrichment of genes associated with MS genetic susceptibility loci for genes expressed in lymphoid and myeloid cells further strengthens the evidence on the role of several immune cell types in MS pathogenesis[2].

The main therapeutic mechanisms currently applied to MS treatment include modulating the function of immune cells, depleting subtypes of these cells, or retaining them in the periphery and preventing them from entering the CNS[3]. However, none of these interventions provide definitive treatment, highlighting our limited understanding of the molecular defects in immune cells involved in MS. Previous transcriptomic studies on whole blood or peripheral blood mononuclear cells (PBMC) have prioritized a number of genes and pathways in relation to immune cell dysfunction in MS[4–6]. However, the inference of cell-type-specific defects from these samples that contain variable proportions of different cell types remains a suboptimal approach.

More recently, data collection from purified peripheral immune cell types has started to address this problem and improve our understanding of the cell-type-specific immune dysfunction associated with MS[7–10]. Here, overcoming some of the limitations of the previous studies (including limited sample sizes, treatment-related confounding effects, and few number of simultaneously-studied cell types), we aimed to disentangle shared and distinct gene expression changes (both at the individual gene level and gene networks) in 6 immune cell types and states implicated in MS. Using a large dataset of 1176 transcriptomes from 227 individuals, we assessed MS-associated transcriptional changes in naïve and memory helper T cells and classical monocytes, each in primary and stimulated states. We then complemented our analyses with replication studies, computational drug screen, and *in vitro* validation experiments. Our results, (1) provide cell-type-specific information on genes and pathways perturbed in MS; (2) identify and replicate coordinated gene expression changes associated with MS; and (3) identify and validate a connection between coordinated transcriptional changes in MS T cells and therapeutic effect of glucocorticoids.

Results

Data generation
MS participants were selected from the Comprehensive Longitudinal Investigation of Multiple Sclerosis at the Brigham and Women's Hospital (CLIMB) study[11] using the following selection criteria: (1) age 18-60 years old, (2) a diagnosis of MS fulfilling 2010 McDonald criteria, (3) a relapsing-remitting disease course at the time of sampling, (4) being untreated for >6 months or on treatment with glatiramer acetate (GA) for >1 year at the time of sampling, (5) no evidence of disease activity in the prior 6 months, (6) no steroid use in the preceding 30 days, and (7) an Expanded Disability Status Scale (EDSS) score between 0 to 3. Of note, we restricted the inclusion of treated MS individuals to those receiving GA in order to minimize the confounding effects of treatment on immune cell gene expression[12]. Healthy participants were from the PhenoGenetic Project, part of the ImmVar study[13,14].

A prospectively collected, cryopreserved vial of PBMC was accessed for each participant. PBMC were thawed and sorted using flow cytometry, and three cell populations were retained: $CD14^+$ monocytes as well as $CD4^+$ naïve and memory T cells. Each of the three cellular samples were split into two aliquots, with one aliquot remaining unstimulated while the other aliquot was placed in culture and stimulated for 24 hours before RNA was extracted for sequencing. Anti-CD3/CD28 antibodies were used for T cell stimulation in order to mimic stimulation by antigen-presenting cells[15], and TNF-α was used for monocyte activation given its implication in MS pathology[16]. Thus, six RNA samples (3 cell types, unstimulated and stimulated) were collected from each cryopreserved PBMC sample (Figure 1). All RNA samples were processed using the Smartseq2 protocol to produce cDNA and a transcriptomic profile using sequencing. A rigorous pre-processing pipeline was instituted after the experimental pipeline was concluded (see Methods), and, ultimately, data were retained from 209 participants of European ancestry. These include 42 healthy participants as well as 167 individuals with relapsing-remitting MS with no evidence of disease activity at the time of sampling. The latter included 69 untreated and 98 GA-treated participants. The demographic characteristics of the participants are presented in Table 1.

Table 1. Demographic characteristics of the participants.

study	cell type	diagnosis (MS/HD)	age (mean ±SD)	sex (F/M)	MS medication
main study	six cell-type-states	167 MS, 42 HD	39.5 ±7.5	155/54	69 untreated, 98 glatiramer acetate
replication #1	PBMC	29 MS, 36 HD	37.4 ±9.7	43/22	29 glatiramer acetate
replication #2	monocyte-derived macrophages	28 MS, 11 HD	37.9 ±10.9	34/5	11 untreated, 17 treated*
in vitro experiment	PBMC	5 MS			

*treatments: n=4 dimethyl fumarate, n=3 natalizumab, n=3 fingolimod, n=2 rituximab, n=2 IFNβ, n=1 teriflunomide, n=1 methotrexate, n=1 glatiramer acetate

The effects of diagnosis and treatment on T cell and monocyte gene expression in MS
For each of the primary (unstimulated) and stimulated cell types, we first performed transcriptome-wide differential gene expression analysis between (1) untreated MS and healthy participants, (2) GA-treated MS and healthy participants, and (3) untreated and GA-treated MS participants. The number of significant differentially expressed genes (adjusted $p <0.05$) in each comparison is shown in Figure 2A. Two main conclusions can be drawn from these simple, descriptive analyses:

(1) There is little or no significant difference in gene expression (range: 0-7 significant genes) between MS (either untreated or GA-treated) and healthy participants in stimulated CD4$^+$ T cells and monocytes. On the other hand, unstimulated cell types showed higher number of differentially expressed genes. Interestingly, memory T (Tmem) cells, which are produced from prior *in vivo* activation of naïve T cells (Tn), had lower number of differentially expressed genes in comparison to their naïve counterparts.

(2) We did not observe notable difference in gene expression in any of the studied primary and stimulated cell types between untreated and GA-treated MS participants (range: 0-5 significant genes; Supplementary Table 1). This was further supported by the high correlations between *t*-statistics from the transcriptome-wide untreated MS vs. healthy and GA-treated MS vs. healthy comparisons (Figure 2B, Supplementary Figure 1). These results corroborate previous literature on the lack of transcriptional difference between untreated and GA-treated MS in PBMC[12]. As a result of these observations, we decided to combine data from untreated and GA-treated MS participants into one group (referred to as the MS group), and take advantage of the larger sample size for performing gene expression comparisons between MS and healthy participants.

T cell and monocyte genes differentially expressed in MS
Performing transcriptome-wide differential gene expression analysis between MS (combined group) and healthy participants, we identified 704 genes that were differentially expressed (adjusted $p < 0.05$) in at least one of the 6 cell-type-states (Figure 2C, Supplementary Table 2, Supplementary Data). The vast majority of the identified differentially expressed genes (>99%) had a nominal $p < 0.05$ with a consistent direction of effect in both untreated MS vs. healthy and GA-treated MS vs. healthy comparisons in their relevant cell-type-states. However, only 50% of the genes had an adjusted $p < 0.05$ in each of the two separate comparisons (Figure 2C), highlighting the effect of the larger sample size in identifying larger number of differentially expressed genes.

We then performed differential gene expression analysis on RNA-seq data available from a smaller number of PBMC samples (n=65; 29 GA-treated MS and 36 healthy participants, from which, 6 of the MS and 29 of the healthy individuals had paired data in the 6 cell-type-states dataset; details in the Methods, Table 1). Of the 704 identified genes, we replicated the association of 12 genes (adjusted $p < 0.05$), and 170 additional genes had a nominal $p < 0.05$ in the PBMC samples (Figure 2C, Supplementary Table 2), suggesting a broader replication of our results in this small replication data set. However, the lower correlations between *t*-statistics from the PBMC analysis and those from the purified cell types - which are part of the PBMC mixture - suggests that the differences observed in primary cell types in MS cannot be fully captured using PBMC data, and vice versa (Supplementary Figure 2).

Another observation from our data was that although a large percentage (72%) of the Tmem differentially expressed genes were also differentially expressed in Tn cells, but this was not the case between Tn cells and monocytes (Figure 2D). Of the >200 differentially expressed genes in MS identified in Tn cells and >400 differentially expressed genes in monocytes, only 23 genes were shared between these two cell types (Table 2). Among these genes, *ZBTB16* and *HMGB2* were differentially expressed in all 3 primary cell types, in addition to PBMC. *ZBTB16* was also differentially expressed in stimulated monocytes (Figure 2E). *ZBTB16* (Zinc finger and BTB

domain-containing 16), also known as *PLZF* (Promyelocytic Leukemia Zinc Finger) or *ZNF145*, is previously reported to be among the top markers associated with MS and a number of other autoimmune diseases: hypomethylation in a *ZBTB16* enhancer region is reported in PBMC in MS[17], and *ZBTB16* downregulation is observed in systemic lupus erythematosus (SLE) B cells[18] and type1 diabetes (T1D) PBMC[19]. Our results on the downregulation of *ZBTB16* in T cells, monocytes, and PBMC in MS are in line with the previous literature on this gene, and suggest an important role for *ZBTB16* in dysregulated immune response.

Table 2. Genes associated with MS in both T cells and monocytes.

HGNC symbol	Ensembl ID	chromosome	t stats. Tn	t stats. Tmem	t stats. Mono	t stats. PBMC
ACTR3	ENSG00000115091	2	**3.74**	3.2	**5.95**	2.84
BRD7	ENSG00000166164	16	**3.46**	-0.02	**4.87**	0.94
CASS4	ENSG00000087589	20	**-4.74**	**-4.89**	**-3.48**	-2.56
CCND3	ENSG00000112576	6	**-6.35**	**-4.69**	**-5.01**	-3.08
CIRBP	ENSG00000099622	19	**-3.48**	-1.9	**-4.65**	-2.29
DDIT4	ENSG00000168209	10	**-5.53**	-2.93	**-5.73**	-3.83
DDX58	ENSG00000107201	9	**4.06**	3.43	**3.32**	2.97
FKBP5	ENSG00000096060	6	**-5.44**	**-5.06**	**-4.93**	-3.59
HMGB2	ENSG00000164104	4	**-4.01**	**-4.07**	**-3.56**	**-3.94**
KLF9	ENSG00000119138	9	**-4.86**	-3.62	**-4.73**	-2.19
LCP2	ENSG00000043462	5	**3.39**	2.39	**3.22**	0.12
NDUFAF5	ENSG00000101247	20	**-4.34**	-1.8	**-3.65**	0.48
SESN3	ENSG00000149212	11	**-3.39**	**-3.89**	**4.63**	-1.96
SH2D3C	ENSG00000095370	9	**3.6**	3.22	**3.24**	2.09
SLA	ENSG00000155926	8	**-4.01**	**-4.16**	**-3.99**	-2.79
SNN	ENSG00000184602	16	**-4.14**	-3.07	**-3.47**	**-4.09**
TIPARP	ENSG00000163659	3	**-3.68**	-3.54	**-4.28**	-0.18
TSC22D3	ENSG00000157514	X	**-3.98**	-3.15	**-4.36**	-2.16
TUBA4A	ENSG00000127824	2	**-4.79**	-2.89	**-3.97**	-2.14
TXLNA	ENSG00000084652	1	**4.16**	3.06	**4.19**	2.37
VSIR	ENSG00000107738	10	**-4.42**	**-4.3**	**-4.27**	-3.67
ZBTB16	ENSG00000109906	11	**-6.9**	**-4.45**	**-7.27**	**-4.12**
ZNF564	ENSG00000249709	19	**-3.98**	-3.17	**-3.47**	-1.9

Bold font represents adjusted *p* <0.05.
PBMC samples are from the replication study.

Finally, we performed pathway enrichment analysis on the differentially expressed genes from each cell-type-state in order to identify shared and distinct pathways involved in immune dysregulation in T cells and monocytes in MS. The top enriched gene ontology (GO)[20] biological processes (adjusted *p* <0.05) for T cells and monocytes were T cell activation and myeloid cell activation, respectively. No enriched GO pathways were found for genes from the stimulated monocytes. Additional pathway enrichment analysis using MsigDB hallmark gene sets[21] identified significant enrichment for several hallmark pathways among the differentially expressed genes from Tn cells and monocytes (Figure 2F, Supplementary Table 3), several of which were previously implicated in MS[16,22]. However, our analysis provides further information on the cell-type-specificity of their involvement in MS pathology. Pathways such as PI3K/AKT/mTOR and mTORC1 signaling, inflammatory response, and transplant rejection were enriched among differentially expressed genes from both cell types. On the other hand, pathways such as IFN-γ response and complement system were enriched among differentially

expressed genes from Tn cells, but not monocytes; and genes from TNF-α NF-κB signaling, IL2 STAT5 signaling, hypoxia, and glycolysis and gluconeogenesis pathways were enriched among genes from monocytes, but not Tn cells.

T cell and monocyte co-expressed gene modules associated with MS
To complement our gene-level results, we elected to complete analyses after implementing a dimension reduction approach to help boost our statistical power to discover novel patterns of perturbed transcriptomic responses in MS. Specifically, we performed weighted gene co-expression network analysis (WGCNA)[23] to (1) identify sets of co-expressed genes (gene modules) in each of the 6 cell-type-state transcription profiles, and (2) determine the association of the identified gene modules with MS. Given that biological processes often involve activation and/or suppression of several genes in concert with each other, the agnostic identification of groups of co-expressed genes in our data could potentially identify novel pathways that may be missing from the above-used predefined gene sets (i.e., GO biological processes and MsigDB hallmark).

Our WGCNA analysis identified a total of 197 co-expressed gene modules from all 6 cell-type-states. Of note, our modules were set to include both positively- and negatively- correlated genes. The list of genes belonging to each module and the top 10 GO gene sets enriched in each module can be found in Supplementary Data. We note that at least one identified module from each cell-type-state was highly associated with sex (Supplementary Table 4) and mainly consisted of known sex-associated genes from X and Y chromosomes. This observation supports the biological relevance of our identified modules.

Of our 197 gene modules, 6 modules were significantly associated with MS (adjusted $p <0.05$, Figure 3A, Supplementary Table 4). The association analysis was performed between the eigengene from each module (i.e., the first principal component of the expression of the genes in each module) and MS diagnosis. The hub gene of each MS-associated module (i.e., a highly connected gene from each module, identified as part of the WGCNA analysis) is shown in Figure 3A and is used for module annotation throughout the rest of the manuscript. Secondary association analysis using module eigengenes demonstrated that the 6 MS-associated modules were significantly different in both untreated MS vs. healthy and GA-treated MS vs. healthy comparisons; and no module showed significant difference between untreated and GA-treated MS participants (Supplementary Table 4). These findings are in line with our observations from the transcriptome-wide association analysis on the similarity between untreated and GA-treated MS transcription profiles.

Comparing the list of genes from the MS-associated modules, we found that the Tn TNFSF8 module (n=54 genes) and the Tmem FKBP5 module (n=30 genes) shared a large number of genes (n=11). Correlation analysis between the module eigengenes further supported the similarity between these modules, as these modules were the only modules that were highly correlated (Spearman's ρ >0.5, $p <0.01$) both among MS and healthy participants (Figure 3B). The two modules were the top associated modules with MS ($p =2.8\times10^{-17}$ and 1.4×10^{-9}, respectively), and they were both enriched for genes from GO annotation's T cell activation pathway.

Figure 3C further visualizes the association between individual genes from each MS-associated module and MS diagnosis (adjusted *p*-value and log fold change from the transcriptome-wide association analysis; differentially expressed genes are labeled). In addition to recapitulating some of the previous enrichment results from the differentially expressed genes, hallmark enrichment analysis on genes from the MS-associated modules provided support for the involvement of additional pathways such as oxidative phosphorylation and adipogenesis in MS. This suggests that although these pathways were not significantly enriched among the differentially expressed genes, the coordinated subthreshold change in the expression of their genes in MS could be detected using the co-expression analysis. Both of these pathways were previously implicated in MS pathology[24–26]. Additional enrichment analysis using genes associated with MS susceptibility genetic loci[2] revealed a nominal enrichment for these genes in the Tn TNFSF8 module (*p* =0.02, Figure 3C), suggesting that the transcriptional changes in this module may provide a link between genetic susceptibility to MS and MS onset.

Replication of module associations with MS in independent data
We used available RNA-seq data from independent PBMC and macrophage samples from MS and healthy participants in order to replicate the association between our modules and MS. Given that PBMC consist of both lymphocytes and monocytes, we used the PBMC data for replication analysis for all 6 MS-associated modules. However, the macrophage data were only used to assess replication for the 3 monocyte associations.

The PBMC data used for the replication study were part of the previously mentioned PBMC data used in the transcriptome-wide association analysis. However, we excluded participants that had samples in the 6 cell-type-state study so that only independent participants would be assessed in the replication analysis. Module eigengenes (i.e., the first principal component) were calculated and used for the association analysis. We found significant evidence of replication for the Tn TNFSF8 module (n=24 GA-treated MS vs. n=12 healthy participants, *p* =0.01). Additionally, we observed suggestive evidence that the Tmem FKBP5 module (n=24 GA-treated MS vs. n=10 healthy participants, *p* =0.09) may also replicate if larger sample sizes are assessed (Figure 3D).

The macrophage data used for the replication of the MS-associated monocyte modules were from a separate study of monocyte-derived macrophages, which was performed and analyzed by independent investigators[25]. CD14[+] monocytes from MS and healthy participants were isolated from peripheral blood and differentiated into macrophages through *in vitro* exposure to GM-CSF (details in the Methods, Table 1). The eigengene for the stimulated monocyte DBNL module was significantly different in this dataset which consists of healthy (n=11) and both untreated (n=11) and treated (n=17) MS participants: *p* =0.02 for the comparison to untreated MS and *p* =0.05 for treated MS (Figure 3E).

Given our moderate sample sizes in these replication efforts and the fact that the cell types from both of our replication datasets were different from the cell types used in the discovery analysis, the other non-significant results do not rule out that the nonreplicated modules are truly perturbed in MS. Future dedicated larger studies will have to answer that question. However, our replication analyses did return significant results for certain modules in both T cells and monocytes; the T cell TNFSF8/FKBP5 modules and the monocyte DBNL module are therefore prioritized as signatures of transcriptional perturbation in the asymptomatic phase of MS (i.e., in

clinical remission). They therefore offer tools for further investigation of the effects of immunomodulatory treatments now routinely used in MS.

Computational drug screen for reversing MS-associated modules
We then queried the Connectivity Map (CMap)[27] database to identify compound classes with reverse transcriptional signatures to each of the MS-associated modules. CMap contains data from the transcriptional effects of ~5,000 small-molecule compounds, each tested in multiple cell lines (although none of which are of leukocyte origin). The top significant compound classes identified for each module are shown in Figure 4A. Intriguingly, glucocorticoid receptor agonists were found as the top compound class for the Tn TNFSF8 and Tmem FKBP5 modules. Among the prioritized compound classes, glucocorticoid receptor agonists are the only drugs that are currently used in clinical management of MS. Our finding suggests that this class of drugs might exert part of their therapeutic effect by upregulating these T cell modules that we identified and validated as being downregulated in MS.

Given this finding, we searched the literature for studies on transcriptional effects of glucocorticoids. We took two gene sets from two separate studies[28,29] on *in vitro* effects of the glucocorticoid drug prednisolone on gene expression in immune cells in order to assess the enrichment of our modules for glucocorticoid response genes. The two studies were performed using different cell types (primary leukemic cells from acute lymphoblastic leukemia participants[28] and PBMC from healthy individuals[29]), and the two sets of differentially expressed genes after glucocorticoid treatment (n=51 and 82 genes, respectively) had only 6 genes in common (Supplementary Table 5). However, the enrichment analysis using both gene sets demonstrated similar results: among all 197 modules, Tn TNFSF8 and Tmem FKBP5 modules were the only modules with significant enrichment for glucocorticoid response genes (adjusted p <0.05, Figure 4B). Combining the genes from the two sets resulted in even more significant results ($p = 2.5 \times 10^{-13}$ and 5.9×10^{-7} for Tn TNFSF8 and Tmem FKBP5 modules, respectively). Among the MS-associated modules, we also found a nominal significance for enrichment in glucocorticoid response genes in monocyte GABARAPL1 module ($p = 0.004$ for the combined gene set).

Given that MS relapse is treated with glucocorticoids, the majority of MS participants have a positive history of previous glucocorticoid treatment. However, none of our MS participants had experienced a relapse in <3 months prior to blood sampling. Given that the biological half-life of glucocorticoids is <72 hours[30], and the transcriptional effects of oral glucocorticoids are shown to reverse to baseline in <48 hours[29,31], the perturbation of glucocorticoid response genes in our MS samples is unlikely to be a long-lasting effect of previous glucocorticoid treatment. This is further supported by a lack of significant difference in the Tn TNFSF8 and Tmem FKBP5 module eigengenes between MS participants who had a relapse in the last 3-12 months (n=16) vs. in >12 months (n=127) prior to blood sampling ($p = 0.68$ and 0.84, respectively; Supplementary Figure 3). On the other hand, the fact that the genes in common between the MS-associated modules and the glucocorticoid response gene sets showed an opposite direction of effect (i.e., they were all downregulated in MS and upregulated in response to glucocorticoid treatment), suggests an intrinsic defect in glucocorticoid response in MS. This is in line with previous literature on the decreased glucocorticoid sensitivity observed in MS PBMC[32,33].

Validation of glucocorticoid effect on gene expression in MS
In the previous sections, our analyses identified genes belonging to MS-associated modules that are downregulated in MS and are shown to be upregulated in response to *in vitro* glucocorticoid treatment in two non-MS studies[28,29]. Given the differences between glucocorticoid response in leukemic cells[28] and cells from healthy participants[29], one can assume that glucocorticoid response in MS may also be to some extent different. Hence, we decided to assess the effect of glucocorticoid treatment on the expression of genes from Tn TNFSF8 and Tmem FKBP5 modules in samples from MS participants. Our experiment on the effect of *in vitro* exposure to the glucocorticoid drug dexamethasone on PBMC samples from MS participants (n=5) demonstrated a significant increase in the expression of T cell hub genes *TNFSF8* and *FKBP5* across all tested doses (all p <0.01, Figure 4C). Unlike *FKBP5*, *TNFSF8* was not part of our two glucocorticoid response gene sets[28,29]. However, the upregulation of both hub genes in response to dexamethasone in samples from MS participants provides support for the mechanism through which glucocorticoids might exert part of their therapeutic effect in MS.

T cell and monocyte MS-associated glucocorticoid response genes
As a follow up analysis to our finding on the enrichment of glucocorticoid response genes in MS-associated modules, we sought to investigate whether similar enrichment could be observed in all MS-associated genes in T cells and monocytes. Our analysis revealed significant enrichment for glucocorticoid response genes (combined set, in addition to *TNFSF8*) among differentially expressed genes in all 3 primary cell types (p =9.9×10^{-8}, 3.9×10^{-6}, and 3.1×10^{-11} for Tn, Tmem, and monocytes, respectively). However, we observed an interesting difference in the pattern of enrichment between T cells and monocytes: the majority of the differentially expressed glucocorticoid response genes in T cells were co-expressed and belonged to a single MS-associated module, while this was not the case for monocytes (Figure 4D). This explains the lack of a prominent glucocorticoid responsive MS-associated module in monocytes, and on the other hand suggests that glucocorticoid treatment acts via a single pathway in MS T cells. This observation further provides human support for results from a previous knockout study[34] on a mouse model of MS, in which the therapeutic effect of glucocorticoids was demonstrated to be dependent on the presence of glucocorticoid receptors in T cells but not myeloid cells.

Gene and module associations with MS susceptibility, severity, and progression
In addition to investigating associations between individual genes and gene modules with MS diagnosis, we performed exploratory analyses within the MS samples in association with several MS-related variables: (1) MS susceptibility assessed using MHC, non-MHC, and total MS polygenic scores; (2) MS severity measured using the Expanded Disability Status Scale (EDSS), brain parenchymal fraction (BPF) from magnetic resonance imaging, and T2 hyperintense white matter lesion burden; and (3) MS progression assessed using change in EDSS over 4 years of follow up, and the slope of change in brain parenchymal fraction and white matter lesion volume. None of our 197 modules showed significant association with any of the outcomes (all adjusted p >0.05). This might be explained by our small sample size for identifying heterogeneity in the disease and the low variability within our MS participant population who were mostly in the early phase of their disease. However, our gene-level analyses identified a few significant associations (Supplementary Table 6). The most prominent associations were observed between the MS MHC polygenic score and the expression of MHC class II genes *HLA-DRB1* and *HLA-DRB5*. These associations were found in all cell types except for Tn cells, and

are in line with previous literature on the effect of the MS top HLA risk allele HLA-DRB1[*]1501 on the expression of these genes[35]. Other associations were less significant and need to be replicated in independent MS datasets.

Discussion

Our study of transcriptional profiles from three cell types in both unstimulated and stimulated conditions provides new insights into perturbation of transcriptional programs in certain immune cells implicated in MS. Using a relatively large cell-type-state-resolved transcriptomic dataset of MS and healthy participants (1,075 transcriptomic profiles from 209 individuals), we identified shared and distinct changes in individual genes, biological pathways, and co-expressed gene sets in MS T cells and monocytes. We replicated part of our results in independent data from PBMC and monocyte-derived macrophages. Further, we complemented our results by performing a computational drug screen and validating its findings using independent enrichment analyses and *in vitro* experiments.

Our analyses demonstrated that primary T cells and monocytes purified from peripheral blood show largely distinguishing transcriptomic signatures in MS, supporting their use as accessible cell models for studying MS immunopathology and its treatment. On the other hand, despite the large number of MS-associated genes in primary T cells and monocytes and their enrichment for T cell and myeloid activation biological pathways, we did not find similarly large differences in our *in vitro* stimulated cell types in MS: few genes were found to be differentially expressed in MS in TNF-α stimulated monocytes, and none were in anti-CD3/CD28 stimulated T cells. This observation suggests that *in vitro* stimulation of T cells with anti-CD3/CD28 antibodies does not provide a proper transcriptomic model for studying MS T cell dysfunction[33,36]. Further, it highlights the fact that more accurate T and myeloid cell activation models are required for MS-related investigations.

Our transcriptome-wide association studies on data from unstimulated cell types identified >200 MS-associated genes in T cells and >400 genes in monocytes. The majority of these genes show a cell-type-specific association with MS. However, results from pathway enrichment analysis suggest that in addition to affecting distinct pathways in different cell types, part of the T cell- and monocyte- specific perturbations converge on the same biological pathways across both cell types. These shared pathways include PI3K/AKT/mTOR and mTORC1 signaling[22], transplant rejection, and inflammatory response. Similar finding was observed between MS-associated T cell MYL6 and stimulated monocyte TMEM179B modules: despite sharing few genes, both modules are enriched for oxidative phosphorylation and adipogenesis pathways[24–26]. These observations demonstrate that shared pathways can be affected through distinct patterns of transcriptional changes across different cell types, and may need to be assessed in a cell-type-specific manner when searching for treatment strategies.

Our dimension reduction approach of identifying co-expressed gene modules complemented the results of our gene-wise studies and identified 6 MS-associated modules. Each module contained a subset of the MS-associated genes with additional co-expressed genes, and therefore provided data-driven information on coordinated gene expression perturbations associated with the disease. The associations for two T cell and one monocyte modules were replicated in independent samples from other immune cells, increasing confidence in their relevance to MS. Follow-up analysis using computational drug screen suggested that the replicated MS-associated T cell modules could be reversed using glucocorticoid receptor agonists. This finding was further validated by the significant enrichment of these modules for glucocorticoid response genes

extracted from two independent studies, in addition to our own *in vitro* experiment on the effects of glucocorticoid treatment on gene expression in MS.

Most MS patients have a positive history of treatment with glucocorticoids during relapse. However, (1) the opposite direction of gene expression effects observed in MS and glucocorticoid treatment[28,29] (i.e., downregulation in MS and upregulation in response to glucocorticoids, and vice versa), (2) the short half-life of glucocorticoid effects on gene expression[29,31], (3) the long intervals between the last relapse and blood sampling in our MS participants, and (4) the lack of association between module expressions and closer relapse history, altogether argue against the confounding effect of previous glucocorticoid treatment in our data, and in favor of a true biological mechanism for the therapeutic effect of glucocorticoids in MS. Thus, our findings, for the first time reveal the coordinated perturbation of glucocorticoid response genes in MS T cells, and provide a novel connection between transcriptional effects of glucocorticoids and MS immune pathology.

Impairment in the hypothalamic-pituitary-adrenal (HPA) axis has long been documented in MS: Individuals with MS have elevated serum adrenocorticotropic hormone (ACTH), increased cerebrospinal fluid and serum cortisol (the main human glucocorticoid), and impaired response to dexamethasone suppression and dexamethasone-suppressed corticotrophin-releasing hormone tests, coupled with decreased glucocorticoid receptor affinity and glucocorticoid sensitivity in PBMC[32,33,37]. These differences are found in all clinical subsets of MS, including individuals with primary-progressive disease course, who have no relapse and glucocorticoid treatment history[32]. The downregulation of glucocorticoid response genes in MS T cells and monocytes observed in our study can likely be explained by the decreased sensitivity to glucocorticoids. However, this needs to be validated in future studies where HPA axis hormones and glucocorticoid sensitivity are measured simultaneous with transcriptome. The exact mechanisms underlying the decreased glucocorticoid sensitivity are also unclear. Previous studies have suggested that a decrease in the number of glucocorticoid receptors is unlikely to be the cause, as the density of glucocorticoid receptors is not affected in MS PBMC[32]. Similarly, the expression of *NR3C1*, the gene encoding the glucocorticoid receptor, was not different in our MS samples. However, we observed differential expression in a number of other genes involved in the formation or function of the glucocorticoid receptor complex: *HSP90B1* (encoding a chaperone for glucocorticoid receptor) in upregulated in MS T cells, *FKBP5* (encoding a co-chaperone of HSP90) in downregulated in MS T cells and monocytes, and *HMGB2* (encoding a non-histone chromosomal protein which facilitates DNA binding of glucocorticoid receptor) is downregulated in MS T cells, monocytes, and PBMC. Whether these changes are linked to the mechanism underlying decreased glucocorticoid sensitivity in MS needs to be investigated in future studies.

The ER22/23EK polymorphism of the glucocorticoid receptor gene is previously associated with decreased glucocorticoid sensitivity and faster MS progression (i.e., shorter time to EDSS=6)[38]. The suggested relation between glucocorticoid sensitivity and disease progression was not replicated in our data, as we did not observe an association between the glucocorticoid-responsive T cell modules and MS clinical and MRI severity and progression measures. However, our results might have been affected by the limited follow up period available for our participants, as only a few of them had progressed to EDSS=6 during the 4-year period.

Additional studies have suggested associations between increased cortisol levels, decreased glucocorticoid sensitivity, and MS activity (i.e., clinical relapse or the presence of gadolinium-enhancing lesions on MRI)[32,39,40]. None of our participants were experiencing clinical relapse at the time of blood sampling, and we did not have available data on other biomarkers of disease activity[41,42] such as cerebrospinal fluid neurofilament light chain (NfL) levels and gadolinium-enhancing MRI lesions. However, the relation between markers of MS activity and glucocorticoid-responsive T cell modules needs to be investigated in future studies using longitudinal measurements. A significant association with MS activity may have important implications for the utility of the identified modules as molecular indicators for treatment modification or glucocorticoid administration.

Although our work benefited from a relatively large sample size, minimization of treatment confounding effects, cell type- and state-specific data generation, complementary analytical approaches, and replication and validation experiments, our study had several limitations, some of which were discussed above. These included the limitations of the studied *in vitro* stimulation models and the unavailability of measures assessing HPA axis impairment and MS activity. Another major limitation of our study was that it did not include data on other peripheral immune cell types implicated in MS pathology such as B cells, plasma cells, and cytotoxic T cells[16]. Further, the use of bulk RNA-seq data from purified $CD4^+$ T cells and $CD14^+$ monocytes might have masked more fine-grained associations that could potentially be observed in subtypes of these cells. Technological advances helping with feasibility of large-scale and high-resolution single-cell RNA-seq studies can alleviate both of these issues in future investigations. Finally, the enrichment of the glucocorticoid-responsive T cell TNFSF8 module for MS susceptibility genes suggests a causal role for glucocorticoid response dysfunction in MS pathology. However, additional studies are required to elucidate the cascade of pathological vs. compensatory molecular events involved in glucocorticoid response pathology in MS, and to elaborate on the serial vs. parallel nature of transcriptional changes observed in different cell types.

Notwithstanding the limitations, our study has helped resolve cell-type-specific perturbations in transcriptomic programs in MS helper T cells and monocytes. Additionally, our approach of combining data-driven identification of coordinated transcriptional changes with computational drug screen has helped provide a novel explanation for the therapeutic effect of glucocorticoids in connection with MS T cell pathology. Our dataset can continue to be used by the scientific community as a resource for replication or meta-analysis studies, or further analyzed in conjunction with other molecular datasets for hypothesis testing or discovery. Our analytical approach can also be used as a working example for disentangling the mechanisms underlying other disease-related conditions. Finally, given the adverse effects associated with chronic use of glucocorticoids[43], our findings can help ground future studies of dose optimization, compound modification[44] or drug combination with the purpose of improving treatment strategies to achieve higher disease-specific and fewer off-target effects.

Methods

Participants

Multiple sclerosis (MS) participants were from the Comprehensive Longitudinal Investigation of Multiple Sclerosis at the Brigham and Women's Hospital (CLIMB) study[11]. CLIMB is a natural history observational study of MS, in which participants undergo semi-annual neurological examinations and annual magnetic resonance imaging (MRI). The participants also donate blood, from which peripheral blood mononuclear cells (PBMC) are extracted and cryopreserved. In 2016-2017, cryopreserved PBMC samples from participants meeting the following criteria were pulled from the archive for sample processing: (1) age 18-60 years old at the time of sampling, (2) a diagnosis of MS based on the 2010 McDonald criteria, (3) a relapsing-remitting disease course, (4) being untreated for >6 months or on treatment with glatiramer acetate (GA) for >12 months at the time of sampling, (5) no evidence of disease activity in the 6 months prior to sampling, (6) no steroid use in the 30 days preceding sampling, and (7) an Expanded Disability Status Scale (EDSS) score between 0 to 3 at the time of sampling.

Healthy participants were from the PhenoGenetic Project, part of the ImmVar study[13,14]. PBMC samples were from participants 18-60 years old who did not have any immune, neurologic, metabolic, or cardiovascular disease, and were not under treatment for any acute or chronic medical conditions.

Both studies were approved by the Institutional Review Board of the Brigham and Women's Hospital, and all participants had signed a written informed consent. PBMC samples from 227 participants who fulfilled the criteria (181 MS and 46 healthy participants) were used for further processing. 90% of samples were collected between 2008-2014.

PBMC sampling

PBMC were collected prospectively from participants using the Immune Tolerance Network (https://www.immunetolerance.org/) protocol to ensure data quality and to minimize variation among samples. In short, fresh blood was processed within 4 hours using the Ficoll procedure to extract PBMC. The PBMC were then resuspended in fetal bovine serum with 10% DMSO and cryopreserved in liquid nitrogen.

T cell and monocyte isolation and stimulation

PBMC samples meeting study criteria were pulled from the sample archive and thawed in batches. Cell viability was assessed by acridine orange/propidium iodide staining. Naïve and memory CD4$^+$ T cells and CD14$^+$ monocytes were isolated on a BD FACSAria flow cytometer using fluorochrome labelled anti-CD3 (1:50 dilution, 2 μl to 100 μl staining volume containing 1×10^6 cells), anti-CD4 (1:50 dilution), anti-CD45RA (1:33 dilution), anti-CD45RO and anti-CD14 (1:xx dilution) antibodies. Cells from each of the 3 cell types were then split in half and placed in culture with serum-free X-Vivo medium. Stimulation was performed for one half using 2 μg/mL anti-CD3 and 1 μg/mL anti-CD28 antibodies for T cells and xx μg/mL TNF-α for monocytes. Cells from both stimulated and unstimulated conditions were collected after 24 hours.

RNA-seq data generation and preprocessing

RNA was extracted from all primary and *in vitro* stimulated samples using xx kit and sent for sequencing. Whole transcriptome 25-bp paired-end sequencing was performed for 1176 samples using the Broad Institute HiSeq 2500 platform to an average depth of 15 million reads per sample. Processing was performed according to the Broad Institute RNA-seq pipeline for the GTEx Consortium[45]. Briefly, RNA sequence reads were aligned to the GRCh38/hg38 genome reference using STAR[46], quality control was performed using RNA-SeQC[47] and quantification of gene expression levels were performed using RSEM[48].

Prior to grouping samples from each cell-type-state for the rest of the preprocessing and analyses, data from participants who were from non-European ancestry (determined based on self-reported ethnicity and/or genetic data; details below; n=17) or had incomplete or mismatching demographic information (n=1) were excluded. Subsequent processing was performed on data from each group, separately, as follows. Transcripts with low expression values (average TPM <2) were removed. TPM values were log-transformed and quantile-normalized. Samples with outlier principal component (PC) values (>3 standard deviation on PCs that explained >5% of the variance) were further excluded. Expression levels were adjusted for the effects of technical confounding variables, i.e., library preparation batch and RNA-seq quality metrics that were associated with the first 10 PCs of expression data, such as the number of aligned reads, percentage of coding, intronic, and intergenic bases, median coverage variability, and median 3' and 5' biases. Residual expression values were used for the association analyses.

Genetic data preprocessing and imputation
Genome-wide genotyping data were available for 218 participants as part of larger batches of previously genotyped CLIMB and ImmVar samples. Genotyping was performed using Illumina MEGA[EX], Infinium OmniExpressExome, and Affymetrix 6.0 arrays. Preprocessing of the genetic data was performed using PLINK[49]. SNPs with call rate <95% were removed. Multidimensional scaling was performed using HapMap3 reference data. Samples from individuals of non-European ancestry and ethnic outliers with >3 standard deviation difference from the European samples were identified and both their RNA-seq and genetic data were removed (n=17). Additionally, genetic data from samples with genotyping rate <90%, mismatch between recorded sex and genetic sex, and low or high heterozygosity rates (>3 standard deviation) were excluded (n=4). No remaining samples were from related individuals (all pi-hat <0.125). Imputation was performed for each genotyping batch separately. Whole-genome imputation was performed using the Michigan Imputation Server[50] and the Haplotype Reference Consortium (HRC) panel v1.1. Imputed data from the study participants were then merged, and SNPs with low imputation quality (R^2 <0.8) were removed. SNPs, amino acids and HLA alleles in the MHC region were imputed using SNP2HLA[51] and the Type 1 Diabetes Genetics Consortium (T1DGC) HLA reference panel.

MS susceptibility polygenic scores
MS polygenic scores were calculated as described previously[52]. Briefly, genome-wide significant results from the 2019 International Multiple Sclerosis Genetics Consortium (IMSGC) MS genome-wide association study[2] (or if not available, tag SNPs in LD >0.8) were used for polygenic score calculation. MS MHC polygenic scores were calculated as the sum of the imputed dosages of 31 MHC markers associated with MS multiplied by their effect sizes (i.e.,

log odds ratio). Similarly, MS non-MHC polygenic scores were weighted scores calculated from the imputed dosages of 193 autosomal non-MHC MS-associated SNPs (7 loci did not have high-quality tag SNPs in our imputed dataset, and therefore were not included in polygenic score calculation). Finally, MS total polygenic scores were calculated as the sum of MHC and non-MHC polygenic scores.

EDSS outcomes
179 MS participants had available EDSS data within 3 months of the blood sampling. From these participants, 132 were followed up for ≥4 years after the blood sampling visit and had at least 4 available longitudinal EDSS measurements (median=21 data points per person including visits prior to the time of blood sampling). In order to overcome the EDSS measurement variability and minimize the effect of temporary outlier measurements, smoothing splines were fit to longitudinal EDSS data using R smooth.spline function, with equivalent degree of freedom set to 4 (Supplementary Figure 4). EDSS progression was defined as the difference between smoothed EDSS measurements at year 4 and the time of blood sampling.

MRI outcomes
Proton density-, T2- and T1-weighted brain MRI was performed for MS participants. MS lesions and brain tissue compartments (gray matter, white matter, and cerebrospinal fluid) were segmented using template-driven segmentation and partial volume artifact correction (TDS+)[53] method. Results underwent quality control and manual correction where necessary. Brain parenchymal fraction (BPF) was calculated as the sum of gray and white matter volumes divided by the intracranial volume (i.e., the sum of gray matter, white matter, and cerebrospinal fluid volumes).

120 MS participants had available MRI data acquired using 1.5T scanners within 1 year of the blood sampling (MRI performed using 3T scanners were few and were excluded from this study). BPF and log-transformed lesion volume from these participants were used as surrogate quantitative markers for brain atrophy and white matter demyelination, respectively. From these participants, 93 had at least 4 available longitudinal MRI data acquired using 1.5T scanners during their follow-up period (median=6 data points per person including visits prior to the time of blood sampling). The rate of progression in brain atrophy and white matter demyelination were calculated for these participants as the linear slope of change in BPF and log-transformed lesion volume over the follow-up period, respectively (Supplementary Figure 4).

Transcriptome-wide association studies
Transcriptome-wide differential expression analyses between diagnostic and treatment groups were performed using limma[54] robust regression modeling. All comparisons included age and sex as covariates. Adjustment for multiple comparisons was performed for the genes tested in each cell-type-state. Associations with false discovery rate (FDR)-adjusted $p < 0.05$ in each comparison were considered significant.

Within-MS transcriptome-wide association studies were performed using the following quantitative outcome variables: MHC, non-MHC, and total MS susceptibility polygenic scores; baseline EDSS, BPF, and lesion volume; and longitudinal change in EDSS, and the rate of change in BPF and lesion volume. Outlier outcome variables were winsorized in case of $|>3|$

standard deviation difference from the mean. Association analyses included additional covariates depending on the outcome variable: all within-MS associations accounted for the effects of disease duration and treatment group, in addition to age and sex; association analyses on longitudinal outcomes (i.e., 4-year change in EDSS, and the rate of change in BPF and lesion volume) additionally accounted for the effects of their related baseline measurements. FDR-adjusted *p*-values from each transcriptome-wide study underwent an additional level of adjustment for the 9 tested outcomes for each cell-type-state.

Co-expressed gene modules identification and analyses

R package WGCNA[23] (weighted gene co-expression network analysis) was used to identify modules of co-expressed genes in data from each cell-type-state. WGCNA partitions the transcriptome into modules of highly co-expressed genes, where each gene is either part of a co-expression module or stays unassigned. Therefore, not all genes are assigned a module, and each gene can only belong to one module. Biweight midcorrelation (bicor) with outlier removal >5 percentile was used to avoid sensitivity to outlier expression values. Modules were set to be unsigned (i.e., contained both positively and negatively correlated genes). Minimum module size (number of genes in each identified module) was set to 20. Module eigengenes (first principal component of the genes belonging to each module) were calculated using WGCNA and used for association analyses with MS diagnosis and MS-related outcomes. Associations were assessed using robust regression modeling. Similar to the transcriptome-wide association studies, quantitative MS-related outcomes were winsorized when necessary, and covariates were included in the model depending on the outcome variable. Adjustment for multiple comparisons was performed using Bonferroni method for the tested modules for each outcome.

Gene set enrichment analyses

Gene ontology (GO)[20] and MsigDB hallmark[21] gene set enrichment analyses were performed using FUMA[55]. Minimum overlap with each gene set was set to 3 genes, and results were reported at 5% FDR. Enrichment analyses for the other gene sets (i.e., glucocorticoid response and MS susceptibility-associated genes) were performed using hypergeometric test. The two sets of glucocorticoid response genes were obtained from [28,29]. Genes associated with MS susceptibility loci were obtained from [2].

Compound screen using Connectivity Map (CMap)

The Connectivity Map (CMap)[27] database contains data on the transcriptional effects of ~5,000 small-molecule compounds, each tested in multiple cell lines. The database can be queried using the CLUE (CMap and LINCS Unified Environment, https://clue.io/) platform. We queried the database (v1.3) using genes from each MS-associated module to identify compound classes with reverse transcriptional similarity to each module. Results with FDR-adjusted $p < 1 \times 10^{-4}$ are reported.

Replication dataset #1: PBMC RNA-seq

Similar to the main study, PBMC samples used in this dataset (n=65) were from CLIMB (n=29) and PhenoGenetic (n=36) participants. However, they were retrieved and sequenced separately as part of a larger batch for a different PBMC study. The inclusion/exclusion criteria were similar to the main study, except that the MS participants were all GA-treated. RNA extracted from these samples underwent the same sequencing and preprocessing pipelines as the above.

Age and sex were used as covariates for all association analyses. Transcriptome-wide differential expression analysis between MS and healthy participants was performed using the above-outlined limma pipeline. Eigengene association studies were performed on the subset of samples that did not have paired data with the main study. Linear regression modeling was used to compare eigengene values between diagnostic groups.

Replication dataset #2: macrophage RNA-seq
Data used in this dataset (n=39) were from a study on monocyte-derived macrophages, which was performed and analyzed by independent investigators[25]. Methods for this study are described in details elsewhere[25]. Briefly, the study was approved by the French Ethics committee, and written informed consent was obtained from all study participants. Recruited MS participants fulfilled the diagnostic criteria for MS, and none of the participants had any other inflammatory or neurological disorders. CD14[+] monocytes were isolated from the peripheral blood of MS and healthy participants and differentiated into macrophages through *in vitro* exposure to granulocyte macrophage colony-stimulating factor (GM-CSF, 500 U/ml, ImmunoTools). Cells were lysed after 96 hours, and extracted RNA underwent 75-bp single-end sequencing using a NextSeq 500 sequencer to an average depth of 60 million reads per sample. Data quality control was performed using fastp[56], alignment was performed to the GRCh38/hg38 genome reference using STAR[46], and gene expression quantification was performed using RSEM[48]. Sequencing batch effects were corrected using the removeBatchEffect function of the limma[54] package, and corrected TPM values were used in the replication analyses. Wilcoxon non-parametric sign rank test was used to compare eigengene values between groups.

Dexamethasone experiment
In vitro experiments were conducted on PBMC samples from MS participants recruited at the Columbia University MS Center (n=5). Blood sampling and PBMC isolation were performed using the same protocol as the main study. The isolated cells were incubated for 24 hours with dexamethasone (MicroSource Discovery System, Cat# 01500203) at five concentrations: 0, 10, 50, 75, or 100 µM in DMSO. RNA was isolated from the cells using RNeasy plus micro kit (QIAGEN, Cat# 74034) according to the manufacturer's protocol, and reverse transcribed to cDNA using iScript cDNA synthesis kit (Bio-Rad, Cat# 1708891). TaqMan gene expression assays HS00174286_m1 and HS01561006_m1 were used to quantify the expression of *TNFSF8* and *FKBP5* genes in each sample. Expression levels were normalized to the expression of the housekeeping gene *B2M*. Quantitative RT-PCR was performed using an Applied Biosystems QuantStudio 3 Real-Time PCR System. Differences in gene expression between dexamethasone-treated samples and the vehicle control group were assessed using *t*-test.

Statistical analysis
All statistical tests are described in their respective methods sections. All reported *p*-values are two-sided.

Data availability

The RNA sequencing data (1075 samples from 6 cell-type-states from 209 participants) will be deposited in the Synapse database under accession codes xx [link]. As sensitive human data, these data can be accessed upon request, following the establishment of a Data Use Agreement with Brigham and Women's Hospital, Dr. Howard Weiner. Supplementary Data will be deposited to the Synapse database under accession code xx [link] and will have public access.

Acknowledgements

We are grateful to the participants for their invaluable contribution to the study. This work was supported by funding from xx and Sanofi. Funding sources had no role in the analysis, interpretation of results or preparation of the manuscript.

Author contributions

VKK, HLW and PLD contributed to the study design. PK supervised the data generation. CM and PN contributed to the data generation. HK, IV, DU and NAP contributed to RNA-seq data quality control. TR, HK and DF contributed to data analysis. TR, HK and PLD contributed to the analysis plan. AS performed the dexamethasone experiment and analyzed its data. HT contributed to the dexamethasone experiment. VZ supervised the generation and analysis of macrophage RNA-seq data. JF contributed to the macrophage RNA-seq replication study. CLIMB study data acquisition was supervised by PLD and HLW. PhenoGenetic study data acquisition was supervised by PLD. TR and PLD contributed to the interpretation of results and drafted the manuscript. All authors read and revised the manuscript. PLD supervised the entire study.

Competing interests

Authors declare no competing interests in relation to the study.

References

1. Lassmann, H. Pathogenic mechanisms associated with different clinical courses of multiple sclerosis. *Front. Immunol.* **10**, (2019).
2. International Multiple Sclerosis Genetics Consortium. Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science* **365**, eaav7188 (2019).
3. Reich, D. S., Lucchinetti, C. F. & Calabresi, P. A. Multiple Sclerosis. *N. Engl. J. Med.* **378**, 169–180 (2018).
4. Nickles, D. *et al.* Blood RNA profiling in a large cohort of multiple sclerosis patients and healthy controls. *Hum. Mol. Genet.* **22**, 4194–4205 (2013).
5. Gandhi, K. S. *et al.* The multiple sclerosis whole blood mRNA transcriptome and genetic associations indicate dysregulation of specific T cell pathways in pathogenesis. *Hum. Mol. Genet.* **19**, 2134–2143 (2010).
6. Ratzer, R. *et al.* Gene expression analysis of relapsing-remitting, primary progressive and secondary progressive multiple sclerosis. *Mult. Scler.* **19**, 1841–1848 (2013).
7. Brorson, I. S. *et al.* No differential gene expression for CD4 + T cells of MS patients and healthy controls. *Mult. Scler. J. - Exp. Transl. Clin.* **5**, 205521731985690 (2019).
8. Kim, K. *et al.* Cell type-specific transcriptomics identifies neddylation as a novel therapeutic target in multiple sclerosis. *Brain* **144**, 450–461 (2021).
9. Ingelfinger, F. *et al.* Twin study reveals non-heritable immune perturbations in multiple sclerosis. *Nature* **603**, 152–158 (2022).
10. Gresle, M. M. *et al.* Multiple sclerosis risk variants regulate gene expression in innate and adaptive immune cells. *Life Sci. alliance* **3**, (2020).
11. Gauthier, S. A., Glanz, B. I., Mandel, M. & Weiner, H. L. A model for the comprehensive investigation of a chronic autoimmune disease: the multiple sclerosis CLIMB study. *Autoimmun. Rev.* **5**, 532–6 (2006).
12. Ottoboni, L. *et al.* An RNA Profile Identifies Two Subsets of Multiple Sclerosis Patients Differing in Disease Activity. *Sci. Transl. Med.* **4**, 153ra131-153ra131 (2012).
13. De Jager, P. L. *et al.* ImmVar project: Insights and design considerations for future studies of "healthy" immune variation. *Semin. Immunol.* **27**, 51–7 (2015).
14. Raj, T. *et al.* Polarization of the effects of autoimmune and neurodegenerative risk alleles in leukocytes. *Science* **344**, 519–23 (2014).
15. Trickett, A. & Kwan, Y. L. T cell stimulation and expansion using anti-CD3/CD28 beads. *J. Immunol. Methods* **275**, 251–255 (2003).
16. Dendrou, C. A., Fugger, L. & Friese, M. A. Immunopathology of multiple sclerosis. *Nat. Rev. Immunol. 2015 159* **15**, 545–558 (2015).
17. Souren, N. Y. *et al.* DNA methylation signatures of monozygotic twins clinically discordant for multiple sclerosis. *Nat. Commun.* **10**, (2019).
18. Fan, H. *et al.* Gender differences of B cell signature related to estrogen-induced IFI44L/BAFF in systemic lupus erythematosus. *Immunol. Lett.* **181**, 71–78 (2017).
19. Jia, X. *et al.* Integrated analysis of different microarray studies to identify candidate genes in type 1 diabetes. *J. Diabetes* **9**, 149–157 (2017).
20. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **25**, 25–29 (2000).
21. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set

collection. *Cell Syst.* **1**, 417–425 (2015).

22. Wu, T. & Mohan, C. The AKT axis as a therapeutic target in autoimmune diseases. *Endocr. Metab. Immune Disord. Drug Targets* **9**, 145–150 (2009).

23. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).

24. La Rocca, C. *et al.* Immunometabolic profiling of T cells from patients with relapsing-remitting multiple sclerosis reveals an impairment in glycolysis and mitochondrial respiration. *Metabolism.* **77**, 39–46 (2017).

25. Fransson, J. *et al.* Dysregulated functional and metabolic response in multiple sclerosis patient macrophages correlate with a more inflammatory state, reminiscent of trained immunity. *bioRxiv* 2021.01.13.426327 (2021) doi:10.1101/2021.01.13.426327.

26. Matarese, G., Carrieri, P. B., Montella, S., De Rosa, V. & La Cava, A. Leptin as a metabolic link to multiple sclerosis. *Nat. Rev. Neurol.* **6**, 455–461 (2010).

27. Subramanian, A. *et al.* A Next Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. *Cell* **171**, 1437-1452.e17 (2017).

28. Tissing, W. J. E. *et al.* Genomewide identification of prednisolone-responsive genes in acute lymphoblastic leukemia cells. *Blood* **109**, 3929–3935 (2007).

29. Hu, Y. *et al.* Development of a Molecular Signature to Monitor Pharmacodynamic Responses Mediated by In Vivo Administration of Glucocorticoids. *Arthritis Rheumatol. (Hoboken, N.J.)* **70**, 1331–1342 (2018).

30. Paragliola, R. M., Papi, G., Pontecorvi, A. & Corsello, S. M. Treatment with Synthetic Glucocorticoids and the Hypothalamus-Pituitary-Adrenal Axis. *Int. J. Mol. Sci.* **18**, (2017).

31. Bali, U., Phillips, T., Hunt, H. & Unitt, J. FKBP5 mRNA Expression Is a Biomarker for GR Antagonism. *J. Clin. Endocrinol. Metab.* **101**, 4305–4312 (2016).

32. Ysrraelit, M. C., Gaitán, M. I., Lopez, A. S. & Correale, J. Impaired hypothalamic-pituitary-adrenal axis activity in patients with multiple sclerosis. *Neurology* **71**, 1948–1954 (2008).

33. Van Winsen, L. M. L. *et al.* Sensitivity to glucocorticoids is decreased in relapsing remitting multiple sclerosis. *J. Clin. Endocrinol. Metab.* **90**, 734–740 (2005).

34. Wüst, S. *et al.* Peripheral T cells are the therapeutic targets of glucocorticoids in experimental autoimmune encephalomyelitis. *J. Immunol.* **180**, 8434–8443 (2008).

35. Alcina, A. *et al.* Multiple sclerosis risk variant HLA-DRB1*1501 associates with high expression of DRB1 gene in different human populations. *PLoS One* **7**, (2012).

36. Paulos, C. M. *et al.* The inducible costimulator (ICOS) is critical for the development of human T(H)17 cells. *Sci. Transl. Med.* **2**, (2010).

37. DeRijk, R. H., Eskandari, F. & Sternberg, E. M. Corticosteroid resistance in a subpopulation of multiple sclerosis patients as measured by ex vivo dexamethasone inhibition of LPS induced IL-6 production. *J. Neuroimmunol.* **151**, 180–188 (2004).

38. van Winsen, L. M. L. *et al.* Glucocorticoid receptor gene polymorphisms associated with more aggressive disease phenotype in MS. *J. Neuroimmunol.* **186**, 150–155 (2007).

39. Gold, S. M. *et al.* Dynamic development of glucocorticoid resistance during autoimmune neuroinflammation. *J. Clin. Endocrinol. Metab.* **97**, (2012).

40. Schumann, E. M. *et al.* Activity of the hypothalamic-pituitary-adrenal axis in multiple sclerosis: correlations with gadolinium-enhancing lesions and ventricular volume. *Ann. Neurol.* **51**, 763–767 (2002).

41. Novakova, L. *et al.* Monitoring disease activity in multiple sclerosis using serum neurofilament light protein. *Neurology* **89**, 2230–2237 (2017).

42. Rosso, M. *et al.* Temporal association of sNfL and gad-enhancing lesions in multiple sclerosis. *Ann. Clin. Transl. Neurol.* **7**, 945–955 (2020).

43. Yasir, M., Goyal, A. & Sonthalia, S. Corticosteroid Adverse Effects. *StatPearls* (2021).

44. Sundahl, N., Bridelance, J., Libert, C., De Bosscher, K. & Beck, I. M. Selective glucocorticoid receptor modulation: New directions with non-steroidal scaffolds. *Pharmacol. Ther.* **152**, 28–41 (2015).

45. GTEx Consortium, K. G. *et al.* Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* **348**, 648–60 (2015).

46. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

47. DeLuca, D. S. *et al.* RNA-SeQC: RNA-seq metrics for quality control and process optimization. *Bioinformatics* **28**, 1530–2 (2012).

48. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

49. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).

50. Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat. Genet.* **48**, 1284–1287 (2016).

51. Jia, X. *et al.* Imputing Amino Acid Polymorphisms in Human Leukocyte Antigens. *PLoS One* **8**, e64683 (2013).

52. Roostaei, T. *et al.* Proximal and distal effects of genetic susceptibility to multiple sclerosis on the T cell epigenome. *Nat. Commun. 2021 121* **12**, 1–12 (2021).

53. Wu, Y. *et al.* Automated segmentation of multiple sclerosis lesion subtypes with multichannel MRI. *Neuroimage* **32**, 1205–1215 (2006).

54. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

55. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

56. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

Figure legends

Figure 1. Summary of the generated data and the analyses performed in the study. (A) 1176 transcriptome data were generated from 227 individuals and 6 cell-type-states. (B) Cell-type-state-specific RNA-seq data were analyzed to identify MS-associated genes, pathways, and co-expressed gene modules. Replication, computational drug screen, and validation studies were also performed.

Figure 2. Results from transcriptome-wide association studies. (A) Number of significant genes (adjusted p <0.05) in transcriptome-wide association studies comparing each pair of the 3 participant groups (untreated MS, GA-treated MS, and healthy individuals) in each of the 6 studied cell-type-states. (B) Correlations between t-statistics from the transcriptome-wide association studies comparing the 2 MS groups with healthy individuals. Significant genes in any of the associations are shown in red. (C) Number of significant genes associated with MS in each cell-type-state are shown in black bars. Significance of the associations between each of these genes and MS in comparisons performed between each MS group and healthy participants, and also in comparisons between MS and healthy PBMC samples are shown in colors: adjusted p <0.05 with consistent direction of effect is shown in red, nominal p <0.05 with consistent direction of effect is shown in orange, and others are shown in gray. (D) Number of shared and distinct genes associated with MS in different cell types. (E) Expression of *ZBTB16* in each of the 6 cell-type-states, in addition to PBMC. (F) Pathways enriched in MS-associated genes in T cells and monocytes. Shared pathways are shown in blue. UNT-MS: untreated MS, GA-MS: GA-treated MS, HD: healthy donor.

Figure 3. Results from weighted gene co-expression network analysis. (A) Co-expression modules associated with MS and their hub genes. Horizontal black line represents the significance level (adjusted p =0.05). (B) Correlations between MS-associated module eigengenes in MS and healthy participants, separately. The two highly correlated modules in both MS and healthy individuals are highlighted using the dashed black square. (C) Associations between the genes in each of the 6 MS-associated modules and MS are visualized using volcano plots. Red dots represent genes belonging to each module. Horizontal dashed line represents the significance level (adjusted p =0.05). Differentially expressed genes from each module are labeled. Green circles represent genes that are also associated with genetic susceptibility to MS (T$_H$ naïve TNFSF8 module: *GGNBP2*, *LIMK2*, *NDFIP1*, *PIK3IP1*, *ZBTB38*; T$_H$ memory FKBP5 module: *ETS1*, *NR1D1*; monocyte GABARAPL1 module: *CXCR4*, *MPV17L2*; stimulated monocyte TMEM179B module: *ECD*; stimulated monocyte DBNL module: *ANXA7*, *RNFT1*). Enriched MsigDB hallmark gene sets are also shown for each module. (D) Replication of the association between T cell TNFSF8/FKBP5 modules and MS using independent PBMC samples. (E) Replication of the association between stimulated monocyte DBNL module and MS using independent data from monocyte-derived macrophages.

Figure 4. Relation between MS-associated genes/modules and glucocorticoid response genes. (A) Results from querying Connectivity Map database: top compound classes predicted to reverse changes in each MS-associated module. Glucocorticoid receptor agonists (highlighted in blue) are the only therapeutic class currently in clinical use. (B) Enrichment of the MS-associated T cell TNFSF8/FKBP5 modules for glucocorticoid response gene sets. Volcano plots

show association between the genes in each module and MS diagnosis. Horizontal dashed line represents the significance level (adjusted p =0.05). Orange and blue circles represent glucocorticoid response genes in leukemic cells and healthy PBMC, respectively. Red circles represent genes shared between the two glucocorticoid response gene sets. (C) Increase in the expression of *TNFSF8* and *FKBP5* in MS PBMC in response to *in vitro* treatment with dexamethasone. ** p <0.01, *** p <0.001, **** p <0.0001. VC= vehicle control (DMSO, 0 μM dexamethasone). (D) Genes with opposite direction of effect in association with MS (adjusted p <0.05) and in response to glucocorticoids. Black font represents genes belonging to the identified MS-associated modules (T_H naïve TNFSF8, T_H memory FKBP5, and monocyte GABARAPL1 modules, respectively).

Supplementary Figure 1. Correlations between t-statistics from the transcriptome-wide association studies comparing the 2 MS groups with healthy individuals.

Supplementary Figure 2. Correlations between t-statistics from the transcriptome-wide association studies comparing MS and healthy individuals in T cells and monocytes vs. PBMC. Orange dots: MS-associated genes in T cells or monocytes (adjusted p <0.05). Blue dots: MS-associated genes in PBMC (adjusted p <0.05). Black dots: MS-associated genes shared between the two comparisons. Horizontal and vertical dashed lines: nominal significance level (p =0.05). Spearman's ρ from left to right: 0.17, 0.18, and 0.26, respectively.

Supplementary Figure 3. No difference in the T cell TNFSF8/FKBP5 module eigengenes between MS participants who had a relapse in the last 3-12 months vs. in >12 months prior to blood sampling.

Supplementary Figure 4. Example changes in EDSS, BPF, and log-transformed lesion volume over the follow up period are visualized for three of the MS participants. Circles represent individual measurements. MRI measurements shown in gray are from 1.5T MR scans. X-axis units are in days. Purple lines represents the fitted models. Vertical blue lines represent the time of blood sampling. Vertical blue lines represent clinical relapses.

Tables

Table 1. Demographic characteristics of the participants.

Table 2. Genes associated with MS in both T cells and monocytes.

Supplementary Table 1. Genes differentially expressed between untreated and GA-treated MS.

Supplementary Table 2. Genes differentially expressed between MS and healthy participants in at least one of the six cell-type-states.

Supplementary Table 3. Enrichment of MS-associated genes in T cells and monocytes for MsigDB hallmark gene sets.

Supplementary Table 4. Associations between module eigengenes and outcome variables.

Supplementary Table 5. Glucocorticoid response genes.

Supplementary Table 6. Genes associated with MS outcome variables in each of the six cell-type-states.

Figure 1

A

Primary cell types

In vitro stimulation

T_H naïve

T_H naïve stim.
[anti-CD3/CD28]

T_H memory

T_H memory stim.
[anti-CD3/CD28]

Monocyte

Monocyte stim.
[TNF-α]

PBMC
181 MS
46 HC

**RNA-seq of
6 cell-type-states
from each individual**

B

1. TWAS

2. WGCNA

3. Replication

4. CMap

5. Validation

1. transcriptome-wide
association with MS
2. pathway enrichment

1. identifying modules
(co-expressed genes)
2. association with MS

1. PBMC
2. monocyte-derived
macrophages

screening compounds
for reversing MS-
associated modules

in vitro assessment of
compound effects on
gene expression in MS

Figure 2

Figure 3

A

B

C

T_H naïve MYL6 module

MsigDB Hallmark:
oxidative phosphorylation
adipogenesis

T_H naïve TNFSF8 module

MsigDB Hallmark:
-

T_H memory FKBP5 module

MsigDB Hallmark:
-

Monocyte GABARAPL1 module

MsigDB Hallmark:
TNF-α signaling via NF-κB
hypoxia
inflammatory response

Monocyte stim. TMEM179B module

MsigDB Hallmark:
oxidative phosphorylation
adipogenesis

Monocyte stim. DBNL module

MsigDB Hallmark:
-

D

T_H naïve TNFSF8 module

Replication: PBMC

T_H memory FKBP5 module

Replication: PBMC

E

Monocyte stim. DBNL module

Replication: macrophages

Figure 4

A

Connectivity Map: predicted therapeutic Compound Classes

T_H naïve MYL6 module:
Proteasome inhibitor
MTOR inhibitor
Protein synthesis inhibitor
PI3K inhibitor
ATPase inhibitor
Protein tyrosine kinase inhibitor
Beta-amyloid inhibitor
HIF modulator

T_H naïve TNFSF8 module:
Glucocorticoid receptor agonist

T_H memory FKBP5 module:
Glucocorticoid receptor agonist

Monocyte GABARAPL1 module:
Proteasome inhibitor

Monocyte stim. TMEM179B module:
Proteasome inhibitor
Ubiquitin-specific proteasome inhibitor
ATPase inhibitor
MTOR inhibitor
Protein synthesis inhibitor
PI3K inhibitor
HIF activator
HMGCR inhibitor
HIF modulator
PKC activator
Tubulin inhibitor

Monocyte stim. DBNL module:
ATPase inhibitor
CDK inhibitor
Telomerase inhibitor
Tubulin inhibitor
PKA inhibitor
Cell cycle inhibitor
PI3K inhibitor
Retinoid receptor agonist

B

T_H naïve TNFSF8 module

T_H memory FKBP5 module

C

TNFSF8

FKBP5

D

T_H naïve

↓APBA2, ↓CCND3, ↓FKBP5, ↓KLF9, ↓LIMK2, ↓NDFIP1, ↓SESN1, ↓SLA, ↓TNFSF8, ↓TSC22D3, ↓TXNIP, ↓ZBTB16

T_H memory

↓CCND3, ↓FKBP5, ↓SLA, ↓TNFSF8, ↓TXNIP, ↓ZBTB16

Monocyte

↓CCND3, ↓CEBPD, ↓CRISPLD2, ↓CSNK1D, ↑CX3CR1, ↓FKBP5, ↓FLT3, ↑IL1RN, ↓IRAK3, ↓IRS2, ↓KLF9, ↓MAN2A2, ↑MYOF, ↓PER1, ↓PHC2, ↓SLA, ↓TBC1D8, ↓TSC22D3, ↓ZBTB16, ↓ZFP36L2

↓ downregulated in MS, upregulated in response to glucocorticoids
↑ upregulated in MS, downregulated in response to glucocorticoids

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Days from blood sampling