Causal role of the dorsolateral prefrontal cortex in modulating the balance between Pavlovian

and instrumental systems in the punishment domain

Hyeonjin Kim[1]*, Jihyun K. Hur[2], Mina Kwon[1], Soyeon Kim[1], Yoonseo Zoh[2], Woo-Young

Ahn[1,3]

[1] Department of Psychology, Seoul National University, Seoul, Korea

[2] Department of Psychology, Yale University, New Haven, Connecticut, United States of

America

[3] Department of Brain and Cognitive Sciences, Seoul National University, Seoul, Korea

* Corresponding author

E-mail: wahn55@snu.ac.kr

Abstract

Previous literature suggests that a balance between Pavlovian and instrumental decision-making systems is critical for optimal decision-making. Pavlovian bias (i.e., approach toward reward-predictive stimuli and avoid punishment-predictive stimuli) often contrasts with the instrumental response. Although recent neuroimaging studies have identified brain regions that may be related to Pavlovian bias, including the dorsolateral prefrontal cortex (dlPFC), it is unclear whether a causal relationship exists. Therefore, we investigated whether upregulation of the dlPFC using transcranial current direct stimulation (tDCS) would reduce Pavlovian bias. In this double-blind study, participants were assigned to the anodal or the sham group; they received stimulation over the right dlPFC for 3 successive days. On the last day, participants performed a reinforcement learning task known as the orthogonalized go/no-go task; this was used to assess each participant's degree of Pavlovian bias in reward and punishment domains. We used computational modeling and hierarchical Bayesian analysis to estimate model parameters reflecting latent cognitive processes, including Pavlovian bias, go bias, and choice randomness. Several computational models were compared; the model with separate Pavlovian bias parameters for reward and punishment domains demonstrated the best model fit. When using a behavioral index of Pavlovian bias, the anodal group showed significantly lower Pavlovian bias in the punishment domain, but not in the reward domain, compared with the sham group. In addition, computational modeling showed that Pavlovian bias parameter in the punishment domain was lower in the anodal group than in the sham group, which is consistent with the behavioral findings. The anodal group also showed a lower go bias and choice randomness, compared with the sham group. These findings suggest that anodal tDCS may lead to behavioral suppression or change in Pavlovian bias in the punishment domain, which will help to improve comprehension of the causal neural mechanism.

Author summary

A decision-making bias guided by the Pavlovian system (i.e., approach reward and avoid punishment) is often useful and predominant across species but it is also related to several psychiatric conditions. The dorsolateral prefrontal cortex (dlPFC) is known to be related to such "Pavlovian bias" but it is unclear whether a causal relationship exists between them. Here, we evaluated whether decision-making biases including Pavlovian bias could be modulated by exogenous brain stimulation, transcranial current direct stimulation, over the right dlPFC for 3 successive days. A combination of behavioral analysis and computational modeling revealed that the anodal group had lower Pavlovian bias in the punishment domain compared with the sham group. In addition, the anodal group showed lower go bias and choice randomness than the sham group, which can also hamper instrumental learning. These findings suggest a causal role for the dlPFC in modulating the balance between the Pavlovian and instrumental decision-making systems.

Introduction

Decision-making is governed by multiple systems, including the fundamental Pavlovian and instrumental systems. The Pavlovian system involves a pre-preprogrammed behavioral tendency known as Pavlovian bias (i.e., approaching reward-predictive stimuli and avoiding punishment-predictive stimuli) [1]. In contrast, the instrumental system involves learning the optimal response to each stimulus by evaluating its outcomes without prior preparation. Although the Pavlovian bias has several benefits, it may hamper goal-directed behavior. For example, animals (e.g., pigeons) with strong Pavlovian bias fail to learn to withhold pecking in response to stimuli predictive of food, even when they can receive food

only by withholding pecking [2,3]. Humans are also affected by Pavlovian bias in various

decision-making situations, such as dieting [4,5] or substance abuse [6]. Thus, there is a need

to investigate methods to effectively overcome such bias.

The neural mechanisms that underlie Pavlovian bias are not fully understood, but

some previous research has suggested that the prefrontal cortex plays a pivotal role in

overcoming Pavlovian bias [7–9]. A functional magnetic resonance imaging (fMRI) study of

participants who successfully employed the instrumental system during conflict with the

Pavlovian system found that such individuals showed hyperactivation of the bilateral inferior

frontal gyri while anticipating inhibition [9]. In addition, an electroencephalography study

showed that the activation of the anterior cingulate cortex, as measured by the midfrontal

theta power of the electroencephalogram signal, was associated with overcoming Pavlovian

bias [7]. However, these studies failed to provide conclusive evidence for a causal neural

mechanism, and the brain regions that control Pavlovian bias remained unknown.

We speculated that the dorsolateral prefrontal cortex (dlPFC) might be a key region

involved in controlling Pavlovian bias. The dlPFC has been implicated in higher-level

cognitive control and goal-directed actions [4,5,10–17]. For example, dieters showed

hyperactivation of the dlPFC when they successfully selected healthy food over tasty food [4].

In addition, the dlPFC was important in individuals who valued stimuli in a context-

dependent manner and performed goal-directed behavior to maximize reward [10]. Although

a previous fMRI experiment studying the neural correlates of Pavlovian bias did not identify

the dlPFC as a candidate region [9], the negative results are related to the imaging strategy

used in the study, rather than the lack of a relationship. The imaging was focused on

subcortical structures; the dlPFC regions were not assessed.

In the present study, we evaluated the presence of a causal relationship between the

dlPFC and Pavlovian bias using non-invasive brain stimulation (i.e., transcranial direct

current stimulation [tDCS]). Using tDCS was based on several previous studies of modulating decision-making biases. For example, the competition between the model-based and the model-free systems [18], as well as affective bias of instrumental action [19], were modulated by tDCS targeting the prefrontal cortex.

Overall, we investigated whether anodal tDCS on dlPFC would suppress the Pavlovian bias (sham-controlled); we sought to identify the causal neural mechanism underlying such bias. We applied anodal tDCS over the right dlPFC [20–22] for 3 consecutive days [23–25]. On the third day, we administered a reinforcement learning task known as the orthogonalized go/no-go task, which measured the degree of Pavlovian bias [9]. The task had four conditions; two were Pavlovian-congruent, where go was the action required to win the reward and no-go was the action required to avoid punishment; the two remaining conditions were Pavlovian-incongruent, where go was the action required to avoid punishment and no-go was the action required to win the reward. Participants were required to learn the correct action for each condition to maximize the reward and minimize punishment. We compared the degree of Pavlovian bias across tDCS groups using the difference in behavioral accuracy between Pavlovian-congruent and Pavlovian-incongruent conditions. We also used a model parameter (i.e., Pavlovian bias parameter) estimated by computational modeling and hierarchical Bayesian analysis (HBA) as another index of Pavlovian bias. Under the punishment domain, we found significantly lower Pavlovian bias in the anodal tDCS group than in the sham group.

Materials and methods

Participants

We recruited 39 participants from Seoul National University in Seoul, Korea, using

online and offline advertisements. The experimental protocol was approved by the Seoul

National University Research Ethics Committee and all participants provided informed

consent before participation. Participants were excluded if they were unwilling to participate

in the study, or were not fluent in Korean; they were also excluded if they reported impaired

color discrimination, psychiatric medication use, neurological or psychiatric illness, or any

health conditions that would make them unsuitable for the experiments. In addition,

participants were excluded if they had low-quality data such as sleep during the experiment

or results that indicated an inability to understand the task. Finally, we eliminated participants

with a go-to-win accuracy of < 0.1 because learning failure in the easiest go-to-win condition

indicated a lack of understanding or concentration. In total, data from 17 and 14 participants

in the anodal and sham sessions, respectively, were analyzed (see below for more

information).

Experimental protocol

First, we collected data regarding the participants' basic demographic information

(age and sex) and psychological characteristics. We administered the Structured Clinical

Interview for DSM-5 to detect mental illnesses. In addition, we evaluated the psychological

characteristics of obsession-compulsion (Yale-Brown Obsessive Compulsive Scale),

depression (Beck's Depression Inventory), anxiety (State-Trait Anxiety Inventory), and

impulsivity (Barratt Impulsiveness Scale version 11). The participants visited the laboratory

for 3 consecutive days and repeated the visits to counterbalance the tDCS polarity (six total

sessions). For the first 2 days, participants received tDCS for 20 min; on the third day,

participants performed an orthogonalized go/no-go task after they had received tDCS

stimulation for 20 min. The daily visiting time was matched on a within-participant basis to

remove the confounding effect of circadian rhythm [26,27]. Participants were randomly

assigned to receive anodal or sham stimulation on the first or second 3 days of visits. The first

and second sets of visits were separated by a mean of 24 days. We found significantly better

performance in the second task (see S1 Fig in the supplementary material for more

information), suggesting a practice effect. Therefore, we analyzed behavior data only from

the first task to avoid any potential confounding effects.

154 tDCS stimulation

During each session, tDCS was applied for 20 min using circular sponge electrodes

(size = 25 cm^2) and the Starstim system (Neuroelectrics, Barcelona, Spain). The target

electrode was positioned on the right dlPFC (i.e., F4 according to the 10-10 International 10-

20 electroencephalogram electrode system); the return electrode was positioned on the left

cheek (Fig 1). The stimulation protocol was based on previous studies that used tDCS

targeting dlPFC [20]. The left cheek was selected as the return position to avoid confounding

cortical activation [28–31]. The stimulation included 30 s of ramp-up and ramp-down at the

beginning and end of the stimulation, respectively. During the anodal session, anodal

stimulation to F4 was performed for 19 min between ramp-up and ramp-down stimulations;

however, in the sham session, participants were not stimulated between the ramp-up and

ramp-down stimulations. During the stimulation, participants were instructed to sit with their

gaze fixed on a crosshair on the computer monitor. The double-blind mode in the Starstim

software was used to ensure that all experimenters and participants remained unaware of the

order of polarity. The software blinds the type of current stimulation using a 4-digit password

lock set by a third-party administrator.

We employed some strategies to reduce the potential limitations of tDCS in the

current stimulation protocol. First, the electrode placement and size can affect the spatial

distribution of stimulation [32]. Therefore, we placed a return electrode in an extracephalic

area (i.e., left chick) to minimize the stimulation of other cortical areas and the shunting

effect caused by a short inter-electrode distance [33]. In addition, the effects of tDCS can be

confounded by biological and lifestyle factors [33]. We mitigated such factors by stimulating

participants over 3 consecutive days before the task to produce cumulative and larger effects.

The participants visited the laboratory at the same time (variation of < 3 h) to reduce the

effects of circadian rhythm. Finally, to reduce confounding factors related to the experimental

design, we used a sham-controlled double-blind protocol.

Fig 1. Montage of tDCS. A sponge was placed over the right dlPFC (F4) to stimulate the

brain using weak electric current (2 mA) for 20 min. Another sponge was placed on the left

cheek. This figure was adapted from the protocol summary panel in Starstim software NIC

(copyright notice © Neuroelectrics SLU).

Experimental task

We used the orthogonalized go/no-go task reported by Guitart-Masip et al. (2013)

(Fig 2). At the beginning of each trial, a 1000-ms cue (fractal image) was presented to

indicate one of four conditions; go-to-win reward, go-to-avoid punishment, no-go-to-win

reward, and no-go-to-avoid punishment. After a variable interval of 250–2000 ms, a target

circle appeared for a maximum of 1500 ms, after which the participants responded with go or

no-go within 1000 ms. After 1000 ms, participants received feedback according to their

response and cue condition. The feedback included virtual monetary gain as a reward, a

yellow bar as a neutral outcome, and an electric shock as punishment instead of monetary

loss to maximize the effect of punishment [34]. The optimal response led to a beneficial

outcome for each condition, with a probability of 0.7. Therefore, participants learned the

optimal response to each cue from trial and error. The task included 180 trials, with 45 trials

for each condition.

Fig 2. Orthogonalized go/no-go task. Four types of stimuli were presented. Two stimuli

were Pavlovian-congruent: go action to win reward and no-go action to avoid punishment.

The remaining two stimuli were Pavlovian-incongruent: go action to avoid punishment and

no-go action to win reward. Participants were instructed to maximize reward and minimize

punishment by learning the correct action for each stimulus. Participants were asked to select

an action when a target was presented. The reward was a picture of money, 1000 won

(approximately US$ 1), whereas the punishment was an electric shock to the wrist.

RT: response time, ITI: intertrial interval

Although money is secondary feedback and shock is primary feedback, we decided to

use monetary gain as the reward and electric shock as the punishment based on previous

studies [34–37]. The electric shock was applied to each participant's left wrist. The intensity

of the electric shock (2–12.4mA) was adjusted to cause a "moderately unpleasant" sensation

(5 points on an 11-point Likert scale [i.e., 0 = not at all unpleasant, 10 = very un- pleasant]).

(see supplementary material for more information).

Behavioral data analysis

Behavioral data were analyzed using R [38]. Response accuracy was calculated as the

proportion of correct choices. The difference in accuracy between Pavlovian-congruent and

Pavlovian-incongruent conditions was evaluated using Student's t-test. The behavioral

Pavlovian bias index was evaluated as the difference between the accuracy of Pavlovian-

congruent and Pavlovian-incongruent conditions; it was calculated individually for each

domain. For example, Pavlovian bias index in the punishment domain was calculated by

subtracting the accuracy of the go-to-avoid condition from the accuracy of the no-go-to-avoid

condition.

$$Pavlovian\ bias = (go\_to\_win + no\_go\_to\_avoid) - (no\_go\_to\_win + go\_to\_avoid)$$

$$Pavlovian\ bias\ (reward) = (go\_to\_win) - (no\_go\_to\_win)$$

$$Pavlovian\ bias\ (punishment) = (no\_go\_to\_avoid) - (go\_to\_avoid)$$

Computational modeling

We tested three models. A previous study suggested that Model 1 had the best fit and

consisted of five parameters (Model RW + noise + bias + Pav; Guitart-Masip et al., 2012).

Model 1 calculates the probability of performing (or withholding it) an action in response to

the stimulus in each trial, based on action weights. If a participant successfully learned the

action-reward contingency, the probability of performing a correct action was higher. It is

calculated as follows:

$\quad p(a_1 \mid s) = \left\{ \dfrac{exp(W_t(a_1, s))}{exp(W_t(a_1, s)) + exp(W_t(a_2, s))} \right\} (1 - \xi) + \dfrac{\xi}{2} \ ... \ (1)$

In particular, the go action probability was larger if the W value for go (a_1) was

greater using squashed softmax and for no-go (a_2), vice versa. Here, t is the trial number ($1 \leq$

$t \leq 180$) and s is the stimulus (s $\in$ {1, 2, 3, 4}). The four stimuli indicate four conditions,

respectively: go-to-win reward, go-to-avoid punishment, no-go-to-win reward, and no-go-to-

avoid punishment. In addition, a is the action (a $\in$ {0, 1}), where 1 is go and 0 is no-go. ξ is

the irreducible noise ($0 \leq \xi \leq 1$), where a value closer to 1 indicates random choice less

considering the W value. W(a, s) is the action weight, which is defined as follows:

$$W_t(a, s) = \begin{cases} Q_t(a, s) + b + \pi V_t(s) & if\ a = go \\ Q_t(a, s) & otherwise \end{cases} \quad \text{... (2)}$$

Q(a, s) and V(s) are updated by each trial according to the equations below:

$Q_t(a_t, s_t) = Q_{t-1}(a_t, s_t) + \varepsilon(\rho r_t - Q_{t-1}(a_t, s_t))$ ... (3)

$V_t(s_t) = V_{t-1}(s_t) + \varepsilon(\rho r_t - V_{t-1}(s_t))$ ... (4)

In equation (3), r is the feedback ($r \in \{-1, 0, 1\}$), where 1 is the reward, 0 is neutral,

and -1 is punishment. ε is the learning rate ($0 \leq \varepsilon \leq 1$); If ε is closer to 1, it is more likely to

reflect the previous feedbacks to update Q values. Furthermore, ρ is outcome sensitivity ($0 \leq$

ρ). A larger ρ indicates the participant subjectively exaggerates the outcome value. Using this

process, the Q value converges to the high-probability outcome for each stimulus when the

correct action for the stimulus is accumulated.

In equation (4), the V value is updated in a manner similar to the Q value, but it

converges to the high-probability feedback for each stimulus, regardless of the performed

actions. In equation (2), for the updated Q values when the action was go, the go bias

parameter b and V value multiplied by the Pavlovian bias parameter π ($0 \leq \pi$) were added to

the Q values; they consisted of the W values. A large go bias was correlated with large W(go,

s). When the V value converged to reward, and the action was go, the large Pavlovian bias

parameter was correlated with generally large W(go, reward) and generally small W(no-go,

reward). When the V value converged to punishment and the action was go, the large

Pavlovian bias parameter was correlated with generally small W(go, punishment) and

generally large W(no-go, punishment). This suggests that a large Pavlovian bias parameter
was correlated with greater predisposition to Pavlovian-congruent choices.

Model 2 shares almost all equations and updating rules with Model 1, although it has
distinct feedback sensitivity parameters for reward and punishment cues: ρ_{rew}, and ρ_{pun},
respectively. Therefore, Model 2 contains six parameters. Model 3 shares almost all
equations and updating rules with Model, although it has different Pavlovian bias parameters
for reward and punishment cues: π_{rew} and π_{pun}, respectively. Model 3 contains seven
parameters and was used to test the distinct effect found in behavioral data, where Pavlovian
bias was only significant in the punishment domain.

277 Model parameter estimation using HBA

The model parameters were estimated using HBA [39–41]. HBA has some
advantages over the traditional maximum likelihood estimation (MLE) method. First, HBA
provides estimated parameters as posterior distributions, rather than the point estimates
provided by MLE. The distributions provide additional information, particularly regarding
the uncertainty of estimated values. Second, the hierarchical structure of HBA allows stable
and reliable estimation of individual parameters. Individual-level MLE estimates are often
noisy and unreliable; group-level MLE estimates do not include information concerning
individual differences. In HBA, each individual estimate informs the group estimate
(hyperparameter), and the individual commonalities reflected in the hyperparameter inform
individual estimates. Therefore, individual estimates are more stable and reliable, even when
data are insufficient. Previous studies have found that parameters estimated by HBA are more
accurate than parameters estimated by MLE [42].

We separately fitted the models for anodal and sham groups to make stable and
reliable individual estimates that reflected similarities within each group. HBA was

conducted by hBayesDM (v. 1.1.1) [43] and R Stan (v. 2.21.0) [44]. Stan is a probabilistic

program used for Bayesian modeling; it provides inferences based on Markov chain Monte

Carlo (MCMC) algorithms, such as the Hamiltonian Monte Carlo, for sampling from high-

dimensional parameter spaces. Weakly informative priors were used to reduce their influence

on the posterior distributions [43]. In addition, non-centered parameterization (Matt trick)

was used to optimize the sampling process [45]. We used four independent chains and a

sample size of 4000, including 2000 burn-in samples per chain. The use of four independent

chains ensured that the estimated parameters were stable, despite variations in the starting

points [46]. We also confirmed the accuracy of parameter estimation by inspecting well-

mixed trace plots and the Rhat values (Rhat < 1.1).

Model comparison

We used LOOIC to compare the models [47]. The LOOIC value for each model was

calculated by estimating the out-of-sample prediction accuracy of the fitted models. This

method uses the log-likelihood from posterior simulations of the estimated parameters. We

used R package loo to identify the model with the lowest LOOIC value, which had the best fit

[47].

Group comparison of model parameters

For each group-level parameter, we subtracted the posterior distribution of the sham

group from the posterior distribution of the anodal group for analysis of group-level

differences. Group differences were considered credible when the 95% highest density

intervals of posterior difference distributions did not include the value 0 [48].

Results

Anodal and sham group characteristics

We analyzed data from 31 participants, including the basic demographic information (age and sex), psychiatric symptoms, and psychological characteristics (Table 1). There were no significant group-level differences in terms of demographic, psychiatric, and psychological variables between the sham and the anodal groups. We also measured the perceived side effects of tDCS; we found no differences between groups in terms of itching, skin irritation, skin pain, fatigue, mood disturbance, and visual distortion ($p > 0.05$ for all; see S1 Table in supplementary material). However, the intensity of perceived tingling was significantly higher in the anodal group than in the sham group ($t(91) = -2.12$; $p = 0.04$). In addition, the degrees of headache and difficulty in concentration were significantly higher in the sham group than in the anodal group (headache: $t(91) = 2.84$; $p = 0.006$, concentration: $t(91) = 2.15$; $p = 0.03$). The differences in perceived side effects did not affect the behavioral Pavlovian bias. However, there were significant differences in perceived duration and continuity of stimulation between the sham and anodal groups (S2 Table in supplementary material).

Table 1. Descriptive statistics.

	Sham (N = 14)	Anode (N = 17)	p value
Age	25.071 (3.731)	23.529 (3.484)	0.245
Sex; male	5 (35.7%)	8 (47.1%)	0.524
SCID[a]			
avoidant	2.357 (2.098)	2.294 (1.724)	0.927
dependent	1.286 (1.773)	1.412 (1.228)	0.817
obsessive-compulsive	2.714 (1.816)	3.353 (1.656)	0.315
passive-aggressive	1.000 (1.109)	1.353 (1.967)	0.555
depressive	1.714 (1.858)	2.059 (1.919)	0.618
paranoid	1.500 (1.871)	1.706 (1.929)	0.767

schizotypal	0.857 (1.027)	0.765 (1.480)	0.845
schizoid	0.929 (1.207)	1.412 (1.502)	0.339
histrionic	1.929 (1.269)	2.176 (1.590)	0.640
narcissistic	3.429 (2.503)	3.353 (2.783)	0.938
borderline	1.429 (1.742)	2.941 (3.750)	0.176
antisocial	0.857 (1.406)	0.412 (0.870)	0.289
Y-BOCS[b]			
obsessive	1.357 (2.530)	1.824 (3.206)	0.662
compulsive	4.143 (4.258)	3.688 (3.860)	0.761
BDI[c]			
	3.714 (2.920)	9.176 (12.259)	0.115
STAI-X[d]			
state	38.000 (9.397)	40.941 (9.523)	0.396
trait	37.071 (6.956)	40.176 (10.212)	0.342
BIS11[e]			
cognitive	16.429 (3.322)	16.412 (3.692)	0.990
motor	20.357 (3.934)	19.882 (5.171)	0.780
non-planning	24.571 (5.721)	24.176 (5.637)	0.848

Mean (standard deviation) for continuous variables and count (%) for categorical variables.

[a]SCID: Structured Clinical Interview for DSM-5

[b]Y-BOCS: Yale-Brown Obsessive Compulsive Scale

[c]BDI: Beck's Depression Inventory

[d]STAI-X: State-Trait Anxiety Inventory

[e]BIS 11: Barratt Impulsiveness Scale Version 11

Behavioral results

We used the difference in behavioral accuracy under Pavlovian-congruent and

Pavlovian-incongruent conditions to compare the degree of Pavlovian bias across tDCS

groups. In the punishment domain, the anodal group did not show any significant difference

in behavior under the two punishment conditions (Fig 3A; $t(32) = 0.18$; $p = 0.86$). In contrast,

the sham group exhibited significantly lower accuracy under the Pavlovian-incongruent

punishment condition (e.g., go-to-avoid) than under the Pavlovian-congruent condition (e.g.,

no-go-to-avoid) (Fig 3B; t(26) = -2.31; p = 0.03). Neither group exhibited a significant

difference in accuracy under the two reward conditions (e.g., go-to-win and no-go-to-win).

Consistent with these findings, the behavioral Pavlovian bias index in the punishment domain

was significantly lower in the anodal group than in the sham group (t(29) = -3.09; p = 0.004).

Fig 3. Pavlovian bias in the punishment domain decreased in the anodal session

(accuracy). In the sham group, we found a significant difference in behavioral accuracy

between the punishment conditions, which indicated the presence of Pavlovian bias,

particularly in the punishment domain. This difference was not observed in the anodal group.

The behavioral index of Pavlovian bias in the punishment domain also showed significantly

lower bias in the anodal group than in the sham group. Error bars indicate SEM. *p < 0.05

Computational modeling

We tested three computational models to explain the data. Model 1 was a

reinforcement learning model suggested by Guitart-Masip et al. (2012 which included five

parameters (ξ: irreducible noise; ε: learning rate; ρ: outcome sensitivity; b: go bias; π:

Pavlovian bias). Model 2 was a model with six parameters, including separate feedback

sensitivity parameters for reward and punishment cues (ρ_{rew} and ρ_{pun}), compared to Model

1. Model 3 further separated Pavlovian bias parameters for reward and punishment cues cues

(π_{rew} and π_{pun}) compared with Model 2. We compared the models using the leave-one-out

information criterion (LOOIC) values, which were calculated using leave-one-out cross-

validation [47] (Table 2). Data from the sham and anodal groups were fitted separately.

Model 3, which had separate Pavlovian bias parameters for reward and punishment domains

the best fit (see S2 and S3 Figs in the supplementary material to check one-step ahead

prediction results). Models 2 and 1 were the second and third best-fitting models, respectively.

Thus, we used estimated parameter values from Model 3 for the subsequent analyses.

Table 2. Model comparison (LOOIC).

	Parameters	sham	anode
Model 1	$\xi, \varepsilon, b, \pi, \rho$	1852.5	2168.7
Model 2	$\xi, \varepsilon, b, \pi, \rho_{rew}, \rho_{pun}$	1813.6	2129.7
Model 3	$\xi, \varepsilon, b, \pi_{rew}, \pi_{pun}, \rho_{rew}, \rho_{pun}$	**1769.7**	**2093.0**

375 Lower LOOIC values indicated better model performance.

Model parameters

We calculated the posterior distributions of all group-level parameters from Model 3;

we compared the results between the anodal and sham groups (Table 3, Fig 4). The anodal

group displayed credibly lower irreducible noise (ξ), compared with the sham group. Go bias

(b) was also credibly lower in the anodal group than in the sham group. Finally, the anodal

group had credibly lower Pavlovian bias in the punishment domain (π_{pun}), compared with

the sham group; this is consistent with the behavioral analysis findings that behavioral

Pavlovian bias in the punishment domain was significantly lower in the anodal group than in

the sham group. Increased involvement of the frontal-striatal network after anodal stimulation

might suppress the biases (e.g. Pavlovian bias in the punishment domain, go bias, and choice

randomness), thereby interrupting goal-directed behavior of the instrumental system.

Table 3. Posterior mean (95% HDI highest density interval) of group mean parameters in

anodal and sham groups

	Anode	Sham	Difference
ξ irreducible noise	0.027 [0.009, 0.060]	0.092 [0.056, 0.141]	-0.064 [-0.117, -0.017]
ε learning rate	0.430 [0.316, 0.554]	0.321 [0.224, 0.442]	0.107 [-0.052, 0.264]

b Go-bias	-0.173 [-0.689, 0.334]	1.27 [0.437, 2.46]	-1.45 [-2.72, -0.461]
π_{rew} Pavlovian bias (reward)	0.045 [-0.223, 0.322]	-0.141 [-0.428, 0.351]	0.183 [-0.364, 0.587]
π_{pun} Pavlovian bias (punishment)	-0.063 [-0.212, 0.091]	0.411 [-0.033, 0.855]	-0.473 [-0.939, -0.006]
ρ_{rew} reward sensitivity	12.4 [7.72, 21.8]	12.3 [7.63, 21.4]	0.146 [-9.91, 10.7]
ρ_{pun} punishment sensitivity	6.24 [4.69, 8.31]	8.83 [5.92, 14.0]	-2.58 [-8.02, 1.02]

Fig 4. Pavlovian bias parameter in the punishment domain and other parameters

decreased in the anodal session (modeling parameter). We found lower parameter values

in irreducible noise, go bias, and Pavlovian bias in the punishment domain in the anodal

session, compared with the sham session. The decrease in Pavlovian bias in the punishment

domain is consistent with the behavioral analysis. * 95% highest density interval of the

posterior difference did not include zero.

However, there were no credible differences between groups in terms of other

parameters, such as learning rate (ε), Pavlovian bias in reward domain (π_{rew}), reward

sensitivity (ρ_{rew}), and punishment sensitivity (ρ_{pun}).

Discussion

Our results suggest a causal role of the dlPFC in modulating Pavlovian bias in the

punishment domain. Moreover, we found that other decision-making tendencies (i.e., go bias

and irreducible noise) were also modulated.

We found that anodal stimulation of the dlPFC reduced Pavlovian bias, which might

be related to goal-directed control in the frontal-striatal circuit. The frontal-striatal circuit

connects the prefrontal cortex and striatum (including following key areas: ventral striatum

[nucleus accumbens], dorsal striatum [caudate and putamen], ventromedial prefrontal cortex,

dlPFC, and dorsal anterior cingulate cortex [dACC]) [17,49]. The dlPFC neurons that project

to the striatum may modulate the action-outcome contingency that is encoded and updated in

the striatum and ventromedial prefrontal cortex. Therefore, anodal stimulation over the right

dlPFC may facilitate high-level cognitive control [4,5,12–17] and enhance goal-directed

behavior by suppressing Pavlovian bias. Another possible mechanism is that anodal

stimulation of the dlPFC increases dopamine release in the striatum [50,51]; when the dlPFC

is stimulated, information about instrumental control is transmitted to the striatum, which

responds by increasing dopamine release and overcoming the Pavlovian bias. Furthermore,

the tDCS might lead to increased connectivity between the dlPFC and dACC. The dACC,

along with the dlPFC, plays a critical role in updating action values and modulating the

integration of subjective value and action-outcome contingency. Previous studies showed that

Pavlovian bias was suppressed by frontal midline theta power, an electroencephalography

correlate of dACC [7,8]. Therefore, it is plausible that the tDCS over the dlPFC facilitated the

dACC activation, which would reduce Pavlovian bias.

However, the current study only found suppression of Pavlovian bias in the

punishment domain, not the reward domain. A recent intracranial study also reported that the

electroencephalogram implanted in dlPFC appears to signal punishment learning rather than

reward during the instrumental learning task [52]. The anodal stimulation on dlPFC maybe

facilitated the value updating for the punishment domain and resulted in a significant change

in Pavlovian bias only in the punishment domain. On the other hands, the present findings

contribute to knowledge about aversion-related decision-making in the Pavlovian system

[53–55]. The underlying neural mechanisms of appetitive-related decision-making have been

widely investigated, but the mechanisms that underlie aversive-related decision-making have

received less attention [56]. A recent study found that aversive stimuli were associated with

active escape response or passive avoidance response [54]. The authors suggested that

serotonin might be involved in passive inhibitory responses [57–59], while dopamine might

be involved in active escape responses; this is similar to the active approach response toward

appetitive stimuli [53,60]. Therefore, the current results concerning suppression of Pavlovian

bias in the aversive domain, obtained by connecting behavioral activation and avoidance (go-

to-avoid), might reflect a similar neural process for active escape response. These results are

consistent with previous evidence that increased dopamine release after anodal tDCS of the

dlPFC might suppress Pavlovian bias [26,55]. Future tDCS studies should separate avoidance

and escape trials to further explore the mechanism that underlies suppression of Pavlovian

bias in the punishment domain.

We also observed decreases in go bias and choice randomness in the anodal group.

Because go bias and choice randomness interrupt the goal of maximizing benefit, a similar

mechanism for interrupting goal-directed behavior may exist, as previously discussed.

Increased involvement of the frontal-striatal network (dlPFC, striatum, and dACC) after

electrical stimulation of the dlPFC might lead to the suppression of go bias and choice

randomness. We presume that the instrumental system may gain preference under conflicting

conditions between the instrumental and Pavlovian systems.

Our result showed that decision-making biases were modulated by external

intervention, which may have clinical relevance, particularly for substance misuse and other

addictive behaviors. For example, increased Pavlovian bias has been linked to substance use

and gambling disorders [61,62], while increased go bias, which may reflect impaired

response inhibition, has also been associated with various addictive disorders [63,64]. Choice

randomness (e.g. decision-making noise or inverse temperature) was greater in patients with

cocaine abuse and gambling disorders than in healthy individuals [65,66]. Thus, the current

findings may aid in the development of treatments that can reduce the decision-making biases

implicated in the various psychiatric conditions. Because the current study only included

healthy participants, future studies should include individuals with psychiatric disorders.

One of the potential limitations of the current study is that the second session data

were affected by the task practice effect. Therefore, the overall accuracy of task performance

was significantly higher in the second session than in the first session. To control for the

practice effect, we only analyzed data from the first session and performed between-subject

analyses. The resulting smaller sample size might underpower the current findings. For

example, while our sham group showed similar behavioral patterns reported in the previous

study with healthy controls [9], previously reported Pavlovian bias in the reward domain (go-

to-win versus no-go-to-win) [9] was not replicated, as the bias was trending but not

statistically significant. Future studies should attempt to eliminate the practice effect from the

experimental protocol and increase the statistical power of the study. Also, we note that the

qualitative difference between electric shock and money might the experimental design and

result less consistent with the previous study [9], although electric shock has been used as

feedback alongside monetary gain in some studies [34–37]. Lastly, brain regions other than

dlPFC were not tested. Contrasting the effects of stimulating other brain regions to

stimulating dlPFC might strengthen the causal link between dlPFC and Pavlovian bias.

However, it should be also noted that we used a sham-controlled protocol, which is generally

considered as effective control condition [33,67,68].

In conclusion, our results suggest a causal relationship between non-invasive dlPFC

stimulation and corresponding decision-making behavior. We found reduced Pavlovian bias

in the punishment domain, go bias, and choice randomness after dlPFC facilitation using

anodal stimulation. However, further clarification using neuroimaging techniques is needed

to identify the neural mechanism that underlies the effects of tDCS; efforts are also needed to

determine how biases are modulated by neural changes in the dlPFC and connected brain
networks. In addition, because decision-making biases have been implicated in addictive
disorders, our results have practical implications for the treatment of individuals with such
disorders. Furthermore, only Pavlovian bias in the punishment, but not the reward domain,
was modulated; thus, there is a need for further studies concerning aversive-related decision-
making to explain why behavior related to avoiding an aversive state was only modulated by
tDCS.

Code accessibility

494 The codes are publicly available in the GitHub repository (https://github.com/CCS-
495 Lab/project_tDCS_public).

Acknowledgments

498 We thank all the CCS lab members who provided their valuable contributions throughout
499 various stages of this research.

Supporting Information

S1 Fig. The order effect found in choice accuracy. When comparing the task performance
of the first (third day) and the second time (sixth day) the participants conducted, we found
significantly higher accuracy in the second time. This indicates that the second task
performance is vulnerable to confounding effects caused by the task order. Therefore, we
included only the first task performance to eliminate this potential confounding factor. Error
bars indicate SEM. *$p < 0.05$ (TTF)

S2 Fig. One-step ahead prediction of computational models. We conducted the one-step-

ahead prediction to compare the predicted results and actual choice behaviors. To generate

predictions, we utilized 8000 MCMC samples (2000 samples x 4 chains) drawn from

individual posterior distributions to predict the probability of a "Go" choice in each of the

four conditions, and averaged the predictions with each participant. Model 3 (7-parameter

model) emerged as the best performing model, with a mean correlation coefficient of 0.890.

(TTF)

S3 Fig. Group prediction accuracy of parameters from Model 2 and Model 3. Although

Model 3 was the best model, Model 2 also showed closely comparable prediction values

(mean correlation value = 0.885). The only difference between the two models is the

utilization of separate Pavlovian bias parameters for reward and punishment. Consequently,

we further compared Models 2 and 3 and investigated whether having two parameters for

Pavlovian bias (as in Model 3) would aid in classifying tDCS group membership. By

employing ridge logistic regression and leave-one-out cross-validation, we compared the

classification accuracy of both models. Model 3 (AUC=0.806) demonstrated enhanced

classification accuracy in comparison to Model 2 (AUC=0.710). Error bars indicate SEM.

(TTF)

S1 Table. Side effect. (DOCX)

S2 Table. manipulation check of double-blind design. (DOCX)

S1 File. Extended methods. Includes information on electric shock protocol. (DOCX)

Reference

1. Bach DR, Dayan P. Algorithms for survival: a comparative perspective on emotions. Nat Rev Neurosci. 2017;18: 311–319.

2. Williams DR, Williams H. Auto-maintenance in the pigeon: sustained pecking despite contingent non-reinforcement. J Exp Anal Behav. 1969;12: 511–520.

3. Hershberger WA. An approach through the looking-glass. Anim Learn Behav. 1986;14: 443–451.

4. Hare TA, Camerer CF, Rangel A. Self-control in decision-making involves modulation of the vmPFC valuation system. Science. 2009;324: 646–648.

5. Hare TA, Malmaud J, Rangel A. Focusing attention on the health aspects of foods changes value signals in vmPFC and improves dietary choice. J Neurosci. 2011;31: 11077–11087.

6. Childress AR, Hole AV, Ehrman RN, Robbins SJ, McLellan AT, O'Brien CP. Cue reactivity and cue reactivity interventions in drug dependence. NIDA Res Monogr. 1993;137: 73–95.

7. Cavanagh JF, Eisenberg I, Guitart-Masip M, Huys Q, Frank MJ. Frontal theta overrides pavlovian learning biases. J Neurosci. 2013;33: 8541–8548.

8. Gershman SJ, Guitart-Masip M, Cavanagh JF. Neural signatures of arbitration between Pavlovian and instrumental action selection. PLoS Comput Biol. 2021;17: e1008553.

9. Guitart-Masip M, Huys QJM, Fuentemilla L, Dayan P, Duzel E, Dolan RJ. Go and no-go learning in reward and punishment: Interactions between affect and effect. Neuroimage. 2012;62: 154–166.

10. Rudorf S, Hare TA. Interactions between dorsolateral and ventromedial prefrontal cortex underlie context-dependent stimulus valuation in goal-directed choice. J Neurosci. 2014;34: 15988–15996.

11. Ridderinkhof KR, Ullsperger M, Crone EA, Nieuwenhuis S. The role of the medial frontal cortex in cognitive control. Science. 2004;306: 443–447.

12. Samejima K, Doya K. Multiple representations of belief states and action values in corticobasal ganglia loops. Ann N Y Acad Sci. 2007;1104: 213–228.

13. Gläscher J, Rudrauf D, Colom R, Paul LK, Tranel D, Damasio H, et al. Distributed neural system for general intelligence revealed by lesion mapping. Proc Natl Acad Sci U S A. 2010;107: 4705–4709.

14. Baumgartner T, Knoch D, Hotz P, Eisenegger C, Fehr E. Dorsolateral and ventromedial prefrontal cortex orchestrate normative choice. Nat Neurosci. 2011;14: 1468–1474.

15. MacDonald AW 3rd, Cohen JD, Stenger VA, Carter CS. Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. Science. 2000;288: 1835–1838.

16. Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. Annu Rev Neurosci. 2001;24: 167–202.

17. Averbeck B, O'Doherty JP. Reinforcement-learning in fronto-striatal circuits. Neuropsychopharmacology. 2022;47: 147–162.

18.   Weissengruber S, Lee SW, O'Doherty JP, Ruff CC. Neurostimulation Reveals Context-
Dependent Arbitration Between Model-Based and Model-Free Reinforcement Learning. Cereb
Cortex. 2019;29: 4850–4862.

19.   Ly V, Bergmann TO, Gladwin TE, Volman I, Usberti N, Cools R, et al. Reduced Affective
Biasing of Instrumental Action With tDCS Over the Prefrontal Cortex. Brain Stimul. 2016;9:
380–387.

20.   Smittenaar P, Prichard G, FitzGerald THB, Diedrichsen J, Dolan RJ. Transcranial direct current
stimulation of right dorsolateral prefrontal cortex does not affect model-based or model-free
reinforcement learning in humans. PLoS One. 2014;9: e86850.

21.   Ye H, Chen S, Huang D, Wang S, Luo J. Modulating activity in the prefrontal cortex changes
decision-making for risky gains and losses: a transcranial direct current stimulation study. Behav
Brain Res. 2015;286: 17–21.

22.   Fecteau S, Knoch D, Fregni F, Sultani N, Boggio P, Pascual-Leone A. Diminishing risk-taking
behavior by modulating activity in the prefrontal cortex: a direct current stimulation study. J
Neurosci. 2007;27: 12500–12505.

23.   Cohen Kadosh R, Soskic S, Iuculano T, Kanai R, Walsh V. Modulating Neuronal Activity
Produces Specific and Long-Lasting Changes in Numerical Competence. Curr Biol. 2010;20:
2016–2020.

24.   Alonzo A, Brassil J, Taylor JL, Martin D, Loo CK. Daily transcranial direct current stimulation
(tDCS) leads to greater increases in cortical excitability than second daily transcranial direct
current stimulation. Brain Stimul. 2012;5: 208–213.

25.   Gálvez V, Alonzo A, Martin D, Loo CK. Transcranial direct current stimulation treatment
protocols: should stimulus intensity be constant or incremental over multiple sessions? Int J
Neuropsychopharmacol. 2013;16: 13–21.

26.   Krause B, Cohen Kadosh R. Not all brains are created equal: the relevance of individual
differences in responsiveness to transcranial electrical stimulation. Front Syst Neurosci. 2014;8:
25.

27.   Li LM, Uehara K, Hanakawa T. The contribution of interindividual factors to variability of
response in transcranial direct current stimulation studies. Front Cell Neurosci. 2015;9: 181.

28.   Nitsche MA, Paulus W. Sustained excitability elevations induced by transcranial DC motor
cortex stimulation in humans. Neurology. 2001;57: 1899–1901.

29.   Nitsche MA, Liebetanz D, Lang N, Antal A, Tergau F, Paulus W. Safety criteria for transcranial
direct current stimulation (tDCS) in humans. Clin Neurophysiol. 2003;114: 2220–2; author reply
2222-3.

30.   Nitsche MA, Fricke K, Henschke U, Schlitterlau A, Liebetanz D, Lang N, et al. Pharmacological
modulation of cortical excitability shifts induced by transcranial direct current stimulation in
humans. J Physiol. 2003;553: 293–301.

31.   Im C-H, Park J-H, Shim M, Chang WH, Kim Y-H. Evaluation of local electric fields generated
by transcranial direct current stimulation with an extracephalic reference electrode based on
realistic 3D body modeling. Phys Med Biol. 2012;57: 2137–2150.

32.   Wagner T, Fregni F, Fecteau S, Grodzinsky A, Zahn M, Pascual-Leone A. Transcranial direct
current stimulation: a computer-based human model study. Neuroimage. 2007;35: 1113–1124.

33.   Thair H, Holloway AL, Newport R, Smith AD. Transcranial Direct Current Stimulation (tDCS):
A Beginner's Guide for Design and Implementation. Front Neurosci. 2017;11: 641.

34.   Lawson RP, Seymour B, Loh E, Lutti A, Dolan RJ, Dayan P, et al. The habenula encodes
negative motivational value associated with primary punishment in humans. Proc Natl Acad Sci
U S A. 2014;111: 11858–11863.

35.   Choi JM, Padmala S, Spechler P, Pessoa L. Pervasive competition between threat and reward in
the brain. Soc Cogn Affect Neurosci. 2014;9: 737–750.

36.   Talmi D, Dayan P, Kiebel SJ, Frith CD, Dolan RJ. How humans integrate the prospects of pain
and reward during choice. J Neurosci. 2009;29: 14617–14626.

37.   Vlaev I, Seymour B, Dolan RJ, Chater N. The price of pain and the value of suffering. Psychol
Sci. 2009;20: 309–317.

38.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing; 2020. Available: https://www.R-project.org/

39.   Berger JO. Statistical Decision Theory and Bayesian Analysis. Springer Science & Business
Media; 2013.

40.   Gelman A, Carlin JB, Stern HS, Rubin DB. Bayesian data analysis. Chapman and Hall/CRC;
1995.

41.   Lee MD. How cognitive modeling can benefit from hierarchical Bayesian models. J Math
Psychol. 2011;55: 1–7.

42.   Ahn W-Y, Krawitz A, Kim W, Busmeyer JR, Brown JW. A Model-Based fMRI Analysis with
Hierarchical Bayesian Parameter Estimation. J Neurosci Psychol Econ. 2011;4: 95–110.

43.   Ahn W-Y, Haines N, Zhang L. Revealing Neurocomputational Mechanisms of Reinforcement
Learning and Decision-Making With the hBayesDM Package. Comput Psychiatr. 2017;1: 24–57.

44.   Stan Development Team. RStan: the R interface to Stan. 2022. Available: https://mc-stan.org/

45.   Papaspiliopoulos O, Roberts GO, Sköld M. A General Framework for the Parametrization of
Hierarchical Models. Stat Sci. 2007;22: 59–73.

46.   Vehtari A, Gelman A, Simpson D, Carpenter B, Bürkner P-C. Rank-normalization, folding, and
localization: An improved R^ for assessing convergence of MCMC (with discussion). Bayesian
Anal. 2021;16. doi:10.1214/20-ba1221

47.   Vehtari A, Gelman A, Gabry J. Practical Bayesian model evaluation using leave-one-out cross-
validation and WAIC. Stat Comput. 2017;27: 1413–1432.

48.   Kruschke J. Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan. Academic Press;
2014.

49.   Griffiths KR, Morris RW, Balleine BW. Translational studies of goal-directed action as a
framework for classifying deficits across psychiatric disorders. Front Syst Neurosci. 2014;8: 101.

50.   Fonteneau C, Redoute J, Haesebaert F, Le Bars D, Costes N, Suaud-Chagny M-F, et al. Frontal
Transcranial Direct Current Stimulation Induces Dopamine Release in the Ventral Striatum in
Human. Cereb Cortex. 2018;28: 2636–2646.

51.   Fukai M, Bunai T, Hirosawa T, Kikuchi M, Ito S, Minabe Y, et al. Endogenous dopamine release

under transcranial direct-current stimulation governs enhanced attention: a study with positron
emission tomography. Transl Psychiatry. 2019;9: 115.

52. Gueguen MCM, Lopez-Persem A, Billeke P, Lachaux J-P, Rheims S, Kahane P, et al.
Anatomical dissociation of intracerebral signals for reward and punishment prediction errors in
humans. Nat Commun. 2021;12: 3344.

53. Lloyd K, Dayan P. Safety out of control: dopamine and defence. Behav Brain Funct. 2016;12:
15.

54. Millner AJ, Gershman SJ, Nock MK, den Ouden HEM. Pavlovian Control of Escape and
Avoidance. J Cogn Neurosci. 2018;30: 1379–1390.

55. Paulus MP. Driven by Pain, Not Gain: Computational Approaches to Aversion-Related Decision
Making in Psychiatry. Biol Psychiatry. 2020;87: 359–367.

56. Dayan, Huys. Neurophysiology: Serotonin's many meanings elude simple theories. Elife. 2015.
Available: https://elifesciences.org/articles/7390

57. Graeff FG, Silveira Filho NG. Behavioral inhibition induced by electrical stimulation of the
median raphe nucleus of the rat. Physiol Behav. 1978;21: 477–484.

58. Crockett MJ, Clark L, Robbins TW. Reconciling the role of serotonin in behavioral inhibition
and aversion: acute tryptophan depletion abolishes punishment-induced inhibition in humans. J
Neurosci. 2009;29: 11993–11999.

59. Crockett MJ, Clark L, Apergis-Schoute AM, Morein-Zamir S, Robbins TW. Serotonin
modulates the effects of Pavlovian aversive predictions on response vigor.
Neuropsychopharmacology. 2012;37: 2244–2252.

60. Navratilova E, Porreca F. Reward and motivation in pain and pain relief. Nat Neurosci. 2014.
Available:
https://idp.nature.com/authorize/casa?redirect_uri=https://www.nature.com/articles/nn.3811&cas
a_token=mV6mzYAzdI4AAAAA:iN9Jw421XS2oV0KY-
RPhvi0qotzoCn3K4pqIDqqeHXHrA8RBCySZLavOtFITL-GCF5jI0FBRkyEjWfo_9Gw

61. Everitt BJ, Belin D, Economidou D, Pelloux Y, Dalley JW, Robbins TW. Neural mechanisms
underlying the vulnerability to develop compulsive drug-seeking habits and addiction. Philos
Trans R Soc Lond B Biol Sci. 2008;363: 3125–3135.

62. Flagel SB, Robinson TE, Clark JJ, Clinton SM, Watson SJ, Seeman P, et al. An animal model of
genetic vulnerability to behavioral disinhibition and responsiveness to reward-related cues:
implications for addiction. Neuropsychopharmacology. 2010;35: 388–400.

63. Zilverstand A, Huang AS, Alia-Klein N, Goldstein RZ. Neuroimaging Impaired Response
Inhibition and Salience Attribution in Human Drug Addiction: A Systematic Review. Neuron.
2018;98: 886–903.

64. Spechler PA, Chaarani B, Hudson KE, Potter A, Foxe JJ, Garavan H. Response inhibition and
addiction medicine: from use to abstinence. Prog Brain Res. 2016;223: 143–164.

65. Zhukovsky P, Puaud M, Jupp B, Sala-Bayo J, Alsiö J, Xia J, et al. Withdrawal from escalated
cocaine self-administration impairs reversal learning by disrupting the effects of negative
feedback on reward exploitation: a behavioral and computational analysis.
Neuropsychopharmacology. 2019;44: 2163–2173.

66. Peters J, Vega T, Weinstein D, Mitchell J, Kayser A. Dopamine and Risky Decision-Making in

Gambling Disorder. eNeuro. 2020;7. doi:10.1523/ENEURO.0461-19.2020

67. Ambrus GG, Al-Moyed H, Chaieb L, Sarp L, Antal A, Paulus W. The fade-in--short stimulation-
-fade out approach to sham tDCS--reliable at 1 mA for naïve and experienced subjects, but not
investigators. Brain Stimul. 2012;5: 499–504.

68. Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): a tool for double-
blind sham-controlled clinical studies in brain stimulation. Clin Neurophysiol. 2006;117: 845–
850.

Stimulation

A. Choice accuracy (**anode**)

B. Choice accuracy (**sham**)

C. Pavlovian bias
(punishment domain)

ns   ns

ns   *

*

Go to win   Nogo to win   Go to avoid   Nogo to avoid

Go to win   Nogo to win   Go to avoid   Nogo to avoid

sham   anode

Choice accuracy (0-1)

Choice accuracy (0-1)

(Nogo-to-Avoid - Go-to-Avoid)

Task conditions

Task conditions

Type of stimulation