Enhanced intramyocardial vascular progenitor cell delivery promotes post-myocardial infarction healing by polarizing pro-regenerative neutrophils

Authors and Affiliations

Xuechong Hong [1,2], Allen Chilun Luo [1], Ilias Doulamis [1,2], Nicholas Oh [1,2], Pedro J. del Nido [1,2], Juan M. Melero-Martin [1,2,3], Ruei-Zeng Lin [1,2]

[1] Department of Cardiac Surgery, Boston Children's Hospital, Boston, MA 02115, USA
[2] Department of Surgery, Harvard Medical School, Boston, MA 02115, USA
[3] Harvard Stem Cell Institute, Cambridge, MA 02138, USA

Corresponding author

Ruei-Zeng Lin, Ph.D.
Department of Cardiac Surgery
Boston Children's Hospital
Harvard Medical School
300 Longwood Ave., Enders 345
Boston, MA 02115
Tel.: (617) 919-1905
Fax: (617) 730-0235
E-mail: ruei-zeng.lin@childrens.harvard.edu

Abstract

The success of vascular progenitor cell transplantation to treat myocardial infarction (MI) is primarily limited by the low engraftment of the delivered cells. Contracting myocardium actively pushes therapeutic cells injected in liquid vehicles out of the heart, resulting in low cell retention and poor therapeutic outcome. A clinically applicable biomaterial to improve cell retention is arguably needed to enable optimization of intramyocardial cell delivery. In this study, we used a preclinical MI model by ligating the left anterior descending arteries in mice. We developed a novel therapeutic cell delivery method for MI treatment based on a photocrosslinkable gelatin methacryloyl (GelMA) hydrogel. Human endothelial colony-forming cells (ECFCs) and mesenchymal stem cells (MSCs), a combination of vascular progenitor cells with the capacity to form functional vascular networks after transplantation in vivo, were injected in GelMA precursor solution into the infarcted zone of mouse hearts. The injected area was immediately illuminated by UV light to produce an in-situ photocrosslinked, cell-laden hydrogel. This rapid photopolymerization approach significantly enhanced cell retention in the beating hearts, achieving a sustained presence of the implanted cells and their therapeutic effects. The improved acute cell retention translated to a long-term beneficial post-MI cardiac healing, including stabilizing cardiac functions, preserving viable myocardium, and preventing cardiac fibrosis. Furthermore, this technical advance was used to demonstrate that the mechanism of post-MI cardiac healing involved the polarization of pro-regenerative neutrophils by the delivered vascular cells. Indeed, the engrafted cells polarized recruited bone marrow-derived neutrophils toward a non-inflammatory phenotype via TGFβ signaling, establishing a pro-regenerative microenvironment that favored post-MI cardiac healing. Depletion of host neutrophils canceled the therapeutic benefits produced by cell delivery in the ischemic hearts, indicating that the non-inflammatory, pro-regenerative neutrophils were indispensable mediators of cardiac remodeling. In summary, our novel GelMA hydrogel-based intramyocardial vascular cell delivery approach has the potential to improve the treatment of acute MI.

Introduction

Myocardial infarction (MI) remains the leading cause of morbidity and mortality worldwide [1,2]. MI produces irreversible damage to the heart muscle due to a blockage of coronary arteries. Myocardial necrosis starts rapidly after the onset of ischemia, leaving an infarct zone that contains nonfunctional myocytes and excessive inflammatory cells [3]. MI patients suffer from decreased cardiac outputs, resulting in a downward spiral leading to congestive heart failure [1]. In the clinics, the acute mortality rate of MI has fallen dramatically through current surgical interventions aided by pharmacotherapy and mechanical devices [4]. Notwithstanding the short-term benefit of these interventions, only modest improvements in global cardiac functions were observed in patients six months after an acute MI [5,6]. A significant number of patients go on to develop late sequelae, including heart failure and arrhythmia [2]. For years, considerable effort has been focused on delivering therapeutic genes, growth factors, and cells to promote post-myocardial infarction healing [7–11]. However, these strategies are insufficient partly because of the complexity of the hyper-inflammatory microenvironment in the infarcted myocardium [12–16].

While the human heart is considered a terminally differentiated organ, adult cardiac regeneration by endogenous repair mechanisms is unfeasible with current medical options [17]. Over the last decade, many investigators have tested cell-based therapies to prevent adverse ventricular remodeling and promote recovery of MI in animal models, as well as in clinical trials [8,10,18]. Delivery of therapeutic cells to the heart is performed in two main categories. First, cardiac patches or cell sheets consisting of pluripotent stem cell-derived cardiomyocytes were transplanted to reinforce or replace the injured myocardium aiming to rebuild the loss of heart walls [19]. These cardiac patches are mainly manufactured by seeding cells in biomaterial scaffolds and preconditioned in a bioreactor before surgical implantation. The second approach injected therapeutic cells directly into the infarcted areas to mitigate the detrimental impact of MI [20]. In human patients, delivery of cells could be achieved by intramyocardial or intracoronary injection routes through a catheter-based minimally invasive surgical approach, which allows the early therapeutic intervention in the acute MI phase [9]. Examples of tested adult stem/progenitor cells include cardiosphere-derived cells [21], bone marrow-derived mononuclear cells [22,23], endothelial progenitor cells [24], and mesenchymal stem cells [25]. However, except for cardiomyocytes, the long-term engraftment of injected cells was shown to be unnecessary to exhibit therapeutic effects. Accumulating evidence suggests non-myocyte cell types can provide beneficial effects through paracrine factors aiding cardiac remodeling or the modulation of inflammatory response [7,12,25]. However, the underlying mechanisms are not well-understood.

Delivering vascular progenitor cells is a promising new therapeutic strategy for ischemic cardiomyopathy. Some studies have focused on delivering vascular cells to build new blood vessels and stimulate angiogenesis [26–28]. The concept was to enable rapid restoration of microvascular networks within the ischemic myocardium to mitigate adverse remodeling. For instance, in a rat myocardial ischemia/reperfusion injury (IRI) model, animals injected with human endothelial colony-forming cells (ECFCs) and mesenchymal stem cells (MSCs) in PBS revealed a modest reduction in adverse ventricular remodeling [26]. The quantitative analysis showed a rapid loss of human cells shortly after myocardial injection, suggesting the lack of sufficient cellular retention could be the major reason for the weak effect [26]. Therefore, it is still an open question whether human vascular progenitor cells with vasculogenic capability can mitigate adverse remodeling and promote functional recovery.

Low cell retention and engraftment are the major obstacles to achieving a significant functional benefit in MI, irrespective of the cell type used [29]. Beating hearts actively push therapeutic cells injected in liquid vehicles out of the myocardium, resulting in low acute cell retention between 0.5-10% across several studies with different cell types, animal models, and delivery routes [29–35]. To address this problem, several biomaterials for enhancing cell engraftment have been tested in the last decade [36–39]. In this study, our approach is to use well-characterized Gelatin methacryloyl (GelMA) hydrogel to deliver human ECFCs and MSCs to build neo-vascular networks in the ischemic myocardium. GelMA was synthesized in our previous studies to be a photocrosslinkable, bioadhesive hydrogel following its injection in tissues [40–42]. Cell-laden GelMA

formulation can polymerize rapidly (10 s upon exposure to UV light in the presence of a photoinitiator) in myocardial tissues without compromising the therapeutic cell viability. We proposed that this rapid polymerization would be a critical feature to avoid hydrogel dissemination at the implantation site. Recently, we also demonstrated that this material is fully compatible with ECFC-based vascular morphogenesis [41], and thus we proposed its use for vascular cell therapy in infarct hearts. Here, we demonstrated that GelMA hydrogel can be transmyocardially photocrosslinked and maintain vascular cell retention in infarcted mouse hearts. GelMA-enabled cell delivery promoted infarct healing and adaptive remodeling to preserve cardiac function. Importantly, enhanced cell retention allows us to study and discover the early beneficial effects of human vascular cells in MI for the first time. We found that the enhanced retention of human ECFCs and MSCs enabled the perivascular-endothelial cell interactions to upregulate transforming growth factor beta (TGFβ) expression. TGFβ signaling promoted the polarization of non-inflammatory, pro-regenerative neutrophils, which were shown to be indispensable mediators of beneficial cardiac remodeling.

Results

In situ polymerization of GelMA hydrogel in infarcted myocardium

Transmyocardial polymerization of GelMA was evaluated in vivo using a mouse MI model (**Figure 1**). MI was induced by permanent ligation of the left anterior descending (LAD) arteries (**Figure S1A** and **Video S1**, Supporting Information) in 10-12-week-old SCID mice. A mixture of human vascular progenitor cells (ECFCs and MSCs; $5x10^5$ for each cell type) suspended in 5% (w/v in PBS) GelMA precursor solution was injected into the infarcted myocardium within 10 min of LAD ligation. The total injection volume per heart was 100 μL. We first demonstrated that this injection volume was sufficient to cover the infarcted zone (**Figure S1B**). We then determined that UV light could transmit through the myocardium of mice using our OmniCure S2000 UV lamp (wavelength 320-500 nm; the intensity of 40 mW/cm^2). GelMA polymerization was achieved after just 10 seconds of transmyocardial exposure to UV light (**Video S2**). The general safety of short UV illumination by our UV lamp had been evaluated in a previous study [42]. Our working range of UV light exposure (10 - 45 seconds) was deemed safe for both injected cells and surrounding tissues [42]. Moreover, we did not observe leakage of GelMA solution or swelling of local tissues with the injection volume and crosslinking conditions used.

GelMA hydrogel enhanced cell retention in ischemic hearts

To evaluate the acute myocardial retention of human vascular progenitor cells, we injected $5x10^5$ luciferase-labeled ECFCs (luc-ECFCs) with $5x10^5$ non-labeled MSCs per heart and quantified the bioluminescence intensity in the infarcted area (**Figure 1C**). We compared transmyocardial photocrosslinking of cell-laden GelMA hydrogel following injection into the infarcted zone (referred to as GelMA-Hydrogel or **GelMA-H** group; **Figure 1B**) with the results of cell injection in the same GelMA precursor solution but without UV illumination (GelMA-Solution or **GelMA-S** group). Notably, the GelMA precursor solution remained liquid in physiological conditions (i.e., at 37°C and in contact with blood or tissues) if not crosslinked by UV light [43]. Since cardiac contraction is primarily responsible for the early washout of cells from the injection site [31], we injected the same number of human cells into arrested mouse hearts and performed a bioluminescence measurement after 15 min to obtain a baseline. Thus, bioluminescence signals compared to this baseline were used to estimate the percentage of ECFC retention (**Figure 1D**).

As early as 3 h after MI induction and cell delivery, the presence of ECFCs in the GelMA-S group was rapidly reduced to 20.6 $\pm$ 2.3% (**Figure 1C-D**), a low retention rate compared to previous studies describing cell delivery in PBS [22,31,44]. Cardiac contraction potentially reduces cell retention by extrusion of the liquid injectate during each heartbeat. However, photopolymerization of GelMA hydrogel should, in principle, minimize this cell backwash. Indeed, in the GelMA-H group, ECFC retention was significantly improved to 48.6 $\pm$ 13.6% (3 h, p < 0.05 vs. cells in GelMA-S; **Figure 1D**), revealing a dramatic effect of sealing the injection site on cell

retention. The higher ECFC retention in the GelMA-H group was maintained throughout the 3-day monitoring post-MI (**Figure 1D**).

To confirm that the GelMA-H enhanced the retention of both endothelial and mesenchymal cells, we quantified non-labeled ECFCs and MSCs in infarcted hearts by flow cytometry (**Figure S2**, Supporting Information). Cell delivery in GelMA-H significantly enhanced both ECFC and MSC retentions at 48 h compared to the GelMA-S group (**Figure 1E**; p value = 0.0005 for ECFCs and 0.0728 for MSCs). Compared to the number of injected cells, the GelMA-H group preserved 25.2% of viable ECFCs and 20.8% of MSCs, while the GelMA-S group only retained 4.4% of ECFCs and 12.8% of MSCs. Notably, GelMA hydrogel maintained the engraftment of ECFCs and MSCs close to the original 1:1 ratio, which was optimized to achieve long-term vascularization [45]. Collectively, our data suggest that the rapid in situ photopolymerization of GelMA hydrogel adequately protected the human vascular progenitor cells and reduced the "washout" effect during the acute phase (72 h) following MI.

Human vascular progenitor cells delivered in GelMA hydrogel improve cardiac remodeling

To determine whether the improvement in acute cell retention would impact post-MI cardiac healing, we used echocardiography to evaluate the cardiac functions (**Figure 2 and S3**, Supporting Information). Echocardiography revealed no significant difference in baseline cardiac function between groups measured 1 h before LAD ligation. On day 7 following the induction of MI, left ventricular fractional shortening (LVFS) and left ventricular ejection fraction (LVEF), two measurements of general cardiac function, were higher in the mice treated with human cells delivered in GelMA-H compared to the untreated or GelMA-S-treated animals (**Figure 2B-C;** 30.3 $\pm$ 2.9%, 24.6 $\pm$ 3.2%, 25.2 $\pm$ 2.8% for LVFS; 66.4 $\pm$ 4.5, 56.9 $\pm$ 5.6, 56.8 $\pm$ 4.6 for LVEF, respectively). After treatment, both cardiac functions were significantly higher in the GelMA-H group. Compared with the GelMA-S group, LVEF and LVFS in the GelMA-H group were improved considerably on days 7, 14, and 21 (increased 20.2%, 18.7%, and 19.7% for LVFS; 16.9%, 17.4%, and 17.1% for LVEF). Both untreated and GelMA-S-treated groups showed a continuous decline in LVEF and LVFS from day 3 to 28 post-MI (MI-untreated: reduced 17.6% for LVFS and 13.4% for LVEF; GelMA-S: reduced 19.6% for LVFS and 14.3% for LVEF), indicating a deterioration of cardiac function due to adverse infarct remodeling. There was no significant difference in LVEF and LVFS between untreated or GelMA-S-treated groups, suggesting a lack of therapeutic benefit in groups with low cell retention. In contrast, in the GelMA-H group, cardiac function was generally preserved from day 3 to day 28 (GelMA-H: reduced only 9.5% for LVFS and 6.7% for LVEF).

Heart muscle salvage after MI is the single most important determinant factor for the long-term preservation of cardiac function and survival [2]. We measured the myocardial wall thickness by echocardiography. The GelMA-H group exhibited a significantly higher left ventricular anterior wall thickness (LVAW) compared to the untreated or GelMA-S groups (**Figure 2D**). Again, no improvement was found by comparing GelMA-S and untreated groups, suggesting that high cell retention was required for beneficial cardiac remodeling. The preservation of the myocardium was confirmed by histological analysis of cross-sections in the infarct area after 4 weeks (**Figure 2E**). In mice that received cells in GelMA-H, myocardial thickness significantly improved compared to the untreated or GelMA-S groups (**Figure 2F;** 0.44 $\pm$ 0.05 mm vs. 0.25 $\pm$ 0.06 mm, *p < 0.01). Fluorescent image assessment demonstrated that the number of human microvessels in GelMA-H-treated hearts was significantly higher in the infarcted area than in the GelMA-S group (**Figure 2G-H**). Masson's trichrome staining of cardiac tissues harvested at 4 weeks also confirmed that the GelMA-H group showed significantly reduced myocardial fibrosis compared to the untreated or GelMA-S-treated groups (**Figure 2I-J**). Taken together, our data indicate that improving acute cell retention translates to beneficial cardiac remodeling, including enhanced revascularization, less scar tissue formation, and preservation of overall cardiac functions after MI.

Functional human vasculature in infarct hearts

We sought to investigate the therapeutic mechanism resulting from the delivery of human vascular progenitor cells in GelMA-H. First, we examined the infarcted hearts on day 7 to assess

the formation of human blood vessels [26]. The histological examination carried out by immunofluorescence staining of human-specific vimentin and UEA1 revealed that the extent of vascular network formation differed in the GelMA-H and GelMA-S groups (**Figure 2G**). Significant engraftment of human ECFCs and MSCs in the GelMA-H groups was evidenced by the formation of human-specific vascular networks inside the infarcted myocardium (**Figure 2H**). In the GelMA-S group, only a few human MSCs, but not ECFCs, were stained in the infarcted myocardium. These data suggested that the in situ photocrosslinking of the GelMA hydrogel significantly improved the retention of co-delivered human ECFCs and MSCs. Without the timely gelation of GelMA hydrogel, although some MSCs could engraft, ECFCs failed to participate in the revascularization of the infarcted myocardium.

The formation of functional human vasculature in ischemic tissues might explain parts of the therapeutic benefit observed in the GelMA-H group. However, we noticed significant improvement of LVAW in the GelMA-H group (**Figure 2D**) as early as day 3, and improved cardiac functions by day 7 (LVFS and LVEF; **Figure 2B-C**). Considering that the onset of human blood vessel perfusion takes approximately 7 days in this model of vascular progenitor cell implantation [41,42,46–48], this timing inconsistency between human cell-mediated revascularization and beneficial cardiac remodeling suggested that other therapeutic pathways might be involved in preserving cardiac functions during the early stage of MI.

Gene expression analysis upon human vascular progenitor cell delivery in infarcted hearts

To further elucidate the molecular aspects of the therapeutic benefits produced by ECFC+MSC delivery in GelMA-H, we conducted an exploratory transcriptional analysis using RNA sequencing (RNAseq). The infarcted left ventricle tissues were harvested from GelMA-H and GelMA-S groups on day 2 post-MI to study the change of gene expression profiles in the early stage of treatments. Tissues from uninjured hearts served as controls.

Globally, there were 2,088 differentially expressed genes (DEGs) among the three analyzed groups (**Figure S4A**). Hierarchical clustering analysis (**Figure S4B**) revealed the proximity between two post-MI groups (GelMA-H and GelMA-S) in comparison to uninjured controls. This pattern was confirmed by the principal component analysis (PCA; **Figure S4C).** To gain more insight into the transcriptional difference between GelMA-H and GelMA-S groups, we carried out a pairwise analysis of DEGs. There were 137 genes significantly upregulated in the GelMA-H group whereas 617 genes were downregulated (**Figure S4A**, **Figure S4D,** GelMA-H vs. GelMA-S). Notably, we observed significant upregulation of several genes related to neutrophil development and maturation, including expression of *s100a8, s100a9, elane, ngp, mpo, camp, lef1, prg3, mmp3, and ms4a2*. This was associated with significant upregulation of genes implicated in immature neutrophil trafficking (*sdf, cxcr4*), hematopoiesis (*gata1, hemgn*), and immunomodulation (*il4, il10, tgfb*) in GelMA-H vs. GelMA-S (**Figure S4D-E**). Altogether, the data suggested a potential effect of human ECFCs+MSCs in GelMA-H on the recruitment and polarization of neutrophil subpopulations in the early stage of MI, although it is important to note that the data variation between samples within the GelMA-H group was high.

Recruitment of neutrophils with reparative phenotype in GelMA-H group

RNAseq data analysis suggested a significant difference in neutrophil-related genes between GelMA-H and GelMA-S groups. Previously, we demonstrated that in vivo implanted human vascular progenitor cells could alternatively polarize neutrophils into a non-iumflammatory phenotype, mediating vascular assembly and anastomosis [47]. Moreover, recent studies have shown that pro-regenerative neutrophils (herein termed NR; "R" for regeneration) were implicated in orchestrating tissue repair in various organs [49–59]. However, the role of NR in post-MI healing remains unclear. Therefore, we focused our studies on the polarization of the neutrophils following MI.

We first examine the early-stage host cell infiltration post-MI (**Figure 3**). We analyzed by flow cytometry the recruited mouse CD45[+] leucocytes on day 2 in infarcted myocardium and classified them into Ly6G[+]F4/80[-] neutrophils or Ly6G[-]F4/80[+] macrophages (**Figure 3A**). Infiltration of Ly6G[-]

F4/80$^-$ lymphocytes was negligible, as expected in immunodeficient mice. Also, the presence of murine resident myeloid cells in the uninjured myocardium was low (**Figure 3B**). LAD ligation followed by either GelMA-H or GelMA-S treatment significantly recruited host myeloid cells (**Figure 3B**), which is consistent with the induced innate immune response observed by RNAseq analysis. Moreover, the GelMA-H group recruited more neutrophils and macrophages than the GelMA-S group (**Figure 3B**), which is consistent with our previous observation on the engagement of host myeloid cells by ECFCs and MSCs [47]. We further evaluated neutrophil subpopulations using flow cytometry and classified Ly6G$^+$F4/80$^-$ cells into naïve neutrophils (N0; CXCR2-), pro-inflammatory neutrophils (NI; CXCR2$^+$CD206$^-$), and pro-regenerative neutrophils (NR; CXCR2$^+$CD206$^+$; **Figure 3A and C**). The NI was the dominant subpopulation (83.27% of total neutrophils; **Figure 3D**) in MI mice treated with GelMA-S, which is consistent with previous report that showed pro-inflammatory neutrophils significantly mediating adverse cardiac remodeling in the first 7 days post-MI [52]. In contrast, NIs were considerably reduced (18.02%) in the GelMA-H group, where NRs were significantly more abundant (57.66% of total neutrophils), suggesting a shift toward NR polarization. The GelMA-H group also recruited more naïve neutrophils from the bone marrow than that in the GelMA-S group (24.32% vs. 13.31%). Together, these data suggested that improving human vascular progenitor cell retention could translate into a pro-regenerative host neutrophil polarization during the early stage of MI healing. Since NRs have been previously shown to promote tissue healing by facilitating the resolution of inflammation and stimulating angiogenesis, cell proliferation, and ECM remodeling [47,60–62], the increased abundance of NR in the GelMA-H group could in part explain the beneficial effects observed in cardiac function.

Human vascular cell interaction upregulated NR polarization factor TGFβ1

We next examined if the prolonged retention and interaction between human ECFCs and MSCs in the GelMA-H group resulted in a release of paracrine factors that could promote NR polarization and whether these factors were absent in the GelMA-S group. To this end, we collected conditioned media from ECFC+MSC coculture in GelMA hydrogel over 7 days in vitro (CM^{CO}; **Figure 4A**). Next, murine naïve neutrophils isolated from bone marrow (BM) were cultured in CM^{CO} for 24 h. NR polarization was evidenced by the upregulation of *Il4*, *Vegfa* and *Arg1* gene expressions upon exposure to CM^{CO} (**Figure 4B**). However, the evidence for NR polarization was minimal when BM neutrophils were cultured in basal medium or conditioned media obtained from ECFC or MSC monoculture (CM^{ECFC} or CM^{MSC}), suggesting an important role for ECFC-MSC crosstalk in the polarization of NR.

To gain further insight into the NR polarization factors, we performed a proteomic analysis of the conditioned media from the ECFC+MSC coculture (**Figure 4C-D**). Mixed conditioned media (1:1 mixture of CM^{ECFC}+CM^{MSC}; termed CM^{MIX}) obtained from separate ECFC and MSC monocultures served as a control. The secretion of several growth factors was induced in the ECFC+MSC coculture, including Activin A, IGFBP-1, Endoglin, Amphiregulin, and TGFβ1 (the increase in TGF-β1 was consistent with our previous study that showed it promotes NR polarization [47]). We confirmed at the mRNA level that the ECFC+MSC coculture upregulated TGFβ1 expression only in ECFCs but not in MSCs (**Figure 4E-F**). Moreover, exposing the ECFC+MSC coculture to the Notch inhibitor DAPT (a γ-secretase inhibitor) blocked the induction of TGFβ1 expression (Notch signaling is known to mediate vascular cell interaction [47]). Together, these results suggested that NR polarization was directly related to the engrafted human vascular progenitor cells in the infarcted tissues. Notch-mediated interaction of ECFCs and MSCs could modulate the early innate immune landscape by increasing TGFβ1 secretion by the ECFCs.

Neutrophil depletion abrogates the improvement in cardiac remodeling

To confirm the pivotal role of host neutrophils during cardiac remodeling, we depleted circulating neutrophils in MI mice treated with GelMA-H. To this end, we treated mice from two days before LAD ligation to post-operative day 6 with an anti-Gr1 or IgG (200 µg) control antibody given every 2 days via intraperitoneal injection (**Figure 5A**). This treatment with anti-Gr1 antibody

was effective at depleting circulating neutrophils without altering blood monocytes and lymphocytes (**Figure 5B-C**). Importantly, neutrophil depletion abolished the improved post-MI cardiac functions observed at days 3 and 7 in the GelMA-H group (LVEF and LVFS; **Figure 5D, 5E**) and significantly reduced myocardium thickness at day 7 (LVAW, **Figure 5F**). All three measurements of cardiac functions (LVEF, LVFS, and LVAW) in neutrophil-depleted mice were comparable to those observed in the untreated or GelMA-S groups (**Figure 2**). Together, these findings suggest that depletion of neutrophils leads to excessive loss of myocardium that might explain the progressive worsening of cardiac function. Indeed, host neutrophils were indispensable for the therapeutic effects exhibited by human vascular cells delivered in GelMA-H.

Discussion

An effective early intervention to achieve sufficient cell retention in the infarcted myocardium has important implications for cardiac remodeling. In the present study, we demonstrated that human vascular progenitor cell retention in MI hearts could be significantly improved by delivering the cells in a photocrosslinked GelMA hydrogel. In contrast, cells injected in GelMA precursor solution were rapidly lost from the beating hearts. The improved acute cell retention showed a significant impact on post-MI cardiac remodeling. Echocardiography and histological examination demonstrated that the delivery of ECFC+MSC in the GelMA hydrogel prevented a drastic loss of cardiac functions, preserving viable myocardium, and averting cardiac fibrosis. The delivered human ECFCs and MSCs formed perfused vascular networks in the ischemic microenvironment. However, vascular regeneration was not the sole mechanism leading to cardiac healing. The human vascular cells also mediated the pro-regenerative polarization of recruited neutrophils, which in turn facilitated cardiac remodeling.

Leukocyte infiltration in ischemic areas is a hallmark of MI. The inflammatory response appears a double-edged sword because it is a protective attempt by the organism to initiate the healing process; however, prolonged inflammation can exacerbate scarring and loss of organ function [15,63]. Overwhelming infiltration of innate immune cells has been shown to promote adverse remodeling and cardiac rupture [64]. The successful resolution of inflammation appears as a critical event in the repair of tissue damage, but little is known about the primary mediators of resolution in the infarcted cardiac tissue [65–67]. Neutrophils massively infiltrate the heart soon after MI, where they can promote both tissue repair and damage [14,49]. Emerging evidence indicates that neutrophils comprise two distinct subpopulations with different polarized phenotypes: a canonical pro-inflammatory phenotype (referred to as NI) and an alternative anti-inflammatory, pro-regenerative, phenotype (NR), resembling the well-established 'M1–M2' macrophage paradigm [58,68,69]. Previous reports have suggested that cardiac neutrophils follow a temporal polarization pattern after MI, switching from a pro-inflammatory NI to an anti-inflammatory NR profile [13,50,52,70,71]. However, this switch could take 5-7 days post-MI in rodent models, while excessive inflammatory cells had produced irreversible damage to the surrounding myocardium [52]. Here we showed that an optimized delivery of human vascular progenitor cells could promote the alternative polarization of recruited host neutrophils in the ischemic tissues as early as 2 days post-MI. NRs have been shown to actively contribute to the repair of various tissues, mainly through suppressing pro-inflammatory reactions to facilitate resolution, promoting M2 macrophage polarization, and stimulating angiogenesis, cell proliferation, and ECM remodeling [47,50,51,58]. Indeed, we reason that the early transition to a NR-mediated regenerative response is part of the therapeutic mechanism achieved by human vascular progenitor cells delivered in infarcted hearts.

TGFβ has been shown to promote the NR phenotype [56,68]. NRs in the injury site derive from the TGFβ-mediated polarization of naïve neutrophils (N0) recruited from the bone marrow (BM) [68]. Blocking TGFβ signaling results in a depletion of the NR subpopulation and leads to a loss of vascularization [47]. We have shown that human ECFC+MSC implanted in the subcutaneous space of immunodeficient mice can form robust vascular networks connected with the host circulation within a week [42,48,72,73]. This vascularization process required host neutrophils expressing TGFβ

receptor 2 (TGFBR2) [47]. Human ECFC+MSC implanted in neutrophil-depleted animals (by sub-lethal radiation) completely failed to vascularize; however, an adoptive transfer of BM-derived neutrophils from unirradiated wild-type donor mice fully rescued vascularization [47]. Of note, the adoptive transfer of BM-derived neutrophils lacking TGFBR2 (from *Tgfbr2*$^{-/-}$ mice; *LysM-Cre*::*Tgfbr2/loxP*) failed to rescue vascularization, which underscored the critical role of TGFβ-TGFBR2 signaling in the generation of functional NRs [47]. Here, we further confirmed that the successfully engrafted human ECFCs are a critical source of active NR-polarizing factors. We found that the conditional medium from ECFC+MSC coculture in a GelMA hydrogel strongly promoted NR polarization, evidenced by the up-regulation of NR signature genes, *Arg1*, *Vegfa*, and *Il4*, in mouse BM neutrophils. ECFC-derived TGFβ1 was up-regulated only in the presence of MSCs. Furthermore, when the endothelial-perivascular cell interaction was disrupted by blocking Notch-mediated signaling, ECFCs failed to up-regulate TGFβ1 expression. Taken together, our data suggest that the beneficial effect of delivering human vascular progenitor cells in GelMA hydrogel was partly due to promoting TGFβ1 secretion and engaging host NRs, resulting in a pro-regenerative microenvironment that favors post-MI cardiac healing.

Over the years, numerous preclinical studies and clinical trials have evidenced the challenges of treating heart diseases with cell injections [29]. Systemic infusion of cells has been shown to be inefficient at delivering cells to the heart. It could even cause adverse effects when the injected cells accumulate in other organs, such as the lungs [74]. For local intracoronary injections, the injected cells need to undergo transendothelial migration to reach the infarcted areas [32]. Studies have shown the risks of forming plugs when transplanted cells pass through narrow capillaries, which have diameters similar to cells [29,75]. Thus, those blocked cells become the barrier that prevents other cells from traveling from the coronary vessels to the ischemic zone.

As one of the most straightforward routes to deliver cells to the heart, the intramyocardial injection has been frequently applied in preclinical animal studies and clinical trials [27,76–78]. With a needle or catheter, an intramyocardial injection can precisely deliver a sustainable number of therapeutic cells into pathological tissues in a short period of time. The intramyocardial injection can be performed by puncturing the myocardium from the epicardial side with a needle and injecting cells [29]. The epicardial intramyocardial injection is common in small animal models and was used in this study. However, this method requires open-chest surgery to expose the heart, which limits its clinical application. The endocardial intramyocardial injection also requires penetration of the myocardium, but instead, from the inside of the ventricular wall [78]. In human patients, this method can be achieved by a long catheter threaded through the peripheral blood vessels into the left ventricle, where the cells are injected through the endocardium. In general, this approach is less invasive because of it could avoid open-chest surgery [79]. For this reason, the endocardial intramyocardial injection possesses a higher translational potential [80]. However, it is also challenging to perform in small animal models.

A common hurdle is shared by all current delivery routes: low cell retention due to the "washout" effects of beating hearts. Given that the heart is a contractile pump, the beating of cardiomyocytes squeezes the narrow space in the myocardium and pushes transplanted cells in fluid medium out through the needle track or damaged blood vessels [29]. Many studies had reported a rapid loss of >80% of injected cells within hours after delivery [30,78]. Besides blood flow, the cells can be washed out of the heart by lymphatic drainage. In addition to physical washout from the heart, transplanted cells are lost over time through cell death before conferring a therapeutic benefit to the injured cardiac tissue. Inflammatory cells and damaged cardiomyocytes in the infarcted tissue can release cytokines and reactive oxygen species, which results in transplanted cell apoptosis [81]. Furthermore, the infarcted microenvironment is deficient in oxygen and nutrient supply. As a result, many transplanted cells die in these unfavorable living conditions.

A myriad of studies confirms that cell retention plays a crucial role in the success of cell-mediated cardiac repair [29]. Any delivery route needs to maintain donor cells in the recipient's heart for enough time to send paracrine signals to surrounding damaged heart cells and repair them. Challenges associated with the in vivo cell delivery of currently available vehicles have limited clinical translation of this technology. To improve the retention of therapeutic cells in the heart,

scientists have been focusing on bioengineering strategies, including designing novel biomaterials that protect cells from the washout effect in the heart [7,82].

Several injectable hydrogels have been designed with different base materials and cell type combinations [36]. Previously, synthetic polymers and nanogels were used to encapsulate cells for delivery into mouse and swine models of MI to demonstrate their safety and efficacy [37–39]. Notwithstanding the benefit of these new materials, the translational usage of synthetic materials faced obstacles due to the limited data on their long-term biocompatibility. Various natural extracellular matrix (ECM) products have been extensively tested for intramyocardial cell injection, including fibrin- or collagen-based hydrogels [31]. However, the slow gelation process of these natural hydrogels failed to achieve desirable cell retention. Accordingly, improving the properties of naturally occurring ECMs has become an alternative approach. This can be achieved by the chemical functionalization of proteins, an approach that was proposed precisely to improve the usability of biomaterials [83–86]. Here, we used GelMA synthesized by adding methacrylate groups to the amine-containing side-groups of gelatin, which becomes a photocrosslinkable, bioadhesive hydrogel [40,41]. As a result, GelMA hydrogels exhibit characteristics of both natural and synthetic biomaterials. GelMA is highly soluble in phosphate-buffered saline and can be prepared as a precursor solution (5-20% w/v) at room temperature. Therapeutic cells can be easily suspended in the GelMA precursor solution for needle- or catheter-based injections. Notably, the photocrosslinking reaction produces a GelMA hydrogel within 10 s of UV illumination without compromising encapsulated cell viability. Consequently, the quick gelation process of GelMA hydrogel effectively stops the washing out of cells injected into the hearts, resolving the main physical problem of low cell retention. Furthermore, GelMA hydrogel possesses excellent biocompatibility. In our previous studies, GelMA hydrogel provided an ideal microenvironment for the engraftment and vascularization of human vascular progenitor cells both in vitro and in vivo in immunodeficient mice [41,42]. In particular, GelMA contains gelatin as its backbone, which provides cell-responsive characteristics such as the provision of appropriate cell adhesion sites and proteolytic degradability. Degraded products of GelMA hydrogel are simply gelatin peptides, which are parts of natural ECM components and are non-cytotoxic or non-immunogenic to the hearts [40].

The intensity needed to photocrosslink GelMA hydrogels transmyocardially was analyzed for safety. Transmyocardial exposure to our UV light spectrum (UVA-visible: 320-500 nm) had little observable effect on the human cells and surrounding mouse tissues [42]. We tested these effects previously by exposing human cells and rodent skins to prolonged UV illumination (120 seconds) and showed minimal effects on cell apoptosis [42]. In this study, we achieved sufficient transmyocardial GelMA photocrosslinking with only 10 s of UV illumination. Therefore, our working range of UV light exposure was deemed sufficiently safe. Alternatively, GelMA could be formulated to form a hydrogel with visible light-responsive photoinitiators to eliminate the usage of UV light [87], although the crosslinking time would be significantly longer than with UV light [88]. Also, UV light illumination could be used through a fiber-optic system together with a catheter, as we previously showed in with a photocurable polymer patch to fix septal defects in the beating hearts of pigs [89]. Patches attached to the heart by UV crosslinking showed similar inflammatory response as sutures, with 100% small animal survival, indicating safety and applicability of UV system in minimally invasive delivery in hearts.

In situ, the photocrosslinked GelMA hydrogel functions as a cell delivery vehicle in our study. We had shown previously the implantation of GelMA hydrogel alone containing no cell provoked minimal host cell infiltration, suggesting GelMA is a relatively inert material [41]. A recent study by Ptaszek et al. applied acellular GelMA hydrogel on top of the epicardial surface of the mouse heart at the time of experimental MI [90]. Mice treated with these bioadhesive GelMA patches displayed improved post-MI survival rates and left ventricular contractile function, raising the opposability that our GelMA hydrogel may benefit cardiac healing by providing mechanical support. However, we compared our intramyocardial GelMA-cell injection to their GelMA bioadhesive application and found important differences between the two experimental conditions. In our study, we used minimal UV exposure time, and thus the mechanical properties of GelMA

hydrogel were relatively low compared to the GelMA bioadhesive protocol (elastin modulus, ~2 kPa vs. ~70 kPa). We reason that the disparity came from the fundamental difference in purpose between the two studies. We chose our GelMA crosslinking condition to optimize the viability and vasculogenic potential of the encapsulated human ECFC+MSC. We determined that a soft GelMA hydrogel (2 to 5 kPa) was permissive for the formation of functional vascular networks [41]. In contrast, the study by Ptaszek et al.aimed to reinforce the loss of mechanical strength of the infarcted myocardial wall [90]. They achieved a stiffer GelMA patch through a significantly longer (up to 4 min) UV illumination. For this purpose, we anticipated that our cell-laden GelMA hydrogel injected intramyocardially is unlikely to provide compatible mechanical support, as shown by Ptaszek et al.

Our study had several limitations, including the need to use immunodeficient mice to enable the transplantation of human cells. Nevertheless, although the SCID mice used in this study have a genetic immune deficiency that affects their B and T cells, their myeloid cell lineage is normal. Indeed, we did not notice dysregulated neutrophil behaviors in the absence of functional B and T immunity compared to other studies of neutrophil recruitment post-MI in immunocompetent mice [50,52]. Also, we previously validated the dependency of graft vascularization on host neutrophils using fully immunocompetent hosts by transplanting mouse autologous endothelial cells and MSCs into syngeneic recipients [47]. Therefore, we have demonstrated that neutrophil participation is critical in both syngeneic (immunocompetent recipients) and xenograft (immunodeficient recipients) cell engraftment models. In both cases, lymphocyte presence in the grafts was minimal. Nevertheless, the difference in the immune system between humans and rodents is well recognized, and thus our preclinical data will need to be further validated in human patients.

Another limitation of the mouse MI model is the difficulty performing more clinically relevant ischemic injuries. For instance, we induced MI by permeant ligation of LAD in mouse hearts, but a cardiac ischemia-reperfusion injury (IRI) would have been clinically more relevant [26]. However, performing IRI in small mouse hearts is challenging due to significant variations and problems of reproducibility. Also, the most likely intramyocardial cell delivery route in the clinics would be through a minimally invasive catheter procedure to penetrate the endocardial side. However, it is extremely difficult to perform this procedure in mice and rats. Lastly, the analyses of cardiac functions in mice are limited to imaging-based measurements (echocardiography and magnetic resonance imaging) for live animals or histological examination for end-point experiments. Future studies in large animals could allow for more precise analysis, including the invasive hemodynamics with PV loops for long-term evaluations.

In summary, we have developed a novel therapeutic cell delivery method for MI treatment. Our approach entails the intramyocardial injection of vascular progenitor cells resuspended in a GelMA precursor solution followed by transmyocardial UV illumination, resulting in an in-situ photocrosslinked hydrogel that effectively retains the cells inside the ischemic myocardial tissue. This approach maintained high viability and cell retention, providing a significant advantage over cells injected in liquid or unmodified ECM gels. The improved cell retention allowed us to investigate the therapeutic effects of the human vascular progenitor cells post-MI. We demonstrated that the beneficial myocardial remodeling and stabilization of cardiac functions post-MI was enabled by the engrafted cells via the engagement and polarization of host pro-regenerative neutrophils through TGFβ signaling. This proof-of-concept study warrants further investigations, including scaling up for large animal tests, integrating a catheter-mediated injectate and a UV light delivery system, and potential modifications to include additional cell types such as iPSC-derived cardiomyocytes [91]. We envision that our research provides an opportunity to generate an off-the-shelf cell-based therapy with the potential to treat acute myocardial infarction.

Reference

1. Reed, G. W., Rossi, J. E. & Cannon, C. P. Acute myocardial infarction. *Lancet* 389, 197–210 (2017).

2. Members, W. G. *et al.* Heart Disease and Stroke Statistics—2012 Update. *Circulation* 125, e2–e220 (2012).

3. Westman, P. C. *et al.* Inflammation as a Driver of Adverse Left Ventricular Remodeling After Acute Myocardial Infarction. *J Am Coll Cardiol* 67, 2050–2060 (2016).

4. Tamargo, J. & López-Sendón, J. Novel therapeutic targets for the treatment of heart failure. *Nat Rev Drug Discov* 10, 536–555 (2011).

5. Stone, G. W. *et al.* Comparison of Angioplasty with Stenting, with or without Abciximab, in Acute Myocardial Infarction. *New Engl J Medicine* 346, 957–966 (2002).

6. Montalescot, G. *et al.* Platelet Glycoprotein IIb/IIIa Inhibition with Coronary Stenting for Acute Myocardial Infarction. *New Engl J Medicine* 344, 1895–1903 (2001).

7. Huang, K., Hu, S. & Cheng, K. A New Era of Cardiac Cell Therapy: Opportunities and Challenges. *Adv Healthc Mater* 8, 1801011 (2019).

8. Katarzyna, R. Adult Stem Cell Therapy for Cardiac Repair in Patients After Acute Myocardial Infarction Leading to Ischemic Heart Failure: An Overview of Evidence from the Recent Clinical Trials. *Curr Cardiol Rev* 13, (2017).

9. Dib, N., Khawaja, H., Varner, S., McCarthy, M. & Campbell, A. Cell Therapy for Cardiovascular Disease: A Comparison of Methods of Delivery. *J Cardiovasc Transl* 4, 177–181 (2011).

10. Pavo, N. *et al.* Cell therapy for human ischemic heart diseases: Critical review and summary of the clinical experiences. *J Mol Cell Cardiol* 75, 12–24 (2014).

11. Popescu, S., Preda, M. B., Marinescu, C. I., Simionescu, M. & Burlacu, A. Dual Stem Cell Therapy Improves the Myocardial Recovery Post-Infarction through Reciprocal Modulation of Cell Functions. *Int J Mol Sci* 22, 5631 (2021).

12. Vagnozzi, R. J. *et al.* An acute immune response underlies the benefit of cardiac stem cell therapy. *Nature* 577, 405–409 (2020).

13. Andreadou, I. *et al.* Immune cells as targets for cardioprotection: new players and novel therapeutic opportunities. *Cardiovasc Res* 115, 1117–1130 (2019).

14. Trajano, L. F. & Smart, N. Immunomodulation for optimal cardiac regeneration: insights from comparative analyses. *Npj Regen Medicine* 6, 8 (2021).

15. Fang, L., Moore, X.-L., Dart, A. M. & Wang, L.-M. Systemic inflammatory response following acute myocardial infarction. *J Geriatric Cardiol Jgc* 12, 305–312 (2015).

16. Pluijmert, N. J., Atsma, D. E. & Quax, P. H. A. Post-ischemic Myocardial Inflammatory Response: A Complex and Dynamic Process Susceptible to Immunomodulatory Therapies. *Frontiers Cardiovasc Medicine* 8, 647785 (2021).

17. He, L., Nguyen, N. B., Ardehali, R. & Zhou, B. Heart Regeneration by Endogenous Stem Cells and Cardiomyocyte Proliferation. *Circulation* 142, 275–291 (2020).

18. Shah, V. K. & Shalia, K. K. Stem Cell Therapy in Acute Myocardial Infarction: A Pot of Gold or Pandora's Box. *Stem Cells Int* 2011, 536758 (2011).

19. Mei, X. & Cheng, K. Recent Development in Therapeutic Cardiac Patches. *Frontiers Cardiovasc Medicine* 7, 610364 (2020).

20. Dimmeler, S., Burchfield, J. & Zeiher, A. M. Cell-Based Therapy of Myocardial Infarction. *Arteriosclerosis Thrombosis Vasc Biology* 28, 208–216 (2008).

21. Zhao, Z.-A. *et al.* Lack of Cardiac Improvement After Cardiosphere-Derived Cell Transplantation in Aging Mouse Hearts. *Circ Res* 123, e21–e31 (2018).

22. Fukushima, S. *et al.* Direct Intramyocardial But Not Intracoronary Injection of Bone Marrow Cells Induces Ventricular Arrhythmias in a Rat Chronic Ischemic Heart Failure Model. *Circulation* 115, 2254–2261 (2007).

23. Sürder, D. *et al.* Intracoronary Injection of Bone Marrow–Derived Mononuclear Cells Early or Late After Acute Myocardial Infarction. *Circulation* 127, 1968–1979 (2013).

24. Szmitko, P. E. *et al.* Endothelial Progenitor Cells. *Circulation* 107, 3093–3100 (2003).

25. Gnecchi, M. *et al.* Paracrine action accounts for marked protection of ischemic heart by Akt-modified mesenchymal stem cells. *Nat Med* 11, 367–368 (2005).

26. Kang, K.-T., Coggins, M., Xiao, C., Rosenzweig, A. & Bischoff, J. Human vasculogenic cells form functional blood vessels and mitigate adverse remodeling after ischemia reperfusion injury in rats. *Angiogenesis* 16, 773–784 (2013).

27. Park, S.-J. *et al.* Dual stem cell therapy synergistically improves cardiac function and vascular regeneration following myocardial infarction. *Nat Commun* 10, 3123 (2019).

28. Hong, X. *et al.* Human endothelial colony-forming cells provide trophic support for pluripotent stem cell-derived cardiomyocytes via distinctively high expression of neuregulin-1. *Angiogenesis* 1–18 (2021) doi:10.1007/s10456-020-09765-3.

29. Li, J. *et al.* All Roads Lead to Rome (the Heart): Cell Retention and Outcomes From Various Delivery Routes of Cell Therapy Products to the Heart. *J Am Hear Assoc Cardiovasc Cerebrovasc Dis* 10, e020402.

30. Teng, C. J., Luo, J., Chiu, R. C. J. & Shum-Tim, D. Massive mechanical loss of microspheres with direct intramyocardial injection in the beating heart: Implications for cellular cardiomyoplasty. *J Thorac Cardiovasc Surg* 132, 628–632 (2006).

31. Terrovitis, J. *et al.* Noninvasive Quantification and Optimization of Acute Cell Retention by In Vivo Positron Emission Tomography After Intramyocardial Cardiac-Derived Stem Cell Delivery. *J Am Coll Cardiol* 54, 1619–1626 (2009).

32. Liu, Z. *et al.* Systemic and local delivery of mesenchymal stem cells for heart renovation: Challenges and innovations. *Eur J Pharmacol* 876, 173049 (2020).

33. Menasché, P. Cell therapy trials for heart regeneration — lessons learned and future directions. *Nat Rev Cardiol* 15, 659–671 (2018).

34. Crisostomo, V. *et al.* Dose-dependent improvement of cardiac function in a swine model of acute myocardial infarction after intracoronary administration of allogeneic heart-derived cells. *Stem Cell Res Ther* 10, 152 (2019).

35. Vrtovec, B. *et al.* Effects of Intracoronary CD34+ Stem Cell Transplantation in Nonischemic Dilated Cardiomyopathy Patients. *Circ Res* 112, 165–173 (2013).

36. Garcia, J. R. *et al.* A Minimally Invasive, Translational Method to Deliver Hydrogels to the Heart Through the Pericardial Space. *Jacc Basic Transl Sci* 2, 601–609 (2017).

37. Cui, X. *et al.* NIPAM-based Microgel Microenvironment Regulates the Therapeutic Function of Cardiac Stromal Cells. *Acs Appl Mater Inter* 10, 37783–37796 (2018).

38. Ichihara, Y. *et al.* Self-assembling peptide hydrogel enables instant epicardial coating of the heart with mesenchymal stromal cells for the treatment of heart failure. *Biomaterials* 154, 12–23 (2018).

39. Tang, J. *et al.* Heart Repair Using Nanogel-Encapsulated Human Cardiac Stem Cells in Mice and Pigs with Myocardial Infarction. *Acs Nano* 11, 9738–9749 (2017).

40. Nichol, J. W. *et al.* Cell-laden microengineered gelatin methacrylate hydrogels. *Biomaterials* 31, 5536–5544 (2010).

41. Chen, Y. *et al.* Functional Human Vascular Network Generated in Photocrosslinkable Gelatin Methacrylate Hydrogels. *Adv Funct Mater* 22, 2027–2039 (2012).

42. Lin, R.-Z., Chen, Y.-C., Moreno-Luna, R., Khademhosseini, A. & Melero-Martin, J. M. Transdermal regulation of vascular network bioengineering using a photopolymerizable methacrylated gelatin hydrogel. *Biomaterials* 34, 6785–6796 (2013).

43. Yue, K. *et al.* Synthesis, properties, and biomedical applications of gelatin methacryloyl (GelMA) hydrogels. *Biomaterials* 73, 254–271 (2015).

44. Robey, T. E., Saiget, M. K., Reinecke, H. & Murry, C. E. Systems approaches to preventing transplanted cell death in cardiac repair. *J Mol Cell Cardiol* 45, 567–581 (2008).

45. Melero-Martin, J. M. *et al.* In vivo vasculogenic potential of human blood-derived endothelial progenitor cells. *Blood* 109, 4761–4768 (2007).

46. Kang, K.-T., Lin, R.-Z., Kuppermann, D., Melero-Martin, J. M. & Bischoff, J. Endothelial colony forming cells and mesenchymal progenitor cells form blood vessels and increase blood flow in ischemic muscle. *Sci Rep-uk* 7, 770 (2017).

47. Lin, R.-Z. *et al.* Host non-inflammatory neutrophils mediate the engraftment of bioengineered vascular networks. *Nat Biomed Eng* 1, 0081 (2017).

48. Lin, R.-Z. *et al.* Human endothelial colony-forming cells serve as trophic mediators for mesenchymal stem cell engraftment via paracrine signaling. *Proc National Acad Sci* 111, 10137–10142 (2014).

49. Silvestre-Roig, C., Braster, Q., Ortega-Gomez, A. & Soehnlein, O. Neutrophils as regulators of cardiovascular inflammation. *Nat Rev Cardiol* 17, 327–340 (2020).

50. Horckmans, M. *et al.* Neutrophils orchestrate post-myocardial infarction healing by polarizing macrophages towards a reparative phenotype. *Eur Heart J* 38, 187–197 (2017).

51. Yang, W. *et al.* Neutrophils promote the development of reparative macrophages mediated by ROS to orchestrate liver repair. *Nat Commun* 10, 1076 (2019).

52. Ma, Y. *et al.* Temporal neutrophil polarization following myocardial infarction. *Cardiovasc Res* 110, 51–61 (2016).

53. Leden, R. E. von, Parker, K. N., Bates, A. A., Noble-Haeusslein, L. J. & Donovan, M. H. The emerging role of neutrophils as modifiers of recovery after traumatic injury to the developing brain. *Exp Neurol* 317, 144–154 (2019).

54. Kolaczkowska, E. & Kubes, P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* 13, 159–175 (2013).

55. Blázquez-Prieto, J., López-Alonso, I., Huidobro, C. & Albaiceta, G. M. The Emerging Role of Neutrophils in Repair after Acute Lung Injury. *Am J Resp Cell Mol* 59, 289–294 (2018).

56. Ohms, M., Möller, S. & Laskay, T. An Attempt to Polarize Human Neutrophils Toward N1 and N2 Phenotypes in vitro. *Front Immunol* 11, 532 (2020).

57. Voskuhl, R. A new cell subtype that confers neuroprotection. *Nat Immunol* 21, 1477–1478 (2020).

58. Sas, A. R. *et al.* A new neutrophil subset promotes CNS neuron survival and axon regeneration. *Nat Immunol* 21, 1496–1505 (2020).

59. Hou, Y. *et al.* N2 neutrophils may participate in spontaneous recovery after transient cerebral ischemia by inhibiting ischemic neuron injury in rats. *Int Immunopharmacol* 77, 105970 (2019).

60. Peiseler, M. & Kubes, P. More friend than foe: the emerging role of neutrophils in tissue repair. *J Clin Invest* 129, 2629–2639 (2019).

61. Cai, B. *et al.* N2-Polarized Neutrophils Guide Bone Mesenchymal Stem Cell Recruitment and Initiate Bone Regeneration: A Missing Piece of the Bone Regeneration Puzzle. *Adv Sci* 8, 2100584 (2021).

62. Wang, J. Neutrophils in tissue injury and repair. *Cell Tissue Res* 371, 531–539 (2018).

63. Dick, S. A. & Epelman, S. Chronic Heart Failure and Inflammation. *Circ Res* 119, 159–176 (2016).

64. Frangogiannis, N. G. The Immune System and the Remodeling Infarcted Heart. *J Cardiovasc Pharm* 63, 185–195 (2014).

65. Cochain, C. *et al.* The Chemokine Decoy Receptor D6 Prevents Excessive Inflammation and Adverse Ventricular Remodeling After Myocardial Infarction. *Arteriosclerosis Thrombosis Vasc Biology* 32, 2206–2213 (2012).

66. Halade, G. V. & Lee, D. H. Inflammation and resolution signaling in cardiac repair and heart failure. *Ebiomedicine* 79, 103992 (2022).

67. Halade, G. V., Norris, P. C., Kain, V., Serhan, C. N. & Ingle, K. A. Splenic leukocytes define the resolution of inflammation in heart failure. *Sci Signal* 11, (2018).

68. Fridlender, Z. G. *et al.* Polarization of Tumor-Associated Neutrophil Phenotype by TGF-β: "N1" versus "N2" TAN. *Cancer Cell* 16, 183–194 (2009).

69. Ng, L. G., Ostuni, R. & Hidalgo, A. Heterogeneity of neutrophils. *Nat Rev Immunol* 19, 255–265 (2019).

70. Vafadarnejad, E. *et al.* Dynamics of Cardiac Neutrophil Diversity in Murine Myocardial Infarction. *Circ Res* 127, e232–e249 (2020).

71. Calcagno, D. M. *et al.* SiglecF(HI) Marks Late-Stage Neutrophils of the Infarcted Heart: A Single-Cell Transcriptomic Analysis of Neutrophil Diversification. *J Am Hear Assoc Cardiovasc Cerebrovasc Dis* 10, e019019 (2021).

72. Lin, R.-Z. & Melero-Martin, J. M. Fibroblast growth factor-2 facilitates rapid anastomosis formation between bioengineered human vascular networks and living vasculature. *Methods* 56, 440–451 (2012).

73. Melero-Martin, J. M. *et al.* Engineering Robust and Functional Vascular Networks In Vivo With Human Adult and Cord Blood–Derived Progenitor Cells. *Circ Res* 103, 194–202 (2008).

74. Zhao, R., Su, Z., Wu, J. & Ji, H.-L. Serious adverse events of cell therapy for respiratory diseases: a systematic review and meta-analysis. *Oncotarget* 8, 30511–30523 (2017).

75. Kindi, A. A., Ge, Y., Shum-Tim, D. & Chiu, R. C.-J. Cellular cardiomyoplasty: routes of cell delivery and retention. *Front Biosci* 13, 2421 (2008).

76. Vrtovec, B. *et al.* Comparison of Transendocardial and Intracoronary CD34+ Cell Transplantation in Patients With Nonischemic Dilated Cardiomyopathy. *Circulation* 128, S42–S49 (2013).

77. Hou, D. *et al.* Radiolabeled Cell Distribution After Intramyocardial, Intracoronary, and Interstitial Retrograde Coronary Venous Delivery. *Circulation* 112, I150-6 (2005).

78. Akker, F. van den *et al.* Intramyocardial stem cell injection: go(ne) with the flow. *Eur Heart J* 38, 184–186 (2017).

79. Suncion, V. Y. *et al.* Does Transendocardial Injection of Mesenchymal Stem Cells Improve Myocardial Function Locally or Globally? *Circ Res* 114, 1292–1301 (2014).

80. Sherman, W., Martens, T. P., Viles-Gonzalez, J. F. & Siminiak, T. Catheter-based delivery of cells to the heart. *Nat Clin Pract Card* 3, S57–S64 (2006).

81. Abdelwahid, E. *et al.* Stem cell death and survival in heart regeneration and repair. *Apoptosis* 21, 252–268 (2016).

82. Li, Z., Hu, S. & Cheng, K. Chemical Engineering of Cell Therapy for Heart Diseases. *Accounts Chem Res* 52, 1687–1696 (2019).

83. Cheng, K. *et al.* Functional performance of human cardiosphere-derived cells delivered in an in situ polymerizable hyaluronan-gelatin hydrogel. *Biomaterials* 33, 5317–5324 (2012).

84. Chuang, C.-H., Lin, R.-Z., Melero-Martin, J. M. & Chen, Y.-C. Comparison of covalently and physically cross-linked collagen hydrogels on mediating vascular network formation for engineering adipose tissue. *Artif Cells Nanomed Biotechnology* 46, 1–14 (2018).

85. Kuo, K.-C. *et al.* Bioengineering vascularized tissue constructs using an injectable cell-laden enzymatically crosslinked collagen hydrogel derived from dermal extracellular matrix. *Acta Biomater* 27, 151–166 (2015).

86. Chuang, C.-H. *et al.* Enzymatic regulation of functional vascular networks using gelatin hydrogels. *Acta Biomater* 19, 85–99 (2015).

87. Sharifi, S., Sharifi, H., Akbari, A. & Chodosh, J. Systematic optimization of visible light-induced crosslinking conditions of gelatin methacryloyl (GelMA). *Sci Rep-uk* 11, 23276 (2021).

88. Noshadi, I. *et al.* In vitro and in vivo analysis of visible light crosslinkable gelatin methacryloyl (GelMA) hydrogels. *Biomater Sci-uk* 5, 2093–2105 (2017).

89. Roche, E. T. *et al.* A light-reflecting balloon catheter for atraumatic tissue defect repair. *Sci Transl Med* 7, 306ra149 (2015).

90. Ptaszek, L. M. *et al.* Gelatin Methacryloyl Bioadhesive Improves Survival and Reduces Scar Burden in a Mouse Model of Myocardial Infarction. *J Am Hear Assoc Cardiovasc Cerebrovasc Dis* 9, e014199 (2020).

91. Chong, J. J. H. *et al.* Human embryonic-stem-cell-derived cardiomyocytes regenerate non-human primate hearts. *Nature* 510, 273–277 (2014).

Figure 1. Enhanced intramyocardial retention of human vascular progenitor cells delivered in a photocrosslinked GelMA hydrogel. (A) Schematic depicting the photocrosslinking of GelMA from a precursor solution (GelMA-S) to a hydrogel matrix (GelMA-H) under UV irradiation. (B) Schematic depicting the intramyocardial cell injection. MI was surgically induced by LAD ligation. Human vascular progenitor cells (ECFCs + MSCs) resuspended in GelMA precursor solution were injected into three locations in the infarct area. UV light was immediately applied on top of the infarcted area to polymerize the cell-laden hydrogel (referred to as GelMA-H group). UV irradiation was omitted for the control group to maintain cells injected in GelMA precursor solution (GelMA-S group). (C) Viable cell retention was measured by bioluminescence imaging of human ECFCs expressing a luciferase reporter. (D) Total bioluminescent signals in each animal's chest region were measured on 3, 24, 48, and 72 h post-MI and compared to the baseline (i.e., cells delivered to arrested mouse hearts). (E) Retention of human ECFCs or MSCs was quantified by flow cytometric analysis on 48 h. Data correspond to cell numbers from each harvested MI heart. Data are mean ± s.d.; n = 3 mice per group (indicated by individual dots). *P < 0.05 between GelMA-H and GelMA-S. Statistical methods: two-way ANOVA followed by Bonferroni's post-test analysis (D) and unpaired two-tailed Student's t-tests (E).

Figure 2. Human vascular progenitor cells delivered in GelMA hydrogel improved cardiac remodeling. (A) Diagram of experimental schedule. (B-D) Echocardiography analysis for cardiac function. Echocardiography was performed on anesthetized mice at 3 days and 1, 2, and 4 weeks post-MI following cell delivery in GelMA-H or GelMA-S (n=6 for each group). Mice received LAD ligation, but no cell injection (MI-untreated) served as control (n=6). Baseline data were obtained from mice within 1h before performing LAD ligation. (B) Left ventricular fractional shortening

(LVFS). (C) Left ventricular ejection fraction (LVEF). (D) Left ventricular anterior wall thickness (LVAW). (E) Representative heart horizontal panoramic views (upper) and microscopic images (lower) of myocardial sections stained with H&E. Scale bars, 100 μm. (F) The LV wall thickness in the infarct areas quantified from H&E sections. (G) Histological analysis of the infarcted myocardium on day 7 revealed the presence of perfused human blood vessels only when cells were delivered in GelMA-H group, but not in GelMA-S. Human microvessels were identified by immunostaining for human-specific vimentin (hVIM) and UEA-1 lectin binding. Scale bars, 100 μm. (H) Total perfused human microvessel density quantified in the infarcted myocardium. (I) Representative heart sections stained with Masson's trichrome. Scale bars, 100 μm. (J) The percentage of fibrotic areas in the total LV cross-sections quantified by Masson's trichrome staining. Statistical methods: ANOVA followed by Bonferroni's post-test analysis (B-D) and unpaired two-tailed Student's t-tests (F, H, J). Data present mean $\pm$ s.d. *$P < 0.1$, **$P < 0.05$, ***$P < 0.001$.

Figure 3. Human vascular progenitor cell retention promotes the recruitment of neutrophils with a reparative phenotype. The infarcted myocardial tissue was harvested on day 2 post-MI. Tissues were digested into single-cell suspensions for flow cytometric analysis. (A) Flow cytometry gating strategy to identify recruited myeloid cells. Representative flow cytometric analyses of ischemic myocardium injected with ECFC+MSC in GelMA-S or GelMA-H. MΦ, macrophages; N, neutrophils; mN, mature neutrophils; N0, naïve neutrophils; NR, pro-regenerative neutrophils; NI, pro-inflammatory neutrophils. (B) Quantitative cytometric analyses of murine CD45+ hematopoietic cells, Ly6G$^-$F4/80$^+$ macrophages, and total Ly6G$^+$F4/80$^-$ neutrophils obtained from the ischemic myocardium. The myocardium from uninjured hearts served as controls. (C) Recruited Ly6G$^+$F4/80$^-$ neutrophils were classified into three subpopulations: N0 (CXCR2$^-$), NI (CXCR2$^+$CD206$^-$), and NR (CXCR2$^+$CD206$^+$). (B,C) Data are mean ± s.d.; n = 3 per group; p values between GelMA-S and GelMA-H are listed. (D) Proportions of the three neutrophil subpopulations in each group.

Figure 4. Human vascular progenitor cell crosstalk upregulates NR polarization factor TGFβ1. (A) Conditioned media (CM) were collected from ECFC+MSC coculture in GelMA hydrogel over 7 days in vitro (CM^{CO}). CMs obtained from ECFC or MSC monoculture (CM^{ECFC} or CM^{MSC}) and mixed in 1:1 ratio (termed CM^{MIX}) served as a control. (B) Mouse naïve neutrophils isolated from bone marrow were cultured in basal medium or indicated conditional media for 24 h. Gene expressions of Il4, Vegfa, and Arg1 expressions in mouse naïve neutrophils were measured by the quantitative PCR analyses and compared to the levels in basal medium. Data are mean ± s.d.; n = 3 per group. *$p < 0.05$ between Basal and CM^{co} groups. (C) The secretions of cytokines were measured by proteomic arrays. Selected cytokines are marked with colored outlines. (D) Quantification of cytokine levels was carried out by densitometry. (E) ECFC+MSC were cocultured in GelMA hydrogel for 48 h and separated by magnetic-activated cell sorting (MACS) for gene expression analyses. ECFC or MSC monoculture served as controls. (F) Quantitative PCR analyses of TGFβ1 expression in ECFCs sorted from ECFC+MSC coculture (labeled EC*) or from ECFC monoculture (EC). The same labeling applied to MSC (monoculture) and MSC* (sorted from coculture). DAPT was added to ECFC+MSC coculture to block Notch-mediated perivascular interaction. Data are mean ± s.d.; n = 3 per group; *$p < 0.01$ between indicated groups.

Figure 5. Neutrophil depletion abrogated the improvement in cardiac remodeling of human vascular cell delivery. (A) Experimental schedule: Anti-Gr-1 or IgG control antibody (200 µg each) was given via an intraperitoneal injection every 2 days. ECFC+MSC were delivered in GelMA-H into the ischemic myocardium at day 0. Cardiac functions were measured by echocardiography within 1 h before LAD ligation to obtain the baseline data for each animal. The effect of anti-Gr-1 depletion was monitored on days 3 and 7 post-MI. (B) Representative flow cytometry analyses of circulating leukocytes from the blood of anti-Gr-1-treated or IgG-treated mice on day 2. (C) Quantitative cytometric analyses of circulating leukocytes were obtained from anti-Gr-1-treated or IgG-treated mice. Data are mean ± s.d.; n = 3 per group; *P* values between IgG and Gr1 treatments are listed. (D-F) Echocardiography analysis for cardiac function. Echocardiography was performed on anesthetized mice at days 3 and 7 post-MI following cell delivery in GelMA-H (n=3 for IgG-treated and n=4 for Gr1-treated). Baseline data were obtained from mice within 1h before performing LAD ligation. (D) Left ventricular fractional shortening (LVFS). (E) Left ventricular ejection fraction (LVEF). (F) Left ventricular anterior wall thickness (LVAW). Data are mean ± s.d.; n=3 for IgG-treated and n=4 for Gr1-treated. **P* < 0.1, ***P* < 0.05.