**TITLE**

2    **Cyclin D1·Cdk4 regulates neuronal activity through phosphorylation of GABA$_A$ receptors**

4    **AUTHORS**

Neus Pedraza*[#1], Ma Ventura Monserrat[#1], Francisco Ferrezuelo[1], Jordi Torres-Rosell[1], Neus

6    Colomina[1], David Soto[2], Federico Miguez-Cabello[2], José A. Esteban[3], Joaquim Egea[4], Eloi Garí*[1].

[1]Cell Cycle, Department of Basic Medical Sciences, Institut de Recerca Biomèdica de Lleida

8    (IRBLLEIDA), University of Lleida, Lleida, Spain.

[2]Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Hospital Clínic, Universitat

10    de Barcelona, Barcelona, Spain. Laboratori de Neurofisiologia, Departament de Biomedicina,

Facultat de Medicina i Ciències de la Salut, Institut de Neurociències, Universitat de Barcelona,

12    Barcelona, Spain.

[3]Department of Molecular Neurobiology, Centro de Biología Molecular 'Severo Ochoa',

14    Consejo Superior de Investigaciones Científicas (CSIC)/Universidad Autónoma de Madrid

(UAM), Spain.

16    [4]Molecular and Developmental Neurobiology, Dept. Ciències Mèdiques Bàsiques, Facultat de

Medicina, Universitat de Lleida/IRBLLEIDA, Rovira Roure 80, 25198, Lleida, Spain.

18    *Corresponding authors

[#]Co-authors

**ABSTRACT**

Cyclin D1 (Ccnd1)·Cdk4 complexes drive cell cycle progression through phosphorylation of pRb. Interestingly, Ccnd1 moves to the cytoplasm at the onset of differentiation in neuronal

precursors. However, the cytoplasmic functions and targets of Ccnd1 in post-mitotic neurons are unknown. Here we identify the α4 subunit of gamma-aminobutyric acid (GABA) type A

receptors (GABA$_A$Rs) as an interactor and target of Ccnd1·Cdk4. Ccnd1 binds to an intracellular loop in α4 and, together with Cdk4, phosphorylates the α4 subunit at threonine 423 and serine

431. These modifications increase the activity of α4-containing GABA$_A$Rs, measured in whole-cell patch-clamp recordings, and upregulate its surface levels. In agreement with this role of

Ccnd1·Cdk4 in neuronal signaling, inhibition of Cdk4 decreases synaptic and extrasynaptic currents in the hippocampus of newborn rats. Moreover, *CCND1* knockout mice display an

altered pattern of dendritic spines, according to α4 functions in synaptic pruning. Overall, our findings molecularly link Ccnd1·Cdk4 to GABA$_A$Rs activity in the central nervous system and

highlight a novel role for this G1 cyclin in neuronal signaling.

Keywords: Cdk4/ Cyclin D1/ GABA signaling/ gamma-aminobutyric acid type A receptor subunit alpha4

**INTRODUCTION**

Cyclin D1 (Ccnd1) and its catalytic counterparts, cyclin-dependent kinase 4 (Cdk4) and Cdk6, are main regulators of cell proliferation by promoting the G1-S phase transition of the cell

cycle (reviewed in[1]). Under favorable conditions, mitogenic signals induce the expression of Ccnd1, the formation of Ccnd1·Cdk4/6 complexes and their localization to the nucleus. There,

they phosphorylate the Retinoblastoma repressor protein (pRB), which eventually leads to its inhibition and in turn releases the activity of the E2F transcription factor. This induces the

expression of genes required to go through the next steps of the cell cycle. This pRB-dependent control of cell proliferation is considered the "canonical" function of Ccnd1 and it

has been reported as very relevant in tumor proliferation. Accordingly, Ccnd1 is classified as an oncogene frequently amplified in many types of neoplasia, and different cancer treatments

with Cdk4/6 inhibitors such as Palbociclib have been approved to block this function. However, Ccnd1 also operates through several other non-canonical pathways including some pRB- and

Cdk-independent ones to regulate different cellular processes (reviewed in[2]). Thus, Ccnd1 promotes cell detachment, migration and invasion of normal and tumor cells independently of

pRB status[3]'[4]'[5] by interacting with cytoplasmic and membranous targets such as filamin A, pacsin, RalGTPases and paxillin[6]'[7]'[8]'[9]. For example, Ccnd1·Cdk4 promotes cell invasion

and metastasis through the phosphorylation of paxillin and the activation of the Rac1 pathway in membrane ruffles[9]. Moreover, the expression of a membrane-associated form of Ccnd1

harboring the pharnesylation signal of K-Ras (Ccnd1-CAAX) maximizes the ability of tumor cells to invade and metastasize[10]. Besides cell invasion, membrane-targeted Ccnd1 also

influences cell-signaling responses[11].

The first hint for a role of Ccnd1 in the nervous system came from the analysis of *CCND1*

62  knockout (KO) mice[12], which present certain behaviors (altered leg-clasping reflex) indicative of neurological abnormalities. Since then, several studies have substantiated the involvement

64  of Ccnd1 in neuronal functioning. In neural stem cells, Ccnd1·Cdk4 has been involved in

regulating the length of G1, which is thought to influence the decision of cells to proliferate or

66    differentiate[13]'[14]'[15]. Also, under pathological conditions, nuclear Ccnd1 reactivation/

upregulation in post-mitotic neurons has been observed to promote apoptosis (reviewed

68    in[2]). Ccnd1 is located in the cytoplasm of differentiating post-mitotic cortical neurons and in

neuroblastoma cell lines, although this has been proposed as a mechanism to prevent

70    apoptosis[16] or to promote cell cycle withdrawal[17], respectively. Nevertheless, Ccnd1 could

be exerting an active role in the cytoplasm. Interestingly, Ccnd1 is necessary for NGF-induced

72    neuritogenesis in PC12 pheochromocytoma cell line[18]. Also cytoplasmic expression of

Ccnd1·Cdk4 has been observed in the hippocampus during development[19] where it has been

74    involved in neuronal plasticity[20]. But despite these works, the role and target/s of Ccnd1 in

the nervous system and in particular its role in the cytoplasm of the neurons remains poorly

76    understood.

$\gamma$-aminobutyric acid (GABA) is the major inhibitor neurotransmitter in the central nervous

78    system. Its actions are mediated through different types of GABA receptors: GABA A receptors

(GABA$_A$Rs), which are ligand-gated chloride ion channels, and GABA B receptors (metabotropic

80    G-protein-coupled). Deficits in GABA$_A$R function are increasingly being involved in different

pathologies, such as anxiety, cognitive deficits, depression, epilepsy, schizophrenia and

82    substance abuse[21]. These receptors are also clinically relevant drug targets for anesthetic,

anticonvulsant, anxiolytic, or sedative-hypnotic agents. GABA$_A$Rs mediate both phasic and

84    tonic inhibition: synaptic GABA$_A$Rs are activated transiently by the GABA released from

presynaptic vesicles, whereas extrasynaptic GABA$_A$Rs are activated by low GABA

86    concentrations in the extracellular space and thus mediate tonic inhibition. The tonic inhibitory

currents control neuronal excitability and the strength of synaptic transmission[22].

88    GABA$_A$Rs are heteropentamers that belong to the ligand-gated ion channel superfamily. A total

of 19 subunits (as well as splice variants) have been cloned and grouped into eight subunit

90    classes: $\alpha$ (1-6), $\beta$ (1-3), $\gamma$ (1-3), $\delta$, $\epsilon$, $\pi$, $\theta$ and $\rho$ (1-3). GABA$_A$Rs are usually formed by two $\alpha$, two

β, and one γ or one δ subunits. Each subunit contains a large amino-terminal extracellular

domain and four transmembrane (TM) domains. Between TM3 and TM4 there is a large

intracellular loop (ICL) where regulation by phosphorylation and protein interactions primarily

occurs. The subunit composition dictates its subcellular location and function.

GABA$_A$Rs containing the α4 subunit are expressed in the hippocampus, cortex and thalamus,

and mediate extrasynaptic inhibition in thalamus and dentate gyrus[23]. At the onset of

puberty, α4-β-δ GABA$_A$R expression increases perisynaptically at excitatory synapses in the

hippocampus, contributing to anxiety, synaptic pruning and learning deficits in

mice[24]'[25]'[26]'[27]'[28]. In fact, in the α4-KO mice synaptic pruning is prevented, and

synaptic plasticity and learning abilities are restored[23]'[29]'[30]. In the adult hippocampal

neurogenesis, α4-containing GABA$_A$Rs are involved in proliferation, migration and dendritic

development[31]. Mutations in the GABAergic receptor subunit 4 gene *GABRA4*[32]'[33]'[34]

and reduced protein and mRNA levels of α4[35]'[36] are reported in patients of autism

spectrum disorder. Furthermore, the α4-KO mice show a phenotype compatible with high-

functioning autism[37].

Here we report that Ccnd1 interacts with the α4 subunit of GABA$_A$Rs. Ccnd1·Cdk4 complex

phosphorylates α4 in the intracellular loop between TM3 and TM4 and enhances the efficacy

of α4-containing GABA$_A$Rs, affecting the pattern of dendritic spines. We propose a novel role

for cytoplasmic Ccnd1·Cdk4/6 in regulating α4-containing GABA$_A$Rs in the central nervous

system.

**RESULTS**

**The α4 subunit of GABA$_A$ receptors interacts with cyclin D1**

To investigate the role of Ccnd1 in the nervous system, we performed a yeast two-hybrid

screening using Ccnd1 as a bait and an adult mouse brain library as a prey[38]. We found that

Ccnd1 interacts with the C-terminal half of the α4 GABA$_A$R subunit (Gabra4). We obtained

three independent clones spanning amino acids 354-552, 385-552 and 386-552 (Fig 1A). This

118   region of α4 contains the intracellular loop (ICL) between the TM3 and TM4 transmembrane

domains, which is phosphorylated by protein kinase C (PKC,[39]). To confirm the interaction

120   between the α4 subunit and Ccnd1 we carried out immunoprecipitation (IP) experiments with

FLAG-tagged Ccnd1 and the HA-tagged C-terminal region of α4 (354-552aa) expressed in HEK-

122   293T cells. We observed specific co-IP of the C-terminal domain of α4 in the Ccnd1 IP (Fig 1B).

To test whether there is a direct interaction between Ccnd1 and the α4 subunit, we performed

124   *in vitro* GST-pull-down assays. Fusions of GST-α4 or GST-C-terminal α4 (354-552aa) purified

from *E.coli* were mixed with FLAG-Ccnd1 produced by *in vitro* transcription and translation.

126   Ccnd1 was co-precipitated when GST-α4 or GST-C-terminal α4 were pulled down, but not with

GST alone (Fig 1C). To analyze the endogenous interaction between Ccnd1 and the α4 GABA$_A$R

128   subunit, we immunoprecipitated Ccnd1 from the hippocampus of adult mice, since co-

expression of Ccnd1 and α4 mRNAs in the CA1, CA3 and dentate gyrus regions of the

130   hippocampus is observed in the Allen mouse brain atlas (www.brain-map.org). As shown in

figure 1D, α4 subunit of GABA$_A$R co-immunoprecipitated with endogenous Ccnd1, in a specific

132   manner. Overall, our data indicate that there is a specific and direct interaction between

Ccnd1 and the α4 subunit of GABA$_A$ receptors.

**Figure 1**

*In vitro* interaction

**Figure 1. Ccnd1 interacts with the α4 subunit of GABA$_A$ receptors.** *A) Scheme representing the mouse*

136    *α4 subunit of GABA$_A$Rs (552aa). N terminal (Nt) and C terminal (Ct), intracellular and extracellular*

*domains and the four transmembrane domains (TM1-4) are shown. In a yeast two-hybrid screening,*

138    *Ccnd1 was found to interact with three independent clones of the C-terminal region of the α4 subunit (in*

*yellow), spanning amino acids 354-552, 385-552, 386-552 (the starting amino acids are shown as stars).*

140    *The three clones contained the intracellular loop (ICL) between TM3 and TM4. Putative phosphorylation*

*sites for Cdk (SP or TP) in the ICL are shown in blue. B) HEK-293T cells were transfected with the C*

142    *terminus of α4 subunit (354-552aa, tagged with HA: HA-Ct-α4) and Ccnd1 (tagged with Flag: Flag-*

*Ccnd1). Flag-Ccnd1 was precipitated using αFlag agarose beads (Sigma). Input (IN) and*

144    *immunoprecipitation (IP) samples were analyzed by western blot using anti-HA and anti-Flag antibodies,*

*to detect co-IP of the C terminus of α4 with Ccnd1. C) In vitro translated FLAG-Ccnd1 was incubated with*

146    *recombinant GST or GST fusions of α4 full length (GST-α4) or the C terminus of α4 (GST-Ct-α4) purified*

*from E. coli. Input (IN) and GST pull down (PD) samples were analyzed by western blot using anti-Flag*

148    *antibody to detect Ccnd1 and anti-GST antibody to detect α4. Intact protein fusions are indicated. D)*

*Immunoprecipitation of endogenous Ccnd1 from adult mouse hippocampal extracts was performed*

 *using a rabbit polyclonal antibody against Ccnd1. As a control, a rabbit polyclonal antibody against the*

*Flag epitope (FL) was used. Input (IN) and IP samples were analyzed by western blot to detect Ccnd1 and*

152 *α4 subunit.*

154 **The α4 subunit of GABA$_A$Rs is phosphorylated at T423 and S431 by Ccnd1·Cdk4 complex**

As Ccnd1 is the regulatory subunit of Cdk4/6 kinases, we analyzed the sequence of α4 for

156 putative phosphorylation sites by Cdks. We found five S/TP sites in the sequence of the

intracellular loop (Fig 1A). In an *in vitro* kinase assay, Ccnd1·Cdk4 complex phosphorylated

158 recombinant GST-α4 and this phosphorylation was prevented by inhibiting the complex with

the specific Cdk4/6 inhibitor Palbociclib (Fig 2A). Besides, another subunit of the GABA$_A$Rs, β3,

160 which lacks S/TP sites, was not phosphorylated by the Ccnd1·Cdk4 complex (Fig 2B). In order to

know which residues of α4 were phosphorylated by Ccnd1·Cdk4, we mutated the five

162 intracellular putative Cdk phosphorylation sites to non-phosphorylatable residues (alanine)

generating three different mutant alleles: T423A and S431A (α4$^{T423AS431A}$); S456A and S458A

164 (α4$^{S456AS458A}$); and T500A (α4$^{T500A}$). Neither α4$^{S456AS458A}$ nor α4$^{T500A}$ mutants showed a decrease in

phosphorylation by Ccnd1·Cdk4 (Fig 2C and 2D). However, the non-phosphorylatable mutant

166 of T423 and S431 (α4$^{T423AS431A}$: AA) significantly reduced phosphorylation by Ccnd1·Cdk4 *in*

*vitro*, either with the full-length or the C-terminus construct of α4 (Fig 2C-F). Moreover, we

168 analyzed by mass spectrometry phosphopeptides from the C-terminus of α4 subunit

phosphorylated by Ccnd1·Cdk4, and we found that both T423 and S431 were phosphorylated

170 *in vitro* (S1 Table). Altogether, our results show that Ccnd1·Cdk4 complex is able to

phosphorylate α4 subunit at T423 and S431.

172 To investigate if the α4 subunit is also phosphorylated by Ccnd1·Cdk *in vivo*, we

immunoprecipitated Cdk substrates with an anti-phospho-Threonine-Proline (phosphoTP)

174 antibody and the presence of phosphorylated α4 was assessed by western blot. We transiently

transfected HEK-293T cells with α4, β3 subunits and Ccnd1 or Ccnd1$^{K112E}$, a mutant of Ccnd1

176     that does not form an active complex with Cdk[40]. For the experiments where we co-expressed Ccnd1 or Ccnd1$^{K112E}$, we used a variant that contains the CAAX motif of K-ras to

178     maintain Ccnd1 attached to the cell membrane[10] to improve the phosphorylation of membrane targets. In the phosphoTP IP, we observed an increase in the phosphorylated α4

180     subunit in the presence of Ccnd1-CAAX relative to the presence of Ccnd1$^{K112E}$-CAAX (Fig 2G and 2H). Moreover, when we used the non-phosphorylatable α4 mutant (α4$^{T423AS431A}$: AA), we did

182     not observe higher levels of phosphorylated α4 in the presence of Ccnd1 (Fig 2G and 2H). This data suggests that Ccnd1·Cdk4 can phosphorylate α4 *in vivo* at least at T423.

# Figure 2

**184**

***Figure 2. Ccnd1·Cdk4 phosphorylates α4 at T423 and S431 residues.*** *A) In vitro radioactive kinase assay*

**186**    *using purified recombinant GST-α4 fusion and Ccnd1·Cdk4 (Sigma). Cdk4/6 inhibitor Palbociclib was*

*added at 1 µM concentration. Coomassie blue staining is shown as loading control (Input). B)*

**188**    *Recombinant GST-α4 and GST-β3 fusions were tested for Ccnd1·Cdk4 kinase activity as in A. C) Kinase*

202     **Ccnd1·Cdk4 promotes the activation of α4-containing GABA$_A$ receptors**

Because Ccnd1·Cdk4 phosphorylates the α4 subunit in the intracellular loop, we reasoned that

204     it may also affect the performance of α4-containing GABA$_A$Rs. To study the electrophysiology

of GABA$_A$Rs we carried out whole cell patch-clamp recordings of HEK293 tsA201 cells

206     transiently expressing α4 and β3 subunits (Fig 3A), since functional GABA$_A$ receptors can be

formed in HEK-293T cells by co-expression of these subunits[39]. To measure the behavior of

208     the α4β3 GABA$_A$Rs, 1 μM GABA was applied for 5 seconds every 2 min in these cells (Fig 3A and

S1). Receptor-mediated current amplitude decreased (rundown) over time (Fig 3A and S1A):

210     after 12 minutes of recording, GABA-mediated current was 60 ± 5 % (n=7) of the initial

response (Fig 3A and S1D). Co-expression of Ccnd1-CAAX with α4β3 receptors partially

212     prevented their rundown in a significant manner (Fig 3A and S1B). At 12 minutes after the start

of the experiment, the GABA-mediated current amplitude, in the presence of Ccnd1-CAAX, was

214     80 ± 9 % (n=5) of the initial GABA-mediated response (Fig 3A and S1D). To confirm that

phosphorylation of T423 S431 in α4 is important for this effect, we co-transfected a

216  phosphomimetic α4$^{T423ES431E}$ (α4$^{EE}$) mutant with β3 and observed that it completely obliterates

the rundown of GABA$_A$Rs (Fig 3A and S1C). Thus, our data suggest that Ccnd1 can enhance the

218  efficacy of α4-containing GABA$_A$Rs by modulating the activity of the receptor through the

phosphorylation of α4.

222   ***Figure 3. Ccnd1·Cdk4/6 modulates α4-containing GABA$_A$ receptor activity. A) Ccnd1 or the***

***phosphomimetic allele of α4 (α4$^{EE}$) decreases GABA$_A$ receptor rundown.*** *Whole-cell patch clamp*

224   *configuration for GABA currents recording in HEK-tsA201 cells transfected with α4 and β3 subunits of*

*GABA$_A$Rs. Currents were evoked by rapid application of 1 μM GABA and holding potential was held at*

226   *−60mV. Peak current values at different times of GABA stimulations were analyzed in cells under 3*

*conditions: 1) α4 wt (α4β3, grey circles), 2) α4 wt co-transfected with Ccnd1-CAAX (α4β3+Ccnd1, red*

228   *circles) and 3) α4$^{T423E\ S431E}$ phosphomimetic allele (α4$^{EE}$β3, blue circles). \*p<0.05, α4β3 vs. α4β3+Ccnd1;*

*#p<0.05 α4β3 vs. α4$^{EE}$β3; Unpaired student's t-Test.* ***B and C) Cdk4/6 inhibition by Palbociclib decreases***

230   ***GABA tonic currents and mIPSC amplitude in rat hippocampal slices.*** *GABA currents recordings in CA1*

*neurons: hippocampal slices from newborn rats were cultured in vitro, treated for 2 h with DMSO or*

232   *Palbociclib 2.5 μM, and tonic (extrasynaptic) or mIPSCs (synaptic currents) were measured. B) GABA*

*inhibitory postsynaptic currents were evoked by stimulation of Schaffer collaterals and the tonic GABA$_A$*

234   *receptor-mediated current was measured as the outward shift in holding current following application of*

*the GABA$_A$R antagonist picrotoxin. CA1 pyramidal neurons were recorded under whole-cell voltage-*

236   *clamp at -60 mV. Quantification of the shift in holding current for DMSO- and Palbociclib-treated cells*

*(n=10), represented in a scatter dot plot showing mean values ± S.E.M (\*p<0.05, Wilcoxon-Mann-*

238   *Whitney test). The shift in holding currents, corresponding to tonic currents, is reduced in Palbociclib-*

*treated slices compared to vehicle. C) To measure miniature inhibitory postsynaptic currents (mIPSC), 1*

240   *μM tetrodotoxin was added to block action potentials. Scatter dot plot showing mIPSC amplitude*

*quantifications of DMSO- (n=9) and Palbociclib-treated cells (n=8). Bars show mean values ± S.E.M.*

242   *\*p<0.05, Wilcoxon-Mann-Whitney test.* ***D-F) CCND1 KO increases the maturity of dendritic spines in***

***cortical neurons****. D) Schematic representation of the experimental design for dendritic spine analysis of*

244   *cortical neurons at 20 DIV. Cultured cortical neurons from WT and CCND1 KO mice were transfected at 3*

*DIV with GFP and fixed at 20 DIV. Changes in spine density and morphology were assessed by analyzing*

246   *GFP-positive neurons with NeuronJ (see Materials and Methods). E) Confocal images of representative*

*neurites from WT and CCND1 KO neurons are shown. Scale bar: 10 μm. F) Quantification of the number*

248   *of mushroom spines relative to the number of total spines from WT (n= 59 neurons) and CCND1 KO (n=*

*62) neurons (4 mice/group). Bars show mean values ± S.E.M. \*\*p<0.01, Unpaired student's t-test.*

The α4 subunit is a component of extra-synaptic, high-affinity GABA$_A$Rs, which mediate tonic

252 inhibition. To assess the importance of Ccnd·Cdk4/6 on GABA$_A$Rs in a physiological context, we

measured tonic currents in hippocampal slices of newborn rats. Organotypic hippocampal slice

254 cultures were treated with DMSO (vehicle) or the Cdk4/6 specific inhibitor Palbociclib at 2.5

μM for 2 h (Fig 3B and S2A). GABA inhibitory post synaptic currents were evoked with single

256 voltage pulses in Schaffer collaterals, and tonic currents were recorded from CA1 pyramidal

neurons under whole-cell voltage-clamp configuration. The application of the antagonist

258 picrotoxin (100 μM) causes a decrease in the operative extra-synaptic GABA$_A$ channels, which

can be measured as an outward shift of the holding current[41] (Fig S2B). This shift

260 corresponds to the GABA$_A$R tonic currents (Fig 3B). While the holding current in DMSO-treated

cells was shifted by 21 ± 4 pA, the shift in Palbociclib-treated slices was only 9 ± 3 pA (Fig 3B).

262 Therefore, Ccnd·Cdk4/6 inhibition by acute Palbociclib application causes a significant

decrease in the GABA$_A$R tonic current.

264 Given that the α4 subunit can also form synaptic receptors when clustered together with βxγ2

subunits[42], we tested whether the acute Cdk4/6 inhibition may in turn affect GABA$_A$

266 receptor synaptic function. To this end, we measured miniature inhibitory postsynaptic

currents (mIPSC) in organotypic cultures treated with 2.5 μM Palbociclib for 2 h, in the

268 presence of the action potential blocker tetrodotoxin (1 μM) (Fig S2A). We observed that the

amplitude of the spontaneous miniature responses was reduced after Ccnd·Cdk4/6 inhibitor

270 treatment as compared to the control (DMSO) treatment (-106 ± 20 pA for DMSO vs -51 ± 5 pA

for Palbociclib; Fig 3C and S2C-D). Putting all these electrophysiological experiments together,

272 our results suggest that Ccnd1·Cdk4/6 enhances GABA$_A$R activity in the hippocampus.

Since Ccnd1 regulates α4-containing GABA$_A$ receptors, we asked if the absence of Ccnd1 could

274 disrupt some α4-dependent functions of GABA$_A$Rs *in vivo*. The α4 subunit of GABA$_A$Rs

promotes synaptic pruning (the removal of dendritic spines) in the hippocampus of adolescent

276    mice[27]'[28]. To assess whether Ccnd1 could affect normal dendritic spine development, we analyzed the number and morphology of dendritic spines in cortical neurons from WT and

278    *CCND1* KO mice. Cultured neurons were transfected with GFP at 3 DIV, and dendritic spines were analyzed at 20 DIV, when spines are completely developed (Fig 3D-F and S3). The density

280    of dendritic spines did not depend on Ccnd1 expression (WT= 8.51 ± 0.16 vs *CCND1* KO= 8.28 ± 0.18 spines/10 µm, non-significant, Fig S3B). However, the ratio of the mushroom-shape (more

282    mature) spines relative to total spines was significantly increased in *CCND1* KO neurons compared with neurons from WT animals (0.45 ± 0.01 vs 0.50 ± 0.01, p<0.01, Fig 3F). Overall,

284    our data suggest that Ccnd1·Cdk4/6 enhances α4-dependent GABA$_A$R activity, which may have consequences for normal dendritic spine development.

**Ccnd1 increases the surface levels of the α4 subunit of GABA$_A$ receptors**

Phosphorylation of several subunits of GABA$_A$ receptors can affect their abundance at the cell surface, where they are active[43]. To analyze whether phosphorylation by Ccnd1·Cdk4 may

affect GABA$_A$R localization, we studied the surface levels of the α4 subunit when co-expressed heterologously with Ccnd1 in HEK-293T cells. For this, we transiently expressed α4 and β3 in

these cells and assessed the surface levels of α4 by biotin labeling of surface proteins (Fig 4A). We used the transferrin receptor (TFRC) protein levels as a surface protein control and

quantified the levels of α4 at the surface relative to the levels of α4 in the total cell extract (Fig 4B). We observed a statistically significant increase in the surface levels of α4 (148 ± 20%,

P=0.04, Fig 4A and B) only when Ccnd1-CAAX was also co-expressed. This increase was abolished in the presence of Ccnd1$^{K112E}$-CAAX, as well as when the non-phosphorylatable

mutant α4$^{T423AS431A}$ was used (98 ± 29%, P=0.93 and 107 ± 24%, P=0.82, respectively), indicating that the observed effect depends on T423/S431 phosphorylation by Ccnd1·Cdk4. These results

suggest that Ccnd1 increases the surface levels of the α4 subunit of GABA$_A$R in a kinase-dependent manner.

Figure 4. Ccnd1 increases surface levels of α4 subunit. *A) HEK-293T cells were transfected with β3 and α4-FLAG or the nonphosphorylatable mutant (α4$^{T42A3S431A}$-FLAG: AA). Either HA-Ccnd1-CAAX (D1) or HA-Ccnd1$^{K112E}$-CAAX (D1$^{KE}$) or a control vector was co-transfected. A biotinylation assay was used to purify cell surface proteins. TFRC is used to normalize α4 levels. Representative blots with whole cell extracts and surface proteins are shown. Antibodies used for protein detection are: polyclonal rabbit anti-FLAG to detect α4-FLAG; TFRC, monoclonal mouse anti-TFRC. B) Quantification of the experiment in A as a surface to WCE ratio. Bars show mean values ± S.E.M (n=4 independent experiments). \*p<0.05 (Unpaired student's t-test). C) Schematic model of the effects of Ccnd1·Cdk4 on α4-containing GABA$_A$ receptors: Ccnd1·Cdk4 phosphorylates the α4 subunit, promoting its surface localization and activity, thus increasing synaptic (phasic) and extrasynaptic (tonic) inhibition and altering the maturity of dendritic spines.*

**DISCUSSION**

318 In neurons, Ccnd1·Cdk4 has been involved in promoting proliferation versus differentiation of

stem cells during cortical development[13] and in the adult hippocampus[44]. However, the

320 role of Ccnd1 in the regulation of neuronal signaling has never been investigated before.

Interestingly, *CCND1* KO mice show an abnormal limb-clasping reflex[12], being this alteration

322 an early symptom of neurological defects associated with hippocampal neuron

malfunctioning[45]. In this work, we have unveiled a novel role of Ccnd1 in the stimulation of

324 $GABA_A$ receptor function (Fig 4C). Ccnd1 interacts with the α4 subunit of $GABA_A$R and

Ccnd1·Cdk4 complex phosphorylates T423 and S431 in the intracellular loop of α4. This

326 phosphorylation increases the surface levels of α4 subunit and the efficacy of $GABA_A$R activity

in whole cell patch-clamp recordings. Furthermore, Ccnd·Cdk4/6 inhibition decreases tonic

328 currents and the amplitude of mIPSCs in the hippocampus of newborn rats. Accordingly, the

proportion of mature dendritic spines is increased in cortical neurons from *CCND1* KO mice.

330 Therefore, we propose that Ccnd1·Cdk4 mediated phosphorylation is an important

physiological regulator of $GABA_A$R-mediated inhibition.

332 Ccnd1 has been found in the cytoplasm of postmitotic neurons[16]. As nuclear localization of

Ccnd1 in these cells promotes apoptosis, its cytoplasmic localization has been suggested as a

334 Ccnd1 sequestration mechanism to avoid cell death. However, Ccnd1·Cdk4 has cytoplasmic

functions interacting with membrane-associated proteins such as Filamin A, RalGTPases and

336 Paxillin[6]'[8]'[9]'[46]. It is then possible that Ccnd1·Cdk4 plays an active role in the cytoplasm

of post-mitotic neurons. Supporting this, the mRNA of Ccnd1 has also been found in the

338 synaptic neuropil in the hippocampus[47], suggesting local translation of Ccnd1 in the

dendrites. Also, Cdk4 has been localized in the cytoplasm in adolescent mouse hippocampal

340 neurons[20]. According with a hippocampal cytoplasmic expression of Ccnd1, here we have

detected co-immunoprecipitation of Ccnd1 and α4 in samples from mouse hippocampus

342    where this subunit of the GABA$_A$R is abundantly expressed[23], providing evidences for a

cytoplasmic role of Ccnd1 in neurons.

344    In this work, we have proved that Ccnd1·Cdk4 phosphorylates α4 at T423 and S431 in the

intracellular loop (ICL) *in vitro*. Moreover, in transiently transfected HEK-293T cells, we have

346    detected Ccnd1-specific phosphorylation of α4 *in vivo*. Rundown of GABA$_A$ receptors involves

phosphorylation-dephosphorylation processes[48]'[49]'[50]. The ICL between TM3 and TM4 of

348    several subunits of GABA$_A$Rs is regulated by phosphorylation, affecting the interaction of

GABA$_A$ receptors with other proteins, the stability and the surface localization of the subunits,

350    or the activity of the GABA$_A$ receptors[21]. For instance, PKC phosphorylates the α4 subunit in

the intracellular loop (S443) increasing its stability, membrane insertion and the efficacy of

352    GABA$_A$Rs[39]'[51]. Consistent with these previous data, we have determined by whole-cell

patch-clamp recordings that the expression of Ccnd1 or the phosphomimetic allele of α4

354    (α4$^{T423ES431E}$) caused a decrease in the rundown of α4β3 containing GABA$_A$ receptors in HEK-293

cells. We propose that Ccnd1·Cdk4 enhances the activity of α4-containing GABA$_A$ receptors by

356    promoting the surface accumulation of α4 through its phosphorylation. On the one hand,

Ccnd1 but not the inactive allele Ccnd1$^{K112E}$ is able to promote surface accumulation of α4.

358    Moreover, Ccnd1 does not increase surface levels of the non-phosphorylatable mutant of α4

(α4$^{T423AS431A}$).

360    The adjustment of the surface levels of GABA$_A$R subunits takes place through the regulation of

export and internalization processes. The ICL of β3 subunit is involved in internalization of the

362    receptors by clathrin. There is an atypical site for interaction with clathrin adaptor protein 2

(AP2) (KTHLRRRSSQLK), that contains the phosphorylation site for PKA, PKC and CamKII: it only

364    binds AP2 when dephosphorylated[52]. Such a mechanism could also be happening for α4

subunit, that contains a similar motif (NMRKRTNAL) in the intracellular loop (aa 385-393). In

366    fact, α4 has been described to interact directly with clathrin μ2 by immunoprecipitation[53].

Although T423 and S431 are not found inside the motif, phosphorylation of the ICL and/or the presence of Ccnd1 could be affecting the binding of α4 to clathrin. Another possibility is that the trafficking from the ER to the cytoplasmic membrane is affected. In the α4 and δ subunits there are dibasic or multibasic motifs in homologous positions that reduce export from the ER and increase transport back from the Golgi apparatus to the ER[54]. The mutation of dibasic motif to Ala in α4 enhances surface expression of the receptors. Again, phosphorylation of ICL or interaction with Ccnd1 could affect the association of dibasic sites with trafficking proteins.

Consistently with a role of Ccnd1·Cdk4/6 on GABAergic signaling, we have shown that inhibition of Cdk4/6 in CA1 pyramidal neurons produces a decrease in tonic currents and the amplitude of mIPSCs. Even though $\alpha$4 is mostly extrasynaptical, it can also form synaptic GABA$_A$Rs when assembled with $\gamma$2. Concerning mIPSC amplitude, it is feasible that phosphorylation of $\alpha$4 by Ccnd·Cdk4/6 complexes is still promoting its activity also when the $\alpha$4 subunit is forming $\alpha$4$\beta$x$\gamma$2 receptors at the synapse[55]. This would mean that $\alpha$4 phosphorylation by Ccnd·Cdk4/6 has the same effect irrespective of whether the receptor localizes synaptically or extrasynaptically. By contrast, $\alpha$4-containing synaptic and extrasynaptic receptors show different modulation by PKA and PKC phosphorylation. PKA activation decreases synaptic $\alpha$4 expression while increases extrasynaptic $\alpha$4. Activation of PKC instead promotes $\alpha$4-expression at the synapse and has no effect extrasynaptically[56].

Dendritic spines are the postsynaptic sites where most excitatory synapsis occurs. Their change in number and morphology underlies synaptic plasticity, important for learning and memory processes. GABA, through GABA$_A$Rs activation promotes spine shrinkage and elimination[57]. Specifically, GABA$_A$Rs containing α4βδ subunits have been involved in synaptic pruning at the CA1 and CA3 regions in the hippocampus[27]·[28]: the expression of the α4 subunit of GABA$_A$Rs increases in the hippocampus during adolescence[28] and the density of dendritic spines decreases post-pubertal, while in the *GABRA4* KO mice this pruning does not occur. Therefore, the density of dendritic spines in the hippocampus of adult mice is increased in the

GABRA4 KO mice[37]. The effect of α4 on synaptic pruning is mainly observed in mushroom-

and stubby-shaped dendritic spines[27],[28]. According to a role of Ccnd1 in promoting α4-

containing GABA$_A$Rs activity, we observed an increase in the proportion of mushroom-type

spines in CCND1 KO neurons. Mushroom spines are considered to be one of the mature types

of spines, involved in memory[58]. In this sense, the increase in α4βδ receptors at puberty

impairs learning[25] and the GABRA4 KO mouse has increased spatial memory[37]. In the same

direction, increased expression of Ccnd1 was found in CA1 region of aged learning-impaired

rats[59] and the expression of Ccnd1 in the hippocampus of rats triggers deficits in spatial

working memory[60]. Moreover, GABRA4 KO mice show an autistic-like behavior, and autism

has been associated with an increased number of spines[61]. The effect of Ccnd1 on GABA

signaling and dendritic spines opens the possibility of exploring Ccnd1 functions in diseases

related to the imbalance between inhibitory/excitatory signaling and dendritic spine

abnormalities such as autism spectrum disorders and epilepsy.

**MATERIALS and METHODS**

**Cell culture.** Mycoplasma-free HEK293T cells were obtained from the American Type Culture

Collection. Cells were maintained at 37ºC in a 5% $CO_2$ incubator, and grown in Dulbecco's

modified Eagle's medium (DMEM) supplemented with 10% Fetal Bovine Serum (FBS), 10 U/mL

penicillin/streptomycin (P/S) and 4 mM glutamine. Mycoplasma detection tests were

performed in the Cell culture service (SCT-CC). Transient transfection of vectors was performed

with Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions.

**Expression vectors.** Human CCND1 was fused to three copies of the Flag or HA or one of Red

monomer epitope under the CMV promoter in pcDNA3. Ccnd1-CAAX construct is a fusion of

the 3' end of the K-Ras ORF containing the CAAX motif (GGC TGT GTG AAA ATT AAA AAA TGC

ATT ATA ATG TAA) to the 3' end of the CCND1 ORF[10]. Mouse GABRA4 (IMAGE ID 6828002,

Source BioScience) and Human GABRB3 (IMAGE ID 3871111, Source BioScience) were used to

420   obtain N-terminal GST fusions in pGEX-KG (Clontech) or N-terminal FLAG fusions in pJEN1

(pcDNA3 derived, from Dr. Egea) or cloned into FCIV1 lentiviral vector (from Dr. Encinas).

422   Standard PCR-mediated site-directed mutagenesis was used to obtain the non-

phosphorylatable and phosphomimetic mutants of GABRA4 using the following primers: T423A

424   and           S431A:           CGCCAAATCCATTCAGCAGGGC           and

CCGAAGCTAAGTGAGCCTTAGGCGCGGCTTCAGAAGACTCCTGGAC; S456A and S458A:

426   CGGCCCCCGCTCCTCATGGCACATTGCGGC and CGGCAGATGAAAGACCTCTGGCTGM; T500A:

CGCCTCCTCCCCCTGCTCCAC and CCGCTGACACATTCCCAGCAGC; and T423E and S431E:

428   GCGAACCAAATCCATTCAGCAGGGC                               and

TAGCTAAGTGAGCCTTAGGCTCGGCTTCAGAAGACTCCTGGAC.

430   **Immunoprecipitation assays.** For co-immunoprecipitation experiments, HEK293T cells were

harvested 24 h after transfection. The cells were resuspended in RIPA buffer (50mM Tris

432   pH7.4, 150mM NaCl, 1% NP-40, 0.1% sodium deoxycholate, 0.1% SDS, 5mM EDTA, protease

and phosphatase inhibitors), rocked 1 h at 4ºC, and spun for 5 min at 600 g. The supernatants

434   were incubated with protein G magnetic beads (Dynabeads, Invitrogen) for 30' at 4ºC. After

the preclearing, the samples were incubated with 2 μg of αFLAG M2 antibody (Sigma)

436   overnight and precipitated with protein G dynabeads. IP of anti-phosphoThreonineProline

antibody (#9391, Cell Signaling) was performed similarly, but using protein A dynabeads. The

438   beads were collected and washed three times with 1 mL of cold RIPA buffer, and bound

proteins were separated by SDS-PAGE gels and visualized by western blots.

440   IP of endogenous Ccnd1 was carried out in hippocampus of adult female mice with a

polyclonal antibody (#06‑137, Upstate). Briefly, cleared cell extracts in RIPA buffer (with

442   protease and phosphatase inhibitors) were immunoprecipitated with Protein A linked to

magnetic beads (Invitrogen). An anti-FLAG (#F7425, Sigma) was used as a mock control.

444   Washes were performed with RIPA buffer without SDS.

**Immunoblotting**. For immunoblot, protein samples were resolved by SDS-PAGE, transferred to PVDF membranes (Millipore), and incubated with primary antibodies: anti-Ccnd1 (monoclonal DCS-6, BD biosciences, 1:500), anti-Gabra4 (polyclonal, AB5459, Merck, 1:1000, and polyclonal 12979-1-AP, Proteintech, 1:500), anti-Flag (polyclonal, 11508721, Invitrogen, 1:1000), anti-HA (rat monoclonal 3F10, Roche #11867431001, 1:5000), anti-TFR (monoclonal H68.4, Invitrogen #13‑6800, 1:1,000) and anti-GST (goat polyclonal, Amersham #27‑4577, 1:2000). Appropriate peroxidase-linked secondary antibodies (GE Healthcare UK Ltd) were detected using the chemiluminescent HRP substrate Immobilon Western (Millipore). Chemiluminescence was recorded with a ChemiDoc-MP imaging system (BioRad).

**Production and purification of recombinant proteins**. All of the GST fusions were expressed in *E. coli* BL21 (DE3) by adding 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) to LB broth cultures at a cell density of 0.3 $A_{600}$ and subsequent incubation for 4 h at 30 °C. The proteins were purified using glutathione-Sepharose 4B beads as directed by the supplier (Amersham Biosciences) in 500 µl of lysis buffer containing 50mM HEPES, pH 7.5, 150 mM NaCl, 1 mM EDTA, 0.5% Triton X-100, 10% glycerol, 1 mM DTT, and protease and phosphatase inhibitors. The concentration and purity of substrates were estimated by comparison to protein standards stained with Coomassie Brilliant Blue.

For *in vitro* transcription and translation, FLAG-Ccnd1 was amplified by PCR (forward primer: CGCGCTAATACGACTCACTATAGGGAGACCCAAGCCCATGGGATCAC; reverse primer: TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTGGCTGATCAGCGAGCTCTAG), transcribed *in vitro* with the T7 RNA polymerase (New England Biolabs), and *in vitro* translated with a Rabbit Reticulocyte Lysate system (Promega).

**GST pull-down assay**. For the GST pull-down assay, 400 ng of GST or GST-α4 or GST-Cter-α4 purified from *E. coli* immobilized on Glutathione-Sepharose 4B beads were incubated with Flag-Ccnd1 in binding buffer (20 mM HEPES-KOH, pH 7.5, 150 mM KCl, 5 mM MgCl$_2$, 0.5 mM EDTA, 0.1% NP-40, 1 mM DTT, 1 mM PMSF, 10% glycerol, protease and phosphatase

inhibitors) for 30 min at room temperature. After four washes with the same buffer, the

472    samples were analyzed by SDS-PAGE.

**Kinase assay**. Briefly, 0.2 mg substrate (either GST-α4, GST-Cter-α4 or GST-β3) was mixed with

474    1.5 µL of active Ccnd1-Cdk4 complex purified from baculovirus (Sigma C0620), 10 mM ATP, 7

mCi of γ-$^{32}$P-ATP (PerkinElmer, 3000 Ci/mmol), and either DMSO or 2 µM of the Cdk4/6-

476    inhibitor Palbociclib (Selleckchem, S1116) in 20 µL of kinase buffer (50 mM Tris‑HCl pH 7.5, 10

mM MgCl$_2$, 0.5 mM DTT, 1 mM EGTA and 2.5 mM b-glycerophosphate). This mixture was

478    incubated for 20 min at 30 ºC, then boiled in 2x Laemmli buffer, and separated by

electrophoresis. Phosphorylated proteins were visualized by autoradiography of the dried slab

480    gels.

**MS analysis of phosphopeptides**. The C terminus of α4 fused to GST was used in an *in vitro*

482    kinase assay with Ccnd1·Cdk4 in the presence of ATP or in the absence of ATP as a control.

Samples were subsequently subjected to SDS-PAGE. The gels were stained with Coomassie

484    Brilliant Blue G-250 colloidal (EZBlue Gel Staining Reagent, Sigma). After washing with water,

protein bands of interest were prepared and submitted to Fundació Institut d'Investigació

486    Biomèdica de Bellvitge (IDIBELL) Proteomics Service (Barcelona) for analysis of the

Chymotryptic peptide molecular masses by liquid chromatography-mass spectrometry. Gel

488    slices were manually cut and proteomic service suggested for each band-assay to recover

three slices (low 1, middle 2, and high 3 mobility) as phosphorylation could alter band mobility.

490    Briefly, gel bands were washed with water, ammonium bicarbonate (50mM) and 50%

acetonitrile. Next, samples were reduced by incubation with DTT (10 mM) at 60 ºC for 45 min

492    and alkylated with iodoacetamide (50mM) for 30 min, in the dark. Finally, proteins were

digested with chymotrypsin (5 ng/µL) at 25ºC overnight (Trypsin gold, Promega). Digestion was

494    stopped by addition of 5% formic acid and peptides extracted twice with 70% acetonitrile and

5% formic acid (10 min sonication). Peptide extracts were evaporated to dryness, resuspended

496    with 2% acetonitrile 0.1% formic acid and analyzed by nano-HPLC-MSMS.

**Surface biotinylation.** For surface biotinylation (all steps performed on ice) HEK-293T cells

498  were washed twice with PBS and incubated with 1 mg/mL EZ-link sulfo-NHS-SS-biotin (Pierce),

a nonpermeable biotin, in PBS for 15 min. Following surface labeling, nonconjugated biotin

500  was quenched by washing twice with TBS. Cells were lysed in RIPA buffer (50 mM Tris, pH7.4,

150 mM NaCl, 1% NP-40, 0.1% SDS, 5 mM EDTA, with phosphatase and protease inhibitors),

502  and protein concentrations were determined using a DC protein assay (Bio-Rad). Equal

amounts of protein (usually 1 mg in 500 μL) were incubated overnight with 20 μL NeutrAvidin-

504  coated beads (Pierce) at 4°C with constant rotation. The next day, beads were washed three

times with RIPA buffer and eluted samples were processed for immunoblotting.

506  **Whole-cell currents measurements in tsA201 cells.** Electrophysiological experiments were

done with mycoplasma-free tsA201 cells (kindly provided by Prof. F. Ciruela, Universitat de

508  Barcelona) and purchased from the American Type Culture Collection (ATCC, CRL-3216, RRID:

CVCL_0063). Mycoplasma detection tests were performed with PlasmoTest (Invivogen, code:

510  rep-pt1). Cells were transiently co-transfected with 1 μg total cDNA using polyethylenimine

(PEI) transfection reagent (1 mg/mL) in a 3:1 ratio (PEI:DNA). In all transfections the DNA ratio

512  used was 1:1:2 (plasmid A: plasmid B: plasmid C), where A is α4 or α4$^{T423ES431E}$, B is β3 and C is a

control vector or Ccnd1-CAAX codifying plasmid. Cells were re-plated on poly-D-lysine coated

514  glass coverslips to allow optimal density with isolated cells. All experiments were performed 48

h after transfection.

516  For the measurements of GABA-evoked whole-cell currents in tsA201, cells were visualized

with an inverted microscope (AxioVert A.1, Carl Zeiss) and maintained in extracellular flowing

518  solution at a rate of 60 mL/h. The extracellular solution contained (in mM): 140 NaCl, 5 KCl, 10

HEPES, 11 glucose, 2.5 CaCl$_2$, 1.2 MgCl$_2$ (pH = 7.4 with NaOH; osmolarity 305 mOsm/Kg

520  adjusted with sorbitol). Electrodes were fabricated from borosilicate glass (1.50 mm O.D., 1.16

I.D., Harvard Apparatus) pulled with a P-97 horizontal puller (Sutter Instruments) and polished

522  with a forge (MF-830, Narishige) to a final resistance of 2–5 MΩ. The intracellular solution

contained (in mM): 140 KCl, 2 $MgCl_2$, 2 $CaCl_2$, 10 HEPES, 1.1 EGTA, 2 Mg ATP (pH = 7.2 with

524    KOH; osmolarity 295 mOsm/Kg adjusted with sorbitol).

Macroscopic $GABA_A$-mediated currents were recorded at room temperature (22-25˚C) in the

526    whole-cell configuration from cells positive for Venus ($\alpha$4 in FCIV1) and mCherry (Ccnd1-CAAX

in pcDNA3 Red monomer) fluorescence at a holding membrane potential of -60 mV. Currents

528    were recorded with Axopatch 200B amplifier, filtered at 2 kHz and digitized at 5 kHz using

Digidata 1440A interface with pClamp 10 software (Molecular Devices Corporation). Series

530    resistance was typically 5–20 M$\Omega$, and was monitored at the beginning and at the end of the

experiment. Cells that showed a change in series resistance greater than 20% were rejected.

532    Rapid application of the agonist GABA (1-2 ms exchange) was applied by switching between a

continuously flowing control solution (extracellular solution diluted by 5%) and a GABA-

534    containing solution (1 µM GABA and 2.5mg/mL of sucrose diluted in extracellular solution).

Solution switching was achieved by piezoelectric translation of a theta-barrel application tool

536    made from borosilicate glass (1.5 mm O.D.; Sutter Instruments) mounted on a piezoelectric

translator (P-601.30; Physik Instrumente). Agonist was applied at 2 minutes interval and the

538    magnitude of the peak current was measured ($I_{GABA}$). Electrophysiological recordings were

analyzed using IGOR Pro (Wavemetrics Inc.) with NeuroMatic (Jason Rothman, UCL).

540    **Preparation of organotypic hippocampal slice cultures.** Organotypic hippocampal slice

cultures were prepared as previously described[62]. Briefly, whole brain of newborn rats

542    (postnatal days 5-7) were dissected and placed in ice-cold $Ca^{2+}$-free dissection solution (10 mM

D-glucose, 4 mM KCl, 26 mM $NaHCO_3$, 8% sucrose, 5 mM $MgCl_2$, 1:1000 Phenol Red) saturated

544    with 5% $CO_2$ / 95% $O_2$, enabling the brain to chill for 1 min. Hippocampi were then isolated

with the help of a magnifying glass, cut in slices (400 µm) in the same solution using a tissue

546    chopper (Stoelting), and maintained at 35.5 °C in culture on permeable membranes in a

medium containing 20% horse serum, 1 mM $CaCl_2$, 2 mM $MgSO_4$, 1 mg/L insulin, 0.0012%

548     ascorbic acid, 30 mM Hepes, 13 mM D-glucose, and 5.2 mM $NaHCO_3$. Culture medium was

replaced with fresh one every 2-3 days. The slices were used at 4-8 days *in vitro*.

550     **Whole-cell recordings of $GABA_A$ receptor-mediated currents.** Prior to the electrophysiological

recordings, slices were treated for 2 hours with either DMSO (control solution) or 2.5 μM

552     Palbociclib (Cdk4/6 inhibitor) (Selleckchem). After that, slices were placed in a chamber and

perfused with artificial cerebrospinal fluid [aCSF: 119 mM NaCl, 2.5 mM KCl, 1 mM $NaH_2PO_4$,

554     26 mM $NaHCO_3$, 11 mM glucose, 1.2 mM $MgCl_2$, 2.5 mM $CaCl_2$; osmolarity adjusted to 290

mOsm; pH7.5] supplemented with 100 μM AP5 (NMDA receptor antagonist), 10 μM CNQX

556     (AMPA receptor antagonist), 1 μM strychnine (glycine receptor antagonist), 4 μM 2-

chloroadenosine (to reduce neurotransmitter release and stabilize synaptic transmission) and

558     DMSO or Palbociclib as external solution gassed with 5% $CO_2$ / 95% $O_2$ at 29 °C in the

electrophysiology set-up. Patch recording pipettes (4-6 MΩ) were pulled from thin-walled

560     borosilicate capillary glass (World Precision Instruments [WPI], Sarasota, FL) on a P-2000 laser

electrode puller (Sutter Instrument, San Rafael, CA) and filled with high chloride internal

562     solution (178 mM CsCl, 10 mM Hepes, 2.5 mM $MgCl_2$, 4 mM $Na_2ATP$, 0.4 mM $Na_3GTP$, 10 mM

sodium phosphocreatine, 0.6 mM EGTA; pH adjusted to 7.2, osmolarity adjusted to 290

564     mOsm). GABA inhibitory postsynaptic currents were evoked with single voltage pulses (200 μs,

<30 V) delivered through platinum-iridium bipolar electrodes (FHC, Bowdoin, ME, USA) placed

566     on Schaffer collaterals. CA1 pyramidal neurons were recorded under whole-cell voltage-clamp

at -60 mV using Multiclamp 700A amplifier (Molecular Devices, San Jose, CA, USA). At least 10

568     min of stable baseline synaptic responses and holding currents were recorded. The tonic

$GABA_A$ receptor-mediated current was measured as the outward shift in holding current

570     following application of picrotoxin (100 μM). Synaptic responses were used to monitor

effective inhibition of $GABA_A$Rs by picrotoxin. To measure miniature inhibitory postsynaptic

572     currents (mIPSC), 1 μM tetrodotoxin was added to the aCSF to block action potentials.

Recordings were obtained with the gap-free mode of pClamp and mIPSCs were analyzed with

574   the event detection of pCLAMP software (Molecular Devices).

**Cortical neuron culture.** For primary culture of cortical neurons, cortexes were dissected from

576   E15.5 *CCND1* KO or WT mice in ice-cold HBSS-MHPS (Hank's Balanced Salt Solution, 10 mM

MgSO$_4$, 10 mM HEPES pH 7.2 and 10 U/mL P/S). After dissection, cortexes were incubated with

578   papain solution (1 mg/mL papain in HBSS-MHPS) at 37ºC for 23 min. Enzymatic digestion was

inhibited by washing the tissue 3 times with 10 mg/mL trypsin inhibitor (Fisher) in HBSS-MHPS.

580   Additionally, one more wash was done with Neurobasal medium supplemented with 2% B27, 2

mM Glutamax and 1 mM sodium pyruvate (NB27, Fisher) at room temperature. The tissue was

582   triturated by passing it through a flame-polished Pasteur pipette (<20 times). After the

mechanical disaggregation, the supernatant containing mostly single cells was centrifuged at

584   650 rpm for 4 min. The pellet was resuspended in NB27 and the cells were counted in a

Neubauer chamber. Cells were seeded at low density (25000 cells/cm$^2$) with NB27 medium in

586   poly-D-lysine (0.5 mg/mL) and laminin (5 µg/mL)-coated plates, in 1:1 conditioned medium.

One third of the culture medium is replaced with fresh one every 3-4 days. Primary cortical

588   neurons were transfected at 3 DIV using Lipofectamine 2000 (Invitrogen) following the

manufacturer's recommendations and the proportion 0.3 µg DNA: 1.2 µL Lipofectamine2000.

590   The media was changed 4 hours after transfection and cells were fixed at 20 DIV.

**Genotyping of *CCND1* knockout mice.** *CCND1* knockout mice[12] were obtained from Charles-

592   River. Transgenic mice were maintained in a mixed background of C57BL/6 and SV129.

Embryo's tails were collected and warmed at 95 ºC NaOH 50 mM for 45 min. After

594   neutralization with Tris 0.15 M pH8.5, the samples were then centrifuged for 1 minute at

10000 rpm. Supernatant containing genomic DNA was used for CCND1 genotyping by PCR. The

596   following primers were used: a common primer CTGTCGGCGCAGTAGCAGAGAGCTACAGAC, a

CCND1   primer   CGCACAGGTCTCCTCCGTCTTGAGCATGGC   and   a   neomycin   primer

598   CTAGTGAGACGTGCTACTTCCATTTGTCACG. PCR was performed in a Thermal Cycler (T100 Bio-

Rad) and PCR products were analyzed in a 1.5 % agarose gel followed by Ethidium Bromide

600   staining. The expected band pattern is 249 bp for WT, and 394 bp for KO alleles.

**Immunofluorescence and spine analysis**. Cortical neurons were fixed using 4% PFA and 4%

602   sucrose in PBS for 15 min at room temperature and then washed with PBS. Neurons were

permeabilized for 5 min with 0.1% Triton X-100 in PBS and blocked with 3% BSA in PBS.

604   Primary antibody α-GFP (A11120, ThermoFisher) was diluted 1:200 in 0.3% BSA in PBS.

Proteins were detected by incubation with secondary antibody Alexa 488 rabbit anti-mouse

606   (A11059, ThermoFisher). Images were acquired with an Olympus FV1000 confocal microscope

using the following parameters: stacks of 10 slices were imaged every 0.37 μm, under a 60X

608   objective. Laser power and PMT values were kept constant throughout images and conditions.

Immunofluorescence quantification was performed using NeuronJ of ImageJ. Spine density

610   and morphology analysis of WT and *CCND1* KO cortical neurons was performed on 3D

reconstructions of confocal z series acquired using a 2x zoom. Spines were counted as

612   mushroom-type if the spine head was wider than the spine neck. All analysis was done blind to

the experimental condition.

614   **Statistical analyses**. Different statistical analyses were used in this work. In every analysis, the

minimum level of statistical significance was a p-value equal to 0.05. The significance level is

616   represented in each graph as indicated. In order to evaluate possible differences between two

experimental groups a t-test was performed. When there was a multiple comparison, one-way

618   ANOVA with a post-hoc test (Tukey) was performed. In whole-cell recordings in ts201A cells,

comparisons between groups were done using the parametric student's t-test. For tonic

620   currents comparison, Wilcoxon test was performed and in mIPSC amplitude, Mann-Whitney

and Kolmogorov-Smirnov tests were done. Microsoft Excel and GraphPad Prism v5.0

622   (GraphPad) were used for statistical analysis and graphical representation.

**Ethical considerations.** All procedures with animals follow the protocols approved by the

624   Institutional Committee of Care and Use of Animals (Comitè Institucional de Cura i Ús

d'Animals), and experiments were approved by the Ethics Committee of the University of

626    Lleida (CEEA 03-02/19). Animals were housed and maintained in the animal facility of the

University of Lleida with 12h:12h light/dark cycle and food/water available *ad libitum*. For

628    electrophysiological experiments, all biosafety procedures and animal care protocols were

approved by the bioethics committee from the Consejo Superior de Investigaciones Cientificas

630    and were performed according to the guidelines set out in the European Community Council

Directives (2010/63/EU, 86/609/EEC). All efforts were made to minimize the number of

632    animals and their suffering.

634    **COMPETING INTEREST STATEMENT**

The authors declared that they have no conflict of interest.

**ACKNOWLEDGEMENTS**

638    We are grateful to M. Encinas for plasmids. We thank Sònia Rius for technical assistance and

members of the CYC lab for helpful considerations. The cell culture experiments were

640    performed in the Cell Culture Scientific & technical Service from Universitat de Lleida. This

work was funded by the Spanish Ministry of Innovation and Science MICINN (PID2019-

642    104859GB-I00 to E.G. and PID2020-117651RB to J.A.E.) and by Generalitat de Catalunya (2017-

SGR-569). M. Ventura Monserrat was supported by a predoctoral fellowship of University of

644    Lleida.

646    **AUTHORS CONTRIBUTION**

Neus Pedraza: conceptualization, supervision, validation, investigation, methodology, writing-

648    original draft, writing – review & editing; Ma Ventura Monserrat: validation, investigation,

methodology; Francisco Ferrezuelo: funding acquisition, data curation, writing – review &

650    editing; Jordi Torres: funding acquisition, supervision, writing – review & editing; Neus

Colomina: funding acquisition, methodology; David Soto: funding acquisition, investigation, validation, methodology; Fede Miguez: investigation, methodology; JA Esteban: supervision, validation, methodology, funding acquisition; Joaquim Egea: funding acquisition, conceptualization, methodology; Eloi Garí: conceptualization, supervision, funding acquisition, investigation, writing – review & editing.

## REFERENCES

1.  Sherr CJ, Roberts JM. Living with or without cyclins and cyclin-dependent kinases. Genes Dev. 2004;18: 2699–2711. doi:10.1101/gad.1256504

2.  Hydbring, Per, Malumbres M, Sicinski P. Non-canonical functions of cell cycle cyclins and cyclin-dependent kinases. Nat Publ Gr. 2016 [cited 17 Feb 2022]. doi:10.1038/nrm.2016.27

3.  Neumeister P, Pixley FJ, Xiong Y, Xie H, Wu K, Ashton A, et al. Cyclin D1 Governs Adhesion and Motility of Macrophages. Mol Biol Cell. 2003;14. doi:10.1091/mbc.02-07-0102

4.  Li Z, Wang C, Jiao X, Lu Y, Fu M, Quong AA, et al. Cyclin D1 Regulates Cellular Migration through the Inhibition of Thrombospondin 1 and ROCK Signaling. Mol Cell Biol. 2006;26: 4240–4256. doi:10.1128/mcb.02124-05

5.  Cemeli T, Guasch-Vallés M, Nàger M, Felip I, Cambray S, Santacana M, et al. Cytoplasmic cyclin D1 regulates glioblastoma dissemination. J Pathol. 2019;248: 501–513. doi:10.1002/path.5277

6.  Zhong Z, Yeow WS, Zou C, Wassell R, Wang C, Pestell RG, et al. Cyclin D1/cyclin-dependent kinase 4 interacts with filamin A and affects the migration and invasion potential of breast cancer cells. Cancer Res. 2010;70: 2105–2114. doi:10.1158/0008-5472.CAN-08-1108

7.  Meng H, Tian L, Zhou J, Li Z, Jiao X, Li WW, et al. Cell Cycle PACSIN 2 represses cellular

migration through direct association with cyclin D1 but not its alternate splice form

678     cyclin D1b. Cell Cycle. 2011;10: 73–81. doi:10.4161/cc.10.1.14243

8.      Fernández RMH, Ruiz-Miró M, Dolcet X, Aldea M, Garí E. Cyclin D1 interacts and

680     collaborates with Ral GTPases enhancing cell detachment and motility. Oncogene.

        2011;30: 1936–1946. doi:10.1038/onc.2010.577

682  9. Fusté NP, Fernández-Hernández R, Cemeli T, Mirantes C, Pedraza N, Rafel M, et al.

        Cytoplasmic cyclin D1 regulates cell invasion and metastasis through the

684     phosphorylation of paxillin. Nat Commun. 2016;7. doi:10.1038/ncomms11581

10.     Fusté NP, Castelblanco E, Felip I, Santacana M, Fernández-Hernández R, Gatius S, et al.

686     Characterization of cytoplasmic cyclin D1 as a marker of invasiveness in cancer.

        Oncotarget. 2016;7: 26979–26991. doi:10.18632/oncotarget.8876

688  11. Chen K, Jiao X, Ashton A, Rocco A Di, Pestell TG, Sun Y, et al. The membrane-associated

        form of cyclin D1 enhances cellular invasion. Oncogenesis. 2020;9. doi:10.1038/s41389-

690     020-00266-y

12.     Sicinski P, Donaher JL, Parker SB, Li T, Fazeli A, Gardner H, et al. Cyclin D1 provides a link

692     between development and oncogenesis in the retina and breast. Cell. 1995;82: 621–

        630. doi:10.1016/0092-8674(95)90034-9

694  13. Lange C, Huttner WB, Calegari F. Cdk4/CyclinD1 Overexpression in Neural Stem Cells

        Shortens G1, Delays Neurogenesis, and Promotes the Generation and Expansion of

696     Basal Progenitors. Cell Stem Cell. 2009;5: 320–331. doi:10.1016/j.stem.2009.05.026

14.     Lim S, Kaldis P. Loss of Cdk2 and Cdk4 induces a switch from proliferation to

698     differentiation in neural stem cells. Stem Cells. 2012;30: 1509–1520.

        doi:10.1002/stem.1114

700  15. Miyashita S, Owa T, Seto Y, Yamashita M, Aida S, Sone M, et al. Cyclin D1 controls

        development of cerebellar granule cell progenitors through phosphorylation and

702     stabilization of ATOH1. EMBO J. 2021;40: 1–17. doi:10.15252/embj.2020105712

16.  Sumrejkanchanakij P, Tamamori-Adachi M, Matsunaga Y, Eto K, Ikeda MA. Role of cyclin D1 cytoplasmic sequestration in the survival of postmitotic neurons. Oncogene. 2003;22: 8723–8730. doi:10.1038/sj.onc.1206870

17.  Sumrejkanchanakij P, Eto K, Ikeda MA. Cytoplasmic sequestration of cyclin D1 associated with cell cycle withdrawal of neuroblastoma cells. Biochem Biophys Res Commun. 2006;340: 302–308. doi:10.1016/j.bbrc.2005.11.181

18.  Marampon F, Casimiro M, Fu M, Powell M, Popov V, Lindsay J, et al. Nerve Growth factor regulation of cyclin D1 in PC12 cells through a p21RAS extracellular signal-regulated kinase pathway requires cooperative interactions between Sp1 and nuclear factor-kappaB. Mol Biol Cell. 2008;19: 2566–78. doi:10.1091/mbc.e06-12-1110

19.  Schmetsdorf S, Gärtner U, Arendt T. Expression of cell cycle-related proteins in developing and adult mouse hippocampus. Int J Dev Neurosci. 2005;23: 101–112. doi:10.1016/j.ijdevneu.2004.07.019

20.  Li C, Li X, Chen W, Yu S, Chen J, Wang H, et al. The different roles of cyclinD1-CDK4 in STP and mGluR-LTD during the postnatal development in mice hippocampus area CA1. BMC Dev Biol. 2007;7: 1–9. doi:10.1186/1471-213X-7-57

21.  Nakamura Y, Darnieder LM, Deeb TZ, Moss SJ. Regulation of GABAARs by phosphorylation. Adv Pharmacol. 2015;72: 97–146. doi:10.1016/bs.apha.2014.11.008

22.  Comenencia-Ortiz E, Moss SJ, Davies PA. Phosphorylation of GABA A receptors influences receptor trafficking and neurosteroid actions. Psycopharmacology. 2014;231: 3453–3465. doi:10.1007/s00213-014-3617-z

23.  Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, et al. GABAA receptor α4 subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and the action of gaboxadol. Proc Natl Acad Sci U S A. 2006;103: 15230–15235. doi:10.1073/pnas.0604304103

24.  Shen H, Gong QH, Aoki C, Yuan M, Ruderman Y, Williams K, et al. Reversal of

neurosteroid effects at α4β2δ GABAA receptors triggers anxiety at puberty. Nat

730       Neurosci. 2007;10: 469–477. doi:10.1038/nn1868

25.    Shen H, Sabaliauskas N, Sherpa A, Fenton AA, Aoki C, Smith SS. A Critical Role for α4βδ

732       GABAA Receptors in Shaping Learning Deficits at Puberty in Mice. Science (80- ).

2010;327: 1515–1518. doi:10.1126/science.1184245.A

734  26.    Afroz S, Shen H, Smith SS. α4βδ GABAA receptors reduce dendritic spine density in CA1

hippocampus and impair relearning ability of adolescent female mice: Effects of a GABA

736       agonist and a stress steroid. Neuroscience. 2017;347: 22–35.

doi:10.1016/j.neuroscience.2017.01.051

738  27.    Afroz S, Parato J, Shen H, Smith SS. Synaptic pruning in the female hippocampus is

triggered at puberty by extrasynaptic GABAA receptors on dendritic spines. Elife.

740       2016;5: 1–23. doi:10.7554/eLife.15106

28.    Parato J, Shen H, Smith SS. α4βδ GABA A Receptors Trigger Synaptic Pruning and

742       Reduce Dendritic Length of Female Mouse CA3 Hippocampal Pyramidal Cells at

Puberty. Neuroscience. 2019;398: 23–36. doi:10.1016/j.neuroscience.2018.11.032

744  29.    Shen H, Sabaliauskas N, Yang L, Aoki C, Smith SS. Role of α4-containing GABAA

receptors in limiting synaptic plasticity and spatial learning of female mice during the

746       pubertal period. Brain Res. 2017;1654: 116–122. doi:10.1016/j.brainres.2016.01.020

30.    Moore MD, Cushman J, Chandra D, Homanics GE, Olsen RW, Fanselow MS. Trace and

748       contextual fear conditioning is enhanced in mice lacking the α4 subunit of the GABAA

receptor. Neurobiol Learn Mem. 2010;93: 383–387. doi:10.1016/j.nlm.2009.12.004

750  31.    Duveau V, Laustela S, Barth L, Gianolini F, Vogt KE, Keist R, et al. Spatiotemporal

specificity of GABAA receptor-mediated regulation of adult hippocampal neurogenesis.

752       Eur J Neurosci. 2011;34: 362–373. doi:10.1111/j.1460-9568.2011.07782.x

32.    Ma DQ, Whitehead PL, Menold MM, Martin ER, Ashley-Koch AE, Mei H, et al.

754       Identification of significant association and gene-gene interaction of GABA receptor

subunit genes in autism. Am J Hum Genet. 2005;77: 377–388. doi:10.1086/433195

756   33.   Collins AL, Ma D, Whitehead PL, Martin ER, Wright HH, Abramson RK, et al.

Investigation of autism and GABA receptor subunit genes in multiple ethnic groups.

758   Neurogenetics. 2006;7: 167–174. doi:10.1007/s10048-006-0045-1

34.   Griswold AJ, Van Booven D, Cuccaro ML, Haines JL, Gilbert JR, Pericak-Vance MA.

760   Identification of rare noncoding sequence variants in gamma-aminobutyric acid A

receptor, alpha 4 subunit in autism spectrum disorder. Neurogenetics. 2018;19: 17–26.

762   doi:10.1007/s10048-017-0529-1

35.   Fatemi SH, Reutiman TJ, Folsom TD, Rooney RJ, Patel H, Thuras PD. mRNA and Protein

764   Levels for GABAAα4, α5, β1 and GABABR1 Receptors are Altered in Brains From

Subjects With Autism. J Autism Dev Disord. 2010;40: 612–624. doi:10.1007/s10803-

766   009-0924-z.mRNA

36.   Fatemi S, Reutiman T, Folsom TD, Thuras PD. GABA A Receptor Downregulation in

768   Brains of Subjects with Autism. J Autism Dev Disord. 2009;39: 223–230.

doi:10.1007/s10803-008-0646-7

770   37.   Fan C, Gao Y, Liang G, Huang L, Wang J, Yang X, et al. Transcriptomics of Gabra4

knockout mice reveals common NMDAR pathways underlying autism, memory, and

772   epilepsy. Mol Autism. 2020;11: 1–17. doi:10.1186/s13229-020-0318-9

38.   Ruiz-Miró M, Colomina N, Fernández RMH, Garí E, Gallego C, Aldea M. Translokin

774   (Cep57) Interacts with Cyclin D1 and Prevents Its Nuclear Accumulation in Quiescent

Fibroblasts. Traffic. 2011;12: 549–562. doi:10.1111/J.1600-0854.2011.01176.X

776   39.   Abramian AM, Comenencia-Ortiz E, Vithlani M, Tretter EV, Sieghart W, Davies PA, et al.

Protein kinase C phosphorylation regulates membrane insertion of GABA A receptor

778   subtypes that mediate tonic inhibition. J Biol Chem. 2010;285: 41795–41805.

doi:10.1074/jbc.M110.149229

780   40.   Baker GL, Landis MW, Hinds PW. Multiple functions of D-type cyclins can antagonize

pRb-mediated suppression of proliferation. Cell Cycle. 2005;4: 329–337.

782      doi:10.4161/cc.4.2.1485

41.    Bright DP, Smart TG. Methods for recording and measuring tonic GABAA receptor-

784      mediated inhibition. Front Neural Circuits. 2013;5: 193. doi:10.3389/fncir.2013.00193

42.    Tang X, Hernandez CC, Macdonald RL. Modulation of spontaneous and GABA-evoked

786      tonic α4β3δ and α4β3γ2L GABAA receptor currents by protein kinase A. J Neurophysiol.

2010;103: 1007–1019. doi:10.1152/jn.00801.2009

788  43.    Connelly WM, Errington AC, Giovanni G Di, Crunelli V. Metabotropic regulation of

extrasynaptic GABAA receptors. Front Neural Circuits. 2013;7: 1–8.

790      doi:10.3389/fncir.2013.00171

44.    Artegiani B, Lindemann D, Calegari F. Overexpression of cdk4 and cyclinD1 triggers

792      greater expansion of neural stem cells in the adult mouse brain. J Exp Med. 2011;208:

937–948. doi:10.1084/jem.20102167

794  45.    Yamada M, Takeshita T, Miura S, Murata K, Kimura Y, Ishii N, et al. Loss of hippocampal

CA3 pyramidal neurons in mice lacking STAM1. Mol Cell Biol. 2001;21: 3807–19.

796      doi:10.1128/MCB.21.11.3807-3819.2001

46.    Fernández-Hernández R, Rafel M, Fusté NP, Aguayo RS, Casanova JM, Egea J, et al. Cell

798      Cycle Cyclin D1 localizes in the cytoplasm of keratinocytes during skin differentiation

and regulates cell-matrix adhesion) Cyclin D1 localizes in the cytoplasm of keratinocytes

800      during skin differentiation and regulates cell-matrix adhesion. Cell Cycle. 2013;12:

2510–2517. doi:10.4161/cc.25590

802  47.    Cajigas IJ, Tushev G, Will TJ, Tom Dieck S, Fuerst N, Schuman EM. The Local

Transcriptome in the Synaptic Neuropil Revealed by Deep Sequencing and High-

804      Resolution Imaging. Neuron. 2012;74: 453–466. doi:10.1016/j.neuron.2012.02.036

48.    Stelzer A, Kay A, Wong R. GABAA-receptor function in hippocampal cells is maintained

806      by phosphorylation factors. Science (80- ). 1988;241: 339–41.

doi:10.1126/science.2455347

808    49.    Chen QX, Stelzer A, KAY AR, Wong RKS. GABAA receptor function is regulated by

phosphorylation in acutely dissociated guinea-pig hippocampal neurones. J Physiol.

810    1990;420: 207–221.

50.    Gyenes M, Wang Q, Gibbs TT, Farb DH. Phosphorylation factors control

812    neurotransmitter and neuromodulator actions at the gamma-aminobutyric acid type A

receptor. Mol Pharmacol. 1994;46: 542–549.

814    51.    Abramian AM, Comenencia-Ortiz E, Modgil A, Vien TN, Nakamura Y, Moore YE, et al.

Neurosteroids promote phosphorylation and membrane insertion of extrasynaptic

816    GABAA receptors. Proc Natl Acad Sci U S A. 2014;111: 7132–7137.

doi:10.1073/pnas.1403285111

818    52.    Kittler JT, Chen G, Honing S, Bogdanov Y, McAinsh K, Arancibia-Carcamo IL, et al.

Phospho-dependent binding of the clathrin AP2 adaptor complex to GABA A receptors

820    regulates the efficacy of inhibitory synaptic transmission. Proc Natl Acad Sci U S A.

2005;102: 14871–14876. doi:10.1073/pnas.0506653102

822    53.    Gonzalez C, Moss SJ, Olsen RW. Ethanol promotes clathrin adaptor-mediated

endocytosis via the intracellular domain of δ-containing GABAA receptors. J Neurosci.

824    2012;32: 17874–17881. doi:10.1523/JNEUROSCI.2535-12.2012

54.    Bracamontes JR, Li P, Akk G, Steinbach JH. Mutations in the Main Cytoplasmic Loop of

826    the GABA A Receptor a 4 and d Subunits Have Opposite Effects on Surface Expression.

Mol Pharmacol. 2014;86: 20–27. doi:10.1124/mol.114.092791

828    55.    Wafford KA, Thompson SA, Thomas D, Sikela J, Wilcox AS, Whiting PJ. Functional

Characterization of Human γ-Aminobutyric AcidA Receptors Containing the a4 Subunit.

830    Mol Pharmacol. 1996.

56.    Bohnsack JP, Carlson SL, Morrow AL. Differential regulation of synaptic and

832    extrasynaptic α4 GABA(A) receptor populations by protein kinase A and protein kinase

C in cultured cortical neurons. Neuropharmacology. 2016;105: 124–132.

834    doi:10.1016/j.neuropharm.2016.01.009

57.    Hayama T, Noguchi J, Watanabe S, Takahashi N, Hayashi-Takagi A, Ellis-Davies GCR, et

836    al. GABA promotes the competitive selection of dendritic spines by controlling local Ca

2+ signaling. Nat Neurosci. 2013;16: 1409–1416. doi:10.1038/NN.3496

838    58.    Bourne J, Harris KM. Do thin spines learn to be mushroom spines that remember? Curr

Opin Neurobiol. 2007;17: 381–6. doi:10.1016/j.conb.2007.04.009

840    59.    Burger C, López MC, Feller JA, Baker H V, Muzyczka N, Mandel RJ. Changes in

transcription within the CA1 field of the hippocampus are associated with age-related

842    spatial learning impairments. Neurobiol Learn Mem. 2007;87: 21–41.

doi:10.1016/j.nlm.2006.05.003

844    60.    Wu K, Li S, Bodhinathan K, Meyers C, Chen W, Campbell-Thompson M, et al. Enhanced

expression of Pctk1, Tcf12 and Ccnd1 in hippocampus of rats: Impact on cognitive

846    function, synaptic plasticity and pathology. Neurobiol Learn Mem. 2012;97: 69–80.

doi:10.1016/j.nlm.2011.09.006

848    61.    Penzes P, Cahill ME, Jones KA, VanLeeuwen J-E, Woolfrey KM. Dendritic spine pathology

in neuropsychiatric disorders. Nat Neurosci. 2011;14: 285–93. doi:10.1038/nn.2741

850    62.    Gähwiler B, Campogna M, Debanne D, McKinney R, Thompson S. Organotypic slice

cultures: a technique has come of age. Trends Neurosci. 1997;20: 471–477.

852    doi:10.1016/S0166-2236(97)01122-3

| Accession | Description | Score | Coverage | # Proteins | # Unique Peptides | # Peptides | MW (kDa) |
|---|---|---|---|---|---|---|---|
| Q9D6F4 | GBRA4_MOUSE | 309,37 | 33,70 | 1 | 16 | 17 | 60,8 |

| Sample | Peptide sequence | Modifications | phoSite Probab | q-Value | MH+ (Da) | ΔM (ppm) | RT (min) |
|---|---|---|---|---|---|---|---|
| ATP | TSAVQESSEATPK | | | 0,002 | 1334,64 | -1,09 | 15,99 |
| | TSAVQESSEAtPK | T11(Phospho) | T(11): 100,0 | 0,012 | 1414,61 | -1,91 | 15,92 |
| | AHLASSPNPFSR | | | 0,038 | 1283,64 | -3,28 | 19,81 |
| | AHLASsPNPFSR | S6(Phospho) | S(6): 98,3 | 0,131 | 1363,61 | -5,17 | 20,50 |
| no ATP | TSAVQESSEATPK | | | 0,0000 | 1334,64 | -0,91 | 16,33 |
| | AHLASSPNPFSR | | | 0,0000 | 1283,64 | -3,57 | 20,01 |

**S1 Table. The α4 subunit of GABA$_A$ receptors is phosphorylated at T423 and S431 by Ccnd1·Cdk4.** *The GST-Ct-α4 wild type was used in an in vitro kinase assay with Ccnd1·Cdk4, in the presence or absence of ATP, and the Chymotryptic peptide molecular masses were analyzed by liquid chromatography-mass spectrometry. MH+: proton adduct; ΔM: mass difference; RT: retention time; q-value: adjusted p-value. Both T423 and S431 of α4 subunit (highlighted in red) were found to be phosphorylated by Ccnd1·Cdk4.*

866 ***S1 Figure. Ccnd1 or the phosphomimetic allele of α4 (α4$^{EE}$) decreases GABA$_A$ receptor rundown.*** *A)*

*Representative recording of whole-cell currents evoked by rapid application of 1 μM GABA (solid line*

868 *above the trace) to HEK-tsA201 cells expressing wild-type α4β3 GABA$_A$Rs. The GABA-evoked current at*

*the beginning of the recording (time 0 min) and the response after 14 minutes (time 14 min) are shown.*

870 *Dashed line denotes the zero current. Holding potential was held at −60mV. B) Same as A, but from cells*

*co-expressing wild-type α4β3 GABA$_A$Rs plus Ccnd1-CAAX. Note that the rundown of the peak current at*

872 *14 minutes is smaller than in condition A. C) Same as A, but from cells co-expressing phosphomimetic*

*(T423E S431E) GABA$_A$ receptor α4$^{EE}$β3. D) Bar graph showing the mean ± S.E.M. of the relative current at*

874 *t=12 minutes compared with current at time=0 for cells expressing wild type α4β3 GABA$_A$Rs (n=7), α4β3*

*GABA$_A$Rs plus Ccnd1-CAAX (n=5) or for α4$^{EE}$β3 GABA$_A$Rs (n=6). Open circles denote single experimental*

876 *values. \*p<0.05, α4β3 vs. α4β3+Ccnd1-CAAX; #p<0.05 α4β3 vs. α4$^{EE}$β3; Unpaired student's t-Test.*

*S2 Figure. Cdk4/6 inhibition by Palbociclib decreases GABA tonic currents and mIPSC amplitude in rat*

880　*hippocampal slices. A) Schematic representation of the experimental design for GABA currents*

*recordings in CA1 neurons: hippocampal slices from newborn rats were cultured in vitro, treated for 2 h*

882　*with DMSO or Palbociclib 2.5 μM, and tonic (extrasynaptic) or mIPSCs (synaptic currents) were*

*measured. B) GABA inhibitory postsynaptic currents were evoked by stimulation of Schaffer collaterals*

884　*and the tonic GABA$_A$ receptor-mediated current was measured as the outward shift in holding current*

*following application of the GABA$_A$R antagonist picrotoxin. CA1 pyramidal neurons were recorded under*

886　*whole-cell voltage-clamp at -60 mV. Effects of picrotoxin (100 μM) on holding currents recorded in*

888    *DMSO- (grey rounds) or Palbociclib-treated (red squares) hippocampal slices are represented: the shift in holding currents is shown as double-headed arrows for DMSO (grey) and Palbociclib (red). Recordings are aligned at the time of picrotoxin addition (t=0). Each point represents the mean value of holding*

890    *current at a specific time for 10 cells. C) To measure miniature inhibitory postsynaptic currents (mIPSC), 1 µM tetrodotoxin was added to block action potentials. Representative traces of mIPSC of DMSO- or*

892    *Palbociclib-treated cells are shown. E) Cumulative distribution of mIPSC amplitude from CA1 pyramidal neurons treated with either Palbociclib or control vehicle, as indicated. "n" represents number of*

894    *miniature responses recorded from nine (DMSO) or eight (Palbociclib) cells. (*p<0.001, Kolmogorov-Smirnov test).*

**wt**          **CCND1 KO**

*S3 Figure. CCND1 KO increases the maturity of dendritic spines in cortical neurons. A) Dendritic spine*

908 *analysis of WT and CCND1 KO mice cortical neurons at 20 DIV. Cultured cortical neurons were*

*transfected at 3 DIV with GFP and fixed at 20 DIV. Representative confocal images of WT and CCND1 KO*

910 *neurons. Scale bar: 10 µm. Changes in spine density and morphology were assessed by analyzing GFP-*

*positive neurons with NeuronJ (see Materials and Methods). B) Quantification of the number of total*

912 *dendritic spines relative to 10 µm of neurite, from WT (n= 59 neurons) and CCND1 KO (n= 62) neurons (4*

*mice/group). Bars show mean values ± S.E.M. ns: non-significant, Unpaired student's t-test.*