Phospholipids Diffusion on the Surface of Model Lipid Droplets

Shima Asfia[a], Ralf Seemann[a], Jean-Baptiste Fleury[a,1]

[a]*Universitat des Saarlandes, Experimental Physics and Center for Biophysics, 66123 Saarbruecken, Germany*

Abstract

Lipid droplets (LD) are suborganelles localized in the membrane of the Endoplasmic Reticulum (ER) that play an important role in metabolic functions. They consist of a core of neutral lipids surrounded by a monolayer of phosphoplipids and proteins resembling to an oil-in-water emulsion droplet. Many studies have been focused on the biophysical properties of these LDs. However, despite numerous efforts, we are lacking information on the mobility of phospholipids on the surface of LDs surface, although they may play a key role in the protein distribution. In this article, we developed a microfluidic setup that allows the formation of a triolein-buffer interface decorated with a phospholipid monolayer. Using this setup, we measured the motility of phospholipid molecules by performing Fluorescent Recovery After Photobleaching (FRAP) experiments for different lipidic compositions. The results of the FRAP measurements reveal that the motility of phospholipids is controlled by the monolayer packing decorating the interface.

Keywords: `Phospholipid, Lipid droplet, Diffusion, FRAP`

1. Introduction

Lipid droplets are suborganelles present in the Endoplasmic Reticulum (ER) membrane [1, 2]. They attract many interest due to their role in several diseases

---

*Corresponding author
Email address:* `jean-baptiste.fleury@physik.uni-saarland.de` (Jean-Baptiste Fleury)

like cancer and metabolic problems [3, 4, 5]. One of their functions is to store
lipids and to regulate their consumption based on mechanisms that are not yet
understood [6]. Lipid droplets (LDs) are composed of a core of neutral lipids and
are surrounded by a phospholipid monolayer [7, 8]. Their biogenesis is assumed
to occur in three consecutive steps [2, 9]. First, neutral lipids are synthesized in
the ER membrane. Then, they accumulate and segregate into the ER bilayer
core to form a droplet with a lens shape [2, 10, 11]. And finally, they grow
through the accumulation of lipids until budding and sometimes pinch-off from
the ER membrane [2, 12, 13, 14, 15]. Many of the biophysical properties of
LDs have been investigated over the last decade using model systems [6]. As
example, droplet interface bilayers (DiB), advanced free-standing bilayer (AFB),
and giant unilamellar vesicles (GUVs) have been extensively used to reconstitute
artificial and natural LDs in such lipid membranes [16, 17, 18, 19, 20, 21].
After successful reconstitution of LDs in a bilayer, the morphology of LDs was
studied as function of the phospholipidic composition of the DiBs, GUVs or
AFB systems [18, 19, 20]. It has been shown that the shape of LDs in a bilayer
is determined by the balance of surface forces at the contact line between the
embedded-LDs and the bilayer [18, 19, 20]. Thus, the associated contact angle
of LDs can be predicted using wetting theory [11, 22]. Accordingly, numerical
simulations based on continuum models were able to confirm many experimental
results obtained on microscopic LDs [15, 18, 23, 24, 25, 26, 27].

In addition to the equilibrium shape determined by a force balance, non-
equilibrium phenomena related to LDs have also attracted attention in recent
years. A key example is protein partitioning, which is used to understand the
dynamic distribution of specific proteins between the monolayer covering the LD
and the bilayer in which the LDs are embedded [6, 16]. Indeed, protein parti-
tioning seems highly dependent of the physical properties of the lipid monolayer
(packing, fluidity) that surrounds the LD surface [18, 17]. However, to the best
of our knowledge, there are no experimental measurements on phospholipid
properties such as motility at the LD surface. One reason for this lack of knowl-
edge could be the small size of the reconstituted LDs in the artificial systems

(DiB, AFB, GUVs), which make FRAP experiments and extraction of precise diffusion coefficients challenging [17, 28]. In consequence, there are few experimental studies in the literature that address diffusion [24]. Interestingly, this problem is not limited to experimental studies but, also, occurs in theoretical studies with numerical simulations [18, 25, 29].

In this article, we experimentally investigate the motility of phospholipids covering the interface between an aqueous and an oily phase. For this purpose, we have developed a microfluidic setup that enables the formation of a triolein-buffer interface decorated with phospholipids and have performed FRAP experiments directly at this interface (Fig. 1). Here, the planar phospholipid monolayer at the triolein-buffer interface mimics the surface of a LD. We used triolein as neutral lipid because it is an important component of the biological LD [5] and varied the molecular composition of the monolayer to assess the influence of the most common molecules that make up the LD surface in living cells. These molecules are mainly phospholipids and cholesterol. Because native biological LDs may contain cholesterol ester [30, 31], we also investigated the influence of LD-core composition on the phospholipid motility by adding some cholesterol ester (chol-ester) to the neutral lipid (triolein). We also tested the possible influence of a relevant human protein, Perilin-2 [30, 31] (PL2), on phospholipid motility with and without chol-ester in the LD core. Our measurements show that the motility of amphililic lipids is directly influenced by the packing of these molecules coating the interface.

2. Results and Discussion

Liquid droplets (LDs) originate in the ER membrane, and the phospholipid monolayer that decorates LDs is therefore composed mainly of phospholipids, sterols and proteins [33, 34, 35, 36, 37], with phospholipids making up the vast majority and proteins and sterols being present in only a small percentage. Phospholipids can be distinguished in two families, saturated and unsaturated. Accordingly, we start our discussion with the most common lipids and pre-

Figure 1: A) Sketch of the microfluidic device. B) Schematic of the phospholipid decorated triolein-buffer interface. C) Example of a FRAP measurement; the time $t = 0$ corresponds to the moment when the bleaching was stopped. D) Experimental intensity data (green dots) as obtained from a FRAP measurement plotted together with the Soumpasis equation fitted to these data (dashed line) [32].

sented the results sorted by lipid families: unsaturated phospholipids, saturated phospholipids, and sterols.

Unsaturated Phospholipids: DOPC, DOPE, N-DOPE, POPC, LPC. A relevant biophysical composition for a phospholipid monolayer covering LDs localized in the ER membrane can be approximated by a DOPC/DOPE mixture with a molar ratio between 1:1 and 3:1 [7, 8, 17, 18]. For completeness, we vary the composition from pure DOPC to a pure DOPE monolayer. The extracted diffusion constants are presented in Fig. 2 and Table 1. The diffusion constant of a pure DOPC monolayer is $D \approx 8~\mu m^2 \cdot s^{-1}$, i.e. close to, or slightly slower than, the DOPC fluidity observed in giant vesicles [28, 37]. For increasing DOPE content, the fluidity monotonically increases reaching $D \approx 17~\mu m^2 \cdot s^{-1}$ for a pure DOPE monolayer. This trend can be easily understood considering the lipid packing that results from the geometry of these two molecules. DOPC can be approximated as a cylinder, while the shape of DOPE is closer to a truncated cone [38]. Thus, the corresponding critical packing parameter $\rho = V/a_0 \cdot l$ is $\rho_{\mathrm{DOPC}} \approx 1$

Figure 2: Diffusion constants D for several saturated phospholipids (DOPC, DOPE, N-DOPE, POPC, LPC) and mixtures of DOPC and DOPE at a triolein-buffer interface extracted from FRAP experiments. Schematic representations of the various lipids are provide for PE (red head), PC (yellow head) where pink tails indicate one unsaturation.

for a monolayer consisting only of DOPC molecules and $0.5 < \rho_{\mathrm{DOPE}} < 1$ for a monolayer consisting only of DOPE molecules [38]. For a flat interface, a DOPC monolayer obviously has a higher lipid packing than a DOPE monolayer. In consequence, at equivalent surface coverage, the average motility of a single DOPE molecule is larger than that of a single DOPC molecule. Accordingly, packing density and mobility of DOPC and DOPE mixtures interpolate between the limits of a pure DOPC and DOPE monolayer, respectively. As the increased presence of lipids at the triolein-buffer interface lowers the surface tension, differences in surface coverage can also be confirmed by differences in surface tension measured by the hanging drop method as $\gamma_{\mathrm{DOPC}} \approx 1.5$ mN/m and $\gamma_{\mathrm{DOPE}} \approx 2.3$ mN/m, for DOPC and DOPE, respectively. These surface tension values are consistent with expectations based on geometry dependent lipid packing and the experimentally observed lipid motility.

To confirm the influence of the lipid packing on lipid motility, we repeated

the experiments described above with monolayers of N-DOPE, POPC and LPC molecules. Compared to DOPE, has N-DOPE an additional methyl group in the PE head, thus this molecule is slightly less conical than DOPE but also less cylindrical than DOPC (Fig. 2). The measured diffusion constant is $D \approx 17 \, \mu m^2 \cdot s^{-1}$ and surface tension is $\approx 2 \, \mathrm{mN/m}$ for a pure N-DOPE monolayer. As expected from the N-DOPE geometry, are both diffusion constant and surface tension rather comparable to pure DOPE monolayer.

POPC has the same hydroplilic head as DOPC and the same length of the hydrophobic tail, but only one unsaturation in its tail [34, 35]. Thus, due to this little difference, a POPC monolayer is expected to present a similar packing than a DOPC monolayer [27, 33, 39, 40]. And consistently with the geometry of the molecules and the expected monolayer packing, we measured the diffusion constant and the surface tension of POPC as $D \approx 7 \mu m^2 \pm 1 \cdot s^{-1}$ and $\gamma \approx 1.3 \pm 0.3 \, \mathrm{mN/m}$, that are very similar to those of a pure DOPC monolayer, see Fig. 2.

LPC is a single tail lipid with one unsaturation in its tail and a conical geometry. The measured diffusion constant for a pure LPC monolayer of $D \approx 7 \mu m^2 \cdot s^{-1}$ and its surface tension is $\gamma \approx 1.3 \, \mathrm{mN/m}$, see Fig. 2. Thus, this LPC appears to have a motility close to POPC and DOPC. This result is a little surprising, as *a priori*, we may have expected to measure a stronger difference between a monolayer composed of lipids with a single tail or a double tail. However, the correlation between low surface tension resembling high lipid packing and low lipid motility still holds true.

Saturated phospholipids (DPPC, DSPA) and Sphingolipids (SM). As described in the previous section, the surface of LDs consists mainly of lipids that may be unsaturated or saturated phospholipids [41]. In this section, we explore the motility of saturated phospholipids (DPPC, SM) on the surface of model lipid droplets. Due to the absence of unsaturations, the packing of unsaturated (neutral) phospholipids may be expected to be denser than that of unsaturated phospholipids and so the corresponding motility is expected to be lower than

that of DOPC. However, the degree of saturation is not the only interaction that controls the monolayer packing. As example, electrostatic interactions can also strongly influence the packing of lipid monolayers. For this reason, we also explore below the mixture between saturated DSPA molecules having an negatively charged hydrophilic head group and uncharged and unsaturated DOPC molecules. In addition to saturated lipids, we will also consider the case of sphingolipids.

DPPC is a saturated lipid with a cylindrical shape that self-assembles in an highly packed monolayer [42]. The measured diffusion constant for a pure DPPC monolayer is $D \approx 5.5 \ \mu\text{m}^2 \cdot \text{s}^{-1}$, which is a little slower to that of unsaturated DOPC and POPC phospholipids, that are also expected to be densely packed (see, Fig. 3). The high DPPC packing at the triolein-buffer interface is also confirmed by the low surface tension, which is $\gamma \approx 1.1$ mN/m, also very similar to surface tensions values for DOPC and POPC. In addition, DPPC can also condense at room temperature, which is another reason for the reduced fluidity [43].

DSPA is a saturated double-tail lipid with a head that is negatively charged and with a conical shape. Due to electrostatic repulsion, the lipid packing of a pure PA monolayer is smaller than that of other lipids [44]. In addition to a large motility of $D \approx 20 \ \mu m^2 \times s^{-1}$ of a pure DSPA monolayer, the comparatively low packing density also leads to a comparatively large surface tension at the triolein-buffer interface, which was measured to be $D \approx 2.6$ mN/m, see table 1. Mixtures of DOPC and DSPA molecules show motilities intermediate between those of a pure DOPC and a pure DSPA monolayer. Presumably due to the repulsion of the charged DSPA molecules, the measured diffusion constants are already quite large at a DSPA content of 25% and increase even further with increasing DSPA content, see Fig. 3.

Sphingolipids are a family of lipids that are synthesized in the (ER) membrane, where they can be found in low amounts, as well as at the trans Golgi [45, 46, 47]. Due to their possible role of sphingolipids in the LDs biogenesis, they have attracted a growing attention on recent years [45, 46, 47]. Sphingomyelin

Figure 3: Diffusion constants D for unsaturated phospholipids (DSPA, DPPC, and SM) and mixtures of DOPC and DSPA at a triolein/buffer interphase extracted from FRAP experiments in comparison to values from DOPC. A schematic representation is provide for PC lipids (red head), SM (green head) and the charged PA (light blue head).

(SM) has a PC head with a fatty acid group and a tail that has a single satura-
tion in contact with the head. Thus, we study this lipids with other saturated
lipids. SM molecule has a geometry close to a cylindrical molecule. The rela-
tively low diffusion constant of a pure SM monolayer of $D \approx 8 \ \mu m^2 \cdot s^{-1}$, and the
corresponding low surface tension $\gamma \approx 1.2$ mN/m, Fig.3, indicate that despite of
the slightly conical shape of the SM, the associated lipid packing is dense, similar
to an assembly of cylindrical molecules, in agreement with literature [38, 48].

Sterol, Sterol ester and Perilipin-2. Sterols like cholesterol (chol) and cholesteryl
ester (chol ester) are also important components of LDs [7, 8]. However, these
molecules differ from previous lipids, as they are unable to form a lipid bilayer.
As example, chol-ester is not a surface active molecule, it is a component of the
LD core and not of the LD surface. Nevertheless, these sterols are important
lipids and it is interesting to investigate their influence on the phospolipids
monoloayer properties. Besides, we also explore the effect of a LD protein
(PLP-2) on the diffusion properties of phospholipids.

Cholesterol (Chol) is present in the ER membrane only in small amounts,
namely below 6 % percent. Nevertheless, Chol may play a role in metabolic
diseases that are involving LDs [4, 49]. Chol drastically alters the biophysical
properties of the plasma membrane and thus, it is interesting to investigate the
case of a lipid monolayer enriched with Chol. It is known that Chol increases
both the bilayer tension and the bilayer fluidity, at least at low temperatures [50,
51, 52]. The measured surface tension for a DOPC monolayer with 10% Chol is
≈ 1.7 mN/m, while the surface tension of a DOPE monolayer with 10% Chol
is ≈ 2.5 mN/m, i.e. the surface tension for both phospholipids are increased
by about 10% compared to pure DOPC or DOPE monolayer, respectively, see
table 1. The observed increase in surface tension indicates a lower packing
density of lipids, which is expected to increase lipid diffusion. In fact, the
measured diffusion constant for a DOPC monolayer with 10% Chol gives a
value of $\approx 10 \ \mu m^2 \cdot s^{-1}$, while that of a DOPE monolayer with 10% Chol gives
$\approx 20 \ \mu m^2 \cdot s^{-1}$, see Fig. 4, i.e. the diffusion constant also increased by about

Figure 4: Diffusion constants D extracted from FRAP experiments for cholesterol containing DOPC, DOPE, POPC and DPPC monolayer at triolein-buffer interface, and DOPC and DOPE monolayer at cholesterol-ester containing triolein-buffer interface with and without PLP2. Schematic representation is provide for cholesterol and DOPC.

10% compared to pure DOPC or DOPE monolayer. This suggests that the increase in surface tension and lipid diffusion is indeed due to decreased lipid packing.

Cholesteryl ester, a dietary lipid, is an ester of cholesterol found in LDs [53]. This molecule differs from the previous molecules in that it is not localized at the surface of LDs but it is only present in the LD core. This is interesting because cholesterol ester appears to affect the recruitment of some LD proteins, such as PLP2, when they enrich the LD composition [30, 31]. The surface tension of a DOPC monolayer at the interface between triolein-buffer with 15% cholesteryl ester in triolein was measured to be $\gamma \approx 1.5$ mN/m, i.e. the same value as for triolein not containing cholesteryl ester. Similarly, we also measured the surface tension for a DOPE monolayer to be ≈ 2.3 mN/m no matter if the triolein contains 15% cholesteryl ester or not. The corresponding measured diffusion constant yields a value of $D \approx 8\mu m^2 \cdot s^{-1}$ for a DOPC monolayer with 15% cholesteryl ester in triolein, and a diffusion constant $D \approx 17\mu m^2 \cdot s^{-1}$ for DOPE monolayer with 15% cholesteryl ester in triolein, see Fig. 4. From this, we can conclude that cholesteryl ester molecules remain in the triolein phase and hardly change the surface properties of the lipid decorated triolein-buffer interface. Similarly, we observe no measurable change in the surface tension of the phospholipid-decorated triolein-buffer interface or in the diffusion constant compared with a pure DOPC or a pure DOPE monolayer, respectively, when the aqueous buffer contains 1 μM of PLP2, regardless of whether the triolein phase contains 15% cholesteryl ester or not. Thus, either the lipid packing is hardly affected by the proteins or their concentration at the LD surface was not high enough to produce a notable effect, cf. Fig. 4.

3. Conclusion

We investigated the diffusion properties of lipids on the surface of model lipid droplets. For that, we produced a model lipid droplet interface made of a triolein-buffer interface in a microfluidic chamber that is decorated with a

Table 1: Table summarizing the measured surface tension values γ and diffusion constants D for triolein-buffer interfaces decorated with monolayers made of a single lipidic component. More results including lipid mixtures are available in Figs. 2,3 & 4.

Lipids	DOPC	DOPE	POPC	DPPC	DSPA	SM
γ (mN/m)	1.5±0.3	2.3±0.3	1.3±0.3	1.1±0.3	2.6±0.3	1.2±0.3
D (μm^2/s)	8.1±1	17.0±1	6.8±1	5.4 ±1	20.2±1	7.9 ±1

lipid monolayer of variable composition. FRAP experiments were conducted directly at this interface and the corresponding diffusion constant for the lipids was obtained as function of the molecular composition of the lipid monolayer. Based on geometrical (cylindrical, conical) and chemical (electrostatic charge, saturation) properties of the respective lipids, we argued that the lipid packing is the key parameter to understand the measured results. In this context, monolayers with high lipid packing have low lipid motility. And monolayers with low lipid packing have high lipid motility. This reasoning was supported by surface tension measurements as high lipid packing means low interfacial tension and low lipid packing means high interfacial tension. By extrapolation, this study provides an estimate of defect density on the LD surface for the lipids composition tested. This point is important because defects seem to play a key role in protein partitioning on LD surface [17].

4. Methods

Molecules. For the preparation of molecular monolayer that cover the model lipid droplets, i.e. the triolein-buffer interface, the following molecules were purchased from Avanti Polar Lipids: 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), 1,2-distearoyl-sn-glycero-3-phosphate (sodium salt) (DSPA), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), 1-(10Z-heptadecenoyl)-2-hydroxy-sn-glycero-3-phosphocholine (LPC), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), 1,2-dioleoyl-sn-glycero-

3-phosphoethanolamine-N-methyl (N-DOPE), cholesterol (Chol), N-stearoyl-D-erythro-sphingosylphosphorylcholine (SM), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), Cholesteryl ester (Chol ester). Glyceryl trioleate (triolein) and squalene were purchased from Sigma-Aldrich. Recombinant human denaturtared perilipin 2 protein (PL2) was purchased from Abcam (ab181932). (FITC-DOPE) 1,2-dioleoyl-sn-glycerol-phospho-ethanolamine Fluorescein was purchased from ChemCruz (Santa Cruz Biotechnology).

Interfacial Tension Measurements. The interfacial tension of phospholipid decorated triolein-buffer interfaces was determined by the pendant drop method using a contact angle measurement device OCA 20 (DataPhysics, Germany). For this method, a transparent cuvette is filled with the lower density fluid, here the triolein-phospholipid mixture ($\rho_{oil} = 0.91$ g/cm^3) and a pendant drop containing a defined volume of the denser fluid, here the buffer solution ($\rho_{buffer} = 0.998$ g/cm^3), is produced from a hollow needle immersed in this liquid. While gravity drags the droplet down, buoyancy and surface forces keep the droplet in place. Based on a contour fit to the pendant droplet, the software SCA 20 extracts both size and shape of the droplet and automatically computes the corresponding interfacial tension. This procedure is most reliable when the droplet size is close to detachment from the needle, and we thus set the droplet volume to (2.0 ± 0.5) μl.

μChip Fabrication and Monolayer Formation. Using computer software (Autocad), we created a drawing in the form of a cylinder with 3 mm diameter, which was produced using a 3D printer. This cylinder was positioned in the center of a glass petri dish. After setting up the mold, mixed and degassed PDMS (Sylgard 184 – Dow Corning) was poured onto the mold with a resulting layer thickness of about 3 mm so that the cylinder is not fully immersed in the PDMS. After curing the PDMS at 60 °C for 6-7 hours, the cylinder was removed and square pieces with the cylindrical hole in the center were cut from the PDMS and removed from the petri dish. The final device was fabricated by attaching the square PDMS piece to a PDMS coated glass cover slip (Sylgard 184 spin

coated for 5 min at 3000 rpm) using plasma bonding (Diener electronics). The thus fabricated device was placed it on a hot plate at 60 °C for 1 hour to achieve higher bonding strength.

To produce the triolein-lipid mixture, a specific amount of lipid was dissolved in Ethanol and mixed with 8% DOPE-FITC (dissolved in chloroform). This solution was then dried under vacuum. Then triolein was added to the dried components and mixed in an ultrasonic bath.

To form the monolayer, 6 μl of the as-prepared triolein-lipid mixture was first placed in the cylindrical opening of the PDMS micro chip, and then 20 μl of 150 M KCl buffer was added on top. Within seconds, a phospholipid monolayer forms at the triolein-buffer interface, with a composition similar to the lipid composition of the originally prepared triolein solution.

FRAP Experiments. The mobility of phospholipid monolayers at triolein-buffer interfaces were determined by fluorescence recovery after photobleaching (FRAP) measurements using an inverted Zeiss microscope, Axio Observer Z7. FITC-DOPE (495 nm/515 nm) is used as a dye for our FRAP experiments due to its very low photostability. For the FRAP measurements, a pinhole (diameter of 150 μm - Edmund Optics) was placed between the microscope tray and the bottom of the microfluidic device. Once the lipid monolayer is formed, the microscope is focused on the center of the monolayer using a $40\times$ objective and blue LED light (469/38 nm) is emitted continuously for 30 to 60 s at maximum power (zeiss colibri 7) to bleach a spot with a diameter of about 150 μm. Subsequently, the pinhole is removed and the $40\times$ objective is changed to a $10\times$ objective and fluorescence recovery is recorded with a frame rate of 1 s using the same blue wavelength excitation but now with the lowest illumination power ($\approx 1\%$). The fluorescence recovery, i.e. the fluorescent intensity of the bleached area was extracted automatically using the software ImageJ. The diffusion constants were obtained via a Soumpasis fit [32] to the intensity data using Origin8 (OriginLab). For information, we choose to performed a maximum of 2 FRAP experiments by phospholipid monolayer. After two measurements, we clean the

chip (ethanol, acetone) and replace the chemical to form a fresh triolein-buffer interface.

References

[1] Y. Guo, K. R. Cordes, R. V. Farese, T. C. Walther, Lipid droplets at a glance, J Cell Sci 122 (6) (2009) 749–752, publisher: The Company of Biologists Ltd Section: Cell Science at a Glance. doi:10.1242/jcs.037630.
URL https://jcs.biologists.org/content/122/6/749

[2] T. C. Walther, J. Chung, R. V. Farese, Lipid Droplet Biogenesis, Annual Review of Cell and Developmental Biology 33 (1) (2017) 491–510, _eprint: https://doi.org/10.1146/annurev-cellbio-100616-060608. doi:10.1146/annurev-cellbio-100616-060608.
URL https://doi.org/10.1146/annurev-cellbio-100616-060608

[3] P. T. Bozza, J. P. B. Viola, Lipid droplets in inflammation and cancer, Prostaglandins, Leukotrienes and Essential Fatty Acids (PLEFA) 82 (4) (2010) 243–250. doi:10.1016/j.plefa.2010.02.005.
URL https://www.sciencedirect.com/science/article/pii/S0952327810000499

[4] G. Onal, O. Kutlu, D. Gozuacik, S. Dokmeci Emre, Lipid Droplets in Health and Disease, Lipids in Health and Disease 16 (1) (2017) 128. doi:10.1186/s12944-017-0521-7.
URL https://doi.org/10.1186/s12944-017-0521-7

[5] S. Xu, X. Zhang, P. Liu, Lipid droplet proteins and metabolic diseases, Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease 1864 (5, Part B) (2018) 1968–1983. doi:10.1016/j.bbadis.2017.07.019.
URL https://www.sciencedirect.com/science/article/pii/S0925443917302454

[6] R. Dhiman, S. Caesar, A. R. Thiam, B. Schrul, Mechanisms of protein targeting to lipid droplets: A unified cell biological and biophysical perspective, Semin. Cell Dev. Biol.`doi:10.1016/j.semcdb.2020.03.004`.

[7] Y. Guo, T. C. Walther, M. Rao, N. Stuurman, G. Goshima, K. Terayama, J. S. Wong, R. D. Vale, P. Walter, R. V. Farese, Functional genomic screen reveals genes involved in lipid-droplet formation and utilization, Nature 453 (7195) (2008) 657–661. `doi:10.1038/nature06928`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2734507/`

[8] M. A. Welte, A. P. Gould, Lipid droplet functions beyond energy storage, Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids 1862 (10, Part B) (2017) 1260–1272. `doi:10.1016/j.bbalip.2017.07.006`.
URL `http://www.sciencedirect.com/science/article/pii/S1388198117301373`

[9] J. A. Olzmann, P. Carvalho, Dynamics and functions of lipid droplets, Nat Rev Mol Cell Biol 20 (3) (2019) 137–155. `doi:10.1038/s41580-018-0085-z`.
URL `http://www.nature.com/articles/s41580-018-0085-z`

[10] H. Khandelia, L. Duelund, K. I. Pakkanen, J. H. Ipsen, Triglyceride Blisters in Lipid Bilayers: Implications for Lipid Droplet Biogenesis and the Mobile Lipid Signal in Cancer Cell Membranes, PLOS ONE 5 (9) (2010) e12811, publisher: Public Library of Science. `doi:10.1371/journal.pone.0012811`.
URL `https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0012811`

[11] M. J. Greenall, C. M. Marques, Hydrophobic droplets in amphiphilic bilayers: a coarse-grained mean-field theory study, Soft Matter 8 (12) (2012) 3308. `doi:10.1039/c2sm07193b`.
URL `http://xlink.rsc.org/?DOI=c2sm07193b`

[12] K. D. Chapman, J. M. Dyer, R. T. Mullen, Biogenesis and functions of lipid droplets in plants, J Lipid Res 53 (2) (2012) 215–226. `doi:10.1194/jlr.R021436`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3269164/`

[13] V. Choudhary, N. Ojha, A. Golden, W. A. Prinz, A conserved family of proteins facilitates nascent lipid droplet budding from the ER, J Cell Biol 211 (2) (2015) 261–271, publisher: The Rockefeller University Press. `doi:10.1083/jcb.201505067`.
URL `https://rupress.org/jcb/article/211/2/261/38599/A-conserved-family-of-proteins-facilitates-nascent`

[14] A. Masedunskas, Y. Chen, R. Stussman, R. Weigert, I. H. Mather, Kinetics of milk lipid droplet transport, growth, and secretion revealed by intravital imaging: lipid droplet release is intermittently stimulated by oxytocin, MBoC 28 (7) (2017) 935–946, publisher: American Society for Cell Biology (mboc). `doi:10.1091/mbc.e16-11-0776`.
URL `https://www.molbiolcell.org/doi/10.1091/mbc.e16-11-0776`

[15] V. Zoni, V. Nieto, L. J. Endter, H. J. Risselada, L. Monticelli, S. Vanni, To Bud or Not to Bud: A Perspective on Molecular Simulations of Lipid Droplet Budding, Front. Mol. Biosci. 6, publisher: Frontiers. `doi:10.3389/fmolb.2019.00124`.
URL `https://www.frontiersin.org/articles/10.3389/fmolb.2019.00124/full`

[16] A. R. Thiam, I. Dugail, Lipid droplet–membrane contact sites – from protein binding to function, J Cell Sci 132 (12), publisher: The Company of Biologists Ltd Section: Review. `doi:10.1242/jcs.230169`.
URL `https://jcs.biologists.org/content/132/12/jcs230169`

[17] L. Caillon, V. Nieto, P. Gehan, M. Omrane, N. Rodriguez, L. Monticelli, A. R. Thiam, Triacylglycerols sequester monotopic membrane proteins to

lipid droplets, Nature Communications 11 (1) (2020) 3944, number: 1 Publisher: Nature Publishing Group. `doi:10.1038/s41467-020-17585-8`.
URL `https://www.nature.com/articles/s41467-020-17585-8`

[18] S. Puza, S. Caesar, C. Poojari, M. Jung, R. Seemann, J. S. Hub, B. Schrul, J.-B. Fleury, Lipid Droplets Embedded in a Model Cell Membrane Create a Phospholipid Diffusion Barrier, Small n/a (n/a) 2106524, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/smll.202106524. `doi:10.1002/smll.202106524`.
URL `https://onlinelibrary.wiley.com/doi/abs/10.1002/smll.202106524`

[19] A. Chorlay, A. R. Thiam, An Asymmetry in Monolayer Tension Regulates Lipid Droplet Budding Direction, Biophysical Journal 114 (3) (2018) 631–640, publisher: Elsevier. `doi:10.1016/j.bpj.2017.12.014`.
URL `https://www.cell.com/biophysj/abstract/S0006-3495(17)35094-4`

[20] A. Chorlay, L. Monticelli, J. V. Ferreira, K. B. M'barek, D. Ajjaji, S. Wang, E. Johnson, R. Beck, M. Omrane, M. Beller, P. Carvalho, A. R. Thiam, Membrane Asymmetry Imposes Directionality on Lipid Droplet Emergence from the ER, Developmental Cell 50 (1) (2019) 25–42.e7, publisher: Elsevier. `doi:10.1016/j.devcel.2019.05.003`.
URL `https://www.cell.com/developmental-cell/abstract/S1534-5807(19)30374-0`

[21] A. Chorlay, A. Santinho, A. R. Thiam, Making Droplet-Embedded Vesicles to Model Cellular Lipid Droplets, STAR Protocols 1 (3) (2020) 100116. `doi:10.1016/j.xpro.2020.100116`.
URL `https://www.sciencedirect.com/science/article/pii/S2666166720301039`

[22] P. G. de Gennes, Wetting: statics and dynamics, Rev. Mod. Phys. 57 (3) (1985) 827–863, publisher: American Physical Society. `doi:10.1103/`

RevModPhys.57.827.

URL https://link.aps.org/doi/10.1103/RevModPhys.57.827

[23] V. Zoni, R. Khaddaj, P. Campomanes, A. R. Thiam, R. Schneiter, S. Vanni, Pre-existing bilayer stresses modulate triglyceride accumulation in the ER versus lipid droplets, eLife 10 (2021) e62886, publisher: eLife Sciences Publications, Ltd. doi:10.7554/eLife.62886.
URL https://doi.org/10.7554/eLife.62886

[24] A. Bacle, R. Gautier, C. L. Jackson, P. F. J. Fuchs, S. Vanni, Interdigitation between Triglycerides and Lipids Modulates Surface Properties of Lipid Droplets, Biophysical Journal 112 (7) (2017) 1417–1430. doi:10.1016/j.bpj.2017.02.032.
URL https://www.sciencedirect.com/science/article/pii/S0006349517302461

[25] G. Bussi, D. Donadio, M. Parrinello, Canonical sampling through velocity rescaling, J Chem Phys 126 (1) (2007) 014101. doi:10.1063/1.2408420.

[26] M. J. Abraham, T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, E. Lindahl, GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers, SoftwareX 1-2 (2015) 19–25. doi:10.1016/j.softx.2015.06.001.
URL https://www.sciencedirect.com/science/article/pii/S2352711015000059

[27] L. Vamparys, R. Gautier, S. Vanni, W. F. D. Bennett, D. P. Tieleman, B. Antonny, C. Etchebest, P. F. J. Fuchs, Conical Lipids in Flat Bilayers Induce Packing Defects Similar to that Induced by Positive Curvature, Biophysical Journal 104 (3) (2013) 585–593. doi:10.1016/j.bpj.2012.11.3836.
URL https://www.sciencedirect.com/science/article/pii/S0006349512051429

[28] P. Heo, S. Ramakrishnan, J. Coleman, J. E. Rothman, J.-B. Fleury, F. Pincet, Highly Reproducible Physiological Asymmetric Membrane with Freely Diffusing Embedded Proteins in a 3D-Printed Microfluidic Setup, Small 15 (21) (2019) 1900725, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/smll.201900725. `doi:10.1002/smll.201900725`.
URL `https://www.onlinelibrary.wiley.com/doi/abs/10.1002/smll.201900725`

[29] C. Tempra, O. H. S. Ollila, M. Javanainen, Accurate Simulations of Lipid Monolayers Require a Water Model With Correct Surface Tension`doi:10.26434/chemrxiv-2021-ws71h`.
URL `https://chemrxiv.org/engage/chemrxiv/article-details/6152338087a02d9ab04af36a`

[30] Y. Li, P. Khanal, F. Norheim, M. Hjorth, T. Bjellaas, C. A. Drevon, J. Vaage, A. R. Kimmel, K. T. Dalen, Plin2 deletion increases cholesteryl ester lipid droplet content and disturbs cholesterol balance in adrenal cortex, Journal of Lipid Research 62 (2021) 100048. `doi:10.1016/j.jlr.2021.100048`.
URL `https://www.sciencedirect.com/science/article/pii/S0022227521000286`

[31] K. Hsieh, Y. K. Lee, C. Londos, B. M. Raaka, K. T. Dalen, A. R. Kimmel, Perilipin family members preferentially sequester to either triacylglycerol-specific or cholesteryl-ester-specific intracellular lipid storage droplets, J Cell Sci 125 (17) (2012) 4067–4076. `doi:10.1242/jcs.104943`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3482316/`

[32] D. M. Soumpasis, Theoretical analysis of fluorescence photobleaching recovery experiments., Biophys J 41 (1) (1983) 95–97.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1329018/`

[33] P. J. Quinn, F. Joo, L. Vigh, The role of unsaturated lipids in membrane

structure and stability, Prog Biophys Mol Biol 53 (2) (1989) 71–103. `doi: 10.1016/0079-6107(89)90015-1`.

[34] N. Kučerka, F. A. Heberle, J. Pan, J. Katsaras, Structural Significance of Lipid Diversity as Studied by Small Angle Neutron and X-ray Scattering, Membranes 5 (3) (2015) 454–472, number: 3 Publisher: Multidisciplinary Digital Publishing Institute. `doi:10.3390/membranes5030454`.
URL `https://www.mdpi.com/2077-0375/5/3/454`

[35] W. Ding, M. Palaiokostas, G. Shahane, W. Wang, M. Orsi, Effects of High Pressure on Phospholipid Bilayers, J. Phys. Chem. B 121 (41) (2017) 9597–9606, publisher: American Chemical Society. `doi:10.1021/acs.jpcb. 7b07119`.
URL `https://doi.org/10.1021/acs.jpcb.7b07119`

[36] G. Onal, O. Kutlu, D. Gozuacik, S. Dokmeci Emre, Lipid Droplets in Health and Disease, Lipids Health Dis 16 (1) (2017) 128. `doi:10.1186/s12944-017-0521-7`.
URL `http://lipidworld.biomedcentral.com/articles/10.1186/s12944-017-0521-7`

[37] R. Dimova, C. Marques, C. Marques, The Giant Vesicle Book, CRC Press, 2019. `doi:10.1201/9781315152516`.
URL `https://www.taylorfrancis.com/books/e/9781315152516`

[38] J. N. Israelachvili, Intermolecular and Surface Forces, 3rd Edition, Academic Press Inc, Burlington (Mass.), 2011.

[39] M. Raja, Do Small Headgroups of Phosphatidylethanolamine and Phosphatidic Acid Lead to a Similar Folding Pattern of the K+ Channel?, J Membr Biol 242 (3) (2011) 137–143. `doi:10.1007/s00232-011-9384-4`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3146712/`

[40] G. Cevc, Phospholipids Handbook, CRC Press, 1993, google-Books-ID: 2ZxaucDNWHcC.

[41] The ins and outs of endoplasmic reticulum-controlled lipid biosynthesis, EMBO reports 18 (11) (2017) 1905–1921, publisher: John Wiley & Sons, Ltd. `doi:10.15252/embr.201643426`.
URL `https://www.embopress.org/doi/full/10.15252/embr.201643426`

[42] L. Qiao, A. Ge, Y. Liang, S. Ye, Oxidative Degradation of the Monolayer of 1-Palmitoyl-2-Oleoyl-sn-Glycero-3-Phosphocholine (POPC) in Low-Level Ozone, J. Phys. Chem. B 119 (44) (2015) 14188–14199, publisher: American Chemical Society. `doi:10.1021/acs.jpcb.5b08985`.
URL `https://doi.org/10.1021/acs.jpcb.5b08985`

[43] K. Nag, J. Perez-Gil, A. Cruz, K. M. Keough, Fluorescently labeled pulmonary surfactant protein C in spread phospholipid monolayers, Biophys J 71 (1) (1996) 246–256. `doi:10.1016/S0006-3495(96)79221-4`.

[44] W. Kulig, H. Korolainen, M. Zatorska, U. Kwolek, P. Wydro, M. Kepczynski, T. Róg, Complex Behavior of Phosphatidylcholine–Phosphatidic Acid Bilayers and Monolayers: Effect of Acyl Chain Unsaturation, Langmuir 35 (17) (2019) 5944–5956, publisher: American Chemical Society. `doi:10.1021/acs.langmuir.9b00381`.
URL `https://doi.org/10.1021/acs.langmuir.9b00381`

[45] A. L. McIntosh, S. M. Storey, B. P. Atshaves, Intracellular Lipid Droplets Contain Dynamic Pools of Sphingomyelin: ADRP Binds Phospholipids with High Affinity, Lipids 45 (6) (2010) 465–477. `doi:10.1007/s11745-010-3424-1`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3065392/`

[46] D. K. Breslow, Sphingolipid Homeostasis in the Endoplasmic Reticulum and Beyond, Cold Spring Harb Perspect Biol 5 (4) (2013) a013326. `doi:10.1101/cshperspect.a013326`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3683901/`

[47] G. M. Deevska, M. N. Nikolova-Karakashian, The expanding role of sph-
ingolipids in lipid droplet biogenesis, Biochim Biophys Acta Mol Cell Biol
Lipids 1862 (10 Pt B) (2017) 1155–1165. `doi:10.1016/j.bbalip.2017.`
`07.008`.

[48] F. M. Goñi, A. Alonso, Biophysics of sphingolipids I. Membrane properties
of sphingosine, ceramides and other simple sphingolipids, Biochimica et
Biophysica Acta (BBA) - Biomembranes 1758 (12) (2006) 1902–1921.
`doi:10.1016/j.bbamem.2006.09.011`.
URL `https://www.sciencedirect.com/science/article/pii/`
`S0005273606003579`

[49] D. Casares, P. V. Escribá, C. A. Rosselló, Membrane Lipid Composition:
Effect on Membrane and Organelle Structure, Function and Compartmen-
talization and Therapeutic Avenues, Int J Mol Sci 20 (9) (2019) 2167.
`doi:10.3390/ijms20092167`.
URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6540057/`

[50] Y. Guo, M. Werner, R. Seemann, V. A. Baulin, J.-B. Fleury, Tension-
Induced Translocation of an Ultrashort Carbon Nanotube through a Phos-
pholipid Bilayer, ACS Nano 12 (12) (2018) 12042–12049, publisher: Amer-
ican Chemical Society. `doi:10.1021/acsnano.8b04657`.
URL `https://doi.org/10.1021/acsnano.8b04657`

[51] F. Lolicato, R. Saleppico, A. Griffo, B. Pokrandt, H.-M. Müller, H. Ewers,
H. Hähl, J.-B. Fleury, R. Seemann, B. Brügger, K. Jacobs, I. Vattulainen,
W. Nickel, Cholesterol promotes both head group visibility and clustering
of PI(4,5)P2 driving unconventional secretion of Fibroblast Growth Factor
2, Tech. rep., bioRxiv, section: New Results Type: article (Apr. 2021).
`doi:10.1101/2021.04.16.440132`.
URL `https://www.biorxiv.org/content/10.1101/2021.04.16.`
`440132v1`

[52] The Effect of Cholesterol on the Lateral Diffusion of Phospholipids in Oriented Bilayers | Elsevier Enhanced Reader. doi:10.1016/S0006-3495(03)70033-2.

URL https://reader.elsevier.com/reader/sd/pii/S0006349503700332?token=D174135284A6750C4C8C01396F72C8C4681860FF9AAC469D6C4BDD009DC31A8F

[53] G. R. Phillips, S. E. Hancock, S. H. J. Brown, A. M. Jenner, F. Kreilaus, K. A. Newell, T. W. Mitchell, Cholesteryl ester levels are elevated in the caudate and putamen of Huntington's disease patients, Sci Rep 10 (2020) 20314. doi:10.1038/s41598-020-76973-8.

URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7680097/

24