Quantifying the movement, behavior, and environmental context of group-living animals using drones and computer vision

Benjamin Koger[1,2,3], Adwait Deshpande[1,2,3], Jeffrey T. Kerby[4,5,6], Jacob M. Graving[7,1,2,3], Blair R. Costelloe[1,2,3]#, Iain D. Couzin[1,2,3]#

1. Department of Collective Behavior, Max Planck Institute of Animal Behavior, Konstanz, Germany
2. Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany.
3. Department of Biology, University of Konstanz, Konstanz, Germany.
4. Aarhus Institute of Advanced Studies, Aarhus University, Aarhus, Denmark.
5. Neukom Institute for Computational Science, Dartmouth College, Hanover, USA.
6. Section for Ecoinformatics and Biodiversity, Department of Biology, Aarhus University, Aarhus, Denmark.
7. Advanced Research Technology Unit, Max Planck Institute of Animal Behavior, Konstanz, Germany

Co-senior authors

Corresponding authors: B. Koger (benkoger@gmail.com) and I.D. Couzin (icouzin@ab.mpg.de)

Keywords: behavioral tracking, computer vision, drones, environmental reconstruction, gelada monkey, pose, posture, video analysis, wildlife, zebra,

Abstract

1. Methods for collecting animal behavior data in natural environments, such as direct

observation and bio-logging, are typically limited in spatiotemporal resolution, the

number of animals that can be observed, and information about animals' social and

physical environments.

2. Video imagery can capture rich information about animals and their environments,

but image-based approaches are often impractical due to the challenges of

processing large and complex multi-image datasets and transforming resulting data,

such as animals' locations, into geographic coordinates.

3. We demonstrate a new system for studying behavior in the wild that uses drone-

recorded videos and computer vision approaches to automatically track the location

and body posture of free-roaming animals in georeferenced coordinates with high

spatiotemporal resolution embedded in contemporaneous 3D landscape models of

the surrounding area.

4. We provide two worked examples in which we apply this approach to videos of

gelada monkeys and multiple species of group-living African ungulates. We

demonstrate how to track multiple animals simultaneously, classify individuals by

species and age-sex class, estimate individuals' body postures (poses), and extract

environmental features, including topography of the landscape and animal trails.

5. By quantifying animal movement and posture, while simultaneously reconstructing a

detailed 3D model of the landscape, our approach opens the door to studying the

sensory ecology and decision-making of animals within their natural physical and

social environments.

Introduction

Studying animals in the wild is essential for understanding how environments shape behavior, and how behavior affects individual fitness. The impacts of animals' behaviors also extend beyond the individual to broader ecological scales: the cumulative behavior of many interacting individuals underpins emergent group-level behaviors such as collective sensing (Berdahl et al., 2013), migration (Flack et al., 2018), and self-assembly (Lutz et al., 2021); mediates population growth and decline (Bro-Jørgensen et al., 2019); and drives ecological processes such as seed dispersal, pollination, and trophic cascades (Atkins et al., 2019; Costa-Pereira et al., 2022). While conservation initiatives commonly focus on preserving phenomena at these broader scales (Bro-Jørgensen et al., 2019) a mechanistic understanding of the individual-level animal behaviors that drive such phenomena is often critical to the success of conservation efforts, including habitat restoration (Hale et al., 2020), wildlife translocation (Berger-Tal et al., 2020), preservation of ecosystem services (Mortelliti, 2022), and human-wildlife conflict interventions (Pekarsky et al., 2021). Thus, the on-going biodiversity crisis (Ceballos et al., 2020) has created an urgent need to understand the drivers and diversity of animal behavioral patterns and how these are influenced by anthropogenic impacts (Buchholz et al., 2019).

Connecting individual behavior to broader scale phenomena requires detailed data on the behavior of large numbers of animals, as well as their interactions with social partners and their biotic and abiotic environments (Costa-Pereira et al., 2022). For example, understanding the collective movement decisions of animal groups requires data on individual movements and how these relate to the concurrent actions of social partners and salient environmental features (Strandburg-Peshkin et al., 2017). Likewise, predicting the resilience of animal-mediated seed dispersal to environmental change requires knowledge of the foraging preferences and movement patterns of dispersers and how these behaviors vary between individuals and across environmental conditions (Mortelliti, 2022; Pollux,

2017). The challenge of collecting such detailed and multifaceted datasets constitutes a

major barrier to integrating behavioral ecology – and its individual-level focus – into the

broader fields of ecology and conservation, where practitioners primarily rely on species or

populations as the units of behavioral analysis (Bro-Jørgensen et al., 2019; Costa-Pereira et

al., 2022). The rapid development of animal-borne data-logging devices (known as bio-

loggers) in recent decades has transformed our ability to collect high-resolution data on

individual animals (Kays et al., 2015), and has fueled calls to incorporate within-species

variation into conservation decisions and investigations of ecological processes (Bro-

Jørgensen et al., 2019; Mortelliti, 2022; Shaw, 2020). However, beyond the normal

constraints of biologging (Hughey et al., 2018), integrating bio-logging data from individual

animals with the concurrent social and ecological context remains a significant challenge

(Costa-Pereira et al., 2022; Hughey et al., 2018).

Aerial video-based observation, in combination with machine learning-based image

processing tools, is emerging as a promising approach for generating the high-resolution

movement datasets necessary for quantitative, multi-scale studies of wildlife behavior

(Hughey et al., 2018; Tuia et al., 2022). Image-based tracking methods, reviewed in Dell et

al. (2014) in which the location of multiple individuals (Walter & Couzin, 2021) and possibly

individual body parts (Graving et al., 2019; Mathis et al., 2018; Nath et al., 2019; Pereira et al.,

2020) are tracked over the course of a video observation, originated in laboratory settings

where overhead cameras enabled continuous monitoring with minimal visual occlusions.

Crucially, controlled recording conditions enabled the use of automated tracking solutions

making large-scale video-based studies tractable. In recent years, attempts have been

made to replicate this approach in field settings, often by using drones to collect aerial

imagery of study subjects. Initially, complex visual environments and a lack of controlled

lighting and filming conditions prevented automated image-based tracking, leading

researchers to rely on alternative approaches, such as manually extracting data from videos

(Sprogis et al., 2020) analyzing a subset of still frames (Inoue et al., 2019) or approximating

animal movement paths using the drone's position (Raoult et al., 2018). However, recent

developments in deep learning based approaches (Zhao et al., 2019), have simplified the

automatic detection of objects in complex scenes, enabling detection of animals and

possibly tracking of their movement and body posture in imagery captured with drones

(Corcoran et al., 2021) as well as camera traps, smartphones, and thermal, infrared and

conventional cameras (Tuia et al., 2022) .

An unresolved challenge is the projection of animal locations throughout an

observation from the pixel coordinates of the camera to geographic coordinates and

standard units of distance (Haalck et al., 2020), a task that is complicated by the complex

topography of natural landscapes and the movement of drone-mounted cameras. Multi-

camera 3D imaging can surmount these issues, but such systems are logistically and

computationally difficult to deploy (Francisco et al., 2020). Instead, previous drone-based

behavioral studies have estimated distances using animal body-lengths and approximated

the ground as locally flat (Ringhofer et al., 2020; Torney et al., 2018). While appropriate in

some landscapes, these approaches are not generalizable and preclude fusion of animal

movement data with external georeferenced datasets and analysis of data in standard

units.

While recording focal animals, cameras simultaneously capture imagery of the

surrounding environment, which can potentially be processed to quantify habitat structure,

detect landscape features, or assess resource quality. Previous studies have manually

extracted concurrent behavioral and environmental information from imagery captured with

camera traps (Hofmeester et al., 2020; C. Sun et al., 2021) and animal-borne cameras

(Thompson et al., 2015). But, while aerial photography is widely used for environmental

monitoring (Manfreda et al., 2018), to our knowledge, the simultaneous recording of the

environment and animal behavior remains unexplored with overhead filming methods.

Outside of concurrent imagery, current methods for integrating movement and behavior

with environmental data involve fusing datasets from different sources (e.g. animal location

data from biologgers and satellite imagery), or using secondary sensors (such as animal-

borne cameras) mounted on biologging tags alongside geolocators (Williams et al., 2020).

But these approaches are limited by mismatches between the scale and capture time of

different data types (in the case of data fusion), or limited sampling range of sensors (in the

case of animal-borne cameras, which have limited fields of view and are prone to

obstruction) (Costa-Pereira et al., 2022; Naganuma et al., 2021). Processing video-based

observations to generate concurrent behavioral and environmental datasets at consistent

spatiotemporal scales could facilitate novel studies on the behavioral interactions of

animals with their social, biotic and abiotic environments, thereby generating a more

holistic understanding of the individual-level drivers of higher-order ecological processes

and patterns (Costa-Pereira et al., 2022).

Here we describe a method for using aerial video and computer vision to collect

high-resolution georeferenced locational and behavioral data on free-ranging animals

without capturing or tagging them. In our approach, we use drones to record overhead

video of focal animals and subsequently use a deep learning-based pipeline to

automatically locate and track all individuals in the video. We use Structure-from-Motion

(SfM) techniques to reconstruct the 3D topography of the surrounding habitat (D'Urban

Jackson et al., 2020) which, combined with further image processing, can accurately

transform animal movement data into geographic coordinate systems independent of

camera movement or landscape smoothness. We simultaneously generate additional

behavioral and environmental information, including estimates of each animal's body

posture (pose) and landscape features such as animal trails. This method, illustrated in Fig.

1, is applicable to a wide range of species, generates behavioral data at sub-second and

sub-meter resolution, and produces synchronous information about the surrounding

physical and biotic environment. The use of inexpensive consumer drones that can be

flexibly redeployed promotes larger sample sizes and replication of studies without

incurring additional equipment costs. Although not without its own limitations, such as

possible animal disturbance by drones and limits on observation time (see *Limitations and*

Considerations), this approach has many advantages that make it a powerful complement

to existing methodologies. Our approach thus has the potential to broaden the scope of

behavioral ecology to encompass questions and systems that are unsuitable for direct

observation or bio-logging approaches in isolation.

Figure 1. Overview of the processing pipeline for extracting movement, behavioral, and

landscape data from aerial drone footage of wildlife. Numbered steps correspond to the

numbered sections in the text. First, the animals of interest are video recorded from above (Step 1).

Next, an object detection algorithm is used to localize each animal in every video frame (Step 2), and

these locations are then linked across frames to generate movement trajectories in pixel coordinates

(Step 3). In parallel, anchor frames are selected from the footage and used to build a 3D model of the
landscape and estimate the locations of the drone across the observation. Camera locations at
anchor frames and local visual features are combined to estimate camera locations for all frames
allowing the transformation of the animal trajectories from Step 3 into geographic coordinate space
(Step 4). Optionally, further analyses can be performed to extract more detailed behavioral and
landscape information, for example through localization of specific body parts (keypoints) for each
individual (Step 5) and landscape feature detection (Step 6).
153

154 **Methods**

155
Below we outline the major steps of our methodological approach (Fig. 1). Full details of

each step are given in the Supplement. To illustrate our processing pipeline, we provide

two worked examples with complete code and data available on GitHub

(https://github.com/benkoger/overhead-video-worked-examples). We encourage readers to

explore these examples and to modify the code for their own datasets.

In the first example, we apply our method to multiple African ungulate species. We

recorded ungulate groups at Ol Pejeta and Mpala Conservancies in Laikipia, Kenya over

two field seasons, from November 2 to 16, 2017 and from March 30 to April 19, 2018. In

total, we recorded thirteen species, but in our example we analyze a single observation of

the endangered Grevy's zebra (*Equus grevyi*) (Rubenstein et al., 2016). We flew DJI

Phantom 4 Pro drones [DJI, Shenzhen, China] at an altitude of 80m above the takeoff point.

We deployed two drones sequentially in overlapping relays to achieve a continuous 50

minute observation. Drones were piloted manually, and the drone's position was adjusted

as necessary to keep all herd members in frame. While the zebras seemed undisturbed in

the example observation and in most other observations, in cases where the animals

appeared agitated or fled (as a result of detecting the drone, or our presence via some

other cue(s)), we ended the observation. Our method is agnostic to the exact drone-type

used, and is scalable to the higher-resolution video made possible by employing more

recent models.

In the second worked example, we process video recordings of grassland-dwelling

gelada monkeys (*Theropithecus gelada*). Aerial video recordings of gelada monkeys were

provided by the Guassa Gelada Research Project. The recordings were collected between

October 16, 2019 and February 28, 2020 at the Guassa Community Conservation Area in

the Ethiopian highlands. Drones were piloted manually, taking off initially several hundred

meters away from the herd and ascending to 100m above ground level before slowly

moving above the herd and descending to 70m. This study herd was habituated to close-

proximity research observers and construction machinery from nearby roadwork, and

appeared undisturbed by the drone apart from an initial increase in glance rate which

decreased with exposure. Flights lasted between 5 and 20 minutes. Geladas were recorded

with a DJI Mavic 2 Pro [DJI, Shenzhen, China].

In each worked example, we generate movement trajectories for each animal in the

example videos and 3D models of the surrounding landscape. In the ungulate example, we

also track body keypoints and analyze landscape imagery to detect animal trails, which

zebras may follow while moving across the landscape. For the geladas, we train our

detection model (Step 2), to distinguish between adult males and other individuals.

Step 1. Video Recording

193
Videos should be recorded from above the animals of interest with the camera pointing

directly down (nadir view). The visibility of the animals is of vital importance and will be

affected by the habitat type, video resolution and characteristics of the animals themselves.

As a general rule, videos in which the animals are easily detectable by humans will

significantly ease processing. Although many camera platforms can be used to capture the

footage, drone-mounted cameras will likely be the most common approach and

subsequently we use "drone" and "camera" interchangeably. While our general approach

may work for fixed wing drones that are constantly in motion, this method was designed for

copter style drones that can transition between periods of hovering and movement. For

specific points to consider when planning drone-based data collection, see *Limitations and*

Considerations, below and, for more details, *Step 1* in the Supplement.

Step 2. Detection

207
Animal detection and localization in each video frame is accomplished with deep

convolutional neural networks (CNNs), which we build, train and deploy to predict localizing

bounding boxes (see Table 1 for definitions of bolded terms) for all individuals in all frames

of each video. We first manually annotate frames from the video footage to build image sets

for training the model and evaluating its performance. **Annotation** can be tedious, and the

content of the images will strongly affect model performance; therefore, it is important to

carefully consider the best annotation strategy to achieve high information value for each

annotation while minimizing human labor (see Supplement section 2.1).

To train a deep learning-based object detection model, researchers must first

choose an appropriate software framework and a specific model to train within that

framework (see Supplement section 2.2). We use the Detectron2 API within the PyTorch

framework (Paszke et al., 2019; Wu et al., 2019) in our worked examples, but the user may

choose a different framework depending on their level of coding proficiency or prior

experience with other programming libraries. For simple use cases with clearly-visible

individuals, many common models can be readily re-configured for the researchers' data.

Users with more challenging footage may need to make more considered choices and

possibly incorporate more recent high-performance algorithms. Generally, users should

choose models that have been **pretrained** on general image datasets and that incorporate

image augmentation, as these steps increase training efficiency and require smaller

annotated training sets. See Table S1 for annotation statistics and model performance

metrics for the datasets and trained models used in the worked examples.

After the model is trained, all frames from all videos can be automatically processed.

For each video frame the model generates bounding box coordinates, predicted object

classes and confidence scores for every detected object (Fig. 2). We initially take the mean

of the coordinates of the bounding box corners as an individual's location in the frame,

which we subsequently use for tracking.

Figure 2. Predicted bounding boxes for two example video frames using the trained models

from our worked examples. A) In the ungulates example, the model distinguishes between five

classes (zebra, impala, buffalo, waterbuck, and other). The animals in dark blue bounding boxes are

Grevy's (*Equus grevyi)* and plains zebras (*E. burchelli*) and animals in white bounding boxes are

impala (*Aepyceros melampus*). B) In the gelada monkey example, we distinguish between species

and age-sex class (human-observer, adult-male gelada, and other-gelada). Yellow boxes are

predicted adult-males while light blue boxes are a mix of females and juveniles.

Step 3. Tracking

Tracking, or linking positions across video frames, allows us to generate trajectories for all

detected individuals in the pixel coordinate system of the video. To match individual

locations across consecutive frames, we use a modified version of the Hungarian algorithm

(Kuhn, 1955), which finds the pairing of trajectories and new positions that minimizes the

total distance between all pairs. We have incorporated additional distance- and time-based

rules for connecting detections to tracks that make the algorithm more robust to missing or

false detections by the detection model or from individuals entering or leaving the camera's

field of view during the observation (see Step 3, in the Supplement).

The initial process of linking positions is automated but can result in multiple partial

trajectories for a single individual in the case of environmental occlusions and other

detection issues. For these instances, we provide a **graphical user interface (GUI)** for easy

track validation and error correction. This allows the user to obtain, with limited manual

effort, human-verified continuous trajectories of all individuals in each video within the

pixel-based coordinate system of the video frame, which can then be transformed to a

geographic coordinate system.

Figure 3. Trajectories extracted from drone videos. Trajectories of A) a gelada monkey group over a period of 5 minutes and B) a Grevy's zebra herd over a period of 50 minutes, 24 seconds plotted in video coordinates. Drone and animal movement are entangled and the position of the animals on the earth is unknown. In C) and D), the same trajectories are plotted in geographic coordinates and embedded in a reconstructed landscape. The landscape reconstruction includes visual information as shown in the upper parts of map C) and D) as well as topographic information as shown at the bottom . E) The trajectories have submeter human-validated error between observed locations in video frames and embedded map locations (0.17m median, interquartile range 0.11-0.23m from 100 validated gelada locations; 0.40m median, interquartile range 0.26m-0.58m from 350 validated zebra

locations). In A and C, light blue tracks denote adult male geladas. In B and D, two tracks are

highlighted in yellow and pink for easier comparison. In E, the horizontal line shows the median.

Step 4. Landscape Reconstruction and Geographic Coordinate Transformation

Transforming trajectories into precise geographic coordinates allows us to disentangle the

movement of the tracked animals over variable terrain and the movement of the drone (Fig.

3), and also allows the user to analyze the resulting movement data in standard units and in

relation to external georeferenced data sources, such as satellite imagery. To achieve an

accurate transformation, we must quantify both the topography of the landscape over

which the animals are moving and the location of the camera relative to that landscape over

the course of the observation (see Supplemental Fig. S1 for an illustration). We obtain this

information by first estimating background movement in the video and then using this

information to automatically select a subset of frames (typically numbering a few hundred,

depending on the extent of camera motion) that capture different but overlapping views of

the observation area. We then feed these **"anchor frames"** into **Structure-from-Motion**

(SfM) software (Supplement section 4.1). This builds a detailed 3D model of the landscape

surrounding each observation and also calculates the location of the camera when each

anchor frame was captured. To accurately locate and scale the model in geographic space,

we georeference it using either **ground control points** collected in the field or information

from the drone's onboard GPS sensor (Supplement section 4.2). We track visual features in

each frame to estimate local camera movement between anchor frames, allowing us to

calculate the camera position for every frame in the video (Supplement section 4.3). We

then transform animal positions from the pixel coordinates of the frames into the

geographic coordinates of the 3D model by projecting rays from the estimated camera

location to the surface of the landscape model (Supplement section 4.4). In our examples,

this method yields movement trajectories with sub-meter mean and median error (Fig. 3E);

we provide a GUI that allows the user to efficiently assess the accuracy of their own track

locations (Supplement section 4.5).

Step 5. Body-part Keypoint Detection

302
A major advantage of image-based techniques is that each image contains much more

behaviorally-relevant information than does position alone. If the video resolution is

sufficient, the user can, for example, use existing tools for markerless pose estimation (e.g.

SLEAP (Pereira et al., 2020), DeepLabCut (Mathis et al., 2018; Nath et al., 2019),

DeepPoseKit (Graving et al., 2019)) to extract positions of user-defined body parts on each

detected animal, generating time-varying postural information for tracked individuals (Step

5 in the Supplement; Fig. 5B). Such postural data are particularly amenable to automated

behavioral annotation, and other downstream tasks, because the user can pair the

(relatively) low dimensional keypoint information with human verified annotations from the

raw video.

Figure 5. Additional landscape and behavioral data generated beyond animal location. A) An
SfM-generated 3D triangular mesh landscape model. B) Examples of keypoints detected on zebras.
Nine user-defined keypoints (snout, head, base of the neck, left and right shoulder, left and right
hindquarters, tail base and tip of the tail) are tracked using DeepPoseKit (Graving et al. 2019). C)
Combining landscape data and animal body and head location allows the visualization of animals'
possible visual fields in their landscapes. Here, the red object is the focal zebra, the white rays show
its maximum possible visual field, and the green and white polygons represent bushes and
conspecifics, respectively. D) A SfM-generated 2D orthomosaic image showing predicted animal trail
presence score. Predictions are generated by a CNN trained on separate orthomosaics that were
annotated by three independent human annotators. See worked examples for more detail. The black
box indicates the area shown in E and F, which provide detailed views of the orthomosaic and the
animal trail predictions, respectively. The color bar applies to both D and F.

Step 6. Landscape Quantification

328
The SfM software used in Step 4 generates multiple forms of landscape data that provide

valuable environmental context for the behavior of the observed individuals, including 2D

rasters encoding elevation data, an **orthomosaic image**, and 3D **point clouds** and

triangular meshes (Fig. 5A, E). These outputs can be used directly or further processed to

extract information regarding local landscape features and topography, classify habitat

types, or even estimate the visual fields of tracked animals. In the ungulate worked example

we apply pixel-wise classification algorithms, known as **semantic segmentation**, to

orthomosaic images to automatically detect possible animal trails, which have been shown

to influence the movement of certain savannah dwelling animals *(Cusack et al., 2015;*

Strandburg-Peshkin et al., 2017) and are hypothesized *(Kashetsky et al., 2021)* to influence

animal movement across a range of contexts (Fig. 5D, F). Further possibilities are explored

in the section *Analysis, Extensions, & Applications*.

Limitations and Considerations

Here we discuss the logistics of using drones to capture behavioral data, the suitability of

different research questions to this approach, and the coding skills necessary to implement

this method. For required computing resources, please see the Supplement.

Ethical, logistical and legal considerations

It is important to consider the potential impact of the drone on the focal animals

when planning future research. Behavioral and physiological responses to drone flights can

negatively impact wildlife (Ditmer et al., 2015; Weimerskirch et al., 2018), and may lead to

biased or misleading results in behavioral studies (Duporge et al., 2021). Most species that

respond to drones seem to be primarily affected by the sound of the drone (Duporge et al.,

2021; McEvoy et al., 2016). Pilots may be able to reduce disturbance by choosing quieter

drone models, using low-noise propellers, launching the drone far away from target

animals, approaching from downwind, and flying at higher altitudes. While still an emerging

field of study, Duporge et al. (2021) and others (Bevan et al., 2018; Christiansen et al., 2016;

Mulero-Pázmány et al., 2017) offer guidelines for flying drones near a variety of species. It is

typically prudent to perform preliminary flights prior to data collection in order to assess the

animals' response to the drone and establish appropriate protocols.

When choosing a study environment, factors such as extreme temperatures, dust,

wind, precipitation and fog can reduce visibility, equipment longevity and flight

performance. Furthermore, landscapes (such as open water and snow fields) without

abundant and persistent visible landmarks are not suitable for SfM methods, and may

preclude transformation of animal trajectories into geographic coordinates (Step 4). SfM

height mapping accuracy is also sensitive to vegetation structure and motion (such as tall

grass in the wind), which may introduce errors in those parts of the landscape relative to

static regions (D'Urban Jackson et al., 2020). Additionally, the real-world geospatial

accuracy of the final data is ultimately determined by the quality of the 3D landscape

models. For sub-meter accuracy one must either be able to place ground control points in

the landscape of interest or have access to existing georeferenced landscape models or

imagery.

Finally, researchers must be aware of and abide by legal regulations regarding

drone operations in any prospective study area both in terms of formal permissions and

licenses required to deploy drones as well as limitations on in-flight maneuvers. Particular

rules vary by location and are continually evolving but can include limits on flight altitudes,

distances, or locations (i.e. airports, national parks, or certain government areas) and

requirements for maintaining visual contact with drones and keeping safe distances from

people and structures. Beyond legal requirements, researchers should also ensure that

projects do not negatively impact local communities. For a thorough discussion of ethical

drone use, see Duffy et al. (2018).

Research question suitability

In determining whether image-based data collection is appropriate for a research

question, researchers should consider the required data resolution and spatial and

temporal scales of all targeted study behaviors. Animal groups that are spread over large

areas may be impossible to fully capture at an appropriate resolution, especially if

automated tracking or individual posture data are required. Deploying multiple drones

simultaneously can increase spatial coverage, but complicates flight protocols and data

processing. Additionally, spatially-expansive behaviors like long-distance hunts may draw

the drone beyond the legal operation distance from stationary pilots (Creel & Creel, 1995).

Nocturnal or crepuscular animals would require imaging via thermal or high-sensitivity

cameras, which imposes further constraints to the spatial, and temporal, resolution of video

that can be obtained. Nocturnal operations also often require additional operation

permissions and specialized pilot training.

The timescale over which behaviors occur is also important. Drone battery efficiency

has rapidly improved over the years, but flight times are still relatively short (currently 30-45

minutes for widely-used models). While observation time can be extended with sequentially

overlapping flights by pairs of drones, theoretically allowing limitless observation times, the

practical duration of observations are still limited by battery supply, pilot fatigue, and the

movement of the focal animals away from the operational site. Recording rare or

unpredictable behaviors, such as predation events, requires deploying drones in the right

place at the right time. Incorporating external observational or biologging data could help

predict such behaviors and inform the location and timing of deployments. Relatedly,

deploying a small number of geolocators on individuals in the target population could allow

researchers to more reliably locate rare species or repeatedly target focal individuals and

associated conspecifics for drone-based observation.

Programming proficiency

All code provided in the worked examples is written in the Python programming language.

We expect limited additional programming will be required to apply this code to new

datasets that are similar in scope to the examples. However, we expect many researchers

will want to adapt our base code to better suit their needs, which will require some degree

of programming skill, depending on the functionality required. Furthermore, since this

method generates large volumes of high-resolution and high-dimensional data, certain

programming capabilities will prove essential for effective visualization and analysis (see

below for possible analysis directions).

Analysis, Extensions, & Applications

With recent advances in imaging technology and video analysis, drone-based behavioral

observation is poised to become a widely-used approach in the study of wildlife ecology.

Here we expand on the potential of this approach by identifying avenues of research that

are particularly well-suited to this approach, outlining possible data analysis approaches,

and discussing future extensions of drone-based observation.

Potential research questions

Drone-based observation and our analytical pipeline open the door to several types of

questions that can be particularly difficult to explore using existing methods, such as direct

observation, camera trapping, or biologging:

1) Our approach allows for precise quantification of the positions of all observed

435   individuals and environmental features, thereby lending itself to exploring *spatially-*

436   *explicit social behavioral processes*. For example, geladas, plains zebras, and many

437   other animals form multi-level societies, which are reflected in individual association

438   patterns and spatial configurations of groups (Couzin, 2006; Couzin & Laidre, 2009;

439   Grueter et al., 2020; Papageorgiou et al., 2019). These in turn likely affect

440   information flow and collective decision-making, which can potentially be explored

441   by tracking behavioral change and movement patterns.

442   2) The ability to collect high-resolution behavioral data on all observation subjects

443   simultaneously is conducive to exploring questions relating to *dynamic collective*

444   *processes*. For example, prey species exhibit vigilance behavior that mediates

445   intragroup information transfer and collective detection of predators (Lima & Zollner,

446   1996; Lin et al., 2020). Individuals may adjust their vigilance behavior in response to

447   the behavior of group mates, as well as various other social and environmental

448   factors (Beauchamp, 2015; Chen et al., 2021; Pays et al., 2007), but it is challenging

449   to track the behavioral states of multiple individuals simultaneously using non-

450   image-based observational methods.

3) The ability to observe animals and their surroundings concurrently allows for more

detailed understanding of *animal-environment interactions*. For example, biologging

studies have found that roadways strongly affect wildlife movement, leading to the

construction of over- or underpasses to facilitate safe crossing of roadways and

promote landscape connectivity (Shepard et al., 2008; Simpson et al., 2016). Aerial

cameras could potentially be used to monitor these structures to determine how the

presence, speed, and visual characteristics of vehicles affects animals' use of these

structures.

To the extent that these examples can be addressed using other methodologies, they will

require multiple sensors, fusion of diverse data streams, and/or considerable effort by

human observers. In contrast, drone-based observation allows for all the relevant data to

be collected simultaneously using a single sensor.

Data analysis

Our method provides a rich set of biologically-relevant features, but these data must

be analyzed to address the research questions of interest. There exists a rich body of

literature for exploring, visualizing, and analyzing animal trajectory data (Patterson et al.,

2017; Seidel et al., 2018). The sub-second and sub-meter animal positions and detailed

landscape data produced by this method allow researchers to use step-selection-type

approaches to evaluate movement decisions across spatial and temporal scales, from

actual individual steps to movement decisions at larger-scales (Fieberg et al., 2021).

Tracking body keypoint locations (Step 5) can directly provide information about

behaviorally-relevant body parts, such as head direction and body orientation. Depending

on the position of the animal(s) within each frame, correctly interpreting animals' relative

keypoint patterns may require taking lens geometry into account (see Supplemental Fig. S3

for details). When possible to estimate, this head and body information can be combined

with the 3D landscape models (Step 4) to reconstruct estimates of each animal's visual field

(Aben et al., 2018).

Future extensions

During detection (Step 2), if given sufficient image resolution, it may be possible to

visually identify individual animals within and across observations. Individual recognition

opens the door to studies of individual behavioral variation across time and contexts, and

the role of individual behavior in driving processes at the population and community levels

(Costa-Pereira et al., 2022). Beyond laboratory settings (Walter & Couzin, 2021) individual

recognition in wild populations is increasingly feasible (Norman et al., 2017; Tuia et al.,

2022). Alternatively, individuals could be identified using ground-level photographs or direct

observation, and then these identities could be manually linked to individuals in the aerial

recordings.

A further advance would be to use temporal keypoint data, or body posture

trajectories (Fig. 1 - Step 5), to define finescale behavioral labels. While similar to the

problem of accelerometer-based behavioral classification, a vision-based approach

provides the added benefit of ground-truth videos for validation (Brown et al., 2013; Wang

et al., 2015). Both supervised (Bohnslav et al., 2021) and unsupervised/self-supervised

(Berman et al., 2014) approaches, or a combination of both (J. J. Sun et al., 2021), could be

applied to achieve automated behavioral annotations describing the behavioral states of

individuals and groups.

Beyond data processing, advancements with drones' on board automated visual

tracking (Islam et al., 2019) and the ability to automatically coordinate flight among multiple

drones (Zhou et al., 2022) could help to streamline complex operations, reduce the risk of

human error, and also facilitate further observation techniques such as multiview 3D

posture tracking (Tallamraju et al., 2019). Drones could thus be deployed to autonomously

find and record individuals of a target species. While these are exciting future steps, current

regulations in many countries would prevent the use of these methods without special

permissions or certifications. Thus, regulatory rather than technological restrictions may be

the most substantial barriers to large-scale automated observation.

Synergizing with other remote-sensing methods

There is great potential in combining this approach with existing remote sensing

methods. In the context of biologging, drones could be deployed at key times or locations

of interest to provide high-resolution behavioral snapshots of tagged individuals along with

their social and environmental context. Image-based data could also be used to guide

biologging study design choices. For example, one could downsample the high resolution

image-based data to determine the optimal sampling frequency for bio-logging studies,

using the videos to verify that the resulting data capture the target behaviors. Recording the

behavior of instrumented animals could also provide sensor ground truth data and aid in

the development of more energy efficient, behaviorally activated sensors (Korpela et al.,

2020; Yu et al., 2021).

The geolocated animal movement and environmental data generated by drone-

based methods can be combined with additional multi-modal remote sensing data to

explore the interplay between animals and important environmental features. For example,

higher accuracy canopy height and landscape models, such as those generated by

emerging workflows of SfM, LiDAR, or their fusion (D'Urban Jackson et al., 2020), combined

with animal behavioral data generated from our method, could enable studies of herbivore

impacts on vegetation via consumption or trampling effects, or allow higher precision visual

field calculations in difficult-to-model environments like tall grass. Additionally, SfM models,

such as those generated in our method, can be combined with various other remote

sensing data to enable calculation of microclimate variability (Duffy et al., 2021; Maclean,

2019) water flow and saturation (Koci et al., 2020), or resource quality and accessibility

(Jennewein et al., 2021) which would also be informative for analyzing recorded animal

behavior.

Conclusion

We present a method that allows researchers to study animal behavior in its natural social

and environmental context in a non-invasive and scalable way. Our approach is

independent of specific species and can be deployed across a range of study systems

making this a powerful and versatile tool for many researchers across fields. Importantly, as

researchers work to understand the relationship between animals and their landscapes in a

changing world, this method, which simultaneously records both at high resolution, is

poised to be an important new way of observing the natural world.

Term	Definition
anchor frames	The subset of video frames in our method input into the structure from motion software to generate the 3D landscape model and camera location information.
annotation	The process of labeling objects of interest in training imagery, for example by drawing bounding boxes around individual animals.
bounding box	A rectangle enclosing an object of interest in an image. Bounding boxes can be drawn by the user as a form of annotation, or can be generated by detection models to denote the predicted location of an object of interest.
graphical user interface (GUI)	A means of viewing or inputting data that relies on graphical elements (e.g. buttons) rather than coding inputs.
ground control point (GCP)	Locations or landmarks with known real-world geographic coordinates. These points are used to georeference the landscape model generated from the anchor frames.
image augmentation	A technique for increasing the effective size of the training set by modifying each training image each time it is shown to the model. Modifications, including blurring, rotating, and adjusting contrast, and are intended to mimic the diversity of images in the entire dataset.
orthomosaic	Two-dimensional composite image generated by the SfM software

	that approximate the appearance of 3D structures from an overhead viewpoint.
point cloud	A set of 3D points often also containing a color value that can be used to represent landscapes or objects in space.
pretraining	A feature of many common deep learning models where the model has been initially trained on large datasets consisting of generic imagery of common scenes and objects. This allows the model to learn the basic, universal aspects of imagery before being fine-tuned on the user's specific dataset.
semantic segmentation	The process of labeling every pixel in an image as one of a set number of object classes.
structure from motion (SfM)	A technique for 3D reconstruction in which two-dimensional images from various overlapping viewpoints are used to define the geometry of the target structure (here, the landscape surrounding the observed animals).
triangular mesh	A 3D surface model created from a point cloud by connecting triads of points to create flat triangular surfaces.

Table 1. Glossary of terms. Defined terms are bolded at first appearance in the text.

Authors contributions

Benjamin Koger, Blair Costelloe and Iain Couzin developed the idea and goals of the

method. Benjamin Koger, Blair Costelloe and Jeffrey Kerby coordinated and conducted

data collection. Benjamin Koger, Blair Costelloe, Adwait Desphande, Jacob Graving and

Jeffrey Kerby curated the data. Benjamin Koger designed the analytical pipeline; wrote the

code in the worked examples and performed all aspects of model development, including

training and validation. Jacob Graving advised on the design, organization, and

development of earlier versions of the code. Blair Costelloe and Iain Couzin provided

supervision and secured financial support. Blair Costelloe, Iain Couzin and Jeffrey Kerby

administered and contributed material resources to the project. Benjamin Koger and Blair

Costelloe wrote the initial manuscript draft and prepared the figures. All authors edited and

revised the manuscript and gave final approval for publication.

Acknowledgements

Figures 1 and 4 were created by Mike Costelloe. We thank Mike Costelloe, Sven Lauke and

Felicitas Oehler for assistance with collecting and annotating the ungulate data. We thank

Guassa Gelada Research Project field managers Odin Bernardo and Nate Redon for

collecting the gelada drone data, and project directors, camp staff members, and the local

community for their support. We thank Anne Scharf for sharing code used to plan DJI

mapping flights. We thank Jean Luo and David Basili for contributions to prior versions of

the animal trail detection and track correction GUI. B.R.C. received support from the

European Union's Horizon 2020 research and innovation program under the Marie

Sklodowska-Curie grant agreement No. 748549. B.R.C. acknowledges support from the

University of Konstanz's Investment Grant program. J.T.K acknowledges support from the

Neukom Institute for Computational Science at Dartmouth College and the European

Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie

grant agreement No. 754513 and the Aarhus University Research Foundation. B.K., I.D.C,

J.M.G., B.R.C., and A.D. acknowledge support from the Deutsche Forschungsgemeinschaft

(DFG, German Research Foundation) under Germany's Excellence Strategy – 'Centre for

the Advanced Study of Collective Behaviour' EXC 2117-422037984. I.D.C gratefully

acknowledges support from the Office of Naval Research (ONR) grant N00014-19-1-2556

and the European Union's Horizon 2020 research and innovation program under the Marie

Sklodowska-Curie grant agreement No. 860949. A.D. acknowledges the support by the

Swiss National Science Foundation (Early PostDoc Fellowship: P2NEP3_200190). J.M.G.

and B.R.C. acknowledge support from NVIDIA Corporation's Academic Hardware Grant

Program.

Permissions and Ethics Statement

We imported and operated drones in Kenya with permission from the Kenya Civil Aviation

Authority (import permits: KCAA/RPA/PERMIT-2017-0006, KCAA/RPA/PERMIT-2017-0007,

KCAA/RPA/PERMIT-2017-0008, KCAA/ASSR/RPA/PERMIT-0016,

KCAA/ASSR/RPA/PERMIT-0017, KCAA/ASSR/RPA/PERMIT-0018; authorization numbers:

KCAA/OPS/2117/4 Vol. 2 (80), KCAA/OPS/2117/4 Vol. 2 (81), KCAA/OPS/2117/5 (86),

KCAA/OPS/2117/5 (87); operator certificate numbers: RPA/TP/0005, RPA/TP/0000-0009).

We conducted research in Kenya with permission from the Kenyan National Commission

for Science, Technology and Innovation (research permits: NACOSTI/P/17/59088/15489,

NACOSTI/P/59088/21567) and in affiliation with the Kenya Wildlife Service. All research

activities pertaining to the ungulate data, including drone operation, were performed with

the knowledge and support of management and security staff at our field sites, Ol Pejeta

Conservancy and Mpala Research Centre. Data collection protocols for the ungulate data

were reviewed and approved by Ethikrat, the independent ethics council of the Max Planck

Society. All research activities pertaining to gelada monkeys, including drone operation,

were undertaken with the knowledge and approval of the Guassa Community Conservation

Area leadership and under the approval of a memorandum of understanding between the

Ethiopian Wildlife Conservation Authority and the Guassa Gelada Research Project.

Data availability statement

Code for the worked examples is available on GitHub at

https://github.com/benkoger/overhead-video-worked-examples. All data for the worked

examples are archived on EDMOND at

https://edmond.mpdl.mpg.de/privateurl.xhtml?token=9c1a978a-21f3-4843-bebe-

40c296bffc73. DOIs and stable links to these resources will be generated upon final

acceptance of the manuscript.

References

Aben, J., Pellikka, P., & Travis, J. M. J. (2018). A call for viewshed ecology: Advancing our understanding of the ecology of information through viewshed analysis. *Methods in Ecology and Evolution*, *9*(3), 624–633. https://doi.org/10.1111/2041-210X.12902

Atkins, J. L., Long, R. A., Pansu, J., Daskin, J. H., Potter, A. B., Stalmans, M. E., Tarnita, C. E., & Pringle, R. M. (2019). Cascading impacts of large-carnivore extirpation in an African ecosystem. *Science*, *364*(6436), 173–177. https://doi.org/10.1126/science.aau3561

Beauchamp, G. (2015). Visual obstruction and vigilance: A natural experiment. *Journal of Avian Biology*, *46*(5), 476–481. https://doi.org/10.1111/jav.00612

Berdahl, A., Torney, C. J., Ioannou, C. C., Faria, J. J., & Couzin, I. D. (2013). Emergent Sensing of Complex Environments by Mobile Animal Groups. *Science*, *339*(6119), 574–576. https://doi.org/10.1126/science.1225883

Berger- Tal, O., Blumstein, D. T., & Swaisgood, R. R. (2020). Conservation translocations: A review of common difficulties and promising directions. *Animal Conservation*, *23*(2), 121–131. https://doi.org/10.1111/acv.12534

Berman, G. J., Choi, D. M., Bialek, W., & Shaevitz, J. W. (2014). Mapping the stereotyped behaviour of freely moving fruit flies. *Journal of The Royal Society Interface*, *11*(99), 20140672. https://doi.org/10.1098/rsif.2014.0672

Bevan, E., Whiting, S., Tucker, T., Guinea, M., Raith, A., & Douglas, R. (2018). Measuring behavioral responses of sea turtles, saltwater crocodiles, and crested terns to drone disturbance to define ethical operating thresholds. *PLOS ONE*, *13*(3), e0194460. https://doi.org/10.1371/journal.pone.0194460

Bohnslav, J. P., Wimalasena, N. K., Clausing, K. J., Dai, Y. Y., Yarmolinsky, D. A., Cruz, T., Kashlan, A. D., Chiappe, M. E., Orefice, L. L., Woolf, C. J., & Harvey, C. D. (2021). DeepEthogram, a machine learning pipeline for supervised behavior classification from raw pixels. *ELife*, *10*, e63377. https://doi.org/10.7554/eLife.63377

Bro-Jørgensen, J., Franks, D. W., & Meise, K. (2019). Linking behaviour to dynamics of

populations and communities: Application of novel approaches in behavioural ecology

to conservation. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *374*(1781), 20190008. https://doi.org/10.1098/rstb.2019.0008

Brown, D. D., Kays, R., Wikelski, M., Wilson, R., & Klimley, A. P. (2013). Observing the

unwatchable through acceleration logging of animal behavior. *Animal Biotelemetry*,

1(1), 20. https://doi.org/10.1186/2050-3385-1-20

Buchholz, R., Banusiewicz, J. D., Burgess, S., Crocker-Buta, S., Eveland, L., & Fuller, L. (2019).

Behavioural research priorities for the study of animal response to climate change.

Animal Behaviour, *150*, 127–137. https://doi.org/10.1016/j.anbehav.2019.02.005

Ceballos, G., Ehrlich, P. R., & Raven, P. H. (2020). Vertebrates on the brink as indicators of

biological annihilation and the sixth mass extinction. *Proceedings of the National*

Academy of Sciences, *117*(24), 13596–13602. https://doi.org/10.1073/pnas.1922686117

Chen, C.-W., Whiting, M. J., Yang, E.-C., & Lin, S.-M. (2021). Do I stay or do I go? Shifts in perch

use by lizards during morning twilight suggest anticipatory behaviour. *Biology Letters*,

17(10), 20210388. https://doi.org/10.1098/rsbl.2021.0388

Christiansen, F., Rojano-Doñate, L., Madsen, P. T., & Bejder, L. (2016). Noise Levels of Multi-

Rotor Unmanned Aerial Vehicles with Implications for Potential Underwater Impacts

on Marine Mammals. *Frontiers in Marine Science*, *3*.

https://www.frontiersin.org/articles/10.3389/fmars.2016.00277

Corcoran, E., Winsen, M., Sudholz, A., & Hamilton, G. (2021). Automated detection of wildlife

using drones: Synthesis, opportunities and constraints. *Methods in Ecology and*

Evolution, *12*(6), 1103–1114. https://doi.org/10.1111/2041-210X.13581

Costa- Pereira, R., Moll, R. J., Jesmer, B. R., & Jetz, W. (2022). Animal tracking moves

community ecology: Opportunities and challenges. *Journal of Animal Ecology*, 1365-

2656.13698. https://doi.org/10.1111/1365-2656.13698

Couzin, I. D. (2006). Behavioral Ecology: Social Organization in Fission–Fusion Societies.

Current Biology, *16*(5), R169–R171. https://doi.org/10.1016/j.cub.2006.02.042

Couzin, I. D., & Laidre, M. E. (2009). Fission–fusion populations. *Current Biology*, *19*(15), R633–

R635. https://doi.org/10.1016/j.cub.2009.05.034

Creel, S., & Creel, N. M. (1995). Communal hunting and pack size in African wild dogs, Lycaon

pictus. *Animal Behaviour*, *50*(5), 1325–1339. https://doi.org/10.1016/0003-

3472(95)80048-4

Cusack, J. J., Dickman, A. J., Rowcliffe, J. M., Carbone, C., Macdonald, D. W., & Coulson, T.

(2015). Random versus Game Trail-Based Camera Trap Placement Strategy for

Monitoring Terrestrial Mammal Communities. *PLOS ONE*, *10*(5), e0126373.

https://doi.org/10.1371/journal.pone.0126373

Dell, A. I., Bender, J. A., Branson, K., Couzin, I. D., de Polavieja, G. G., Noldus, L. P. J. J., Pérez-

Escudero, A., Perona, P., Straw, A. D., Wikelski, M., & Brose, U. (2014). Automated

image-based tracking and its application in ecology. *Trends in Ecology & Evolution*,

29(7), 417–428. https://doi.org/10.1016/j.tree.2014.05.004

Ditmer, M. A., Vincent, J. B., Werden, L. K., Tanner, J. C., Laske, T. G., Iaizzo, P. A., Garshelis, D.

L., & Fieberg, J. R. (2015). Bears show a physiological but limited behavioral response

to unmanned aerial vehicles. *Current Biology*, *25*(17), 2278–2283.

https://doi.org/10.1016/j.cub.2015.07.024

Duffy, J. P., Anderson, K., Fawcett, D., Curtis, R. J., & Maclean, I. M. D. (2021). Drones provide

spatial and volumetric data to deliver new insights into microclimate modelling.

Landscape Ecology, *36*(3), 685–702. https://doi.org/10.1007/s10980-020-01180-9

Duffy, J. P., Cunliffe, A. M., DeBell, L., Sandbrook, C., Wich, S. A., Shutler, J. D., Myers-Smith, I.

H., Varela, M. R., & Anderson, K. (2018). Location, location, location: Considerations

when using lightweight drones in challenging environments. *Remote Sensing in*

Ecology and Conservation, *4*(1), 7–19. https://doi.org/10.1002/rse2.58

Duporge, I., Spiegel, M. P., Thomson, E. R., Chapman, T., Lamberth, C., Pond, C., Macdonald,

D. W., Wang, T., & Klinck, H. (2021). Determination of optimal flight altitude to minimise

acoustic drone disturbance to wildlife using species audiograms. *Methods in Ecology*

and Evolution, *12*(11), 2196–2207. https://doi.org/10.1111/2041-210X.13691

D'Urban Jackson, T., Williams, G. J., Walker-Springett, G., & Davies, A. J. (2020). Three-

dimensional digital mapping of ecosystems: A new era in spatial ecology. *Proceedings*

of the Royal Society B: Biological Sciences, *287*(1920), 20192383.

https://doi.org/10.1098/rspb.2019.2383

Fieberg, J., Signer, J., Smith, B., & Avgar, T. (2021). A 'How to' guide for interpreting

parameters in habitat-selection analyses. *Journal of Animal Ecology*, *90*(5), 1027–

1043. https://doi.org/10.1111/1365-2656.13441

Flack, A., Nagy, M., Fiedler, W., Couzin, I. D., & Wikelski, M. (2018). From local collective

behavior to global migratory patterns in white storks. *Science*, *360*(6391), 911–914.

https://doi.org/10.1126/science.aap7781

Francisco, F. A., Nührenberg, P., & Jordan, A. (2020). High-resolution, non-invasive animal

tracking and reconstruction of local environment in aquatic ecosystems. *Movement*

Ecology, *8*(1), 27. https://doi.org/10.1186/s40462-020-00214-w

Graving, J. M., Chae, D., Naik, H., Li, L., Koger, B., Costelloe, B. R., & Couzin, I. D. (2019).

DeepPoseKit, a software toolkit for fast and robust animal pose estimation using deep

learning. *ELife*, *8*, e47994. https://doi.org/10.7554/eLife.47994

Grueter, C. C., Qi, X., Zinner, D., Bergman, T., Li, M., Xiang, Z., Zhu, P., Migliano, A. B., Miller, A.,

Krützen, M., Fischer, J., Rubenstein, D. I., Vidya, T. N. C., Li, B., Cantor, M., & Swedell, L.

(2020). Multilevel Organisation of Animal Sociality. *Trends in Ecology & Evolution*,

35(9), 834–847. https://doi.org/10.1016/j.tree.2020.05.003

Haalck, L., Mangan, M., Webb, B., & Risse, B. (2020). Towards image-based animal tracking in

natural environments using a freely moving camera. *Journal of Neuroscience*

Methods, *330*, 108455. https://doi.org/10.1016/j.jneumeth.2019.108455

Hale, R., Blumstein, D. T., Mac Nally, R., & Swearer, S. E. (2020). Harnessing knowledge of

animal behavior to improve habitat restoration outcomes. *Ecosphere*, *11*(4).

https://doi.org/10.1002/ecs2.3104

Hofmeester, T. R., Young, S., Juthberg, S., Singh, N. J., Widemo, F., Andrén, H., Linnell, J. D. C.,

& Cromsigt, J. P. G. M. (2020). Using by-catch data from wildlife surveys to quantify

climatic parameters and timing of phenology for plants and animals using camera

traps. *Remote Sensing in Ecology and Conservation*, *6*(2), 129–140.

https://doi.org/10.1002/rse2.136

Hughey, L. F., Hein, A. M., Strandburg-Peshkin, A., & Jensen, F. H. (2018). Challenges and

solutions for studying collective animal behaviour in the wild. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *373*(1746), 20170005.

https://doi.org/10.1098/rstb.2017.0005

Inoue, S., Yamamoto, S., Ringhofer, M., Mendonça, R. S., Pereira, C., & Hirata, S. (2019). Spatial

positioning of individuals in a group of feral horses: A case study using drone

technology. *Mammal Research*, *64*(2), 249–259. https://doi.org/10.1007/s13364-018-

0400-2

Islam, M. J., Hong, J., & Sattar, J. (2019). Person-following by autonomous robots: A

categorical overview. *The International Journal of Robotics Research*, *38*(14), 1581–

1618. https://doi.org/10.1177/0278364919881683

Jennewein, J. S., Eitel, J. U. H., Joly, K., Long, R. A., Maguire, A. J., Vierling, L. A., & Weygint, W.

(2021). Estimating integrated measures of forage quality for herbivores by fusing

optical and structural remote sensing data. *Environmental Research Letters*, *16*(7),

075006. https://doi.org/10.1088/1748-9326/ac09af

Kashetsky, T., Avgar, T., & Dukas, R. (2021). The Cognitive Ecology of Animal Movement:

Evidence From Birds and Mammals. *Frontiers in Ecology and Evolution*, *9*.

746   https://www.frontiersin.org/articles/10.3389/fevo.2021.724887

747 Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye on

748   life and planet. *Science*, *348*(6240). https://doi.org/10.1126/science.aaa2478

749 Koci, J., Sidle, R. C., Jarihani, B., & Cashman, M. J. (2020). Linking hydrological connectivity to

750   gully erosion in savanna rangelands tributary to the Great Barrier Reef using

751   structure- from- motion photogrammetry. *Land Degradation & Development*, *31*(1), 20–

752   36. https://doi.org/10.1002/ldr.3421

753 Korpela, J., Suzuki, H., Matsumoto, S., Mizutani, Y., Samejima, M., Maekawa, T., Nakai, J., &

754   Yoda, K. (2020). Machine learning enables improved runtime and precision for bio-

755   loggers on seabirds. *Communications Biology*, *3*(1), Article 1.

756   https://doi.org/10.1038/s42003-020-01356-8

757 Kuhn, H. W. (1955). The Hungarian Method for the assignment problem. *Naval Research*

758   *Logistics Quarterly*, *2*, 83–97.

759 Lima, S. L., & Zollner, P. A. (1996). Anti-predatory vigilance and the limits to collective

760   detection: Visual and spatial separation between foragers. *Behavioral Ecology and*

761   *Sociobiology*, *38*(5), 355–363. https://doi.org/10.1007/s002650050252

762 Lin, B., Foxfoot, I. R., Miller, C. M., Venkatamaran, V. V., Kerby, J. T., Bechtold, E. K., Kellogg, B.

763   S., Nguyen, N., & Fashing, P. J. (2020). Leopard predation on gelada monkeys at

764   Guassa, Ethiopia. *American Journal of Primatology*, *82*(2), e23098.

765   https://doi.org/10.1002/ajp.23098

766 Lutz, M. J., Reid, C. R., Lustri, C. J., Kao, A. B., Garnier, S., & Couzin, I. D. (2021). Individual error

767   correction drives responsive self-assembly of army ant scaffolds. *Proceedings of the*

768   *National Academy of Sciences*, *118*(17), e2013741118.

769   https://doi.org/10.1073/pnas.2013741118

770 Maclean, I. M. D. (2019). Predicting future climate at high spatial and temporal resolution.

771   *Global Change Biology*, *26*(2), 1003–1011. https://doi.org/10.1111/gcb.14876

Manfreda, S., McCabe, M. F., Miller, P. E., Lucas, R., Pajuelo Madrigal, V., Mallinis, G., Ben Dor,

E., Helman, D., Estes, L., Ciraolo, G., Müllerová, J., Tauro, F., De Lima, M. I., De Lima, J.

L. M. P., Maltese, A., Frances, F., Caylor, K., Kohv, M., Perks, M., … Toth, B. (2018). On

the Use of Unmanned Aerial Systems for Environmental Monitoring. *Remote Sensing*,

10(4), Article 4. https://doi.org/10.3390/rs10040641

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M.

(2018). DeepLabCut: Markerless pose estimation of user-defined body parts with deep

learning. *Nature Neuroscience*, *21*(9), 1281–1289. https://doi.org/10.1038/s41593-018-

0209-y

McEvoy, J. F., Hall, G. P., & McDonald, P. G. (2016). Evaluation of unmanned aerial vehicle

shape, flight path and camera type for waterfowl surveys: Disturbance effects and

species recognition. *PeerJ*, *4*. https://doi.org/10.7717/peerj.1831

Mortelliti, A. (2022). The importance of animal behavior for ecosystem services. *Trends in*

Ecology & Evolution, S0169534722002762. https://doi.org/10.1016/j.tree.2022.10.009

Mulero-Pázmány, M., Jenni-Eiermann, S., Strebel, N., Sattler, T., Negro, J. J., & Tablado, Z.

(2017). Unmanned aircraft systems as a new source of disturbance for wildlife: A

systematic review. *PLOS ONE*, *12*(6), e0178448.

https://doi.org/10.1371/journal.pone.0178448

Naganuma, T., Tanaka, M., Tezuka, S., M.J.G. Steyaert, S., Tochigi, K., Inagaki, A., Myojo, H.,

Yamazaki, K., & Koike, S. (2021). Animal- borne video systems provide insight into the

reproductive behavior of the Asian black bear. *Ecology and Evolution*, *11*(14), 9182–

9190. https://doi.org/10.1002/ece3.7722

Nath, T., Mathis, A., Chen, A. C., Patel, A., Bethge, M., & Mathis, M. W. (2019). Using

DeepLabCut for 3D markerless pose estimation across species and behaviors. *Nature*

Protocols, *14*(7), 2152–2176. https://doi.org/10.1038/s41596-019-0176-0

Norman, B. M., Holmberg, J. A., Arzoumanian, Z., Reynolds, S. D., Wilson, R. P., Rob, D., Pierce,

S. J., Gleiss, A. C., de la Parra, R., Galvan, B., Ramirez-Macias, D., Robinson, D., Fox, S.,

Graham, R., Rowat, D., Potenski, M., Levine, M., Mckinney, J. A., Hoffmayer, E., …

Morgan, D. L. (2017). Undersea Constellations: The Global Biology of an Endangered

Marine Megavertebrate Further Informed through Citizen Science. *BioScience*, *67*(12),

1029–1043. https://doi.org/10.1093/biosci/bix127

Papageorgiou, D., Christensen, C., Gall, G. E. C., Klarevas-Irby, J. A., Nyaguthii, B., Couzin, I. D.,

& Farine, D. R. (2019). The multilevel society of a small-brained bird. *Current Biology*,

29(21), R1120–R1121. https://doi.org/10.1016/j.cub.2019.09.072

Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z.,

Gimelshein, N., Antiga, L., Desmaison, A., Kopf, A., Yang, E., DeVito, Z., Raison, M.,

Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., … Chintala, S. (2019). PyTorch: An

Imperative Style, High-Performance Deep Learning Library. In H. Wallach, H.

Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, & R. Garnett (Eds.), *Advances in*

Neural Information Processing Systems 32 (pp. 8024–8035). Curran Associates, Inc.

http://papers.neurips.cc/paper/9015-pytorch-an-imperative-style-high-performance-

deep-learning-library.pdf

Patterson, T. A., Parton, A., Langrock, R., Blackwell, P. G., Thomas, L., & King, R. (2017).

Statistical modelling of individual animal movement: An overview of key methods and

a discussion of practical challenges. *AStA Advances in Statistical Analysis*, *101*(4),

399–438. https://doi.org/10.1007/s10182-017-0302-7

Pays, O., Renaud, P.-C., Loisel, P., Petit, M., Gerard, J.-F., & Jarman, P. J. (2007). Prey

synchronize their vigilant behaviour with other group members. *Proceedings of the*

Royal Society B: Biological Sciences, *274*(1615), 1287–1291.

https://doi.org/10.1098/rspb.2006.0204

Pekarsky, S., Schiffner, I., Markin, Y., & Nathan, R. (2021). Using movement ecology to evaluate

the effectiveness of multiple human-wildlife conflict management practices. *Biological*

Conservation, *262*, 109306. https://doi.org/10.1016/j.biocon.2021.109306

Pereira, T. D., Tabris, N., Li, J., Ravindranath, S., Papadoyannis, E. S., Wang, Z. Y., Turner, D. M.,

McKenzie-Smith, G., Kocher, S. D., Falkner, A. L., Shaevitz, J. W., & Murthy, M. (2020).

SLEAP: Multi-animal pose tracking (p. 2020.08.31.276246).

https://doi.org/10.1101/2020.08.31.276246

Pollux, B. J. A. (2017). Consistent individual differences in seed disperser quality in a seed-

eating fish. *Oecologia*, *183*(1), 81–91. https://doi.org/10.1007/s00442-016-3749-4

Raoult, V., Tosetto, L., & Williamson, J. E. (2018). Drone-based high-resolution tracking of

aquatic vertebrates. *Drones*, *2*(4), 37. https://doi.org/10.3390/drones2040037

Ringhofer, M., Go, C. K., Inoue, S., S. Mendonça, R., Hirata, S., Kubo, T., Ikeda, K., & Yamamoto,

S. (2020). Herding mechanisms to maintain the cohesion of a harem group: Two

interaction phases during herding. *Journal of Ethology*, *38*(1), 71–77.

https://doi.org/10.1007/s10164-019-00622-5

Rubenstein, D., Low Mackey, B., Davidson, Z. D., Kebede, F., & King, S. R. B. (2016). Equus

grevyi. *The IUCN Red List of Threatened Species*, eT7950A89624491.

Seidel, D. P., Dougherty, E., Carlson, C., & Getz, W. M. (2018). Ecological metrics and methods

for GPS movement data. *International Journal of Geographical Information Science*,

32(11), 2272–2293. https://doi.org/10.1080/13658816.2018.1498097

Shaw, A. K. (2020). Causes and consequences of individual variation in animal movement.

Movement Ecology, *8*(1), 12. https://doi.org/10.1186/s40462-020-0197-x

Shepard, D. B., Kuhns, A. R., Dreslik, M. J., & Phillips, C. A. (2008). Roads as barriers to animal

movement in fragmented landscapes. *Animal Conservation*, *11*(4), 288–296.

https://doi.org/10.1111/j.1469-1795.2008.00183.x

Simpson, N. O., Stewart, K. M., Schroeder, C., Cox, M., Huebner, K., & Wasley, T. (2016).

Overpasses and underpasses: Effectiveness of crossing structures for migratory

ungulates. *The Journal of Wildlife Management*, *80*(8), 1370–1378.

850   https://doi.org/10.1002/jwmg.21132

851 Sprogis, K. R., Videsen, S., & Madsen, P. T. (2020). Vessel noise levels drive behavioural

852   responses of humpback whales with implications for whale-watching. *ELife*, *9*,

853   e56760. https://doi.org/10.7554/eLife.56760

854 Strandburg-Peshkin, A., Farine, D. R., Crofoot, M. C., & Couzin, I. D. (2017). Habitat and social

855   factors shape individual decisions and emergent group structure during baboon

856   collective movement. *ELife*, *6*, e19505. https://doi.org/10.7554/eLife.19505

857 Sun, C., Beirne, C., Burgar, J. M., Howey, T., Fisher, J. T., & Burton, A. C. (2021). Simultaneous

858   monitoring of vegetation dynamics and wildlife activity with camera traps to assess

859   habitat change. *Remote Sensing in Ecology and Conservation*, *7*(4), 666–684.

860   https://doi.org/10.1002/rse2.222

861 Sun, J. J., Kennedy, A., Zhan, E., Anderson, D. J., Yue, Y., & Perona, P. (2021). Task

862   Programming: Learning Data Efficient Behavior Representations. *2021 IEEE/CVF*

863   *Conference on Computer Vision and Pattern Recognition (CVPR)*, 2875–2884.

864   https://doi.org/10.1109/CVPR46437.2021.00290

865 Tallamraju, R., Price, E., Ludwig, R., Karlapalem, K., Bülthoff, H. H., Black, M. J., & Ahmad, A.

866   (2019). Active Perception Based Formation Control for Multiple Aerial Vehicles. *IEEE*

867   *Robotics and Automation Letters*, *4*(4), 4491–4498.

868   https://doi.org/10.1109/LRA.2019.2932570

869 Thompson, I. D., Wiebe, P. A., Mallon, E., Rodgers, A. R., Fryxell, J. M., Baker, J. A., & Reid, D.

870   (2015). Factors influencing the seasonal diet selection by woodland caribou (Rangifer

871   tarandus tarandus) in boreal forests in Ontario. *Canadian Journal of Zoology*, *93*(2),

872   87–98. https://doi.org/10.1139/cjz-2014-0140

873 Torney, C. J., Lamont, M., Debell, L., Angohiatok, R. J., Leclerc, L.-M., & Berdahl, A. M. (2018).

874   Inferring the rules of social interaction in migrating caribou. *Philosophical Transactions*

875   *of the Royal Society B: Biological Sciences*, *373*(1746), 20170385.

876   https://doi.org/10.1098/rstb.2017.0385

877   Tuia, D., Kellenberger, B., Beery, S., Costelloe, B. R., Zuffi, S., Risse, B., Mathis, A., Mathis, M.

878   W., van Langevelde, F., Burghardt, T., Kays, R., Klinck, H., Wikelski, M., Couzin, I. D.,

879   van Horn, G., Crofoot, M. C., Stewart, C. V., & Berger-Wolf, T. (2022). Perspectives in

880   machine learning for wildlife conservation. *Nature Communications*, *13*(1), 792.

881   https://doi.org/10.1038/s41467-022-27980-y

882   Walter, T., & Couzin, I. D. (2021). TRex, a fast multi-animal tracking system with markerless

883   identification, and 2D estimation of posture and visual fields. *ELife*, *10*, e64000.

884   https://doi.org/10.7554/eLife.64000

885   Wang, Y., Nickel, B., Rutishauser, M., Bryce, C. M., Williams, T. M., Elkaim, G., & Wilmers, C. C.

886   (2015). Movement, resting, and attack behaviors of wild pumas are revealed by tri-axial

887   accelerometer measurements. *Movement Ecology*, *3*(1), 2.

888   https://doi.org/10.1186/s40462-015-0030-0

889   Weimerskirch, H., Prudor, A., & Schull, Q. (2018). Flights of drones over sub-Antarctic seabirds

890   show species- and status-specific behavioural and physiological responses. *Polar*

891   *Biology*, *41*(2), 259–266. https://doi.org/10.1007/s00300-017-2187-z

892   Williams, H. J., Taylor, L. A., Benhamou, S., Bijleveld, A. I., Clay, T. A., Grissac, S., Demšar, U.,

893   English, H. M., Franconi, N., Gómez- Laich, A., Griffiths, R. C., Kay, W. P., Morales, J. M.,

894   Potts, J. R., Rogerson, K. F., Rutz, C., Spelt, A., Trevail, A. M., Wilson, R. P., & Börger, L.

895   (2020). Optimizing the use of biologgers for movement ecology research. *Journal of*

896   *Animal Ecology*, *89*(1), 186–206. https://doi.org/10.1111/1365-2656.13094

897   Wu, Y., Kirillov, A., Massa, F., Lo, W.-Y., & Girshick, R. (2019). *Detectron2*.

898   https://github.com/facebookresearch/detectron2

899   Yu, H., Deng, J., Nathan, R., Kröschel, M., Pekarsky, S., Li, G., & Klaassen, M. (2021). An

900   evaluation of machine learning classifiers for next-generation, continuous-ethogram

901   smart trackers. *Movement Ecology*, *9*(1), 15. https://doi.org/10.1186/s40462-021-00245-

x

Zhao, Z.-Q., Zheng, P., Xu, S.-T., & Wu, X. (2019). Object Detection With Deep Learning: A

Review. *IEEE Transactions on Neural Networks and Learning Systems*, *30*(11), 3212–

3232. https://doi.org/10.1109/TNNLS.2018.2876865

Zhou, X., Wen, X., Wang, Z., Gao, Y., Li, H., Wang, Q., Yang, T., Lu, H., Cao, Y., Xu, C., & Gao, F.

(2022). Swarm of micro flying robots in the wild. *Science Robotics*, *7*(66), eabm5954.

https://doi.org/10.1126/scirobotics.abm5954

911