Acetaminophen production in the edible, filamentous cyanobacterium *Arthrospira platensis*

Jacob M. Hilzinger[1], Skyler Freidline[1], Divya Sivanandan[1], Ya-Fang Cheng[2], Shunsuke

Yamazaki[1,3], Douglas S. Clark[4,5], Jeffrey M. Skerker[1], Adam P. Arkin[1,6,*]

Author Affiliations:

1-Department of Biological Engineering, University of California-Berkeley, Berkeley, CA

94720, USA

2-QB3-Berkeley, University of California, Berkeley, CA 94720, USA

3-Ajinomoto Co., Inc., 1-1 Suzuki-cho, Kawasaki, Kanagawa 210-8681, Japan

4-Department of Chemical and Biomolecular Engineering, University of California-Berkeley,

Berkeley, CA 94720, USA

5-Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National

Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

6-Environmental Genomics and Systems Biology Division, Lawrence Berkeley National

Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

*- Correspondence should be addressed to A.P.A. (aparkin@lbl.gov)

Keywords: Cyanobacteria, Spirulina, *Arthrospira platensis*

Abstract

Spirulina is the common name for the edible, non-heterocystous, filamentous

cyanobacterium *Arthrospira platensis* that is grown industrially as a food supplement, animal

feedstock, and pigment source. Although there are many applications for engineering this

organism[1–3], until recently no genetic tools or reproducible transformation methods have been

published. While recent work showed the production of single domain antibodies for oral

delivery, no publicly available genetic toolkit was provided[4]. Here, we establish a genetic toolkit

and reproducible method for the transformation of *A. platensis* and engineer this bacterium to

produce acetaminophen as proof-of-concept for small molecule production in an edible host from

CO_2, H_2O, and light. This work opens *A. platensis* to the wider scientific community for future

engineering as a functional food for nutritional enhancement, modification of organoleptic traits,

and production of pharmaceuticals for oral delivery.

Introduction

Arthrospira platensis, commonly referred to and sold commercially as spirulina, is an

edible, non-heterocystous, filamentous cyanobacterium of the order *Oscillatoriales*, with species

of this genus having been used as traditional food sources in Mexico and Chad for centuries[5,6]. *A.*

platensis is considered safe for human consumption and is designated as Generally Recognized

as Safe (GRAS) by the U.S. Food and Drug Administration (GFA GRN No. 417); it has

exceptionally high protein content relative to plants and eukaryotic microalgae[1], and is grown

commercially as a food, pigment source, and animal feedstock[1,6]. Given its enticing food

properties and established commercial growth, as well as applications driven by a photosynthetic

metabolism, there is keen interest in genetically modifying this bacterium for enhanced nutrition,

modified organoleptic traits, improved growth properties, and pharmaceutical production[1,2,4].

While there are clear applications for sustainable biomanufacturing on Earth, the edible

nature and photosynthetic metabolism of *A. platensis* makes it well-suited as an alternate food

source and microbial host for on-demand pharmaceutical production for *in situ* resource

utilization (ISRU)-based missions for human space exploration and habitation[7–9]. As humans

delve deeper into space via the Moon and eventually to Mars and beyond, the need to reduce

mission launch cost, reduce risk, and anticipate pharmaceutical degradation induced by radiation

becomes paramount, further increasing the need for ISRU-driven food and pharmaceutical

production[3,7,10–13]. NASA and ESA have both established the utility of this organism for space

exploration via their biological life support projects, CELSS and MELISSA, respectively, which

used *A. platensis* to photoautotrophically recycle NO_3^- and CO_2 for production of O_2 and

biomass for astronaut consumption[14,15]. Expanding upon its established life support properties to

further produce functional foods for nutrition and pharmaceutical delivery will further increase

the utility of this organism for deep space missions.

A. platensis has historically remained recalcitrant to the development of genetic tools and

a reproducible transformation method[16–19]. Potential reasons range from its multicellularity,

polyploidy, and motility, which combine to inhibit colony formation on plates, and a genome

that encodes a high number of restriction modification systems of types I-IV in addition to

several CRISPR/Cas systems[20,21]. However, in a landmark study, Jester *et al.* (2022) recently

engineered *A. platensis* for antibody production targeting campylobacter infections, and showed

that their orally-delivered strain prevents disease in mice and is safe for human consumption[4].

While this work provided a transformation method, a robust, publicly available genetic toolkit

for pathway engineering of *A. platensis* remains unavailable. This limitation, combined with the

general difficulty of performing genetic manipulations in this bacterium, have so far prevented *A.*

platensis from being engineered for small molecule production.

Here, we develop two reproducible transformation methods based on electroporation and

natural competence, use these methods to develop and characterize a genetic toolkit for

downstream pathway engineering, and engineer *A. platensis* to produce acetaminophen as proof-

of-concept for small molecule production derived from CO_2, H_2O, and light in an edible host for

bioavailable, oral drug delivery.

73　

Results

We initially developed an electroporation method inspired by the patent literature[22] (see

Methods for details). Due to its filamentous and motile nature, *A. platensis* does not form

colonies on plates, and we were unable to obtain filament or cell counts. Thus, we were unable to

quantify plating or transformation efficiency, and were therefore dependent on the binary output

of growth or no growth, as has been observed in *Phormidium lacuna*[23]. Based on the predictions

of Nies *et al.* (2020), we confirmed natural competence in *A. platensis*, and established this

method as our standard transformation procedure.

Cyanobacteria have a wide range of genome copy numbers per cell[24], with reported copy

numbers of the *Oscillatoriales* ranging from 20-90 copies per cell in *Phormidium lacuna*[23] to

nearly 700 in *Trichodesmium erythreaum* IMS 101[25]. For all transformations resulting in

biomass under selection, we observed that transformant biomass encoded the native locus in

addition to the exogenous DNA (data not shown), suggesting an initial integration into the

genome without full segregation through the genome and/or filament. Transformant biomass was

subcultured until no native locus could be detected via PCR and/or genome sequencing. This

took approximately 3-4 months on agar plates and 2-3 months in liquid culture (data not shown).

Jester *et al.* (2022) reported a segregation time of 8-10 weeks for *A. platensis*[4], further supporting

our observations that exogenous DNA must be selected over long timeframes through a

polyploid genome and multi-cell filament.

We established a series of working parts in *A. platensis* NIES-39: one antibiotic

resistance marker, three genomic loci for insertion of exogenous DNA, four endogenous and five

exogenous promoters, and one fluorescent reporter (Supplemental Table 1). All parts were

integrated into a series of suicide vectors for homologous recombination with the genome

(Supplemental Table 2). To screen multiple loci as genomic integration sites, we chose neutral

site I (NSI; NIES39_Q01230), a commonly used site in other cyanobacteria for insertion of

exogenous DNA[26], along with two loci we expected to be involved in motility to potentially

create non-motile strains to aid in plating and genetics.

Twitching motility in cyanobacteria is dependent on Type IV pili and is light regulated

resulting in phototaxis towards or away from light sources[27,28]. Filamentous cyanobacteria from

the *Oscillatoriales* and certain species of the *Nostocales* employ surface-dependent gliding

motility, with the oscillin protein from *Phormidium uncinatum* being essential for this type of

motility[27,29]. We targeted *pilA* (NIES39_C03030), a central component of the Type-IV pilin

system, and an oscillin homolog (NIES39_A01430) as integration sites for exogenous DNA and

motility ablation. We knocked out these three loci with the codon optimized version of *aadA*,

aadA.co, which encodes for streptomycin/spectinomycin (Sm/Sp) resistance (Fig. 1;

Supplemental Table 2). The ΔNSI::P*pilA*-*aadA.co*, Δoscillin::*aadA.co*, and Δ*pilA*::*aadA.co*

strains were confirmed by whole genome sequencing (Fig. 1A). Of note, Δ*pilA*::*aadA.co* mutants

display a "star" morphology when plated on agar, while Δoscillin::*aadA.co* mutants display a

"string" morphology (Fig. 1). This implies that both genes are involved in motility and/or colony

formation on plates.

Deletion of the NSI locus does not result in morphological or growth differences between

the wild type (WT) strain as expected (Fig. 1). Light-based motility assays showed that the WT

and ΔNSI::P*pilA*-*aadA.co* strains moved into the illuminated area of the plate within seven days,

while the Δoscillin::*aadA.co* and Δ*pilA*::*aadA.co* strains remained in the shade where they were

initially spotted (Fig. 1C). While the Δ*pilA*::*aadA.co* strain did not expand beyond its initial spot,

the Δoscillin::*aadA.co* mutant moved in a small halo around the initial spot, indicating that

motility is severely limited by, but not entirely dependent on, the loss of oscillin. This indicates

that both *pilA* and oscillin genes contribute to phototaxis in *A. platensis*.

To test an expression system in *A. platensis*, we designed a series of vectors

(Supplemental Table 2) that resulted in mutant strains at the *pilA* locus that had one of three

native, strong promoters (P_{psaA}, P_{cpcB}, or P_{rbcL})[30] or one of five J23-series promoters driving the

expression of codon optimized *eYFP* (*eYFP.co*) downstream of the native *pilA* promoter driving

the expression of *aadA.co* followed by the native *aadA* terminator (Fig. 2A). The fluorescent

output of each strain was measured, showing orders of magnitude higher fluorescence in the

eight reporter strains than in the WT (Fig. 2B). The J23-series promoters displayed a 4-fold

dynamic range in *A. platensis* compared to a 120-fold dynamic range in both *Synechococcus*

elongatus UTEX 2973 and *Synechocystis* sp. 6803[31]. Although methodological differences

between these studies may explain some of this difference, these results highlight the need for

further development and characterization of *A. platensis* specific genetic parts. Fluorescence was

manually checked under a fluorescence microscope to confirm expression of eYFP.co in the

reporter strains; no fluorescence was detected in the WT strain (Fig. 2C).

For the biosynthesis of acetaminophen (APAP; Fig. 3A), chorismate is converted into

para-aminobenzoic acid (PABA) by the multienzyme complex PabABC, composed of

aminodeoxychorismate synthase (pabAB) and 4-amino-4-deoxychorismate lyase (pabC). PABA

is converted into 4-aminophenol by the 4-aminobenzoate hydrolase (4ABH) from *Agaricus*

bisporus, which is subsequently converted to APAP by N-hydroxyarylamine O-acetyltransferase

(NhoA) from *Escherichia coli*[32]. A series of 38 suicide vectors encoding variations on the

synthetic APAP biosynthesis pathway with codon optimized genes (Fig. 3A) were designed and

screened in *A. platensis* (Supplemental Table 3). Of these, 23 resulted in transformant biomass

that displayed the WT morphology under selection at 1 µg/ml Sm/Sp; no transformants for any

vector could be recovered at higher selection strengths.

For every transformation, a negative control was run to ensure that the selection killed

WT cells, and pJMH024 was transformed as a positive control to ensure that cells were

transformable. PCR screens of genomic DNA isolated from transformant biomass indicated the

presence of *pilA*, *aadA.co*, *4ABH*, and *nhoA* (data not shown), suggesting an initial integration of

the suicide vector into the genome. However, no transformant biomass resulting from any of

these vectors grew after subculturing into the same growth conditions. The only way we could

recover a fully segregated strain, APA16206 (Fig. 3B), was under 16h light:8h dark cycles

illuminated with ~20-30 µmol m^{-2} s^{-1} (µEi) of photosynthetically active radiation (see Methods

for details) for three months until star morphology appeared. Tufts of star morphology were

subcultured into fresh selective medium under the same conditions. Upon reaching mid to late

log phase, biomass was subcultured and grown under 24h light illuminated with ~70-100 µEi.

Biomass was regularly subcultured under these conditions upon reaching mid to late log phase

until fully segregated (Supplemental Fig. 1).

As PABA supplementation was previously shown to increase APAP titer[32], we screened

APAP production in the WT and APA16206 strains with and without the addition of the

precursor compound, PABA (Fig. 3C-E; Supplemental Fig. 2). The WT strain did not produce

APAP under any condition, while the mutant produced APAP only when PABA was added to

the medium (Fig. 3C). The APAP titer and production rate in APA16206 were dependent on the

amount of PABA supplied, with titer and rate increasing with increasing PABA concentrations.

The maximal titer and rate were 2.9 mg L^{-1} and 0.018 mg L^{-1} hr^{-1}, respectively, when supplied

with 2.74 g L^{-1} PABA. Growth rate was slowed by increasing PABA concentration, but did not

impact the final biomass yield, except for two of the replicates supplemented with 2.74 g L^{-1}

PABA, which prematurely reached stationary phase and died (Fig. 3E). The APAP production

rate appeared dependent on the growth rate.

Discussion

Here we independently establish the reproducible transformation of *A. platensis* via

electroporation and natural competence, characterize an initial genetic toolkit for pathway

engineering, and produce APAP as a proof-of-concept pharmaceutical for oral delivery. Our

methods help unlock this bacterium for future engineering efforts for sustainable

biomanufacturing on Earth and ISRU-based food and pharmaceutical production in space by

providing a reproducible blueprint. Building on recent work[4,23], we help establish genetic

tractability in the *Oscillatoriales,* an order of cyanobacteria that have been historically

understudied through genetic methods. Members of this order, such as the diazotrophic genus

Trichodesmium, are major contributors to the biological nitrogen cycle.

180   While our open-source genetic toolkit and proof-of-concept biosynthesis of APAP is an

important step for metabolically engineering *A. platensis*, continued improvement in the genetic

tools available is paramount to decrease the design-build-test-learn cycle time in this bacterium.

Crucial to this endeavor is the development of new promoters to overcome the limited dynamic

range we observed, which may be achieved through the characterization of alternate native

promoters, synthetic promoters engineered from native promoters, or the establishment of

inducible promoters[33]. Operationalizing a CRISPR/Cas system[34] would provide a locus-

independent counterselection that may decrease the segregation time, as the Cas enzyme would

be expected to cut all, or most, genome copies present in the cell, , and would allow for

installation of more complex pathways. Developing self-replicating vectors for the expression of

pathways or a CRISPR/Cas system would bypass the need for genome integration, and allow for

rapid screening of pathway topologies and simpler CRISPR/Cas-based genome engineering.

192   APAP is the first reported exogenous small molecule produced in *A. platensis* and

193   provides a milestone in the long-term efforts to genetically engineer this organism. Given that a

194   dose of APAP is 325 mg, our maximum titer of 2.9 mg/L would require ~112 L for a single dose,

195   assuming complete bioavailability. Thus, higher titer will need to be achieved to make

196   manufacturing a dose of APAP feasible if oral delivery via spirulina is the goal. To this end,

197   there are several routes that should be explored.

198   Of the 38 APAP pathway topologies we transformed into *A. platensis*, we could only

recover one fully segregated mutant that, in turn, could only be recovered under day/night cycles.

This was in stark contrast to the creation of other mutants reported here, which proceeded

without difficulty, and suggests that enzyme toxicity from 4ABH and/or NhoA, due to their

products or the enzymes themselves, inhibited integration and/or segregation of the pathway into

and through the genome. Several additional, unknown chromatography peaks were observed in

APA16206 when PABA was supplied (Fig. 3C), and may contribute to the observed pathway

toxicity. These unknown peaks eluted at similar times as PABA and APAP, implying a similar

chemical structure. These peaks may be due to enzyme promiscuity of 4ABH and/or NhoA, or

from APAP degradation products; these peaks are not from the intermediate 4-aminophenol as

the standard for this molecule produced two peaks at elution times of 2.9 and 3.3 min (data not

shown). If these compounds are produced from enzyme promiscuity, engineering a strain to

produce only APAP would likely increase the titer of APAP by avoiding flux of PABA into

these accessory compounds. This could be done through enzyme engineering, removing

biosynthesis pathways from the genome for substrates that these enzymes may promiscuously

convert, or screening 4ABH and/or NhoA homologs for improved selectivity towards APAP

production. Decreasing the potential pathway toxicity through use of weaker promoters, enzyme

scaffolding, or enzyme engineering for higher product selectivity may make this pathway more

engineerable for future iterations towards a higher titer.

PABA was not detected in either strain without PABA supplementation. This may be due

to natively low flux through PABA or a high rate of PABA turnover. However, while we were

able to identify a homolog for PabAB (NIES39_E04160), we were unable to identify a homolog

for characterized PabC proteins[35–37] in the *A. platensis* genome. While there may be an

uncharacterized PabC, there is a chance that PABA is not natively produced by *A. platensis*,

which would be unexpected as PABA is a precursor for folate. Establishing and/or increasing

endogenous PABA biosynthesis could increase APAP titer, or, crucially, allow for production

without PABA supplementation. As the APAP yield from supplied PABA was ~0.1%, a

sufficient intracellular production rate of PABA may be easily achievable to reach APAP

production rates observed with supplementation. This may also alleviate the observed toxicity

from PABA when supplied at high extracellular concentrations (Fig. 3D). Engineering *A.*

platensis for sufficient PABA production would also remove the economic burden of PABA

supplementation for scale-up.

The genetic toolkit and methods we provide here pave the way for the diversification of

commodity chemicals and biologics to be produced in *A. platensis* beyond APAP and single

domain antibodies[4]. Decisions on which molecules to produce would benefit from a combination

of technoeconomic and life cycle analyses targeting high value products such as pharmaceuticals

and cannabinoids. This work further opens *A. platensis* as an engineerable, functional food.

Modification of organoleptic traits and nutritional properties may lead to more palatable and

nutritious strains for improved consumption by humans or for animal feedstocks. Improved

genetic tools along with host optimization; for example, through the removal of restriction

modification systems, genome copy number control, and morphological control, may further

improve the engineerability of this organism. The present work is thus an important step towards

realizing the potential of an industrially grown, edible, photosynthetic organism for the oral

delivery of pharmaceuticals and nutritional compounds.

Methods

Growth and transformation of *Arthrospira platensis*. A list of strains used in this study can be

found in Supplemental Table 2. WT *A. platensis* NIES-39, obtained from the NIES culture

collection, was grown in Zarrouk's Medium (ZM)[38] at 30°C under ~70-100 µmol photons m^{-2} s^{-1}

(µEi) of photosynthetically active radiation produced by cool, white LEDs (compact LED bars

from Photon Systems Instruments controlled by their LC 200 light controller) with shaking at

200 rpm unless stated otherwise. Working streptomycin/spectinomycin (Sm/Sp) concentrations

ranged from 1-100 µg/ml.

For electroporations, WT *A. platensis* NIES-39 cells were grown to early log phase

(OD_{750} ~0.2-0.3), washed three times with room temperature, sterile water and concentrated to

150 µg Chl *a* ml^{-1} in sterile water. Chl *a* concentration was measured based on the methods of

Meeks and Castenholz (1971)[39]. For each electroporation reaction, 400 µl concentrated cells

were mixed with 10 µg plasmid on ice, transferred to a cold 2 mm electroporation cuvette, and

electroporated at 600 V with a time constant of ~6-7 ms^{-1} on a BTX Gemini electroporator. Cells

were recovered in a final volume of 1 ml ZM, and grown overnight (~15-20 hours) at 30°C under

~20-30 µEi with shaking. The following day, cells were washed once with ZM + Sm/Sp[1], and

plated onto selective 1.5% Bacto agar (BD #214010) plates or into selective liquid ZM in

multiwell (96-, 24-, 12-, and/or 6-well) plates. Biomass was selectively passaged onto plates and

into liquid media, and monitored via PCR and/or Illumina sequencing until fully segregated.

For natural transformations, WT *A. platensis* NIES-39 cells were grown to early log

phase (OD_{750} ~0.2-0.3) and concentrated to 150 µg Chl *a* ml^{-1} in ZM. 100 µl concentrated cells

were mixed with 2.5 µg plasmid DNA, and incubated for 45 min at room temperature. Cells

were recovered in a final volume of 1 ml ZM, and grown overnight (~15-20 hours) at 30°C under

~20-30 µEi with shaking. Recovered cells were added to 99 ml ZM + Sm/Sp (1-10 µg/ml) in

250-ml flasks, and incubated under ~20-30 µEi until growth appeared. Biomass was subcultured

under increasing concentrations of Sm/Sp (up to 100 µg/ml) until fully segregated as monitored

by PCR.

Genomic DNA Extraction. gDNA was extracted as follows: 1-ml aliquots of culture were

harvested by centrifugation. Pelleted cells were resuspended in 300 µl ELB (20 mM Tris-HCl, 2

mM Na-EDTA, 1.2% Triton-X). Cell suspension was incubated at room temperature for 30 min

with 2,500 units of Ready-Lyse lysozyme (Lucigen #R1804M) and 200 µg RNase A (Qiagen

#19101). An additional 2,500 units of lysozyme were added and incubated for 30 min at room

temperature. 15 µl of Proteinase K (20 mg/ml; Qiagen #19157) were added to the suspension and

incubated at 65°C for approximately one hour. 400 µl phenol:chloroform:isoamyl alcohol

(25:24:1 v/v; Sigma #P3803) were mixed with the cell suspension, and centrifuged for 5 min at

21,130 x g. 30 µl of 3M sodium acetate + 0.1M EDTA were added to the aqueous phase,

followed by 660 µL cold 100% ethanol, and centrifuged for 15 min at 21,130 x g. Supernatant

was removed, and the pellet was washed once with 1 mL 70% EtOH. Following removal of the

wash, the pellet was air dried for 5 minutes prior to resuspension in nuclease-free TE buffer

(Thermo Scientific #J75793).

Next Generation Sequencing. Library preparation and 250 bp paired-end sequencing on an

Illumina HiSeq 4000 was completed by the QB3 Genomics, UC Berkeley, Berkeley, CA,

RRID:SCR_022170.

Data analysis. Raw Illumina NovaSeq sequencing reads were processed as follows. Burrows-

Wheeler Alignment tool (BWA)[40] was used to index the reference genome AP011615, align

sequencing reads and generate a sam file. Samtools[41] was used to convert the data to a bam file

and then sort and index the alignments. Picard's DownsampleSam

(https://broadinstitute.github.io/picard/) was used to downsample the BAM files to ~10% of the

total depth for visualization in Geneious. Geneious was used to visualize the alignments and

generate figures.

Motility assays. ZM solidified with 0.5% agar was poured into each lane of a 4-well plate.

Cultures of each strain to be assayed were inoculated to a starting OD_{750} of 0.05 and allowed to

grow until mid-log phase ($OD_{750} = 0.5 - 0.7$). A volume of each culture equal to 750 µL $OD_{750} =$

1.0 was pelleted by centrifugation and the supernatant was removed. The cell pellets were

collected and dispensed at one end of an agar lane in the 4-well plate. The plate was sealed and

imaged immediately. The plate was moved to a 30°C room and placed directly under cool, white

LEDs at a flux of 50 µEi. The half of the plate containing the cell pellets was placed in a

protective covering to prevent direct exposure to light and encourage growth to the far end of the

plate. The plate was imaged every 24 hours for 7 days.

Vector design and synthesis. A list of vectors can be found in Supplemental Table 2. All genes

were codon optimized, purged of the restriction modification (RM) sites (see below),

synthesized, and cloned by Genscript Corp. The remaining parts were synthesized and cloned by

Genscript Corp. The purged RM sites can be found in Supplemental Table 4. This list was

compiled from literature and homology searches[20,42,43].

Fluorescence microscopy. Mid-log phase cultures were imaged on a Zeiss Observer D1

fluorescence microscope. Chlorophyll autofluorescence was imaged under excitation at 587 nm

and emission at 610 nm and eYFP fluorescence was measured under excitation at 495 nm and

emission at 527 nm. Images were falsely colored using Zeiss Zen Pro version 3.5.

Quantification of promoter strengths. *A. platensis* WT and mutant strains were each grown in

eight wells of triplicate 96-well plates under ~20-30 µEi without shaking at 30°C. OD_{750} and

eYFP fluorescence (excitation at 485 nm and emission at 535 nm) were measured daily on a

Tecan Spark M10 plate reader, and these values were normalized to blank measurements.

HPLC. Acetaminophen (APAP) and para-aminobenzoic acid (PABA) were quantified on a

Shimadzu Prominence HPLC system equipped with a photodiode array detector for UV

detection. Compounds were separated on a Zorbax StableBond Plus C18 5 µm with detection at

240 nm. The elution protocol was based on a gradient method using (A) 0.1% phosphoric acid

and (B) methanol as follows: 0 min - 95% A, 5% B; 15 min 100% B; 16-20 min 95% A, 5% B at

a flow rate of 1 ml min^{-1}.

For sample preparation, 1 ml culture was transferred to a BeadBug bead beating tube

(Millipore Sigma #Z763748) and vortexed on high for 10 minutes. Samples were then

centrifuged for 10 min at 14k rpm. 800 µl supernatant were filtered through 13 mm 0.2 µm

PVDF syringe filters (Pall #4406). Standards were prepared in volumetric flasks using APAP

(Sigma #A5000) and PABA (Sigma #A9878), and serially diluted in ZM. The 4-aminophenol

(Sigma #A71328) standard was prepared similarly using methanol as the solvent.

Acknowledgements

This work was supported by the Center for the Utilization of Biological Engineering in Space

(CUBES, https://cubes.space), a NASA Space Technology Research Institute (grant number

NNX17AJ31G) and by NIH S10 OD018174 Instrumentation Grant. We thank Kyle Sander,

Aaron Berliner, Kelly Wetmore, Kelsey Hern, and Yolanda Huang for useful discussions and

support.

Author Contributions

J.M.H., J.M.S., and A.P.A. conceived the project. J.M.H., S.F., J.M.S., and A.P.A. designed

experimental work. J.M.H., S.F., D.S., S.Y., and J.M.S. performed experimental work. J.M.H.,

S.F., J.M.S., and A.P.A. analyzed data. Y.C. developed the HPLC method. J.M.H., S.F., J.M.S.,

D.S.C., and A.P.A wrote the manuscript.

Supplementary Data

Plasmid maps for plasmids listed in Supplementary Tables 2 and 3 are available at

https://doi.org/10.6084/m9.figshare.20085284.

References

1.   Torres-Tiji, Y., Fields, F. J. & Mayfield, S. P. Microalgae as a future food source.

Biotechnol. Adv. **41**, 107536 (2020).

2.   Furmaniak, M. A. *et al.* Edible cyanobacterial genus *Arthrospira*: Actual state of the art in

cultivation methods, genetics, and application in medicine. *Front. Microbiol.* **8**, 1–21

(2017).

3.   Berliner, A. J. *et al.* Towards a Biomanufactory on Mars. **8**, 1–20 (2020).

4.   Jester, B. W. *et al.* Development of spirulina for the manufacture and oral delivery of

protein therapeutics. *Nat. Biotechnol.* **40**, 1-9 (2022).

5.   Ciferri, O. Spirulina, the Edible Microorganism. *Microbiol. Rev.* **47**, 551–578 (1983).

6.   Vonshak, A. *Spirulina plantesis (Arthrospira): Physiology, Cell-biology and*

Biotechnology. (2002).

7.   Nangle, S. N. *et al.* The case for biotech on Mars. *Nature Biotechnology* **38**, 401–407

(2020).

8.   McNulty, M. J. *et al.* Molecular pharming to support human life on the moon, mars, and

beyond. *Crit. Rev. Biotechnol.* **41**, 849–864 (2021).

9.   Mapstone, L. J., Leite, M. N., Purton, S., Crawford, I. A. & Dartnell, L. Cyanobacteria

and microalgae in supporting human habitation on Mars. *Biotechnol. Adv.* **59**, 107946

(2022).

10. Menezes, A. A., Montague, M. G., Cumbers, J., Hogan, J. A. & Arkin, A. P. Grand

challenges in space synthetic biology. *J. R. Soc. Interface* **12**, 20150803 (2015).

11. Menezes, A. A., Cumbers, J., Hogan, J. A. & Arkin, A. P. Towards synthetic biological

approaches to resource utilization on space missions. *J. R. Soc. Interface* **12**, 20140715

(2015).

12. Blue, R. S. *et al.* Supplying a pharmacy for NASA exploration spaceflight: challenges and

current understanding. *npj Microgravity* **5**, 14 (2019).

13. Blue, R. S. *et al.* Limitations in predicting radiation-induced pharmaceutical instability

during long-duration spaceflight. *npj Microgravity* **5**, 15 (2019).

14. Tadros, M. G. Characterization of Spirulina Biomass for CELSS Diet Potential. *NASA*

Contract. **2**–**501**, 53 (1988).

15. Gòdia, F. *et al.* MELISSA: A loop of interconnected bioreactors to develop life support in

Space. *J. Biotechnol.* **99**, 319–330 (2002).

16. Toyomizu, M., Suzuki, K., Kawata, Y., Kojima, H. & Akiba, Y. Effective transformation

of the cyanobacterium Spirulina platensis using electroporation. *J. Appl. Phycol.* **13**, 209–

214 (2001).

17. Kawata, Y., Yano, S., Kojima, H. & Toyomizu, M. Transformation of Spirulina platensis

strain C1 (Arthrospira sp. PCC9438) with Tn5 transposase-transposon DNA-cation

liposome complex. *Mar. Biotechnol.* **6**, 355–363 (2004).

18. Jeamton, W., Dulsawat, S., Tanticharoen, M., Vonshak, A. & Cheevadhanarak, S.

Overcoming intrinsic restriction enzyme barriers enhances transformation efficiency in

Arthrospira platensis C1. *Plant Cell Physiol.* **58**, 822–830 (2017).

19. Dehghani, J. *et al.* Stable transformation of *Spirulina* (*Arthrospira*) *platensis*: a promising

microalga for production of edible vaccines. *Appl. Microbiol. Biotechnol.* **102**, 9267–9278

(2018).

20. Fujisawa, T. *et al.* Genomic structure of an economically important cyanobacterium,

Arthrospira (*Spirulina*) *platensis* NIES-39. *DNA Res.* **17**, 85–103 (2010).

21. Silas, S. *et al.* On the origin of reverse transcriptase- using CRISPR-Cas systems and their

hyperdiverse, enigmatic spacer repertoires. *MBio* **8**, e00897-17 (2017).

22. Takeuchi, R. & Roberts, J. Targeted Mutagenesis in Spirulina. (2017).

23. Nies, F., Mielke, M., Pochert, J. & Lamparter, T. Natural transformation of the

filamentous cyanobacterium *Phormidium lacuna*. *PLoS One* **15**, 1–18 (2020).

24. Griese, M., Lange, C. & Soppa, J. Ploidy in cyanobacteria. *FEMS Microbiol. Lett.* **323**,

124–131 (2011).

25. Sargent, E. C. *et al.* Evidence for polyploidy in the globally important diazotroph

Trichodesmium. *FEMS Microbiol. Lett.* **363**, 1–7 (2016).

26. Taton, A. *et al.* Broad-host-range vector system for synthetic biology and biotechnology

in cyanobacteria. *Nucleic Acids Res.* **42**, e136 (2014).

27. Brahamsha, B. & Bhaya, D. Motility in Unicellular and Filamentous Cyanobacteria. *The*

Cell Biology of Cyanobacteria 233–262 (2013).

28. Schuergers, N., Mullineaux, C. W. & Wilde, A. Cyanobacteria in motion. *Curr. Opin.*

Plant Biol. **37**, 109–115 (2017).

29. Hoiczyk, E. & Baumeister, W. Oscillin, an extracellular, Ca2+-binding glycoprotein

essential for the gliding motility of cyanobacteria. *Mol. Microbiol.* **26**, 699–708 (1997).

30. Wang, B., Wang, J. & Meldrum, D. R. Application of synthetic biology in cyanobacteria

and algae. *Front. Microbiol.* **3**, 1–15 (2012).

31. Vasudevan, R. *et al.* Cyanogate: A modular cloning suite for engineering cyanobacteria

based on the plant moclo syntax. *Plant Physiol.* **180**, 39–55 (2019).

32.   Anderson, J. C. Biosynthetic production of acetaminophen, p-aminophenol, and p-
aminobenzoic acid. (2017).

33.   Behle, A., Saake, P., Germann, A. T., Dienst, D. & Axmann, I. M. Comparative Dose-
Response Analysis of Inducible Promoters in Cyanobacteria. *ACS Synth. Biol.* **9**, 843–855
(2020).

34.   Ungerer, J. & Pakrasi, H. B. Cpf1 Is A Versatile Tool for CRISPR Genome Editing
Across Diverse Species of Cyanobacteria. *Sci. Rep.* **6**, 1–9 (2016).

35.   Green, J. M., Merkel, W. K. & Nichols, B. P. Characterization and sequence of
Escherichia coli *pabC*, the gene encoding aminodeoxychorismate lyase, a pyridoxal
phosphate-containing enzyme. *J. Bacteriol.* **174**, 5317–5323 (1992).

36.   Satoh, Y., Kuratsu, M., Kobayashi, D. & Dairi, T. New gene responsible for para-
aminobenzoate biosynthesis. *J. Biosci. Bioeng.* **117**, 178–183 (2014).

37.   de Crécy-Lagard, V., El Yacoubi, B., de la Garza, R. D., Noiriel, A. & Hanson, A. D.
Comparative genomics of bacterial and plant folate synthesis and salvage: Predictions and
validations. *BMC Genomics* **8**, 1–15 (2007).

38.   Madkour, F. F., Kamil, A. E. W. & Nasr, H. S. Production and nutritive value of Spirulina
platensis in reduced cost media. *Egypt. J. Aquat. Res.* **38**, 51–57 (2012).

39.   Meeks, J. C. & Castenholz, R. W. Growth and photosynthesis in an extreme thermophile,
Synechococcus lividus (*Cyanophyta*). *Arch. Mikrobiol.* **78**, 25–41 (1971).

40.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

41.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, 1–4 (2021).

42.   Shiraishi, H. & Tabuse, Y. The *Apl* I Restriction-Modification System in an Edible
Cyanobacterium, *Arthrospira* (*Spirulina*) *platensis* NIES-39, Recognizes the Nucleotide
Sequence 5′-CTGCAG-3′. *Biosci. Biotechnol. Biochem.* **77**, 782–788 (2013).

43.   Roberts, R. J., Vincze, T., Posfai, J. & Macelis, D. REBASE-a database for DNA
restriction and modification: Enzymes, genes and genomes. *Nucleic Acids Res.* **43**, D298–
D299 (2015).

Figure 1. Confirmation and phenotyping of deletion mutants at three genomic loci. (**A**) Downsampled Illumina coverage for each genomic region that was swapped for *aadA.co*. (**B**) Images showing colony morphology on agar plates for the WT and each deletion strain. (**C**) Images depicting colony position on agar plates at Day 0 (top) and Day 7 (bottom) for the WT and each deletion strain. Strains were plated in the dark, and light was available to the upper half of the plate.

Figure 2. Fluorescent reporter and promoter characterization. (**A**) Schematic of the mutant genotypes for *eYFP.co* expression at the *pilA* locus: P$_{pilA}$ drives expression of *aadA.co*, which is terminated by T$_{aadA}$. Downstream, one of three native promoters drives expression of the *eYFP.co* gene. (**B**) Relative promoter strength for three native *A. platensis* promoters and the five J23-series promoters. Error bars represent standard deviations of technical replicates. (**C**) Fluorescence microscopy of the Δ*pilA*::*aadA.co*-P*psaA*-*eYFP.co* and WT strains showing Chl *a* autofluorescence of filaments under excitation at 587 nm, and YFP fluorescence under excitation at 485 nm.

Figure 3. Figure 3. Acetaminophen (APAP) production in *A. platensis* APA16206. (A) The pathway for APAP production from para-aminobenzoic acid (PABA) uses the enzyme 4ABH to convert PABA to 4-aminophenol, which is then converted to APAP by NhoA. (B) Schematic for the APAP production pathway at the *pilA* locus in APA16206. (C) HPLC chromatograms showing peaks eluting between 6 and 10.2 min for PABA and APAP standards along with APA16206 and WT strains with and without PABA supplementation. (D) APAP titer over time in APA16206 when supplemented with 0.69, 1.37, and 2.74 g L^{-1} PABA. (E) Growth of APA16206 with and without PABA supplementation at 0, 0.69, 1.37, and 2.74 g L^{-1}.