Pseudomonas aeruginosa transcriptome analysis in a cystic fibrosis lung model reveals metabolic changes accompanying biofilm maturation and increased antibiotic tolerance over time.

Niamh E. Harrington[1,2]*, Freya Allen[1,3], Ramón Garcia-Maset[4], Freya Harrison[1]*

1: School of Life Sciences, Gibbet Hill Campus, The University of Warwick, Coventry, CV4 7AL, UK.

2: Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, L69 7ZB, UK.

3: Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham, Birmingham, B15 2TT, UK.

4: Warwick Medical School, Gibbet Hill Campus, The University of Warwick, Coventry, CV4 7AL, UK.

* Co-corresponding authors: Niamh.Harrington@liverpool.ac.uk, F.Harrison@warwick.ac.uk

Keywords: Biofilm, chronic infection, cystic fibrosis, *ex vivo* model, *Pseudomonas aeruginosa,* RNA sequencing, transcriptome, quorum sensing, antimicrobial resistance, sulfur metabolism

Abstract

The opportunistic pathogen *Pseudomonas aeruginosa* forms biofilm infections in the lungs

of people with the genetic condition cystic fibrosis (CF) that can persist for decades. There

are numerous *P. aeruginosa* lifestyle changes associated with chronic biofilm infection cued

by the CF lung environment. These include a loss of virulence, metabolic changes, and

increased antimicrobial tolerance. We have investigated *P. aeruginosa* PA14 biofilm

infection over 7 d in an *ex vivo* pig lung (EVPL) model for CF, previously shown to facilitate

formation of a clinically-relevant *P. aeruginosa* biofilm structure with gene expression

comparable to human infection. We extracted and sequenced *P. aeruginosa* RNA from

EVPL-associated biofilms 24 h, 48 h, and 7 d after infection, and compared gene expression

between sequential time points. We also investigated tolerance to polymyxins across these

three time points. Our results indicate that the EVPL model can maintain a *P. aeruginosa*

biofilm population, which exhibits increased antibiotic tolerance, for at least 7 d. Differential

expression of antimicrobial resistance-associated genes was not observed, however there

was significant upregulation of sulfur metabolism and phenazine biosynthesis-related genes,

as well as maintenance of a structured biofilm. These findings provide further insight into

the incidence of increased *P. aeruginosa* antibiotic tolerance during infection of the CF lung,

and the gene expression changes that arise as chronic infection establishes.

Introduction

Pseudomonas aeruginosa chronic biofilm infections of the cystic fibrosis (CF) lung are

characterised by physiological adaptations to the environment over years of infection,

leading to reduced phenotypic diversity and loss of virulence in late stage infections [1].

Specific lifestyle changes have been shown to improve *P. aeruginosa* persistence, including

loss of swimming motility, impaired pyoverdine production, and increased adherence [2]. *P.*

aeruginosa adaptation is driven by the selection pressures of the unique infection

environment, including osmotic, nitrosative, and oxidative stress, as well as interactions

with the microbiome and co-infecting pathogens, and frequent antibiotic exposure [3]. *P.*

aeruginosa chronic infections in the CF lung are almost impossible to eradicate once formed

due to the high level of antibiotic resistance exhibited, contributed to by the formation of a

biofilm matrix that protects the bacterial cells from not only the host immune response, but

also antibiotics [4].

P. aeruginosa anaerobic respiration during chronic infection has also been associated with

increased antibiotic tolerance [5]. There is a build-up of thick mucus in the CF lung that is

not effectively cleared, which results in mucus plugs that block the airways. Alongside the

oxygen reduction caused by an abundance of active host immune cells [6], this creates an

oxygen gradient with regions of low oxygen availability (hypoxic regions) or no oxygen

(anoxic regions) [7, 8]. *P. aeruginosa* has been found to occupy a hypoxic niche in the CF

lung, and the oxygen gradient is then further steepened by cell proliferation [9]. Although

there has been research to show that *P. aeruginosa* is able to use an oxygen supply in the

lumen to respire via microaerobic respiration at the surface of the mucus layer [10],

anaerobic respiration is believed to be the predominant mechanism of respiration as growth

is mainly in hypoxic regions [5].

The mechanisms of respiration, driven by the CF lung environment, also have an impact on

the changes in *P. aeruginosa* growth and physiology over time. For example, sulfur

metabolism has been linked to *P. aeruginosa* growth in anoxic conditions. Traditionally

sulfur starvation has been considered the cause of sulfur metabolism upregulation, however

a study of *P. aeruginosa* PA14 growth in synthetic cystic fibrosis sputum media (SCFM)

found that upregulation of sulfur metabolism in PA14 anaerobic biofilms was induced by the

lack of oxygen [11]. There is actually an increase in sulfation of the airway mucins in CF

compared with a non-CF lung [12, 13], in part caused by inflammation [14], that chronically

adapted *P. aeruginosa* can use. It has also been shown that there is enough inorganic sulfate

in CF sputum to facilitate growth even without the use of mucin as a source [15], thus sulfur

is in abundance in this environment.

Quorum sensing (QS) largely controls the phenotypic changes observed in the *P. aeruginosa*

biofilm population during infection. It is involved in the development of a biofilm with

mature architecture and is estimated to influence expression of approximately 10% of all *P.*

aeruginosa genes [16]. Expression of QS-related genes declines over time in CF, linked to the

reduced virulence factor production observed in chronic infection [17]. However, QS-

associated gene expression in *in vitro* models has been shown to be distinct from CF sputum

[18]. Our previous RNA-sequencing analysis of *P. aeruginosa* in an *ex vivo* pig lung (EVPL)

model of CF infection demonstrated that this growth environment may more accurately

capture *in vivo* QS regulated gene expression than *in vitro* growth in CF sputum-mimicking

medium [19]. The model uses EVPL bronchiolar tissue surrounded by synthetic cystic fibrosis

sputum media (SCFM) to recapitulate the CF lung environment [20]. We now build upon our

previously published work to investigate *P. aeruginosa* PA14 gene expression in the EVPL

model biofilm over 7 days, to determine the changes in expression that drive the formation

of a chronic biofilm infection in the CF lung.

Extended analysis of an RNA-sequencing data set, which we partially explored in our

previous study [19], showed that there are distinct *P. aeruginosa* transcriptomes at 24 h, 48

h, and 7 d in the EVPL-associated biofilm. Through differential expression analysis and Kyoto

encyclopedia of genes and genomes (KEGG) pathway enrichment analysis, we identified a

number of genes and pathways that significantly change in expression over 7 d. We also

investigated *P. aeruginosa* PA14 tolerance to two clinically-relevant antibiotics, colistin and

polymyxin B, and found that tolerance increases over time. This is not accompanied by

upregulation of specific antimicrobial resistance (AMR) genes but by other factors, including

upregulation of sulfur metabolism and the formation and maintenance of a biofilm. Hence,

these results provide new insight into the gene expression changes that occur as *P.*

aeruginosa antibiotic tolerance increases in the CF lung.

Results

P. aeruginosa PA14 was grown as a biofilm on the surface of EVPL bronchiolar tissue for 24

h, 48 h, and 7 d. The biofilm was recovered from three tissue pieces, from each of two

independent pig lungs, per time point. *P. aeruginosa* RNA was then extracted, and RNA

sequencing was performed. The colony forming units (CFU) lung^{-1} were reported in our

previous study, in which we compared PA14 gene expression in the EVPL model with SCFM

in vitro growth at 24 h and 48 h [19]. In the present study, we carried out differential

expression analysis to determine which *P. aeruginosa* PA14 genes that were significantly

differentially expressed ($P < 0.05$, log$_2$ fold change (log$_2$FC) $\geq$ |1.5|) over 24 h, 48 h, and 7 d

in the EVPL-associated biofilm. Each contrast was between two time points, and the earliest

time point was used as the baseline for comparison. For example, the contrast between 24

h and 48 h determined the log$_2$FC in expression of genes at 48 h compared with 24 h,

showing the change in expression from 24 h to 48 h.

P. aeruginosa PA14 had a distinct transcriptome in the EVPL biofilm at 24 h, 48 h, and 7 d,

with the majority of genes significantly overexpressed over time.

Principal component analysis (PCA) of PA14 gene expression at each time point identified a

distinct transcriptome dependent on the time *P. aeruginosa* was grown on the EVPL tissue

for; samples from the same time point clustered together (Figure 1). The first two principal

components captured most of the variance between samples; principal component three

only accounted for 6% of the variance and its inclusion did not change the pattern observed

(Figure S1). There was also no clear difference in samples from different lungs, indicating

that these were true differences between time points and were not caused by lung

variation. *P. aeruginosa* grown for 24 h appeared to be the most distinct in principal

component 1, which accounted for 43% of the variance (Figure 1), showing that *P.*

aeruginosa PA14 gene expression in the EVPL-associated biofilm may have been more

similar at 48 h and 7 d than 24 h. This was also observed with hierarchical clustering analysis

and Pearson's correlation coefficient analysis (Figure S2).

Figure 1. Principal component analysis (PCA) plot of the *Pseudomonas aeruginosa* PA14 transcriptome, considering all genes (n = 5829), in the *ex vivo* pig lung (EVPL) model biofilm at 24 h, 48 h, and 7 d. RNA was sequenced from the lung-associated biofilm at each time point from three repeats, from each of two independent pig lungs, shown by individual data points. Each time point is shown by a different shaped data point, and the individual pig lungs are shown by different colours (see key). The 95% confidence ellipses are also shown.

Gene expression was then compared between pairs of time points to identify genes that were significantly over- or underexpressed over time. There were more significantly overexpressed genes than underexpressed genes when comparing sequential time points, with the greatest number of significantly overexpressed genes at 7 d compared with 24 h (267 genes; Figure 2A). Furthermore, the 24 h to 48 h contrast had the greatest number of unique significant differentially expressed genes (DEGs) (Figure 2B). This further indicates

that there was a greater difference in gene expression at 24 h compared with 48 h and 7 d,

than from 48 h to 7 d.

Figure 2. The number of *Pseudomonas aeruginosa* PA14 significant differentially expressed genes (DEGs) (*P* <

0.05, log$_2$ fold change ≥ |1.5|) in each time contrast in the *ex vivo* pig lung (EVPL) model biofilm. RNA was

sequenced from three tissue-associated biofilms from each of two independent pig lungs, and differential

expression analysis was performed comparing three time points: 24 h, 48 h, and 7 d. **(A)** Bar charts showing

the number of significantly overexpressed genes (pink, left bar) and significantly underexpressed genes

(yellow, right bar) in each time contrast. Significant DEGs were determined at 48 h compared with 24 h (24 h

to 48 h), at 7 d compared with 48 h (48 h to 7 d), and at 7 d compared with 24 h (24 h to 7 d). **(B)** A Venn

diagram of the number of significant DEGs. The shared genes were either significantly overexpressed or

underexpressed in all contrasts compared. Genes that were significant in multiple contrasts but in opposing

directions were not considered to be shared.

Sulfur metabolism was upregulated from 24 h to 7 d in the EVPL-associated biofilm.

An overview of the functional implications of the significant DEGs found was investigated

using KEGG enrichment analysis. The analysis identified KEGG pathways that were

significantly enriched ($P < 0.05$) for each time contrast. Although overall there was least

difference in gene expression from 48 h to 7 d (Figure 2), some KEGG pathways were found

to be significantly enriched (see Figure S3); these were predominantly associated with the

upregulation of phenazine biosynthesis. There were no *P. aeruginosa* PA14 KEGG pathways

significantly enriched from 24 h to 48 h, however there were four pathways significantly

enriched from 24 h to 7 d: sulfur metabolism, ATP-binding cassette (ABC) transporters,

degradation of aromatic compounds, and nitrogen metabolism. The majority of the

significant DEGs associated with three of these pathways were overexpressed, indicating

that they were upregulated over 7 d (Figure 3).

 Figure 3. Significantly enriched (*P* < 0.05) *Pseudomonas aeruginosa* PA14 Kyoto encyclopedia of genes and

182 genomes (KEGG) pathways in the *ex vivo* pig lung-associated biofilm at 7 d compared with 24 h. Each bar

shows the percentage of genes associated with each pathway that were significantly differentially expressed (*P*

< 0.05, log$_2$ fold change ≥ |1.5|). The overexpressed genes (left bar) and underexpressed genes (right bar) are

show by different colour bars (see key). *P. aeruginosa* PA14 biofilms associated with three tissue pieces from

each of two independent pig lungs were studied at each time point.

The KEGG pathway with the greatest percentage of associated genes significantly

differentially expressed from 24 h to 7 d was sulfur metabolism (Figure 3); 47.06% of sulfur

metabolism genes were significantly overexpressed in the EVPL biofilm by 7 d. This included

the sulfate transport genes *cysA*, *cysW*, and *cysT* (3.04, 2.87, and 3.14 log$_2$FC respectively).

There was only one associated gene significantly underexpressed at 7 d: the sulfite/nitrate

reductase PA14_10550 (-1.56 log$_2$FC). The majority of the significantly differentially

expressed ABC transporter genes were also involved in sulfur metabolism. Additionally,

three of the ABC transporter genes significantly overexpressed from 24 h to 7 d were part of

the phosphate-specific transporter PstSCAB: *pstB* (1.52 log$_2$FC), *pstA* (1.83 log$_2$FC), and *pstC*

(1.88 log$_2$FC).

Nitrogen metabolism was the only significantly enriched KEGG pathway where more

significant DEGs were underexpressed than overexpressed from 24 h to 7 d (Figure 3).

However, it was not clear whether this pathway was upregulated or downregulated over

time as a similar number of genes were significantly overexpressed and underexpressed

(8.82% and 14.71% respectively). Expression of genes in the *P. aeruginosa* nitrogen

metabolism operons responsible for a shift towards anaerobic respiration in CF sputum (*nor*,

nir, *nap*, and *nos*, [21]) was investigated, but there was no clear pattern of upregulation or

downregulation (Table S1).

There were seven gene expression profiles in the EVPL *P. aeruginosa* PA14 biofilm

population, associated with a number of KEGG pathways including sulfur metabolism.

The gene expression profiles of the significant DEGs were then determined, to investigate

how gene expression changed over the three time points in the EVPL-associated biofilm.

Expression profiles were identified based on when expression changes were shared, unique,

or non-significant between the time point comparisons. Table 1 describes each of the seven

expression profiles found for *P. aeruginosa* PA14 in the EVPL-associated biofilm from 24 h to

48 h, to 7 d.

 Table 1. *Pseudomonas aeruginosa* **PA14 gene expression profiles identified in the *ex vivo* pig lung-associated**

 biofilm at 24 h, 48 h, and 7 d. Differential expression analysis was performed using three repeats from each of

 two independent pig lungs. Contrasts were performed at 48 h compared with 24 h, 7 d compared with 48 h,

 and 7 d compared with 24 h. The changes in expression of each significant differentially expressed gene found

 in any contrast was investigated across all three time points, to determine the expression changes over time.

 The seven different expression profiles found were separated into groups, described in the table. The graphical

 representations demonstrate the changes in expression of genes over the three time points. Below each graph

 is the number of significant differentially expressed genes found to follow that expression profile in the lung-

 associated *P. aeruginosa* PA14 biofilm.

Profile number	Gene expression profile	Graphical representation
1	Significant change in expression 24 h to 48 h, then a significant change in the **same** direction 48 h to 7 d. Results in significant change 24 h to 7 d.	3 genes 1 gene
2	Significant change in expression 24 h to 48 h, which is **maintained** 48 h to 7 d (no significant change). Results in significant change 24 h to 7 d.	80 genes 68 genes
3	No significant change in expression 24 h to 48 h, then a significant change 48 h to 7 d. Results in significant change 24 h to 7 d.	66 genes 20 genes
4	Significant change in expression 24 h to 48 h, then a small but not significant change in the **opposite** direction 48 h to 7 d. Results in no significant change 24 h to 7 d.	157 genes 79 genes
5	Small but not significant change in expression 24 h to 48 h, then a small but not significant change in the **same** direction 48 h to 7 d. Results in overall significant change 24 h to 7 d.	118 genes 79 genes
6	Small but not significant change in expression 24 h to 48 h, then a significant change in the **opposite** direction 48 h to 7 d. Results in no significant change 24 h to 7 d.	20 genes 51 genes

| 7 | Significant change in expression 24 h to 48 h, then a significant change in the **opposite** direction 48 h to 7 d. Results in no significant change 24 h to 7 d. |

 |

____

There were four genes that were significantly differentially expressed in the same direction

from 24 h to 48 h, and 48 h to 7 d (Figure 2B), which followed expression profile 1 (Table 1).

Further investigation of these genes did not indicate any specific pathways following this

steep increase or decrease in expression (Table S2). The greatest number of significant DEGs

followed expression profile 4 (Table 1). These genes were only significantly differentially

expressed 24 h to 48 h, then a non-significant change in expression in the opposite direction

48 h to 7 d resulted in no overall significant difference 24 h to 7 d (Table 1).

KEGG enrichment analysis was performed for each of the gene expression profiles in the *P.*

aeruginosa PA14 EVPL-associated biofilm populations. Significantly enriched KEGG pathways

($P < 0.05$) were found for three of the profiles (Figure 4). Consistent with the previous

finding, sulfur metabolism was found to be significantly enriched and the associated genes

followed two expression profiles: 2 and 5. These genes were overexpressed following both

expression profiles; either upregulation from 24h to 48 h that was maintained to 7 d (Figure

2A), or a gradual upregulation (Figure 2B). The degradation of aromatic compounds and

benzoate degradation KEGG pathways were similarly significantly enriched following

expression profile 5 (Figure 4B). All significant DEGs in the degradation of aromatic

compounds pathway were part of the benzoate degradation pathway. Arginine biosynthesis

was also significantly enriched following profile 5 (Figure 4B); two of the significantly

associated underexpressed genes were arginine fermentation enzymes [22]: *arcB* (-2.25

log$_2$FC) and *arcC* (-2.16 log$_2$FC), which have been shown to be induced under stress [23, 24].

Figure 4. Significantly enriched (*P* < 0.05) *Pseudomonas aeruginosa* PA14 Kyoto encyclopedia of genes and

genomes (KEGG) pathways found for gene expression profiles in the *ex vivo* pig lung-associated biofilm over

24h, 48 h, and 7 d. The expression of significantly differentially expressed genes over the three time points

was investigated and separated into seven different profiles, three of which were found to involve sufficient

genes associated with KEGG pathways for pathways to be significantly enriched. The bars show the percentage

of significant differentially expressed genes associated with each enriched pathway (*P* < 0.05, log$_2$ fold change

≥ |1.5|); those overexpressed (left bar) and underexpressed (right bar) are shown by different colours (see

key). The graphical representation of the gene expression profile is shown in the top right of each graph; the

left graph shows the profile of overexpressed genes, and the right graph shows the profile of underexpressed genes. Three repeats from each of two independent pig lungs were studied at each time point. **(A)** Significantly enriched KEGG pathway following expression profile 2, where there was a significant change 24 h to 48 h, which was maintained 48 h to 7 d resulting in a significant change 24 h to 7 d. **(B)** Significantly enriched KEGG pathways following profile 5, where there was a significant change in expression in the direction specified 24 h to 7 d, caused by non-significant changes in the same direction 24 h to 48 h, and 48 h to 7 d. **(C)** Significantly enriched KEGG pathways following profile 6, where there was a significant change in the direction specified at 7 d compared with 48 h.

Gene expression profile 6 was composed of genes that were only significantly differentially expressed from 48 h to 7 d (Table 1). Three KEGG pathways were significantly enriched following this expression profile, including phenazine biosynthesis and QS which were upregulated (Figure 4C). All the significant differentially expressed genes associated with phenazine biosynthesis were also associated with the QS pathway.

P. aeruginosa PA14 expression of genes associated with antimicrobial resistance was reduced over time in the EVPL biofilm but tolerance to colistin and polymyxin B increased.

We previously found significant differential expression of AMR genes when _P. aeruginosa_ PA14 was grown in the EVPL model compared with SCFM _in vitro_, at both 24 h and 48 h [19]. We were therefore interested in investigating whether changes in AMR gene expression occurred in the EVPL-associated biofilm over 7 d infection. Analysis of AMR gene expression found three genes significantly underexpressed at either 48 h or 7 d compared with 24 h (Table 2), and no genes significantly overexpressed. The full AMR gene list studied is shown in Table S3. The _brc-1_ gene, which encodes a major facilitator superfamily (MFS) transporter

associated with bicyclomycin resistance [25], was significantly underexpressed from 24 h to

7 d. The other two significant DEGs found were significantly underexpressed from 24 h to 48

h. This included *mexG* and *arnA*, which encode a transmembrane protein associated with

multiple efflux pumps and a bifunctional UDP-glucuronic acid decarboxylase/UDP-4-amino-

4-deoxy-L-arabinose formyltransferase [3], respectively. More specifically, the enzyme

encoded by *arnA* is involved in lipid A modifications, linked to cationic antimicrobial peptide

(CAMP) resistance [26, 27].

Table 2. *Pseudomonas aeruginosa* PA14 antibiotic resistance associated genes found to be significantly

differentially expressed in at least one contrast in the *ex vivo* pig lung-associated biofilm between 24 h, 48 h,

and 7 d. Differential expression analysis was performed on three repeats from each of two independent pig

lungs at 48 h compared with 24 h, 7 d compared with 48 h, and 7 d compared with 24 h. The gene expression

group denotes the expression profile the gene belonged to (see Table 1). The $\log_2$ fold change ($\log_2$FC) in the

contrast where the gene was significantly differentially expressed is shown. A gene was considered

significantly differentially expressed if $\log_2$FC ≥ |1.5| and $P < 0.05$.

Locus tag	Gene name	Gene product	Gene expression group	Time: $\log_2$FC
PA14_09540	*mexG*	Transmembrane protein	4	24 h to 48 h: -1.52
PA14_18350	*arnA*	Bifunctional UDP-glucuronic acid decarboxylase/UDP-4-amino-4-deoxy-L-arabinose formyltransferase	4	24 h to 48 h: -1.59
PA14_10470	*bcr-1*	MFS transporter	5	24 h to 7 d: -2.49

To determine how AMR gene expression compared to phenotype, *P. aeruginosa* PA14 was

grown as a biofilm on the EVPL tissue for 24 h, 48 h or 7 d then exposed to one of two

clinically relevant polymyxins, polymyxin B or colistin, for a further 24 h. The CFU lung^{-1} was

determined and compared with PA14 EVPL biofilms treated with phosphate-buffered saline

(PBS) for the second 24 h incubation as a negative control (Figure 5). Samples initially grown

for 24 h prior to treatment did not exhibit any growth when exposed to 256 µg ml^{-1} colistin

(Figure 5A). After 48 h initial biofilm growth, some samples were able to survive following

colistin exposure (Figure 5B). However, after 7 d growth followed by colistin exposure, the

P. aeruginosa PA14 biofilm populations were able to tolerate 256 µg ml^{-1} colistin and

maintain a CFU lung^{-1} comparable to the untreated samples (Figure 5C). A similar pattern

was observed following 256 µg ml^{-1} polymyxin B treatment; PA14 biofilm tolerance

increased with time following initial infection (Figure 5), and 7 d biofilms were consistently

able to survive treatment (Figure 5C). The pattern of increased polymyxin tolerance with

biofilm age is a contrast to the reduced expression of AMR-associated genes observed,

suggesting indirect factors may be involved.

Figure 5. The colony forming units per lung tissue piece (CFU lung^{-1}) for the *Pseudomonas aeruginosa* PA14

tissue-associated biofilm of different ages following no antibiotic treatment, treatment with 256 µg ml^{-1}

colistin, or 256 µg ml^{-1} polymyxin B. Three replicates from each of two independent pig lungs, shown by

different shaped data points (see key), were used. The *P. aeruginosa* PA14 biofilms were grown in the *ex vivo*

pig lung (EVPL) model for **(A)** 24 h **(B)** 48 h **(C)** 7 d, then exposed to either phosphate-buffered saline (PBS) as a

negative control (no antibiotic; pink), 256 µg ml^{-1} colistin (blue), or 256 µg ml^{-1} polymyxin B (yellow) for a

further 24 h. The CFU lung^{-1} was then determined and is shown on the graphs, data points below the dashed

horizontal line represent samples where no CFU were detected. The asterisks represent a significant difference

between treatments determined using post-hoc Tukey HSD tests following ANOVA (* = $P < 0.05$). **(A)** Day 1

ANOVA: treatment $F_{2,12} = 30.32$, $P < 0.01$; lung $F_{1,12} = 0.15$, $P = 0.71$; interaction $F_{2,12} = 0.04$, $P = 0.96$. **(B)** Day 2

ANOVA: treatment $F_{2,12}$ = 10.53, P < 0.01; lung $F_{1,12}$ = 5.76, P = 0.03; interaction $F_{2,12}$ = 1.26, P = 0.32. **(C)** Day 7

ANOVA: treatment $F_{2,12}$ = 14.51, P < 0.01; lung $F_{1,12}$ = 4.12, P = 0.07; interaction $F_{2,12}$ = 2.30, P = 0.14.

There was significantly reduced expression of Pel biosynthesis genes from 24 h to 48 h in

the EVPL-associated biofilm, but a dense, structured biofilm was clear at 48 h.

QS and biofilm formation pathways were then investigated, as both are involved in the

establishment of a chronic *P. aeruginosa* infection in the CF lung and may have been

associated with the increased antibiotic tolerance observed. As shown in Figure 6, there was

no overall difference in QS gene expression from 24 h to 48 h. This was similarly observed

from 48 h to 7 d (Figure S4), and from 24 h to 7 d (Figure S5). Other genes associated with *P.*

aeruginosa PA14 virulence factor production were also investigated, including those

involved in motility, type III secretion, type VI secretion, and alginate biosynthesis. However,

there was no significant differential expression found in the lung-associated biofilm over the

three time points for any of these processes.

Figure 6. Diagram of the gene expression changes in the *Pseudomonas aeruginosa* PA14 Kyoto encyclopedia of genes and genomes (KEGG) biofilm formation and quorum sensing (QS) pathways in the *ex vivo* pig lung-associated biofilm from 24 h to 48 h. Three repeats from each of two independent pig lungs were studied at each time point. The diagram shows all genes involved in each pathway, including associated signalling

pathways and the Pel exopolysaccharide biosynthesis pathway. The $\log_2$ fold change value from 24 h to 48 h is

shown below each gene name. The genes shown in red were significantly underexpressed, and genes shown in

green were significantly overexpressed (significance: $P < 0.05$, $\log_2$ fold change $\geq |1.5|$). The blocked arrows

represent gene activation, and the dashed arrows show indirect gene activation. The solid, blocked lines

(ending with |, not an arrow) show inhibition, and the dashed, blocked lines show indirect inhibition. The blue

'P' represents phosphorylation.

There were only two genes found to be significantly overexpressed from 24 h to 48 h in

these pathways, including *lecB* (Figure 6). LecB is the second *P. aeruginosa* internal lectin

(PA-IIL) that has been shown to act as a biofilm matrix protein, and is associated with

increased cell and exopolysaccharide retention in the biofilm [28]. The anti-sigma factor

antagonist HsbA, associated with RpoS regulation [29], was also found to be significantly

overexpressed from 24 h to 48 h. However, there were no other genes in this pathway

found to be significant differentially expressed (Figure 6), indicating that this differential

expression may not have had an effect. In contrast, five out of the seven genes in the

operon for biosynthesis of the biofilm exopolysaccharide Pel were found to be significantly

underexpressed from 24 h to 48 h (Figure 6). The other two genes were also

underexpressed but below the $\log_2$FC threshold considered significant. This downregulation

was maintained for the rest of the 7 d, with all genes in the operon significantly

underexpressed at 7 d compared with 24 h (Figure S5).

Scanning electron microscopy (SEM) was then performed to visualise the *P. aeruginosa*

PA14 biofilm formed on the surface of the EVPL tissue at 24 h, 48 h, and 7 d, to determine

the phenotypic implications of the differential expression. Uninfected tissue was also

imaged as a negative control (Figure S6). A biofilm could not be imaged at 24 h, however

there was a dense, structured *P. aeruginosa* PA14 biofilm clear at 48 h despite significantly

reduced expression of genes for biosynthesis of the biofilm exopolysaccharide Pel (Figure 7).

This was consistent with previous work performed using the EVPL model [20]. The biofilm

population was maintained at 7 d, where the *P. aeruginosa* PA14 cells remained clearly

visible (Figure S7).

A. Lung 3: *P. aeruginosa* PA14 (48 h)

B. Lung 4: *P. aeruginosa* PA14 (48 h)

10 µm

10 µm

C. Lung 3: *P. aeruginosa* PA14 (48 h)

D. Lung 4: *P. aeruginosa* PA14 (48 h)

5 µm

5 µm

376 **Figure 7. Scanning electron micrographs of *Pseudomonas aeruginosa* PA14 biofilms associated with *ex vivo***

377 **pig lung (EVPL) bronchiolar tissue 48 h after infection.** Images were taken from biofilms formed on tissue

378 from two independent pig lungs: **(A, C)** lung 3 **(B, D)** lung 4. Scale bars are shown. Images were taken using a

379 Zeiss Gemini electron microscope with an InLens detector, at 1kV, and analysed using Omero5.6 [30]

Discussion

Extended analysis of an RNA-sequencing dataset [19] identified distinct transcriptomes of the *P. aeruginosa* PA14 biofilm population associated with EVPL bronchiolar tissue over 7 d infection. Differential expression analysis was performed comparing gene expression at 24 h, 48 h, and 7 d, and no significant changes in expression of key regulatory genes were found. There was however upregulation of a number of non-stress related KEGG pathways, and the majority of significant DEGs were overexpressed. This indicated that the biofilm was maintained for 7 d and the population was not dying. The greatest difference in expression was between 24 h and the two later time points: 48 h and 7 d. SEM showed that a dense biofilm had formed by 48 h, with *P. aeruginosa* cells at 7 d that did not exhibit visible indications of stress. These findings demonstrate that the EVPL model provides an environment for *P. aeruginosa* biofilm formation that not only phenotypically exhibits characteristics of a mature biofilm at 7 d as previously described [20], but also maintains the *P. aeruginosa* PA14 population for 7 d infection.

Sulfur metabolism was found to have the highest percentage of significant DEGs associated of all the significantly enriched KEGG pathways over time. It was significantly upregulated from 24 h to 7 d, and the associated genes followed two different expression profiles. There was with a gradual increase in expression from 24 h to 7 d, or a significant increase 24 h to 48 h that was maintained up to 7 d. More specifically, all genes in the sulfur transporter operon associated with cysteine biosynthesis: *cysTWA*, were found to be significantly overexpressed 24 h to 7 d. In *E. coli,* upregulation of this operon has been linked to sulfate starvation [31]. However, the role of these genes in *P. aeruginosa* biofilms is less

understood. Investigation of the sulfate starvation response in *P. aeruginosa* PAO1 and a

cystic fibrosis clinical isolate found genes involved in cysteine biosynthesis, including

cysTWA, were only weakly upregulated when sulfate was limited [32]. As outlined, *P.*

aeruginosa PA14 upregulation of sulfur metabolism in SCFM has actually been linked to a

shift towards anaerobic respiration [11]. There have also been links made with iron

starvation and sulfur metabolism. In fact, during iron starvation proteins involved in cysteine

biosynthesis and sulfur assimilation are reduced [33]. Therefore, upregulation of sulfur

metabolism in the EVPL biofilm may have been a result of a change of lifestyle, such as

anaerobic respiration or iron availability, rather than sulfur starvation. Thus, these findings

may further support the progression of *P. aeruginosa* biofilm infection in the EVPL model as

capturing the lifestyle changes observed in CF. However, sulfur and oxygen availability in the

EVPL biofilm should be determined in the future to confirm this hypothesis.

Among the pathways upregulated was also phenazine biosynthesis; the associated genes

were significantly overexpressed from 48 h to 7 d. More specifically, genes involved in

biosynthesis of the phenazine virulence factor pyocyanin were significantly overexpressed.

Pyocyanin acts as a signalling molecule for *P. aeruginosa* PA14 by upregulating efflux pump

genes, and genes involved in iron acquisition and redox. Additionally, when *P. aeruginosa*

PA14 biofilms have been grown in flow cells, phenazines, specifically pyocyanin, have been

shown to play a role in the formation of a biofilm with defined architecture, and phenazine-

deficient mutants were unable to form a structured biofilm [34]. Therefore, upregulation of

this pathway from 48 h to 7 d in the PA14 EVPL-associated biofilm may be linked to biofilm

maturation once it has formed at 48 h, leading to the *in vivo*-like architecture previously

observed at 7 d [20].

Pyocyanin is also believed to link the *Pseudomonas* quinolone signal (PQS) QS system with

expression of the efflux pump MexGHI-OpmD [35], which we previously found to be

significantly underexpressed in the PA14 EVPL biofilm compared with *in vitro* SCFM growth

at 48 h [19]. All three *P. aeruginosa* QS systems are involved in the regulation of phenazine

biosynthesis operons, including production of phenazine-1-carboxylic acid (PCA) and

pyocyanin [36]. The upregulation of phenazine biosynthesis thus indicates that there were

significant changes in expression of QS-regulated pathways over time in the EVPL biofilm.

Following this, and our previous work indicating that the EVPL biofilm may more accurately

recapitulate *P. aeruginosa* QS gene expression in CF sputum than other *in vitro* models [19],

P. aeruginosa PA14 changes in expression of genes in the three QS pathways was

investigated: LasI/R, RhlI/R, and PQS. There were no significant differences found in the

expression of genes in these pathways for any of the time contrasts in the EVPL biofilm,

suggesting changes in the QS systems did not occur over 7 d. Other regulatory pathways

involved in lifestyle changes, including the switch from acute to chronic infection, were also

investigated however no clear differences in expression were found either.

There were also no significant changes in virulence factor gene expression found. It is known

that as part of adaptation to the CF lung and formation of a chronic infection, mutations in

P. aeruginosa virulence pathways are selected for and virulence factor production is

significantly reduced [37]. It is hypothesised that *P. aeruginosa* may have already switched

off virulence factor production by 24 h when it was forming the established biofilm

visualised at 48 h, hence no change in gene expression was observed. This is supported by

the findings in our previous study, in which we found two of the major protease-encoding

genes *lasA* and *lasB* were significantly underexpressed in the EVPL-associated biofilm

compared with SCFM *in vitro* growth at both 24 h and 48 h [19]. This may also explain why

no significant differential expression was observed in regulatory pathways including QS.

Future work looking at the accumulation of mutations over 7 d in the EVPL-associated *P.*

aeruginosa biofilm is required to confirm this hypothesis.

There were however significant changes in expression of biofilm-associated genes found in

the EVPL PA14 biofilm population over 7 d. Pel biosynthesis was significantly downregulated

from 24 h to 48 h, then maintained at the same expression level from 48 h to 7 d, which

resulted in significant downregulation from 24 h to 7 d. This suggested that *P. aeruginosa*

PA14 production of Pel was reduced after 24 h of infection, then kept at the same level from

48 h onwards. Pel is a biofilm exopolysaccharide that is essential to biofilm development by

P. aeruginosa PA14 and is only expressed during biofilm growth [38, 39]. It must be

considered that Pel was significantly underexpressed in the EVPL biofilm at 48 h and 7 d

compared with 24 h. Thus, our results do not indicate that there was not any Pel being

produced at the later time points, just less than the initial stage of infection in the EVPL

model. This may have been caused by upregulation of Pel biosynthesis as the biofilm was

established at 24 h, which was reduced once the biofilm was formed at 48 h, and then

maintained at this lower level over the remaining 5 d. This hypothesis was supported by

SEM images of the biofilm, which showed a dense, structured *P. aeruginosa* biofilm already

present at 48 h.

Taking this into account, following formation of a biofilm, gene expression analysis indicated

that the focus may have shifted from production of the biofilm matrix to biofilm

maintenance and maturation from 48 h to 7 d. Significant upregulation of two lectin genes

was also observed in the lung-associated biofilm from 24 h to 7 d, and from 24 h to 48 h.

Both have been associated with a mature biofilm; *lecA* is indicated to play a role in the

formation of a mature biofilm architecture [40], and *lecB* has been shown to be involved in

increased cell and exopolysaccharide retention in biofilms [28]. This retention may also have

been associated with the reduction in Pel production, as less new Pel would be required if

the Pel already produced was being retained. This proposed maintenance of a mature

biofilm structure was also accompanied by increased antimicrobial tolerance.

In our first study of *P. aeruginosa* PA14 gene expression in the EVPL model compared with *in*

vitro SCFM growth, we found that the EVPL growth environments cued differential

expression of a number of AMR genes despite no antibiotic exposure, at both 24 h and 48 h

[19]. However only three genes predicted to be involved in AMR were found to be

differentially expressed over 7 d in the EVPL biofilm, suggesting that this differential

expression of resistance-associated genes cued by the environment is unlikely to be

dynamic and remains consistent throughout 7 d infection.

The significantly differentially expressed AMR genes that were found included significant

reduction in expression of two individual antibiotic resistance genes from 24 h to 48 h: *arnA*

and *bcr-1*, associated with resistance to CAMPs [27] and bicyclomycin [25] respectively.

However, the *P. aeruginosa* PA14 biofilm population grown for 7 d in the EVPL model was

able to tolerate treatment with 256 µg ml^{-1} colistin or 256 µg ml^{-1} polymyxin B at 7 d,

despite not being able to following 24 h growth. Thus, there were other mechanisms

involved by 7 d that resulted in increased *P. aeruginosa* polymyxin tolerance. It is

hypothesised that the formation of a biofilm, of which the matrix facilitates antibiotic

resistance [4], may have resulted in other mechanisms of antibiotic resistance being

downregulated as they were not required. This would provide explanation as to why

expression of AMR genes was reduced once a biofilm had formed, and the matrix was no

longer being rapidly produced. This finding would further support the use of biofilm

disrupters to improve treatment regimens, as the *P. aeruginosa* cells within the biofilm may

not be expressing genes associated with antibiotic resistance and are thus more susceptible

to treatments.

However, it must be considered that there were other mechanisms of antibiotic resistance

that were found to be significantly overexpressed in the EVPL biofilm, which are associated

with potential environmental pressures. Phosphate transporters were significantly

overexpressed from 24 h to 7 d, indicative of phosphate stress. Under phosphate stress *P.*

aeruginosa has been shown to undergo lipid modifications that confer resistance to

polymyxin B. There is a transition from phospholipids to glycolipids, which confer protection

against polymyxin B [41]. Therefore, this differential expression may also be responsible, or

involved in, the increased antibiotic tolerance observed. Future investigation of *P.*

aeruginosa lipid modifications in the EVPL model environment would provide further

insight. This analysis has however indicated that these pathways may be induced, thus

further understanding of lipid modifications cued by the EVPL model may highlight novel

treatment targets.

In summary, we have demonstrated that there are changes in the *P. aeruginosa* lung-

associated biofilm population in the EVPL model over 7 d, however the model is able to

maintain the population for this time with no obvious indicators of stress. We have

identified an increase in polymyxin tolerance that is accompanied by an upregulation in

sulfur metabolism, and maintenance of a structured biofilm, thus providing further insight

into *P. aeruginosa* antibiotic tolerance in CF. Introduction of other factors that influence *P.*

aeruginosa gene expression in the CF lung into the EVPL model, such as other microbes,

may reveal other changes in the biofilm population over time that facilitate survival in these

conditions. Further understanding of these mechanisms will provide better understanding

as to how to eradicate or prevent chronic *P. aeruginosa* infections.

Materials and Methods

Bacterial strain. The *Pseudomonas aeruginosa* PA14 wild type strain used for this work was

originally from the University of Washington [42, 43], and was grown on Luria-Bertani (LB)

agar (Melford Laboratories) for 24 h at 37 °C prior to all experiments.

Synthetic cystic fibrosis sputum medium. Synthetic cystic fibrosis sputum medium (SCFM)

was made based on the original protocol [44]. Glucose was substituted with deionised water

as previous work using the EVPL model demonstrated that in the presence of lung tissue,

the absence of glucose did not affect *P. aeruginosa* growth but did facilitate endogenous

lung bacterial species growth [45].

EVPL dissection and infection. Pig lung bronchiolar tissue was dissected and subsequently

infected following the previously published method [19, 20, 46]. Two independent lungs

were used per experiment, and three individual tissue pieces per lung. All lungs used for this

work were provided by Steve Quigley and Sons Butchers Ltd. (Cubbington, United Kingdom),

and were dissected and infected on the day of arrival from the abattoir.

EVPL antibiotic treatment. Two different antibiotics were tested: colistin and polymyxin B,

both prepared as master stocks as follows; colistin (Acros Organics) was prepared to 10 mg

ml^{-1} in dH$_2$O, and polymyxin B (Sigma-Aldrich) was prepared to 50 mg ml^{-1} in dH$_2$O. The

master stocks of each antibiotic were further diluted in SCFM, in 15 ml Falcon tubes, to the

desired concentration of 256 µg ml^{-1} for use with the EVPL model.

The EVPL grown biofilms were treated with each antibiotic as previously described [46].

Following incubation at 37 °C for either 24 h, 48 h, or 7 d, *P. aeruginosa* PA14 infected pig

lung tissue pieces were removed from the individual wells of a 24-well plate using flame

sterilised forceps. Each tissue piece was washed in 500 µl PBS in individual wells of a 48-well

plate to remove non-biofilm associated bacterial cells. The tissue pieces were then

transferred to individual wells of a 48-well plate, each containing 500 µl of either 256 µg ml^{-1}

colistin or 256 µg ml^{-1} polymyxin B. A replicate control, non-antibiotic treated set of tissue

pieces from each lung per experiment, were transferred to individual wells of a 48-well plate

each containing 500 µl PBS. All plates were covered with a UV sterilised Breathe-Easier®

sealing membrane (Diversified Biotech), and incubated stationary at 37 °C for 24 h.

EVPL biofilm recovery and determination of bacterial load. Bacteria was recovered from

the EVPL tissue pieces, and the bacterial load determined following the published method

[19, 20, 46]. Briefly, following the final incubation for both RNA extractions and antibiotic

treatments, tissue pieces were transiently washed in 500 µl PBS in individual wells of a 48-

well plate to remove non-biofilm bacterial cells. Each tissue piece was then transferred into

individual, sterile homogenisation tubes (Fisherbrand) containing 1 g of 2.38 mm metal

beads (Fisher Scientific) and 1 ml PBS. The tissue-containing bead tubes were placed in a

FastPrep-24 5G (MP Biomedicals), and bead beaten for 40 s at 4 m s^{-1} to recover bacterial

cells from the lung-associated biofilm.

The lung homogenate was then used to determine the biofilm bacterial load associated with

the EVPL tissue. An aliquot of each homogenate was serially dilute in PBS in a 96-well plate

to final volumes of 200 µl. The dilutions were then plated in triplicate on LB agar and

incubated overnight at 37 °C. The number of colonies of the two highest dilutions that were

countable were determined and used to calculate the colony forming units (CFU) lung^{-1} for

each sample.

RNA sequencing work. All RNA samples used for this work have been published as part of

another analysis, and thus the same protocol was followed for extraction, precipitation,

DNA removal, and sequencing [19]. In the initial publication, differential expression in the

EVPL biofilm compared with the lung-associated planktonic growth in surrounding SCFM

media, and *in vitro* SCFM planktonic cultures was explored but differential expression across

the time course of infection in the biofilm populations was not investigated.

RNA extraction and precipitation. Briefly, following biofilm recovery after either 24 h, 48 h,

or 7 d incubation, 1 ml of the lung homogenate was added to 2 ml DNA LoBind tubes

(Eppendorf) alongside 500 µl sterile killing buffer (20 mM Tris-HCl pH 7.5, 5 mM MgCl$_2$, 20

mM NaN$_3$). The samples were then centrifuged for 1 min at 13,000 rpm, snap frozen, then

stored at -80 °C for a minimum of 1 h. Subsequently, samples were defrosted on ice and the

supernatant was carefully discarded without disturbing the pellet. The pellet was re-

suspended in 600 µl sterile LETS buffer (0.1 M LiCl, 0.01 M Na_2EDTA, 0.01 M Tris-HCl pH 7.5,

0.2% SDS) then transferred to 2 ml lysing matrix B tubes (MP Biomedicals) and bead beat in

a FastPrep-24 5G for three cycles: 6 m s^{-1} for 40 s then 5 min incubation on ice. All samples

were then incubated until they reached room temperature, centrifuged at 13,000 rpm for

10 min, then 600 µl 125:24:1 Phenol Chloroform Isoamyl alcohol (PCI) pH 4.5 (Invitrogen)

was added to each tube. The tubes were vortexed at ~14,000 rpm for 5 min, then

centrifuged for 5 min at 15,000 rpm at 4 °C. The top layer that had formed in each sample

was transferred to individual 2 ml RNase-free tubes (Sarstedt Ltd), and 1 volume of 125:24:1

PCI pH 4.5 was also added. The above 5 min vortex, centrifugation, and transfer steps were

repeated then a volume of 24:1 Chloroform Isoamyl alcohol (Sigma-Aldrich) was added. All

samples were centrifuged a further time for 5 min at 15,000 rpm, then the top layer was

transferred to individual 2 ml RNase-free tubes. A 0.1 volume of 3 M $NaCH_3COO$ pH 5.2 and

1 volume of isopropanol was added to each tube, the tubes were inverted, and stored at -20

°C overnight.

All samples were defrosted on ice and centrifuged for 15 min at 15,000 rpm, and the

supernatant was removed. The pellets were each re-suspended in 1 volume 70% (v/v)

ethanol then centrifuged at 4 °C for 15 min at 15,000 rpm, and the supernatant was

removed. The pellets were dried for ~15 min, re-suspended in 50 µl RNase-free water,

incubated on ice for 3 h, incubated at room temperature for 30 min, then snap frozen and

stored at – 80 °C.

DNA removal. To remove DNA, each sample was defrosted and 50 µl aliquots of maximum

concentration 200 µg ml^{-1} were transferred into PCR tubes. A 9:1 ratio of 10X DNase I buffer

(Invitrogen) was added to each aliquot, followed by 2 µl DNase I (Invitrogen), then the

samples were incubated for 30 min at 37 °C. Another 2 µl DNase I was added to each

reaction and the incubation was repeated. A volume of ethanol was added, then the

samples were transferred into Zymo-Spin ICC columns, and the Direct-zol RNA MiniPrep Plus

kit (Zymo Research) was followed as outlined previously [19]. DNA removal was confirmed

using 16S PCR and gel electrophoresis.

RNA quality check and sequencing. The final RNA concentration was determined using the

Qubit RNA broad range (BR) assay kit, and quality confirmed with the Agilent 2100

bioanalyzer system and RNA 6000 Pico Kit (Agilent). Depletion of any bacterial or

mammalian rRNA and Illumina library preparation for strand-specific RNA-sequencing were

then performed by Genewiz. Sequencing was performed on an Illumina Novaseq 150 bp

paired-end run.

RNA-sequencing bioinformatic analyses. All reads were quality checked using FastQC

v0.11.8 [47] initially and following every processing step. Read trimming was performed

using Trimmomatic v0.38 [48] and a minimum read length threshold of 25 bp [18]. Residual

eukaryotic and bacterial rRNA were filtered out of the reads using SortMeRNA v2.1b [49],

then the remaining reads were mapped to the pig genome (*Sus scrofa*: NCBI,

GCF_000003025.6) using HISAT2 v2.1.0 [50]. All reads that aligned were removed using

Seqtk v1.3-r106 [51], and the remaining reads were aligned to the *P. aeruginosa* UCBPP-

PA14 genome (NCBI, GCF_000014625.1) using BWA v0.7.17-r1188 aligner with the MEM

algorithm [52]. The reads mapped to coding sequences were counted using the

"featureCounts", part of the Rsubread v2.0.1 R package [53], against the *P. aeruginosa*

UCBPP-PA14 annotation from pseudomonas.com [3]. Count data was normalised using the

rlog transformation function, part of the R package DESeq2 v1.26.0 [54]. Principal

component analysis (PCA) was then performed using the "plotPCA" function within DESeq2,

using genes (n = 5,829), and 95% confidence ellipses added using ggpubr v0.4.0 [55].

Differential expression analysis comparing the three time points: 24 h, 48 h, and 7 d, was

performed using the R package DESeq2 [54]. The threshold for significant differential

expression was an adjusted *P* value < 0.05 (Benjamini-Hochberg procedure to control the

false discovery rate) and log2 fold change ≥ |1.5|. A Venn diagram of the numbers of

differentially expressed genes was produced using VennDiagram v1.7.0 R package [56].

KEGG pathway enrichment analysis was then performed using the differentially expressed

genes using the "enrichKEGG" function from the clusterProfiler v3.14.3 R package [57] and

the organism KEGG code "pau" for *P. aeruginosa* UCBPP-PA14. A KEGG pathway was

considered significantly enriched if the adjusted *P* value < 0.05 (Benjamini-Hochberg

procedure). The antimicrobial resistance genes studied were sourced from

pseudomonas.com [3], using the Comprehensive Antibiotic Resistance Database (CARD)

predictions [58]. Genes associated with the specific pathways: quorum sensing and biofilm

formation, were sourced from the KEGG database [59-61].

Scanning electron microscopy. Following EVPL dissection, infection with *P. aeruginosa*

PA14, and incubation at 37 °C for either 24 h, 48 h, or 7 d, lung tissue samples were fixed in

0.5 ml 2.5% glutaraldehyde solution in PBS, in a 24-well plate at 4 °C overnight. Uninfected

tissue was also prepared as a negative control for *P. aeruginosa* infection.

Subsequently the 2.5% glutaraldehyde solution was removed, and the tissue samples were

rinsed with PBS three times. Each tissue piece was then transferred to individual wells of a

new 24-well plate and passed through an ethanol gradient at the following concentrations

(v/v): 20%, 50%, 70%, 90%, 100%, and a further 100%, each for 1 h. The samples were

transferred to individual wells of a further 24-well plate, and 0.5 ml hexamethyldisilazane

(Electronic grade, 99+%) (HDMS) (VWR International Ltd) was added. The plate was

incubated at room temperature for 3 h, then the HDMS solution was discarded. The tissue

pieces were transferred to a new 24-well plate and left to dry in a laminar flow cabinet for 1

h. Double-sided copper tape (Agar scientific) was added to the scanning electron microscopy

sample holders (Agar scientific), then the tissue samples were placed on top. All samples

were then sputtered with carbon by using a carbon coater (SC7620 Mini Sputter

Coater/Glow Discharge System) and immediately imaged using a Zeiss Gemini scanning

electron microscope with an InLens detector, at a voltage of 1 kV. The data was analysed

using Omero5.6 [30].

Data presentation and statistical analyses. All bar graphs were produced using DataGraph

v4.7.1 [62] on Mac OS Mojave v10.14.6. Analysis of variance (ANOVA) was performed on

antibiotic treatment CFU data and post-hoc Tukey HSD tests. All statistical analyses including

ANOVA and differential expression analysis were performed using RStudio v1.2.1335

(www.rstudio.com) on Mac OS Mojave v10.14.6.

Data availability. A full list of the significant DEGs from the three contrasts reported are

provided as a data supplement. The raw sequence data is available at the European

Nucleotide Archive (accession number: PRJEB48552).

Acknowledgements

We would like to thank all at Steve Quigley & Sons butchers for donating the pig lungs used

for this work, and Leo Eberl for the bacterial strain used. We would also like to thank Jenna

Lam for her help with the RNA extraction process and Blessing Anonye for her guidance. We

also acknowledge the help of the Media Preparation Facility in the School of Life Sciences at

the University of Warwick, with special thanks to Cerith Harries and Caroline Stewart, and

the Electron Microscopy Research Technology Platform at the University of Warwick for use

of the Zeiss Gemini scanning electron microscope. This work was supported by an MRC New

Investigator Research Grant (grant number MR/R001898/1) awarded to F.H., a PhD

studentship from the BBSRC Midlands Integrative Biosciences Training Partnership (MIBTP)

awarded to N.E.H., and the University of Warwick as part of F.A.'s MBio degree in

Biomedical Science.

Author Contributions

N.E.H. and F.H. designed research; N.E.H., F.A., and R.G.M. performed research; N.E.H. and

F.A. analyzed data; N.E.H. and F.A. wrote the paper; N.E.H., F.A., R.G.M., and F.H. edited the

paper.

Competing Interests

The authors declare no conflict of interest.

References

1.   Rossi, E., et al., *Pseudomonas aeruginosa adaptation and evolution in patients with*

cystic fibrosis. Nature Reviews Microbiology, 2021. **19**(5): p. 331-342.

2.   Wardell, S.J.T., et al., *Genome evolution drives transcriptomic and phenotypic*

adaptation in Pseudomonas aeruginosa during 20 years of infection. Microbial

Genomics, 2021. **7**(11).

3.   Winsor, G.L., et al., *Enhanced annotations and features for comparing thousands of*

Pseudomonas genomes in the Pseudomonas genome database. Nucleic Acids

Research, 2016. **44**(D1): p. D646-D653.

4.   Høiby, N., et al., *Antibiotic resistance of bacterial biofilms.* International Journal of

Antimicrobial Agents, 2010. **35**(4): p. 322-332.

5.   Schobert, M. and D. Jahn, *Anaerobic physiology of Pseudomonas aeruginosa in the*

cystic fibrosis lung. International Journal of Medical Microbiology, 2010. **300**(8): p.

549-556.

6.   Kolpen, M., et al., *Polymorphonuclear leucocytes consume oxygen in sputum from*

chronic Pseudomonas aeruginosa pneumonia in cystic fibrosis. Thorax, 2010. **65**(1): p.

57-62.

7.   Filkins, L.M. and G.A. O'Toole, *Cystic Fibrosis Lung Infections: Polymicrobial, Complex,*

and Hard to Treat. PLOS Pathogens, 2015. **11**(12): p. e1005258.

8.   Cowley, E.S., et al., *Pediatric Cystic Fibrosis Sputum Can Be Chemically Dynamic,*

Anoxic, and Extremely Reduced Due to Hydrogen Sulfide Formation. mBio, 2015. **6**(4):

p. e00767-15.

9.   Worlitzsch, D., et al., *Effects of reduced mucus oxygen concentration in airway*

Pseudomonas infections of cystic fibrosis patients. Journal of Clinical Investigation,

2002. **109**(3): p. 317-325.

10. Alvarez-Ortega, C. and C.S. Harwood, *Responses of Pseudomonas aeruginosa to low*

oxygen indicate that growth in the cystic fibrosis lung is by aerobic respiration. Mol

Microbiol, 2007. **65**(1): p. 153-65.

11. Tata, M., et al., *RNASeq Based Transcriptional Profiling of Pseudomonas aeruginosa*

PA14 after Short- and Long-Term Anoxic Cultivation in Synthetic Cystic Fibrosis

Sputum Medium. PLOS ONE, 2016. **11**(1): p. e0147811.

12. Cheng, P.W., et al., *Increased sulfation of glycoconjugates by cultured nasal epithelial*

cells from patients with cystic fibrosis. Journal of Clinical Investigation, 1989. **84**(1): p.

68-72.

13. Xia, B., et al., *Altered O-glycosylation and sulfation of airway mucins associated with*

cystic fibrosis. Glycobiology, 2005. **15**(8): p. 747-775.

14. Lamblin, G., et al., *Human airway mucin glycosylation: A combinatory of*

carbohydrate determinants which vary in cystic fibrosis. Glycoconjugate Journal,

2001. **18**(9): p. 661-684.

15.   Robinson, C.V., et al., *Desulfurization of mucin by Pseudomonas aeruginosa:*

760   *influence of sulfate in the lungs of cystic fibrosis patients.* Journal of Medical

761   Microbiology, 2012. **61**(12): p. 1644-1653.

762   16.   Hogardt, M. and J. Heesemann, *Microevolution of Pseudomonas aeruginosa to a*

763   *Chronic Pathogen of the Cystic Fibrosis Lung*, in *Between Pathogenicity and*

764   *Commensalism*. 2011, Springer Berlin Heidelberg. p. 91-118.

765   17.   Jiricny, N., et al., *Loss of Social Behaviours in Populations of Pseudomonas aeruginosa*

766   *Infecting Lungs of Patients with Cystic Fibrosis.* PLoS ONE, 2014. **9**(1): p. e83124.

767   18.   Cornforth, D.M., et al., *Pseudomonas aeruginosa transcriptome during human*

768   *infection.* PNAS, 2018. **115**(22): p. E5125-E5134.

769   19.   Harrington, N.E., J.L. Littler, and F. Harrison, *Transcriptome Analysis of Pseudomonas*

770   *aeruginosa Biofilm Infection in an Ex Vivo Pig Model of the Cystic Fibrosis Lung.*

771   Applied and Environmental Microbiology, 2022. **88**(3).

772   20.   Harrington, N.E., E. Sweeney, and F. Harrison, *Building a better biofilm - Formation of*

773   *in vivo-like biofilm structures by Pseudomonas aeruginosa in a porcine model of*

774   *cystic fibrosis lung infection.* Biofilm, 2020. **2**.

775   21.   Rossi, E., et al., *High-resolution in situ transcriptomics of Pseudomonas aeruginosa*

776   *unveils genotype independent patho-phenotypes in cystic fibrosis lungs.* Nature

777   Communications, 2018. **9**(1).

778   22.   Schreiber, K., et al., *Anaerobic Survival of Pseudomonas aeruginosa by Pyruvate*

779   *Fermentation Requires an Usp-Type Stress Protein.* Journal of Bacteriology, 2006.

780   **188**(2): p. 659-668.

23. Eschbach, M., et al., *Long-Term Anaerobic Survival of the Opportunistic Pathogen*

Pseudomons aeruginosa via Pyruvate Fermentation. Journal of Bacteriology, 2004.

186(14).

24. Mercenier, A., et al., *Regulation of enzyme synthesis in the arginine deiminase*

pathway of Pseudomonas aeruginosa. Journal of Bacteriology, 1980. **144**(1).

25. Fonseca, E.L., et al., *Full characterization of the integrative and conjugative element*

carrying the metallo-β-lactamase blaSPM-1 and bicyclomycin bcr1 resistance genes

found in the pandemic Pseudomonas aeruginosa clone SP/ST277. Journal of

Antimicrobial Chemotherapy, 2015. **70**(9): p. 2547-2550.

26. Roy, S., et al., *Transcriptional expression of PmrB and arnA within polymyxin-*

resistant nosocomial isolates of Pseudomonas aeruginosa from India. Journal of

Microbiology and Infectious Diseases, 2018. **8**(3): p. 61-68.

27. Gutu, A.D., et al., *Pseudomonas aeruginosa High-Level Resistance to Polymyxins and*

Other Antimicrobial Peptides Requires cprA, a Gene That Is Disrupted in the PAO1

Strain. Antimicrobial Agents and Chemotherapy, 2015. **59**(9): p. 5377-5387.

28. Passos Da Silva, D., et al., *The Pseudomonas aeruginosa lectin LecB binds to the*

exopolysaccharide Psl and stabilizes the biofilm matrix. Nature Communications,

2019. **10**(1).

29. Bouillet, S., et al., *Connected partner-switches control the life style of Pseudomonas*

aeruginosa through RpoS regulation. Scientific Reports, 2019. **9**(1).

30. Goldberg, I.G., et al., *The Open Microscopy Environment (OME) Data Model and XML*

file: open tools for informatics and quantitative analysis in biological imaging.

Genome Biology, 2005. **6**(5): p. R47.

31.   Gyaneshwar, P., et al., *Sulfur and Nitrogen Limitation in Escherichia coli K-12: Specific*

805   *Homeostatic Responses.* Journal of Bacteriology, 2005. **187**(3): p. 1074-1090.

806   32.   Tralau, T., et al., *Transcriptomic Analysis of the Sulfate Starvation Response of*

807   *Pseudomonas aeruginosa.* Journal of Bacteriology, 2007. **189**(19): p. 6743-6750.

808   33.   Nelson, C.E., et al., *Proteomic Analysis of the Pseudomonas aeruginosa Iron*

809   *Starvation Response Reveals PrrF Small Regulatory RNA-Dependent Iron Regulation*

810   *of Twitching Motility, Amino Acid Metabolism, and Zinc Homeostasis Proteins.*

811   Journal of Bacteriology, 2019. **201**(12).

812   34.   Ramos, I., et al., *Phenazines affect biofilm formation by Pseudomonas aeruginosa in*

813   *similar ways at various scales.* Research in Microbiology, 2010. **161**(3): p. 187-191.

814   35.   Dietrich, L.E.P., et al., *The phenazine pyocyanin is a terminal signalling factor in the*

815   *quorum sensing network of Pseudomonas aeruginosa.* Molecular Microbiology, 2006.

816   **61**(5): p. 1308-1321.

817   36.   Higgins, S., et al., *Differential Regulation of the Phenazine Biosynthetic Operons by*

818   *Quorum Sensing in Pseudomonas aeruginosa PAO1-N.* Frontiers in Cellular and

819   Infection Microbiology, 2018. **8**.

820   37.   Smith, E.E., et al., *Genetic adaptation by Pseudomonas aeruginosa to the airways of*

821   *cystic fibrosis patients.* Proceedings of the National Academy of Sciences, 2006.

822   **103**(22): p. 8487-8492.

823   38.   Colvin, K.M., et al., *The Pel Polysaccharide Can Serve a Structural and Protective Role*

824   *in the Biofilm Matrix of Pseudomonas aeruginosa.* PLoS Pathogens, 2011. **7**(1): p.

825   e1001264.

826   39.   Marmont, L.S., et al., *PelA and PelB proteins form a modification and secretion*

827   *complex essential for Pel polysaccharide-dependent biofilm formation in*

Pseudomonas aeruginosa. Journal of Biological Chemistry, 2017. **292**(47): p. 19411-

19422.

40.   Diggle, S.P., et al., *The galactophilic lectin, LecA, contributes to biofilm development*

in Pseudomonas aeruginosa. Environmental Microbiology, 2006. **8**(6): p. 1095-1104.

41.   Jones, R.A., et al., *Phosphorus stress induces the synthesis of novel glycolipids in*

Pseudomonas aeruginosa that confer protection against a last-resort antibiotic. The

ISME Journal, 2021. **15**(11): p. 3303-3314.

42.   Liberati, N.T., et al., *An ordered, nonredundant library of Pseudomonas aeruginosa*

strain PA14 transposon insertion mutants. Proceedings of the National Academy of

Sciences, 2006. **103**(8): p. 2833-2838.

43.   Liberati, N.T., et al. *PA14 transposon insertion mutant library*. 2006 [cited 2022

10/05/2022]; Available from: http://pa14.mgh.harvard.edu/cgi-bin/pa14/home.cgi.

44.   Palmer, K.L., L.M. Aye, and M. Whiteley, *Nutritional Cues Control Pseudomonas*

aeruginosa Multicellular Behavior in Cystic Fibrosis Sputum. Journal of Bacteriology,

2007. **189**(22): p. 8079-8087.

45.   Harrison, F. and S.P. Diggle, *An ex vivo lung model to study bronchioles infected with*

Pseudomonas aeruginosa biofilms. Microbiology, 2016. **162**(10): p. 1755-1760.

46.   Harrington, N.E., et al., *Antibiotic Efficacy Testing in an Ex vivo Model of*

Pseudomonas aeruginosa and Staphylococcus aureus Biofilms in the Cystic Fibrosis

Lung. Journal of Visualized Experiments, 2021(167).

47.   Andrews, S. *FastQC: a quality control tool for high throughput sequence data.* 2018;

Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

48.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina*

sequence data. Bioinformatics, 2014. **30**(15): p. 2114-2120.

49.   Kopylova, E., L. Noé, and H. Touzet, *SortMeRNA: fast and accurate filtering of*

853   *ribosomal RNAs in metatranscriptomic data.* Bioinformatics, 2012. **28**(24): p. 3211-

854   3217.

855   50.   Kim, D., et al., *Graph-based genome alignment and genotyping with HISAT2 and*

856   *HISAT-genotype.* Nature Biotechnology, 2019. **37**(8): p. 907-915.

857   51.   Li, H. *A toolkit for processing sequences in FASTA/Q formats.* 2015; Available from:

858   https://github.com/lh3/seqtk.

859   52.   Li, H., *Aligning sequence reads, clone sequences and assembly contigs with BWA-*

860   *MEM.* ArXiv, 2013. **00**.

861   53.   Liao, Y., G.K. Smyth, and W. Shi, *The R package Rsubread is easier, faster, cheaper*

862   *and better for alignment and quantification of RNA sequencing reads.* Nucleic Acids

863   Research, 2019. **47**(8): p. e47-e47.

864   54.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*

865   *dispersion for RNA-seq data with DESeq2.* Genome Biology, 2014. **15**(12).

866   55.   Kassambara, A. *ggpubr: "ggplot2" based publication ready plots. R package version*

867   *0.4.0.* 2020; Available from: https://cran.r-project.org/package=ggpubr.

868   56.   Chen, H., *VennDiagram: Generate High-Resolution Venn and Euler Plots. R package*

869   *version 1.7.0.* 2021.

870   57.   Yu, G., et al., *ClusterProfiler: an R package for comparing biological themes among*

871   *gene clusters.* OMICS, 2012. **16**: p. 284-287.

872   58.   Alcock, B.P., et al., *CARD 2020: antibiotic resistome surveillance with the*

873   *comprehensive antibiotic resistance database.* Nucleic Acids Research, 2019.

874   59.   Kanehisa, M. and S. Goto, *KEGG: Kyoto Encyclopedia of Genes and Genomes.* Nucleic

875   Acids Research, 2000. **28**(1): p. 27-30.

876   60.   Kanehisa, M., *Toward understanding the origin and evolution of cellular organisms.*

877   Protein Science, 2019. **28**(11): p. 1947-1951.

878   61.   Kanehisa, M., et al., *KEGG: integrating viruses and cellular organisms.* Nucleic Acids

879   Research, 2021. **49**(D1): p. D545-D551.

880   62.   Adalsteinsson, D. and P. Schultz, *The DataGraph Manual.* 2020.