How long does it take to cue a *Pseudomonas aeruginosa* chronic infection phenotype in the lab? Insights from transcriptome analysis in a cystic fibrosis lung model.

Niamh E. Harrington[1,2*], Freya Allen[1,3], Ramón Garcia Maset[4,5], Freya Harrison[1*]

1: School of Life Sciences, Gibbet Hill Campus, The University of Warwick, Coventry, CV4 7AL, UK.

2: Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, L69 7ZB, UK.

3: (Current address) Institute of Microbiology and Infection and Department of Microbes, Infection and Microbiomes, School of Infection, Inflammation and Immunology, College of Medicine and Health, University of Birmingham, Birmingham, B15 2TT, UK.

4: Warwick Medical School, Gibbet Hill Campus, The University of Warwick, Coventry, CV4 7AL, UK.

5: (Current address) Department of Engineering Science, University of Oxford, Parks Road, Oxford, OX1 3PJ, UK

* Co-corresponding authors: Niamh.Harrington@liverpool.ac.uk, F.Harrison@warwick.ac.uk

Keywords: Biofilm, chronic infection, cystic fibrosis, *ex vivo* model, *Pseudomonas*

22   *aeruginosa,* RNA sequencing, transcriptome, quorum sensing, antimicrobial resistance,

23   sulfur metabolism

Abstract

The opportunistic pathogen *Pseudomonas aeruginosa* forms biofilm infections in the lungs of people with the genetic condition cystic fibrosis (CF) that can persist for decades. There are numerous *P. aeruginosa* lifestyle changes associated with chronic biofilm infection cued by the CF lung environment. These include a loss of virulence, metabolic changes and increased antimicrobial tolerance. We have investigated *P. aeruginosa* PA14 biofilm infection over 7 d in an *ex vivo* pig lung (EVPL) model for CF, previously shown to facilitate formation of a clinically relevant *P. aeruginosa* biofilm structure with expression of key genes comparable to human infection. We have compared *P. aeruginosa* gene expression between sequential time points: 24 h, 48 h and 7 d post infection, and investigated tolerance to polymyxins. Our results demonstrate that the EVPL model can maintain a *P. aeruginosa* biofilm population, which exhibits increased antibiotic tolerance, for at least 7 d. Differential expression of antimicrobial resistance-associated genes was not observed, however there was significant upregulation of sulfur metabolism and maintenance of a structured biofilm. Our findings provide further insight into the increased *P. aeruginosa* antibiotic tolerance during chronic infection of the CF lung, and suggest we can cue aspects of chronic infection in 7 d under the right lab conditions.

Introduction

Pseudomonas aeruginosa chronic biofilm infections of the cystic fibrosis (CF) lung are, in part, characterised by their physiological adaptations to the environment. This is driven by

environmental pressures such as osmotic, nitrosative and oxidative stress, as well as

interactions with the commensal microbiome and co-infecting pathogens, and antibiotic

exposure [1]. Lifestyle changes have been shown to improve *P. aeruginosa* persistence in

the lung, including loss of swimming motility, impaired pyoverdine production, increased

adherence and formation of a biofilm matrix that protects the bacterial cells from not only

the host immune response, but also antibiotics [2, 3]. Quorum sensing (QS) largely controls

the phenotypic changes observed in the biofilm population, and is estimated to influence

expression of approximately 10% of all *P. aeruginosa* genes [4]. Initial physiological changes

that induce a chronic, antibiotic-tolerant biofilm phenotype are followed by evolutionary

changes in the population over years of infection. Late stage infections exhibit reduced

phenotypic diversity and loss of virulence compared with early stage infections [5]. The

reduction in virulence factor production is linked with a decline in expression of QS-related

genes [6]. Once established, chronic *P. aeruginosa* infections are almost impossible to

eradicate due to high levels of antibiotic resistance and tolerance, contributed to by the

protective biofilm matrix.

Studies have shown that current *in vitro* models do not accurately represent critical aspects

of *P. aeruginosa* chronic infection; in particular, QS-associated gene expression in CF sputum

is not well captured [7]. However, our previous work has indicated that an *ex vivo* pig lung

(EVPL) model may more accurately capture *in vivo* QS regulated gene expression over a

short infection time (48 h) [8]. Our study focused on *P. aeruginosa* gene expression at 24 h

and 48 h post infection (PI) in the EVPL model compared with *in vitro* synthetic cystic fibrosis

sputum media (SCFM) growth. We found that the EVPL bronchiolar tissue-associated biofilm

formed at 48 h exhibited key characteristics of *in vivo* chronic infection not observed *in vitro*

[8].

We now aim to build upon these findings to determine whether the EVPL model is viable for

an extended infection time of 7 days, as a longer infection period may more accurately

replicate chronic CF infections. We previously showed *P. aeruginosa* biofilms to have a

clinically realistic, mature structure in the model at 7 days post infection [9]. We have now

performed differential expression analysis and Kyoto Encyclopedia of Genes and Genomes

(KEGG) pathway enrichment analysis. This confirms that the biofilm population was

maintained at 7 d with limited stress and allows us to better understand expression changes

in established biofilms and further knowledge of *P. aeruginosa* gene expression in later

stage infections. We identified changes in several genes and pathways, including

upregulation of sulfur metabolism and biofilm maintenance, that accompanied increased

antibiotic tolerance over time. Our findings provide new insight into *P. aeruginosa* gene

expression changes in an established biofilm infection and their potential implications for

treatment success, as well as highlighting a viable extended time point for use of the EVPL

model to study chronic *P. aeruginosa* biofilm infection in CF.

Methods

Bacterial strain. The *Pseudomonas aeruginosa* PA14 wild type strain used for this work was

originally from the University of Washington [10, 11], and was grown on Lysogeny Broth (LB)

agar (Melford Laboratories) for 24 h at 37 °C prior to all experiments.

95 **Synthetic cystic fibrosis sputum medium.** SCFM was made based on the original recipe by

96 Palmer *et al.*, now commonly referred to as "SCFM1" [12]. Glucose was removed from the

97 recipe as previous work using the EVPL model demonstrated that in the presence of lung

98 tissue, the absence of glucose did not affect *P. aeruginosa* growth but did facilitate

99 endogenous lung bacterial species growth [13].

101 **EVPL dissection and infection.** Pig lung bronchiolar tissue was dissected and subsequently

102 infected following the previously published method [8, 9, 14]. Two independent lungs were

103 used per experiment, with three individual tissue pieces used per lung. All lungs used for

104 this work were provided by Steve Quigley and Sons Butchers Ltd. (Cubbington, United

105 Kingdom) and were dissected and infected on the day of arrival from the abattoir.

107 **EVPL antibiotic treatment.** Two different antibiotics were tested: colistin and polymyxin B,

108 both prepared as master stocks as follows: colistin (Acros Organics) was prepared to 10 mg

109 ml^{-1} in dH$_2$O, and polymyxin B (Sigma-Aldrich) was prepared to 50 mg ml^{-1} in dH$_2$O. The

110 master stocks of each antibiotic were further diluted in SCFM, in 15 ml Falcon tubes, to the

111 desired concentration of 256 µg ml^{-1} for use with the EVPL model.

113 The EVPL grown biofilms were treated with each antibiotic as previously described [14].

114 Following incubation at 37 °C for either 24 h, 48 h, or 7 d, *P. aeruginosa* PA14 infected pig

115 lung tissue pieces were removed from the individual wells of a 24-well plate using flame

116 sterilised forceps. Each tissue piece was washed in 500 µl PBS in individual wells of a 48-well

117 plate to remove non-biofilm associated bacterial cells. The tissue pieces were then

118 transferred to individual wells of a 48-well plate, each containing 500 µl of either 256 µg ml^{-1}

colistin or 256 µg ml^{-1} polymyxin B. A replicate control, non-antibiotic treated set of tissue

pieces from each lung per experiment, were transferred to individual wells of a 48-well plate

each containing 500 µl PBS. All plates were covered with a UV sterilised Breathe-Easier®

sealing membrane (Diversified Biotech), and incubated stationary at 37 °C for 24 h.

EVPL biofilm recovery and determination of bacterial load. Bacteria were recovered from

the EVPL tissue pieces, and the bacterial load determined following the published method

[8, 9, 14]. Briefly, following the final incubation for both RNA extractions and antibiotic

treatments, tissue pieces were transiently washed in 500 µl PBS in individual wells of a 48-

well plate to remove non-biofilm bacterial cells. Each tissue piece was then transferred into

individual, sterile homogenisation tubes (Fisherbrand) containing 1 g of 2.38 mm metal

beads (Fisher Scientific) and 1 ml PBS. The tissue-containing bead tubes were placed in a

FastPrep-24 5G (MP Biomedicals), and bead beaten for 40 s at 4 m s^{-1} to recover bacterial

cells from the lung-associated biofilm.

The lung homogenate was then used to determine the biofilm bacterial load associated with

the EVPL tissue. An aliquot of each homogenate was serially dilute in PBS in a 96-well plate

to final volumes of 200 µl. The dilutions were then plated in triplicate on LB agar and

incubated overnight at 37 °C. The number of colonies of the two highest dilutions that were

countable were determined and used to calculate the colony forming units (CFU) lung^{-1} for

each sample.

RNA extraction and sequencing, and relationship to previously published work. All RNA

samples used for this work were collected as part of a bigger study, and thus the protocol

for extraction, precipitation, DNA removal and sequencing are outlined in the original

publication [8].

RNA-sequencing bioinformatic analyses. All reads were quality checked using FastQC

v0.11.8 [15] initially and following every processing step. Read trimming was performed

using Trimmomatic v0.38 [16] and a minimum read length threshold of 25 bp [7]. Residual

eukaryotic and bacterial rRNA were filtered out of the reads using SortMeRNA v2.1b [17],

then the remaining reads were mapped to the pig genome (*Sus scrofa*: NCBI,

GCF_000003025.6) using HISAT2 v2.1.0 [18]. All reads that aligned were removed using

Seqtk v1.3-r106 [19], and the remaining reads were aligned to the *P. aeruginosa* UCBPP-

PA14 genome (NCBI, GCF_000014625.1) using BWA v0.7.17-r1188 aligner with the MEM

algorithm [20]. The reads mapped to coding sequences were counted using the

"featureCounts", part of the Rsubread v2.0.1 R package [21], against the *P. aeruginosa*

UCBPP-PA14 annotation from pseudomonas.com [1]. Count data was normalised using the

rlog transformation function, part of the R package DESeq2 v1.26.0 [22]. Principal

component analysis (PCA) was then performed using the "plotPCA" function within DESeq2,

using genes (n = 5,829), and 95% confidence ellipses added using ggpubr v0.4.0 [23].

Differential expression analysis comparing the three time points: 24 h, 48 h, and 7 d, was

performed using the R package DESeq2 [22]. The threshold for significant differential

expression was an adjusted *P* value < 0.05 (Benjamini-Hochberg procedure to control the

false discovery rate) and log2 fold change ≥ |1.5|. A Venn diagram of the numbers of

differentially expressed genes was produced using VennDiagram v1.7.0 R package [24].

KEGG pathway enrichment analysis was then performed using the differentially expressed

genes using the "enrichKEGG" function from the clusterProfiler v3.14.3 R package [25] and

the organism KEGG code "pau" for *P. aeruginosa* UCBPP-PA14. A KEGG pathway was

considered significantly enriched if the adjusted *P* value < 0.05 (Benjamini-Hochberg

procedure). The antimicrobial resistance genes studied were sourced from

pseudomonas.com [1], using the Comprehensive Antibiotic Resistance Database (CARD)

predictions [26]. Genes associated with the specific pathways: quorum sensing and biofilm

formation, were sourced from the KEGG database [27-29].

Scanning electron microscopy. Following EVPL dissection, infection with *P. aeruginosa*

PA14, and incubation at 37 °C for either 24 h, 48 h, or 7 d, lung tissue samples were fixed in

0.5 ml 2.5% glutaraldehyde solution in PBS, in a 24-well plate at 4 °C overnight. Uninfected

tissue was also prepared as a negative control for *P. aeruginosa* infection.

Subsequently the 2.5% glutaraldehyde solution was removed, and the tissue samples were

rinsed with PBS three times. Each tissue piece was then transferred to individual wells of a

new 24-well plate and passed through an ethanol gradient at the following concentrations

(v/v): 20%, 50%, 70%, 90%, 100%, and a further 100%, each for 1 h. The samples were

transferred to individual wells of a further 24-well plate, and 0.5 ml hexamethyldisilazane

(Electronic grade, 99+%) (HDMS) (VWR International Ltd) was added. The plate was

incubated at room temperature for 3 h, then the HDMS solution was discarded. The tissue

pieces were transferred to a new 24-well plate and left to dry in a laminar flow cabinet for 1

h. Double-sided copper tape (Agar scientific) was added to the scanning electron microscopy

sample holders (Agar scientific), then the tissue samples were placed on top. All samples

were then sputtered with carbon by using a carbon coater (SC7620 Mini Sputter

Coater/Glow Discharge System) and immediately imaged using a Zeiss Gemini scanning

electron microscope with an InLens detector, at a voltage of 1⬚kV. The data was analysed

using Omero5.6 [30].

Data presentation and statistical analyses. All bar graphs were produced using DataGraph

v4.7.1 [31] on Mac OS Mojave v10.14.6. Analysis of variance (ANOVA) was performed on

antibiotic treatment CFU data and post-hoc Tukey HSD tests. All statistical analyses including

ANOVA and differential expression analysis were performed using RStudio v1.2.1335

(www.rstudio.com) on Mac OS Mojave v10.14.6.

Data availability. A full list of the significant DEGs from the three contrasts reported are

provided as a data supplement. The raw sequence data is available at the European

Nucleotide Archive (accession number: PRJEB48552).

Results

As reported in our previous study [8], we grew *P. aeruginosa* PA14 as a biofilm associated

with the surface of EVPL bronchiolar tissue from two independent pig lungs for 24 h, 48 h

and 7 d, and performed RNA sequencing from the recovered tissue-associated biofilm

population. Our previous work confirmed that PA14 was still metabolically active 7 d PI,

using an ATP-based viability assay (ref. [8] Figure S3), and the bacterial load was maintained;

recovered CFU lung^{-1} 7 d PI were comparable to 24 h and 48 h PI (ref. [8] Figure S4). In the

present study, we have extended our previous work to investigate *P. aeruginosa* PA14 gene

expression changes in the EVPL tissue-associated biofilm 24h – 7d PI (P < 0.05, $\log_2$ fold

change ($|\log_2 FC|$) ≥ 1.5). For all comparisons the earlier time point was used as the baseline.

Expression of *P. aeruginosa* genes significantly changed over 7 d in the EVPL biofilm.

PCA of PA14 gene expression identified distinct transcriptomes at 24h, 48h and 7d PI (Figure

S1). There was no clear difference between samples from different lungs, indicating

variation between time points was much larger than between-lung variation. *P. aeruginosa*

grown for 24 h was the most distinct from other time points, supported by hierarchical

clustering and Pearson's correlation coefficient analyses (Figure S1; Figure S2). Differential

expression analysis similarly showed the greatest difference at 24 h vs 48h or 7d PI, with the

smallest number of significantly differentially expressed genes (DEGs) arising between 48 h

and 7 d (Figure 1).

Overall, more genes were found to be significantly overexpressed than underexpressed

between sequential time points (Figure 1A), suggesting gene expression increased over time

in the EVPL biofilm. While some genes showed changes in expression over all three time

points, there were also unique DEGs identified between each time point comparison (Figure

1B). This indicated there were different dynamics in gene expression changes across the

three time points.

Figure 1. The number of _Pseudomonas aeruginosa_ PA14 significant differentially expressed genes (DEGs) (_P_ <

0.05, |log$_2$ fold change| ≥ 1.5) in each time contrast in the _ex vivo_ pig lung (EVPL) model biofilm. RNA was

sequenced from three tissue-associated biofilms from each of two independent pig lungs, per time point. **(A)**

The number of significantly overexpressed genes (pink, left bar) and significantly underexpressed genes

(yellow, right bar) in each time contrast. Significant DEGs were determined at 48 h compared with 24 h

(showing change from 24 h to 48 h), at 7 d compared with 48 h (48 h to 7 d), and at 7 d compared with 24 h

(24 h to 7 d). **(B)** A Venn diagram of the number of significant DEGs. Shared genes were either significantly

overexpressed or underexpressed in all contrasts compared. Genes that were significant in multiple contrasts

but in opposing directions were not considered shared.

Changes in *P. aeruginosa* gene expression followed seven different profiles with

functional implications.

To further investigate the gene expression changes over time in the EVPL-associated biofilm,

expression profiles were determined based on shared, unique and non-significant changes

between time point comparisons. We classified the DEGs into seven profiles based on the

direction and magnitude of their change in expression 24 - 48h and 48h - 7d, outlined in

Table 1.

Table 1. *Pseudomonas aeruginosa* PA14 gene expression profiles identified in the *ex vivo* pig lung-associated

biofilm at 24 h, 48 h and 7 d. The graphical representations demonstrate the changes in expression of genes

over the three time points. Below each graph is the number of significant differentially expressed genes found

to follow that expression profile.

Profile number	Gene expression profile	Graphical representation
1	Significant change in expression 24 h to 48 h, then a significant change in the **same** direction 48 h to 7 d. Results in significant change 24 h to 7 d.	3 genes — 1 gene
2	Significant change in expression 24 h to 48 h, which is **maintained** 48 h to 7 d (no significant change). Results in significant change 24 h to 7 d.	80 genes — 68 genes
3	No significant change in expression 24 h to 48 h, then a significant change 48 h to 7 d. Results in significant change 24 h to 7 d.	66 genes — 20 genes
4	Significant change in expression 24 h to 48 h, then a small but not significant change in the **opposite** direction 48 h to 7 d. Results in no significant change 24 h to 7 d.	157 genes — 79 genes
5	Small but not significant change in expression 24 h to 48 h, then a small but not significant change in the **same** direction 48 h to 7 d. Results in overall significant change 24 h to 7 d.	118 genes — 79 genes
6	Small but not significant change in expression 24 h to 48 h, then a significant change in the **opposite** direction 48 h to 7 d. Results in no significant change 24 h to 7 d.	20 genes — 51 genes
7	Significant change in expression 24 h to 48 h, then a significant change in the **opposite** direction 48 h to 7 d. Results in no significant change 24 h to 7 d.	29 genes — 15 genes

Four genes were found to be significantly differentially expressed in the same direction 24 h

- 48 h, and then 48 h - 7 d (Figure 1B), classified as expression profile 1 (Table 1), however

there was no functional association between these genes (Table S2). A second profile where

gene expression changed in the same direction over 7 d but at a slower rate was identified,

resulting in a significant change 24 h - 7 d (profile 5; Table 1). KEGG enrichment analysis

found four pathways significantly enriched ($P < 0.05$) for this profile, including degradation

of aromatic compounds and benzoate degradation (Figure 2B). All significant DEGs in the

degradation of aromatic compounds pathway were linked to benzoate degradation.

Arginine biosynthesis was also significantly enriched (Figure 2B), and two of the associated

significant DEGs that reduced in expression over time were arginine fermentation enzymes

[32]: *arcB* (-2.25 $\log_2$ fold change (FC)) and *arcC* (-2.16 $\log_2$FC), which have been shown to be

induced under stress [33, 34]. Additionally, sulfur metabolism was significantly enriched for

this profile as well as expression profile 2 (Figure 2), where there was a significant increase

in expression 24 h - 48 h that was maintained for 7 d (Table 1).

Figure 2. Significantly enriched (*P* < 0.05) *Pseudomonas aeruginosa* PA14 Kyoto Encyclopedia of Genes and

Genomes (KEGG) pathways found for gene expression profiles in the *ex vivo* pig lung-associated biofilm over

24h, 48 h and 7 d. The expression of significantly differentially expressed genes over the three time points was

investigated and separated into seven different profiles, three of which were found to involve sufficient genes

associated with KEGG pathways for pathways to be significantly enriched. The bars show the percentage of

significant differentially expressed genes associated with each enriched pathway (*P* < 0.05, |$\log_2$ fold change| ≥

1.5); those overexpressed (left bar) and underexpressed (right bar) are shown by different colours (see key).

The graphical representation of the gene expression profile is shown in the top right of each graph; the left

graph shows the profile of overexpressed genes, and the right graph shows the profile of underexpressed

genes. **(A)** Significantly enriched KEGG pathway following expression profile 2. **(B)** Significantly enriched KEGG

pathways following profile 5. **(C)** Significantly enriched KEGG pathways following profile 6.

Gene expression profile 6 was the only other group where there were significantly enriched

KEGG pathways ($P < 0.05$), involving a significant change in expression 48 h - 7 d (Table 1;

Figure 2C). This included phenazine biosynthesis and QS (Figure 2C); all phenazine

biosynthesis significant DEGs were also associated with the QS pathway.

Sulfur metabolism was upregulated over 7 d in the EVPL-associated biofilm.

As expression changes 24 h - 7 d appeared to have the greatest functional implications,

KEGG enrichment analysis was performed on all significant DEGs that arose in this

comparison. There were no *P. aeruginosa* PA14 pathways significantly enriched 24 h - 48 h,

and those found between 48 h and 7 d were comparable to expression profile 6 (Figure 2C;

Figure S3). Four pathways were significantly enriched 24 h - 7 d in the EVPL-associated

biofilm ($P < 0.05$): sulfur metabolism, ATP-binding cassette (ABC) transporters, degradation

of aromatic compounds and nitrogen metabolism (Figure 3).

Figure 3. Significantly enriched (*P* < 0.05) *Pseudomonas aeruginosa* PA14 Kyoto encyclopedia of genes and

genomes (KEGG) pathways in the *ex vivo* pig lung-associated biofilm from 24 h to 7 d. Each bar shows the

percentage of genes associated with each pathway that were significantly differentially expressed (*P* < 0.05,

|log$_2$ fold change| ≥ 1.5). The overexpressed genes (left bar) and underexpressed genes (right bar) are show by

different colour bars (see key).

The greatest percentage of associated significant DEGs was for sulfur metabolism (Figure 3):

47.06% of sulfur metabolism genes were significantly overexpressed in the EVPL biofilm 24 h

- 7 d, suggesting this pathway was upregulated. This included sulfate transport genes *cysA*,

cysW and *cysT* (3.04, 2.87, and 3.14 log$_2$FC respectively). There was only one associated

gene significantly reduced in expression at 7 d: the sulfite/nitrate reductase PA14_10550 (-

1.56 log$_2$FC). The majority of the significantly differentially expressed ABC transporter genes

were involved in sulfur metabolism. However, three phosphate-specific transporter genes

were also significantly overexpressed: *pstB* (1.52 log$_2$FC), *pstA* (1.83 log$_2$FC), and *pstC* (1.88

log$_2$FC).

Nitrogen metabolism was the only significantly enriched pathway where more significant

DEGs decreased in expression 24 h - 7 d than increased (Figure 3). Expression of genes in the

P. aeruginosa nitrogen metabolism operons responsible for a shift towards anaerobic

respiration in CF sputum (*nor*, *nir*, *nap*, and *nos*, [35]) was investigated, but no clear, shared

pattern of upregulation or downregulation was observed (Table S1).

Antimicrobial tolerance increased in the *P. aeruginosa* PA14 EVPL-associated biofilm over

7 d despite limited antimicrobial resistance gene expression changes.

Our previous work highlighted significant differential expression of antimicrobial resistance

(AMR) genes in the *P. aeruginosa* PA14 EVPL-associated biofilm compared with *in vitro*

growth at both 24 h and 48 h [8]. Therefore, we next aimed to investigate whether any of

the changes in gene expression found over 7 d were linked to AMR.

Three AMR genes were found to be significantly reduced in expression over time in the EVPL

biofilm (Table 2) and none were found to be significantly increased. The full AMR gene list

investigated is shown in Table S3. The *brc-1* gene, encoding a major facilitator superfamily

(MFS) transporter associated with bicyclomycin resistance [36], was significantly reduced in

expression 24 h - 7 d. The other two significant DEGs: *mexG* and *arnA*, which encode a

transmembrane protein associated with multiple efflux pumps and a bifunctional UDP-

glucuronic acid decarboxylase/UDP-4-amino-4-deoxy-L-arabinose formyltransferase [1],

were significantly reduced in expression 24 h - 48 h. The enzyme encoded by *arnA* is

involved in lipid A modifications, linked to cationic antimicrobial peptide (CAMP) resistance

[37, 38], and was previously shown to be significantly overexpressed in the EVPL-associated

biofilm compared with *in vitro* SCFM growth at 24 h [8].

Table 2. *Pseudomonas aeruginosa* PA14 antibiotic resistance associated genes found to be significantly

differentially expressed in at least one contrast in the *ex vivo* pig lung-associated biofilm between 24 h, 48 h

and 7 d. The gene expression group denotes the expression profile the gene belonged to, with the

corresponding expression plot (Table 1). The $\log_2$ fold change ($\log_2$FC) in the contrast where the gene was

significantly differentially expressed is shown. A gene was considered significantly differentially expressed if

$|\log_2$ fold change$| \geq 1.5$ and $P < 0.05$.

Locus tag	Gene name	Gene product	Gene expression group	Time: $\log_2$FC
PA14_09540	*mexG*	Transmembrane protein	4	24 h to 48 h: -1.52
PA14_18350	*arnA*	Bifunctional UDP-glucuronic acid decarboxylase/UDP-4-amino-4-deoxy-L-arabinose formyltransferase	4	24 h to 48 h: -1.59
PA14_10470	*bcr-1*	MFS transporter	5	24 h to 7 d: -2.49

To determine whether reduction in expression of the CAMP resistance-associated gene *arnA*

had a phenotypic effect, *P. aeruginosa* PA14 was grown as a biofilm on the EVPL tissue for

24 h, 48 h or 7 d, and exposed to each of two clinically relevant CAMPs, colistin and

polymyxin B, for a further 24 h. *P. aeruginosa* grown as a biofilm for 24 h could not be

recovered following treatment with 256 µg ml^{-1} colistin (Figure 4A). However, after 48 h

initial growth, PA14 was inconsistently recovered (Figure 4B). By 7 d, all biofilm populations

were able to tolerate 256 µg ml^{-1} colistin (Figure 4C). This pattern was comparable to the

response to 256 µg ml^{-1} polymyxin B (Figure 4). The increased polymyxin tolerance was a

contrast to the limited changes in AMR gene expression observed, and the significantly

reduced expression of *arnA*. This may be linked to persistence of non-dividing cells in the

biofilm population. As we do not know the rate of *P. aeruginosa* division in the EVPL model,

it may be possible that there is an early increase in efflux pump and *arnA* expression that

modifies the cell surface to a CAMP-tolerant phenotype, which is maintained. Alternatively,

there may be other phenotypic factors driving increased tolerance, such as the established

biofilm providing protection against CAMPs.

Figure 4. The colony forming units per lung tissue piece (CFU lung^{-1}) for the *Pseudomonas aeruginosa* PA14

ex vivo pig lung tissue-associated biofilm of different ages following no antibiotic treatment, treatment with

256 µg ml^{-1} colistin, or 256 µg ml^{-1} polymyxin B. Three replicates from each of two independent pig lungs,

shown by different shaped data points (see key), were used. The *P. aeruginosa* PA14 biofilms were grown in

the *ex vivo* pig lung (EVPL) model for **(A)** 24 h **(B)** 48 h **(C)** 7 d, then exposed to either phosphate-buffered

saline (PBS) as a negative control (no antibiotic; pink), 256 µg ml^{-1} colistin (blue), or 256 µg ml^{-1} polymyxin B

(yellow) for a further 24 h. The CFU lung^{-1} was then determined and is shown on the graphs, data points below

the dashed horizontal line represent samples where no CFU were detected. The asterisks represent a

significant difference between treatments determined using post-hoc Tukey HSD tests following ANOVA (* = *P*

< 0.05). **(A)** Day 1 ANOVA: treatment $F_{2,12}$ = 30.32, *P* < 0.01; lung $F_{1,12}$ = 0.15, *P* = 0.71; interaction $F_{2,12}$ = 0.04, *P*

= 0.96. **(B)** Day 2 ANOVA: treatment $F_{2,12}$ = 10.53, *P* < 0.01; lung $F_{1,12}$ = 5.76, *P* = 0.03; interaction $F_{2,12}$ = 1.26, *P* =

0.32. **(C)** Day 7 ANOVA: treatment $F_{2,12}$ = 14.51, *P* < 0.01; lung $F_{1,12}$ = 4.12, *P* = 0.07; interaction $F_{2,12}$ = 2.30, *P* =

0.14.

A mature, structured *P. aeruginosa* biofilm was maintained for 7 d on the surface of the

EVPL tissue, associated with reduced expression of Pel biosynthesis.

Therefore, to further explore the increase in antimicrobial tolerance observed over 7 d,

gene expression changes in QS and biofilm formation pathways were investigated. Both are

involved in the establishment of antibiotic tolerant, chronic *P. aeruginosa* infections in the

CF lung, and the EVPL biofilm has been shown to capture these aspects of *in vivo* infection at

48 h PI [8]. There were no overall differences in QS gene expression in any of the time point

comparisons (Figure 5; Figure S4; Figure S5). As our previous study demonstrated that the

expression of QS-regulated genes in the EVPL-associated biofilm at 48 h PI is comparable to

human infection [19], this indicates a representative biofilm infection was maintained for 7

d (Figure S4). Other genes associated with *P. aeruginosa* PA14 virulence factor production

were also investigated, including those involved in motility, type III secretion, type VI

secretion, and alginate biosynthesis, however no significant differential expression was

found.

Figure 5. Pathway map of the gene expression changes in the *Pseudomonas aeruginosa* PA14 Kyoto Encyclopedia of Genes and Genomes (KEGG) biofilm formation and quorum sensing (QS) pathways in the *ex vivo* pig lung-associated biofilm from 24 h to 7 d. The diagram shows all genes involved in each pathway, including associated signalling pathways and the Pel exopolysaccharide biosynthesis pathway. The $\log_2$ fold change value from 24 h to 7 d is shown below each gene name. The genes shown in red were significantly underexpressed, and genes shown in green were significantly overexpressed (significance: $P < 0.05$, $|\log_2$ fold change$| \geq 1.5$). The blocked arrows represent gene activation, and the dashed arrows show indirect gene activation. The solid, blocked lines (ending with |, not an arrow) show inhibition, and the dashed, blocked lines show indirect inhibition. The blue 'P' represents phosphorylation.

In contrast, all genes in the operon for biosynthesis of the biofilm exopolysaccharide Pel

were found to be significantly reduced in expression 24 h - 7 d (Figure 5), five of which were

also significantly reduced 24 h - 48 h (Figure S5). This suggests that the biofilm matrix was

formed in the first 24 h, and maintained for 7 d. To investigate this hypothesis, the biofilm

formed at each of these time points was visualised using scanning electron microscopy

(SEM). Uninfected tissue was imaged as a negative control (Figure S6). A biofilm could not

be imaged at 24 h, however a dense, structured *P. aeruginosa* PA14 biofilm was clear at 48

h (Figure 6), consistent with previous work [9]. The biofilm population was maintained at 7

d, where a very dense biofilm of *P. aeruginosa* PA14 cells was observed (Figure 7).

A. Lung 5: *P. aeruginosa* PA14 (48 h)   **B.** Lung 6: *P. aeruginosa* PA14 (48 h)

C. Lung 5: *P. aeruginosa* PA14 (48 h)   **D.** Lung 6: *P. aeruginosa* PA14 (48 h)

Figure 6. Scanning electron micrographs of *Pseudomonas aeruginosa* PA14 biofilms associated with *ex vivo*

pig lung (EVPL) bronchiolar tissue 48 h after infection. Images were taken from biofilms formed on tissue

from two independent pig lungs: **(A, C)** lung 5 **(B, D)** lung 6, showing biofilm aggregates of cells. The thicker

fibrous structures are consistent with the appearance of collagen fibrils under SEM, while the finer fibrous

structures may be individual collagen fibres and/or elastin fibres – both major components of the connective

tissue in bronchioles. Scale bars are shown. Images were taken using a Zeiss Gemini electron microscope with

an InLens detector, at 1kV, and analysed using Omero5.6 [30].

Figure 7. Scanning electron micrographs of *Pseudomonas aeruginosa* **PA14 biofilms associated with** *ex vivo*

pig lung (EVPL) bronchiolar tissue 7 d post infection Scale bars are shown. Images were taken using a Zeiss

Gemini electron microscope with an InLens detector, at 1kV, and analysed using Omero5.6 [30]

Discussion

We have used RNA sequencing data [8] to investigate *P. aeruginosa* biofilm infection in the

EVPL model over 7 d. We identified upregulation of several non-stress related pathways,

including sulfur metabolism, and demonstrated increased antibiotic tolerance. SEM

confirmed that a dense, mature biofilm was formed by 48 h that was maintained at 7 d PI,

with no visible indications of stress. Our findings show that the EVPL model provides an

environment for *P. aeruginosa* biofilm formation that once a clinically relevant biofilm has

formed at 48 h [8], it can be maintained as a chronic-like infection for 7 d.

Sulfur metabolism had the highest percentage of significant DEGs associated of all the

significantly enriched KEGG pathways over time. More specifically, all genes in the sulfur

transporter operon associated with cysteine biosynthesis: *cysTWA*, were found to be

significantly overexpressed 24 h - 7 d. In *E. coli,* upregulation of this operon has been linked

to sulfate starvation [39]. However, investigation of the sulfate starvation response in *P.*

aeruginosa PAO1 and a cystic fibrosis isolate found cysteine biosynthesis genes were only

weakly upregulated when sulfate was limited [40]. *P. aeruginosa* PA14 upregulation of

sulfur metabolism in SCFM has actually been linked to a shift towards anaerobic respiration

[41], which is associated with increased antibiotic tolerance [42]. There have also been links

made with iron starvation and sulfur metabolism. However, during iron starvation, proteins

involved in cysteine biosynthesis and sulfur assimilation are reduced [43].

Additionally, during infection of the CF lung, there is an increase in sulfation of the airway

mucins compared with a non-CF lung [44, 45], in part caused by inflammation [46], that

chronically adapted *P. aeruginosa* can use. It has also been shown that there is enough

inorganic sulfate in CF sputum to facilitate *P. aeruginosa* growth without mucin as a source

[47], therefore sulfur starvation is unlikely in this environment. Therefore, upregulation of

sulfur metabolism in the EVPL biofilm was more likely a result of a change in lifestyle, such

as anaerobic respiration, than nutrient starvation. Hence, our findings may further support

the progression of *P. aeruginosa* biofilm infection in the EVPL model as capturing the

lifestyle changes observed in CF. However, sulfur and oxygen availability in the EVPL biofilm

should be determined in the future to confirm this hypothesis.

Our previous work demonstrated that the EVPL biofilm may more accurately recapitulate *P.*

aeruginosa QS gene expression in CF sputum than other *in vitro* models [8], so *P. aeruginosa*

PA14 changes in expression of genes in the three QS pathways over 7 d were investigated:

LasI/R, RhlI/R, and PQS, and no significant changes were found. Other regulatory pathways

involved in lifestyle changes, including the switch from acute to chronic infection, were also

investigated however no clear differences in expression were found either. There were also

no significant changes in virulence factor gene expression. It is known that as part of

adaptation to the CF lung and formation of a chronic infection, virulence factor production is

significantly reduced [48]. Therefore, *P. aeruginosa* may have already switched off virulence

factor production by 24 h as it formed the established biofilm visualised at 48 h, hence no

change in gene expression was observed. This is supported by the findings in our previous

study, in which we found two of the major protease-encoding genes *lasA* and *lasB* were

significantly underexpressed in the EVPL-associated biofilm compared with SCFM *in vitro*

growth at both 24 h and 48 h [8]. This may also explain why no significant differential

expression was observed in regulatory pathways including QS. Additionally, adaptation to

the CF lung is also linked with mutations in *P. aeruginosa* virulence pathways [48], therefore

future work looking at the accumulation of mutations over 7 d in the EVPL-associated *P.*

aeruginosa biofilm would provide further insight.

In contrast, significant changes in the expression of biofilm-associated genes were found in

the EVPL PA14 biofilm population over 7 d. Pel biosynthesis was significantly downregulated

24 h - 48 h, then maintained at the same expression level 48 h - 7 d, which resulted in

significant downregulation 24 h - 7 d. This indicated that Pel production was reduced after

24 h of infection, then maintained from 48 h onwards. Pel is a biofilm exopolysaccharide

essential for PA14 biofilm development and is only expressed during biofilm growth [49, 50].

Whilst our results do not confirm there was not any Pel being produced at the later time

points, they indicate less than the initial stage of infection in the EVPL model. This may have

been caused by upregulation of Pel biosynthesis as the biofilm was established at 24 h,

which was reduced once the biofilm was formed at 48 h and maintained at this lower level

over the remaining 5 d. This hypothesis was supported by SEM images of the biofilm, which

showed a dense, structured *P. aeruginosa* biofilm already present at 48 h.

Taking this into account, following formation of a biofilm, gene expression analysis indicated

that the focus may have shifted from production of the biofilm matrix to biofilm

maintenance and maturation 48 h - 7 d. Significant upregulation of two lectin genes was

also observed in the lung-associated biofilm 24 h - 7 d, and 24 h - 48 h. Both have been

associated with a mature biofilm; *lecA* is indicated to play a role in the formation of a

mature biofilm architecture [51], and *lecB* has been shown to be involved in increased cell

and exopolysaccharide retention in biofilms [52]. This retention may also have been

associated with the reduction in Pel production, as less new Pel would be required if the

exopolysaccharide already produced was being retained. The proposed maintenance of a

mature biofilm structure was also accompanied by increased antimicrobial tolerance.

In our first study of *P. aeruginosa* PA14 gene expression in the EVPL model compared with *in*

vitro SCFM growth, we found that the EVPL growth environments cued differential

expression of a number of AMR genes despite no antibiotic exposure, at both 24 h and 48 h

[8]. However, only three genes involved in AMR were differentially expressed over 7 d in the

EVPL biofilm, suggesting differential expression of resistance-associated genes cued by the

environment is unlikely to be dynamic and remains consistent throughout 7 d infection. The

significantly differentially expressed AMR genes that were found included significant

reduction in expression of two individual antibiotic resistance genes: *arnA* and *bcr-1*,

associated with resistance to CAMPs [38] and bicyclomycin [36] respectively.

Despite these findings, increased tolerance to colistin and polymyxin B was observed at 7 d,

compared with 24 h and 48 h. It is hypothesised that the formation of a biofilm, of which

the matrix facilitates antibiotic resistance [2], may have resulted in other mechanisms of

antibiotic resistance being downregulated as they were not required. This would provide

explanation as to why expression of AMR genes was reduced once a biofilm had formed,

and the matrix was no longer being rapidly produced. This finding would further support the

use of biofilm disrupters to improve treatment regimens, as the *P. aeruginosa* cells within

the biofilm may not be expressing genes associated with antibiotic resistance and are thus

more susceptible to treatments.

However, it must also be considered that there were other indirect mechanisms of

antibiotic resistance that were found to be significantly overexpressed in the EVPL biofilm,

which are associated with potential environmental pressures. Phosphate transporters were

significantly overexpressed from 24 h to 7 d, indicative of phosphate stress. Under

phosphate stress *P. aeruginosa* has been shown to undergo lipid modifications that confer

resistance to polymyxin B. There is a transition from phospholipids to glycolipids, which

confer protection against polymyxin B [53]. Therefore, this differential expression may also

be responsible for or involved in, the increased antibiotic tolerance observed. Future

investigation of *P. aeruginosa* lipid modifications in the EVPL model environment would

further understanding. This analysis has however indicated that these pathways may be

induced, thus investigation of lipid modifications cued by the EVPL model may highlight

novel treatment targets.

In summary, we have demonstrated that there are changes in the *P. aeruginosa* lung-

associated biofilm transcriptome in the EVPL model over 24h - 7 d of infection. We have

identified an increase in polymyxin tolerance 48h - 7d that is accompanied by an

upregulation of sulfur metabolism and phosphate transport, rather than AMR-associated

genes. Whilst further work is required to fully determine the chronic-like phenotype formed

by 7 d PI, our results highlight a key further time point to address future research questions

related to chronic lung infection using the EVPL model. Introduction of other factors that

influence *P. aeruginosa* gene expression in the CF lung, such as other microbes, may reveal

other changes in the biofilm population over time that facilitate survival in these conditions

in vivo. Further understanding of these mechanisms will provide better understanding to

eradicate or prevent chronic *P. aeruginosa* infections in the CF lung.

Acknowledgements

We would like to thank all at Steve Quigley & Sons butchers for donating the pig lungs used

for this work, and Leo Eberl for the bacterial strain used. We would also like to thank Jenna

Lam for her help with the RNA extraction process and Blessing Anonye for her guidance. We

also acknowledge the help of the Media Preparation Facility in the School of Life Sciences at

the University of Warwick, with special thanks to Cerith Harries and Caroline Stewart, and

the Electron Microscopy Research Technology Platform at the University of Warwick for use

of the Zeiss Gemini scanning electron microscope. This work was supported by an MRC New

Investigator Research Grant (grant number MR/R001898/1) awarded to F.H., a PhD

studentship from the BBSRC Midlands Integrative Biosciences Training Partnership (MIBTP)

awarded to N.E.H., a PhD studentship from the MRC Doctoral Training Programme in

Interdisciplinary Biomedical Research awarded to R.G.M., and the University of Warwick as

part of F.A.'s MBio degree in Biomedical Science.

Author Contributions

567   N.E.H. and F.H. designed research; N.E.H., F.A., and R.G.M. performed research; N.E.H. and

568   F.A. analyzed data; N.E.H. and F.A. wrote the paper; N.E.H., F.A., R.G.M., and F.H. edited the

569   paper.

Conflicts of Interest

573   The authors declare no conflict of interest.

References

1.   Winsor, G.L., et al., *Enhanced annotations and features for comparing thousands of*
Pseudomonas genomes in the Pseudomonas genome database. Nucleic Acids
Research, 2016. **44**(D1): p. D646-D653.
2.   Høiby, N., et al., *Antibiotic resistance of bacterial biofilms.* International Journal of
Antimicrobial Agents, 2010. **35**(4): p. 322-332.
3.   Wardell, S.J.T., et al., *Genome evolution drives transcriptomic and phenotypic*
adaptation in Pseudomonas aeruginosa during 20 years of infection. Microbial
Genomics, 2021. **7**(11).
4.   Hogardt, M. and J. Heesemann, *Microevolution of Pseudomonas aeruginosa to a*
Chronic Pathogen of the Cystic Fibrosis Lung, in *Between Pathogenicity and*
Commensalism. 2011, Springer Berlin Heidelberg. p. 91-118.

5.   Rossi, E., et al., *Pseudomonas aeruginosa adaptation and evolution in patients with*
cystic fibrosis. Nature Reviews Microbiology, 2021. **19**(5): p. 331-342.

6.   Jiricny, N., et al., *Loss of Social Behaviours in Populations of Pseudomonas aeruginosa*
Infecting Lungs of Patients with Cystic Fibrosis. PLoS ONE, 2014. **9**(1): p. e83124.

7.   Cornforth, D.M., et al., *Pseudomonas aeruginosa transcriptome during human*
infection. PNAS, 2018. **115**(22): p. E5125-E5134.

8.   Harrington, N.E., J.L. Littler, and F. Harrison, *Transcriptome Analysis of Pseudomonas*
aeruginosa Biofilm Infection in an Ex Vivo Pig Model of the Cystic Fibrosis Lung.
Applied and Environmental Microbiology, 2022. **88**(3).

9.   Harrington, N.E., E. Sweeney, and F. Harrison, *Building a better biofilm - Formation of*
in vivo-like biofilm structures by Pseudomonas aeruginosa in a porcine model of
cystic fibrosis lung infection. Biofilm, 2020. **2**.

10.   Liberati, N.T., et al., *An ordered, nonredundant library of Pseudomonas aeruginosa*
strain PA14 transposon insertion mutants. Proceedings of the National Academy of
Sciences, 2006. **103**(8): p. 2833-2838.

11.   Liberati, N.T., et al. *PA14 transposon insertion mutant library*. 2006 [cited 2022
10/05/2022]; Available from: http://pa14.mgh.harvard.edu/cgi-bin/pa14/home.cgi.

12.   Palmer, K.L., L.M. Aye, and M. Whiteley, *Nutritional Cues Control Pseudomonas*
aeruginosa Multicellular Behavior in Cystic Fibrosis Sputum. Journal of Bacteriology,
2007. **189**(22): p. 8079-8087.

13.   Harrison, F. and S.P. Diggle, *An ex vivo lung model to study bronchioles infected with*
Pseudomonas aeruginosa biofilms. Microbiology, 2016. **162**(10): p. 1755-1760.

14.   Harrington, N.E., et al., *Antibiotic Efficacy Testing in an Ex vivo Model of*
Pseudomonas aeruginosa and Staphylococcus aureus Biofilms in the Cystic Fibrosis
Lung. Journal of Visualized Experiments, 2021(167).

15.   Andrews, S. *FastQC: a quality control tool for high throughput sequence data.* 2018;
Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

16.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina*
sequence data. Bioinformatics, 2014. **30**(15): p. 2114-2120.

17.   Kopylova, E., L. Noé, and H. Touzet, *SortMeRNA: fast and accurate filtering of*
ribosomal RNAs in metatranscriptomic data. Bioinformatics, 2012. **28**(24): p. 3211-
3217.

18.   Kim, D., et al., *Graph-based genome alignment and genotyping with HISAT2 and*
HISAT-genotype. Nature Biotechnology, 2019. **37**(8): p. 907-915.

19.   Li, H. *A toolkit for processing sequences in FASTA/Q formats.* 2015; Available from:
https://github.com/lh3/seqtk.

20.   Li, H., *Aligning sequence reads, clone sequences and assembly contigs with BWA-*
MEM. ArXiv, 2013. **00**.

21.   Liao, Y., G.K. Smyth, and W. Shi, *The R package Rsubread is easier, faster, cheaper*
and better for alignment and quantification of RNA sequencing reads. Nucleic Acids
Research, 2019. **47**(8): p. e47-e47.

22.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
dispersion for RNA-seq data with DESeq2. Genome Biology, 2014. **15**(12).

23.   Kassambara, A. *ggpubr: "ggplot2" based publication ready plots. R package version*
0.4.0. 2020; Available from: https://cran.r-project.org/package=ggpubr.

24.   Chen, H., *VennDiagram: Generate High-Resolution Venn and Euler Plots. R package*
version 1.7.0. 2021.

25.   Yu, G., et al., *ClusterProfiler: an R package for comparing biological themes among*
635   *gene clusters.* OMICS, 2012. **16**: p. 284-287.
636   26.   Alcock, B.P., et al., *CARD 2020: antibiotic resistome surveillance with the*
637   *comprehensive antibiotic resistance database.* Nucleic Acids Research, 2019.
638   27.   Kanehisa, M. and S. Goto, *KEGG: Kyoto Encyclopedia of Genes and Genomes.* Nucleic
639   Acids Research, 2000. **28**(1): p. 27-30.
640   28.   Kanehisa, M., *Toward understanding the origin and evolution of cellular organisms.*
641   Protein Science, 2019. **28**(11): p. 1947-1951.
642   29.   Kanehisa, M., et al., *KEGG: integrating viruses and cellular organisms.* Nucleic Acids
643   Research, 2021. **49**(D1): p. D545-D551.
644   30.   Goldberg, I.G., et al., *The Open Microscopy Environment (OME) Data Model and XML*
645   *file: open tools for informatics and quantitative analysis in biological imaging.*
646   Genome Biology, 2005. **6**(5): p. R47.
647   31.   Adalsteinsson, D. and P. Schultz, *The DataGraph Manual.* 2020.
648   32.   Schreiber, K., et al., *Anaerobic Survival of Pseudomonas aeruginosa by Pyruvate*
649   *Fermentation Requires an Usp-Type Stress Protein.* Journal of Bacteriology, 2006.
650   **188**(2): p. 659-668.
651   33.   Eschbach, M., et al., *Long-Term Anaerobic Survival of the Opportunistic Pathogen*
652   *Pseudomons aeruginosa via Pyruvate Fermentation.* Journal of Bacteriology, 2004.
653   **186**(14).
654   34.   Mercenier, A., et al., *Regulation of enzyme synthesis in the arginine deiminase*
655   *pathway of Pseudomonas aeruginosa.* Journal of Bacteriology, 1980. **144**(1).
656   35.   Rossi, E., et al., *High-resolution in situ transcriptomics of Pseudomonas aeruginosa*
657   *unveils genotype independent patho-phenotypes in cystic fibrosis lungs.* Nature
658   Communications, 2018. **9**(1).
659   36.   Fonseca, E.L., et al., *Full characterization of the integrative and conjugative element*
660   *carrying the metallo-β-lactamase blaSPM-1 and bicyclomycin bcr1 resistance genes*
661   *found in the pandemic Pseudomonas aeruginosa clone SP/ST277.* Journal of
662   Antimicrobial Chemotherapy, 2015. **70**(9): p. 2547-2550.
663   37.   Roy, S., et al., *Transcriptional expression of PmrB and arnA within polymyxin-*
664   *resistant nosocomial isolates of Pseudomonas aeruginosa from India.* Journal of
665   Microbiology and Infectious Diseases, 2018. **8**(3): p. 61-68.
666   38.   Gutu, A.D., et al., *Pseudomonas aeruginosa High-Level Resistance to Polymyxins and*
667   *Other Antimicrobial Peptides Requires cprA, a Gene That Is Disrupted in the PAO1*
668   *Strain.* Antimicrobial Agents and Chemotherapy, 2015. **59**(9): p. 5377-5387.
669   39.   Gyaneshwar, P., et al., *Sulfur and Nitrogen Limitation in Escherichia coli K-12: Specific*
670   *Homeostatic Responses.* Journal of Bacteriology, 2005. **187**(3): p. 1074-1090.
671   40.   Tralau, T., et al., *Transcriptomic Analysis of the Sulfate Starvation Response of*
672   *Pseudomonas aeruginosa.* Journal of Bacteriology, 2007. **189**(19): p. 6743-6750.
673   41.   Tata, M., et al., *RNASeq Based Transcriptional Profiling of Pseudomonas aeruginosa*
674   *PA14 after Short- and Long-Term Anoxic Cultivation in Synthetic Cystic Fibrosis*
675   *Sputum Medium.* PLOS ONE, 2016. **11**(1): p. e0147811.
676   42.   Schobert, M. and D. Jahn, *Anaerobic physiology of Pseudomonas aeruginosa in the*
677   *cystic fibrosis lung.* International Journal of Medical Microbiology, 2010. **300**(8): p.
678   549-556.
679   43.   Nelson, C.E., et al., *Proteomic Analysis of the Pseudomonas aeruginosa Iron*
680   *Starvation Response Reveals PrrF Small Regulatory RNA-Dependent Iron Regulation*

of Twitching Motility, Amino Acid Metabolism, and Zinc Homeostasis Proteins. Journal of Bacteriology, 2019. **201**(12).

44.   Cheng, P.W., et al., *Increased sulfation of glycoconjugates by cultured nasal epithelial cells from patients with cystic fibrosis.* Journal of Clinical Investigation, 1989. **84**(1): p. 68-72.

45.   Xia, B., et al., *Altered O-glycosylation and sulfation of airway mucins associated with cystic fibrosis.* Glycobiology, 2005. **15**(8): p. 747-775.

46.   Lamblin, G., et al., *Human airway mucin glycosylation: A combinatory of carbohydrate determinants which vary in cystic fibrosis.* Glycoconjugate Journal, 2001. **18**(9): p. 661-684.

47.   Robinson, C.V., et al., *Desulfurization of mucin by Pseudomonas aeruginosa: influence of sulfate in the lungs of cystic fibrosis patients.* Journal of Medical Microbiology, 2012. **61**(12): p. 1644-1653.

48.   Smith, E.E., et al., *Genetic adaptation by Pseudomonas aeruginosa to the airways of cystic fibrosis patients.* Proceedings of the National Academy of Sciences, 2006. **103**(22): p. 8487-8492.

49.   Colvin, K.M., et al., *The Pel Polysaccharide Can Serve a Structural and Protective Role in the Biofilm Matrix of Pseudomonas aeruginosa.* PLoS Pathogens, 2011. **7**(1): p. e1001264.

50.   Marmont, L.S., et al., *PelA and PelB proteins form a modification and secretion complex essential for Pel polysaccharide-dependent biofilm formation in Pseudomonas aeruginosa.* Journal of Biological Chemistry, 2017. **292**(47): p. 19411-19422.

51.   Diggle, S.P., et al., *The galactophilic lectin, LecA, contributes to biofilm development in Pseudomonas aeruginosa.* Environmental Microbiology, 2006. **8**(6): p. 1095-1104.

52.   Passos Da Silva, D., et al., *The Pseudomonas aeruginosa lectin LecB binds to the exopolysaccharide Psl and stabilizes the biofilm matrix.* Nature Communications, 2019. **10**(1).

53.   Jones, R.A., et al., *Phosphorus stress induces the synthesis of novel glycolipids in Pseudomonas aeruginosa that confer protection against a last-resort antibiotic.* The ISME Journal, 2021. **15**(11): p. 3303-3314.