1

3   Cavity geometry shapes overall ant colony organization through spatial limits but workers

4   maintain fidelity zones

6   Greg T. Chism[a*], William Nichols[b], Anna Dornhaus[c*]

10   [a] School of Information, University of Arizona, Tucson, AZ, U.S.A.
11   [b] School of Plant Sciences, University of Arizona, Tucson, AZ, U.S.A.
12   [c] Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, U.S.A.
13

14   * Corresponding Author
15   E-mail: dornhaus@email.arizona.edu
16

Abstract

Many animals inhabit nests that protect them from adverse environments. However, the effects

of living in a built or found structure are not limited to protection: the physical space can shape

and organize behavior, particularly in self-organized collective systems. In addition, the

geometry of nest space may not be under the animal's control, raising the question whether

animals can compensate for the effects that unexpected or suboptimal geometries may have.

Here we examine how the shape of a nest cavity affects spatial organization of colonies in the ant

Temnothorax rugatulus, a species that adapted to nest cavities of unmodifiable internal

dimensions, since they inhabit rock crevices with rigid walls. We show that the emerging spatial

relationships of workers, brood, queens, and young alates, as well as their relationships and

distances to significant points in the nest, are all significantly influenced by nest shape, with the

brood distributions most affected. However, we also found that the size of worker spatial fidelity

zones, i.e. the areas in the nest that individual workers occupy and that may be key regulators of

division of labor, are overall not affected by nest shape. These findings indicate that ants may

actively regulate which areas of a nest they occupy, and that they may compensate for effects of

nest architecture constraints. Physical properties of nests can thus influence the organization of

ant colonies, highlighting the need to explore spatial constraints as a direct influence on the

organization, movement, and communication of evolved or engineered self-organized systems.

Introduction

Social animal architectures benefit individuals by opening a wider range of suitable habitats that

they, and their kin, can successfully inhabit. These architectures, i.e., built environmental

modifications, range from simple depressions in the ground (e.g., social crab burrows, Laidre et

al., 2018) to massive above and below ground structures that span meters across (e.g.,

Macrotermes termites, Harris 1956). This diversity is unified by the impact each architecture can

have on the social behavior of their occupants: social weaver bird nests promote altruism through

spatial and social clustering (van Dijk et al., 2015), kin cohabitation increases with burrow size

in social crabs (Laidre et al., 2018, Laidre 2019), and reproductive division of labor is facilitated

by both naked mole rat burrows (Tofts and Franks 1992; Faulkes and Bennett 2001) and social

insects nests (Wilson and Kinne 1990, Wilson 1992). Ant nests are particularly interesting

because of both their representation in nearly every environment, and their diversity in form and

function within these environments (Wilson and Kinne 1990, Wilson 1992).

Here we are interested in how the geometric shape of such built architectures influences the

organization of the colony within. Ants have been demonstrated to be distributed non-uniformly

and non-randomly across their internal nest space in many cases (Sendova-Franks and Franks

1995; Sendova-Franks and Franks 1999; Tschinkel 1999; Tschinkel 2005). For example, colony

members may be concentrated in specific sections of the nest (Tschinkel 1999; Tschinkel 2005);

and workers may use chemical 'road-signs' to navigate nest space and differentiate chambers

with different functions (Heyman et al., 2017). However, some complex spatial relationships in

the nest may be predicted by random walk models (Sendova-Franks and Van Lent 2002,

Davidson and Gordon 2017), implying that the placement of colony members may not always be

an explicitly adaptive strategy. In addition, many ants exploit pre-existing natural structures as

nests (e.g., *Temnothorax* ants: Prebus 2017, *Cephalotes* turtle ants that occupy unmodifiable

wood nests: Powell 2008, and rock dwelling *Rhytidoponera metallica* ponerine ants: Thomas

2002), which means that the geometry of the available space may not be under the ants' control.

We therefore ask how the internal nest space may influence the spatial organization of workers,

queens, and brood in the nest.

It has become clear over the last decade that the pattern of interactions in and allowed by social

insect nests may affect colony organization. For example, worker interactions and recruitment

increase in nest chambers that are highly connected to other nest chambers (Pinter-Wollman

2015; Vaes et al., 2020), information processing and exploitation of the environment decreases in

nests with multiple entrances (Lehue and Detrain 2019; Lehue and Detain 2020; Lehue et al.,

2020a,b; Lehue and Detrain 2020), while panicked workers bottleneck when evacuating through

a single nest entrance (Burd et al., 2010; Wang and Song 2016). Further, worker density in a nest

increases the likelihood that colonies become polydomous in the ant *Temnothorax rugatulus*

(Cao 2013). Social interactions and heterogeneity in spatial distributions in the nest are closely

linked (Guo et al., 2020). Information, such as alarm signals, may be spread through worker

physical contact, in which case the spatial arrangement of workers critically affects information

flow (Guo et al., 2022). Further, queen(s) and brood may be positioned in nests so that they are

both more accessible from other nest spaces and possibly more defensible during nest invasions

(Varoudis et al., 2018). Ant colonies further reduce disease spread through social distancing:

disease spread is reduced by adaptively clustered (rather than globally random) worker

interactions, and patterns of interactions thus may confer organizational immunity (Stroeymeyt et

al., 2018; Pusceddu et al., 2021; Naug and Smith 2007). It thus seems clear that worker (and

other colony member) interactions and density may affect colony-level performance in essential

tasks both inside and outside the nest; and thus, properties of the inhabited space, which are

likely to affect such patterns of interaction and spatial distribution, are in turn critical to

understand collective colony function.

One context in which spatial relationships appear particularly important is task allocation and

division of labor. Although this may vary between tasks and species, proximity of individuals to

areas where particular work is needed seems to be a major factor affecting task allocation

(Leitner and Dornhaus 2019). Workers do not move through the nest randomly, and are often

faithful to particular, often quite small, mini-'territories' in the nest called 'spatial fidelity zones'.

Spatial fidelity zones have been demonstrated in multiple species, and the location of these zones

often correlates with worker task (Franks and Tofts 1994; Sendova-Franks and Franks 1994;

Sendova-Franks and Franks 1995; Sendova-Franks and Franks 1999; Powell and Tschinkel

1999; Jandt and Dornhaus 2009). The size of these fidelity zones varies with task, location in the

nest, or specialization of workers (e.g. smaller fidelity zones in workers performing brood care,

Sendova-Franks and Franks 1995; Jandt and Dornhaus 2009, or inactive workers, Charbonneau

et al., 2017a, Leitner and Dornhaus 2019).

Nest-related collective behavior has been well studied in the ant genus *Temnothorax*. Rock-

dwelling *Temnothorax* have been the subject of studies involving house-hunting (Franks et al.,

2002; Dornhaus et al., 2004; Sasaki and Pratt 2013; Sasaki et al., 2015), collective nest building

(Franks et al., 1992; Franks and Deneubourg 1997; Aleksiev et al., 2007a; Aleksiev et al., 2007b;

Aleksiev et al., 2007c; DiRienzo and Dornhaus 2017), and risk-taking behaviors relating to

ecological nest site availability (Bengston and Dornhaus 2014; Bengston and Dornhaus 2015;

Bengston et al., 2017), among other topics. We used the rock ant *Temnothorax rugatulus* to

examine the influence of nest shape on its occupants. *T. rugatulus* colonies dwell in pre-existing

rock-crevices, producing single-chambered nests that are easily replicated and manipulated in a

laboratory setting (Charbonneau and Dornhaus 2015). Worker spatial position in *Temnothorax*

unifasciatus (a con-gener to *T. rugatulus* with similar rock-dwelling nesting behavior) nests

directly influences the task they perform, such as brood care near the brood pile (Sendova-Franks

and Franks 1995), and *T. unifasciatus* workers will return to their spatial positions in relation to

one another in new nests (Sendova-Franks and Franks 1994). This species thus provides a

suitable model system, in which we know that worker interactions and spatial arrangement are

relevant to the colony, but also are feasible to comprehensively monitor and manipulate; and

furthermore, ants in this genus likely have to adapt frequently to nest geometries that are largely

beyond their control.

We investigate the influence of internal nest geometry on colony organization in *T. rugatulus*

colonies for two artificial nest shapes designed to differ markedly in physical accessibility: a

'circle' and a 'tube' nest (Fig. 1). We quantify the distribution of all types of individuals in the

nests, and then ask how the location of the nest entrance and the centroid of the brood

distribution relate to where colony members are located in the nest. We also examine how the

distributions are affected by overall colony density in the nest. Finally, we determined whether

the spatial fidelity zones of individual workers are different across nest shapes, and how they

relate to the location of nest and colony features.

Methods

Colony collections

We collected 20 queenright colonies of *Temnothorax rugatulus* from rocky semi-steep slopes on

the Santa Catalina Mountains (GPS: 32.395, -110.688), USA, Pima County, Arizona from July

to October 2017 and February to May 2018. We found all of our colonies in a pine-juniper zone

(altitude approximately 2500m) inhabiting granite rock-crevices where entire colonies were

collected by aspiration after prying open the nest cavity. The second half of the colonies (IDs 11-

20) were collected during the colonies' reproductive period, which allowed us to examine alate

spatial organization (alates being winged ants, i.e. new queens and males). The first ten colonies

had only workers and (non-alate, i.e. reproductive) queens and brood, whereas seven of the latter

ten colonies additionally had alates. We did not identify or account for the presence of

microgynes (diminutive but possibly reproductive queens) in our study and only identified

obvious queens (significant size and color difference compared to workers and proximity to a

cluster of her brood). Microgynes were likely present however, given that they have been shown

to be prevalent in higher elevation populations of this species (Choppin et al., 2021). Microgynes

were most likely categorized as workers in our study. They are likely to be few in number if they

occur and behave and are treated differently than macrogynes (regular, large-size reproductive

queens) by workers (fed more but groomed less, Negroni et al., 2021).

Initial housing and care

We initially placed all colonies in generic artificial nests resembling their natural nest sites in

rock crevices. These nests consisted of a 2-mm-thick piece of cardboard (75 mm x 50 mm x 2

mm) sandwiched between two glass panes (76.2 mm x 50.8 mm x 0.5 mm), with a 2 mm x 2 mm

entrance at the center of one long side leading to an open nest space (35 mm x 25 mm)

(Charbonneau et al., 2015; Fig. A1). We gave colonies food and water *ad libitum*, refreshed

weekly, for the duration of their housing: water was given through water-filled, cotton-ball-

stopped 5 ml plastic tubes, and food was given through 2 ml microcentrifuge tubes of honey

water with a concentration of 1/4 teaspoon honey per 50ml water, and 1/8 (approximately 0.075

g) of a fresh-frozen cockroach (*Nauphoeta cinerea*). We kept colonies on a 12:12 hour light

cycle (8 a.m. to 8 p.m.), constant temperature (approximately 21-24 °C). We placed all artificial

nests in open-top plastic containers (11.1 cm x 11.1 cm x 3.3 cm) with walls lined with 'insect-a-

slip' (BioQuip 2871A, 'fluon') to prevent escape. We individually marked CO_2-anesthetized

workers with four identifying marks with multicolored paints (Testor's Pactra® paint): one each

on the head and prosoma, and two on the gaster (Charbonneau and Dornhaus 2015). We marked

ants three to five days before the colony was placed into experimental nests.

Experimental setup

We used two nest types, a circle and a tube shape laser-cut from the 2mm thick cardboard, for

the experimental phase (Fig. 1). We gave colonies comparable nest densities by scaling the

internal area of the nests to colony size (number of workers). We produced nests with high

worker density, by examining the size of a *Temnothorax rugatulus* colony that utilized nearly all

the available nest space in a pre-experimental nest (248 workers, Fig. A1) and doubling the area

per worker to permit flexible nest space use (0.033 mm^2 per worker). We further produced a low

nest density treatment by doubling the nest area allocated to each worker compared to that first

treatment, producing a density half as dense as the high-density treatment (0.066 mm^2 per

worker). It is known that colonies favor nests that allow low in-nest worker densities (Franks et

al., 2006; Visscher 2007; Mitrus 2015). However, these populations are thought to be nest-site-

limited (Foitzik et al., 2004; Bengston and Dornhaus 2015), such that under conditions of high

nest competition, colonies may often nest in sub-optimal nests or be prevented from expanding

into multiple nests despite high density.

Figure 1. Nest shapes used for this study: (a) circle and (b) tube nests. Internal nest area was

always scaled to the number of workers in the colony. The nest entrance is in the middle lower

edge of the image in both cases. A green background was used to make the brood items (white)

more visible. Pictured here are high-density treatment nest shapes for one colony.

Experimental timeline

Each colony experienced both nest shapes and both density treatments. We first randomly

assigned each colony one of the nest shapes, such that ten of our colonies experienced the circle

nest first and assigned the other ten colonies the tube nest first. We then forced an emigration

into the new assigned nest by offering that nest and removing the top glass pane from their

current nest, which prompts the ants to emigrate into the new cavity. After this emigration, we

allowed the ants to acclimate to the new nest for three days, and then photographed colonies in

this nest each day for 16 days. Once the 'nest assignment I' phase of 16 days was completed, we

forced an emigration into the other nest type, and repeated the three-day acclimation and 16-day

monitoring phase ('nest assignment II'). Photos were taken with a digital HD SLR camera

(Nikon D7000 with 60 mm lens). We used the first 10 photos from each nest assignment for

analyses excluding only images that were problematic (e.g. out of focus).

Photo analysis

We assigned cartesian coordinates to each colony member on the digital images and standardized

(i.e. translated them back into real distances in mm by using the bottom-left and top-right outside

corners of the nest as reference coordinates) them with the image analysis software *Fiji*

(Schindelin et al., 2012).

Nest sections and colony member densities

We divided each nest into eight equal-area sections from the nest entrance to the back of the nest

(1-8, see Fig. A2a-b). We chose not to use more than eight bins because these nest section

assignments were likely large enough to capture and segregate different worker tasks. We

determined all colony member densities by calculating what proportion of total observed

individuals of that type on the same image were found in that section.

Scaled distances in the nest

Distance to the nest entrance: We calculated the shortest linear distance *within the available nest*

cavity from each colony member to the nest entrance. In the circle nest, we straightforwardly

calculated each colony member's distance to the nest entrance (per a scaled reference x and y

coordinate) using the formula: sqrt((colony member x - entrance x)^2 + (colony member y -

entrance y)^2), with a small adjustment for the nest entrance tunnel. In the tube nest however, a

direct (i.e., straight-line) path would not reflect the actual walking distance. Thus, we calculated

the distance as the shortest possible line within the nest cavity to the entrance (see Fig. A3 in

supplementary information for details and illustration).

Distance to the brood center: We calculated the brood center as the average brood x and y

coordinates in the nest section with the most brood items in each observation. This was done

because calculating the overall centroid of brood across the entire nest would often take this

'center' outside of the nest space boundaries in the tube nest. We then found the distance from

each mobile colony member to this brood center (similar to the calculation for distance to the

nest entrance, this was calculated as straight-line distance in circle nests, and as walking distance

in the tube nest; for details see Fig. A3 in supplementary information). *Distance scaling*: Actual

possible distances varied across nests since the nest dimensions were scaled to colony size to

standardize worker density across colonies (see above). To be able to compare relative

distributions across colonies, we scaled all calculated distances to nest size, by setting the

shortest distance from the back of the tube nest shape to the entrance as 1. These 'scaled'

distances are thus only relative to the size of that colony's tube nest, not relative to each nest

shape. Since the farthest point from the entrance in the circle nest is never as far as the farthest

point in an equal-area tube nest, the maximum 'scaled' distance in the circle nest for any colony

is thus 0.34 (i.e. 34% of the maximum available distance in the tube nest).

Site fidelity

We generated twenty-four total zones by again dividing each of the eight nest sections described

above into three (illustrated in Fig. A2). We assigned each observation of each worker with

identifiable unique color markings to one of these zones. We defined 'occurrence zone size' for

each worker as the total number of nest zones that the worker was observed in (out of the 24 total

possible zones). We defined 'spatial fidelity zones' as the number of zones in which the worker

was observed for at least 15% of its observations (note that zones, as nest sections, are all equal

area for the same colony but reflect different amount of area across colonies because the total

nest size is scaled to colony size). We also analyzed the true area (in cm^2) of site fidelity zones

(by adding the areas of the respective nest zones) in supplemental materials. We used this

method instead of, for example, a minimum convex polygon (Jandt and Dornhaus 2009;

Charbonneau, et al., 2017a) to ensure that we did not include areas that were not part of the nest

cavity in the 'tube'-shaped nest. We only included workers that were observed at least seven (out

of ten possible) times (136 out of 941 possible workers in the high-density treatment and 247 out

of 838 possible workers from the low-density treatment). This loss occurred in part because

although all workers were initially individually painted, many lost two or more of their four

color-combination marks.

While the 7 observations cutoff and the 15% threshold are both somewhat arbitrary, we reasoned

that they provide a sensible combination given the total number of observations available. For a

worker with between 7 and 10 observations, every zone it is observed in at least twice will be

counted towards its spatial fidelity zone (2/7=29%), but zones in which it is observed only once

will not be counted (1/7=14%). This method of calculating spatial fidelity thus enables us to

identify zones the worker is likely to return to and constitutes a compromise between excluding

too many observations/workers and reliability of the calculated zone sizes.

Data processing

We conducted all data processing using the statistical software R (*v4.3.1*; R Core Team 2017)

and RStudio (*v2023.06.1+524*; Allaire 2012), specifically the Tidyverse language ('tidyverse'

v1.3.1: Wickham et al., 2019). All original data and analysis scripts are publicly available on

GitHub at https://github.com/Gchism94/NestArchOrg (Chism 2022).

Colony distributions: We used linear mixed effects models to examine colony member

distributions (proportion in each nest section) in our experimental nests using the R package

'lme4' (*v1.1-27.1*; Bates et al., 2014), where *P* values were calculated through the R package

'lmerTest' (*v3.1-3*; Kuznetsova et al., 2017). We report the relationships between colony

member distributions across nest sections (quadratic term) and nest shape though the interactions

term (nest * nest section^2), and report how nest density affects this relationship through a three-

way interaction term (Nest * Nest section^2 * Nest density). Our linear mixed effects models here

and below (unless noted otherwise) had colony identification (hereby termed 'Colony ID') as a

random effect, and by comparing the variation explained by the fixed effects alone (marginal R^2)

and with the random effect included (conditional R^2), we determined the amount of variation that

Colony ID explained (marginal and conditional R^2 values calculated through the R package

'sjPlot' *v2.8.14*; Lüdecke 2023).

Scaled distances: We used linear mixed effects models to examine the relationships between nest

shape and individual scaled distances from the nest entrance and the center of the brood,

including the interaction of nest shape and density (Nest * Nest density).

Worker site fidelity: We used linear mixed effects models to test the effect of nest shape, density,

colony size, and colony ID on worker spatial fidelity and occurrence zone sizes. We also

consider the true zone sizes (in cm^2).

Worker site fidelity and distances in the nest: We used the same linear mixed effects models as

above to test the effect of either worker mean scaled distance to the nest entrance and nest shape,

and a separate linear mixed effects models to test the effect of worker mean scaled distance to

the brood center and nest shape on site fidelity. We additionally tested the interaction terms

between nest shape and colony size, nest shape and mean scaled distance to the nest entrance,

and colony size and nest density.

Results

Overall colony distributions in the two nest shapes

All types of colony members' distributions were significantly affected by nest shape.

Figure 2. The distribution of all types of ants in the two nest shapes (top row: circle nest, bottom

row: tube nest, pooled across all 400 photos from 20 colonies). Colors indicate the average

number of neighbors for each item within 0.16 of the longest distance in tube nest.

Most prominently, brood typically formed a distinct, dense, oval cluster in the circular nest, often

in the center or towards the back of the nest. In contrast, in the tube nest, brood was typically

found across the entire length of the nest and arranged in several smaller clusters, which led to

more brood (and workers) being located nearer the entrance of the nest (Fig. 2). We analyzed

both how ants were distributed in the nest sections, and thus how they were divided up across the

available space in the nest (details in Table 1), and the distribution of their actual distances from

the nest entrance (details in Table A1) and center of the brood (understood to reflect the

meaningful center of the colony; details in Table A2). Broadly, nest shape affected all colony

member distributions across nest sections except that of queens (Fig. 3: LME; n=3192

proportions in nest sections for workers, corresponding to 10 colonies over 4 treatments with 8

sections and 10 replicates each; similar numbers for brood and queens, n=456 for alates; all

p<0.01 for nest shape as a main effect, except for queens p=0.12). Density overall had no main

effect on distribution across nest sections, except for brood (p<0.03; Fig. 3), and 'Day', i.e. time

since emigration into the nest, also had no effects (all p>0.05), indicating that colonies settled

into their distributions by the beginning of data collection (3 days since emigration). Colony ID,

as the only random effect included in our model, did not explain any variation (identical

marginal and conditional R^2 values, Table 1).

Distances from the nest entrance

For both brood and queens, there were significant linear and quadratic effects of 'nest section'

(all p<0.001), indicating that both distributed with an intermediate peak along the axis from the

front to the back of the nest, whereas there were no main effects (linear or quadratic) of nest

section on worker distributions (p=0.85 and p=0.71), indicating workers were largely evenly

distributed across the nest space (as can also be seen in Fig. 2 and 3). However, nest shape

interacted significantly with nest section for worker distributions, and with the quadratic effect of

nest section for brood - indicating a change in the shape of the worker and brood distributions

across the extent of the nest with nest geometry.

When analyzing the specific, individual, scaled distances of each ant from the nest entrance (Fig.

4), we found that nest shape again influenced all colony members' distributions (including that

of queens: LME, n=30247 locations of workers, 59459 locations of brood items, 1041 locations

of queens, 1006 locations of alates, all p<0.001, Table A1): all were on average further from the

nest entrance in the tube nest. This result is not very surprising since the tube nest allows ants to

be located much further from the entrance than space is available in the circle nest. However, it

illustrates that while there are also effects of relative spread over the nest as stated in the last

section, there are obvious effects on real distances between ants when nest geometry allows

those to be larger (illustrated in Fig. 4). We also found that worker density in the nest did not

affect distances to the entrance (of any ant types, all p>0.05), and male alates were closer than

queen alates to the nest entrance (p<0.001 for the main effect of sex for alates). Over time,

workers and brood tended to move further, and alates closer to the nest entrance (effect of Day,

all p<0.001).

Linear Mixed Effects: colony member proportions within nest sections

Coeffcient	Workers					Brood					Queens					Alates				
	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P
Intercept	0.145	0.009	3181	16.566	**<0.001**	-0.127	0.013	3173	-9.489	**<0.001**	-0.161	0.031	3181	-5.157	**<0.001**	0.363	0.060	446	6.073	**<0.001**
NestSect	-0.006	0.004	3181	-1.438	0.151	0.090	0.006	3173	14.888	**<0.001**	0.118	0.014	3181	8.331	**<0.001**	-0.083	0.027	446	-3.028	**0.003**
NestSect2	0.000	0.000	3181	0.294	0.769	-0.006	0.001	3173	-9.596	**<0.001**	-0.010	0.001	3181	-6.479	**<0.001**	0.006	0.003	446	1.924	0.055
Nest	0.107	0.010	3181	10.486	**<0.001**	0.086	0.016	3173	5.487	**<0.001**	0.073	0.037	3181	2.003	**0.045**	-0.192	0.069	446	-2.787	**0.006**
Density	0.000	0.003	3181	0.000	1.000	0.000	0.004	3173	0.000	1.000	0.000	0.009	3181	0.000	1.000					
Day	0.000	0.000	3181	0.000	1.000	0.000	0.001	3173	0.000	1.000	0.000	0.001	3181	0.000	1.000	-0.000	0.003	446	-0.000	1.000
Corner	0.019	0.004	3181	5.502	**<0.001**	0.027	0.005	3173	5.019	**<0.001**	0.029	0.013	3181	2.261	**0.024**	-0.054	0.023	446	-2.340	**0.020**
NestSect:Nest	-0.040	0.006	3181	-7.358	**<0.001**	0.009	0.008	3173	1.016	0.310	0.001	0.020	3181	0.048	0.962	0.052	0.037	446	1.414	0.158
NestSect2:Nest	0.003	0.001	3181	4.316	**<0.001**	-0.005	0.001	3173	-5.974	**<0.001**	-0.004	0.002	3181	-1.701	0.089	-0.001	0.004	446	-0.150	0.881

Random Effects

σ^2	0.01		0.01	0.07	0.03
τ_{00}	0.00 Colony		0.00 Colony	0.00 Colony	0.00 Colony
N	20 Colony		20 Colony	20 Colony	7 Colony
Observations	3192		3184	3192	456
Marginal R2 / Conditional R2	0.235 / 0.235		0.285 / 0.285	0.070 / 0.070	0.097 / 0.097

Table 1. Proportions of colony members in each nest section related to nest shape and physical

properties. Here and below sections (NestSect) are numbered beginning at the nest entrance (1)

and end at the back of the nest (8); a quadratic term for nest section is included to allow for a

unimodal distribution shape. Density is the amount of area allocated for each worker (high =

0.033144 cm^2, low = 0.066288 cm^2). Day is the observation day (starting 3 days after colonies

moved into the nest). Corner is the presence of a bend in the tube nest or the corners around the

entrance tunnel in the circle nest in the respective nest section. The random effect Colony ID is

colony identity. Bold *P* values indicate significance. Model formula in R: PropColonyMember ~

poly(NestSect, degree = 2, raw = TRUE)*Nest + Density + Day + Corner + (1 | Colony ID)

Figure 3. While workers are fairly evenly distributed across all available nest sections, brood and

queens are clustered in intermediate areas in both nest shapes. Shown here are the proportion of

the respective ant types across the equal-area nest sections: workers (a, b), brood (c, d), queens

(e, f), and alates (g) in the tube (red) and circle (blue) nests and when at low (right column) or

high density (left column). Nest sections scale with colony size, worker density is kept constant.

Boxes represent first and third quartiles, the bar within boxes is the median, and whiskers show

the data range. For corresponding statistics see Table 1.

Figure 4. Actual distances of workers and brood from the nest entrance scaled only by nest size,

by setting the farthest point of the tube nest as 1. When comparing these distances to the

distributions across sections as in Fig. 3, it becomes clear how colonies in tube nests are spread

out more from the entrance and from each other relative to the circle nest (where the range of

farthest possible distances from the entrance is indicated by the gray lines). Shown are worker (a,

b) and brood (c, d) distributions for the circle (blue) and tube (red) nests, and for high (a, c) and

low (b, d) worker density treatments. N: workers = 30247 distance measurements, brood =

59459, queens = 1041, alates = 1006, over a total of 400 photos of 20 colonies. Colonies are

sorted in order of increasing colony (and nest) size. For statistical analysis see Table A1.

Distances from the center of the brood

As with distance to the entrance, we found that workers, queens, and alates were farther from the

brood center in the tube nest (all $p<0.01$, Fig. 5, Table A2). We also found that male alates were

found farther from the brood center than queen alates (Fig. A5d, Table A2), and that neither

density nor time ('Day') had an effect (Table A2). Therefore, elongated nest shapes like our tube

nest likely force colony members to real distances farther from unique points such as the nest

entrance, the center of the brood, and the queen.

Figure 5. Workers are often dispersed farther from the center of the brood pile in the tube nest

than what is possible in the circle nest (scaled only by colony size, by setting the farthest point of

the tube nest as 1). Distributions are shown across high (a) and low (b) nest densities. N = 26733

observations of worker ants. For statistical analysis see Table A2.

The effect of corners

The two nest shapes in our study were designed to maximally differ in their diameter/area ratio,

i.e. elongation along a length axis, which simultaneously made one nest narrow, possibly

limiting ants from assessing density in all parts of the nest. However, since we arranged both

nests to be contained within our standard 2x3inch nest size, this also caused the tube nest to be

'bent', i.e. to form corners. The circle nest also has slight heterogeneities where the actual

'circle' is met by the short entrance tunnel. We found that these corners, which might have

affected movement, affected spatial distributions of workers, brood, queens, and alates in the

nests (with more workers, brood, and queens in 'corners', and fewer alates: all $p<0.05$ for the

effect of 'nest section including a corner' on proportion of individuals in that section, Table 1).

Small differences between colonies in spatial distributions

Colony identity was included as a random effect in the statistical models for both proximity to

the nest entrance (Table A1) and to the center of the brood (Table A2). The impact of random

effects can be estimated from the difference between marginal and conditional R^2 values; here,

we found that colony ID explained very little of the variation among worker distances (1.5% for

the distance to the entrance, 2.2% for distances to the center of the brood), but more of the

variation among for brood and queens (4.8% for distance of brood to entrance; 6.3% and 7.8%

for of variation in distance to entrance and brood center, respectively, for queens), but a larger

amount for the distances of alates from the entrance (17.2% and 0.6% respectively). This may

reflect that the positions of alates are more idiosyncratic by colony and depend on the sex of the

alates.

Worker site fidelity is a function of colony size and density, not geometry…

Despite these significant impacts of nest geometry on the distances of colony members to each

other and the distributions particularly of brood, we did not find any impact of nest shape on

worker spatial fidelity (LME, p=0.96 for scaled zone sizes measured as number of nest sections,

p=0.16 for actual zone sizes in cm^2, N=384 worker spatial fidelity zones from a total of 81

unique workers, Fig. 6, Table 2) or occurrence (p=0.94 for scaled, p=0.10 for actual) zone sizes.

Instead, the main predictor of how dispersed an individual's movement through the nest were

seemed to be the total nest size (which we scaled with colony size to keep a controlled worker

density in our experiment; Fig. 7): colony size did not predict scaled zone sizes but did strongly

correlate with absolute zone sizes, and also interacted significantly with density in determining

absolute zone sizes (different slopes in Fig. 7 a vs b and c vs d). This (the strong effect of

'Colony size' but not density, and the effect on absolute zone area but not proportion of the nest

used) is consistent with the idea that it is available area, not the number of individuals, that

matters. The average distance of the worker's location from the nest entrance or brood center

impacted neither fidelity zone nor occurrence zone sizes (Tables 2 and A3, Fig. 6). Colonies

differed only very slightly in their workers' zone sizes (Colony ID explained 6-7% of variation

in scaled zone sizes, and only 1% of variation in absolute zone sizes, Table 2).

There was also a significant interaction effect of nest shape and colony size for the absolute size

of occurrence zones (p=0.03); as seen in Fig. 7, occurrence zone sizes increase more slowly with

the number of individuals (~total available area) in the tube nest than the circle nest. Since the

same was not found for the more conservatively defined fidelity zones (p=0.40), it is possible

that this is because workers in the tube nest wandered into zones outside their typical fidelity

zone more rarely in the tube nest, particularly in large nests.

Linear Mixed Effects: individual worker site fidelity in each nest shape

Coeffcient	Scaled Fidelity Zone					Fidelity Zone (cm^2)					Scaled Occurrence Zone					Occurrence Zone (cm^2)				
	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P
Intercept	1.772	0.401	374	4.413	**<0.001**	0.053	0.109	374	0.483	0.629	5.498	0.592	374	9.286	**<0.001**	-0.222	0.183	374	-1.214	0.225
Nest	-0.024	0.442	374	-0.055	0.957	-0.179	0.128	374	-1.395	0.164	0.608	0.644	374	0.944	0.346	0.343	0.205	374	1.675	0.095
ColonySize	-0.003	0.002	374	-1.201	0.231	0.002	0.001	374	3.093	**0.002**	0.000	0.004	374	0.132	0.895	0.009	0.001	374	8.128	**<0.001**
MeanScaledDist	0.980	1.592	374	0.616	0.539	-0.042	0.462	374	-0.092	0.927	2.488	2.322	374	1.072	0.285	1.167	0.737	374	1.584	0.114
Density	-0.489	0.373	374	-1.311	0.191	-0.030	0.088	374	-0.337	0.737	0.113	0.560	374	0.202	0.840	0.052	0.165	374	0.316	0.752
Nest:ColonySize	0.001	0.002	374	0.243	0.808	0.001	0.001	374	0.849	0.396	-0.004	0.003	374	-1.307	0.192	-0.002	0.001	374	-2.152	**0.032**
Nest:MeanScaledDist	-0.852	1.663	374	-0.512	0.609	0.240	0.483	374	0.497	0.619	-2.318	2.426	374	-0.955	0.340	-1.118	0.770	374	-1.452	0.147
ColonySize:Density	0.002	0.003	374	0.721	0.472	0.002	0.001	374	2.507	**0.013**	0.001	0.005	374	0.239	0.811	0.008	0.001	374	6.074	**<0.001**

Random Effects

σ^2	0.70	0.06	1.50	0.15
τ_{00}	0.05 Colony	0.00 Colony	0.11 Colony	0.01 Colony
N	19 Colony	19 Colony	19 Colony	19 Colony
Observations	384	384	384	384
Marginal R^2 / Conditional R^2	0.026 / 0.085	0.341 / 0.353	0.020 / 0.087	0.803 / 0.813

Table 2. Size of worker spatial fidelity zones and occurrence zones related to nest shape (Nest), nest and colony size (ColonySize), density treatment (Density), and distance of the average position from the nest entrance (MeanScaledDist). Scaled zone sizes are as a proportion of total nest area, other zone sizes are in actual area. The random effect Colony ID is colony identity. Bold *P* values indicate significance. Model formula in R: Zone ~ Nest * ColonySize + Nest * MeanScaledDist + ColonySize * Density + (1 | ColonyID)

Figure 6. Worker spatial fidelity and occurrence zone sizes (in nest sections out of 24 total) were

constant across the two nest shapes and did not differ with density, distance to the nest entrance

(see Table 2 for statistics) or distance to the brood center (see Table A3 for statistics). Fidelity

zones: sections in which workers spent at least 15% of observations: a, b; Occurrence occurrence

zones: all sections in which a worker was observed: c, d. N: 383 marked ants across 19 colonies

had sufficient observations to be included in the analysis (74 in circle/high density, 148 in

circle/low density, 63 in tube/high density, 99 in tube/low density).

Figure 7. Worker spatial fidelity (a, b) and occurrence (c, d) absolute zone sizes (in cm^2)

increased with number of workers (i.e. colony size and thus total nest area, since all colonies

were given nests of a size that was scaled to worker number). Nest shape (tube: red, circle: blue)

and worker density (high: a, c; low: b, d) did not have significant main effects, although density

interacted significantly with colony size. Only for occurrence zone size was there a significant

interaction between colony size and nest shape (difference in slope in c/d). Scatter plot points are

jittered by 5 x-axis units. Lines represent significant linear relationships (solid lines for circle

nest, dashed lines for tube nest; Table 2). N: 384 ants as in Fig. 6.

…But geometry drives the location of spatial fidelity zones relative to points in the nest

Nest shape reverses the relationship between worker spatial fidelity zone locations and important

points in the nest space. In the circle nest, workers who are close to the center of the brood pile

are far from the entrance and vice versa; in the tube nest on the other hand, the opposite is the

case: workers close to the entrance are also close to brood, and workers in the back of the nest

are far from both the entrance and the center of the brood (LME, $p<0.001$ for interaction of nest

shape and distance to the entrance on distance to the center of the brood, using the center of each

worker's spatial fidelity zone; Fig. 8, Table 3). Only a small portion of variation (3%, Table 3)

was explained by the random effect of colony identity.

Figure. 8. Worker spatial fidelity zones nearer the nest entrance are further away from the center

of the brood in the circle nest (b), in agreement with previous literature using similar nest shapes;

however, in tube-shaped nests (a), we find the opposite relationship: workers who are close to

the entrance are also close to brood. Overall distances are longer for the circle nest (as it does not

extend as far, see also Fig. 4). Worker density did not affect locations of spatial fidelity zones.

For statistics, see Table 3.

Linear Mixed Effects: worker scaled distance to brood center vs. scaled distance to nest entrance

	Mean To Brood				
Coeffcient	Est.	SE	df	T	P
Intercept	0.133	0.025	377	5.375	**<0.001**
MeanScaledDist	-0.332	0.132	377	-2.521	**0.012**
Nest	-0.079	0.030	377	-2.639	**0.009**
Density	0.015	0.013	377	1.151	0.250
MeanScaledDist:Nest	0.686	0.137	377	4.988	**<0.001**

Random Effects	
σ^2	0.00
τ_{00} Colony	0.00
N Colony	19
Observations	384
Marginal R^2 / Conditional R^2	0.604 / 0.634

Table 3. The relationship of worker spatial fidelity zone locations relative to the nest entrance

and relative to the center of the brood (see also Fig. 8).

Discussion

Available nest geometry influences the spatial organization of colonies in the rock-dwelling ant

Temnothorax rugatulus. In our circle-shaped nest, as in previous studies using similar 'open'

nest geometries, worker ants distributed themselves approximately evenly across the available

nest space, while brood and queens were clustered at an intermediate distance from the entrance,

and alates tended to be near the entrance. However, in the constricted space of the tube nest,

brood were more evenly distributed, all types of colony members were at further distances from

specific points (such as the nest entrance and the center of the brood), and workers, brood, and

queen locations were affected by heterogeneities of the nest geometry such as corners. The tube

geometry thus may have reduced global interaction, and may have prevented the clear distinction

between nest areas with respect to function that is apparent in the circle nest (e.g. a distinct brood

area separate from the space containing alates). Interestingly, overall worker density did not

affect these patterns. In contrast to these pervasive effects of nest geometry on distributions and

relative distances of colony members to each other, we found that individual worker spatial

fidelity zone sizes remained constant across nest geometries. That is, the area of the nest that is

used by each worker remains remarkably consistent as a proportion of total nest area across both

nest shapes and nest sizes (scaled with colony size). In absolute terms, both conservative spatial

fidelity zones (areas workers repeatedly visit) and general occurrence zones (all areas with

observations for a specific worker) were largely determined by total available nest area (scaled

with colony size, i.e. worker number), but did not change with nest geometry. One exception to

this is that we found a significant interaction of colony size and nest shape in their effect on

absolute occurrence zone (but not fidelity zone) sizes - this may indicate that at large nest and

colony sizes, workers in circle nests make occasional forays into more different sections of the

nests than workers in tube nests (even as their core spatial fidelity zones remain unchanged).

Does it matter how workers, brood, and queens are distributed across the space of a nest? Our

results show that both nest size and nest geometry can affect parameters such as the clustering of

colony features, the distances of individuals from key points, and heterogeneity in local density.

Previous work has shown that in *Temnothorax* ants, nest area plays a key role in nest site

selection (Pratt and Pierce 2001; Mitrus 2015), and workers actively gauge the size of

prospective nests (Pratt 2005; Pavlic et al., 2021). It has also been demonstrated that worker

density matters, as it can elevate worker energy expenditure (Cao and Dornhaus 2008) and lead

to increased foraging and scouting rates and induce polydomy (Cao 2013). Additionally, with

respect to geometry, certain structural features can alleviate consequences of high nest density

like traffic jams during nest evacuation (Burd et al., 2010; Shiwakoti et al., 2014; Wang and

Song 2016). While *Temnothorax* ants generally nest in preexising cavities, the size and geometry

of which may constrain them, they can make some modifications, both by adding internal

surrounding walls (Franks et al., 1992; Franks and Deneubourg 1997; Aleksiev et al., 2007a;

Aleksiev et al., 2007b; Aleksiev et al., 2007c; DiRienzo and Dornhaus 2017), and because

inactive colony members might act as temporary barriers, similar to how army ants produce

walls to regulate trail traffic (Baudier and Pavlic, 2020). The purpose of such modifications is

not yet clear, but it suggests that these ants might actively alter the available area or shape of

their nests. We also know in general that ants may also regulate contact rates as part of strategies

for task allocation and information exchange (Pacala et al., 1996; Pinter-Wollman et al., 2012;

Pinter-Wollman et al., 2013; Lehue et al., 2020b), and that contact rates can be used to estimate

colony or group sizes (Gordon 1992; Pratt 2005; Dornhaus and Franks 2006). Given that our

study suggests that nest geometry may impact local density and distribution of workers relative

to brood and queens, it is thus likely that both worker interaction rates and the interactions

between workers and brood, and thus colony function generally, are also impacted. Future

studies should closely examine how such interactions are affected by nest area and geometry, as

well as possible heterogeneities in local densities and relative positions of different colony

members.

While global spatial distributions and relationships across workers, brood, and queens were

significantly affected by nest size and geometry, we found that nest shape did not affect either

the size of workers' spatial fidelity zones (a median of ~4-8% of total nest area) nor the overall

area used by each worker, which we termed 'occurrence zones' (~25% of total nest area, i.e.

approximately 6x the size of the fidelity zones for our method and sample size). We found that

the absolute size of these zones (~0.1 - 4 cm^2 for fidelity zones) was primarily affected by the

total nest size, which was artificially scaled with colony size. We can thus not definitively

separate the effects of colony size (number of workers) and available nest area; however, since

density had no main significant effect, it is likely that it is simply total nest area that determines

spatial fidelity zone size. This is interesting since it indicates that workers somehow 'divide up'

the available nest area among them, but not simply by aiming for a particular worker density. It

is possible that workers move until achieving a constant density (interaction rate) over time.

While our study provides these clues, it does not fully demonstrate the mechanism used by

workers in determining either the placement or the size of their spatial fidelity zone.

The placement and size of worker spatial fidelity zones are thought to affect task allocation and

specialization across workers, by virtue of determining which task items workers are likely to

interact with, in *Temnothorax* (Franks and Tofts 1994; Leitner and Dornhaus 2019) and other

social insects (Sendova-Franks and Franks 1995; Jandt and Dornhaus 2009; Powell and

Tschinkel 1999). Given this potential importance of spatial fidelity zones for colony

organization, it is thus possible that the ants in our study were actively regulating spatial fidelity

zone size to match their needs independently of nest geometry. Such resilience could potentially

be an evolved strategy to adapt to varying nest conditions, a hypothesis supported by the

ecological diversity of *T. rugatulus* colonies across North America (Bengston and Dornhaus

2015) and the fact that in some populations, colonies face strong nest competition and thus

constraints on nest selection (Bengston and Dornhaus 2015). In contrast to previous findings in

the ant *Temnothorax unifasciatus* and the bumble bee *Bombus impatiens*, we did not find spatial

fidelity zone size to change systematically across the nest space (e.g. larger zones further from

the nest center, Sendova-Franks and Franks 1995; Jandt and Dornhaus 2009). However, our

observations are in line with earlier studies on *T. rugatulus*, which showed that although inactive

workers typically have small spatial fidelity zones and stay near the nest center, this pattern

changes during their active phases (Charbonneau et al., 2017b).

Our study addresses the general principle of how external, spatial configuration can affect the

behavior and function of adaptive systems. The importance of real space and physical constraints

on ant colonies specifically (Pinter-Wollman et al., 2017) and more widely on self-organized and

collective systems both engineered and biological (Araújo et al., 2023) is increasingly being

recognized, and has wide-ranging implications (Araújo et al., 2023; Pinter-Wollman et al., 2017;

Pinter-Wollman et al., 2018). Two processes are particularly important. First, confinement, i.e. in

this case physical barriers, can affect self-organized pattern formation; and beyond just forcing a

collective system to obey boundaries, can lead to entirely different outcomes with regard to

relationships and functionality, even in counterintuitive ways (Schäfer et al., 2022). Second,

stigmergy, i.e. the encoding of organizationally relevant information in the external environment,

is a key principle that allows efficient and robust organization in many natural systems. In this

study, we show that the brood clustering and the spatial separation of a dense brood area from

other nest areas (e.g. the area near the entrance and area containing alates) disappears when ants

are inhabiting a nest shape that, while offering the same total area, limits and defines traffic and

interactions across nest sections. This, first, demonstrates the role that confinement can play in

structuring or destroying self-organized patterns. Second, if, as hypothesized for these ants, task

specialization is organized by worker spatial fidelity zones overlapping with only specific task

stimuli, this stigmergic organization is upended by different nest shapes that prevent ants from

separating out different task areas. Both confinement and stigmergy thus drive important aspects

of self-organized function and we demonstrate here that they can be significantly impacted by

external spatial constraints.

References

Allaire, J. (2012). *RStudio: integrated development environment for R*. Boston, MA, 770(394),

165-171. URL: http://www.rstudio.com/

Aleksiev, A. S., Longdon, B., Christmas, M. J., Sendova-Franks, A. B., and Franks, N. R.

(2007a). Individual choice of building material for nest construction by worker ants and the

collective outcome for their colony. *Animal Behaviour*, *74*(3), 559-566.

https://doi.org/10.1016/j.anbehav.2006.12.019

Aleksiev, A. S., Sendova-Franks, A. B., and Franks, N. R. (2007b). Nest 'moulting' in the ant

Temnothorax albipennis. *Animal Behaviour*, *74*(3), 567-575.

https://doi.org/10.1016/j.anbehav.2006.12.023

Aleksiev, A. S., Sendova-Franks, A. B., and Franks, N. R. (2007c). The selection of building

material for wall construction by ants. *Animal Behaviour*, *73*(5), 779-788.

https://doi.org/10.1016/j.anbehav.2006.06.014

Araújo, N. A., Janssen, L. M., Barois, T., Boffetta, G., Cohen, I., Corbetta, A., ... and Volpe, G.

(2023). Steering self-organisation through confinement. *Soft matter*, *19*(9), 1695-1704.

https://doi.org/10.1039/D2SM01562E

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2014). Fitting linear mixed-effects models

using lme4. *arXiv* preprint arXiv:1406.5823. https://doi.org/10.18637/jss.v067.i01

Baudier, K. M., and Pavlic, T. P. (2020). Incidental interactions among Neotropical army-ant

colonies are met with self-organized walls of ants (Hymenoptera: Formicidae).

Myrmecological News, *30*. https://doi.org/10.25849/myrmecol.news_030:251

Bengston, S. E., and Dornhaus, A. (2014). Be meek or be bold? A colony-level behavioural

syndrome in ants. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1791),

20140518. https://doi.org/10.1098/rspb.2014.0518

Bengston, S. E., and Dornhaus, A. (2015). Latitudinal variation in behaviors linked to risk

tolerance is driven by nest-site competition and spatial distribution in the ant *Temnothorax*

rugatulus. *Behavioral Ecology and Sociobiology*, 69(8), 1265-1274.

https://doi.org/10.1007/s00265-015-1939-4

Bengston, S. E., Shin, M., and Dornhaus, A. (2017). Life-history strategy and behavioral type:

risk☐tolerance reflects growth rate and energy allocation in ant colonies. *Oikos*, 126(4), 556-

564. https://doi.org/10.1111/oik.03527

Burd, M., Shiwakoti, N., Sarvi, M., and Rose, G. (2010). Nest architecture and traffic flow: large

potential effects from small structural features. *Ecological Entomology*, *35*(4), 464-468.

https://doi.org/10.1111/j.1365-2311.2010.01202.x

Cao, T. T., and Dornhaus, A. (2008). Ants under crowded conditions consume more energy.

Biology Letters, *4*(6), 613-615. https://doi.org/10.1098/rsbl.2008.0381

Cao, T. T. (2013). High social density increases foraging and scouting rates and induces

polydomy in *Temnothorax ants*. *Behavioral Ecology and Sociobiology*, 67(11), 1799-1807.

https://doi.org/10.1007/s00265-013-1587-5

Charbonneau, D., and Dornhaus, A. (2015). Workers 'specialized' on inactivity: behavioral

consistency of inactive workers and their role in task allocation. *Behavioral Ecology and*

Sociobiology, *69*(9), 1459-1472. https://doi.org/10.1007/s00265-015-1958-1

Charbonneau, D., Hillis, N., and Dornhaus, A. (2015). 'Lazy' in nature: ant colony time budgets

show high 'inactivity' in the field as well as in the lab. *Insectes Sociaux*, 62(1), 31-35.

https://doi.org/10.1007/s00040-014-0370-6

Charbonneau, D., Poff, C., Nguyen, H., Shin, M. C., Kierstead, K., and Dornhaus, A. (2017a).

Who are the "lazy" ants? The function of inactivity in social insects and a possible role of

constraint: inactive ants are corpulent and may be young and/or selfish. *Integrative and*

Comparative Biology, 57(3), 649-667. https://doi.org/10.1093/icb/icx029

Charbonneau, D., Sasaki, T., and Dornhaus, A. (2017b). Who needs 'lazy' workers? Inactive

workers act as a 'reserve' labor force replacing active workers, but inactive workers are not

replaced when they are removed. *PLoS One*, 12(9), e0184074.

https://doi.org/10.1371/journal.pone.0184074

Chism. G. (2022). Gchism94/NestArchOrg: Research Compendium for DOI:

https://doi.org/10.1101/2022.06.30.498314 (v1.0.0). Zenodo.

https://doi.org/10.5281/zenodo.6828919

Choppin, M., Graf, S., Feldmeyer, B., Libbrecht, R., Menzel, F., and Foitzik, S. (2021). Queen

and worker phenotypic traits are associated with colony composition and environment in

Temnothorax rugatulus (Hymenoptera: Formicidae), an ant with alternative reproductive

strategies. *Myrmecological News*, *31*. https://doi.org/10.25849/myrmecol.news_031:061

Davidson, J. D., and Gordon, D. M. (2017). Spatial organization and interactions of harvester

ants during foraging activity. *Journal of The Royal Society Interface*, 14(135), 20170413.

https://doi.org/10.1098/rsif.2017.0413

DiRienzo, N., and Dornhaus, A. (2017). *Temnothorax rugatulus* ant colonies consistently vary in

nest structure across time and context. *PLoS One*, *12*(6), e0177598.

https://doi.org/10.1371/journal.pone.0177598

Dornhaus, A., Franks, N. R., Hawkins, R. M., and Shere, H. N. S. (2004). Ants move to improve:

colonies of *Leptothorax albipennis* emigrate whenever they find a superior nest site. *Animal*

Behaviour, 67(5), 959-963. https://doi.org/10.1016/j.anbehav.2003.09.004

Dornhaus, A., and Franks, N. R. (2006). Colony size affects collective decision-making in the

ant Temnothorax albipennis. *Insectes Sociaux*, 53(4), 420-427.

https://doi.org/10.1007/s00040-006-0887-4

Faulkes, C. G., and Bennett, N. C. (2001). Family values: group dynamics and social control of

reproduction in African mole-rats. *Trends in Ecology and Evolution*, 16(4), 184-190.

https://doi.org/10.1016/S0169-5347(01)02116-4

Foitzik, S., Backus, V. L., Trindl, A., and Herbers, J. M. (2004). Ecology of *Leptothorax* ants:

impact of food, nest sites, and social parasites. *Behavioral Ecology and Sociobiology*, *55*,

484-493. https://doi.org/10.1007/s00265-003-0718-9

Franks, N. R., Wilby, A., Silverman, B. W., and Tofts, C. (1992). Self-organizing nest

construction in ants: sophisticated building by blind bulldozing. *Animal Behaviour*, 44, 357-

375. https://doi.org/10.1016/0003-3472(92)90041-7

Franks, N. R., and Tofts, C. (1994). Foraging for work: how tasks allocate workers. *Animal*

Behaviour, *48*(2), 470-472. https://doi.org/10.1006/anbe.1994.1261

Franks, N. R., and Deneubourg, J. L. (1997). Self-organizing nest construction in ants: individual

worker behaviour and the nest's dynamics. *Animal Behaviour*, 54(4), 779-796.

https://doi.org/10.1016/0003-3472(92)90041-7

Franks, N. R., Pratt, S. C., Mallon, E. B., Britton, N. F., and Sumpter, D. J. (2002). Information

flow, opinion polling and collective intelligence in house–hunting social insects.

Philosophical Transactions of the Royal Society of London B: Biological Sciences,

357(1427), 1567-1583. https://doi.org/10.1098/rstb.2002.1066

Franks, N. R., Dornhaus, A., Best, C. S., and Jones, E. L. (2006). Decision making by small and

large house-hunting ant colonies: one size fits all. *Animal behaviour*, *72*(3), 611-616.

https://doi.org/10.1016/j.anbehav.2005.11.019

Gordon, D. M. (1992). Nest relocation in harvester ants. *Annals of the Entomological Society of*

America, 85(1), 44-47. https://doi.org/10.1093/aesa/85.1.44

Guo, X., Chen, J., Azizi, A., Fewell, J., and Kang, Y. (2020). Dynamics of social interactions, in

the flow of information and disease spreading in social insects colonies: effects of

environmental events and spatial heterogeneity. *Journal of Theoretical Biology*, *492*, 110191.

https://doi.org/10.1016/j.jtbi.2020.110191

Guo, X., Lin, M. R., Azizi, A., Saldyt, L. P., Kang, Y., Pavlic, T. P., and Fewell, J. H. (2022).

Decoding alarm signal propagation of seed-harvester ants using automated movement

tracking and supervised machine learning. *Proceedings of the Royal Society B*, *289*(1967),

20212176. https://doi.org/10.1098/rspb.2021.2176

Harris, W. V. (1956). Termite mound building. *Insectes Sociaux*, *3*(2), 261-268. doi:

https://doi.org/10.1007/BF02224306

Heyman, Y., Shental, N., Brandis, A., Hefetz, A., and Feinerman, O. (2017). Ants regulate

colony spatial organization using multiple chemical road-signs. *Nature Communications*,

8(1), 1-11. https://doi.org/10.1038/ncomms15414

Jandt, J. M., and Dornhaus, A. (2009). Spatial organization and division of labour in the

bumblebee *Bombus impatiens. Animal Behaviour,* 77(3), 641-651.

https://doi.org/10.1007/s00265-011-1244-9

Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. (2017). lmerTest package: tests in

linear mixed effects models. *Journal of Statistical Software*, 82(1), 1-26.

https://doi.org/10.18637/jss.v082.i13

Laidre, M. E., Wellborn, G. A., and Thiel, M. (2018). *Evolutionary ecology of burrow*

construction and social life (pp. 279-301). New York, NY: Oxford University Press.

Laidre, M. E. (2019). Architectural modification of shells by terrestrial hermit crabs alters social

dynamics in later generations. *Ecology*, 100(9), e02767. https://doi.org/10.1002/ecy.2767

Lehue, M., and Detrain, C. (2019). What's going on at the entrance? A characterisation of the

social interface in ant nests. *Behavioural Processes*, 160, 42-50.

https://doi.org/10.1016/j.beproc.2018.12.006

Lehue, M., and Detrain, C. (2020). Foraging through multiple nest holes: An impediment to

collective decision-making in ants. *PLoS One*, 15(7), e0234526.

https://doi.org/10.1371/journal.pone.0234526

Lehue, M., Collignon, B., and Detrain, C. (2020a). Multiple nest entrances alter foraging and

information transfer in ants. *Royal Society Open Science*, 7(2), 191330.

https://doi.org/10.1098/rsos.191330

Lehue, M., Detrain, C., and Collignon, B. (2020b). Nest entrances, spatial fidelity, and foraging

patterns in the red ant *Myrmica rubra*: a field and theoretical study. *Insects*, 11(5), 317.

https://doi.org/10.3390/insects11050317

Leitner, N., and Dornhaus, A. (2019). Dynamic task allocation: how and why do social insect

workers take on new tasks?. *Animal Behaviour*, 158, 47-63.

https://doi.org/10.1016/j.anbehav.2019.09.021

Lüdecke D (2023). _sjPlot: Data Visualization for Statistics in Social Science_. R package

version 2.8.14, <https://CRAN.R-project.org/package=sjPlot >.

Mitrus, S. (2015). The cavity-nest ant *Temnothorax crassispinus* prefers larger nests. *Insectes*

Sociaux, 62(1), 43-49. https://doi.org/10.1007/s00040-014-0372-4

Naug, D., and Smith, B. (2007). Experimentally induced change in infectious period affects

transmission dynamics in a social group. *Proc. R. Soc. B*, *274*, 61-65.

https://doi.org/10.1098/rspb.2006.3695

Negroni, M. A., Stoldt, M., Oster, M., Rupp, A. S., Feldmeyer, B., and Foitzik, S. (2021). Social

organization and the evolution of life-history traits in two queen morphs of the ant

Temnothorax rugatulus. *Journal of Experimental Biology*, *224*(7), jeb232793.

https://doi.org/10.1242/jeb.232793

Pacala, S. W., Gordon, D. M., and Godfray, H. C. J. (1996). Effects of social group size on

information transfer and task allocation. *Evolutionary Ecology*, 10(2), 127-165.

https://doi.org/10.1007/BF01241782

Pavlic, T. P., Hanson, J., Valentini, G., Walker, S. I., and Pratt, S. C. (2021). Quorum sensing

without deliberation: biological inspiration for externalizing computation to physical spaces

in multi-robot systems. *Swarm Intelligence*, *15*(1-2), 171-203.

https://doi.org/10.1007/s11721-021-00196-4

Pinter-Wollman, N., Hubler, J., Holley, J. A., Franks, N. R., and Dornhaus, A. (2012). How is

activity distributed among and within tasks in *Temnothorax* ants?. *Behavioral Ecology and*

Sociobiology, 66(10), 1407-1420.

Pinter-Wollman, N., Bala, A., Merrell, A., Queirolo, J., Stumpe, M. C., Holmes, S., and Gordon,

D. M. (2013). Harvester ants use interactions to regulate forager activation and availability.

Animal Behaviour, 86(1), 197-207. https://doi.org/10.1007/s00265-012-1396-2

Pinter-Wollman, N. (2015). Nest architecture architectures the collective behaviour of harvester

ants. *Biology Letters*, *11*(10), 20150695. https://doi.org/10.1098/rsbl.2015.0695

Pinter-Wollman, N., Fiore, S. M., and Theraulaz, G. (2017). The impact of architecture on

collective behaviour. *Nature Ecology & Evolution*, *1*(5), 0111.

https://doi.org/10.1038/s41559-017-0111

Pinter-Wollman, N., Jelić, A., and Wells, N. M. (2018). The impact of the built environment on

health behaviours and disease transmission in social systems. *Philosophical Transactions of*

the Royal Society B: Biological Sciences, *373*(1753), 20170245.

https://doi.org/10.1098/rstb.2017.0245

Powell, S. (2008). Ecological specialization and the evolution of a specialized caste in

Cephalotes ants. *Functional Ecology*, *22*(5), 902-911. https://doi.org/10.1111/j.1365-

2435.2008.01436.x

Powell, S., and Tschinkel, W. R. (1999). Ritualized conflict in *Odontomachus brunneus* and the

generation of interaction-based task allocation: a new organizational mechanism in ants.

Animal Behaviour, *58*(5), 965-972. https://doi.org/10.1006/anbe.1999.1238

Pratt, S. C., and Pierce, N. E. (2001). The cavity-dwelling ant *Leptothorax curvispinosus* uses

nest geometry to discriminate between potential homes. *Animal Behaviour*, 62(2), 281–287.

https://doi.org/10.1006/anbe.2001.1777

Pratt, S. C. (2005). Quorum sensing by encounter rates in the ant *Temnothorax albipennis*.

Behavioral Ecology, 16(2), 488-496. https://doi.org/10.1093/beheco/ari020

Prebus, M. (2017). Insights into the evolution, biogeography and natural history of the acorn

ants, genus *Temnothorax* Mayr (hymenoptera: Formicidae). *BMC Evolutionary Biology*,

17(1), 1-22. https://doi.org/10.1186/s12862-017-1095-8

Pusceddu, M., Cini, A., Alberti, S., Salaris, E., Theodorou, P., Floris, I., and Satta, A. (2021).

Honey bees increase social distancing when facing the ectoparasite *Varroa destructor*.

Science Advances, *7*(44), eabj1398. doi: https://doi.org/10.1126/sciadv.abj1398

R Core Team (2017). *R: A language and environment for statistical computing*. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Sasaki, T., and Pratt, S. C. (2013). Ants learn to rely on more informative attributes during

decision-making. *Biology Letters*, 9(6), 20130667. https://doi.org/10.1098/rsbl.2013.0667

Sasaki, T., Colling, B., Sonnenschein, A., Boggess, M. M., and Pratt, S. C. (2015). Flexibility of

collective decision making during house hunting in *Temnothorax* ants. *Behavioral Ecology*

and Sociobiology, *69*(5), 707-714. https://doi.org/10.1093/beheco/arr007

Schäfer, B., Pesch, T., Manik, D., Gollenstede, J., Lin, G., Beck, H. P., ... and Timme, M. (2022).

Understanding Braess' paradox in power grids. *Nature Communications*, *13*(1), 5396.

https://doi.org/10.1038/s41467-022-32917-6

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., ... and

Tinevez, J. Y. (2012). Fiji: an open-source platform for biological-image analysis. *Nature*

methods, 9(7), 676. https://doi.org/10.1038/nmeth.2019

Sendova-Franks, A. B., and Franks, N. R. (1994). Social resilience in individual worker ants and

its role in division of labour. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, 256(1347), 305-309. https://doi.org/10.1098/rspb.1994.0085

Sendova-Franks, A. B., and Franks, N. R. (1995). Spatial relationships within nests of the ant

Leptothorax unifasciatus (Latr.) and their implications for the division of labour. *Animal*

Behaviour, 50(1), 121-136. https://doi.org/10.1006/anbe.1995.0226

Sendova-Franks, A. B., and Franks, N. R. (1999). Self-assembly, self-organization and division

of labour. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*,

354(1388), 1395-1405. https://doi.org/10.1098/rstb.1999.0487

Sendova-Franks, A. B., and Van Lent, J. (2002). Random walk models of worker sorting in ant

colonies. *Journal of Theoretical Biology*, 217(2), 255-274.

https://doi.org/10.1006/jtbi.2002.3011

Shiwakoti, N., Sarvi, M., and Burd, M. (2014). Using non-human biological entities to

understand pedestrian crowd behaviour under emergency conditions. *Safety Science*, 66, 1-8.

https://doi.org/10.1016/j.ssci.2014.01.010

Stroeymeyt, N., Grasse, A. V., Crespi, A., Mersch, D. P., Cremer, S., and Keller, L. (2018).

Social network plasticity decreases disease transmission in a eusocial insect. *Science*,

362(6417), 941-945. doi: https://doi.org/10.1126/science.aat4793

Thomas, M. L. (2002). Nest site selection and longevity in the ponerine ant *Rhytidoponera*

metallica (Hymenoptera, Formicidae). *Insectes Sociaux*, 49(2), 147-152.

https://doi.org/10.1007/s00040-002-8294-y

Tofts, C., and Franks, N. R. (1992). Doing the right thing: ants, honeybees and naked mole-rats.

Trends in Ecology and Evolution, 7(10), 346-349. https://doi.org/10.1016/0169-

5347(92)90128-X

Tschinkel, W. R. (1999). Sociometry and sociogenesis of colonies of the harvester ant,

Pogonomyrmex badius: distribution of workers, brood and seeds within the nest in relation to

colony size and season. *Ecological Entomology*, 24(2), 222-237.

https://doi.org/10.1673/2006_06_32.1

Tschinkel, W. R. (2005). The nest architecture of the ant, *Camponotus socius*. *Journal of Insect*

Science, *5*(1). https://doi.org/10.1093/jis/5.1.9

Vaes, O., Perna, A., and Detrain, C. (2020). The effect of nest topology on spatial organization

and recruitment in the red ant *Myrmica rubra*. *The Science of Nature*, 107, 1-14.

https://doi.org/10.1007/s00114-020-01675-0

van Dijk, R. E., Covas, R., Doutrelant, C., Spottiswoode, C. N., and Hatchwell, B. J. (2015).

Fine-scale genetic structure reflects sex-specific dispersal strategies in a population of

sociable weavers (*Philetairus socius*). *Molecular Ecology*, 24(16), 4296-4311.

https://doi.org/10.1111/mec.13308

Varoudis, T., Swenson, A. G., Kirkton, S. D., and Waters, J. S. (2018). Exploring nest structures

of acorn dwelling ants with X-ray microtomography and surface-based three-dimensional

visibility graph analysis. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *373*(1753), 20170237. https://doi.org/10.1098/rstb.2017.0237

Visscher, P. K. (2007). Group decision making in nest-site selection among social insects.

Annual Review of Entomology, 52, 255-275.

https://doi.org/10.1146/annurev.ento.51.110104.151025

Wang, S., and Song, W. (2016). Experimental study of ant movement in a straight passageway

under stress conditions. *Journal of Insect Behavior*, *29*(6), 735-743.

https://doi.org/10.1007/s10905-016-9593-x

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D. A., François, R., ... and

Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4(43), 1686.

https://doi.org/10.21105/joss.01686

Wilensky, U. (1999). *NetLogo*. http://ccl.northwestern.edu/netlogo/. Center for Connected

Learning and Computer-Based Modeling, Northwestern University, Evanston, IL.

Wilson, E. O. (1992). The effects of complex social life on evolution and biodiversity. *Oikos*, 13-

18. https://doi.org/10.2307/3545511

Wilson, E. O., and Kinne, O. (1990). *Success and dominance in ecosystems: the case of the*

social insects (Vol. 2, pp. I-XXI). Oldendorf/Luhe: Ecology Institute.

Appendix

Table A1. Relationships between colony members scaled distance to the nest entrance and nest

shape and physical properties. Sex is male or female alates, and ratio is the proportion of male

alates in the observation. Formula: ScaledDist ~ Nest*Density + Day + (1 | Colony ID); Formula

(alates): Formula: ScaledDist ~ Nest + Day + Sex + SexRatio (1 | Colony ID)

Table A2. Relationships between mobile colony members scaled distance to the brood center and

nest shape and physical properties. The brood center is a colony's brood centroid in each

observation. Distance to the brood center is the absolute value of the difference between the

scaled distances of the brood center from each worker to the entrance and is scaled such that 1 is

the back of the tube nest. Formula: ToBrood~ Nest*Density + Day + (1 | Colony ID); Formula

(alates): Formula: ToBrood ~ Nest + Day + Sex + SexRatio (1 | Colony ID)

Table A3. The relationship between worker site fidelity and scaled distance to the brood center.

Scaled fidelity zone size is the summation of all zones that a worker was found in for at least

15% of its observations, and scaled occurrence zone size counts all zones with any observations

(and divide by the total of 24 possible, equal-area zones; worker must have at least 7

observations). True zone size (cm^2) multiplies this number by the internal area of the nest (note

that nest size was scaled to worker number and thus differs between colonies). Formula: Zone ~

Nest * ColonySize + Nest * MeanToBrood + Density * ColonySize + (1 | Colony)

Figure A1. The reference photo used to determine the nest area allocated to each worker in a

colony. The colony contains 248 individuals. The internal area is approximately $4.11mm^2$,

producing 0.017 mm^2 for each worker, which was doubled to 0.033 mm^2 to promote more

flexible space usage. This value represents the high nest density treatment and was doubled to

produce the low nest density treatment: 0.066 mm^2 for each worker in a colony.

Figure A2. The eight equal-area nest sections for the circle (a) and tube (b) nests (black lines)

used to determine the densities of colony members through the nest, and the twenty-four equal

area zones used to calculate worker site fidelity in the circle (c) and tube (d) nests. Two zone

sizes were used because the larger sections potentially capture tasks throughout the nest whereas

the smaller sections capture the smaller grain worker spatial fidelity zones.

Figure A3. Visual example of the shortest distance from an example worker (red dots) to an

example brood center (white dots) in the circle (a) and tube (b) nests. We found each colony

member's distance to the nest section closest to the entrance using reference coordinates for that

nest section, where we then added the shortest distance from that nest section to the entrance

through reference coordinates. In the circle nest, distance to the entrance = sqrt((colony member

x - entrance x)^2 + (colony member y - entrance y)^2), in the tube nest, where direct (i.e. straight-

line) path was often not possible within the available nest space, distance to the entrance =

sqrt((colony member x - nest section entrance x)^2 + (colony member y - nest section entrance

y)^2) + reference distance, where the nest section entrance is the closest accessible facing toward

the direction of the entrance (Fig. A3b). Correcting corner cuts at the entrance: Near the entrance

of both nests (in nest section 1), a colony member's distance to the nest entrance can cut through

the corner where the entrance tunnel opens into the nest (Fig. A3c). We solved this by first

determining whether a colony member's distance would cut through the corner and then

assigning an alternative distance to the entrance where necessary.

Subfigure (c): the criteria used to determine if a colony member required an alternative linear

distance to the nest entrance near the corner formed from the nest entrance tunnel opening into

the nest (black solid lines). The red dot represents a colony member, and the black dot represents

the entrance. The dashed lines represent the hypotenuses between the colony member and the

corner (red) and the corner and the entrance (black). The arcs represent angles, and the black

square represents 90°. The solid red polygon indicates nest space that would require this type of

alternative calculation of distance to the nest entrance.

Figure A4. When showing the actual distances of queens (a, b) and alates (c, d) from the nest

entrance (scaled only by colony size, by setting the farthest point of the tube nest as a distance of

1) instead of sections as in Fig. 3, it becomes clear how colony reproductive members in tube

nests (red) are spread out more from the entrance and from each other relative to the circle nest

(blue). This observation holds true regardless of high (a) and low (b) nest densities. Alate males

are also found to be closer to the nest entrance than queen alates (d). Sample sizes were 1178

queen observations across 20 colonies, and 1006 alate observations across 7 low density

treatment colonies.

Figure A5. Both queens and alates are found farther from the center of the brood pile (centroid)

in the tube nest (red) than what is possible in the circle nest (blue) (scaled only by colony size, by

setting the farthest point of the tube nest as 1). Differences in queen distributions are shown

across high (a) and low (b) nest densities; alates were only present in the low-density treatment.

Alate males are found to be farther from the brood centroid than young queens (female alates, d).

Sample sizes were 1080 queen observations from 400 photos across 20 colonies in each density

treatment, and 676 alate observations across 7 low density treatment colonies.

Linear Mixed Effects: colony member proportions within nest sections

Coeffcient	Workers					Brood					Queens					Alates				
	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P
Intercept	0.145	0.009	3181	16.566	**<0.001**	-0.127	0.013	3173	-9.489	**<0.001**	-0.161	0.031	3181	-5.157	**<0.001**	0.363	0.060	446	6.073	**<0.001**
NestSect	-0.006	0.004	3181	-1.438	0.151	0.090	0.006	3173	14.888	**<0.001**	0.118	0.014	3181	8.331	**<0.001**	-0.083	0.027	446	-3.028	**0.003**
NestSect2	0.000	0.000	3181	0.294	0.769	-0.006	0.001	3173	-9.596	**<0.001**	-0.010	0.001	3181	-6.479	**<0.001**	0.006	0.003	446	1.924	0.055
Nest	0.107	0.010	3181	10.486	**<0.001**	0.086	0.016	3173	5.487	**<0.001**	0.073	0.037	3181	2.003	**0.045**	-0.192	0.069	446	-2.787	**0.006**
Density	0.000	0.003	3181	0.000	1.000	0.000	0.004	3173	0.000	1.000	0.000	0.009	3181	0.000	1.000					
Day	0.000	0.000	3181	0.000	1.000	0.000	0.001	3173	0.000	1.000	0.000	0.001	3181	0.000	1.000	-0.000	0.003	446	-0.000	1.000
Corner	0.019	0.004	3181	5.502	**<0.001**	0.027	0.005	3173	5.019	**<0.001**	0.029	0.013	3181	2.261	**0.024**	-0.054	0.023	446	-2.340	**0.020**
NestSect:Nest	-0.040	0.006	3181	-7.358	**<0.001**	0.009	0.008	3173	1.016	0.310	0.001	0.020	3181	0.048	0.962	0.052	0.037	446	1.414	0.158
NestSect2:Nest	0.003	0.001	3181	4.316	**<0.001**	-0.005	0.001	3173	-5.974	**<0.001**	-0.004	0.002	3181	-1.701	0.089	-0.001	0.004	446	-0.150	0.881

Random Effects

	Workers	Brood	Queens	Alates
σ^2	0.01	0.01	0.07	0.03
τ_{00}	0.00 Colony	0.00 Colony	0.00 Colony	0.00 Colony
N	20 Colony	20 Colony	20 Colony	7 Colony
Observations	3192	3184	3192	456
Marginal R2 / Conditional R2	0.235 / 0.235	0.285 / 0.285	0.070 / 0.070	0.097 / 0.097

Linear Mixed Effects: individual worker site fidelity in each nest shape

Coeffcient	Scaled Fidelity Zone					Fidelity Zone (cm^2)					Scaled Occurrence Zone					Occurrence Zone (cm^2)				
	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P	Est.	SE	df	T	P
Intercept	1.772	0.401	374	4.413	**<0.001**	0.053	0.109	374	0.483	0.629	5.498	0.592	374	9.286	**<0.001**	-0.222	0.183	374	-1.214	0.225
Nest	-0.024	0.442	374	-0.055	0.957	-0.179	0.128	374	-1.395	0.164	0.608	0.644	374	0.944	0.346	0.343	0.205	374	1.675	0.095
ColonySize	-0.003	0.002	374	-1.201	0.231	0.002	0.001	374	3.093	**0.002**	0.000	0.004	374	0.132	0.895	0.009	0.001	374	8.128	**<0.001**
MeanScaledDist	0.980	1.592	374	0.616	0.539	-0.042	0.462	374	-0.092	0.927	2.488	2.322	374	1.072	0.285	1.167	0.737	374	1.584	0.114
Density	-0.489	0.373	374	-1.311	0.191	-0.030	0.088	374	-0.337	0.737	0.113	0.560	374	0.202	0.840	0.052	0.165	374	0.316	0.752
Nest:ColonySize	0.001	0.002	374	0.243	0.808	0.001	0.001	374	0.849	0.396	-0.004	0.003	374	-1.307	0.192	-0.002	0.001	374	-2.152	**0.032**
Nest:MeanScaledDist	-0.852	1.663	374	-0.512	0.609	0.240	0.483	374	0.497	0.619	-2.318	2.426	374	-0.955	0.340	-1.118	0.770	374	-1.452	0.147
ColonySize:Density	0.002	0.003	374	0.721	0.472	0.002	0.001	374	2.507	**0.013**	0.001	0.005	374	0.239	0.811	0.008	0.001	374	6.074	**<0.001**

Random Effects

σ^2	0.70					0.06					1.50					0.15				
τ_{00}	0.05 Colony					0.00 Colony					0.11 Colony					0.01 Colony				
N	19 Colony					19 Colony					19 Colony					19 Colony				
Observations	384					384					384					384				
Marginal R^2 / Conditional R^2	0.026 / 0.085					0.341 / 0.353					0.020 / 0.087					0.803 / 0.813				

Linear Mixed Effects: worker scaled distance to brood center vs. scaled distance to nest entrance

Coeffcient	Mean To Brood				
	Est.	SE	df	T	P
Intercept	0.133	0.025	377	5.375	**<0.001**
MeanScaledDist	-0.332	0.132	377	-2.521	**0.012**
Nest	-0.079	0.030	377	-2.639	**0.009**
Density	0.015	0.013	377	1.151	0.250
MeanScaledDist:Nest	0.686	0.137	377	4.988	**<0.001**

Random Effects

σ^2	0.00
$\tau_{00 \ Colony}$	0.00
$N_{\ Colony}$	19
Observations	384
Marginal R^2 / Conditional R^2	0.604 / 0.634