Nanopore-only assemblies for genomic surveillance of the global priority drug-

resistant pathogen, *Klebsiella pneumoniae*

Ebenezer Foster-Nyarko[1], Hugh Cottingham[2], Ryan R. Wick[2], Louise M. Judd[2], Margaret M.

C. Lam[2], Kelly L. Wyres[2], Thomas D. Stanton[1], Kara K. Tsang[1], Sophia David[3], David M.

Aanensen[3], Sylvain Brisse[4], Kathryn E. Holt[1, 2]

[1] Department of Infection Biology, London School of Hygiene and Tropical Medicine, London,

United Kingdom

[2] Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne,

VIC, 3004, Australia.

[3] Centre for Genomic Pathogen Surveillance, Big Data Institute, Li Ka Shing Centre for

Health Information and Discovery, Nuffield Department of Medicine, Oxford University,

Oxford OX3 7LF, United Kingdom

[4] Institut Pasteur, Université Paris Cité, Biodiversity and Epidemiology of Bacterial

Pathogens, Paris, France

Background

Oxford Nanopore Technologies (ONT) sequencing has rich potential for genomic

epidemiology and public health investigations of bacterial pathogens, particularly in low-

resource settings and at the point of care, due to its portability and affordability. However,

low base-call accuracy has limited the reliability of ONT data for critical tasks such as

antimicrobial resistance (AMR) and virulence gene detection and typing, serotype prediction

and cluster identification. Thus, Illumina sequencing remains the standard for genomic

surveillance despite higher capital and running costs.

Methods

We tested the accuracy of ONT-only assemblies for common applied bacterial genomics

tasks (genotyping and cluster detection, implemented via Kleborate, Kaptive and

Pathogenwatch), using data from 54 unique *Klebsiella pneumoniae* isolates. ONT reads

generated via MinION with R9.4 flowcells were basecalled using three alternative models

(Fast, High-accuracy (HAC) and Super-accuracy (SUP), available within ONT's Guppy

software), assembled with Flye and polished using Medaka. Accuracy of typing using ONT-

only assemblies was compared with that of Illumina-only and hybrid ONT+Illumina

assemblies, constructed from the same isolates as reference standards.

Results

The most resource-intensive ONT-assembly approach (SUP basecalling, with or without

Medaka polishing) performed best, yielding reliable capsule (K) type calls for all strains

(100% exact or best matching locus), reliable multi-locus sequence type (MLST) assignment

(98.3% exact match or single-locus variants), and good detection of acquired AMR genes

and mutations (88% – 100% correct identification across the various drug classes).

Distance-based trees generated from SUP+Medaka assemblies accurately reflected overall

genetic relationships between isolates; however, the definition of outbreak clusters from

ONT-only assemblies was problematic. HAC basecalling + Medaka polishing performed

similarly to SUP basecalling without polishing, and polishing introduced errors into HAC- or

Fast-basecalled assemblies. Therefore, we recommend investing compute resources into

basecalling (SUP model) over polishing, where compute resources and/or time are limiting.

Conclusions

Overall, our results show that MLST, K type and AMR determinants can be reliably identified

with ONT-only data. However, cluster detection remains challenging with this technology.

Keywords

Klebsiella pneumoniae, nanopore sequencing, basecalling, benchmarking, AMR, serotyping,

MLST, phylogenetic clustering, genomic surveillance, bacterial pathogens

Introduction

Klebsiella pneumoniae is a highly versatile species widely linked with multi-drug resistant (MDR) healthcare-associated infections and hypervirulent community-acquired infections [1, 2]. It contributes a third of all healthcare-associated Gram-negative infections globally and is a leading cause of Gram-negative invasive diseases, second to *Escherichia coli* [1-3]. Worryingly, the acquisition rate of antimicrobial resistance (AMR) determinants among *K. pneumoniae* has continued to rise rapidly over the last decades—particularly resistance to third-generation cephalosporins and carbapenems. The Global Research on AntiMicrobial resistance (GRAM) study estimated that *K. pneumoniae* was one of six pathogens that caused more than 250,000 deaths associated with AMR in 2019 [4]. In that study, *K. pneumoniae* was the leading Gram-negative pathogen in sub-Saharan Africa, contributing 20% of the deaths attributable to AMR [4]. At the same time, data aggregated from 151 studies encompassing 26 countries in sub-Saharan Africa indicate that *K. pneumoniae* is the top Gram-negative and second most important cause of neonatal sepsis [3]. *K. pneumoniae* infections are also characterised by a high case fatality rate (18-49%) [1, 3, 5, 6]. Consequently, *K. pneumoniae* has been flagged in the World Health Organisation's highest category of priority pathogens, for which the development of new antibiotics is urgently needed [7].

Accumulating genomic evidence has highlighted the emergence of hybrid MDR-hypervirulent clones, driven by the acquisition of mobile genetic elements [5, 6, 8-10]. This phenomenon represents a significant challenge to treating infections caused by *K. pneumoniae* and signals the urgent need for appropriate surveillance systems to identify MDR-hypervirulent clones and their genetic elements. WHO recommends the robust surveillance of AMR as a critical part of the Global Action Plan on AMR [11]. However, classical approaches to surveillance have relied on antimicrobial susceptibility testing and low resolution genotyping methods, which yield little information on the evolution and spread of AMR within bacterial populations.

AMR surveillance in critical pathogens such as *K. pneumoniae* has been dramatically

enhanced by the recent democratisation of whole-genome sequencing (WGS), primarily by

short-read (Illumina) sequencing. WGS is currently the gold standard approach for public

health pathogen surveillance, facilitating cluster identification and genotyping of clinically

relevant loci, including AMR, virulence traits, and antigen prediction [12-15]. Furthermore, as

few labs perform *K. pneumoniae* serology, prediction of capsular (K) and lipopolysaccharide

(O) antigen serotypes and their population distribution relies on inference from K and O

antigen biosynthesis loci captured in whole-genome sequence data [16, 17]. Genomic

surveillance for K and O types is of increasing interest, as these antigens are the main target

for alternative control measures such as vaccines, phage therapy, and monoclonal antibody

therapy [18-21]. Thirteen O antigen locus (OL) and 162 K antigen locus (KL) types are

described to date [16]. Unfortunately, the gold standard Illumina short-read sequencing

platforms result in fragmented genome assemblies, which cannot accurately resolve

plasmids and other mobile genetic elements that drive the dissemination of AMR

determinants via horizontal and lateral gene transfer. Fragmentation of Illumina genome

assemblies also complicates K and O typing, resulting in up to a third of Illumina-based

genomes being untypeable [22]. Long-read sequencing platforms, such as Pacific

Biosciences and Oxford Nanopore Technologies (ONT), can overcome these limitations, as

they yield longer reads (often exceeding 10 kbp, with current maximal read lengths reaching

1 Mbp) that are capable of resolving structural variations, long repeat regions, and genomic

copy-number alterations [23-25]. ONT is increasingly adopted to generate high-quality hybrid

Illumina-plus-long-read assemblies for bacteria, owing to its flexibility and low capital cost

compared to PacBio [26, 27]; however, high read-level error rates have precluded precluded

the adoption of ONT devices as a standalone platform for pathogen genomic epidemiology

studies or public health investigations [28]. The situation has shifted somewhat during the

COVID-19 pandemic as ONT has been widely adopted for virus sequencing and outbreak

analysis, including in settings where Illumina platforms are not available and are unlikely to

become so due to capital costs, reagent access, and other logistical issues [29]. There is,

therefore, now an unparalleled opportunity to harness ONT sequencing for genomic

epidemiology and surveillance of bacterial pathogens, which is particularly attractive for

AMR-associated pathogens such as *K. pneumoniae,* where resolving plasmids and other

highly variable genomic components is highly beneficial.

The current state-of-the-art ONT basecaller, Guppy, utilises the Fast, High-accuracy (HAC)

and Super-accuracy (SUP) models. The Fast model, as the name indicates, is the fastest of

the three algorithms but at the cost of accuracy. SUP is the most accurate and slowest; the

speed of the basecaller being a factor of the number of parameters in the neural network

model. HAC is intermediate between Fast and SUP and about six times slower than the Fast

model (https://github.com/rrwick/August-2019-consensus-accuracy-update#basecalling).

ONT polishing, utilising Medaka—a neural network-based tool that generates consensus

sequences by aligning the individual sequence reads against a draft assembly [30]—is

rapid, highly effective and believed to improve sequence accuracy significantly [31]. ONT

continuously refines these algorithms to improve raw reads and consensus accuracy [32].

However, despite continuous improvement over the years, ONT is still susceptible to non-

trivial numbers of base-call errors, which can impact the identification of clinically important

features such as acquired virulence and AMR genes [33, 34]. For example, a mean global

error rate of ~6% is reported for reads with quality scores of at least 10—corresponding to

94% raw read accuracy and >99.99% consensus accuracy with the primary nanopore in use

currently (R9.4.1 chemistry), although this varies by the organism [35] and is improving with

new chemistries such as R10 [36]. For a 5 Mbp genome, such as *K. pneumoniae,* we

previously reported roughly 3,000 errors (corresponding to at least 337 substitutions per

genome assembly) [37]; therefore, ONT-only assemblies are prone to erroneous SNP calls,

which can potentially hamper outbreak investigations [38]. The performance of ONT-only

assemblies in identifying AMR and virulence traits against the gold-standard (hybrid

assemblies) is unclear but is gaining attention [39, 40], and is pertinent to inform the

research community working with long-read data on what levels of accuracy to expect with

current basecalling algorithms.

Here, we use a collection of 54 clinical isolates of *K. pneumoniae* with matched Illumina and

ONT data [41] to assess the performance of ONT-only data when used for the prediction of

AMR determinants, virulence factors, K/O typing and phylogenetic clustering analysis.

Current read-based genotyping tools for Illumina data (such as ARIBA [42] or SRST2 [43])

are not optimised to be used efficiently on noisy long reads like those produced by ONT.

However, assembly-based tools such as Pathogenwatch [44], Kleborate [17, 45] and Kaptive

[16] can in principle be applied to any assembly regardless of the sequencing platform,

allowing for direct comparisons and facilitating the straightforward application of common

genotyping workflows to data generated from different sequencing instruments deployed for

pathogen surveillance. Furthermore, Pathogenwatch provides a free and accessible online

analysis platform, suitable for non-informatics specialists such as public health or clinical

microbiologists and epidemiologists, which, together with the growing network of ONT-

equipped labs could facilitate the widespread acceleration of genomic surveillance for AMR

pathogens. We therefore focused our investigation on the performance of these tools

(Pathogenwatch, Kleborate and Kaptive) for genomic surveillance of *K. pneumoniae* using

ONT-only assemblies.

Methods

Bacterial isolates and sequencing

We utilised 54 *K. pneumoniae* that were previously isolated at the Alfred Health Clinical

Microbiological Diagnostic Laboratory in Melbourne, Australia, as part of a year-long

prospective study of isolates from clinical infections (hospital-wide) and screening swabs (in

intensive care and geriatric wards) in the Alfred Health network [41, 46-48]. The accession

numbers for all genome data analysed in this study are provided in **Supplementary Table 1**.

Isolates were sequenced first via Illumina HiSeq 2500, generating 125 bp paired-end reads,

as described previously [46]. Later, selected isolates were subcultured, and fresh DNA

extracted for long-read sequencing using GenFind (Beckman Coulter) kits (full protocol

available under DOI: 10.17504/protocols.io.p5mdq46). ONT libraries were prepared using

the ligation protocol with barcoding kits to multiplex 12 to 24 isolates per run (barcode kits

EXP-NBD104 and EXP-NBD114) and sequenced on a MinION device with R9.4.1 flowcells

as previously described [26].

Basecalling and assembly

Fast5 files generated by the ONT MinION were initially basecalled with Guppy v4.0.14 [49]

on the Monash University M3 MASSIVE GPU cluster using the "Fast" (r9.4.1_450bps_fast)

and "High-accuracy", aka HAC (r9.4.1_450bps_hac) models for each sample. The "Super-

accuracy" (r9.4.1_450bps_sup) model was later used when it was released in Guppy v5.0.7

[49, 50]. The basecalled FASTQ files were then concatenated into a single file per

basecalled sequence run and demultiplexed into individual FASTQ files (one per sample)

with the qcat command-line tool v1.1.0 [51] based on the barcode sequences. As a quality

control step before assembly, we first filtered out poor-quality reads using fastp v0.20.1 [52]

(short reads) and Filtlong v0.2.1 [53] (long reads) at a sequence similarity threshold of 90-

95%, retaining only the reads with a minimum length of 1 kbp and excluding the worst 5-10%

of the reads (for FASTQ files less than 200 MB, a keep_percent of 95% was used, while a

keep_percent of 90% was applied to read files of size greater than 200 MB).

Three separate approaches were employed to generate genome assemblies (**Figure 1**).

First, Unicycler v0.4.9b [54] was used to create Illumina-only assemblies for all the study

isolates using default settings. When given short reads, Unicycler performs genome

assembly using SPAdes v3.14.0 [55] but includes extra fine-tuning steps such as filtering out

low-depth contigs and thus low-level contamination.

Next, we used Flye v2.8.1 [56] to generate ONT-only assemblies from basecalled reads

(three read sets per genome, called with the three different basecallers). Flye was chosen

based on our previous benchmarking of algorithms for ONT-only assemblies of bacterial

genomes [57]. Each ONT assembly was then polished with ONT reads using Medaka

(v1.4.3) models r941_min_fast_g303 and r941_min_high_g360 [30], resulting in two ONT-

only assemblies per basecalling model (polished vs unpolished); i.e., a total of six ONT-only

assemblies per sample.

Finally, we produced two kinds of hybrid Illumina+ONT assemblies: short-read-first and long-

read-first. Short-read-first hybrid assemblies were generated using Unicycler v0.4.9b, which

starts by building a short-read assembly graph with SPAdes v3.14.0, then uses the

corresponding long reads (HAC basecalled, in this case) to scaffold the genome, and finally

runs Pilon v1.23 [58] in an attempt to fill gaps, correct bases and fix misassemblies using the

short reads [26, 54] (HAC basecalled reads were used for the short-read-first assemblies as

these were generated before the SUP basecalling model became available [50]). To

generate long-read-first assemblies [59], we used Flye v2.8-1 to produce ONT-only

assemblies for the set of reads basecalled with the HAC and SUP-accuracy basecalling

models, followed by long-read polishing with Medaka [30] to repair any residual errors using

ONT long reads, then finally short-read polishing using Illumina reads and Pilon v1.24

(following the recommendations noted at [60]). Thus, altogether, we produced ten

assemblies per sample, encompassing reads derived from the three separate basecalling

models (**Figure 1**). QUAST v5.0.0 [de6973bb] [61] and CheckM v1.1.10 [62] were used to

assess the quality and completion of the hybrid assemblies (See **Supplementary File 1**,

available at https://figshare.com/s/6026855223031e769d8a, (DOI:

10.6084/m9.figshare.19745608).

We designate the ONT-only assemblies, with and without Medaka ONT-read polishing, as

"Fast", "Fast+Medaka", "HAC", "HAC+Medaka", "SUP", "SUP+Medaka". Based on the

recommendations of Wick et al. ([60] and [63]), we consider the long-read-first hybrid

assembly derived from SUP basecalled reads as the most accurate assembly and designate

this as the reference sequence, against which the performance of other assemblies is

benchmarked. We hereafter use the term "hybrid" to refer to the short-read-first (Unicycler)

hybrid assemblies, to distinguish them from long-read-first hybrid assemblies (otherwise

referred to as "SUP+Medaka+pilon" and "HAC+Medaka+pilon").

Genotyping

Genome assemblies were uploaded to Pathogenwatch v2.3.1 [44] where Kleborate v2.2 [45]

and Kaptive v2.0 [16] were automatically deployed to call multi-locus sequence types (STs)

using the 7-locus scheme [64], capsular polysaccharide (K) and lipopolysaccharide O locus

types and serotype predictions, acquired virulence traits including the siderophores

aerobactin (*iuc*), yersiniabactin (*ybt*) and salmochelin (*iro*), the genotoxin colibactin (*clb*) and

the hypermucoidy locus (*rmpADC*). Pathogenwatch also deploys Kleborate to identify

established AMR determinants (acquired genes and chromosomal mutations) [45] for the

following antimicrobial classes: aminoglycosides, carbapenems, third-generation

cephalosporins, third-generation cephalosporins plus β-lactamase inhibitors, colistin,

fluoroquinolones, fosfomycin, penicillins, penicillins + β-lactamase inhibitors, phenicols,

sulfonamides, tetracyclines, tigecycline and trimethoprim.

Clustering

We downloaded the Pathogenwatch pairwise distance matrix and corresponding neighbour-

joining tree for the full set of assemblies. The distance matrix is available at

https://figshare.com/s/6026855223031e769d8a (DOI: 10.6084/m9.figshare.19745608), and

the tree for interactive viewing at Microreact

(https://microreact.org/project/sUrpBsvXi1aiKD7ssPv9pu-nanopore-only-assemblies-for-

genomic-surveillance-of-klebsiella-pneumoniae). Pathogenwatch calculates pairwise SNP

distances between genomes based on a concatenated alignment of 1,972 genes

(2,172,367bp) that make up the core gene library for *K. pneumoniae* in Pathogenwatch and

infers a neighbour-joining tree from the resulting pairwise distance matrix [44]. Here, we

assessed the feasibility of identifying potential nosocomial transmission clusters using these

distance matrices. Several studies have proposed thresholds in the range of 21-25 genome-

wide SNPs for identifying nosocomial transmission clusters of *K. pneumoniae* [65-67].

However, as Pathogenwatch calls SNPs only in 1,972 core genes and not genome-wide, we

compared the SNP distances calculated by Pathogenwatch with genome-wide SNP counts

obtained by mapping short reads to a reference genome to determine the equivalent cut-off

for clustering analysis using Pathogenwatch distances. To do this, we used the genome-

wide SNP alignment generated previously for n=270 *K. pneumoniae* isolated at Alfred

Health, based on mapping of Illumina reads to the *K. pneumoniae* NTUH-K2044 reference

genome using the RedDog pipeline [68] (see full details in [41]). Pairwise SNP counts were

extracted using snp-dist [69]. Assemblies for these 270 genomes (assembled from Illumina

reads *de novo* using SPAdes optimised with Unicycler v0.4.74, see full details in [41]) were

uploaded to Pathogenwatch, and the pairwise distance matrix was downloaded and

compared against that generated from RedDog. We then used R to fit a linear regression

model for Pathogenwatch distances as a function of genome-wide mapping-based SNP

distances (see **Supplementary Figure 1**). This indicated a Pathogenwatch distance

threshold of 10 SNPs would be approximately equivalent to the established genome-wide

distance threshold of 25. These thresholds assume accurate basecalling from Illumina data.

To ascertain a corresponding threshold distance using ONT-only data, we compared

pairwise Pathogenwatch distances calculated using ONT-only (SUP+Medaka) assemblies

vs Illumina assemblies, for pairs of strains linked via probable transmission clusters. Using R

to fit a linear regression model indicated an ONT-only Pathogenwatch distance of 50 SNPs

283 would approximate the Illumina-based Pathogenwatch distance of 10 SNPs or genome-wide

284 distance of 25 SNPs (see **Figure 6**).

286 We compared the topologies of neighbour-joining trees generated from Pathogenwatch

287 distance matrices calculated using SUP+Medaka, HAC+Medaka or Fast+Medaka

288 assemblies against the reference tree (calculated from hybrid SUP+Medaka+pilon

289 assemblies), using the tanglegram function in the R package dendextend v1.15.2 to

290 generate comparative tree plots and calculate entanglement coefficients. We also used the

291 phytools package v1.0-3 in R to compute the Robinson-Foulds distance [70, 71] between

292 tree topologies, which represents a sum of the number of partitions inferred by the first tree

293 but not the second tree and that inferred by the second tree but not the first tree.

295 ***Data analysis and visualisation***

296 Data were analysed and visualised using R v4.1.0. Linear regression was done using the lm

297 function in base R, forcing the line through the origin. Genotyping and clustering results were

298 visualised and compared using R packages ape v5.5 [72], cowplot v1.1.1 [73], data.table

299 v1.14.2 [74], DECIPHER v2.20.0 [75], dendextend v1.15.2 [76], dplyr v1.0.7 [77], ggtree

300 v3.0.4 [78], ggpubr v0.4.0 [79], gridExtra v2.3 [80], janitor 2.1.0 [81], kableExtra v1.3.4 [82],

301 phytools v1.0.3 [71], tidyverse v1.3.1 [83], treeio v1.16.2 [84], treemap v2.4-3 [85] and stringr

302 v1.4.0 [86].

304 **Results and Discussion**

305 We investigated the accuracy of ONT-only assemblies for identifying ST, K and O antigen

306 loci, AMR determinants, plasmid- and ICE*Kp*-borne virulence factors and phylogenetic

307 clusters, compared with assemblies generated via Illumina or hybrid methods that combine

308 Illumina and ONT reads (**Figure 1**). We used a diverse set of 54 unique *K. pneumoniae*

309 isolates (see **Methods**), spanning thirty STs and displaying a wide range of K types, O

310 types, AMR and virulence profiles (see **Supplementary Table 1**). The genome sizes were in

the range 5,075,945 bp – 6,163,371 bp, and spanned G+C content of 49.6% to 57.7%

(**Supplementary Figure 2**). Notably, the choice of ONT basecalling model did not appear to

have a significant impact on the N50 or G+C content. Full details of sequence quality and

assembly metrics are presented in **Supplementary Table 1** and **Supplementary File 1**.

MLST and clustering analysis

Perfect ST calls require exact sequence matches at all 7 MLST loci; however, ST is often

treated as a proxy for lineage, which can sometimes be correctly identified based on single-

locus or double-locus variants of the correct ST (i.e., where 6/7 or 5/7 of the MLST loci have

exact matches, these are reported as STx-1LV or STx-2LV respectively by Kleborate). The

Illumina-only and SUP+Illumina hybrid assemblies gave identical MLST results for all

genomes. Compared to this gold-standard result, for ONT-only assemblies, the proportion of

correct ST calls was 87.3% for SUP basecalled assemblies with or without polishing; 78.2%

for HAC+Medaka; and 32.7% for non-polished HAC (see **Figure 2A**). Fast basecalled

assemblies showed poor results (20% correct with polishing, 0 without). ONT-only

assemblies based on SUP basecalled reads reliably identified lineage in all cases when

allowing for single- or double-locus variant calls. Assemblies based on HAC basecalling also

performed well, with 96.3% (HAC+Medaka) and 90.7% (HAC) matching the true ST within 0-

2 locus variants, respectively. Interestingly, for isolate KSB1_10J, the SUP+Medaka

assembly yielded two loci mismatches with the expected ST323, while the SUP assembly

without polishing yielded a single mismatch (i.e., a more accurate result).

Pathogenwatch calculates pairwise SNP distances (hereafter referred to as PW distances)

across a set of 1,972 *K. pneumoniae* core genes and constructs a neighbour-joining tree

based on these distances. **Figure 3** shows a PW-distance tree for all 540 assemblies

analysed here. PW distances between ONT-only assemblies vs their corresponding hybrid

reference ranged from 2–882 SNPs (median 17) for SUP+Medaka to 198–3390 SNPs

(median 374) for unpolished Fast basecalled assemblies (**Figure 4**). Accordingly, the tree

shows clear clustering of alternative assemblies for the same isolate, with SUP and HAC

assemblies clustering closely with their corresponding hybrid references and Fast basecalled

assemblies being distant relatives (see **Figures 3-4,** and interactive version of the tree in

Microreact, [https://microreact.org/project/5chGLxaT1eVHKrThyc4b4J-nanopore-only-

assemblies-for-genomic-surveillance-of-the-global-priority-drug-resistant-pathogen-

klebsiella-pneumoniae]).

We then constructed assembly method-specific neighbour-joining trees for the set of 54

isolates, using PW distance matrices constructed from either reference assemblies, or SUP,

HAC or Fast basecalled and polished ONT-only assemblies. The trees based on SUP or

HAC ONT-only assemblies yielded generally similar topologies to the reference tree

(entanglement coefficients 0.31 and 0.35, Robinson-Foulds distances 20 and 22,

respectively), but the Fast basecalled ONT-only tree was more divergent (entanglement

coefficient 0.96, Robinson-Foulds distance 102; see **Figure 5** and **Supplementary Figures**

3-4). However, clustering by ST was evident in all trees regardless of basecalling and

assembly method.

Pairwise SNP distances can also be used directly, in the absence of trees, as a rule-in/rule-

out indicator for potential transmission clusters. A consensus has emerged recently that a

pairwise distance exceeding a threshold of 21–25 genome-wide SNPs is a reasonable basis

for ruling out recent transmission of *K. pneumoniae* [65-67]. Based on comparisons using

Illumina assemblies for n=270 unique *K. pneumoniae* clinical isolates, we estimate this

equates to a PW distance of 10 SNPs (see **Methods** and **Supplementary Figure 1**). **Figure**

6 shows PW distances calculated using reference assemblies, or SUP+Medaka ONT-only

assemblies, for four groups of isolates in our data set that have previously been identified as

part of nosocomial transmission clusters (based on a combined analysis of patient

movement data and Illumina sequence, using genome-wide SNPs called from read mapping

to an in-cluster reference sequence) [46, 87]. Clusters of ST29, ST231 and ST491 had

pairwise PW distances 0-2 using reference genomes, but much higher PW distances using

ONT-only assemblies (ranges 6-10, 4-4, and 6-20, respectively using SUP+Medaka; 13-23,

10-16, 19-28, respectively using SUP without polishing). Cluster ST323 was more diverse,

with PW distances of 3-34 between reference genomes and 58-659 using ONT-only

(excluding one outlier strain, KSB1_10J, the distances were 3-23 for reference assemblies,

58-118 for SUP+Medaka and 83-183 for SUP). Linear regression of polished ONT-only PW

distances vs Illumina PW distances yielded a slope of ~5 (adjusted r^2=0.89, see **Figure 6**),

suggesting a relaxed PW distance threshold of ~50 could be suitable for calling transmission

clusters using ONT-only (SUP+Medaka) data, although the relationship does seem to vary

by ST (see **Figure 6**). We therefore conclude Pathogenwatch analyses of ONT-only

assemblies are quite reliable for identification of lineage or ST (as described above), and

thus could be reliably used to 'rule out' isolates of distinct lineages from suspected

transmission clusters. However, more work is needed to establish suitable analytical

methods for calculating and interpreting smaller distances within lineages, as is required to

differentiate recent transmission chains from coincidental but independent infections with the

same lineage.

K and O locus typing

In *K. pneumoniae*, K and O locus types vary in their gene content, and typing via the Kaptive

software is achieved by searching genomes for full-length reference K or O loci and

confirming the presence of the expected genes within the best-matching locus [16, 17, 88]. K

locus (KL) and O locus (OL) calls were in perfect agreement between Illumina and hybrid

assemblies. Compared to these results as reference, all ONT-only assemblies correctly

identified the best-matching KL type. However, the confidence reported by the Kaptive typing

tool—which depends on the detection of intact genes (i.e., open reading frames) in the K

locus and is thus expected to be impacted by basecall accuracy—varied widely for the

different assemblies (see **Figure 2C**). The distribution of missing gene count, stratified by

confidence, is shown in **Figure 2E**; this supports the expectation that the lack of confidence

in KL calls is due to disruption of open reading frames in the locus, presumably due to

basecall errors that result in frameshifts or stop codons in the expected coding genes.

Figure 2G shows that nucleotide similarity (compared to KL reference sequences) along the

K locus was quite high for all assemblies, with nucleotide divergence well below the Kaptive

threshold for a confident call (≤5% divergence). This confirms that even small numbers of

basecall errors are enough to disrupt open reading frames and thus reduce confidence in

genotype calls.

The results were similar for O typing, although some incorrect OL calls were made with the

Fast assemblies (1.85% each with Fast and Fast+Medaka; red colours in **Figure 2D**). A

correct O1 prediction requires the detection of an extra gene (*wbbY*) outside the O locus [17,

89]; if this open reading frame is disrupted, O1 may be miscalled as O2. Unsurprisingly, the

discrepancies with the Fast OL calls arose from O1 being erroneously reported as O2, due

to frameshift mutations arising from basecalling/assembly errors affecting the unlinked *wbbY*

gene.

AMR determinants

Kleborate screens for acquired AMR genes by interrogating each input assembly against a

species-aware modified version of the Comprehensive Antibiotic Resistance Database [45].

A total of 270 acquired genes were identified across the 54 reference genome assemblies

(median 9 per genome, range 0-18). These acquired genes were mostly identified correctly

from SUP+Medaka (95.8%), SUP (84.8%), HAC+Medaka (82.7%) and HAC (60.3%)

assemblies, with lower recovery rates from Fast+Medaka (53.2%) and Fast (25.7%)

assemblies. Kleborate also screens for mutations in chromosomal core genes known to be

associated with AMR [45]. Here, reportable mutations were identified from reference

assemblies in the *gyrA* and *parC* genes (substitutions associated with fluoroquinolone

resistance, n=9 isolates), in *bla*SHV (substitutions associated with extended spectrum β-

lactamase activity, n=27 genomes) and in *ompK35* and *ompK36* (truncations associated with

reduced susceptibility to β-lactams, n=2 genomes). Truncations in *pmrB* and *mgrB*

(associated with colistin resistance) are screened by Kleborate, but these genes were intact

in all reference assemblies. The substitutions in *gyrA*, *parC* and *bla*SHV were recovered

quite accurately from the SUP+Medaka (91.7%) and SUP (88.9%) assemblies, but less so

using HAC+Medaka (83.3%), HAC (75%), Fast+Medaka (55.6%) and Fast (44.4%)

assemblies (see **Figure 7a-b**). The two *omp* gene truncations were identified by all

assembly methods, however, truncations of *omp*, *pmrB* and *mgrB* were also falsely identified

in many cases (7.4% with SUP+Medaka, 7.4% SUP, 14.8% HAC+Medaka, 9.3% HAC,

77.8% Fast+Medaka, 27.8% Fast; see **Figure 7c-d**).

As the presence of AMR determinants in genomes is often used to predict or explain

resistance to clinically relevant drugs, we also considered the impact of ONT basecalling and

assembly method on the task of identifying which isolates are likely non-susceptible to

specific drug classes. **Table 1** shows the proportion of genomes that would be correctly

classified as non-susceptible due to the detection of AMR determinants known to be present

in the reference genome. SUP+Medaka assemblies performed well, with 88% – 100% of

genomes with AMR determinants correctly identified as such across the various drug

classes. HAC+Medaka also showed good detection rates (60% - 100%), while SUP results

were more variable, ranging from 42-100% detection rate (see **Table 1**).

Acquired virulence

Kleborate reports the presence of five major acquired virulence loci, including the ICE*Kp*-

encoded siderophore yersiniabactin (*ybt*) and genotoxin colibactin (*clb*), plus three loci that

are typically plasmid-borne and commonly linked with invasive infections caused by

hypervirulent *K. pneumoniae*, namely, siderophores aerobactin (*iuc*) and salmochelin (*iro*),

and the hypermucoidy locus *rmpADC.* Each of these loci comprises several genes, which

are included in locus-specific MLST schemes for each [45, 90, 91]. Each locus will be

reported, and an ST (or the closest match) assigned following the same logic to assign STs

as for seven-gene MLST, if >50% of the genes contained in the locus are detected.

Lineages of each virulence locus are inferred based on virulence STs, with each lineage pre-

assigned to a set of virulence STs [90, 91]. Acquired virulence genes detected in our

reference genomes include *ybt*, *clb,* and *iuc. Iuc* and *clb* were present in only two isolates

each (*iuc* in INF079 and INF151, *clb* in INF079 and INF341), while *ybt* was detected in

twenty isolates (36.4%). When present, the Illumina and hybrid assemblies correctly

detected both *iuc* and *clb,* including correct lineage assignment. For the ONT-only

assemblies, SUP (±Medaka) and HAC+Medaka correctly reported the *iuc* and *clb* STs and

lineages, while the HAC and Fast (±Medaka) detected the loci as present but failed to

identify a ST match and thus could not identify a lineage. Similarly, for *ybt*, the Illumina and

hybrid assemblies correctly reported the presence or absence of the *ybt* locus, and

accurately predicted the lineage when the locus was present (100% each). However, for

ONT-only assemblies, only the SUP (±Medaka) matched the Illumina and hybrid in uniformly

detecting *ybt* and assigning the correct lineage (**Figure 2B**). The other ONT-only assemblies

all reported the presence or absence of *ybt* but failed to call the ST and thus to report a

lineage (HAC+Medaka, 7.4%; HAC, 20.4%; Fast+Medaka, 31.5%, Fast, 35.2%). Notably,

there were no false positives (i.e., reporting the locus present when absent), nor any false

negatives (i.e., failing to detect presence of the locus when present).

Limitations and future directions

An artefact of our sample collection is that there were few virulence plasmid-positive strains,

so we could not assess other virulence genes apart from yersiniabactin. However, it is clear

from the available results that genotyping based on detection of gene presence/absence is

generally quite reliable using assemblies inferred from SUP-basecalled ONT reads, hence

virulence gene detection is expected to be well tolerated generally. There were also very few

strains in our collection with genuine AMR-associated truncations, so the call rate for these

could not be accurately established. However, clearly, basecall errors frequently disrupt

open reading frames, such that AMR-related truncations are subject to relatively high false-

positive call rates. The same issue also affects the confidence of K and O locus typing:

whilst the correct locus is almost always detected, the reported confidence of the call

depends on the detection of open reading frames for all genes in the locus, which is rarely

achieved using ONT data alone.

Our study considered *K. pneumoniae* only. However, the basecalling, assembly and

genotyping tasks explored here are commonly applied across other bacterial pathogens. In

particular, since *Enterobacterales* share common methylation patterns (which is a major

determinant of basecalling accuracy) and AMR mechanisms, the results presented here can

be considered informative as to the general accuracy of ONT-only analysis of the family

more broadly.

Notably, laboratory and informatics methods for ONT sequencing are under constant

development, e.g., new flowcells have recently been released with new (R10) pores;

basecalling models can be trained on a larger or species-specific dataset, and assembly and

polishing methods are under active development [92, 93]. These developments are expected

to lead to improvements in the accuracy of ONT-only assemblies and therefore the accuracy

of genotyping from such assemblies [36]. Therefore, our results should be considered a

baseline from which improvements are expected to accrue. In addition, it is likely that the

task of pairwise SNP calling could be more accurately performed using read-based variant

calling approaches, rather than the assembly-based methods used here. However, there is

currently a lack of SNP-calling tools designed specifically for bacterial ONT data (notably

those trained on human data do not perform well on bacterial data due to differences in

methylation [37]), so exploring this was beyond the scope of the current study which focused

on the use of a consistent assembly-based approach to all analysis tasks (implemented in

the freely accessible online Pathogenwatch platform).

Despite these limitations, our study provides a timely update on the level of accuracy that

can be achieved with currently well-established and widely available protocols, including (i)

R9.4 pores (which have been widely and stably used for several years now); (ii) out-of-the-

box basecalling with Guppy, which can be done in real-time, and on-board with devices like

Mk1C or GridION; (iii) a simple, rapid low-resource assembly with Flye (assessed in several

benchmarking studies as the best singular assembler for ONT-only); and (iv) the well-

established and stable Medaka polisher.

Conclusions

Overall, our results show that MLST, K/O locus type, virulence and AMR determinants can

be reliably identified from ONT-only genome assemblies. However, pairwise SNP distance

estimation was less reliable and thus we propose that ONT-only analysis should be

considered reliable only for rule-out of potential clusters using MLST-based lineage

assignments, rather than being the sole trigger for specific actions that are usually based on

a high suspicion of transmission (e.g., enhanced containment procedures in hospital

settings), as it currently lacks the sensitivity needed for public health or infection control

investigations. Where compute resources and/or time are limiting, our data indicate that

compute resources are best directed towards basecalling (SUP model) over polishing.

Funding information

This work was supported by the Bill and Melinda Gates Foundation, Seattle (OPP1175797,

KlebNet Project to DMA, KEH, SB). The funding bodies had no role in study design or in

data collection, analysis and interpretation.

Author contributions

Conceptualisation: KEH. Methodology and Software: KEH, KLW, MMCL, RRW. Formal

Analysis and Visualisation: EFN, HC, KLW, MMCL, KEH. Data curation: KEH, HC.

Supervision: KEH. Funding: KEH. Writing – Original Draft Preparation: EFN, KEH. All

authors read and approved the final manuscript.

Conflicts of interest

The authors declare they have no conflicts of interest.

Ethical statement

Ethical approval for the collection and sequencing of clinical isolates was granted by the

Alfred Hospital Ethics Committee, Melbourne, Australia (Project numbers #550/12 and

#526/13).

References

1.   Meatherall, B.L., et al., *Incidence, risk factors, and outcomes of Klebsiella*

pneumoniae bacteremia. Am J Med, 2009. **122**(9): p. 866-73.

2.   Anderson, D.J., et al., *Bloodstream infections in community hospitals in the 21st*

century: a multicenter cohort study. PLoS One, 2014. **9**(3): p. e91713.

3.   Vading, M., et al., *Invasive infection caused by Klebsiella pneumoniae is a disease*

affecting patients with high comorbidity and associated with high long-term mortality.

PLoS One, 2018. **13**(4): p. e0195258.

4.   Antimicrobial Resistance Collaborators, *Global burden of bacterial antimicrobial*

resistance in 2019: a systematic analysis. Lancet, 2022. **399**(10325): p. 629-655.

5.   Jung, Y., et al., *Differences in characteristics between healthcare-associated and*

community-acquired infection in community-onset Klebsiella pneumoniae

bloodstream infection in Korea. BMC Infect Dis, 2012. **12**: p. 239.

6.   Giske, C.G., et al., *Clinical and economic impact of common multidrug-resistant*

gram-negative bacilli. Antimicrob Agents Chemother, 2008. **52**(3): p. 813-21.

7.   World Health Organisation, *Prioritization of pathogens to guide discovery, research*

and development of new antibiotics for drug-resistant bacterial infections, including

tuberculosis. 2017.

8.   Navon-Venezia, S., et al., *Klebsiella pneumoniae: a major worldwide source and*

shuttle for antibiotic resistance. FEMS Microbiol Rev, 2017. **41**(3): p. 252-275.

9.   Temkin, E., et al., *Estimating the number of infections caused by antibiotic-resistant*

Escherichia coli and Klebsiella pneumoniae in 2014: a modelling study. Lancet Glob

Health, 2018. **6**(9): p. e969-e979.

10.   European Centre for Disease Prevention and Control, *Antimicrobial resistance*

(EARS-Net). ECDC. Annual epidemiological report for 2014, 2018.

11.   World Health Organisation, *Antimicrobial resistance. Draft global action plan on*

antimicrobial resistance. Geneva, 2015. Available from:

https://apps.who.int/iris/handle/10665/193736. Accessed 10 November 2021.

12.   Cassini, A., et al., *Attributable deaths and disability-adjusted life-years caused by*

infections with antibiotic-resistant bacteria in the EU and the European economic

area in 2015: a population-level modelling analysis. Lancet Infect Dis, 2019. **19**(1): p.

56-66.

13.   Nagaraj, G., et al., *High-resolution genomic profiling of carbapenem-resistant*

Klebsiella pneumoniae isolates: A multicentric retrospective indian study. Clin Infect

Dis, 2021. **73**(Supplement_4): p. S300-S307.

14.   Saavedra, S.Y., et al., *Complexity of genomic epidemiology of carbapenem-resistant*

Klebsiella pneumoniae isolates in Colombia urges the reinforcement of whole

genome sequencing-based surveillance programs. Clin Infect Dis, 2021.

73(Supplement_4): p. S290-S299.

15.   Aanensen, D.M., et al., *Implementing whole-genome sequencing for ongoing*

surveillance of antimicrobial resistance: Exemplifying insights into Klebsiella

pneumoniae. Clin Infect Dis, 2021. **73**(Suppl_4): p. S255-S257.

16.   Lam, M.M.C., et al., *Kaptive 2.0: updated capsule and lipopolysaccharide locus typing for the.* Microb Genom, 2022. **8**(3).

17.   Wyres, K.L., et al., *Identification of Klebsiella capsule synthesis loci from whole genome data.* Microb Genom, 2016. **2**(12): p. e000102.

18.   de Sousa, J.A.M., et al., *Modular prophage interactions driven by capsule serotype select for capsule loss under phage predation.* ISME J, 2020. **14**(12): p. 2980-2996.

19.   Feldman, M.F., et al., *A promising bioconjugate vaccine against hypervirulent.* Proc Natl Acad Sci U S A, 2019. **116**(37): p. 18655-18663.

20.   Ravinder, M., et al., *A Synthetic carbohydrate-protein conjugate vaccine candidate against Klebsiella pneumoniae serotype K2.* J Org Chem, 2020. **85**(24): p. 15964-15997.

21.   Campbell, W.N., et al., *Immunogenicity of a 24-valent Klebsiella capsular polysaccharide vaccine and an eight-valent Pseudomonas O-polysaccharide conjugate vaccine administered to victims of acute trauma.* Clin Infect Dis, 1996. **23**(1): p. 179-81.

22.   Wyres, K.L., et al., *Genomic surveillance for hypervirulence and multi-drug resistance in invasive Klebsiella pneumoniae from South and Southeast Asia.* Genome Med, 2020. **12**(1): p. 11.

23.   Ashton, P.M., et al., *MinION nanopore sequencing identifies the position and structure of a bacterial antibiotic resistance island.* Nat Biotechnol, 2015. **33**(3): p. 296-300.

24.   Loman, N.J. and M.J. Pallen, *Twenty years of bacterial genome sequencing.* Nat Rev Microbiol, 2015. **13**(12): p. 787-94.

25.   Arredondo-Alonso, S., et al., *On the (im)possibility of reconstructing plasmids from whole-genome short-read sequencing data.* Microb Genom, 2017. **3**(10): p. e000128.

26.   Wick, R.R., et al., *Completing bacterial genome assemblies with multiplex MinION sequencing.* Microb Genom, 2017. **3**(10): p. e000132.

613 27. De Maio, N., et al., *Comparison of long-read sequencing technologies in the hybrid*

614   *assembly of complex bacterial genomes.* Microb Genom, 2019. **5**(9).

615 28. Ben Khedher, M., et al., *Application and challenge of 3rd generation sequencing for*

616   *clinical bacterial studies.* Int J Mol Sci, 2022. **23**(3).

617 29. Madhumita Murgia, *Pandemic puts Oxford Nanopore 'on the map'.* Financial Times,

618   2021. Accessed 11 April 2022.

619 30. Technologies, O.N. *Medaka.* 08 June 2022]; Available from:

620   https://github.com/nanoporetech/medaka.

621 31. Lee, J.Y., et al., *Comparative evaluation of Nanopore polishing tools for microbial*

622   *genome assembly and polishing strategies for downstream analysis.* Sci Rep, 2021.

623   **11**(1): p. 20740.

624 32. Wan, Y.K., et al., *Beyond sequencing: machine learning algorithms extract biology*

625   *hidden in Nanopore signal data.* Trends Genet, 2022. **38**(3): p. 246-257.

626 33. Vasiljevic, N., et al., *Developmental validation of Oxford Nanopore Technology*

627   *MinION sequence data and the NGSpeciesID bioinformatic pipeline for forensic*

628   *genetic species identification.* Forensic Sci Int Genet, 2021. **53**: p. 102493.

629 34. Chen, Z., D.L., et al., Erickson, and J. Meng, *Polishing the Oxford Nanopore long-*

630   *read assemblies of bacterial pathogens with Illumina short reads to improve genomic*

631   *analyses.* Genomics, 2021. **113**(3): p. 1366-1377.

632 35. Delahaye, C. and J. Nicolas, *Sequencing DNA with nanopores: Troubles and biases.*

633   PLoS One, 2021. **16**(10): p. e0257521.

634 36. Sanderson, N., et al., *Comparison of R9.4.1/Kit10 and R10/Kit12 Oxford Nanopore*

635   *flowcells and chemistries in bacterial genome reconstruction.* bioRxiv, 2022: p.

636   2022.04.29.490057.

637 37. Wick, R.R., et al., *Performance of neural network basecalling tools for Oxford*

638   *Nanopore sequencing.* Genome Biol, 2019. **20**(1): p. 129.

639 38. Steinig, E., et al., *Phylodynamic inference of bacterial outbreak parameters using*

640   *Nanopore sequencing.* Mol Biol Evol, 2022. **39**(3).

39.   Khezri, A., et al., *Hybrid assembly provides improved resolution of plasmids,*

antimicrobial resistance genes, and virulence factors in Escherichia coli and

Klebsiella pneumoniae clinical isolates. Microorganisms, 2021. **9**(12).

40.   Boostrom, I., et al., *Comparing long-read assemblers to explore the potential of a*

sustainable low-cost, low-infrastructure approach to sequence antimicrobial resistant

bacteria with Oxford Nanopore sequencing. Front Microbiol, 2022. **13**: p. 796465.

41.   Gorrie C.L., et al., *Genomic dissection of the bacterial population underlying*

Klebsiella pneumoniae infections in hospital patients: insights into an opportunistic

pathogen. Nat Commun, 2022. **13**(1): p. 3017.

42.   Hunt, M., et al., *ARIBA: rapid antimicrobial resistance genotyping directly from*

sequencing reads. Microb Genom, 2017. **3**(10): p. e000131.

43.   Inouye, M., et al., *SRST2: Rapid genomic surveillance for public health and hospital*

microbiology labs. Genome Med, 2014. **6**(11): p. 90.

44.   Argimón, S., et al., *Rapid genomic characterization and global surveillance of*

Klebsiella using Pathogenwatch. Clin Infect Dis, 2021. **73**(Suppl_4): p. S325-S335.

45.   Lam, M.M.C., et al., *A genomic surveillance framework and genotyping tool for*

Klebsiella pneumoniae and its related species complex. Nat Commun, 2021. **12**(1):

p. 4188.

46.   Gorrie, C.L., et al., *Antimicrobial-resistant Klebsiella pneumoniae carriage and*

infection in specialized geriatric care wards linked to acquisition in the referring

hospital. Clin Infect Dis, 2018. **67**(2): p. 161-170.

47.   Hawkey, J., et al., *ESBL plasmids in Klebsiella pneumoniae: diversity, transmission,*

and contribution to infection burden in the hospital setting. medRxiv, 2021.

48.   Wyres, K.L., et al., *Genomic surveillance of antimicrobial resistant bacterial*

colonisation and infection in intensive care patients. BMC Infect Dis, 2021. **21**(1): p.

683.

49.   Oxford Nanopore Technologies, *Guppy v4.0.14.* Available from:

https://github.com/nanoporetech/pyguppyclient. Accessed 08 June 2022.

50.   Oxford Nanopore Technologies, *Guppy v5.0.7 release note (21st May 2021).*

670   Accessed 01 June 2021.

671   51.   Oxford Nanopore Technologies, *qcat v1.1.0.* Available from:

672   https://github.com/nanoporetech/qcat. Accessed 10 March 2022.

673   52.   Chen, S., et al., *fastp: an ultra-fast all-in-one FASTQ preprocessor.* Bioinformatics,

674   2018. **34**(17): p. i884-i890.

675   53.   Wick, R.R., et al., *Unicycler: Resolving bacterial genome assemblies from short and*

676   *long sequencing reads.* PLoS Comput Biol, 2017. **13**(6): p. e1005595.

677   54.   Wick, R.R., et al., *Unicycler: Resolving bacterial genome assemblies from short and*

678   *long sequencing reads.* PLoS Comput Biol, 2017. **13**(6): p. e1005595.

679   55.   Bankevich, A., et al., *SPAdes: a new genome assembly algorithm and its*

680   *applications to single-cell sequencing.* J Comput Biol, 2012. **19**(5): p. 455-77.

681   56.   Kolmogorov, M., et al., *Assembly of long, error-prone reads using repeat graphs.* Nat

682   Biotechnol, 2019. **37**(5): p. 540-546.

683   57.   Wick, R.R. and K.E. Holt, *Benchmarking of long-read assemblers for prokaryote*

684   *whole genome sequencing.* F1000Res, 2019. **8**: p. 2138.

685   58.   Walker, B.J., et al., *Pilon: an integrated tool for comprehensive microbial variant*

686   *detection and genome assembly improvement.* PLoS One, 2014. **9**(11): p. e112963.

687   59.   Zhang, P., et al., *Comparison of de novo assembly strategies for bacterial genomes.*

688   Int J Mol Sci, 2021. **22**(14).

689   60.   Wick, R.R. *Guide to bacterial genome assembly: long read polishing*. Available from:

690   https://github.com/rrwick/Trycycler/wiki/Guide-to-bacterial-genome-assembly#12-

691   long-read-polishing. Accessed 12 February 2022.

692   61.   Gurevich, A., et al., *QUAST: quality assessment tool for genome assemblies.*

693   Bioinformatics, 2013. **29**(8): p. 1072-5.

694   62.   Parks, D.H., et al., *CheckM: assessing the quality of microbial genomes recovered*

695   *from isolates, single cells, and metagenomes.* Genome Res, 2015. **25**(7): p. 1043-55.

696 63. Wick, R.R. and Holt, K.E., *Polypolish: Short-read polishing of long-read bacterial*

697   *genome assemblies.* PLoS Comput Biol, 2022. **18**(1): p. e1009802.

698 64. Diancourt, L., et al., *Multilocus sequence typing of Klebsiella pneumoniae nosocomial*

699   *isolates.* J Clin Microbiol, 2005. **43**(8): p. 4178-82.

700 65. Gorrie, C.L., et al., *Gastrointestinal carriage is a major reservoir of Klebsiella*

701   *pneumoniae infection in intensive care patients.* Clin Infect Dis, 2017. **65**(2): p. 208-

702   215.

703 66. Sherry, N.L., et al., *Genomics for molecular epidemiology and detecting transmission*

704   *of carbapenemase-producing Enterobacterales in Victoria, Australia, 2012 to 2016.* J

705   Clin Microbiol, 2019. **57**(9).

706 67. David, S., et al., *Epidemic of carbapenem-resistant Klebsiella pneumoniae in Europe*

707   *is driven by nosocomial spread.* Nat Microbiol, 2019. **4**(11): p. 1919-1929.

708 68. Holt, K.E., *RedDog.* Available from: https://github.com/katholt/RedDog. Accessed 10

709   March 2022.

710 69. Seemann T, *snp-dist.* Available from: https://github.com/tseemann/snp-dists.

711   Accessed 12 April 2022.

712 70. Robinson, D.R., Foulds, L. R., *Comparison of phylogenetic trees*, in *Mathematical*

713   *Biosciences ,* . 1981. p. 131-147.

714 71. Revell, L.J., *phytools: an R package for phylogenetic comparative biology (and other*

715   *things).* Methods Ecol Evol, 2012. **3**(2): p. 217-223.

716 72. Paradis, E. and Schliep, K., *ape 5.0: an environment for modern phylogenetics and*

717   *evolutionary analyses in R.* Bioinformatics, 2019. **35**(3): p. 526-528.

718 73. Wilke, C.O., *cowplot: Streamlined plot theme and plot annotations for 'ggplot2'.*

719   Available from https://wilkelab.org/cowplot/. Accessed 10 January 2022.

720 74. Dowle, M., et al., *data.table: Extension of 'data.frame'.* Available from

721   https://rdatatable.gitlab.io/data.table/. Accessed 03 April 2022.

722 75. Wright, E., *Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R.*

723   The R Journal, 2016. 8: p. 352-359.

76. Galili, T., *dendextend: an R package for visualizing, adjusting and comparing trees of*

hierarchical clustering. Bioinformatics, 2015. **31**(22): p. 3718-20.

77. Wickham, H., et al. *dplyr: A grammar of data manipulation*. Available from:

https://dplyr.tidyverse.org, https://github.com/tidyverse/dplyr. Accessed 07 June

2022.

78. Yu, G., *Using ggtree to visualize data on tree-like structures.* Curr Protoc

Bioinformatics, 2020. **69**(1): p. e96.

79. Kassambara, A., et al. *ggpubr: 'ggplot2' based publication ready plots*. Available

from: https://rpkgs.datanovia.com/ggpubr/. Accessed 10 February 2022.

80. Auguie, B. *gridExtra: Miscellaneous functions for "Grid" graphics*. Available from:

https://cran.r-project.org/web/packages/gridExtra/gridExtra.pdf. Accessed 06 June

2022.

81. Firke, S., et al., *janitor: Simple tools for examining and cleaning dirty data*. Available

from: https://cran.r-project.org/web/packages/janitor/index.html. Accessed 03 June

2022.

82. Zhu, H., et al., *kableExtra: Construct complex table with 'kable' and pipe syntax*.

Available from

http://haozhu233.github.io/kableExtra/, https://github.com/haozhu233/kableExtra.

Accessed 08 June 2022.

83. Wickham, H. and G. Grolemund. *Tidyverse: R packages for data science*. Available

from: https://www.tidyverse.org/. Accessed 08 June 2022.

84. Wang, L.G., et al., *Treeio: An R package for phylogenetic tree input and output with*

richly annotated and associated data. Mol Biol Evol, 2020. **37**(2): p. 599-603.

85. Tennekes, M. and P. Ellis. *treemap: Treemap visualization*. Available from:

https://cran.r-project.org/web/packages/treemap/treemap.pdf. Accessed 04 January

2022.

86.   Wickham, H. *stringr: Simple, consistent wrappers for common string operations*. 11

February 2022]; Available from: http://stringr.tidyverse.org,

https://github.com/tidyverse/stringr. Accessed 05 June 2022.

87.   Gorrie, C.L., et al., *Genomic dissection of Klebsiella pneumoniae infections in*

hospital patients reveals insights into an opportunistic pathogen. Nat Commun, 2022.

13(1): p. 3017.

88.   Wick, R.R., et al., *Kaptive Web: User-friendly capsule and lipopolysaccharide*

serotype prediction for Klebsiella genomes. J Clin Microbiol, 2018. **56**(6).

89.   Follador, R., et al., *The diversity of Klebsiella pneumoniae surface polysaccharides.*

Microb Genom, 2016. **2**(8): p. e000073.

90.   Lam, M.M.C., et al., *Genetic diversity, mobilisation and spread of the yersiniabactin-*

encoding mobile element ICEKp in Klebsiella pneumoniae populations. Microb

Genom, 2018. **4**(9).

91.   Lam, M.M.C., et al., *Tracking key virulence loci encoding aerobactin and salmochelin*

siderophore synthesis in Klebsiella pneumoniae. Genome Med, 2018. **10**(1): p. 77.

92.   Wick, R.R., et al., *Trycycler: consensus long-read assemblies for bacterial genomes.*

Genome Biol, 2021. **22**(1): p. 266.

93.   Sereika, M., et al., *Oxford Nanopore R10.4 long-read sequencing enables near-*

perfect bacterial genomes from pure cultures and metagenomes without short-read

or reference polishing. bioRxiv, 2021: p. 2021.10.27.466057.

Figures, legends and table

Figure 1. Genome assembly flow diagram. We generated ten assemblies for each of 54 *Klebsiella pneumoniae* isolates, which were sequenced on both Illumina and ONT MinION (numbered 1 to 10). ONT reads were basecalled using three alternative models (via Guppy v4.0.14): fast, high-accuracy (HAC), and super-accuracy. The resulting basecalled read sets were each subjected to assembly using Flye (v2.8.1), generating three ONT-only assemblies per isolate. These assemblies were each then polished with the corresponding ONT read set (using Medaka models r941_min_fast_g303 and r941_min_high_g360) to generate a further three ONT-only polished assemblies per isolate, totalling 6 ONT-only assemblies (3 polished, 3 unpolished) per isolate. For comparison, we used Illumina data to generate Illumina-only and ONT/Illumina hybrid assemblies for each isolate. Illumina reads were assembled using SPAdes v3.14.0 to generate an Illumina-only assembly. A short-read-first hybrid assembly

was generated using Unicycler v0.4.9b and then polished using Pilon v1.24. Finally, two long-read-first hybrid assemblies were generated by polishing the ONT-only HAC and super-accuracy models using Illumina reads (via pilon v1.24). *SPAdes was run via Unicycler v0.4.9b. ∞The Fast and HAC basecalling utilised Guppy v4.0.14 while SUP basecalling was done using Guppy v5.0.7.

Figure 2. Genotyping accuracy for MLST, yersiniabactin and K/O loci using different assemblies. Each panel summarises accuracy of genotyping results across n=54 isolates, for each type of assembly (one per row as labelled on the left—numbered 1 to 10 as in **Figure 1**, and ordered based on overall performance), compared to the reference assembly (i.e., long-read first hybrid assembly using Flye and Medaka to assemble and polish with super-accurate

basecalled reads, then polished with Illumina; last row per panel). (**A**) Multi-locus sequence type (ST) calling; 1LV, 2LV, 3LV, 4LV indicates the correct ST was predicted but the allele call was incorrect for 1, 2, 3, or 4 alleles. (**B**) Detection and typing of yersiniabactin; 'correct call, no determinants present' indicates the *ybt* locus was absent in the reference assembly and was (correctly) not detected in the test assembly; 'correct call, determinants present' indicates the *ybt* locus was present in the reference assembly and this was correctly identified and subtyped (to lineage level) in the test assembly; 'determinants present, called as unknown', indicates the *ybt* locus was present in the reference assembly, and this was correctly identified as present but could not be subtyped in the test assembly. (**C**) K locus and (**D**) O locus typing; 'correct call' indicates the correct locus was identified with confidence ranking of good or better; 'non-confident call ("best-match"), correct' indicates that the correct locus was identified but with low or no confidence. (**E**) and (**F**) show the number of open reading frames that are encoded in the reference K and O loci but were not detected in the test assemblies (labelled as 'missing genes' by the Kaptive genotyper, although in this case the nucleotide sequences are present but basecall errors result in disruption of the open reading frames). (**G**) and (**H**) show the nucleotide divergence between each assembly and the reference K and O loci, across the length of the detected loci.

Figure 3. Neighbour-joining tree representing the phylogenetic relationships among the 540 genome assemblies, derived from n=54 *K. pneumoniae* isolates as outlined in Figure 1. The tree was constructed from the Pathogenwatch distance matrix (i.e. based on SNPs called in 1972 core genes) and is midpoint-rooted. Tips are coloured by assembly type, as indicated by the inset legend; note the reference assembly (i.e. long-read first hybrid assembly using Flye and Medaka to assemble and polish with super-accurate basecalled reads, then polished with Illumina) is coloured black and all other Illumina-containing genomes are grouped and coloured purple. Alternate assemblies of the same genome tended to cluster together but are non-zero branches because the sequences are non-identical due to basecalling errors. ONT/Illumina hybrid assemblies of the same isolated clustered more closely than ONT-only assemblies. Assemblies derived from the Fast model (±medaka, blue colours) consistently formed outliers separated by long branches, due to high rates of basecalling errors. Most of the study isolates belong to distinct *K. pneumoniae* lineages and

are unrelated, except for sixteen isolates belonging to four transmission clusters (see **Figure 6**), these clades are highlighted in green and labelled by multi-locus sequence type (ST).

Figure 4. SNPs called vs reference assembly. A violin plot depicting the distribution of SNPs between each assembly vs the gold-standard hybrid reference assembly for the fifty-four study genomes based on a concatenated alignment of 1,972 genes (2,172,367bp) that make up the core gene library for *K. pneumoniae* in Pathogenwatch. Assembly types are numbered 1 to 10 as in **Figure 1**. Data points corresponding to ONT-only assemblies are coloured red. Within the cluster of ten assemblies for each sample, the SUP and HAC were consistently separated by a reasonable SNP threshold (≤10 SNPs); however, the Fast assemblies tended to form outliers (>100 SNPs). For clustering, only the SUP+Medaka assemblies reliably identified outbreak clusters (≤25 genome-wide SNPs threshold, which corresponds to ≤11 SNPs within the Pathogenwatch *K. pneumoniae* core genome typing scheme). HAC+Medaka polishing performed similarly to SUP without polishing, and polishing introduced errors into HAC- or Fast-basecalled assemblies.

Figure 5. Comparison of trees constructed from ONT-only vs reference assemblies.
Trees are midpoint-rooted neighbour-joining trees inferred from Pathogen pairwise distance matrices calculated from ONT-only (SUP+medaka) assemblies or ONT / Illumina long-read-first hybrid reference assemblies. Trees are shown as a tanglegram, which attempts to render the tree visualisations to maximise the alignment of tips, and links tips with lines (coloured to highlight shared subtrees), to facilitate comparison of topology between the two trees. The tanglegram yields an entanglement coefficient of 0.33 and a Robinson-Foulds distance of 20, indicating a good alignment. Coloured lines join matching tip labels for common subtrees, while grey lines indicate subtrees that are not common between the two trees. Discordant branch lengths on the ONT-only tree arise from basecalling errors, despite a good alignment.

C. Pairwise Pathogenwatch (PW) distances for known transmission clusters, calculated using different assemblies

ST29	INF249	INF250	INF322
INF249	0	1	1
INF250	8 (18)	0	0
INF322	10 (23)	6 (13)	0

ST231	INF281	INF298	INF299
INF281	0	0	1
INF298	4 (14)	0	1
INF299	4 (16)	4 (10)	0

ST491	INF344	INF355	INF361
INF344	0	1	1
INF355	6 (19)	0	2
INF361	16 (32)	20 (28)	0

ST323	INF277	INF302	INF329	KSB1_10J	KSB1_4E	KSB1_7E	KSB2_1B
INF277	0	16	23	34	21	20	21
INF302	58 (89)	0	11	22	9	8	9
INF329	91 (128)	93 (142)	0	29	16	15	16
KSB1_10J	551 (608)	541 (613)	578 (659)	0	21	20	27
KSB1_4E	68 (83)	60 (92)	99 (134)	547 (605)	0	3	14
KSB1_7E	80 (102)	76 (105)	103 (146)	555 (621)	72 (95)	0	13
KSB2_1B	89 (128)	85 (136)	118 (183)	562 (634)	97 (138)	105 (152)	0

Figure 6. Comparison of pairwise Pathogenwatch distances between n=54 isolates, calculated from ONT-only vs Illumina assemblies. Y-axis (**A** and **B**) shows pairwise distances between isolates, calculated using SUP+Medaka assemblies for all isolates; x-axis shows pairwise SNP distances between isolates, calculated using Illumina assemblies for all isolates (based on the core gene set of 1,972 genes within the Pathogenwatch *Klebsiella pneumoniae* core genome scheme). Each data point represents a pair of isolates, coloured to indicate isolate pairs belonging to the same transmission cluster (as per inset legend). (**A**) All isolate pairs; line shows y=x. (**B**) Zoom in to transmission cluster isolates with ONT-based pairwise distances ≤150; linear regression line is shown (solid line), adjusted R^2=0.8923 indicating very good model fit, slope=5.1. (**C**) Pairwise Pathogenwatch (PW) distances for known transmission clusters, calculated using different assemblies. PW distances calculated from reference assemblies are highlighted in orange; those calculated from ONT-only with SUP basecalling are highlighted in blue (numbers in brackets are those without Medaka polishing). Distances below the clustering threshold for PW distances, i.e., ≤11, are bolded.

Figure 7. Detection of AMR-associated mutations in core chromosomal genes. Each panel summarises accuracy of genotyping results across n=54 isolates, for each type of assembly (one per row as labelled on the left—numbered 1 to 10 as in **Figure 1**, and ordered based on overall performance), compared to the reference assembly (i.e. long-read first hybrid assembly using Flye and Medaka to assemble and polish with super-accurate basecalled reads, then polished with Illumina; last row per panel). **(A)** Detection of fluoroquinolone resistance-associated mutations (GyrA-83I, ParC-80I, GyrA-83Y, GyrA-87N and GyrA-87Y). **(B)** Detection of mutations in *bla*SHV that are associated with a change in enzyme activity (35Q, 146V, 238S and 156D). **(C)** Detection of colistin resistance-associated mutations (disruption of MgrB or PmrB). **(D)** Detection of mutations in outer membrane porins associated with change in carbapenem susceptibility (truncation of OmpK35 or OmpK36, GD or TD insertions in OmpK36 loop 3).

Supplementary Figure 1. Comparison of Pathogenwatch distances with genome-wide mapping-based SNP distances, across n=270 diverse *Klebsiella pneumoniae* clinical isolates using Illumina data. Y-axis shows Pathogenwatch pairwise distances (based on the core gene set of 1,972 genes within the Pathogenwatch *Klebsiella pneumoniae* core genome scheme) using Illumina assemblies; x-axis shows genome-wide mapping-based pairwise SNP distances (based on all variant sites for which alleles were called in ≥95% genomes, when mapping Illumina reads to the NTUH-K2044 reference genome as described in reference 40). Data points represent pairwise distances, coloured to indicate pairs of genomes with the same (red) or different (blue) 7-locus sequence type. (**A**) All pairwise distances. (**B**) Zoom in to data points with pairwise genome-wide SNP distances ≤200; linear regression line is shown (solid line), adjusted R^2=0.9914 indicating very good model fit, slope=0.432; dashed line shows x=25 genome-wide SNPs, which is a common threshold used to identify putative transmission clusters. According to the linear regression, this is equivalent to a Pathogenwatch distance of 10.8 SNPs (i.e., threshold n=10).

Supplementary Figure 2. Distributions of sequencing metrics stratified by assembly type. Assembly types are numbered 1 to 10 as in **Figure 1**. (**A**) Genome size. (**B**) N50. N50 distribution for Illumina is not shown as it generates draft assemblies with N50 < 1 Mbp (range, 69194 – 612419; median, 274529) (**C**) Contig count. Similar to panel B, contig count distribution for Illumina is not shown as it generates draft assemblies with >20 contigs (range 32 – 315; median, 92) (**D**) %GC content. The y-axis shows the various assemblies under comparison, with the gold standard hybrid ONT / Illumina reference assembly (SUP+Medaka+pilon); while the x-axis displays the genome size, N50, contig count and %GC content distribution, respectively.

Supplementary Figure 3. A tanglegram depicting the alignment between a Neighbor-Joining tree produced from HAC+medaka assemblies and the ONT / Illumina hybrid reference tree. The tree yields an entanglement coefficient of 0.35 and a Robinson-Foulds distance of 22, indicating a good alignment. However, several clades are discordant with the reference tree, represented by the dotted lines. Coloured lines join matching tip labels for common subtrees, while grey lines indicate subtrees that are not common between the two trees. Errors in the ONT-only assembly result in discordant branch lengths between the two trees.

Supplementary Figure 4. A tanglegram depicting the alignment between a Neighbor-Joining tree produced from Fast+medaka assemblies and the ONT / Illumina hybrid reference tree. The tree yields an entanglement coefficient of 0.96 and a Robinson-Foulds distance of 102, indicating a very poor alignment. Coloured lines join matching tip labels for common subtrees, while grey lines indicate subtrees that are not common between the two trees. Dotted lines represent distinct subtree present on either tree. Errors in the ONT-only assembly result in discordant branch lengths between the two trees.

Table 1: Accuracy of identifying non-susceptibility to specific drugs/classed based on detection of known AMR determinants

	Aminoglycosides	Carbapenems	3rd gen. cephalosporins	Fluoroquinolones	Phenicols	Sulfamethoxazole	Trimethoprim	Tetracycline
Reference genomes with AMR determinants N (% of total)	26 (48.1%)	6* (11.1%)	25 (46.3%)	^27 (50.0%)	17 (31.5%)	26 (48.1%)	24 (44.4%)	20 (37.0%)
ONT-only assemblies with AMR determinants **N (% of reference genome positives)**								
SUP+Medaka	**24 (92.3%)**	**6 (100%)**	**22 (88%)**	26 (96.3%)	**16 (94.1%)**	26 (100%)	24 (100%)	20 (100%)
SUP	11 (42.3%)	**6 (100%)**	21 (84%)	**27 (100%)**	14 (82.4%)	25 (96.2.3%)	24 (100%)	11 (55%)
HAC+Medaka	20 (76.9%)	**6 (100%)**	15 (60%)	**27 (100%)**	14 (82.4%)	24 (92.3%)	21 (87.5%)	15 (75%)
HAC	2 (7.7%)	**6 (100%)**	19 (76.0%)	**27 (100%)**	14 (82.4%)	8 (30.8%)	14 (58.3%)	2 (10%)
Fast+Medaka	3 (11.5%)	**6 (100%)**	8 (32%)	17 (63.0%)	13 (76.5%)	21(80.9%)	19 (79.2%)	4 (20%)
Fast	1 (3.8)	5 (71%)	6 (24%)	13 (48.1%)	12 (70.6%)	4 (15.4)	8 (33.3%)	1 (5%)

Reference assemblies are long-read-first ONT/Illumina hybrid assemblies generated using SUP-basecalled ONT data. *Carbapenem resistance determinants detected were mainly acquired genes (n=4 genomes) than porin mutations (n=2 genomes). ^Fluoroquinolone resistance mutations detected were mainly acquired genes (n=18 genomes) rather than point mutations (n=9 genomes). Details of resistance determinants detected in each reference genome are given in **Supplementary Table 1**. The highest percentages of correctly identified AMR determinants per antibiotic class are **bolded**.