Title: Habitual aerobic exercise evokes fast and persistent adaptation during split-belt walking

Authors: Sarah A. Brinkerhoff[1], *Natalia Sánchez[2], *Jaimie A. Roper[1]

*Indicates co-senior authorship, these authors contributed equally to this work

[1]School of Kinesiology, Auburn University, Auburn, Alabama, United States of America

[2]Department of Physical Therapy, Chapman University, Irvine, California, United States of

America

Corresponding author:

Sarah A. Brinkerhoff

Email: sbrinkerhoff@auburn.edu

Keywords: biomechanics, motor learning, gait adaptation, energetics, exercise

Summary Statement:

Young adults who regularly participate in aerobic exercise respond more quickly to the onset of

a split-belt perturbation and continue to adapt their gait pattern throughout the duration of the

split-belt exposure.

List of Abbreviations:

SLA = step length asymmetry

Abstract:

Changing movement patterns in response to environmental perturbations is a critical aspect of

gait and is related to reducing the energetic cost of the movement. Aerobic exercise improves

energetic capacity for submaximal exercise and may affect how people adapt movement to

reach an energetic minimum. The purpose of this study was to determine whether differences in

recreational aerobic exercise training influence gait adaptation in healthy young adults. Young

adults who habitually exercised (n=19) and young adults who did not habitually exercise (n=13)

walked on a split-belt treadmill with one belt moving twice the speed of the other belt for 10

minutes. Step length asymmetry (SLA) and mechanical work done by each leg were measured

over each stride. Nonlinear mixed effects models compared the time course of adaptation

between Habitual Exercisers and Non-exercisers, and *t*-tests compared net work at the end of

adaptation between Habitual Exercisers and Non-exercisers. Compared to Non-exercisers,

Habitual Exercisers initially adapted SLA and positive work by the fast leg quickly, and

continued to modify SLA over the duration of split-belt treadmill walking. Young adults who

regularly participate in aerobic exercise respond more quickly to the onset of a perturbation, and

throughout the perturbation they continue to explore movement strategies, which might be

related to reduction of energetic cost.

Introduction

Adapting walking patterns in response to environmental perturbations is a critical aspect

of locomotion. When a person encounters a perturbation — such as when walking on a boat

rocking on the water or walking on an icy surface — they must adapt their walking patterns to

avoid falling, which can be achieved using different strategies. How individual factors might

influence walking adaptation strategies remains to be determined.

A common approach to study walking adaptation is split-belt treadmill walking, in which

the belts under each leg move at different speeds (1,2). The changes in the asymmetry between

left and right step lengths, or step length asymmetry (SLA) is one measure used to track

adaptation. SLA robustly changes during split-belt walking, shows an aftereffect after the split-

belt perturbation is removed, is observable with the unaided eye, and is sensitive to

experimental manipulations (3–7). In line with upper extremity motor adaptation (8), adaptation

of SLA during split-belt walking occurs at two distinct rates – a fast component that adapts

rapidly and a slow component that adapts more gradually (9–12). From work in both upper and

lower extremity motor tasks, the two timescales of motor adaptation may derive from two

separate (but not necessarily independent) processes

Indeed, Seethapathi and colleagues posited that the component that adapts rapidly is

driven by balance optimization, whereas the component that adapts more gradually is driven by

energetic cost optimization (13). Consistent with these findings, experimental studies have

shown that SLA adaptation during split-belt walking occurs in parallel with reductions in the work

generated by the legs; gait adaptation results in decreasing positive work and increasing

negative work done by the legs, especially the leg on the fast belt (14,15). As the legs reduce

positive work done and increase negative work done, energetic cost decreases concomitantly

(14,16,17). This complements the fact that doing negative work is less energetically costly than

doing positive work (18,19). By reducing work done by the legs, people are likely gradually

adapting towards some lower energetic cost, as determined by Seethapathi et. al.(13) and

previously empirically suggested (20–22). Similarly, a study by Park et al showed that during

early adaptation to split belt walking, individuals increase whole body angular momentum, which

is inversely proportional to balance (23). Therefore, these experimental findings also support the

idea of balance and energetic cost as the two distinct processes driving gait adaptation.

The question then arises — do individual factors that affect energy consumption affect

the ability to adapt gait during split-belt walking? The multisystem benefits of aerobic exercise

are well-known and include improved physical function, cognition, quality of life, and reduced

risk of cardiovascular disease and all-cause mortality (24). Moreover, the ability to move at a

minimum energetic cost of transport at a given speed is a function of aerobic training

experience, such that people with more aerobic training reach a minimum cost of transport while

running, but those who engage in less aerobic training do not (25). Along with an ability to reach

an energetic minimum, aerobically trained adults have a higher maximal oxygen uptake

(VO_2max) (26), a better exercise economy (26,27), and higher lactate and ventilatory thresholds

(26). These training-induced adaptations would increase capacity for exercise and enable

aerobically trained individuals to tolerate submaximal exercise for longer. If we posit that people

adapt to split-belt walking in order to reduce energetic cost, then those who engage in more

aerobic exercise may reach an energetic minimum quicker than those who engage in less

exercise (25); conversely, those who engage in more aerobic exercise may more gradually

reach an energetic minimum, due to a higher tolerance for submaximal exercise (26,27). These

two hypotheses indicate that fitness might influence adaptation toward more energetically

economical movement patterns.

The purpose of this study was to determine whether differences in weekly amount of

exercise influence gait adaptation in healthy young adults. We hypothesized that amount of

exercise would affect gait adaptation, and this effect would primarily be driven by differences in

the slow component of adaptation, given the role of energetics in shaping the rate of adaptation

of the slow component (13,20–22) and the well-established effects of exercise on energetics

(25–27). Our hypothesis is based on two opposing mechanisms which we will test here: 1)

Habitual Exercisers would adapt faster than Non-exercisers to reach a minimum energetic cost

quickly (25), evidenced by faster reduction in SLA and work done by the legs; 2) Habitual

Exercisers would have a higher tolerance for the submaximal exercise of treadmill walking than

Non-exercisers, evidenced by more gradual reduction in SLA and work done by the legs. To test

these competing mechanisms, we analyzed the adaptation of SLA and the positive and negative

work done by each leg while Habitual Exercisers and Non-exercisers walked on a split-belt

treadmill. We also hypothesized that at the end of adaptation, Habitual Exercisers would

perform more-negative net work than Non-exercisers, by reducing positive work done by the

legs and increasing negative work done by the legs in order to reduce overall energetic cost

(14,16,17). A significant effect of regular exercise participation on gait adaptation would support

the idea that individual factors that influence energetics can also influence walking adaptation.

Our findings will provide insights into the multisystem positive effects of exercise, including

walking adaptation, and will provide directions for exercise rehabilitation research.

Methods

Participants

We recruited a convenience sample of young adults ages 19-35 for participation in this

study. Participants were excluded if they reported cardiovascular, pulmonary, renal, metabolic,

vestibular, or neurologic disorders; any lower-extremity injuries or surgeries in the past 12

months; or a prior anterior cruciate ligament injury. The Auburn University Institutional Review

Board approved all procedures and all participants completed written informed consent prior to

participating (Protocol #18-418 MR 1811). Data were collected in the Locomotor and

Movement Control Lab at Auburn University.

Experimental Protocol

Participants first completed a self-report exercise questionnaire. This questionnaire

asked how many times per week participants currently exercised at a moderate-to-vigorous

intensity, for how many minutes per session, and for how many consecutive weeks. Based on

their responses, participants were grouped into one of two groups according to the American

College of Sports Medicine Physical Activity Guidelines for Americans: Habitual Exercisers (at

least 150 minutes per week of exercise for at least the last three months) or Non-exercisers

(less than 150 minutes per week of exercise for at least the last three months) (24). The

questionnaire also asked participants which leg they kicked a ball with – the reported kicking leg

was determined the dominant leg and was always set on the fast belt during adaptation.

After completing the questionnaire, participants walked on the treadmill. Participants

held onto side handrails for the entirety of treadmill walking. The instructions given to

participants noted that they should use the handrails for stability, and not to offload bodyweight.

First, participants determined their typical and their fastest comfortable walking speeds on the

treadmill using a modified protocol by Dingwell and Marin (28). The treadmill was accelerated to

a relatively slow speed, and the speed slowly increased until the participant reported that the

current speed was their typical walking speed or was their fastest comfortable speed. This was

repeated twice for typical walking speed, and twice for fastest comfortable walking speed. The

prompt for typical walking speed was, "tell me when you reach your typical walking speed." The

prompt for fastest comfortable speed was, "tell me when you reach the fastest speed that you'd

be comfortable walking for ten minutes." The participant's fastest comfortable speed was set as

the fast belt speed, and half of the fastest comfortable speed was set as the slow belt speed.

The split-belt walking protocol is shown in Figure 1. After finding their preferred typical

and fast speeds, participants walked for three minutes at their typical walking speed, followed by

three minutes at their fastest comfortable speed, followed by three minutes at half of the fast

speed (slow speed). The last tied walking condition — three minutes of walking at the slow

speed — was considered the baseline condition for data analysis. Then, participants walked for

ten minutes with the belt speeds split — the belt under the dominant leg moved at participants'

fastest comfortable speed, and the belt under the non-dominant leg moved at the slow speed.

Figure 1. Split-belt walking protocol. A) An oblique and birds-eye view of the split-belt

treadmill. B) First, the participant walked for 3 minutes at their self-selected typical

walking speed. Then, they walked for 3 minutes at their self-selected fastest comfortable

walking speed. Then, they walked for 3 minutes at half of their fast speed. Finally, they

walked for 10 minutes with the belt under their dominant leg at the fast speed, and the

belt under their non-dominant leg at the slow speed.

Data Analysis

Kinematic data were recorded at 100 Hz from reflective markers placed bilaterally on the

anterior superior iliac spine and the lateral malleoli of the ankles, using a 17-camera motion

capture system (VICON; Vicon Motion Systems Ltd, Oxford, United Kingdom). Kinematic data

were lowpass filtered with a 4^{th} order Butterworth filter with a cutoff frequency of 6 Hz. Ground

reaction force data were obtained for each individual leg using an instrumented split-belt

treadmill, recorded at 1000 Hz from two separate force plates. Force data were lowpass filtered

with a 4^{th} order Butterworth filter with a cutoff frequency of 20 Hz.

Step length was calculated as the distance between the ankle markers along the

anterior-posterior walking axis at foot strike. SLA was calculated and normalized to stride length

as follows:

$$Step\ length\ Asymmetry = \frac{Step\ length_{fast} - Step\ length_{slow}}{Step\ length_{fast} + Step\ length_{slow}}$$

Here, *Step Length$_{fast}$* is the step length when the leg on the fast belt strikes the belt, and

Step Length$_{slow}$ is the step length when the leg on the slow belt strikes the belt. A negative SLA

indicates that the leg on the slow belt is taking a longer step than the leg on the fast belt, and an

SLA of zero indicates that the legs are taking steps of equal length.

We calculated positive and negative mechanical work generated by the legs using a

custom MATLAB program. We used the point-mass model to estimate mechanical work

generated by the legs on the treadmill and on the center of mass (12,14,15). In brief, we

segmented force data into strides (15) and calculated the center of mass velocities in each

direction as the time integral of the center of mass accelerations. Next, we calculated the

instantaneous power generated by each leg for each stride as the instantaneous sum of the dot

product of the ground reaction force and the center of mass velocity and the dot product of the

force applied to the respective belt and the belt speed. We then calculated the total positive and

total negative work performed by each leg as the time integral of the positive or negative portion

of the total instantaneous power over the stride cycle. We calculated work rate by dividing each

work measure by stride duration. Finally, we calculated the net work rate by the legs at the end

of adaptation as the time integral of the power divided by stride cycle, averaged over the last

100 strides.

Statistical Analysis

All statistical analyses were conducted in R (29). We used Welch Two Sample t-tests to

compare age, height, leg length, mass, and fast-leg belt speed between groups. We also used

Pearson's Chi-squared tests with Yates' continuity correction to assess the difference in

proportion of males and females in each group, and to assess the difference in proportion of

people who walked with the right leg on the fast belt in each group. We analyzed SLA, positive

and negative work rate by the fast legs, and positive and negative work rate by the slow leg

using mixed effects nonlinear regression models with the 'nlme' package in R (30). We

truncated the data for all participants to minimum number of steps taken by a participant (823

steps) so that all participants would weight equally on each step.

Detailed description of the statistical models can be found in the supporting information.

In brief, for each of the work rate outcome measures and for SLA, we built-up models to two-

exponent models — assuming that adaptation of the outcome measures occurred over two

timescales — that included a fixed effect for group (eqs. 1 and 2). Here, c is the estimated

plateau if strides went to infinity; a_f is the initial value of the fast component of adaptation; r_f is

the growth rate of the fast component of adaptation; a_s is the initial value of the slow component

of adaptation; r_s is the growth rate of the slow component of adaptation. All five model

parameters (c, a_f, r_f, a_s, r_s) were allowed to differ for Habitual Exercisers vs. Non-exercisers.

$$SLA = (c + a_f * e^{-\frac{step}{r_f}} + a_s * e^{-\frac{step}{r_s}}) \sim Group + (c|ID) \qquad 1$$

$$Work\ rate = \left(c + a_f * e^{-\frac{stride}{r_f}} + a_s * e^{-\frac{stride}{r_s}}\right) \sim Group + (c|ID) \qquad 2$$

We compared these two-exponent models with fixed effects for group to models without

fixed effects for group, and to one-exponent models assuming adaptation occurs over a single

timescale. All models and outcomes can be found in the supporting information. The simplest,

best-fitting model for each measure was the final model that we interpreted, such that a lower

Akaike Information Criterion (AIC) and Bayesian Information Criteria (BIC) and a ΔAIC > 2

indicated a better fit to the data (31,32). All models were fit with a maximum likelihood

estimation and all models contained a random effect such that the outcome measure's

estimated plateau was allowed to vary by participant. If the final model included a fixed effect for

group, the significance (or lack thereof) of the effect of group on the equation variables in the

best-fitting models was evaluated using Welch's t-test with an *a priori* alpha level of 0.05. If the

final model included a fixed effect for group, initial adaptation values were calculated for each

variable as the average over the first five strides and were compared between group using

Welch's Two Sample t-test.

We used Welch's Two Sample t-test between groups to compare the net work rate by

the legs at the end of adaptation, to determine if there was a difference between exercise

groups in the ability to gain assistance from the treadmill.

Results

Participants

Thirty-two participants completed the study (6 female, age 22 (SD 3) yrs, mass 70.3 (SD

14.6) kg, height 171.82 (SD 9.75) mm). Based on the self-report exercise questionnaire, 19

participants were placed in the Habitual Exercisers group and 13 were placed in the Non-

exercisers group. The two groups did not significantly differ by sex, height, mass, leg on the fast

belt, or fast belt speed (Habitual Exercisers = 1.49 (0.21) m/s; Non-exercisers = 1.57 (0.16)

m/s), but did significantly differ by age where Habitual Exercisers were slightly older (Habitual

Exercisers 22 (SD 4) yrs; Non-exercisers 20 (SD 1) yrs, $t(21.827)=-2.127$, $p=0.045$).

Demographic and belt speed data are provided in Table 1.

Table 1. Participant demographics in Mean (SD) [range].

	Non-exercisers	Habitual Exercisers	Total
n	*13*	*19*	*32*
Age (yr)	20 (1) [19,22]	22 (4) [19,32]	22 (3) [19,32]
Sex	3 F	3 F	6 F
Height (mm)	1706 (99) [1540,1910]	1727 (98) [1540,1920]	1718 (97) [1540,1920]
Mass (kg)	66.5 (11.9) [48,89]	72.9 (16.0) [50,116]	70.3 (14.6) [48,116]
Fast leg	3 right	3 right	6 right
Fast belt speed (m/s)	1.57 (0.16) [1.25,1.95]	1.49 (0.21) [1.15,1.98]	1.52 (0.20) [1.15,1.98]
Minutes/week exercise	38 (44) [0,120]	268 (157) [150,750]	175 (168) [0,750]
Exercise modality	Run, cycle, basketball, tennis	Run, cycle, basketball, tennis, CrossFit[®], dance, boxing	—

235   Note. Sex: F = female. Type of exercise participated in reflects the exercise modalities of people
236   who participated in any exercise every week (regardless of minutes per week of exercise).
237

General Timescales of Adaptation

The best-fitting model for each outcome measure was the two-exponent model

containing a fixed effect for group (eqs. 1 and 2, Table 2), indicating that SLA and work rate by

the legs adapted over two distinct timescales and was affected by exercise. All candidate

models for each outcome measure, and the final model fits to each participant's data, can be

found in the supporting information.

Table 2. Final mixed effects models for SLA, positive and negative work of the fast leg, and
positive and negative work of the slow leg.

	Coefficient	SLA	Positive Work Rate		Negative Work Rate	
			Fast Leg	Slow Leg	Fast Leg	Slow Leg
Non-exercisers	c (plateau)	-0.043 *** (0.010)	0.54 *** (0.12)	0.20 (0.17)	-0.28 ** (0.10)	-0.48 ** (0.16)
	a_f (fast component initial value)	-0.097 *** (0.007)	0.38 *** (0.06)	0.28 *** (0.03)	0.02 (0.02)	-0.60 *** (0.06)
	r_f (fast component growth rate)	32.465 *** (3.751)	5.19 *** (1.47)	7.56 *** (1.33)	31.94 (48.41)	6.26 *** (1.18)
	a_s (slow component initial value)	-0.017 *** (0.008)	0.66 ** (0.03)	0.32 * (0.15)	0.03 (0.04)	-0.66 *** (0.03)
	r_s (slow component growth rate)	162.122 ** (69.910)	86.45 *** (5.92)	629.28 (423.13)	1.12 (3.53)	101.41 *** (8.35)
Habitual Exercisers	c (plateau)	-0.033 (0.014) p=0.475	0.68 (0.16) p=0.359	0.36 (0.18) p=0.362	-0.24 (0.14) p=0.754	-0.68 (0.21) p=0.341
	a_f (fast component initial value)	-0.078 [†] (0.009) p=0.032	0.56 [†] (0.07) p=0.009	0.55 [†] (0.04) P<0.001	-0.15 [†] (0.04) p<0.001	-0.831 [†] (0.07) p=0.002
	r_f (fast component growth rate)	11.628 [†] (3.929) p<0.001	6.78 (1.74) p=0.362	3.45 [†] (1.36) p=0.003	1.86 (48.41) p=0.534	6.76 (1.38) p=0.718
	a_s (slow component	-0.060 [†] (0.008)	0.36 [†] (0.03)	0.05 (0.15)	0.12 (0.07)	-0.40 [†] (0.03)

initial value)	*p<0.001*	*p<0.001*	*p=0.064*	*p=0.194*	*p<0.001*
r_s (slow component growth rate)	396.341 [†] (78.306) *p=0.003*	157.47 [†] (23.30) *p=0.002*	321.16 (579.39) *p=0.595*	433.6 (304.94) *p=0.157*	156.07 [†] (23.47) *p=0.20*
AIC	-106168.7	2021.38	-12838.49	-12350.91	4848.64
BIC	-106070.6	2111.25	-12748.6	-12261.04	4938.51
Log-likelihood	53096.36	-998.69	6431.24	6187.46	-2412.32
Number of observations	26336	13216	13216	13216	13216
Number of participants	32	32	32	32	32
SD: ID (*c*)	0.038	0.435	0.231	0.347	0.572
SD: Residual (*c*)	0.032	0.259	0.147	0.150	0.288

Note. The best-fitting model to the data for all five outcome measures was the two-
exponent model with a fixed effect on all variables by group. All models contain a random
plateau on participant. AIC = Akaike Information Criterion; BIC = Bayesian Information Criterion.
Values in parentheses are standard errors of the fixed effects. Asterisks indicate significant
effects where *p<0.05, **p<0.01, ***p<0.001. Daggers on coefficients in the Habitual Exercisers
group indicate coefficients that are significantly different between Habitual Exercisers and Non-
exercisers, and all group-difference p-values are shown.
254

Step Length Asymmetry

Compared to Non-exercisers, Habitual Exercisers adapted SLA more gradually. The

groups differed in every model-estimated coefficient except for SLA plateau. Overall, Habitual

Exercisers and Non-exercisers did not differ in initial SLA (Habitual Exercisers = -0.144, Non-

exercisers = -0.137, *t*(28.763)=0.283, *p*=0.779)). Habitual Exercisers adapted the fast

component of SLA more quickly than Non-exercisers did (Habitual Exercisers r_f = 12 steps,

Non-exercisers r_f = 32 steps), seen in the quicker achievement of the "elbow" by Habitual

Exercisers in Figure 2. Conversely, Habitual Exercisers adapted the slow component of SLA

more slowly than Non-exercisers did (Habitual Exercisers r_s = 396 steps, Non-exercisers r_s =

162 steps), seen in the gradual change in SLA after the elbow in Habitual Exercisers in Figure

2. Notably, the variability in SLA at each step between subjects appeared much larger across

Habitual Exercisers than across Non-exercisers (Fig. 2). For better understanding of SLA

adaptation, Figure 2B shows the averaged individual step lengths for fast and slow legs.

The model explained some of the variance in SLA plateau (Table 2, SD: Residual (c) =

0.032). Variation between participants accounted for a large portion of the variance in SLA

plateau (Table 2, SD: ID (c) = 0.038).

Figure 2. Step length asymmetry adapts differently over 823 steps of split-belt

treadmill walking. (A) Step length asymmetry adaptation. Shaded areas indicate the

standard deviation. Solid lines indicate the model fit to the data. Dashed lines indicate

the model-estimated plateau for each group. Blue=Habitual Exercisers(n=19),

orange=Non-exercisers (n=13). (B) Average step lengths for the fast and slow legs

during adaptation, for each group.

Positive Work Rate by the Fast Leg

Compared to Non-exercisers, Habitual Exercisers continued to gradually adapt positive

work rate by the fast leg over the entire trial. The groups differed in the initial values and the

slow-component growth rate (Figure 3A). Overall, Habitual Exercisers and Non-exercisers did

not differ in initial positive work rate by the fast leg (Habitual Exercisers = 1.44 W/kg, Non-

exercisers = 1.45 W/kg, t(23.484)=0.051, p=0.960). Habitual Exercisers adapted the slow

component of positive work rate by the fast leg more slowly than Non-exercisers did (Fig 3A)

(Habitual Exercisers r_s = 157 strides, Non-exercisers r_s = 86 strides).

The model explained much of the variance in fast-leg positive work rate plateau (Table

2, SD: Residual (c) = 0.259). Variation between participants accounted for a large portion of the

variance in fast-leg positive work rate plateau (Table 2, SD: ID (c) = 0.435).

Figure 3. Positive work rate by the legs on the fast (A) and slow (B) belts over 823

steps of split-belt treadmill walking. Shaded areas indicate the standard deviation of

each group. Solid lines indicate the model fit to the data. Dashed lines indicate the

model-estimated plateau for each group. Blue= Habitual Exercisers (*n*=19),

orange=Non-exercisers (*n*=13).

Negative Work Rate by the Fast Leg

Habitual Exercisers and Non-exercisers did not differ in adaptation of the negative work

rate by the fast leg. There were no significant adaptation in negative work rate by the fast leg, in

either group (Fig 4A).

The model explained much of the variance in fast-leg negative work rate plateau (Table

2, SD: Residual (*c*) = 0.150). Variation between participants accounted for a large portion of the

variance in fast-leg negative work rate plateau (Table 2, SD: ID (*c*) = 0.347).

Figure 4. Negative work rate by the legs on the fast (A) and slow (B) belts over 823

steps of split-belt treadmill walking. Shaded areas indicate the standard deviation of

each group. Solid lines indicate the model fit to the data. Dashed lines indicate the

model-estimated plateau for each group. Blue= Habitual Exercisers (*n*=19),

orange=Non-exercisers (*n*=13).

Positive Work Rate by the Slow Leg

Overall, Habitual Exercisers and Non-exercisers did not differ in initial positive work rate

by the slow leg (Habitual Exercisers = 0.701 W/kg, Non-exercisers = 0.722 W/kg,

t(23.462)=0.147, *p*=0.885). Habitual Exercisers adapted the fast component of positive work

rate by the slow leg more quickly than Non-exercisers did (Fig 3B) (Habitual Exercisers r_f = 3

strides, Non-exercisers r_f = 8 strides). Both groups converged on a plateau by about 10 strides,

and the groups did not differ in their estimated plateaus.

The model explained much of the variance in slow-leg positive work rate plateau (Table

2, SD: Residual (c) = 0.147). Variation between participants accounted for a large portion of the

variance in slow-leg positive work rate plateau (Table 2, SD: ID (c) = 0.231).

Negative Work Rate by the Slow Leg

Compared to Non-exercisers, Habitual Exercisers adapted the slow component of

negative work rate by the slow leg more quickly (Figure 4B). Overall, Habitual Exercisers and

Non-exercisers did not differ in initial negative work rate by the slow leg (Habitual Exercisers = -

1.664 W/kg, Non-exercisers = -1.57 W/kg, t(23.717)=0.248, p=0.806). Habitual Exercisers

adapted the slow component significantly slower (Habitual Exercisers r_s = 156 strides, Non-

exercisers r_s = 101 strides). The groups did not differ in their estimated plateaus.

The model explained much of the variance in slow-leg negative work rate plateau (Table

2, SD: Residual (c) = 0.288). Variation between participants accounted for a large portion of the

variance in slow-leg negative work rate plateau (Table 2, SD: ID (c) = 0.572).

Net work rate

The groups did not differ in the total net work rate by the legs (t(28.549) = -0.774, p =

0.445) (Fig. 5). Habitual Exercisers performed 0.166 (SD 0.072) W/kg and Non-exercisers

performed 0.188 (SD 0.085) W/kg at the end of ten minutes of split-belt walking.

Figure 5. Net Work rate by the legs. Orange=Non-exercisers (n=13), Blue= Habitual

Exercisers (n=19).

Discussion

The purpose of this study was to determine the effect of exercise on gait adaptation in

healthy young adults. Our main findings are: 1) Compared to Non-exercisers, Habitual

Exercisers more gradually adapted SLA, positive work rate by the fast leg, and negative work

rate by the slow leg over the entirety of split-belt exposure, 2) Habitual Exercisers initially

adapted SLA and positive work rate by the slow leg quicker than Non-exercisers, 3) Neither

group adapted the negative work rate by the fast leg, and 4) There was no effect of exercise on

net work done by the legs. The findings from this study support the hypothesis that young adults

who habitually exercise have a higher tolerance for the energetically-challenging asymmetric

belt speeds and adapt more gradually than those who do not.

Habitual Exercisers continued to gradually adapt over the entirety of split-belt exposure.

These current findings complement the literature. If the gradual component of gait adaptation is

driven by energetic cost optimization (13), then long-term involvement in regular exercise —

affecting exercise capacity — would affect the gradual component of gait adaptation, as seen

here in both SLA adaptation, and in the reduction in the positive work rate by the fast leg and

negative work rate by the slow leg. Exercise-trained adults have an enhanced exercise capacity

relative to non-exercisers, evidenced by higher maximal oxygen uptake (VO2max) (26); a better

exercise economy (26,27); and higher lactate and ventilatory thresholds(26). A higher exercise

capacity indicates that Habitual Exercisers can tolerate submaximal exercise — such as

treadmill walking — at a given energetic cost for longer than Non-exercisers, consistent with our

results. Habitual Exercisers and Non-exercisers can both reduce positive work by the fast leg

during split-belt adaptation, but Non-exercisers may have a direr need to do so quickly.

Therefore, Non-exercisers would be working at a higher energetic relative effort, resulting in a

stronger incentive to reduce energetic, and therefore mechanical, (12,14), work than the

Habitual Exercisers. Our results contradict the potential mechanism that Habitual Exercisers

would be quicker at reducing energetic cost than less active adults. Thus, the fact that adults

with more exercise sessions per week and more endurance training reach a minimum cost of

transport during running while those with less weekly exercise and less endurance training do

not ever reach a minimum cost of transport (25) does not apply to split-belt adaptation.

Therefore, habitual exercise may allow people to have a higher tolerance for the perturbation

induced by split-belt walking, allowing them to adapt their gait more gradually.

Habitual Exercisers initially adapted SLA and positive work rate by the slow leg quicker

than Non-exercisers. Our findings suggest that young adults who engage in more exercise are

quickly able to modify their gait in response to a continuous perturbation, possibly improving

stability (13) more quickly than those who do not exercise habitually. Indeed, athletes have been

shown to have better static balance than non-athletes (33). Buurke and colleagues investigated

the relationship between mediolateral stability adaptation and metabolic cost adaptation during

9 minutes of 2:1 split-belt walking (20). However, they did not find a relationship between

mediolateral stability and the reduction in metabolic cost. While their finding seems in contrast to

both our findings and the theory that initial rapid gait adaptations are intended to optimize

balance (13), all these findings taken together support a nuanced approach to motor skill

learning. One explanation, supported by the dual-rate model, is that adults prioritize balance

over metabolic cost when first perturbed, to meet the goals of the task (here, the assumed goal

being to stay upright and continue walking). Then, once balance is achieved, metabolic cost

becomes more relevant.

Neither group adapted the negative work rate by the fast leg. Negative work rate by the

fast leg during split-belt adaptation was largely unaffected by exercise. Previous work reported

that over forty-five minutes of adaptation at a 3:1 belt speed ratio, negative work rate by the fast

leg increased prior to positive work rate by the fast leg decreasing (12). Our finding of no

adaptation of negative work rate by the fast leg, by either group contrasts our hypothesis that

exercise capacity affects metabolic adaptation through increasing the negative work rate done

by the legs. A short adaptation period at a smaller belt speed ratio coupled with high variability

across participants could explain the lack of significant change in the negative work rate by the

fast leg in the current study, compared to previous work (12).

Both Habitual Exercisers and Non-exercisers performed net positive work by the end of

ten minutes of gait adaptation. Sánchez and colleagues reported that, at positive values of step

length asymmetry — achieved by enough time spent split-belt walking (12) — the legs perform

net negative work (14). This suggests that people are able to reduce overall positive work and

increase overall negative work performed. In contrast, our sample performed net positive, not

negative, work. The difference in findings between studies is likely due to the shorter adaptation

period (ten versus forty-five minutes) and the smaller split-belt speed ratio (2:1 Vs. 3:1) in the

current study, leading to negative average values of SLA at the end of adaptation. Given

enough time and/or a larger belt speed ratio, our sample would likely have plateaued at positive

SLA values and performed net negative work. Future research should explore if exercise affects

the ability to perform net negative work over prolonged adaptation periods at more extreme

split-belt perturbations.

There are some limitations to this study. First, participants held on to the handrails

during treadmill walking, which likely influenced both balance and mechanical work calculations.

While handrail holding was kept constant across participants and across groups, future studies

should investigate the effect of exercise on gait adaptation without holding on to handrails.

Second, we did not directly measure metabolic energetic cost using expired gas analyses. It is

possible that Habitual Exercisers may have modified SLA and mechanical work rate through

nonmetabolic processes, such as tendon stiffness, without modifying metabolic cost. Our

groupings based on aerobic exercise training – and not including resistance-trained adults –

reasonably allows us to infer that reported differences between groups would be due to

differences in energetic capacity. Future studies examining the metabolic adaptations between

Habitual Exercisers and Non-exercisers is a warranted next-step. Third, we standardized belt

speeds to participants' fastest comfortable walking speeds, which did not differ between groups.

This method of selecting the fast-belt speed has been used previously (34–36), and because

there was not a statistical difference between groups in initial SLA, we know that the groups

were not perturbed differently relative to their baseline, despite different belt speeds. Fourth, we

used self-report measures of exercise participation. Habitual exercise recall has a moderate

validity correlation to actual exercise (0.36) (37) and, despite its drawbacks, is the most

common way to assess physical activity (38–40). Additionally, the presence of group differences

in the current study suggests that self-report exercise habits were keen enough to separate

participants by energetic capacity that would affect gait adaptation. Finally, we did not test

retention of the adapted gait pattern, so we cannot comment on whether exercise affects young

adults' magnitude or persistence of errors during gait pattern washout.

Habitual exercise affects a young adult's response to a novel perturbation. Our results

suggest that people who regularly participate in aerobic exercise respond more quickly to the

onset of a perturbation and continue to explore strategies and search for assistance from the

environment in order to reduce energetic cost. The effect of aerobic exercise training on

movement adaptation likely acts through increasing aerobic capacity, thus reducing relative

effort of a given submaximal movement. On a split-belt treadmill, Habitual Exercisers may

require longer time than Non-exercisers to fully take advantage of the work done by the

treadmill; however, requiring a longer time to adapt is not inherently unfavorable. In fact, if

provided enough time to fine-tune their response to the perturbation, Habitual Exercisers may

plateau at a more-positive SLA, a more-negative net work done by the legs, and therefore

perhaps a lower energetic cost than Non-exercisers. The current results provide preliminary

evidence that Habitual Exercisers may be more able than Non-exercisers to use the work done

by the treadmill to adapt and reduce positive work done by the legs, given enough time.

Acknowledgements: We thank the individuals who participated in the study and the

undergraduate research assistants who assisted with data collection and processing. We also

thank Maxwell Donelan, PhD and Surabhi Simha PhD, for the MATLAB code on which we

based our code to calculate positive and negative work generated by the legs.

448 **Competing Interests:** No competing interests declared.

450 **Funding:** This work was supported by the Auburn University College of Education under Seed

451 Grant [JR18SG to J.A.R.].

References

1. Dietz V, Zijlstra W, Duysens J. Human neuronal interlimb coordination during split-belt locomotion. Exp Brain Res [Internet]. 1994 Oct [cited 2022 Jun 10];101(3). Available from: http://link.springer.com/10.1007/BF00227344

2. Reisman DS, Block HJ, Bastian AJ. Interlimb Coordination During Locomotion: What Can be Adapted and Stored? J Neurophysiol. 2005 Oct;94(4):2403–15.

3. Brinkerhoff SA, Monaghan PG, Roper JA. Adapting gait with asymmetric visual feedback affects deadaptation but not adaptation in healthy young adults. Barbieri FA, editor. PLOS ONE. 2021 Feb 25;16(2):e0247706.

4. Gonzalez-Rubio M, Velasquez NF, Torres-Oviedo G. Explicit Control of Step Timing During Split-Belt Walking Reveals Interdependent Recalibration of Movements in Space and Time. Front Hum Neurosci. 2019 Jul 3;13:207.

5. Long AW, Roemmich RT, Bastian AJ. Blocking trial-by-trial error correction does not interfere with motor learning in human walking. J Neurophysiol. 2016 May 1;115(5):2341–8.

6. Malone LA, Bastian AJ. Thinking About Walking: Effects of Conscious Correction Versus Distraction on Locomotor Adaptation. J Neurophysiol. 2010 Apr;103(4):1954–62.

7. Mariscal DM, Iturralde PA, Torres-Oviedo G. Altering attention to split-belt walking increases the generalization of motor memories across walking contexts. J Neurophysiol. 2020 Apr 1;123:11.

8. Smith MA, Ghazizadeh A, Shadmehr R. Interacting Adaptive Processes with Different Timescales Underlie Short-Term Motor Learning. Ashe J, editor. PLoS Biol. 2006 May 23;4(6):e179.

9. Darmohray DM, Jacobs JR, Marques HG, Carey MR. Spatial and Temporal Locomotor Learning in Mouse Cerebellum. Neuron. 2019 Apr;102(1):217-231.e4.

10. Mawase F, Haizler T, Bar-Haim S, Karniel A. Kinetic adaptation during locomotion on a split-belt treadmill. J Neurophysiol. 2013 Apr 15;109(8):2216–27.

11. Roemmich RT, Long AW, Bastian AJ. Seeing the Errors You Feel Enhances Locomotor Performance but Not Learning. Curr Biol. 2016 Oct;26(20):2707–16.

12. Sánchez N, Simha SN, Donelan JM, Finley JM. Using asymmetry to your advantage: learning to acquire and accept external assistance during prolonged split-belt walking. J Neurophysiol. 2021 Feb 1;125(2):344–57.

13. Seethapathi N, Clark B, Srinivasan M. Exploration-based learning of a step to step controller predicts locomotor adaptation [Internet]. Neuroscience; 2021 Mar [cited 2021 Aug 13]. Available from: http://biorxiv.org/lookup/doi/10.1101/2021.03.18.435986

14. Sánchez N, Simha SN, Donelan JM, Finley JM. Taking advantage of external mechanical
work to reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking.
J Physiol. 2019 Aug;597(15):4053–68.

15. Selgrade BP, Thajchayapong M, Lee GE, Toney ME, Chang YH. Changes in mechanical
work during neural adaptation to asymmetric locomotion. J Exp Biol. 2017 Jan
1;jeb.149450.

16. Donelan JM, Kram R, Kuo AD. Simultaneous positive and negative external mechanical
work in human walking. J Biomech. 2002;35(1):117–24.

17. Donelan JM, Kram R, Kuo AD. Mechanical work for step-to-step transitions is a major
determinant of the metabolic cost of human walking. J Exp Biol. 2002;205:3717–27.

18. Burdett RG, Skrinar GS, Simon SR. Comparison of mechanical work and metabolic energy
consumption during normal gait. J Orthop Res. 1983;1(1):63–72.

19. Umberger BR, Gerritsen KGM, Martin PE. A Model of Human Muscle Energy Expenditure.
Comput Methods Biomech Biomed Engin. 2003 May;6(2):99–111.

20. Buurke TJW, Lamoth CJC, Vervoort D. Adaptive control of dynamic balance in human gait
on a split-belt treadmill. J Exp Biol. 2018;10.

21. Finley JM, Bastian AJ, Gottschall JS. Learning to be economical: the energy cost of walking
tracks motor adaptation: Split-belt adaptation reduces metabolic power. J Physiol. 2013
Feb;591(4):1081–95.

22. Selinger JC, O'Connor SM, Wong JD, Donelan JM. Humans Can Continuously Optimize
Energetic Cost during Walking. Curr Biol. 2015 Sep;25(18):2452–6.

23. Park S, Finley JM. Manual stabilization reveals a transient role for balance control during
locomotor adaptation. J Neurophysiol. 2022 Oct 1;128(4):808–18.

24. Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, et al. The Physical
Activity Guidelines for Americans. JAMA. 2018 Nov 20;320(19):2020.

25. Cher PH, Stewart IB, Worringham CJ. Minimum Cost of Transport in Human Running Is Not
Ubiquitous. Med Sci Sports Exerc. 2015 Feb;47(2):307–14.

26. Jones AM, Carter H. The Effect of Endurance Training on Parameters of Aerobic Fitness:
Sports Med. 2000;29(6):373–86.

27. Morgan DW, Bransford DR, Costill DL, Daniels JT, Howley ET, Krahenbuhl GS. Variation in
the aerobic demand of running among trained and untrained subjects: Med Sci Sports
Exerc. 1995 Mar;27(3):404???409.

28. Dingwell JB, Marin LC. Kinematic variability and local dynamic stability of upper body
motions when walking at different speeds. J Biomech. 2006 Jan;39(3):444–52.

29. R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing [Internet]. Vienna, Austria; 2020. Available from: https://www.R-
project.org/

30. Pinhiero J, Bates D, DebRoy S, Sarkar D, R Core Team. nlme: Linear and Nonlinear Mixed
Effects Models. [Internet]. 2020. Available from: https://CRAN.R-project.org/package=nlme

31. Burnham KP, Anderson DR. Model Selection and Multimodel Inference: A Practical
Information-Theoretic Approach. J Wildl Manag. 2003 Jul;67(3):655.

32. Lohse KR, Shen J, Kozlowski AJ. Modeling Longitudinal Outcomes: A Contrast of Two
Methods. J Mot Learn Dev. 2020 Apr 1;8(1):145–65.

33. Cristina-Elena M, Liliana-Elisabeta R. Aspects Regarding the Level of Coordination Abilities
in Both Athletes and Non-athletes. Procedia - Soc Behav Sci. 2014 Mar;117:162–6.

34. Charalambous CC, Alcantara CC, French MA, Li X, Matt KS, Kim HE, et al. A single
exercise bout and locomotor learning after stroke: physiological, behavioural, and
computational outcomes: Effect of exercise on locomotor learning in chronic stroke
survivors. J Physiol. 2018 May 15;596(10):1999–2016.

35. Helm EE, Tyrell CM, Pohlig RT, Brady LD, Reisman DS. The presence of a single-
nucleotide polymorphism in the BDNF gene affects the rate of locomotor adaptation after
stroke. Exp Brain Res. 2016 Feb;234(2):341–51.

36. Hirata K, Hanawa H, Miyazawa T, Kubota K, Sonoo M, Kokubun T, et al. Adaptive changes
in foot placement for split-belt treadmill walking in individuals with stroke. J Electromyogr
Kinesiol. 2019 Oct;48:112–20.

37. Sallis JF, Saelens BE. Assessment of Physical Activity by Self-Report: Status, Limitations,
and Future Directions. Res Q Exerc Sport. 2000 Jun;71(sup2):1–14.

38. Castillo-Retamal M, Hinckson EA. Measuring physical activity and sedentary behaviour at
work: A review. Work. 2011;40(4):345–57.

39. Corder K, van Sluijs EM, Wright A, Whincup P, Wareham NJ, Ekelund U. Is it possible to
assess free-living physical activity and energy expenditure in young people by self-report?
Am J Clin Nutr. 2009 Mar 1;89(3):862–70.

40. Ishikawa-Takata K, Tabata I, Sasaki S, Rafamantanantsoa HH, Okazaki H, Okubo H, et al.
Physical activity level in healthy free-living Japanese estimated by doubly labelled water
method and International Physical Activity Questionnaire. Eur J Clin Nutr. 2008
Jul;62(7):885–91.

Supporting information captions

Methods S1. A detailed description of the statistical modeling analysis.

Table S2. Model f**its of mixed effects linear models for each outcome measure.**

Figure S3. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of step length asymmetry adaptation.

Figure S4. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the positive work rate of the fast leg.

Figure S5. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the negative work rate of the fast leg.

Figure S6. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the positive work rate of the slow leg.

Figure S7. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the positive work rate of the slow leg.

A

fast belt   slow belt

B

<u>Baseline</u>   <u>Adaptation</u>

Warmup self-selected typical speed

Warmup fast speed

Warmup slow speed

Belt speeds split 2:1 ratio

3 min   3 min   3 min   10 min