1

Habitual exercise evokes fast and persistent adaptation during split-belt walking

Sarah A. Brinkerhoff[1]*, Natalia Sánchez[2#], Jaimie A. Roper[1#]

[1]School of Kinesiology, Auburn University, Auburn, Alabama, United States of America

[2]Department of Physical Therapy, Chapman University, Irvine, California, United States of

America

*Corresponding Author

Email: sbrinkerhoff@auburn.edu

[#]NS and JAR are Joint Senior Authors

Abstract

Changing movement patterns in response to environmental perturbations is a critical aspect of gait and is related to reducing the energetic cost of the movement. Exercise improves energetic capacity for submaximal exercise and may affect how people adapt movement to reach an energetic minimum. The purpose of this study was to determine whether self-reported exercise behavior influences gait adaptation in young adults. Young adults who met the optimal volume of exercise according to the Physical Activity Guidelines for Americans (MOVE; n=19) and young adults who did not meet the optimal volume of exercise (notMOVE; n=13) walked on a split-belt treadmill with one belt moving twice the speed of the other belt for 10 minutes. Step length asymmetry (SLA) and mechanical work done by each leg were measured. Nonlinear mixed effects models compared the time course of adaptation between MOVE and notMOVE, and *t*-tests compared net work at the end of adaptation between MOVE and notMOVE. Compared to notMOVE, MOVE had a faster initial response to the split belt treadmill, and continued to adapt over the duration of split-belt treadmill walking. Young adults who engage in sufficient amounts of exercise responded more quickly to the onset of a perturbation, and throughout the perturbation they continued to explore movement strategies, which might be related to reduction of energetic cost. Our findings provide insights into the multisystem positive effects of exercise, including walking adaptation.

Introduction

Adapting walking patterns in response to environmental perturbations is a critical aspect of locomotion. When a person encounters a perturbation, such as when walking on an icy or uneven surface, they must adapt their walking patterns to avoid falling which can be achieved using different strategies. While prior research suggests that visual feedback [1,2], focus of attention [3,4], and neurological injury [5–7] can affect aspects of walking adaptation, how or if individual factors related to overall physical activity might influence walking adaptation strategies needs further evidence.

A common approach to study walking adaptation is split-belt treadmill walking, in which the belts under each leg move at different speeds [8,9]. The changes in the asymmetry between left and right step lengths, or step length asymmetry (SLA) is one measure used to track how a person's gait pattern adapts in response to a continuous perturbation. As a robust measure of gait adaptation, SLA is observable with the unaided eye, is sensitive to experimental manipulations, and persists even after the split-belt perturbation is removed [1–4,10]. In line with upper extremity motor adaptation [11], adaptation of SLA during split-belt walking occurs at two distinct and interacting timescales – a fast component that adapts rapidly and a slow component that adapts more gradually [12–15]. From work in both upper and lower extremity motor tasks, the two timescales of motor adaptation may derive from two separate (but not necessarily independent) processes.

Indeed, Seethapathi and colleagues posited that the component that adapts rapidly is driven by balance optimization, whereas the component that adapts more gradually is driven by energetic cost optimization [16]. Consistent with these findings, experimental studies have shown that SLA adaptation during split-belt walking occurs in parallel with reductions in the work generated by the legs; gait adaptation results in decreasing positive work and increasing negative work done by the legs, especially the leg on the fast belt [17,18]. As the legs reduce

positive work done and increase negative work done, energetic cost decreases concomitantly

[17,19,20]. This complements the fact that doing negative work is less energetically costly than

doing positive work [21,22]. By reducing work done by the legs, people are likely gradually

adapting towards some lower energetic cost, as determined by Seethapathi et. al. [16] and

previously empirically suggested [23–25]. Similarly, a study by Park et al showed that during

early adaptation to split belt walking, individuals increase whole body angular momentum, which

is inversely proportional to balance [26]. Therefore, these experimental findings also support the

idea of balance and energetic cost as the two distinct processes driving gait adaptation.

The question then arises — do individual factors that affect energy consumption affect

the ability to adapt gait during split-belt walking? The multisystem benefits of exercise are well-

known and include improved physical function, cognition, quality of life, and reduced risk of

cardiovascular disease and all-cause mortality [27]. Moreover, the ability to move at a minimum

energetic cost of transport at a given speed is a function of aerobic training experience, such

that people with more aerobic training reach a minimum cost of transport while running, but

those who engage in less aerobic training do not [28]. Habitual physical activity is generally

known to improve aerobic capacity [29,30]. Exercise-induced adaptations would increase

capacity for exercise and enable trained individuals to tolerate submaximal exercise for longer.

If we posit that people adapt to split-belt walking in order to reduce energetic cost, then those

who engage in more exercise may reach an energetic optimum quicker than those who engage

in less exercise [28]; conversely, those who engage in more exercise may more gradually reach

an energetic optimum, due to a higher tolerance for submaximal exercise [31,32]. These two

hypotheses indicate that exercise habits might influence adaptation toward more energetically

economical movement patterns.

The purpose of this study was to determine whether self-reported exercise behavior

influences gait adaptation in young adults. We hypothesized that amount of self-reported

exercise would affect gait adaptation, and this effect would primarily be driven by differences in

the slow component of adaptation, given the role of energetics in shaping the rate of adaptation

of the slow component [16,23–25] and the well-established effects of exercise on energetics

[28,31,32]. Our hypothesis is based on two opposing mechanisms which we will test here: 1)

Young adults engaging in sufficient weekly exercise (Meets Optimal Volume of Exercise;

MOVE) would adapt faster than those engaging in low or no weekly exercise (does not Meet

Optimal Volume of Exercise; notMOVE) to reach a minimum energetic cost quickly [28],

evidenced by faster reduction in SLA and work done by the legs; 2) MOVE would have a higher

tolerance for the submaximal exercise of treadmill walking than notMOVE, evidenced by more

gradual reduction in SLA and work done by the legs. To test these competing mechanisms, we

analyzed the adaptation of SLA and the positive and negative work done by each leg while

MOVE and notMOVE walked on a split-belt treadmill. We also hypothesized that at the end of

adaptation, MOVE would perform more-negative net work than notMOVE, by reducing positive

work done by the legs and increasing negative work done by the legs in order to reduce overall

energetic cost [17,19,20]. A significant effect of regular exercise participation on gait adaptation

would support the idea that individual factors that influence energetics can also influence

walking adaptation. Our findings will provide insights into the multisystem positive effects of

exercise, including walking adaptation, and will provide directions for exercise rehabilitation

research.

Methods

Participants

We recruited a convenience sample of young adults ages 19-35 for participation in this

study. Participants were excluded if they reported cardiovascular, pulmonary, renal, metabolic,

vestibular, or neurologic disorders; any lower-extremity injuries or surgeries in the past 12

months; or a prior anterior cruciate ligament injury. The Auburn University Institutional Review

Board approved all procedures and all participants completed written informed consent prior to

participating (Protocol #18-418 MR 1811). Data were collected in the Locomotor and

Movement Control Lab at Auburn University.

Experimental Protocol

Participants first completed a self-report exercise behavior questionnaire that consisted

of a modified Godin-Leisure Time Exercise Questionnaire [33] and a custom survey on sport

and exercise modalities. This questionnaire asked how many times per week participants

currently exercised at a moderate-to-vigorous intensity, for how many minutes per session, and

for how many consecutive weeks. Based on their responses, participants were grouped into one

of two groups according to The Physical Activity Guidelines for Americans: MOVE (at least 150

minutes per week of exercise for at least the last three months) or notMOVE (less than 150

minutes per week of exercise for at least the last three months) [27]. The questionnaire also

asked what types of exercise participants were currently engaged in. For this study, we

excluded participants who engaged only in weight training or flexibility exercise, in an attempt to

bias the sample towards aerobic-based activities.

After completing the questionnaire, participants walked on the treadmill. Participants

held onto side handrails for the entirety of treadmill walking. The instructions given to

participants noted that they should use the handrails for stability, and not to offload bodyweight.

First, we determined participants' typical and fastest comfortable walking speeds on the

treadmill using a modified protocol by Dingwell and Marin [34]. Starting at 0.6 m/s, the treadmill

incrementally increased by 0.05 m/s every 4 seconds until participants reported that the current

speed was their typical walking speed, or was their fastest comfortable speed. This was

repeated twice for typical walking, and twice for fast walking. We instructed participants either,

"tell me when you reach your typical walking speed," or, "tell me when you reach the fastest

speed that you'd be comfortable walking for ten minutes." The fastest comfortable speed was

set as the fast belt velocity, and the slow belt velocity was set as half of the fastest comfortable

speed.

The split-belt walking protocol is shown in Figure 1. After finding their typical and fast

speeds, participants warmed up to the treadmill and were familiarized to the belt speeds first by

walking for three minutes at their typical walking speed, second by walking for three minutes at

their fastest comfortable speed, and third by walking for three minutes at half of the fast speed

(slow speed). The last tied walking condition — three minutes of walking at the slow speed —

was considered the baseline condition for data analysis. We did not randomize the order of the

warmup trials, to ensure that the baseline condition was the same across participants. Then,

participants walked for ten minutes with the belt speeds split — the belt under the dominant leg

moved at participants' fastest comfortable speed, and the belt under the non-dominant leg

moved at the slow speed. Leg dominance was determined as the leg that a participant reported

they would use to kick a ball.

Figure 1. Split-belt walking protocol. A) An oblique and birds-eye view of the split-belt

treadmill. B) First, participants warmed up to the treadmill and were familiarized to the

belt speeds first by walking for three minutes at their typical walking speed, second by

walking for three minutes at the fast speed, and third by walking for three minutes at the

slow speed. Finally, they walked for 10 minutes with the belt under their dominant leg at

the fast speed, and the belt under their non-dominant leg at the slow speed.

Data Analysis

Kinematic data were recorded at 100 Hz from reflective markers placed bilaterally on the

anterior superior iliac spine and the lateral malleoli of the ankles, using a 17-camera motion

capture system (VICON; Vicon Motion Systems Ltd, Oxford, United Kingdom). Kinematic data

were lowpass filtered with a 4th order Butterworth filter with a cutoff frequency of 6 Hz. Ground

reaction force data were obtained for each individual leg using an instrumented split-belt

treadmill, recorded at 1000 Hz from two separate force plates. Force data were lowpass filtered

with a 4th order Butterworth filter with a cutoff frequency of 20 Hz.

Step length was calculated as the distance between the ankle markers along the

anterior-posterior walking axis at foot strike. SLA was calculated and normalized to stride length

as in eq. 1.

$$Step\ length\ Asymmetry = \frac{Step\ length_{fast} - Step\ length_{slow}}{Step\ length_{fast} + Step\ length_{slow}} \qquad 1$$

Here, *Step Length$_{fast}$* is the step length when the leg on the fast belt strikes the belt, and

Step Length$_{slow}$ is the step length when the leg on the slow belt strikes the belt. A negative SLA

indicates that the leg on the slow belt is taking a longer step than the leg on the fast belt, and an

SLA of zero indicates that the legs are taking steps of equal length.

We calculated positive and negative mechanical work generated by the legs across each

stride cycle using a custom MATLAB program. A stride cycle was calculated as the time

between ipsilateral foot-strikes. We used the point-mass model to estimate mechanical work

generated by the legs on the treadmill and on the center of mass [15,17,18]. In brief, we

segmented force data into strides [18] and calculated the center of mass velocities in each

direction as the time integral of the center of mass accelerations. Next, we calculated the

instantaneous power generated by each leg for each stride as the instantaneous sum of the dot

product of the ground reaction force and the center of mass velocity and the dot product of the

force applied to the respective belt and the belt speed. We then calculated the total positive and

total negative work performed by each leg as the time integral of the positive or negative portion

of the total instantaneous power over the stride cycle. We calculated work rate by dividing each

work measure by stride duration. Finally, we calculated the net work rate by the legs at the end

of adaptation as the time integral of the power divided by stride cycle, averaged over the last

100 strides.

Statistical Analysis

All statistical analyses were conducted in R [35]. We used Welch Two Sample t-tests to

compare age, height, leg length, mass, and fast-leg belt speed between groups. We also used

Pearson's Chi-squared tests with Yates' continuity correction to assess the difference in

proportion of males and females in each group, and to assess the difference in proportion of

people who walked with the right leg on the fast belt in each group. We analyzed SLA, positive

and negative work rate by the fast legs, and positive and negative work rate by the slow leg

using mixed effects nonlinear regression models with the 'nlme' package [36]. We truncated the

data for all participants to minimum number of steps taken by a participant (823 steps) so that

all participants would weight equally on each step.

Detailed description of the statistical models can be found in the supporting information.

In brief, for each of the work rate outcome measures and for SLA, we built-up models to two-

exponent models — assuming that adaptation of the outcome measures occurred over two

timescales — that included a fixed effect for group (eqs. 2 and 3). Here, c is the estimated

plateau if strides went to infinity; a_f is the initial value of the fast component of adaptation; r_f is

the growth rate of the fast component of adaptation; a_s is the initial value of the slow component

of adaptation; r_s is the growth rate of the slow component of adaptation. In the models that

included a fixed effect for group, all five model parameters (c, a_f, r_f, a_s, r_s) were allowed to differ

for MOVE vs. notMOVE.

222
$$SLA = (c + a_f * e^{-\frac{step}{r_f}} + a_s * e^{-\frac{step}{r_s}}) \sim Group + (c|ID) \qquad 2$$

223
$$Work\ rate = \left(c + a_f * e^{-\frac{stride}{r_f}} + a_s * e^{-\frac{stride}{r_s}}\right) \sim Group + (c|ID) \qquad 3$$

224  We compared these two-exponent models with fixed effects for group to models without

225 fixed effects for group, and to one-exponent models assuming adaptation occurs over a single

226 timescale. All models and outcomes can be found in the supporting information All models were

227 fitted with a maximum likelihood estimation and all models contained a random effect such that

228 the outcome measure's estimated plateau was allowed to vary by participant. The simplest,

229 best-fitting model for each measure was the final model that we interpreted, such that a lower

230 Akaike Information Criterion (AIC) and Bayesian Information Criteria (BIC) indicated a better fit

231 to the data and log-likelihood tests assessed goodness of fit to the data [37,38]. If the final

232 model included a fixed effect for group, the significance (or lack thereof) of the effect of group on

233 the equation variables in the best-fitting models was evaluated using Welch's t-test with an *a*

234 *priori* alpha level of 0.05. If the final model included a fixed effect for group, initial adaptation

235 values were calculated for each variable as the average over the first five strides and were

236 compared between group using Welch's Two Sample t-test.

237  We used Welch's Two Sample t-test between groups to compare the net work rate by

238 the legs at the end of adaptation, to determine if there was a difference between exercise

239 groups in the ability to gain assistance from the treadmill.

Results

Participants

243  Thirty-seven people participated in this study. We excluded five participants who

244 engaged in weightlifting, yoga, and scuba diving, in an attempt to bias the sample towards

245 aerobic-based activities. Therefore, 32 participants were included in these analyses (Table 1).

246 Based on the self-report exercise questionnaire, 19 participants were placed in the MOVE group

247 and 13 were placed in the notMOVE group. The two groups did not significantly differ by sex,

height, mass, leg on the fast belt, or fast belt speed, but did significantly differ by age where the

MOVE group was slightly older. Demographic and belt speed data are provided in Table 1.

Table 1. Participant demographics in Mean [range].

	notMOVE	MOVE	p-value	Total
n	13	19	—	32
Age (yr)	20 [19,22]	22 [19,32]	**0.045**	22 [19,32]
Height (cm)	1706 [1540,1910]	1727 [1540,1920]	0.565	1718 [1540,1920]
Mass (kg)	66.5 [48,89]	72.9 [50,116]	0.208	70.3 [48,116]
Fast belt speed (m/s)	1.57 [1.25,1.95]	1.49 [1.15,1.98]	0.231	1.52 [1.15,1.98]
Minutes/week exercise	38 [0,120]	268 [150,750]	**<0.001**	175 [0,750]
Sex	11 female, 2 male	11 female, 8 male	0.225	22 female, 10 male
Fast leg	3 right, 10 left	3 right, 16 left	0.954	6 right, 26 left
Exercise modality	run, cycle, basketball, tennis	run, cycle, basketball, tennis, CrossFit®, dance, boxing	—	—

Note. Type of exercise participated in reflects the exercise modalities of people who participated in any exercise every week (regardless of minutes per week of exercise).

General Timescales of Adaptation

The best-fitting model for each outcome measure was the two-exponent model

containing a fixed effect for group (eqs. 2 and 3, Table 2), indicating that SLA and work rate by

the legs adapted over two distinct timescales and was affected by exercise. All candidate

models for each outcome measure, and the final model fits to each participant's data, can be

found in the supporting information. AIC, BIC, and log-likelihood tests for goodness of fit agreed

for all outcome measures.

Table 2. Final mixed effects models for SLA, positive and negative work of the fast leg, and positive and negative work of the slow leg.

Coefficient	SLA	Positive Work Rate		Negative Work Rate	
		Fast Leg	Slow Leg	Fast Leg	Slow Leg
c (plateau)	-0.043 ***	0.54 ***	0.20	-0.28 **	-0.48 **
	(0.010)	(0.12)	(0.17)	(0.10)	(0.16)
a_f (fast component	-0.097 ***	0.38 ***	0.28 ***	0.02	-0.60 ***

initial value)	(0.007)	(0.06)	(0.03)	(0.02)	(0.06)
r_f (fast component growth rate)	32.465 *** (3.751)	5.19 *** (1.47)	7.56 *** (1.33)	31.94 (48.41)	6.26 *** (1.18)
a_s (slow component initial value)	-0.017 *** (0.008)	0.66 ** (0.03)	0.32 * (0.15)	0.03 (0.04)	-0.66 *** (0.03)
r_s (slow component growth rate)	162.122 ** (69.910)	86.45 *** (5.92)	629.28 (423.13)	1.12 (3.53)	101.41 *** (8.35)
MOVE					
c (plateau)	-0.033 (0.014) p=0.475	0.68 (0.16) p=0.359	0.36 (0.18) p=0.362	-0.24 (0.14) p=0.754	-0.68 (0.21) p=0.341
a_f (fast component initial value)	-0.078 † (0.009) p=0.032	0.56 † (0.07) p=0.009	0.55 † (0.04) P<0.001	-0.15 † (0.04) p<0.001	-0.831 † (0.07) p=0.002
r_f (fast component growth rate)	11.628 † (3.929) p<0.001	6.78 (1.74) p=0.362	3.45 † (1.36) p=0.003	1.86 (48.41) p=0.534	6.76 (1.38) p=0.718
a_s (slow component initial value)	-0.060 † (0.008) p<0.001	0.36 † (0.03) p<0.001	0.05 (0.15) p=0.064	0.12 (0.07) p=0.194	-0.40 † (0.03) p<0.001
r_s (slow component growth rate)	396.341 † (78.306) p=0.003	157.47 † (23.30) p=0.002	321.16 (579.39) p=0.595	433.6 (304.94) p=0.157	156.07 † (23.47) p=0.20
AIC	-106168.7	2021.38	-12838.49	-12350.91	4848.64
BIC	-106070.6	2111.25	-12748.6	-12261.04	4938.51
Log-likelihood	53096.36	-998.69	6431.24	6187.46	-2412.32
Number of observations	26336	13216	13216	13216	13216
Number of participants	32	32	32	32	32
SD: ID (c)	0.038	0.435	0.231	0.347	0.572
SD: Residual (c)	0.032	0.259	0.147	0.150	0.288

Note. Model parameters are given as Coefficient (SEM), *p*-value. The best-fitting model to the data for all five outcome measures was the two-exponent model with a fixed effect on all variables by group. All models contain a random plateau on participant. AIC = Akaike Information Criterion; BIC = Bayesian Information Criterion. Values in parentheses are standard errors of the fixed effects. Asterisks indicate significant effects where *p<0.05, **p<0.01, ***p<0.001. Daggers on coefficients in the MOVE group indicate coefficients that are significantly different between MOVE and notMOVE, and all group-difference p-values are shown.

Step Length Asymmetry

Compared to notMOVE, MOVE adapted SLA more gradually. The groups differed in every model-estimated coefficient except for SLA plateau. Overall, MOVE and notMOVE did not differ in initial SLA (MOVE = -0.144, notMOVE = -0.137, $t(28.763)=0.283$, $p=0.779$)). MOVE adapted the fast component of SLA more quickly than notMOVE (MOVE r_f = 12 steps, notMOVE r_f = 32 steps), seen in the quicker achievement of the "elbow" by MOVE in Figure 2. Conversely, MOVE adapted the slow component of SLA more slowly than notMOVE (MOVE r_s = 396 steps, notMOVE r_s = 162 steps), seen in the gradual change in SLA after the elbow in Figure 2. Notably, the variability in SLA at each step between subjects appeared much larger across MOVE than across notMOVE (Fig. 2). For better understanding of SLA adaptation, Figure 2B shows the averaged individual step lengths for fast and slow legs.

The model explained some of the variance in SLA plateau (Table 2, SD: Residual (c) = 0.032). Variation between participants accounted for a large portion of the variance in SLA plateau (Table 2, SD: ID (c) = 0.038).

Figure 2. Step length asymmetry adapts differently over 823 steps of split-belt treadmill walking. (A) Step length asymmetry adaptation. Shaded areas indicate the standard deviation. Solid lines indicate the model fit to the data. Dashed lines indicate the model-estimated plateau for each group. Blue= MOVE (n=19), orange= notMOVE (n=13). (B) Average step lengths for the fast and slow legs during adaptation, for each group.

Positive Work Rate by the Fast Leg

Compared to notMOVE, MOVE continued to gradually adapt positive work rate by the fast leg over the entire trial. The groups differed in the initial values and the slow-component

growth rate (Figure 3A). Overall, MOVE and notMOVE did not differ in initial positive work rate

by the fast leg (MOVE = 1.44 W/kg, notMOVE = 1.45 W/kg, $t(23.484)=0.051$, $p=0.960$). MOVE

adapted the slow component of positive work rate by the fast leg more slowly than notMOVE did

(Fig 3A) (MOVE r_s = 157 strides, notMOVE r_s = 86 strides).

The model explained much of the variance in fast-leg positive work rate plateau (Table

2, SD: Residual (c) = 0.259). Variation between participants accounted for a large portion of the

variance in fast-leg positive work rate plateau (Table 2, SD: ID (c) = 0.435).

Figure 3. Positive work rate by the legs on the fast (A) and slow (B) belts over 823

steps of split-belt treadmill walking. Shaded areas indicate the standard deviation of

each group. Solid lines indicate the model fit to the data. Dashed lines indicate the

model-estimated plateau for each group. Blue= MOVE (n=19), orange= notMOVE

(n=13).

313   Negative Work Rate by the Fast Leg

MOVE and notMOVE did not differ in adaptation of the negative work rate by the fast

leg. There was no significant adaptation in negative work rate by the fast leg, in either group

(Fig 4A).

The model explained much of the variance in fast-leg negative work rate plateau (Table

2, SD: Residual (c) = 0.150). Variation between participants accounted for a large portion of the

variance in fast-leg negative work rate plateau (Table 2, SD: ID (c) = 0.347).

Figure 4. Negative work rate by the legs on the fast (A) and slow (B) belts over 823

steps of split-belt treadmill walking. Shaded areas indicate the standard deviation of

each group. Solid lines indicate the model fit to the data. Dashed lines indicate the

model-estimated plateau for each group. Blue= MOVE (*n*=19), orange= notMOVE

(*n*=13).

Positive Work Rate by the Slow Leg

Overall, MOVE and notMOVE did not differ in initial positive work rate by the slow leg

(MOVE = 0.701 W/kg, notMOVE = 0.722 W/kg, *t*(23.462)=0.147, *p*=0.885). MOVE adapted the

fast component of positive work rate by the slow leg more quickly than notMOVE did (Fig 3B)

(MOVE r_f = 3 strides, notMOVE r_f = 8 strides). Both groups converged on a plateau by about 10

strides, and the groups did not differ in their estimated plateaus.

The model explained much of the variance in slow-leg positive work rate plateau (Table

2, SD: Residual (*c*) = 0.147). Variation between participants accounted for a large portion of the

variance in slow-leg positive work rate plateau (Table 2, SD: ID (*c*) = 0.231).

Negative Work Rate by the Slow Leg

Compared to notMOVE, MOVE adapted the slow component of negative work rate by

the slow leg more gradually (Figure 4B). Overall, MOVE and notMOVE did not differ in initial

negative work rate by the slow leg (MOVE = -1.664 W/kg, notMOVE = -1.57 W/kg,

t(23.717)=0.248, *p*=0.806). MOVE adapted the slow component significantly slower (MOVE r_s =

156 strides, notMOVE r_s = 101 strides). The groups did not differ in their estimated plateaus.

The model explained much of the variance in slow-leg negative work rate plateau (Table

2, SD: Residual (*c*) = 0.288). Variation between participants accounted for a large portion of the

variance in slow-leg negative work rate plateau (Table 2, SD: ID (*c*) = 0.572).

Net work rate

346   The groups did not differ in the total net work rate by the legs (t(28.549) = -0.774, p =

347   0.445) (Fig. 5). MOVE performed 0.166 (SD 0.072) W/kg and notMOVE performed 0.188 (SD

348   0.085) W/kg at the end of ten minutes of split-belt walking.

350   **Figure 5. Net Work rate by the legs.** Blue= MOVE (n=19), Orange= notMOVE (n=13).

Discussion

The purpose of this study was to determine whether self-reported exercise behavior

influences gait adaptation in young adults. Our main findings are: 1) Compared to notMOVE,

MOVE more gradually adapted SLA, positive work rate by the fast leg, and negative work rate

by the slow leg over the entirety of split-belt exposure, 2) MOVE initially adapted SLA and

positive work rate by the slow leg quicker than notMOVE, 3) Neither group adapted the negative

work rate by the fast leg, and 4) There was no effect of exercise on net work done by the legs.

The findings from this study support the hypothesis that young adults who engage in sufficient

amounts of weekly exercise have a higher tolerance for the energetically-challenging

asymmetric belt speeds and adapt more gradually than those who do not.

MOVE continued to gradually adapt over the entirety of split-belt exposure. These

current findings complement the literature. If the gradual component of gait adaptation is driven

by energetic cost optimization [16], then long-term involvement in regular exercise — affecting

exercise capacity — would affect the gradual component of gait adaptation, as seen here in

both SLA adaptation and in the reduction in the positive work rate by the fast leg and negative

work rate by the slow leg. Exercise-trained adults have an enhanced exercise capacity relative

to non-exercisers, evidenced by higher maximal oxygen uptake (VO2max) [31]; a better

exercise economy [31,32]; and higher lactate and ventilatory thresholds [31]. A higher exercise

capacity indicates that MOVE can tolerate submaximal exercise — such as treadmill walking —

at a given energetic cost for longer than notMOVE, consistent with our results. Our results

suggest that young adults, regardless of exercise habits, can reduce positive work by the fast

leg during split-belt adaptation, but that the amount of exercise practiced weekly and regularly

affects the rate at which positive work rate reduction occurs. Those engaging in suboptimal

amounts of exercise (notMOVE) may have a direr need to reduce positive work by the leg than

those who engage in sufficient amounts of exercise (MOVE), because they may be working at a

higher relative effort, resulting in a stronger incentive to reduce energetic and therefore

mechanical work [15,17]. Our results contradict the potential mechanism that MOVE would be

quicker at reducing energetic cost than notMOVE. While adults who engage in more exercise

per week do reach a minimum cost of transport during running, unlike those engaging in less

exercise [28], our results suggest that adults who engage in more exercise per week do not

have an advantage in reaching a minimum energetic cost during split-belt adaptation. Therefore,

high amounts of regular weekly exercise may allow people to have a higher tolerance for the

perturbation induced by split-belt walking, allowing them to adapt their gait more gradually.

The MOVE group initially adapted SLA and positive work rate by the slow leg quicker

than the notMOVE group. Our findings suggest that young adults who engage in more exercise

are quickly able to modify their gait in response to a continuous perturbation, possibly improving

stability [16] more quickly than those who engage in less exercise. Indeed, athletes perform

better than non-athletes on a balance task after a learning block of multimodal balance training

[39]. Buurke and colleagues investigated the relationship between mediolateral stability

adaptation and metabolic cost adaptation during 9 minutes of 2:1 split-belt walking [23].

However, they did not find a relationship between mediolateral stability and the reduction in

metabolic cost. While their finding seems in contrast to both our findings and the theory that

initial rapid gait adaptations are intended to optimize balance [16], all these findings taken

together support a nuanced approach to motor skill learning. One explanation, supported by the

two-timescale model, is that adults prioritize balance over metabolic cost when first perturbed, to

meet the goals of the task (here, the assumed goal being to stay upright and continue walking).

Then, once balance is achieved, metabolic cost becomes more relevant.

Neither group adapted the negative work rate by the fast leg. Negative work rate by the

fast leg during split-belt adaptation was largely unaffected by exercise. Previous work reported

that over forty-five minutes of adaptation at a 3:1 belt speed ratio, negative work rate by the fast

leg increased prior to positive work rate by the fast leg decreasing [15]. Our finding of no

adaptation of negative work rate by the fast leg, by either group contrasts our hypothesis that

exercise capacity affects metabolic adaptation through increasing the negative work rate done

by the legs. A short adaptation period at a smaller belt speed ratio coupled with high variability

across participants could explain the lack of significant change in the negative work rate by the

fast leg in the current study, compared to previous work [15].

Both the MOVE and notMOVE groups performed net positive work by the end of ten

minutes of gait adaptation. Sánchez and colleagues reported that, at positive values of SLA —

achieved by enough time spent split-belt walking [15] — the legs perform net negative work [17].

This suggests that people are able to reduce overall positive work and increase overall negative

work performed. In contrast, our sample performed net positive, not negative, work. The

difference in findings between studies is likely due to the shorter adaptation period (ten versus

forty-five minutes) and the smaller split-belt speed ratio (2:1 Vs. 3:1) in the current study,

leading to negative average values of SLA at the end of adaptation. Given enough time and/or a

larger belt speed ratio, our sample would likely have plateaued at positive SLA values and

performed net negative work. Future research should explore if exercise affects the ability to

perform net negative work over prolonged adaptation periods at more extreme split-belt

perturbations.

Variability in SLA and positive work by the fast leg appeared larger in the MOVE group

than in the notMOVE group. Abrams and colleagues determined that during 45 minutes of 3:1

split-belt treadmill walking, young adults reduced the variability in SLA over the course of split-

belt treadmill walking (Abram et al., 2022). If the MOVE group had a larger tolerance for

submaximal exercise than did the notMOVE group, then they may also have had a larger

window of tolerance for movement variability during initial adaptation. To determine if active the

MOVE group was indeed more variable than did the notMOVE group, we conducted follow-up

analyses of the within-participant SLA and positive fast-leg work rate variability [40]. We

calculated the standard deviation of each measure in the initial 100 strides, middle 100 strides,

and final 100 strides. Separate student's two-sample t-tests compared variability of each

measure during the initial, middle, and final epochs of gait adaptation between the active and

inactive groups. There was no difference between groups in variability in either measure during

any epoch. Therefore, our sample does not support the theory that adults who exercise more

would employ a more varied hypothesis testing strategy during early adaptation during ten

minutes of split-belt treadmill walking. The larger variation in the standard error of the MOVE

group in SLA and positive fast-leg work rate is therefore likely because the individual responses

to the perturbation within the MOVE group is varied, while individual responses in the notMOVE

group were more homogenous. Further individual factors that can explain the variation in active

individuals' response should be investigated.

There are some limitations to this study. First, participants held on to the handrails

during treadmill walking, which likely influenced both balance and mechanical work calculations.

While handrail holding was kept constant across participants and across groups, future studies

should investigate the effect of exercise on gait adaptation without holding on to handrails.

Second, we did not directly measure metabolic energetic cost using expired gas analyses. It is

possible that the MOVE group may have modified SLA and mechanical work rate through

nonmetabolic processes, such as tendon stiffness, without modifying metabolic cost. Our

groupings based on self-reported exercise training and not including resistance-trained adults

reasonably allows us to infer that reported differences between groups would be due to

differences in energetic capacity. Future studies examining the metabolic adaptations between

those who meet the exercise recommendations and those who do not is a warranted next-step.

Third, we standardized belt speeds to participants' fastest comfortable walking speeds, which

did not differ between groups. This method of selecting the fast-belt speed has been used

previously [41–43], and because there was not a statistical difference between groups in initial

SLA, we know that the groups were not perturbed differently relative to their baseline, despite

different belt speeds. Fourth, we used self-report measures of exercise participation. Habitual

exercise recall has a moderate validity correlation to actual exercise (0.36) [44] and, despite its

drawbacks, is the most common way to assess physical activity [45–47]. Additionally, the

presence of group differences in the current study suggests that self-report exercise habits were

keen enough to separate participants by energetic capacity that would affect gait adaptation,

though it is possible that participants practiced other exercise modalities that they did not

disclose. Finally, we did not test retention of the adapted gait pattern, so we cannot comment on

whether exercise affects young adults' magnitude or persistence of errors during gait pattern

washout.

The amount of self-reported habitual exercise affects a young adult's response to a

novel perturbation. Our results suggest that people who regularly participate in sufficient

exercise respond more quickly to the onset of a perturbation and continue to explore strategies

and search for assistance from the environment in order to reduce energetic cost. The effect of

exercise training on movement adaptation likely acts through increasing aerobic capacity, thus

reducing relative effort of a given submaximal movement. On a split-belt treadmill, MOVE may

require longer time than notMOVE to fully take advantage of the work done by the treadmill;

however, requiring a longer time to adapt is not inherently unfavorable. In fact, if provided

enough time to fine-tune their response to the perturbation, MOVE may plateau at a more-

positive SLA, a more-negative net work done by the legs, and therefore perhaps a lower

energetic cost than notMOVE. The current results provide preliminary evidence that MOVE may

be more able than notMOVE to use the work done by the treadmill to adapt and reduce positive

work done by the legs, given enough time.

Acknowledgements

478 We thank the individuals who participated in the study and the undergraduate research

479 assistants who assisted with data collection and processing. We also thank Maxwell Donelan,

480 PhD and Surabhi Simha PhD, for the MATLAB code on which we based our code to calculate

481 positive and negative work generated by the legs.

Competing Interests

484 No competing interests declared.

References

[1]   Brinkerhoff SA, Monaghan PG, Roper JA. Adapting gait with asymmetric visual feedback

affects deadaptation but not adaptation in healthy young adults. PLOS ONE

2021;16:e0247706. https://doi.org/10.1371/journal.pone.0247706.

[2]   Gonzalez-Rubio M, Velasquez NF, Torres-Oviedo G. Explicit Control of Step Timing

During Split-Belt Walking Reveals Interdependent Recalibration of Movements in Space

and Time. Front Hum Neurosci 2019;13:207. https://doi.org/10.3389/fnhum.2019.00207.

[3]   Malone LA, Bastian AJ. Thinking About Walking: Effects of Conscious Correction Versus

Distraction on Locomotor Adaptation. J Neurophysiol 2010;103:1954–62.

https://doi.org/10.1152/jn.00832.2009.

[4]   Mariscal DM, Iturralde PA, Torres-Oviedo G. Altering attention to split-belt walking

increases the generalization of motor memories across walking contexts. J Neurophysiol

2020;123:11. https://doi.org/10.1152/jn.00509.2019.

[5]   Statton MA, Vazquez A, Morton SM, Vasudevan EVL, Bastian AJ. Making Sense of

Cerebellar Contributions to Perceptual and Motor Adaptation. The Cerebellum

2018;17:111–21. https://doi.org/10.1007/s12311-017-0879-0.

[6]   Finley JM, Long A, Bastian AJ, Torres-Oviedo G. Spatial and Temporal Control Contribute

to Step Length Asymmetry During Split-Belt Adaptation and Hemiparetic Gait.

Neurorehabil Neural Repair 2015;29:786–95. https://doi.org/10.1177/1545968314567149.

[7]   Roper JA, Brinkerhoff SA, Harrison BR, Schmitt AC, Roemmich RT, Hass CJ. Persons

with essential tremor can adapt to new walking patterns. J Neurophysiol 2019;122:1598–

605. https://doi.org/10.1152/jn.00320.2019.

[8]   Dietz V, Zijlstra W, Duysens J. Human neuronal interlimb coordination during split-belt

locomotion. Exp Brain Res 1994;101. https://doi.org/10.1007/BF00227344.

[9] Reisman DS, Block HJ, Bastian AJ. Interlimb Coordination During Locomotion: What Can

be Adapted and Stored? J Neurophysiol 2005;94:2403–15.

https://doi.org/10.1152/jn.00089.2005.

[10] Long AW, Roemmich RT, Bastian AJ. Blocking trial-by-trial error correction does not

interfere with motor learning in human walking. J Neurophysiol 2016;115:2341–8.

https://doi.org/10.1152/jn.00941.2015.

[11] Smith MA, Ghazizadeh A, Shadmehr R. Interacting Adaptive Processes with Different

Timescales Underlie Short-Term Motor Learning. PLoS Biol 2006;4:e179.

https://doi.org/10.1371/journal.pbio.0040179.

[12] Darmohray DM, Jacobs JR, Marques HG, Carey MR. Spatial and Temporal Locomotor

Learning in Mouse Cerebellum. Neuron 2019;102:217-231.e4.

https://doi.org/10.1016/j.neuron.2019.01.038.

[13] Mawase F, Haizler T, Bar-Haim S, Karniel A. Kinetic adaptation during locomotion on a

split-belt treadmill. J Neurophysiol 2013;109:2216–27.

https://doi.org/10.1152/jn.00938.2012.

[14] Roemmich RT, Long AW, Bastian AJ. Seeing the Errors You Feel Enhances Locomotor

Performance but Not Learning. Curr Biol 2016;26:2707–16.

https://doi.org/10.1016/j.cub.2016.08.012.

[15] Sánchez N, Simha SN, Donelan JM, Finley JM. Using asymmetry to your advantage:

learning to acquire and accept external assistance during prolonged split-belt walking. J

Neurophysiol 2021;125:344–57. https://doi.org/10.1152/jn.00416.2020.

[16] Seethapathi N, Clark B, Srinivasan M. Exploration-based learning of a step to step

controller predicts locomotor adaptation. Neuroscience; 2021.

https://doi.org/10.1101/2021.03.18.435986.

[17] Sánchez N, Simha SN, Donelan JM, Finley JM. Taking advantage of external mechanical

work to reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking.

J Physiol 2019;597:4053–68. https://doi.org/10.1113/JP277725.

[18] Selgrade BP, Thajchayapong M, Lee GE, Toney ME, Chang Y-H. Changes in mechanical

work during neural adaptation to asymmetric locomotion. J Exp Biol 2017:jeb.149450.

https://doi.org/10.1242/jeb.149450.

[19] Donelan JM, Kram R, Kuo AD. Simultaneous positive and negative external mechanical

work in human walking. J Biomech 2002;35:117–24. https://doi.org/10.1016/S0021-

9290(01)00169-5.

[20] Donelan JM, Kram R, Kuo AD. Mechanical work for step-to-step transitions is a major

determinant of the metabolic cost of human walking. J Exp Biol 2002;205:3717–27.

[21] Burdett RG, Skrinar GS, Simon SR. Comparison of mechanical work and metabolic energy

consumption during normal gait. J Orthop Res 1983;1:63–72.

https://doi.org/10.1002/jor.1100010109.

[22] Umberger BR, Gerritsen KGM, Martin PE. A Model of Human Muscle Energy Expenditure.

Comput Methods Biomech Biomed Engin 2003;6:99–111.

https://doi.org/10.1080/1025584031000091678.

[23] Buurke TJW, Lamoth CJC, Vervoort D. Adaptive control of dynamic balance in human gait

on a split-belt treadmill. J Exp Biol 2018:10.

[24] Finley JM, Bastian AJ, Gottschall JS. Learning to be economical: the energy cost of

walking tracks motor adaptation: Split-belt adaptation reduces metabolic power. J Physiol

2013;591:1081–95. https://doi.org/10.1113/jphysiol.2012.245506.

[25] Selinger JC, O'Connor SM, Wong JD, Donelan JM. Humans Can Continuously Optimize

Energetic Cost during Walking. Curr Biol 2015;25:2452–6.

https://doi.org/10.1016/j.cub.2015.08.016.

[26]   Park S, Finley JM. Manual stabilization reveals a transient role for balance control during

572   locomotor adaptation. J Neurophysiol 2022;128:808–18.

573   https://doi.org/10.1152/jn.00377.2021.

574   [27]   Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, et al. The

575   Physical Activity Guidelines for Americans. JAMA 2018;320:2020.

576   https://doi.org/10.1001/jama.2018.14854.

577   [28]   Cher PH, Stewart IB, Worringham CJ. Minimum Cost of Transport in Human Running Is

578   Not Ubiquitous. Med Sci Sports Exerc 2015;47:307–14.

579   https://doi.org/10.1249/MSS.0000000000000421.

580   [29]   Kurtze N, Rangul V, Hustvedt B-E. Reliability and validity of the international physical

581   activity questionnaire in the Nord-Trøndelag health study (HUNT) population of men. BMC

582   Med Res Methodol 2008;8:63. https://doi.org/10.1186/1471-2288-8-63.

583   [30]   Mynarski W, Rozpara M, Czapla K, Garbaciak W. Aerobic Capacity of Students with

584   Different Levels of Physical Activity as Assessed by IPAQ. J Hum Kinet 2009;21:89–96.

585   https://doi.org/10.2478/v10078-09-0011-8.

586   [31]   Jones AM, Carter H. The Effect of Endurance Training on Parameters of Aerobic Fitness:

587   Sports Med 2000;29:373–86. https://doi.org/10.2165/00007256-200029060-00001.

588   [32]   Morgan DW, Bransford DR, Costill DL, Daniels JT, Howley ET, Krahenbuhl GS. Variation

589   in the aerobic demand of running among trained and untrained subjects: Med Sci Sports

590   Exerc 1995;27:404???409. https://doi.org/10.1249/00005768-199503000-00017.

591   [33]   Godin G, Shephard RJ. A simple method to assess exercise behavior in the community.

592   Can J Appl Sport Sci 1985;10:141–6.

593   [34]   Dingwell JB, Marin LC. Kinematic variability and local dynamic stability of upper body

594   motions when walking at different speeds. J Biomech 2006;39:444–52.

595   https://doi.org/10.1016/j.jbiomech.2004.12.014.

596   [35]   R Core Team. R: A language and environment for statistical computing. R Foundation for

597   Statistical Computing 2020.

598   [36]   Pinhiero J, Bates D, DebRoy S, Sarkar D, R Core Team. nlme: Linear and Nonlinear Mixed

599   Effects Models. 2020.

600   [37]   Burnham KP, Anderson DR. Model Selection and Multimodel Inference: A Practical

601   Information-Theoretic Approach. J Wildl Manag 2003;67:655.

602   https://doi.org/10.2307/3802723.

603   [38]   Lohse KR, Shen J, Kozlowski AJ. Modeling Longitudinal Outcomes: A Contrast of Two

604   Methods. J Mot Learn Dev 2020;8:145–65. https://doi.org/10.1123/jmld.2019-0007.

605   [39]   Seidel O, Carius D, Kenville R, Ragert P. Motor learning in a complex balance task and

606   associated neuroplasticity: a comparison between endurance athletes and nonathletes. J

607   Neurophysiol 2017;118:1849–60. https://doi.org/10.1152/jn.00419.2017.

608   [40]   Abram SJ, Poggensee KL, Sánchez N, Simha SN, Finley JM, Collins SH, et al. General

609   variability leads to specific adaptation toward optimal movement policies. Curr Biol

610   2022;32:2222-2232.e5. https://doi.org/10.1016/j.cub.2022.04.015.

611   [41]   Charalambous CC, Alcantara CC, French MA, Li X, Matt KS, Kim HE, et al. A single

612   exercise bout and locomotor learning after stroke: physiological, behavioural, and

613   computational outcomes: Effect of exercise on locomotor learning in chronic stroke

614   survivors. J Physiol 2018;596:1999–2016. https://doi.org/10.1113/JP275881.

615   [42]   Helm EE, Tyrell CM, Pohlig RT, Brady LD, Reisman DS. The presence of a single-

616   nucleotide polymorphism in the BDNF gene affects the rate of locomotor adaptation after

617   stroke. Exp Brain Res 2016;234:341–51. https://doi.org/10.1007/s00221-015-4465-8.

618   [43]   Hirata K, Hanawa H, Miyazawa T, Kubota K, Sonoo M, Kokubun T, et al. Adaptive

619   changes in foot placement for split-belt treadmill walking in individuals with stroke. J

620   Electromyogr Kinesiol 2019;48:112–20. https://doi.org/10.1016/j.jelekin.2019.07.003.

[44] Sallis JF, Saelens BE. Assessment of Physical Activity by Self-Report: Status, Limitations,

and Future Directions. Res Q Exerc Sport 2000;71:1–14.

https://doi.org/10.1080/02701367.2000.11082780.

[45] Castillo-Retamal M, Hinckson EA. Measuring physical activity and sedentary behaviour at

work: A review. Work 2011;40:345–57. https://doi.org/10.3233/WOR-2011-1246.

[46] Corder K, van Sluijs EM, Wright A, Whincup P, Wareham NJ, Ekelund U. Is it possible to

assess free-living physical activity and energy expenditure in young people by self-report?

Am J Clin Nutr 2009;89:862–70. https://doi.org/10.3945/ajcn.2008.26739.

[47] Ishikawa-Takata K, Tabata I, Sasaki S, Rafamantanantsoa HH, Okazaki H, Okubo H, et al.

Physical activity level in healthy free-living Japanese estimated by doubly labelled water

method and International Physical Activity Questionnaire. Eur J Clin Nutr 2008;62:885–91.

https://doi.org/10.1038/sj.ejcn.1602805.

Supporting information captions

S1 Table. Model fits of mixed effects linear models for each outcome measure.

Note: All models contain a random plateau on participant. Model 1 contains fixed effects

for the parameters in the one-exponent equation (eq.1), fitted to the whole sample.

Model 2 contains fixed effects for the parameters in the one-exponent equation, fitted by

group. Model 3 contains fixed effects for the parameters in the two-exponent equation

(eq. 2), fitted to the whole sample. Model 4 contains fixed effects for the parameters in

the two-exponent equation, fitted by group. AIC = Akaike Information Criterion; BIC =

Bayesian Information Criterion. The final model for each outcome measure is in bold

(model 4 for all outcome measures).

S2 Fig. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of step length asymmetry adaptation.

S3 Fig. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the positive work rate of the fast leg.

S4 Fig. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the negative work rate of the fast leg.

S5 Fig. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the positive work rate of the slow leg.

S6 Fig. A comparison of the group average model fit (black line) and the individual model fits

(blue line) of the adaptation of the positive work rate of the slow leg.

A

fast belt   slow belt

B

Warmup self-selected typical speed

Warmup fast speed

<u>Baseline</u>
Warmup slow speed

<u>Adaptation</u>
Belt speeds split 2:1 ratio

3 min   3 min   3 min   10 min