Reproductive fitness is associated with female chronotype in a songbird

Authors:

Robyn J. Womack[1#], Pablo Capilla-Lasheras[1#@], Ciara L. O. McGlade[2], Davide M. Dominoni[1], Barbara Helm[1, 3]

[#]: joint first authors

[@]: corresponding author - pacapilla@gmail.com

Affiliations:

[1] School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, G12 8QQ Glasgow, UK

[2] School of Biological Sciences, Queen's University Belfast, BT9 5DL Belfast, UK

[3] Swiss Ornithological Institute, Seerose 1, 6204 Sempach, Switzerland

RJW: robynwomack@outlook.com; PC-L: pacapilla@gmail.com; CLOM: ciara.mcglade123@gmail.com; DMD: davide.dominoni@glasgow.ac.uk BH: barbara.helm@vogelwarte.ch

Running title: Fitness-chronotype associations in a songbird

Keywords: chronotype, reproduction, incubation, rhythms, activity, timing

Abstract

Research on biological rhythms has revealed widespread variation in timing within populations. Repeatable individual chronotypes have been linked to performance in humans but, in free-living species, benefits of chronotype are poorly understood. To address this gap, we investigated fitness correlates of incubation patterns in female songbirds (great tit, *Parus major*) at urban and forest sites. We confirm repeatable chronotypes ($r \geq 0.31$) and show novel links between chronotype and reproductive fitness. In both habitats, females that started activity earlier in the day raised more fledglings. We also observed that forest females started their day at similar times throughout the breeding season, whereas urban females tied their onset of activity closely to sunrise. Our study points to possible mechanisms that underlie chronotype variation and provides sought-after evidence for its relevance to fitness.

Introduction

Due to the rotation of the Earth around its axis, no environments are completely constant across the 24 hour day. Hence, for organisms, appropriate diel timing of activities and physiology relative to environmental cycles is thought to be important for fitness (Kronfeld-Schor & Dayan 2003). Still, inter-individual differences in diel activities can be large, whereby individuals display highly consistent temporal phenotypes called "chronotype" (Roenneberg *et al.* 2003; Alós *et al.* 2017; Helm *et al.* 2017). Chronotype, first defined for laboratory rodents (Ehret 1974), has gained major research importance in human studies, where millions of subjects have been scored (Roenneberg *et al.* 2019). Human chronotype has been associated with genetic variants (e.g., in clock genes), performance and, physical and mental health (Roenneberg *et al.* 2003; Jones *et al.* 2019). For example, in athletes, performance depends on chronotype and can be enhanced by modified wake-up time (Facer-Childs & Brandstaetter 2015).

Interest in chronotype is also rapidly increasing in ecology and evolution (Alós *et al.* 2017; Helm *et al.* 2017; Maury *et al.* 2020), fuelled by remote and automated tracking technology (e.g., transmitters or on-site loggers (Dominoni *et al.* 2013; Graham *et al.* 2017; Maury *et al.* 2020)). Simultaneous data collection from many individuals is paving the way for studying fitness implications of particular chronotypes, the mechanisms that generate them, and the maintenance of inter-individual variation (Hau *et al.* 2017; Helm *et al.* 2017; Martorell-Barceló *et al.* 2018). Still, our understanding of causes and consequences of chronotype in free-living animals has been hindered by shortcomings when assessing chronotype, and by bias in the sex and traits investigated.

First, disentangling contributing factors to variation in chronotype requires engaging with the complexity of diel timing, which is based on circadian rhythms in

interaction with light, further environmental features and state-dependent modifiers

(Helm *et al.* 2017). Chronotype is broadly defined as consistent timing of an individual's

rhythmic features (e.g., activity onset), relative to an external reference and to

conspecifics measured under similar conditions (Helm *et al.* 2017). Choice of temporal

reference is challenging. Human studies usually refer to a fixed time of day (hereafter

called 'absolute time') (Roenneberg *et al.* 2003; Jones *et al.* 2019). In contrast, because

animals often track natural light environments (Bennie *et al.* 2014), most ecological

studies refer to annually and spatially variable aspects of the solar day (e.g., sunrise;

hereafter called 'relative time') (Graham *et al.* 2017; Maury *et al.* 2020). However,

animals typically change their relationship to these solar features over time and space

(Daan & Aschoff 1975) so that, for example, latitude, daylength and life-cycle stage can

bias or alter findings (Shaw & Cresswell 2014; Da Silva & Kempenaers 2017). Therefore,

variation in both absolute and relative timing of activities should be investigated in

parallel to understand variation in chronotype in the wild.

Secondly, investigation of fitness effects must be broadened in scope. Fitness

studies on wild chronotypes have mostly focused on males (but see Maury *et al.*

(2020)), partly due to extra-pair mating and to conspicuous features such as courtship,

song and ornaments (Pagani-Núñez & Senar 2016; Hau *et al.* 2017). For example, avian

observational and experimental studies suggest that early-active males may sire more

extra-pair young than late-active males (Kempenaers *et al.* 2010; Greives *et al.* 2015),

and that such differences could be based on endogenous circadian clocks (Helm &

Visser 2010). However, females are disproportionately more involved in reproductive

activities (Mace 1985), and mating represents but a fraction of factors that shape the

fitness landscape of chronotype. For example, offspring must develop to sexual

maturity, potentially requiring extensive parental care, and adults must survive, forage

and maintain sufficient body condition to generate offspring. Thus, data are needed for

implications of female chronotype, and for diverse life-cycle stages.

To investigate fitness implications of chronotype, birds offer excellent study

opportunities because their behaviours and reproductive outcomes are unusually

tractable. Therefore, in this study, we leverage data from wild birds to i) disentangle

particular factors that contribute to explain variation in chronotype, and ii) newly

document links between chronotype and reproductive fitness. In a well-studied

songbird, the great tit (*Parus major*), we inferred female chronotype through measuring

behaviour during incubation, a critically important post-zygotic stage of avian

reproduction, while monitoring reproductive success (Graham *et al.* 2017; Capilla-

Lasheras 2018; Gwinner *et al.* 2018; Maury *et al.* 2020). Because features of the

environment can influence chronotype, we also included data from two habitat types,

urban and forest, which often affect the diel rhythm of animals (Dominoni *et al.* 2013;

Gaynor *et al.* 2018).

Great tits are female-only intermittent incubators, who spend nights on their

nest, but at daytime alternate between nest attendance (i.e., on-bouts) and foraging

outings (i.e., off-bouts) (Diez-Méndez *et al.* 2021). From small temperature loggers (i.e.,

iButtons) inserted into nests of urban and forest-breeding great tits, we first infer

incubation chronotype of females for both absolute and relative time, and assess

consistency of chronotype across the breeding season. As our measure of chronotype,

we focus on activity onset (inferred from the first incubation off-bout of the day), which

in birds is particularly robust and sometimes associated with male fitness (Kempenaers

et al. 2010; Dominoni *et al.* 2013; Pagani-Núñez & Senar 2016; Graham *et al.* 2017; Hau

et al. 2017), but we also report end and duration of activity. Secondly, we link

incubation chronotype to reproductive outcomes from these same nests to test

associations between fitness and female chronotype. Our research spans three years

and multiple breeding locations in Scotland, ranging from oak forests to urban settings.

Methods

Study populations and field protocols

We studied five nest-box breeding populations of great tits (*Parus major*; nest-box

details: woodcrete material, 260H x 170W x 180D, hole diameter 32 cm, Schwegler,

Germany) during the breeding seasons of 2016, 2017 and 2018 (April to June). Three

study populations were located in ancient deciduous forests, dominated by oak species

(*Quercus* sp.; Scottish Centre for Ecology and the Natural Environment [SCENE; N = 28

nest-boxes included in the study; 56° 7' N, 4° 36' W], Sallochy Forest [N = 8 nest-boxes

included in the study; 56° 7' N, 4° 36' W] and Cashel Forest [N = 31 nest-boxes included

in the study; 56° 6' N, 4° 34' W]). The remaining two populations were situated in an

urban park (Kelvingrove Park [N = 14 nest-boxes included in the study; 55° 52' N, 4° 16'

W]) and a suburban park (Garscube estate [N = 9 nest-boxes included in the study;

55°54' N, 4°19' W]) within the city of Glasgow (UK). Both urban sites contained a

mixture of open land, small shrubs, and sparse woodland with introduced and native

tree species. For further details on the study sites, see Capilla-Lasheras et al. (2017),

Pollock et al. (2017) and Jarrett et al. (2020).

All nest-boxes were checked weekly from April 1 for signs of nest building

activity and egg laying. Once a new completed clutch was detected, we calculated the

date of clutch completion (from the number of eggs present between two consecutive

nest-box visits, assuming that females laid one egg per day). After the estimated earliest

possible date of hatching (assuming a minimum incubation length of 14 days from the

date of clutch completion; Gosler 1993), nest-boxes were checked every other day,

allowing determination of exact date of hatching based on nestling presence. Thirteen

days after hatching, all nestlings within a brood were weighed (± 0.01 g) and ringed for

individual identification (N = 57 broods of 13-day old nestlings). Nest-boxes were

checked again > 21 days after hatching to determine the number and identity of any

dead nestlings remaining in the nest. As our sample size varied slightly per each trait

under investigation (see below), we provide a breakdown of sample size per habitat and

year in Table S1. Sunrise and sunset times at SCENE (56° 7' 46'' N, 4° 36' 46'' W) and

Glasgow (55° 52' 11'' N, 4° 16' 56'' W) were obtained from www.timeanddate.com. Our

data are collected from individual nest-boxes, rather than from identified females. Thus,

some individuals might have been recorded in multiple years. Given that our study was

spread across five sites over three years, the potential bias introduced by this

methodological limitation is expected to be minimal (Table S1).

Incubation temperature data

To quantify incubation behaviour in female great tits, we deployed small temperature

loggers (iButtons DS1922L-F5, Thermochron) inside their nests (Figure S1). We

programmed iButtons to record temperature (± 0.0625 °C) every 3 min (Capilla-

Lasheras 2018). iButtons were placed carefully next to the eggs (after the third egg of

the clutch had been laid), covered with a small piece of white cloth, and attached to the

base of the nest by a green wire anchored by a small fishing weight (Figure S1).

Environmental temperature data

To control for variation in environmental temperature when quantifying incubation behaviour (Capilla-Lasheras 2018), daily mean temperatures for the breeding seasons of 2016, 2017 and 2018 were obtained from the UK Met Office for an area close to our forest sites (Tyndrum [56°25'N, 4°42'W] and city sites (Bishopton [55°54'N, 4°30'W]). We also incorporated daily mean temperatures in our statistical models explaining variation in incubation behaviour (details below).

Ethical statement

All field procedures were carried out under the appropriate Home Office project licence and BTO ringing licence held by Barbara Helm.

Quantification of incubation behaviour

Some individuals removed iButtons from the nest cup and pushed them to the side of the nest-box, so that these iButtons did not record incubation temperature accurately. These cases of failed incubation temperature data collection were identified by visual inspection of the incubation temperature time series blind to factors in the analysis and were removed from the dataset. Our incubation analyses only included days of incubation after the clutch was completed and started no earlier than 15 days before the hatch date. In total, 729 days of iButton incubation temperature recordings from 102 clutches were included in the analysis (sample sizes vary slightly across statistical models; details are given in the result section and Table S1).

Incubation behaviour (e.g., on- and off-bout timing and length) was determined using the R package incR (v1.1.0; Capilla-Lasheras 2018; Gwinner *et al.* 2018), choosing

parameter values for *incRscan* validated for great tit incubation (Capilla-Lasheras 2018;

lower.time=22, *upper.time*=3, *sensitivity*=0.15, *temp.diff*=8, *maxNightVar_accepted*=2).

For each incubating female, we determined: first morning off-bout, last evening on-bout,

and duration of active day (e.g., time difference between the first morning off-bout and

the last evening on-bout) (Capilla-Lasheras 2018).

Data analysis

General modelling procedures

All analyses and visualisations were performed in R (version 4.1.1; R Core Team 2022).

Generalised linear mixed models were employed to explain variation in several

incubation and reproductive traits (see below). For each of these traits, we first built a

global model that contained all explanatory variables and interactions of interest for

each trait (see below). Then, we used likelihood-ratio tests (LRTs) to investigate the

importance of each model predictor. Starting from each global model, nested models

were sequentially built and compared via LRTs until a minimal adequate model was

reached (i.e., a model containing only significant terms). The statistical significance of

terms present in the minimal adequate model was assessed dropping such terms (one

at a time) from the model and comparing this new simpler model against the minimal

adequate model via LRT. The statistical importance of predictors not included in the

minimal adequate model was calculated in a similar way, using LRTs which compared

the minimal adequate with and without the focal predictor. Quadratic terms were only

included in models that also contained linear effects. When two-way interaction terms

were present in a model, the single effect predictors that formed the interaction were

always included. Random effects were present in every model as specified for the

analysis of each response variable (details below). All statistical models were performed

using the R package lme4 (v1.1.29; Bates *et al.* 2015). Linear model residuals were

visually inspected to check the assumption of normality. The R package DHARMa

(v0.4.5; Hartig 2018) was employed to check the normality of residuals in generalised

linear models.

Global models

Incubation behaviour: We analysed absolute and relative onsets and ends of activity. To

account for differences in photoperiod throughout the breeding season, we calculated

relative onset as the time of the first incubation off-bout minus sunrise time for each

day (i.e., positive values represent onset of activity after sunrise, whereas negative vales

indicate an onset of activity earlier than sunrise). Similarly, relative end of activity was

defined daily as the time of the last on-bout minus sunset time (i.e., positive values

represent end of activity after sunset, whereas negative values indicate an end of

activity earlier than sunset).

Global models for onset and end of activity (both absolute and relative metrics)

included as explanatory variables habitat (urban *versus* forest), clutch size (as a

continuous predictor), mean daily temperature (as a continuous predictor), and days

before hatching (as a continuous predictor whose minimum value was zero [i.e.

hatching day], included as a quadratic and a linear term - these terms effectively

modelled within-female variation in onset and end of activity). Additionally, we

controlled for absolute among-nest differences in timing of reproduction by including

the date of incubation start (i.e., clutch completion date) as a fixed effect predictor (in

number of days after April 1; included as a quadratic and a linear term - these terms

effectively modelled among-female [e.g., cross-sectional] variation in onset and end of

activity). We also included the interactions between habitat and days before hatching

(both quadratic and linear terms), and between habitat and incubation start date (both

quadratic and linear terms). Nest-box identity (90-level factor), site (5-level factor) and

year (3-level factor) were included as random effect intercepts. Using the same model

structure, we analysed the duration of the active day of incubating females, defined as

the interval (in absolute time) between the first incubation off-bout and the last on-bout

per day. Consistency (e.g., repeatability) in female onset, end and duration of activity,

was calculated as the proportion of variation explained by between-nest variation in the

linear mixed models presented above, as implemented in the R package *rptR* (Stoffel *et*

al. 2017). Female chronotype was defined as the average within-nest onset of activity,

but we also report end and duration of activity.

Survival of nestlings to fledging and nestling weight: A Poisson generalised linear mixed

model was used to explain variation in the number of nestlings that survived to

fledgling. The probability of total brood failure (i.e., the probability that no nestling

survived to fledging) was modelled using a binomial GLMM. A LMM was employed to

analyse the number of nestlings that survived to fledgling excluding broods were no

nestlings survived. Variation in the average 13-day old nestling weight per brood was

analysed using a LMM. These models included habitat (urban *versus* forest), female

chronotype (see definition above), hatching date (as a continuous variable in days after

January 1; included in the model as a linear and a quadratic term) and clutch size as

fixed effect predictors. The interactions between hatching date and habitat, and

between female chronotype and habitat, were also added. Nest-box identity, site (5-

level factor) and year (3-level factor) were included as random effect intercept.

Results

Correlates and consistency of incubating female chronotype

We recorded nest temperatures in 2016, 2017 and 2018, and analysed 729 days of great tit incubation in 102 clutches (median = 7 days of incubation per clutch; range = 1 - 15 days; see details of sample sizes in Table S1). Urban great tit females laid their eggs and started incubation earlier in the year than forest females, and thus experienced shorter days with later sunrise and earlier sunset (start of incubation date: $mean_{urban} \pm SE = 30^{th}$ April $\pm$ 1.09 days; N_{urban} = 27 clutches; $mean_{forest} \pm SE = 8^{th}$ May $\pm$ 0.61 days; N_{forest} = 75 clutches). Therefore, we detail results separately for the two habitats, although all data were analysed in overarching models.

We found that, on population level, absolute time of activity onset was affected by habitat and by the date when incubation started (interaction 'Incubation start date × habitat': $\chi^2_{df = 1}$ = 11.41, p = 0.001; Table 1; Figure 1a & 1b). Urban females closely tracked the seasonally advancing sunrise time (Figure 1a), but forest females largely ignored this advance and rose at similar absolute time throughout the season (Figure 1b). Whereas early-breeding urban females started activity at a later absolute time than forest females, for late-breeding birds the pattern reversed, so that urban females started their day earlier than forest females (Figure 1a & 1b). In contrast, the mean end of activity was similar in both habitats and became progressively earlier with later incubation start date (Figure S2; Table S2 – S5). Overall, the active day lengthened over the breeding season for urban but shortened for forest females (Figure S3 & Table S6 & S7). Time left until hatching also affected timing, whereas ambient temperature and clutch size did not (Tables 1, S2 – S7).

Relative to sunrise, females started their day on average 28.5 min after sunrise

(SE = 0.91 min; range = [-87 min, 137 min]), but onset of activity depended on habitat

and on the date when incubation started (Figure 1c-1d; Table 2). Females that initiated

incubation later in the year began their day relative to sunrise progressively earlier in

the city, but progressively later in the forest (interaction 'Incubation start date ×

habitat': $\chi^2_{df = 1}$ = 11.44, p = 0.001; Figure 1c-1d). The end of activity relative to sunset

advanced consistently over the season in both habitats ('Incubation start date': $\chi^2_{df = 1}$ =

36.78, p = 0.001; Figure S2c-S2d; Table S4 & S5). Effects of the time left until hatching,

ambient temperature and clutch size were similar to those for absolute time (Tables 1,

S2 – S7).

The population pattern described above was broadly matched by temporal

variation within clutches (i.e., variation between the first and last day of incubation of a

clutch; Tables 1-2; Figures S4 – S5). We identified consistent individual differences in

the time of onset of activity (i.e., female chronotype). Among-female differences

explained 31 % of the variation in absolute onset time (LRT on the nest-box ID random

effect: $\chi^2_{df = 1}$ = 134.30, p < 0.0001; among-female repeatability [95%CI] = 0.31 [0.21,

0.40]). Analyses of relative onset time (i.e., correcting for changes in sunrise time)

yielded similar results, with consistent among-female differences in onset of activity

(repeatability [95%CI] = 0.32 [0.22, 0.41]). We also found consistent among-female

differences in the end of activity, both in absolute (repeatability [95%CI] = 0.25 [0.16,

0.34]) and relative end of activity (repeatability [95%CI] = 0.25 [0.17, 0.34]); and

consistent among-female differences in the duration of the active day (repeatability

[95%CI] = 0.20 [0.11, 0.28]).

Effects of female chronotype on fledging success and pre-fledging weight

We detected substantial variation between broods in the number of nestlings that survived to fledging. Female chronotype, derived from activity onset relative to sunrise, predicted the number of surviving nestlings: the earlier the female chronotype, the more nestlings fledged ($\chi^2_{df=1}$ = 5.49, p = 0.019; Figure 2a; Table 3a). This effect was consistent across habitats (interaction between female chronotype and habitat, $\chi^2_{df=1}$ = 0.43, p = 0.513; Table 3a) and was robust to controlling for clutch size ($\chi^2_{df=1}$ = 19.31, p < 0.0001; Table 3a). The number of surviving chicks was also strongly affected by habitat ($\chi^2_{df=1}$ = 6.82, p = 0.009; Table 3a; urban females fledged 0.66 nestlings less than forest females (i.e., a decrease in surviving nestlings of 48%; Table 4a). Conversely, absolute chronotype of females did not predict the number of surviving nestlings (Table S8).

To investigate the breeding parameters that could have generated the decreasing reproductive success with later female chronotype relative to sunrise, we performed two additional analyses. Firstly, we assessed whether female chronotype was associated with total brood failure, and found no support (effect of female chronotype on the probability that no nestling survives to fledging: $\chi^2_{df=1}$ = 0.15, p = 0.699; Table S9). Secondly, we tested whether female chronotype predicted the number of nestlings surviving to fledging in successful broods (i.e., those that fledged at least one offspring), and confirmed that earlier chronotypes fledged more offspring than later chronotypes ($\chi^2_{df=1}$ = 7.03, p = 0.008; Table S10 & S11).

Mean body mass of 13-day old nestlings was affected by habitat, hatching date and clutch size. Forest nestlings were on average 2.73g (95%CI = [1.66, 3.86]; Table 3b;

Table 4b) heavier than urban nestlings of the same age. In both habitats, pre-fledging

weight was higher for broods that hatched in the middle of the season (Figure 2b), and

nestlings from larger clutches were on average lighter ($\chi^2_{df = 1}$ = 4.10, p = 0.043; Table

3b; Table 4b). Neither relative nor absolute female chronotype affected pre-fledging

weight in either habitat (relative chronotype: $\chi^2_{df = 1}$ = 0.20, p = 0.656; absolute

chronotype: $\chi^2_{df = 1}$ = 0.31, p = 0.580; interaction terms between chronotype and habitat

were non-significant, for relative chronotype: $\chi^2_{df = 1}$ < 0.01, p = 0.939; for absolute

chronotype: $\chi^2_{df = 1}$ < 0.01, p = 0.938).

Discussion

Recent research has identified surprisingly high variation in chronotype of free-living

animals, but determinants and effects of this variation are still largely unclear. Our

study is among the few that have identified fitness correlates of chronotype, in

particular in female animals. We firstly show high repeatability of timing, and thus

corroborate the interpretation of chronotype as a consistent individual trait. We then

show that the chronotype of female great tits, measured during the incubation period,

predicted reproductive success, such that early-rising females raised more offspring to

fledging than late chronotypes.

Early rising may be beneficial as energy stores are depleted, and the ability of

small passerine birds to successfully forage peaks in the early morning, once light

conditions are suitable (Kacelnik 1979; Pagani-Núñez & Senar 2016). However, foraging

in the early morning can be costly because of low ambient temperatures, and may be

constrained by condition of an individual and its endogenous circadian clock.

Balancing costs and benefits of early rising might be intricate during incubation.

For uniparental incubators, self-maintenance is weighed up against maximal offspring

development (Nord & Cooper 2020). This trade-off is heightened during early morning

hours, when incubators must replenish energy stores while the typically low morning

temperatures risk cooling of the eggs. Hence, an incubating female should delay leaving

the nest until she can forage efficiently, and promptly return to the nest. Early rising

may thus indicate superior condition of incubating females, as proposed for courtship

song and provisioning of males (McNamara *et al.* 1987; Murphy *et al.* 2008; Pagani-

Núñez & Senar 2016).

Alternatively, the ability to perform efficiently early in the day could be based on

circadian mechanisms that facilitate an early start, as demonstrated in human athletes

(Vitale & Weydahl 2017). Reproductive advantages due to circadian-based early-rising

have been proposed for male great tits whose circadian rhythm affects extra-pair

paternity (EPP), and have been supported by experiments on the same study species

(Helm & Visser 2010; Greives *et al.* 2015; Hau *et al.* 2017). Great tit chicks with fast

circadian clocks were significantly more likely to be sired through EPP, and males

whose circadian system was pharmacologically delayed lost paternity (Helm & Visser

2010; Greives *et al.* 2015). As in these other studies, our work found benefits for the

early bird, without indicating what benefits or costs, in turn, might arise for late

chronotypes.

A putative circadian basis to early chronotype could involve several mechanistic

features. These include a fast clock (i.e., short free-running period; Helm & Visser 2010),

but also individual variation in sensitivity to light (Brown *et al.* 2008; Helm *et al.* 2017;

Tudorache *et al.* 2018; Jones *et al.* 2019). A contribution of light pathways to the

chronotype - fitness link is indicated by our comparison of absolute and relative timing.

Fitness effects were evident only for chronotype relative to sunrise, but not for absolute

timing. This suggests that for incubating females, shifting their response in relation to

light cues is more important for fitness than maintaining absolute timing.

We also detected unexpected differences in response to sunrise, but not sunset,

between females at urban and forest sites. Forest females started activity at almost

constant times of day, despite the rapid advance of sunrise time over the breeding

season. Conversely, urban females were far more responsive to light and largely tracked

the rapid advance of sunrise. This finding was counter to the expectation that in urban

habitats, where artificial light at night is prevalent, the birds' responsiveness to natural

light would be reduced (Roenneberg *et al.* 2007; Dominoni *et al.* 2013). It is possible

that habitat differences other than light levels contributed to the differences in

behaviour. For related blue tits (*Cyanistes caeruleus*), the same study habitats differed in

quality, with poorer adult state and reproductive success in the city (Capilla-Lasheras *et*

al. 2017; Pollock *et al.* 2017). Thus, some urban great tit females may have foraged at

the earliest opportunity, to replenish their resources. Disentangling effects of the

circadian clock from those of the birds' body condition would require experimental

examination (Dominoni *et al.* 2013; Greives *et al.* 2015).

The only other study we are aware of that examined reproductive success

relative to incubation chronotype did not find such an association (Maury *et al.* 2020).

This investigation differed in several aspects, including use of the European starling

(*Sturnus vulgaris*) as study species. While we cannot explain the different findings, we

speculate that colonial breeding of the studied starlings may have affected

synchronicity, and thereby altered or obscured effects of chronotype (Gwinner 1966;

Menaker & Eskin 1966). In other contexts, fitness implications of chronotype are also

beginning to arise. For example, a recent study on fish showed that under fishery

pressure, chronotype was associated with differential survival (Martorell-Barceló *et al.*

2018). Still, we are far from understanding how variation in chronotype is maintained.

Our study results come with some slight caveats. Firstly, because we report

correlative data from wild birds, we cannot assess whether chronotype was affected by

the local micro-environment, either directly or via differences in individual quality

(Maury *et al.* 2020). Secondly, we have recorded female chronotype only during one life-

cycle stage, incubation, similar to earlier studies on males that considered only

courtship (Murphy *et al.* 2008). Thus, the consistency of chronotype across seasons and

life stages remains to be tested.

Nonetheless, our study strengthens the evidence for variation in chronotype in

free-living animals and provides a sought-after link to reproductive success. We extend

the circadian focus of chronotype studies by indicative findings on light pathways, and

confirm the importance of looking at both, relative and absolute time, as previously

suggested for avian incubation (Shaw & Cresswell 2014). Future challenges, likely

requiring experimental approaches, are a disentangling of effects of endogenous clock

from body condition, and determination of counter-balancing benefits that maintain

variation in chronotype.

Author contributions

RJW and BH conceived of the study. RJW, CLOM, DMD and BH collected the data. PC-L performed all statistical analyses, with input from RJW, CLOM, DMD and BH. PC-L and BH wrote the first draft of the manuscript, with major contributions by RJW. All authors read and revised the manuscript.

Data availability

All R scripts and datasets needed to reproduce the analyses presented in this paper are available at: https://github.com/PabloCapilla/incubation_chronotype. Should the manuscript be accepted, a DOI to this data repository will be provided.

References

Alós, J., Martorell-Barceló, M. & Campos-Candel, A. (2017). Repeatability of circadian behavioural variation revealed in free-ranging marine fish. *Royal Society Open Science*, 4, 160791.

Bates, D., Machler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67, 1–48.

Bennie, J.J., Duffy, J.P., Inger, R. & Gaston, K.J. (2014). Biogeography of time partitioning in mammals. *Proc Natl Acad Sci U S A*, 111, 13727–13732.

Brown, S.A., Kunz, D., Dumas, A., Westermark, P.O., Vanselow, K., Tilmann-Wahnschaffe, A., *et al.* (2008). Molecular insights into human daily behavior. *Proc Natl Acad Sci U S A*, 105, 1602–1607.

Capilla-Lasheras, P. (2018). incR: a new R package to analyse incubation behaviour. *Journal of Avian Biology*, 49, e01710.

Capilla-Lasheras, P., Dominoni, D.M., Babayan, S.A., O'Shaughnessy, P.J., Mladenova, M., Woodford, L., *et al.* (2017). Elevated Immune Gene Expression Is Associated with Poor Reproductive Success of Urban Blue Tits. *Frontiers in Ecology and Evolution*, 5, 64.

Daan, S. & Aschoff, J. (1975). Circadian rhythms of locomotor activity in captive birds and mammals: Their variations with season and latitude. *Oecologia*, 18, 269–316.

Diez-Méndez, D., Sanz, J.J. & Barba, E. (2021). Impacts of ambient temperature and clutch size on incubation behaviour onset in a female-only incubator songbird. *Ibis*, 163, 1056–1071.

Dominoni, D.M., Helm, B., Lehmann, M., Dowse, H.B. & Partecke, J. (2013). Clocks for the city: circadian differences between forest and city songbirds. *Proceedings of the Royal Society B*, 280, 20130593.

Ehret, C.F. (1974). The sense of time: evidence for its molecular basis in the eukaryotic gene-action system. *Adv Biol Med Phys*, 15, 47–77.

Facer-Childs, E. & Brandstaetter, R. (2015). The impact of circadian phenotype and time since awakening on diurnal performance in athletes. *Current Biology*, 25, 518–522.

Gaynor, K.M., Hojnowski, C.E., Carter, N.H. & Brashares, J.S. (2018). The influence of human disturbance on wildlife nocturnality. *Science*, 360, 1232–1235.

Gosler, A. (1993). *The great tit*. Hamlyn.

Graham, J.L., Cook, N.J., Needham, K.B., Hau, M. & Greives, T.J. (2017). Early to rise, early to breed: A role for daily rhythms in seasonal reproduction. *Behavioral Ecology*, 28, 1266–1271.

Greives, T.J., Kingma, S.A., Kranstauber, B., Mortega, K., Wikelski, M., van Oers, K., *et al.*
(2015). Costs of sleeping in: Circadian rhythms influence cuckoldry risk in a
songbird. *Functional Ecology*, 29, 1300–1307.

Gwinner, E. (1966). Entrainment of a circadian rhythm in birds by species-specific song
cycles (Aves, Fringillidae: Carduelis spinus, Serinus serinus). *Experientia (Basel)*,
22, 765.

Gwinner, H., Capilla-lasheras, P., Cooper, C. & Helm, B. (2018). "Green incubation": avian
offspring benefit from aromatic nest herbs through improved parental incubation
behaviour. *Proceedings of the Royal Society B*, 285, 20180376.

Hartig, F. (2018). DHARMa: Residual diagnostics for hierarchical (multi-Level / mixed)
regression models.

Hau, M., Dominoni, D., Casagrande, S., Buck, C.L., Wagner, G., Hazlerigg, D., *et al.* (2017).
Timing as a sexually selected trait⍰: the right mate at the right moment.
Philosophical Transactions of the Royal Society B, 372, 20160249.

Helm, B. & Visser, M.E. (2010). Heritable circadian period length in a wild bird
population. *Proceedings of the Royal Society B*, 277, 3335–3342.

Helm, B., Visser, M.E., Schwartz, W., Kronfeld-Schor, N., Gerkema, M., Piersma, T., *et al.*
(2017). Two sides of a coin: Ecological and chronobiological perspectives of timing
in the wild. *Philosophical Transactions of the Royal Society B*, 372, 20160246.

Jarrett, C., Powell, L.L., McDevitt, H., Helm, B. & Welch, A.J. (2020). Bitter fruits of hard
labour: diet metabarcoding and telemetry reveal that urban songbirds travel
further for lower-quality food. *Oecologia*, 193, 377–388.

Jones, S.E., Lane, J.M., Wood, A.R., van Hees, V.T., Tyrrell, J., Beaumont, R.N., *et al.* (2019).
Genome-wide association analyses of chronotype in 697,828 individuals provides
insights into circadian rhythms. *Nature Communications*, 10, 343.

Kacelnik, A. (1979). The foraging efficiency of great tits (Parus major L.) in relation to
light intensity. *Animal Behaviour*, 27, 237–241.

Kempenaers, B., Borgström, P., Loës, P., Schlicht, E. & Valcu, M. (2010). Artificial night
lighting affects dawn song , extra-pair siring success, and lay date in songbirds.
Current Biology, 20, 1735–1739.

Kronfeld-Schor, N. & Dayan, T. (2003). Partitioning of Time as an Ecological Resource.
Annual Review of Ecology, Evolution, and Systematics, 34, 153–181.

Mace, R. (1985). Importance of female behaviour in the dawn chorus. *Animal Behaviour*,
34, 621–622.

Martorell-Barceló, M., Campos-Candela, A. & Alós, J. (2018). Fitness consequences of fish
circadian behavioural variation in exploited marine environments. *PeerJ*, e4814.

Maury, C., Serota, M.W. & Williams, T.D. (2020). Plasticity in diurnal activity and
temporal phenotype during parental care in European starlings, Sturnus vulgaris.
Animal Behaviour, 159, 37–45.

McNamara, J.M., Mace, R.H. & Houston, A.I. (1987). Optimal daily routines of singing and
foraging in a bird singing to attract a mate. *Behavioral Ecology and Sociobiology*, 20,
399–405.

Menaker, M. & Eskin, A. (1966). Entrainment of circadian rhythms by sound in Passer
domesticus. *Science,* 154, 1579–1581.

Murphy, M.T., Sexton, K., Dolan, A.C. & Redmond, L.J. (2008). Dawn song of the eastern
kingbird: an honest signal of male quality? *Animal Behaviour*, 75, 1075–1084.

Nord, A. & Cooper, C.B. (2020). Night conditions affect morning incubation behaviour
differently across a latitudinal gradient. *Ibis*, 162, 827–835.

Pagani-Núñez, E. & Senar, J.C. (2016). More Ornamented Great Tit Parus major Fathers
Start Feeding Their Offspring More ornamented Great Tit Parus major fathers start
feeding their offspring earlier. *Ardea*, 104, 167–175.

Pollock, C.J., Capilla-Lasheras, P., McGill, R.A.R., Helm, B. & Dominoni, D.M. (2017).
Integrated behavioural and stable isotope data reveal altered diet linked to low
breeding success in urban-dwelling blue tits (Cyanistes caeruleus). *Scientific*
Reports, 7, 5014.

R Core Team. (2022). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.

Roenneberg, T., Kumar, C.J. & Merrow, M. (2007). The human circadian clock entrains to
sun time. *Current Biology*, 17, 44–45.

Roenneberg, T., Pilz, L.K., Zerbini, G. & Winnebeck, E.C. (2019). Chronotype and social
jetlag: A (self-) critical review. *Biology (Basel)*, 8, 54.

Roenneberg, T., Wirz-Justice, A. & Merrow, M. (2003). Life between clocks: Daily
temporal patterns of human chronotypes. *Journal of Biological Rhythms*, 18, 80–90.

Shaw, P. & Cresswell, W. (2014). Latitudinal variation in day length and working day
length has a confounding effect when comparing nest attentiveness in tropical and
temperate species. *Journal of Ornithology*, 155, 481–489.

Da Silva, A. & Kempenaers, B. (2017). Singing from North to South: Latitudinal variation
in timing of dawn singing under natural and artificial light conditions. *Journal of*
Animal Ecology, 86, 1286–1297.

Stoffel, M.A., Nakagawa, S. & Schielzeth, H. (2017). rptR: repeatability estimation and
variance decomposition by generalized linear mixed-effects models. *Methods in*
Ecology and Evolution, 8, 1639-1644.

Tudorache, C., Slabbekoorn, H., Robbers, Y., Hin, E., Meijer, J.H., Spaink, H.P., *et al.* (2018).
Biological clock function is linked to proactive and reactive personality types. *BMC*
Biology, 16, 148.

Vitale, J.A. & Weydahl, A. (2017). Chronotype, Physical Activity, and Sport Performance:
A Systematic Review. *Sports Medicine*, 47, 1859–1868.

Figures

Figure 1. **Mean absolute and relative time of activity onset in urban and forest great tit females**. **(a)** On a population level, urban females advanced their absolute time of activity onset throughout the breeding season, seemingly tracking the seasonal advance of sunrise time (dashed line); whereas **(b)** the mean onset of activity of forest females remained relatively constant throughout the breeding season. Consequently, **(c)** urban females modified their onset of activity relative to sunrise over the breeding season only slightly, while forest females started their activity progressively later relative to sunrise. Points represent raw data, where each point shows the mean timing of one female, while thick solid lines and shaded areas provide mean model predictions ± 1 SE (see model coefficients in Tables 1 and 2).

Figure 2. Effects of relative female chronotype and seasonal time on pre-fledging survival and weight. (**a**) Both in urban and forest nests, progressively later chronotype of the breeding female was negatively associated with the number of nestlings that survived to fledgling (N = 101 broods; Table 3a and 4a). **(b)** Urban nestlings on day 13 of their life were lighter than forest nestlings (N = 57 broods; Table 3b and 4b). Variation in the weight of 13-day old nestlings showed a similar seasonal pattern in forest and city habitats. Points represent raw data, while thick lines and shaded areas provide mean model predictions ± 1 SE (Table 4).

Tables

Table 1. (a) Likelihood-ratio test (LRT) results for predictors explaining variation in absolute time of female onset of activity (i.e., time of first incubation off-bout). Significant terms are highlighted in bold and italic type. LRT results for 'Habitat', 'Incubation start Date[1]' and 'Days before hatching[1]' are not provided as these terms were part of a significant interaction present in the final model ('Incubation start Date[1] × Habitat' and 'Days before hatching[1] × Habitat'). The significance of these three terms is given by the LRT results of the interactions. **(b)** Minimal adequate model coefficients (i.e., after model simplification using likelihood-ratio tests) for absolute onset of activity (model coefficients are given in min after 00:00 h). Model coefficients ('Estimate') are shown along with standard errors ('SE') and 95% confidence intervals ('95% CI' obtained using 500 parametric bootstrap simulations of the model). 'df' = degrees of freedom for LRTs. Superscripts '[1]' and '[2]' refer to linear and quadratic terms, respectively; reference level for Habitat is 'urban'.

Absolute onset of activity (N = 729 days of incubation)
(a) Likelihood-ratio test results

	χ^2	Df	P
Incubation start date[1] × Habitat	*11.41*	*1*	*0.001*
Incubation start date[2] × Habitat	0.82	1	0.364
Days before hatching[1] × Habitat	*7.52*	*1*	*0.006*
Days before hatching[2] × Habitat	1.55	1	0.213
Habitat	-	-	-
Incubation start date [1]	-	-	-
Incubation start date[2]	0.01	1	0.938
Days before hatching[1]	-	-	-
Days before hatching[2]	*4.78*	*1*	*0.029*
Clutch size	0.01	1	0.935
Mean daily temperatures	1.87	1	0.171

(b) Model coefficients

Fixed effects	Estimate	SE	95% CI	
Intercept	329.07	4.243	320.60	337.47
Habitat (forest)	7.367	4.535	-2.01	16.74
Incubation start date [1]	-360.4	66.238	-495.40	-231.50
Days before hatching[2]	-43.42	19.817	-81.98	-7.48
Days before hatching[1]	-33.25	39.318	-112.30	46.60
Incubation start date[1] × Habitat (forest)	293.83	83.825	135.80	459.87
Days before hatching[1] × Habitat (forest)	130.97	47.35	35.57	219.11

Table 2. (**a**) Likelihood-ratio test (LRT) results for predictors explaining variation in
relative time of female onset of activity (i.e., time of first incubation off-bout minus
sunrise time). Significant terms are highlighted in bold and italic type. (**b**) Minimal
adequate model coefficients (i.e., after model simplification using likelihood-ratio tests)
for relative onset of activity. Model coefficients ('Estimate') are shown along with
standard errors ('SE') and 95% confidence intervals ('95% CI' obtained using 500
parametric bootstrap simulations of the model). 'df' = degrees of freedom for LRTs.
Superscripts '1' and '2' refer to linear and quadratic terms, respectively; reference level
for Habitat is 'urban'.

Relative onset of activity (N = 729 days of incubation)

(a) Likelihood-ratio test results

	χ^2	Df	P
Incubation start Date^1 × Habitat	***11.44***	***1***	***0.001***
Incubation start Date^2 × Habitat	1.67	1	0.197
Days before hatching^1 × Habitat	***3.93***	***1***	***0.048***
Days before hatching^2 × Habitat	1.09	1	0.297
Habitat	-	-	-
Incubation start Date^1	-	-	-
Incubation start Date^2	0.3	1	0.584
Days before hatching^1	-	-	-
Days before hatching^2	***6.31***	***1***	***0.012***
Clutch size	< 0.01	1	0.98
Mean daily temperatures	1.49	1	0.222

(b) Model coefficients

Fixed effects	Estimate	SE	95% CI	
Intercept	42.44	11.75	17.95	66.04
Habitat (forest)	-42.79	17.00	-76.02	-9.37
Incubation start date^1	-0.514	0.38	-1.28	0.26
Days before hatching^2	-0.131	0.05	-0.23	-0.03
Days before hatching^1	-0.068	0.84	-1.68	1.50
Incubation start date^1 × Habitat (forest)	1.71	0.49	0.76	2.68
Days before hatching^1 × Habitat (forest)	0.991	0.48	0.07	1.96

592 **Table 3**. Likelihood-ratio test (LRT) results for relative female chronotype and further
593 predictors explaining variation in (**a**) nestling survival to fledging and (**b**) weight of 13-
594 day old nestlings. Significant terms are highlighted in bold and italic type. 'df' = degrees
595 of freedom for LRTs. Superscripts '1' and '2' refer to linear and quadratic terms,
596 respectively.

a   Number of nestlings that survived to fledging (N = 101 broods)

	χ^2	df	P
Hatching date^1 × Habitat	0.02	1	0.885
Hatching date^2 × Habitat	0.03	1	0.865
Habitat	*6.82*	*1*	*0.009*
Relative female chronotype × Habitat	0.43	1	0.513
Relative female chronotype	*5.49*	*1*	*0.019*
Hatching date^1	0.08	1	0.778
Hatching date^2	0.40	1	0.527
Clutch size	*19.31*	*1*	*<0.001*

b   Pre-fledging weight of nestlings (N = 57 broods)

	χ^2	df	P
Hatching date^1 × Habitat	*11.80*	*1*	*0.001*
Hatching date^2 × Habitat	0.09	1	0.766
Habitat	-	-	-
Relative female chronotype × Habitat	0.01	1	0.939
Relative female chronotype	0.20	1	0.656
Hatching date^1	-	-	-
Hatching date^2	*19.69*	*1*	*<0.001*
Clutch size	*4.10*	*1*	*0.043*

Table 4. Minimal adequate model coefficients (i.e., after model simplification using likelihood-ratio tests) for relative female chronotype and further predictors explaining variation in (**a**) number of nestlings surviving to fledging and (**b**) pre-fledging weight of nestlings on day 13. Model coefficients ('Estimate') are shown along with standard errors ('SE') and 95% confidence intervals ('95% CI' obtained using 500 parametric bootstrap simulations of the model). Superscripts '1' and '2' refer to linear and quadratic terms, respectively; reference level for Habitat is 'urban'.

a   Number of nestlings that survived to fledging (N = 101 broods)

Fixed effects	Estimate	SE	95% CI	
Intercept	0.327	0.245	-0.111	0.810
Habitat (forest)	0.394	0.135	0.141	0.651
Relative female's chronotype	-0.007	0.003	-0.013	-0.001
Clutch size	0.145	0.031	0.084	0.203

b   Pre-fledging weight of nestlings (N = 57 broods)

Fixed effects	Estimate	SE	95% CI	
Intercept	16.920	0.697	15.333	18.463
Hatching date^1 × Habitat (forest)	21.053	5.224	9.255	33.048
Habitat (forest)	2.725	0.516	1.657	3.857
Hatching date^1	-16.152	3.604	-24.565	-7.949
Hatching date^2	-10.234	1.916	-14.896	-5.539
Clutch size	-0.164	0.076	-0.336	0.010