Single-component multilayered self-assembling protein nanoparticles presenting glycan-trimmed uncleaved prefusion optimized envelope trimers as HIV-1 vaccine candidates

Yi-Nan Zhang[1#], Jennifer Paynter[1#], Aleksandar Antanasijevic[1,2#], Joel D. Allen[3#], Mor Eldad[1], Yi-Zong Lee[1], Jeffrey Copps[1,2], Maddy Newby[3], Linling He[1], Deborah Chavez[4], Pat Frost[4], Anna Goodroe[4], John Dutton[4], Robert Lanford[4], Christopher Chen[4], Ian A. Wilson[1,2,5], Max Crispin[3], Andrew B. Ward[1,2], and Jiang Zhu[1,6]*

[1]Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, CA 92037, USA.

[2]IAVI Neutralizing Antibody Center, The Scripps Research Institute, La Jolla, CA 92037, USA.

[3]School of Biological Sciences, Highfield Campus, University of Southampton, Southampton, SO17 1BJ, UK.

[4]Southwest National Primate Research Center, Texas Biomedical Research Institute, San Antonio, TX 78227, USA.

[5]The Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, CA 92037, USA.

[6]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA 92037, USA.

[#] These authors contributed equally to this work

* Correspondence and requests for materials should be addressed to:

JZ: Phone: (858) 784-8157; Email: jiang@scripps.edu

ABSTRACT (150 words)

Uncleaved prefusion-optimized (UFO) design can stabilize diverse HIV-1 envelope

glycoproteins (Envs). Single-component, self-assembling protein nanoparticles (1c-SApNP) can

display 8 or 20 trimeric antigens as multivalent vaccines. Here, we characterized the biophysical,

structural, and antigenic properties of 1c-SApNPs that present the BG505 UFO trimer with

wildtype and modified glycans. Trimming the glycan shield improved Env recognition by

broadly neutralizing antibodies (bNAbs) to the CD4 binding site and other major glycan-

containing epitopes. In mice, rabbits, and nonhuman primates, glycan trimming increased the

frequency of vaccine responders (FVR) and steered antibody responses away from

immunodominant glycan holes and glycan epitopes. The mechanism of vaccine-induced

immunity was examined in mice. Compared with the soluble trimer, two large 1c-SApNPs

showed 420 times longer retention, 20-32 times greater presentation on follicular dendritic cell

dendrites, and up-to-4 times stronger germinal center reactions in lymph node follicles. These

findings will inform the next phase of HIV-1 vaccine development.

ONE-SENTENCE SUMMARY (125 characters)

Glycan trimming of HIV-1 Env immunogens improves the vaccine-induced neutralizing

antibody responses in small animals and primates

INTRODUCTION

The HIV-1 envelope glycoprotein (Env) mediates cell entry and is the target of the host humoral immune response. Functional Env is a trimer of noncovalent gp120-gp41 heterodimers, which are generated from the proteolytic cleavage of a gp160 precursor (*1*). In addition to substantial genetic diversity caused by an error-prone reverse transcriptase (*2*), HIV-1 exploits two Env-dependent strategies to evade host immunity: (*i*) Env metastability causes Env misfolding, sheds gp120, and leaves gp41 stumps on the virus (*3*), and (*ii*) host-produced glycans form a dense "glycan shield" to mask the Env surface (*4*). Nonetheless, up to 30% of infected people eventually develop serum neutralization against the majority of circulating HIV-1 strains. Hundreds of broadly neutralizing antibodies (bNAbs) have been isolated from HIV-1-infected donors (*5*) and target seven different epitope regions on Env: the CD4 binding site (CD4bs), three peptide-glycan epitopes (the V2 apex, V3 base, and gp120-gp41 interface), silent face (SF), fusion peptide (FP), and membrane-proximal external region (MPER) (*6, 7*). These bNAbs and the characterization of their epitopes have thus provided a number of templates for "antibody-guided" rational vaccine design (*8-11*).

The design and characterization of native-like Env trimers marked a milestone in modern HIV-1 research (*12, 13*). While gp120-derived constructs are still promising CD4bs immunogens (*14-18*), the ongoing HIV-1 vaccine efforts have largely shifted to Env trimers, as exemplified by SOSIP (*19-22*), native flexibly linked (NFL) (*23-25*), and uncleaved prefusion optimized (UFO) trimers (*26-28*). As the first rational trimer design, SOSIP transformed the HIV-1 vaccine field and generated a wealth of information on Env structure and immunogenicity (*29*). Both SOSIP and NFL are empirical designs that may require additional mutations to improve trimer stability (e.g., SOSIP.v9 (*30*)). In contrast, UFO eliminates the root cause of Env metastability with only

two implementations, UFO and UFO-BG, which have been applied to diverse Envs (*26-28*).

While Env metastability can now be dealt with by these three trimer platforms, the glycan shield

remains a challenge due to its complex roles in Env-bNAb interactions (*31-36*). Liquid

chromatography-mass spectrometry (LC-MS) (*37-39*) and structural analysis by x-ray

crystallography (*40-46*) and cryo-electron microscopy (cryo-EM) (*47-49*) resulted in site-specific

glycan profiles and atomic details of bNAb-glycan interactions, respectively. These studies

placed a premium on mimicking the glycan shield of functional viral Env in vaccine design (*50*).

Although "multistage" vaccine strategies originated from the evolutionary analysis of bNAbs

(*51*), glycans have become a major consideration. Sequential Env immunogens with glycan

deletion and reintroduction were designed to guide B cell maturation toward bNAbs (*52-54*).

Undesirable Env immunodominance is another compounding factor in bNAb elicitation using

soluble trimers, as exemplified by glycan holes and a neoepitope at the trimer base (*55, 56*). The

former (small, exposed protein surfaces within the glycan shield) are the primary target of

autologous tier-2 NAbs in rabbits (*56-65*), whereas the latter elicits non-neutralizing antibodies

(nNAbs) in multiple species (*63, 66, 67*) and causes trimer disassembly during immunization

(*68*). In rodents, glycan epitopes at C3/V4 and C3/V5 accounted for the autologous NAb

response induced by BG505 Env (*69-71*). In nonhuman primates (NHPs), the BG505 SOSIP

trimer elicited an autologous NAb response to C3/465 (*60, 72*). Therefore, the native-like trimer-

induced autologous NAb response appears to be dominated by Env surfaces that either lack or

overexpress glycans, leading to contradictory views on the glycan shield in vaccine design, i.e.,

glycan nativeness (*50*) *vs.* glycan removal or addition (*52-54, 73*). The manipulation of

individual glycans to minimize immunodominance or improve neutralization breadth proved to

be ineffective (*64*) and is unlikely to be a general solution for broadly protective vaccines.

Independent of Env design, the delivery of Env antigens in multimeric forms to mimic

virus-like particles (VLPs) has become a trend in recent vaccine research for HIV-1 (*74-77*) and

other viruses (*78-80*). With virus-like size and shape as well as a dense display of surface

antigens, VLPs can induce a more potent and long-lasting immune response than soluble

antigens (*81-85*). Engineering protein nanoparticles (NPs) to mimic authentic VLPs was the

driving force in recent technological advances in the vaccine field (*80*). Three technologies,

SpyTag/SpyCatcher (termed "SPY") (*86*), two-component NPs (2c-NP) (*87*), and single-

component self-assembling protein NPs (1c-SApNP) (*27, 88, 89*), have each been assessed for

multiple viral targets. SPY utilizes an isopeptide bond within a bacterial adhesin to covalently

link antigens to VLPs or NPs (*90*). The 2c-NP and 1c-SApNP platforms rely on in vitro (cell

free) and in vivo (in producer cells) assembly, respectively. In terms of vaccine production, SPY

and 2c-NP achieve greater versatility at the price of increased complexity and cost, as they both

involve multiple plasmids, expression systems, and purification methods. The 1c-SApNPs

require a highly optimized antigen and NP carrier, which are encoded within a single plasmid, to

achieve manufacturability and quality (*91, 92*). VLPs and NPs are more immunogenic because of

their advantages in antigen trafficking to lymph nodes, high-avidity follicular dendritic cell

(FDC) interactions, antigen-presenting cell (APC) activation, and germinal center (GC)

reactions, which involve both innate and adaptive systems (*77, 93*). A recent study of HIV-1 NP

immunogens suggested that the density of the mannose patches is critical for the FDC targeting

of heavily glycosylated antigens (*94, 95*). Env trimers attached to iron oxide NPs or aluminum

salt have also been tested as multivalent HIV-1 vaccines (*96, 97*).

Here, we displayed the BG505 UFO trimer on multilayered 1c-SApNP platforms (*91, 92*)

as our second-generation vaccine candidates. A panel of Env-NP constructs was first designed

for biochemical, biophysical, and structural characterization to facilitate immunogen selection.

Cryo-EM yielded high-resolution structures for the multilayered E2p and I3-01v9 1c-SApNPs

that each present 20 BG505 UFO trimers, with localized reconstructions obtained for the trimer

at 3.7 and 6.0 Å, respectively. Site-specific glycan analysis and antigenic profiling of the BG505

UFO trimer and 1c-SApNPs bearing wildtype and modified glycans revealed distinct features.

Unexpectedly, endoglycosidase H (endo H) trimming of the glycan shield retained NP binding to

NAbs/bNAbs that require specific glycans for Env recognition and target the CD4bs. Negative-

stain EM revealed differential effects of glycan trimming on the angle of approach for bNAbs

PGT128 and VRC01. Next, we immunized mice and rabbits with UFO trimers and 1c-SApNPs

bearing the wildtype and trimmed glycans, which were formulated with conventional adjuvants.

Glycan trimming diverged NAb responses away from known glycan clusters and glycan holes,

while increasing the frequency of vaccine responders (FVR). The beneficial effect of glycan

trimming on tier 2 NAb elicitation was confirmed in NHP studies. Lastly, we analyzed vaccine

delivery and immune responses at the intra-organ, intercellular, and intracellular levels in the

mouse model. Compared with the soluble trimer, the E2p and I3-01v9 1c-SApNPs showed 420

times longer follicular retention, 20-32 times greater presentation on FDC dendrites, and up-to-4

times stronger GC reactions. Glycan trimming showed little effect on 1c-SApNP trafficking to,

and retention in, lymph nodes. Intact 1c-SApNPs and adjuvants in lymph node tissues were

directly visualized by transmission electron microscopy (TEM). Our study thus presents a

promising strategy to overcome the glycan shield challenges in HIV-1 vaccine development

based on the UFO trimer and 1c-SApNP platforms.

RESULTS

Rational design and characterization of BG505 UFO trimer-presenting 1c-SApNPs

In the UFO design paradigm, the N terminus of heptad repeat 1 ($HR1_N$, aa 547-569) is the trigger

of HIV-1 Env metastability and must be shortened and optimized (*26, 27*). Crystal structures

were determined for clade A, B, and C Envs bearing the $HR1_N$, UFO, and UFO-BG designs (*26,*

27, 71, 98). The $HR1_N$-redesigned BG505 trimer was displayed on ferritin (FR), E2p, and I3-01

NPs (*27, 88*). The reengineered I3-01, when formulated with a Toll-like receptor 3 (TLR3)

agonist adjuvant, elicited a potent tier 2 NAb response to HIV-1 in mice for the first time (*27,*

71). We have also applied these NP platforms to design multivalent immunogens for other

viruses (*89, 91, 92*).

Here we revisited the HIV-1 Env-NP vaccine design based on the high-resolution BG505

UFO trimer structure (*98*) and newly developed "multilayered" 1c-SApNP platforms (*91, 92*).

We first modeled the BG505 UFO trimer (PDB ID: 6UTK) on wildtype FR, E2p, and I3-01,

resulting in particles with diameters of 30.2, 41.7, and 46.1 nm, respectively (**Fig. 1A**). The

redesigned 8-aa $HR1_N$ bend ("NPDWLPDM") forms a 24-Å triangle in the center of each UFO

trimer (**Fig. 1A**, middle). For FR and E2p, the superposition of BG505 gp140 C termini onto the

N termini yielded C_α root-mean-square deviations (RMSDs) of 1.0 and 11.4 Å, respectively. A

3-aa "ASG" linker, with "AS" resulting from the restriction site between the Env and NP genes,

enabled the proper display of Env trimers on FR and E2p. For I3-01, a 10-aa $(G_4S)_2$ linker was

added to overcome the large spacing (50.5 Å) between the N termini of three subunits around

each three-fold axis (*27, 89, 91, 92*). We then examined the multilayered 1c-SApNP design

concept using a modeling approach (**Fig. 1B**). Previously, we incorporated a locking domain

(LD) and a pan-reactive $CD4^+$ T-cell epitope (PADRE) into E2p and I3-01 60-mers via gene

fusion (*91, 92*). PADRE has been shown to improve antibody responses elicited by recombinant

protein vaccines (*99*). We hypothesize that antigens will fold properly on the NP surface upon

self-assembly, while the LD and PADRE may form an inner layer and a hydrophobic core to

stabilize the NP shell and deliver a strong T-helper signal, respectively (**Fig. 1B**, left). E2p-LD4-

PADRE, which was used as a carrier for EBOV GP (*92*), was modeled to visualize the

multilayered structure (**Fig. 1B**, right). Briefly, E2p forms a protein shell of 23.2 nm with a

distance of 12.8 nm across the hollow interior measured at the inward-facing tip (G382). LD4, a

58-aa dimeric protein domain (PDB ID: 2MG4), was fused to the C terminus of E2p at the

dimeric NP-forming interface with a 5-aa G_4S linker, resulting in a smaller protein shell of 14.4

nm when icosahedral symmetry was applied. A 13-aa PADRE with an extended backbone was

fused to the C terminus of LD4 with a 2-aa GS linker, resulting in a hydrophobic core with a

diameter of ~5 nm. Based on structural modeling, a set of multilayered BG505 UFO Env-NP

constructs was designed for experimental validation (*92*).

Nineteen BG505 UFO Env-NP constructs were transiently expressed in 100-ml

ExpiCHO cells, followed by PGT145 purification and size-exclusion chromatography (SEC) on

a Superose 6 column (**Fig. 1C**). Overall, the BG505 UFO trimer can be displayed on wildtype

FR, E2p, and I3-01 NPs with high purity. Similar to our previous study (*92*), seven LDs (1-7)

and five LDs (4-5 and 7-9) were tested for E2p and I3-01, respectively. E2p-LD4 and I3-01-LD7

showed the highest NP yield among various NP-LD combinations. Adding PADRE increased the

NP yield by 5.2- to 8.6-fold relative to the wildtype NPs, measured by ultraviolet (UV)

absorbance at 280 nm. I3-01v9, a variant of I3-01 (*27*), further improved NP yield by 24% based

on the UV value. Seven SEC-purified NPs were analyzed by sodium dodecyl sulfate–

polyacrylamide gel electrophoresis (SDS-PAGE) under reducing conditions (**Fig. 1D**). E2p and

I3-01 registered higher bands than FR on the gel (~150 kDa). These seven NP samples were then

analyzed by blue native polyacrylamide gel electrophoresis (BN-PAGE) (**Fig. 1E**). While FR

displayed a distinctive NP band, none of the E2p and I3-01 NPs migrated down from the

chambers due to their large size. The SEC-purified FR, E2p-LD4-PADRE (E2p-L4P), and I3-

01v9-LD7-PADRE (I3-01v9-L7P) were visualized by negative stain EM (**Fig. 1F**).

Thermostability was assessed by differential scanning calorimetry (DSC) (**Fig. 1G**, **Fig. S1A**).

The melting temperature (T_m) of the three NPs was comparable to that of the BG505 UFO trimer

(66.9-68.3°C *vs.* 68.4°C) (*27*). The multilayered E2p displayed a second peak in the thermogram

(T_{m2} = 74.0°C), suggesting a stepwise unfolding process upon heating. The distribution of

hydrodynamic diameter (D_h) was characterized by dynamic light scattering (DLS) using a

Zetasizer (**Fig. 1H**, **Fig. S1B**). Ferritin and I3-01v9 had an average particle size of 43.7 and 50.8

nm, respectively, whereas E2p displayed a wide distribution (50.8-68.7 nm).

The in vitro assessment of EBOV GP-NP (*92*) and HIV-1 Env-NP constructs

demonstrated the advantages of multilayered protein NPs. In this study, FR, E2p-L4P, and I3-

01v9-L7P were used to present the BG505 UFO trimer as particulate HIV-1 immunogens (**Fig.**

S1C). DSC and DLS suggested that the multilayered E2p may adopt two distinct states in

solution. Nonetheless, the two multilayered SApNPs provide a general platform for designing

VLP vaccines.

Cryo-EM analysis of multilayered E2p and I3-01v9 NPs displaying the BG505 UFO trimer

Cryo-EM played a critical role in determining high-resolution structures of native-like Env

trimers (*47-49*). Recently, cryo-EM was used to characterize 2c-NPs that display diverse Env

trimers (*100-103*). In most cases, nanometer resolution was obtained for the NP backbone, with

4-7 Å resolution achieved for the displayed trimers after localized reconstruction (*101, 102*).

Here, we characterized the two multilayered BG505 UFO trimer-presenting 1c-SApNPs by cryo-

EM.

Prior to cryo-EM, the PGT145/SEC-purified BG505 UFO trimer-presenting E2p-L4P and

I3-01v9-L7P NP samples were analyzed by negative stain EM to confirm their homogeneity

(**Fig. S2A**). In the micrographs, most NPs displayed appropriate size and morphology with only

minimal quantities of free trimers and partially assembled/disassembled particles. We then

applied cryo-EM to characterize the two NP immunogens (**Fig. 2**), with the data processing

protocol described in the Methods section (*101, 102*) and data acquisition parameters listed in

Table S1. The BG505 UFO trimers are attached to the E2p and I3-01v9 scaffolds using peptide

linkers (**Fig. S1C**). Due to conformational flexibility, the antigen-corresponding signal (or

density) was scattered in the 2D classes and initial 3D maps (**Fig. 2A-B**, top left). Consequently,

we were unable to reconstruct high-resolution maps of full BG505 UFO trimer-presenting E2p-

L4P and I3-01v9-L7P NPs. To overcome this problem, we applied a previously developed data

processing method, in which two flexibly linked entities (i.e., the NP scaffold and Env trimer

here) are analyzed independently (*101*). The resulting maps are represented as a transparent gray

mesh (**Fig. 2A-B**). The E2p-L4P NP consists of the E2p shell (based on dihydrolipoyl

transacetylase, PDB ID: 1B5S (*104*)), LD4 layer (PDB ID: 2MG4), and PADRE core (**Fig. 1B**).

Focused refinement of the NP core converged on the E2p shell, resulting in a 3.7 Å EM map; and

the E2p model was then relaxed into the map (**Fig. 2A**, right). The resulting structure displayed

excellent agreement with the crystal structure of this bacterial enzyme (*104*) with a C_α RMSD of

0.9 Å, suggesting a negligible effect of the surface-displayed trimers and encapsulated LD4 layer

and PADRE core on particle assembly. Although additional density at the core of the NP and

along the five-fold axes was observed in the EM map, attempts to resolve their structures were

unsuccessful (**Fig. S2B-C**). Therefore, although LD4 and PADRE can significantly improve NP

expression (**Fig. 1C**), they appear to have a highly flexible structure. Based on the EM data, the

LD4s may not assemble into a tightly knit shell as depicted by computational modeling (**Fig. 1B**)

and likely interact with each other in a less well-organized fashion. The I3-01v9-L7P NP

exhibited inherent flexibility due to the 10-aa GS linkers connecting the surface trimers and NP

backbone, resulting in relatively low-resolution EM maps. Nonetheless, good fits were obtained

by docking the structures of I3-01 (PDB ID: 5KP9 (*105*)) and BG505 UFO trimer (PDB ID:

6UTK (*98*)) into the corresponding maps (**Fig. 2B**).

Altogether, cryo-EM confirmed the correct assembly of two multilayered 1c-SApNPs

and native-like structural features of surface-displayed BG505 UFO trimers. Given the wide

particle size distribution (**Fig. 1H**), we speculate that both our cryo-EM model and the previously

reported crystal structure (*104*) captured the ground state of E2p, in which the N-terminal loop

(A184-S203) is packed against the NP surface and E2p adopts the most compact NP

conformation.

Site-specific glycan analysis of BG505 UFO immunogens with wildtype and modified

glycans

The glycan shield is a defense mechanism used by HIV-1 to escape antibody neutralization (*4*).

However, the identification of glycan-reactive bNAbs (*6*) suggests that it can also be exploited as

a vaccine target (*32, 33, 36*). The structure of the glycan shield has been extensively

characterized to facilitate rational vaccine design (*31*). Here, we determined the site-specific

glycan profiles for BG505 UFO trimers and NPs using LC-MS and a previously established

protocol (*27, 71*).

Wildtype BG505 UFO Env glycosylation (**Fig. 3A**, **Fig. S3A**) is consistent with previous

reports on other BG505 Envs (*27, 38*). The trimer-associated mannose patches (TAMPs) at

N156, N160, N276, and N301 present primarily oligomannose-type glycans, suggesting correct

folding of the Env-NP fusion protein. This processing state is conserved across all samples,

indicating that particulate display does not disrupt glycan processing near the trimer apex. The

intrinsic mannose patch (IMP) around N332 consists of predominantly $Man_9GlcNAc_2$ glycans

(*27, 38*). This pattern was observed for all samples tested, demonstrating that the presentation of

this glycan supersite is unaffected by multivalent NP display. Complex-type glycans can be

found at gp120 N88, N185e, N301/N355 (although high mannose predominates), N398, N462,

and across the N611, N618, and N637 glycans on gp41 (**Fig. S3A**) (*27, 38*). Several potential *N*-

linked glycosylation sites (PNGS) lack an attached glycan at a proportion of the sites. For

example, N133, N137, N185e, and N197 on gp120 are underoccupied, similar to the BG505

SOSIP trimer (*50*). Treatment with swainsonine did not enrich hybrid-type glycans as expected

from glycosidase inhibition, with a nearly identical glycan profile to the wildtype trimer (**Fig.**

3A, **Fig. S3B**). Treatment with kifunensine produced predominantly $Man_9GlcNAc_2$ glycans at

most sites, except for certain sites that were still occupied by $Man_{5-8}GlcNAc_2$ due to

mannosidase trimming (**Fig. 3A, Fig. S3C**).

BG505 UFO trimer-presenting SApNPs behaved similarly in terms of glycan processing

and occupancy (**Fig. 3A**, **Fig. S3D-F**). Moderate differences were noted for N133 and N137,

where FR is less occupied than E2p and I3-01v9, whereas the same FR sample is more occupied

at N185h compared to all other samples. However, soluble trimers and SApNPs showed

markedly different glycan profiles. Variations in glycan occupancy were noted for N133, N137,

N185e, N185h, and N197 on gp120. Trimers were less occupied than SApNPs at all four glycan

sites on gp41. This pattern was most pronounced for N625, showing a ~50% increase in

occupancy for three SApNPs. Since glycan occupancy normally decreases toward the C terminus

(*106*), a possible explanation is that the NP subunit in effect shifts the relative position of gp41

and becomes the new C terminus of the Env-NP fusion protein. Nevertheless, the increased

glycan occupancy on gp41 will likely be beneficial and diminish the elicitation or binding of

nNAbs to the gp41 base and perhaps the NP backbone. The processing of oligomannose-type

glycans also differs between soluble trimers and NPs, especially for sites outside the mannose

patches. Two gp120 sites, N88 and N462, which present complex-type glycans on soluble

trimers, are less processed on NPs (e.g., N88 shifts from 25% to 90% oligomannose-type

glycans). Restricted glycan processing was also observed for the gp41 sites located proximal to

the base of the NP-displayed Envs. The glycan presented at these gp41 sites is $Man_5GlcNAc_2$,

not unprocessed $Man_9GlcNAc_2$ or a complex-type glycan.

Endoglycosidase H (endo H) cleaves the chitobiose core of oligomannose- and hybrid-

type glycans and leaves a single GlcNAc residue attached to the asparagine. Endo H has been

used in the glycan analysis of diverse HIV-1 Envs (*39*). Here, the PGT145/SEC-purified E2p and

I3-01v9 SApNPs were treated by endo Hf (a fusion of endo H and maltose-binding protein

[MBP]) at 25°C for 5 hours and purified again by SEC (**Fig. 3B**). Due to the significant

difference in their molecular weights, endo Hf appears as a well-separated peak in the SEC

profile. However, a rapid enzyme-linked immunosorbent assay (ELISA) using an MBP-specific

mouse antibody detected residual endo Hf in SEC-purified SApNPs (**Fig. S3G**). Reducing SDS-

PAGE showed lower bands for the E2p and I3-01v9 SApNPs after endo H treatment (**Fig. 3C**),

with the structural integrity of glycan-trimmed SApNPs confirmed by negative stain EM (**Fig.**

3D). Comparable T_m values from DSC suggest that glycan trimming has little effect on the

thermostability of SApNPs (**Fig. 3E**). Next, reducing SDS-PAGE was used to analyze E2p and

I3-01v9 SApNPs treated with endo H, which was subsequently removed by the MBP-specific

resin (see Methods), at 25°C *vs.* 37°C after 100°C denaturation (**Fig. 3F**). The denatured material

showed the lowest band on the gel, suggesting that some glycans may be protected by the Env

structure. Lastly, we used ultra-high performance liquid chromatography (UPLC) with in-gel

digestion to determine the global glycosylation status before and after endo H treatment with

subsequent enzyme removal (**Fig. 3G**). Our analysis suggests that all oligomannose-type glycans

on SApNPs can be cleaved by endo H. Notably, almost all Man5-7 glycans were trimmed by

endo H, whereas some Man8-9 glycans remained intact after treatment, with a Man9 signal of

approximately 40% relative to the untreated sample**.**

Antigenicity of BG505 UFO immunogens with wildtype and modified glycans

Native-like Env trimers of diverse origins that contain the modified HR1$_N$ bend or full

UFO/UFO-BG designs have been assessed against panels of bNAbs and nNAbs (*26-28*). Here,

BG505 UFO trimers and SApNPs with wildtype and modified glycans were tested for antibody

binding using bio-layer interferometry (BLI). The previously established 19-antibody panel (*27*)

was extended to include human bNAbs 438-B11, VRC34, and SF12, which target the N332

supersite (*98*), FP (*107*), and SF (*108*), respectively, a potent mouse NAb (M4H2K1) that binds

the C3/V4 region (*71*), and an NHP-derived nNAb (RM20A3) that recognizes the trimer base

(*109*).

The peak antibody-binding signals were summarized according to immunogen valency

and glycan treatment (**Fig. 4A**), with association and dissociation curves plotted for six

concentrations (**Fig. S4**). Analytical sensors were used for soluble trimers (**Fig. 4A**, top).

Overall, the wildtype BG505 UFO Env preferentially bound to (b)NAbs and not nNAbs, except

for V3 tip-directed 19b and 446-52D and base-directed RM20A3 (**Fig. S4A**). Kifunensine

treatment reduced Env binding to (b)NAbs that either interact with or require the accommodation

of glycans (**Fig. S4B**). PGT151, a bNAb that interacts with complex-type glycans at N611 and

N637 (*47, 110-112*), showed undetectable binding to the BG505 UFO Env that displays unprocessed glycans at these two sites. VRC01 binding was made more difficult by the modified glycan barrier around the CD4bs due to the inhibition of ER α mannosidase I in CHO cells. Endo H treatment resulted in an overall reduction of Env binding to glycan-reactive (b)NAbs (**Fig. S4C**). Compared with the apex bNAbs, glycan trimming had a less severe but still noticeable impact on bNAbs to the glycan-V3 supersite (e.g., PGT121 and PGT128), which interact with the GDIR motif and high-mannose glycans at N332 and N301 (*44, 48, 113*). PGT151 barely recognized the glycan-trimmed BG505 UFO Env. Endo H treatment largely retained accessibility of the CD4bs to the VRC01 bNAb (*114, 115*) upon trimming of the glycan barrier surrounding this proteinaceous surface. Notably, glycan modification had little effect on a SF-directed bNAb (SF12) that interacts with N262, N295, and N448 (*108*), as well as the base-directed nNAb, RM20A3 (*109*).

Antigenicity was then determined for three SApNPs displaying wildtype and modified Env glycans, with respective trimers included as a control (**Fig. 4A**, bottom). Quantitation sensors were used in the Octet assays to measure the avidity effect of SApNPs (*91, 92*). Wildtype SApNPs (**Fig. S4D-G**) bound at elevated levels to (b)NAbs that target the V2 apex (×2.5-4.7), N332 supersite (×2.0-3.9), and CD4bs (×1.4-1.6) and at reduced levels to (b)NAbs that recognize the FP (×0.4-0.6) and gp120/gp41 interface (×0.2-0.7). When tested against nNAbs, enhanced binding was only observed for the V3 tip (×2.6-4.0) but not the non-neutralizing CD4bs, CD4-induced (CD4i), gp41, and base epitopes. Thus, the multivalent display can exert a positive influence, as well as structural constraints, on antibody recognition of an Env epitope depending on the location and accessibility of this epitope on the NP surface (*27, 88, 101, 102*). Notably, the wildtype SApNPs only displayed a moderate increase in VRC01 binding with a slow on-rate,

suggesting the CD4bs is more occluded compared to the V2 apex and glycan-V3 supersite on the

NP surface. SApNPs bearing unprocessed $Man_9GlcNAc_2$ glycans exhibited comparable antigenic

profiles to wildtype SApNPs (**Fig. S4H-K**). Similar to the trimer presenting unprocessed

glycans, no PGT151 binding was observed. A similar but less severe effect was noted for

VRC34, which interacts with the N terminus of FP and the complex-type N88 glycan (*107*).

Endo H-treated SApNPs (**Fig. S4L-O**) demonstrated unique antigenic profiles with marked

improvement in VRC01 binding (×2.0-2.3). Slightly lower binding signals were observed for

most glycan-specific NAbs and bNAbs except for two. PG16 binds to a high-mannose glycan at

N160 and a hybrid-type glycan at N173 (*41*). While the wildtype SApNPs bound to PG16 with

signals 3.4-4.6-fold relative to the trimer, glycan trimming showed a greater fold change.

PGT135 requires $ManGlcNAc_2$ at N386, $Man_6GlcNAc_2$ at N332, and $Man_8GlcNAc_2$ at N392 in

Env recognition (*116*). The glycan-trimmed SApNPs showed higher PGT135-binding signals

than the wildtype SApNPs, 2.3-2.8 and 2.0-2.4-fold with respect to their trimers. PGT151

binding was barely detectable for three SApNPs, but increased binding was observed for another

interface bNAb, 35O22 (*117*), and for the FP bNAb, VRC34 (*107*). Two nNAbs to the V3 tip,

19b and 446-52D, bound to glycan-trimmed SApNPs at comparable levels to wildtype SApNPs.

The nNAb RM20A3 (*109*) showed negligible binding to SApNPs independent of their surface

glycans, likely because the trimer base would become nearly inaccessible on the NP surface.

Wildtype E2p was tested in duplicate to estimate the signal variation, which was within 2.5% for

all antibodies with detectable binding, except for SF12 which showed a variation of 5.6% (**Fig.**

S4P). Therefore, experimental variation would have had little effect on the patterns observed in

the BLI analysis.

Negative stain EM was utilized to characterize bNAb interactions with the glycan-

trimmed BG505 UFO trimer (**Fig. 4B**, **Fig. S4Q**). PGT128 binds the V3 base and high-mannose

glycans at N295 (absolutely required) and N301 or N332 (*44, 48, 118*). Previous structures

suggest that the mannose moieties of these glycans serve as anchors to orientate and stabilize the

interaction with PGT128 Fab, which binds the "GDIR" motif with the HCDR2 and HCDR3 tips

(*44, 48*). The removal of mannose groups at these three sites may destabilize PGT128 and affect

its angle of approach. Indeed, one endo H-treated BG505 UFO trimer could bind to one to three

PGT128 Fabs, with less resolved density for the bound Fabs in the 2D class averages (**Fig. 4B**).

Comparison with the cryo-EM model of the BG505 SOSIP/PGT128 complex (EMDB-3121

(*48*)) confirmed that PGT128, albeit with greater flexibility, still bound to the same epitope.

However, the angle of approach could not be determined in this EM analysis. VRC01 targets a

large proteinaceous area of the CD4bs protected by a ring of glycans with the N276 glycan

positioned as a critical barrier to the maturation of VRC01-class bNAbs (*114, 115, 119, 120*).

Trimming the glycan shield can, in principle, improve the CD4bs recognition by VRC01-class

bNAbs. Indeed, we obtained a 12.2 Å-resolution 3D model of the BG505 UFO/VRC01 complex,

which superimposes well with the previous EM models of BG505 SOSIP/VRC01 complexes

(*121, 122*) (**Fig. 4C**).

BLI demonstrated a cross-panel improvement in bNAb binding to SApNPs compared

with soluble trimers, consistent with our previous studies (*27, 89, 91, 92*). Glycan trimming

retained bNAb binding to major glycan epitopes on the NP-displayed Env and improved

recognition of the CD4bs by VRC01-class bNAbs. Negative-stain EM confirmed that bNAbs

PGT128 and VRC01 could still recognize their epitopes on the glycan-trimmed Env trimer.

Altogether, glycan-trimmed SApNPs may provide more effective immunogens due to more

balanced epitope accessibility.

Assessment of HIV-1 UFO Env immunogens with wildtype glycans in mice and rabbits

While BG505 SOSIP and $HR1_N$-redesigned trimers were ineffective in NAb induction in mice

(*27, 55*), a reengineered I3-01 NP displaying the $HR1_N$-redesigned trimer elicited potent tier 2

murine NAbs to the C3/V4 epitope (*27, 71*). Recently, an autologous tier 2 NAb response was

induced in mice by iron oxide NP-attached SOSIP trimers (*96*) and a DNA trimer vaccine (*70*)

using long-boost intervals. In rabbits, Env-induced tier 2 NAb responses mainly target glycan

holes (*56*).

We first assessed wildtype BG505 UFO immunogens formulated with three commonly

used adjuvants in mice. We adopted an immunization protocol used in our previous vaccine

studies (*27, 89, 91, 92, 123*) (**Fig. 5A**). In brief, groups of eight mice were immunized four times

at 3-week intervals. SApNPs (30 μg/dose) were formulated with AddaVax (AV), aluminum

phosphate (AP), and aluminum hydroxide (AH). A non-adjuvanted group and a heterologous

AV×2/AP×2 group were included as a negative control and for comparison, respectively. We

performed TZM-bl neutralization assays using purified mouse immunoglobulin G (IgG) from the

last time point, week 11 (**Fig. 5B**, **Fig. S5A**). Without adjuvant, both SApNPs failed to elicit tier

2 NAb responses to BG505.T332N, but I3-01v9 appeared to be more effective against a tier 1

clade B virus, SF162 (**Fig. S5A**, middle). E2p mixed with AV or AP and I3-01v9 mixed with AP

or AH each showed tier 2 NAb induction in one of the eight mice (**Fig. S5A**, top). However,

switching the adjuvant during the immunization proved ineffective. Adjuvanted SApNPs

induced a robust tier 1 NAb response (**Fig. 5B**, **Fig. S5A**, middle). When tested against a BG505

variant with a C3/V4 epitope knockout mutation, I396R (**Fig. 5C, Fig. S5B**), reduced

neutralization (by 45-82%) was noted for all four mice that generated tier 2 NAb responses,

suggesting that M4H2K1-like NAbs (*71*) may be present in the serum. The remaining

neutralizing activity may be attributed to C3/V5 (*70*) and other Env epitopes. As a negative

control, purified IgG from three tier 2 positive mice was tested against pseudoparticles (pps)

bearing the murine leukemia virus (MLV) Env, MLV-pps, and did not show any neutralization

(**Fig. S5A**, bottom). Borderline neutralization of clade A 398F1 was observed in the TZM-bl

assays against a 12-virus global panel (**Fig. S5C**).

We then tested an extended set of wildtype UFO immunogens in rabbits. Briefly, groups

of six rabbits were immunized three times at weeks 0, 8, and 24 with blood draws at 12 time

points throughout immunization (**Fig. 5D**). All antigens (100 μg/dose) were mixed with AV

except for I3-01v9 SApNPs, which were paired with AP. While the BG505 UFO trimer and its

SApNPs (FR, E2p, and I3-01v9) were the main focus of this rabbit study, a cocktail mixing

equal amounts (33 μg/dose) of SApNPs that present clade A BG505 UFO, clade B H078.14

UFO-BG, and clade C CH505 UFO-BG trimers was created for each NP platform, resulting in

seven groups in total. Longitudinal NAb responses of rabbit sera to BG505.T332N, as measured

by 50% inhibitory dilution (ID_{50}) titers, were examined (**Fig. 5E**). While the overall kinetics

were similar across all groups, the cocktail groups exhibited an NP platform-dependent pattern in

their NAb responses to BG505.T332N. Reducing the BG505 dose to one-third for each cocktail

formulation had negative and negligible effects for FR and I3-01v9, respectively. The tier 2 NAb

response became detectable at week 10 (**Fig. 5E**, above dotted line). The ID_{50} titers reached the

highest point at week 26, with both I3-01v9 groups outperforming the BG505 UFO trimer, FR

cocktail, and both E2p groups (**Fig. 5E**, **Fig. S5D**). The advantage of particulate display was also

evident when a tier 1 clade B SF162 was tested, with a detectable response first observed at week

4 for the two I3-01v9 groups (**Fig. S5E**). Rabbit samples at weeks 0, 2, 10, and 26 displayed no

MLV-pp reactivity (**Fig. S5F**). We then investigated the prevalence of glycan hole NAbs in the

total polyclonal NAb response. Plasma neutralization against four BG505.T332N mutants,

Q130N, S241N, P291T, and T465N as previously described (*71*), was tested for select samples

(**Fig. 5F**). Filling glycan holes partially depleted neutralizing activity, consistent with our

previous study (*71*). The trimer group exhibited a more visible reduction against the 241/289

glycan mutants, whereas all SApNP groups appeared to become more sensitive to the T465N

mutation. Finally, all SApNPs failed to generate a strong cross-clade NAb response to other tier

2 isolates in a 12-virus global panel, and no advantage in breadth was seen for the Env mixture

vaccines (**Fig. S5G**).

Assessment of BG505 UFO Env immunogens with trimmed glycans in mice and rabbits

We investigated how glycan trimming affects Env-elicited NAb responses in mice and rabbits by

testing a homologous regimen ("regimen 1") with glycan-trimmed immunogens in all doses and

a heterologous prime-boost regimen ("regimen 2") with glycan-trimmed and wildtype

immunogens in doses 1-2 and 3-4, respectively (**Fig. 6A**). The ExpiCHO-produced BG505 UFO

trimer was treated with endo Hf and purified by SEC and an additional resin step (see Methods).

Eight and five CHO-K1 cell clones were developed to stably express BG505 UFO trimer-

presenting E2p and I3-01v9 SApNPs, respectively. The SApNPs produced by individual cell

clones were inspected by negative-stain EM (**Fig. S6A**) and then pooled for glycan trimming by

endo H.

In the mouse study, groups of eight BALB/c mice were immunized four times, at three-

week intervals. The BG505 UFO trimer and SApNPs (30 µg/dose) were formulated with AV or

AP adjuvants, with the immunization experiment repeated for some formulation/regimen groups

to confirm their immunogenicity. The frequency of vaccine responders (FVR, the ratio of

subjects that generate an autologous tier 2 NAb response) was calculated for comparison.

Overall, glycan trimming (GT) by endo H substantially improved the FVR for the UFO trimer

(FVR up to 86%; **Fig. 6B**, left), E2p SApNP (up to 88%; **Fig. 6B**, middle), and I3-01v9 SApNP

(up to 50%; **Fig. 6B**, right), in addition to the improved IC_{50} values and less within-group

variation compared with the equivalent formulations using wildtype immunogens (**Fig. S6B-D**).

The use of the AP adjuvant was found to be associated with a higher FVR independent of the

Env immunogens tested, while regimen 2 (two GT primes + two WT boosts) appeared to have a

slight advantage for the two SApNP/AP formulations. Glycan trimming also increased the tier 1

NAb response against clade B SF162 while showing no nonspecific MLV reactivity in TZM-bl

assays (**Fig. S6B-D**). We then tested the tier 2 positive IgG samples from eight groups against

BG505.T332N carrying a C3/V4 epitope knockout (I396R) mutation (**Fig. 6C, Fig. S6E**).

Notably, most samples from the SApNP groups escaped this immunodominant glycan epitope, in

contrast to those in the trimer groups, which still targeted the C3/V4 epitope. Vaccine responders

from the trimer and SApNP groups, subject to IgG sample availability, were further tested

against a 12-virus global panel (**Fig. S6F**). The less stringent IC_{30} titers showed some breadth,

with marginal to appreciable neutralization of 398 F1 (clade A), CNE8 (CRF01 AE), and a few

other isolates of diverse subtypes.

A rapid screening study was performed in rabbits following the mouse protocol (**Fig.**

6A). Four groups of six rabbits were immunized to compare the E2p/AV and I3-01v9/AP

formulations each combined with regimens 1 and 2. While purified IgG at week 0 exhibited a

clean background (**Fig. S6G**), purified IgG at week 11 neutralized autologous tier 2

BG505.T332N with patterns largely resembling the mouse study. (**Fig. 6D, Fig. S6H**). Namely,

BG505 I3-01v9 formulated with AP outperformed E2p paired with AV, and, between the two I3-

01v9 groups, regimen 2 was slightly more effective than regimen 1 with a higher FVR (4 *vs.* 3

respectively; **Fig. 6E**). Similar to the mouse study, purified IgG samples from weeks 0 and 11

were also tested in TZM-bl assays against tier 1 clade B SF162 and the negative control, MLV

(**Fig. S6G and H**). We then analyzed the prevalence of glycan hole NAbs in select rabbit

samples from each group (**Fig. 6F, Fig. S6I**). Interestingly, purified IgG samples from the

BG505 I3-01v9 regimen 2 group exhibited notably more complete escape from the four glycan

holes, supporting our notion that glycan trimming may offer a general solution to overcome

glycan holes in Env immunization. Lastly, select rabbit IgG samples were tested against the 12-

virus global panel (**Fig. S6J**), but the IC_{50}/IC_{30} titers showed less pronounced cross-clade NAb

responses compared with the mouse data (**Fig. S6F**).

Our results revealed the importance of glycan trimming in improving the immunogenicity

of HIV-1 Env vaccines in small animal models. The 50-86% FVR noted for the glycan-trimmed

trimer in mice was rather surprising, given the past failures in tier 2 NAb elicitation by stabilized

BG505 trimers (*27, 55*) and difficulties in improving it (*70, 96*). Glycan trimming not only

enabled antibody access to a broad range of epitopes that would otherwise be occluded by the

glycan shield but also minimized glycan-related immunodominance, such as glycan holes in

rabbits.

Comparison of wildtype and glycan-trimmed BG505 UFO Env-SApNPs in NHPs

NHPs have served as a "gatekeeper" in evaluating HIV-1 Env vaccines (*124-128*). Various

trimer designs, injection routes, and delivery methods have been carefully examined. Native-like

Env trimers, only when formulated with a potent adjuvant, elicited autologous tier 2 NAb

responses in NHPs. The SHIV$_{BG505}$ challenge study demonstrated that NAb titers, but not T cells

or antibody-dependent cell-mediated cytotoxicity (ADCC) activity, correlated with protection

(*125*).

We first tested wildtype SApNPs in NHPs using a regimen modeled after a recent NHP

study of BG505 trimers (*126*) (**Fig. 7A**). Two groups of six rhesus macaques were immunized

three times at weeks 0, 8, and 24, with 13 blood draws throughout the immunization. The E2p

and I3-01v9 SApNPs (200 µg/dose) were formulated with AV and AP, respectively. TZM-bl

neutralization assays were performed to evaluate the NAb response in NHP sera from the last

time point at week 28 (**Fig. 7B**; **Figs. S7A,B**). Overall, wildtype SApNPs failed to elicit an

autologous NAb response to tier 2 clade A BG505.T332N except for one macaque in the

E2p/AV group, while all macaques in both groups showed robust neutralization of tier 1 clade B

SF162 (**Fig. 7B; Fig. S7B**). In the control assays, the pre-immunization samples exhibited little

to no detectable neutralizing activity against HIV-1 (BG505.T332N and SF162) and MLV (**Fig.**

S7A). The poor vaccine immunogenicity observed for wildtype SApNPs is not unexpected. In

previous studies, native-like Env immunogens were often formulated with a potent adjuvant to

induce autologous NAb responses (*124-127*). We next tested glycan-trimmed BG505 SApNPs in

NHPs using two regimens adopted from the mouse study with extended boost intervals (**Fig.**

7C). Four groups of four macaques were immunized four times at weeks 0, 4, 12, and 24 with 9

blood draws throughout the immunization. In this study, E2p and I3-01v9 SApNPs (100

µg/dose) were formulated with the AP adjuvant based on the mouse immunogenicity data (**Fig.**

6B). Overall, glycan trimming appeared to have a positive impact on autologous tier 2 NAb

elicitation compared with wildtype glycans (**Fig. 7D**). Specifically, glycan trimming improved

the FVR for both E2p and I3-01v9 with an FVR of 50% for all four groups except for the I3-

01v9/AP group using regimen 2 (**Fig. 7E**, middle and right); in contrast, wildtype E2p and I3-01v9 induced weak autologous tier 2 NAb responses with FVRs of 16.7% and 0%, respectively (**Fig. 7E,** left). We then analyzed the tier 2 positive NHP sera against the BG505.T332N variant with a T465N mutation in the C3/465 epitope, which is responsible for autologous NAb responses and protection against SHIV$_{BG505}$ challenge in NHPs (*60, 67, 72*) (**Fig. 7F**). Two out of ten samples partially or completely escaped this epitope but still showed tier 2 neutralization. The pre-immunization and week 28 serum samples did not exhibit nonspecific reactivity against MLV in the TZM-bl assays (**Fig. S7C, D**).

Our NHP data suggest that glycan trimming may be a general strategy to improve vaccine-induced NAb responses. The C3/465 epitope remained a main neutralizing target (*60, 67, 72*) after glycan trimming. The finding that glycan-trimmed Env immunogens could be formulated with an aluminum adjuvant to achieve tier 2 NAb elicitation in NHPs is important, because only hints of a tier 2 response were noted for an aluminum-adjuvanted, high-dose BG505 trimer vaccine in humans (*129*). However, the current studies, while promising, were mostly observational. Future NHP studies with optimized matching regimens would be needed to verify these findings.

Distribution, trafficking, and retention of HIV-1 Env trimers and SApNPs in lymph nodes

Following a similar protocol (*123*), we studied the in vivo behavior of the wildtype BG505 UFO trimer and SApNPs E2p and I3-01v9 to understand how they interact with immune cells and induce immune responses. The endo H-treated I3-01v9 SApNP was included to probe the effect of glycan trimming on the immunological characteristics of SApNPs. To induce a robust humoral response, these vaccines must be transported to lymph nodes, accumulate in follicles, retain and present native-like Env, and engage B cell receptors (BCRs) (*130-132*). Here, we first

examined the transport and distribution of BG505 UFO trimer-presenting SApNPs in lymph

nodes via footpad injections (4×10 μg/footpad). Mice were euthanized 12 hours after a single-

dose injection to isolate brachial and popliteal sentinel lymph nodes for histological analysis.

Immunostaining by human bNAbs VRC01(*115*), PGT124 (*133*), and PGDM1400 (*134*) was

used to detect the BG505 UFO Env presented on the SApNPs in lymph node tissue sections (**Fig.**

8A). Consistent with our previous studies of SARS-CoV-2 spike SApNPs (*123*) and ovalbumin-

conjugated gold NPs (*135*), BG505 UFO trimer-presenting SApNPs accumulated in the centers

of lymph node follicles at 12 hours after a single-dose injection (**Fig. 8A**, images on the left; **Fig.**

8B, schematics on the right). Immunostaining images obtained from all three bNAbs demonstrate

similar distributions in lymph nodes (**Fig. 8A-B**, **Fig. S8A-B**). VRC01 (*115*) was selected to

study the trafficking of four BG505 Env immunogens in lymph nodes because of its optimal

signal-to-noise ratio.

We next determined the trafficking and retention patterns of the BG505 UFO trimer and

three SApNPs in lymph node follicles 30 minutes to 8 weeks after a single-dose injection (4×10

μg/footpad) (**Fig. 8C**). The injected dose was normalized to 40 μg of protein per mouse, or 10 μg

protein per footpad. Histological images indicated that both the BG505 UFO trimer and SApNPs

were transported into lymph nodes and accumulated in the subcapsular sinus within 30 minutes

(**Fig. 8C**). The BG505 UFO trimer was trafficked into lymph node follicles at 2 hours and

cleared within 12 hours. In contrast, the three SApNPs first appeared in lymph node follicles at 2

hours, further accumulated at 12 hours, peaked at 48 hours, and remained detectable over a

period of 5-8 weeks (**Fig. 8C**). The VRC01-stained area was quantified in a time-dependent

manner, showing a ~420-times longer retention for the three SApNPs *vs*. the soluble trimer (**Fig.**

8C-D). The area under the curve (AUC) suggested that the exposure of SApNPs in lymph node

follicles is 20-32 times higher than the soluble trimer (**Fig. 8E**). At 48 hours, a significantly

greater accumulation, 46-66 times, was found when the BG505 UFO trimer was presented

multivalently (**Fig. 8F**). These findings are consistent with our previous studies (*123, 135*), in

which small 5-15 nm particles are cleared from the follicles within 48 hours but large 50-100 nm

particles can persist for weeks. The BG505 UFO trimer-presenting SApNPs displayed

distinctively long retention compared with the SARS-CoV-2 spike SApNPs (5-8 weeks *vs.* ~2

weeks), which correlates well with their antigen thermostability (T_m), 68.4°C for the BG505

UFO Env *vs.* 47.6°C for the S2GΔHR2 spike (*27, 91*), respectively. We also noted that glycan

trimming had minimal impact on SApNP distribution, trafficking, and retention in lymph nodes,

in contrast to a recent report (*95*).

Antigen retention and presentation in lymph node follicles are critical for B cell

stimulation and affinity maturation through GC reactions (*130, 136*). FDCs form a network

structure and play roles in antigen collection and presentation in the center of lymph node

follicles (*131, 132*). FDCs retain soluble antigens, immune complexes, virus-like particles,

viruses, and bacteria, and induce GC initiation and maintenance (*131, 135, 137-139*). Based on

our previous studies (*123, 135*), we hypothesized that FDC networks are the major resident

places for these BG505 Env immunogens. To test this hypothesis, we injected mice with the

UFO trimer and three SApNPs through footpads (4×10 μg/footpad). Mice were sacrificed to

collect sentinel lymph nodes at the peak of SApNP accumulation (48 hours) and other timepoints

(30 minutes to 8 weeks) after a single-dose injection (**Fig. 8G**, **Fig. S8C-I**). Lymph node tissues

were stained with bNAb VRC01 (*140*) for BG505 Env (white), anti-CD21 antibodies for FDCs

(green), and anti-CD169 antibodies for subcapsular sinus macrophages (red). The signals of

SApNPs colocalized with FDC (CD21$^+$) networks in lymph node follicles at 48 hours (**Fig. 8G**),

confirming the crucial role of FDC networks in mediating the accumulation and retention of

various Env immunogens in lymph node follicles.

How HIV-1 Env SApNPs interact with FDCs and phagocytic cells in lymph nodes

FDC networks create areas of high antigen density in lymph node follicles to facilitate effective

BCR crosslinking, B cell activation, and GC reactions (*77, 130, 131*). Our TEM analysis

revealed that FDCs could align ovalbumin-conjugated gold NPs (*135*) and SARS-CoV-2 spike-

presenting SApNPs (*123*) on their surfaces or dendrites through interactions between

complement protein 3 (C3) and complement receptor 2 (CR2) (*135*). B cells are embraced by

long FDC dendrites to maximize interactions between multivalently displayed antigens and

BCRs (*123, 135*). Here, we studied the interface between FDC dendrites and B cells to better

understand how FDCs process HIV-1 SApNPs formulated with adjuvants to engage B cells. We

injected mice with AV- or AP-formulated SApNPs through their hind footpads (2×50

μg/footpad) (**Fig. 9A-F**, **Fig. S9A-V**). Fresh popliteal sentinel lymph nodes were isolated at 2, 12

and 48 hours after a single-dose injection and processed immediately for TEM analysis. The

TEM images showed that FDC dendrites embrace and interact with B cells in lymph node

follicles (**Fig. 9A-C**, **Fig. S9A-M**). A large number of intact E2p SApNPs (round-shaped

granules, yellow arrows) and AV particles (oil-in-water nano-emulsion of ~150 nm in size, green

arrows) were aligned on FDC dendrites and B cell surfaces at 2, 12, and 48 hours (**Fig. 9A**, **Fig.**

S9A-C). Notably, few E2p SApNPs were internalized in the endolysosomes of B cells at 48

hours (**Fig. S9C**). I3-01v9 SApNPs (yellow arrows), wildtype and glycan-trimmed alike, were

similarly aligned on the FDC dendrites and B cell surfaces at 2, 12, and 48 hours, without the

colocalization of AP adjuvants (**Fig. 9B-C**, **Fig. S9D-I**). Of note, AV particles alone were

presented on the FDC dendrites 12 hours after injection (**Fig. S9L**).

Phagocytic cells, such as macrophages and dendritic cells in the subcapsular sinus and

medullary sinus of lymph nodes, capture large particles, immune complexes, viruses and bacteria

and transport them to migrating B cells and deposit them on the surface of FDCs via a

complement receptor-dependent mechanism (*131, 132, 141-143*). These immune cell populations

also serve as APCs that can further promote adaptive immunity. Here, we studied the association

between macrophages and adjuvanted SApNPs (**Fig. 9D-F**, **Fig. S9N-V**). E2p SApNPs and AV

adjuvant particles are colocalized and aligned on the surface or inside the endolysosomes of

medullary sinus macrophages at 2 hours (**Fig. 9D**, **Fig. S9N**). Wildtype and glycan-trimmed I3-

01v9 SApNPs behaved similarly to E2p SApNPs, with AP forming visible aggregates inside the

endolysosomes of macrophages and in the extracellular matrix (**Fig. 9E**, **Fig. S9O-U**).

Our results reveal how BG505 UFO trimer-presenting SApNPs associate with adjuvants

in the intercellular and intracellular compartments of lymph nodes. Three SApNPs showed

similar patterns of interaction with FDC networks, B cells, and phagocytic cells. Two common

adjuvants with different chemical properties, AV and AP, interact with cells in lymph nodes

through different immune pathways. FDC networks can retain SApNPs on long FDC dendrites

and present them to naïve B cells, which is an intrinsic feature independent of adjuvants (**Fig.**

S9J-K).

HIV-1 Env SApNPs induce more robust germinal center reactions than soluble trimers

B cell somatic hypermutation, selection and affinity maturation occur in long-lived GCs and lead

to the formation of immune memory and development of (b)NAbs upon vaccination (*138, 144-*

147). Here, we studied whether the long-term retention of BG505 UFO trimer-presenting E2p

and I3-01v9 SApNPs can induce more robust and long-lived GCs in follicles than the soluble

UFO trimer. We first assessed GC reactions induced by wildtype I3-01v9 SApNP at 2 weeks

after a single-dose injection (4×10 μg/footpad). GC reactions, i.e., GC B cells (GL7$^+$) and T

follicular helper (T$_{fh}$) cells (CD4$^+$Bcl6$^+$), in sentinel lymph nodes were characterized by

immunohistological analysis. Robust GCs (GL7$^+$, red) were generated that were attached to FDC

networks (CD21$^+$, green) with clearly organized dark zone (DZ) and light zone (LZ)

compartments in lymph node B cell follicles (B220$^+$, blue) (**Fig. 10A**, left). T$_{fh}$ cells (CD4$^+$Bcl6$^+$,

co-labeled with cyan and red) were mainly located in the LZ to support B cell affinity maturation

and GC maintenance (**Fig. 10A**, right). Next, we applied this analysis to the BG505 UFO-trimer

and three SApNPs at 2, 5, and 8 weeks after a single-dose injection (**Fig. 10B**, **Fig. S10A-C**) and

at 2 and 5 weeks after the boost (**Fig. 10C**, **Fig. S10D, E**). Two metrics were defined to quantify

GC reactions based on immunohistological images: GC/FDC ratio (i.e., whether GC formation is

associated with an FDC network, %) and GC size (i.e., occupied area) (*123*). All four Env

immunogens induced robust GCs at 2 weeks after a single-dose injection (**Fig. 10B**, **Fig. S10A**).

The glycan-trimmed I3-01v9 SApNP induced the largest GCs at 2 weeks (**Fig. 10B**). Following

trimer immunization, the GC/FDC ratio and GC size were small and decreased rapidly over time,

whereas a single dose of SApNPs generated long-lived GCs that persisted for 8 weeks (**Fig. 10B,**

D, **Fig. S10C**). For the soluble trimer, robust GCs can be restored after the boost (**Fig. 10C, E**,

Fig. S10D, E). Overall, the three SApNPs generated larger GCs than the soluble trimer, 2.3-4.2

times larger after one dose (**Fig. 10B, D**) and 0.4-1.1 times larger after the boost (**Fig. 10C, E**),

both measured at week 8.

We next characterized GC reactions by flow cytometry. We collected sentinel lymph

nodes at 2, 5, and 8 weeks after a single-dose footpad injection (**Fig. 10F**, **Fig. S11A, B**) and at 2

and 5 weeks after the boost (**Fig. 10F**, **Fig. S11C**) (4×10 μg/injection). Fresh sentinel lymph

nodes were disaggregated into a single cell suspension and stained with an antibody cocktail.

The GC reactions were analyzed by flow cytometry. The percentage and number of GC B cells

and T_{fh} cells were in line with the immunohistological data (**Fig. 10A-E**). The three SApNPs

outperformed the soluble trimer at 2 weeks after a single-dose injection (**Fig. 10F**). Among the

three SApNPs, the glycan-trimmed I3-01v9 elicited the largest GC B cell and T_{fh} cell

populations. GC reactions peaked at 2 weeks for all tested immunogens and declined over time.

The populations of GC B cells and T_{fh} cells induced by the soluble trimer were barely detectable

at 8 weeks after a single-dose injection (**Fig. 10F**). A boost effectively expanded the populations

of GC B cells and T_{fh} cells but had little effect on their percentages (**Fig. 10G**). In addition to

size-dependent trafficking and retention in lymph node follicles, GC reactions may also be

influenced by adjuvants, considering that AV and AP undergo different mechanisms of cellular

interactions (**Fig. 9**). Overall, the three SApNPs generated 3.4-6.6/0.7-1.8 times more GC B cells

and 0.5-4.3/0.2-1.6 times more T_{fh} cells than the soluble trimer at 8 weeks after the single-

dose/boost injection, respectively (**Fig. 10F, G**). Together, our analysis indicates that large

SApNPs can generate long-lived GCs in lymph nodes more effectively than the soluble trimer,

resulting in a strong Env-specific B cell response.

DISCUSSION

Over the last decade, HIV-1 vaccine research has been driven by advances in the identification of

bNAbs targeting diverse epitopes (*6, 7*) and development of native-like Env trimers (*12, 13*).

Three designs have been established as the leading Env trimer platforms, including SOSIP (*19,*

29), NFL (*23, 25*), and UFO (*26, 27*). Multistage vaccine strategies that follow specific B cell

maturation events have been proposed and tested in knock-in mice carrying human bNAb

(precursor) genes (*51*). However, it is unclear whether such strategies will be relevant to human

vaccination. Recent HIV-1 vaccine research also received a boost from the engineering of

protein particles to mimic VLPs (*74-77, 80*). Various animal models have been used to assess

HIV-1 Env vaccines. Rabbits can readily generate autologous tier 2 NAb responses, which

mainly target glycan holes (*56*) or the trimer base (*73*). The tier 2 NAb responses in wildtype

mice and NHPs are sporadic and often target glycan epitopes on the HIV-1 Env (*67, 70, 71*).

Although bNAb responses were occasionally reported (*52, 58, 127, 148, 149*), it is unclear

whether these are reproducible outcomes of a general vaccine solution or coincidence. Therefore,

despite all advances achieved to date, HIV-1 vaccine development still faces critical challenges

posed by the Env, an atypical viral antigen (*150*).

The glycan shield is at the heart of HIV-1 Env vaccine design (*31-36*). Ideally, an

effective vaccine strategy must overcome conflicting roles of Env glycans, namely, occluding

NAb access to conserved protein epitopes *vs.* participating in NAb-Env interactions. Here, we

aimed to develop such a strategy by combining our trimer stabilization (i.e., the UFO design (*26,*

27)) and particulate display (i.e., the multilayered 1c-SAPNPs (*91, 92*)) platforms. The resulting

BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P SApNPs were thoroughly evaluated both in

vitro and in vivo. Notably, a key element in our current vaccine strategy was glycan trimming.

We hypothesized that trimming structurally less essential glycans would improve epitope

accessibility (e.g., CD4bs) and minimize glycan-related immunodominance (e.g., strain-specific

glycan holes or glycan patches), while retaining Env binding to most glycan-reactive bNAbs.

The success of this strategy may hinge on glycan promiscuity (*118*), which allows various glycan

sites and moieties to substitute for one another in glycan-bNAb interactions. The avidity effect

on the NP surface may further compensate for weakened glycan-bNAb interactions on one

trimmed Env trimer by enabling interactions with surrounding Env trimers. As a result, we

observed rather balanced interactions between glycan-trimmed BG505 Env-SApNPs and bNAbs

that target various protein and glycan epitopes (**Fig. 4**).

Mice were used in our large-scale in vivo evaluation because they are inexpensive and

known to generate tier 2 NAbs to epitopes that are found in human infection (*70, 71*). Rabbits

were tested primarily to probe NAb responses to glycan holes. Most common, but not the most

potent (*124, 126, 151*), adjuvants were chosen because of their safety record in humans. Instead

of bNAb elicitation, we focused on increasing the frequency of vaccine responders (FVR,

defined as the ratio of tier 2 positive subjects in a group), because this is more achievable, and

the low FVR has been a major challenge for trimer vaccines in mice, NHPs, and also humans

(*129*). Glycan-trimmed Env immunogens mixed with AP yielded a much higher FVR in mice

than the wildtype ones, 50-88% *vs.* 12.5%, with reduced glycan-related immunodominance in

mice and rabbits (**Fig. 6**). Our NHP studies confirmed the beneficial effect of glycan-trimming in

a primate model, with the FVR reaching 50% for three of the four aluminum-adjuvanted SApNP

groups (**Fig. 7**). An in-depth mechanistic analysis provided much-needed insights into the

vaccine mode of action. The 5-to-8-week retention of E2p and I3-01v9 SApNPs in follicles

indicates sustained B cell maturation, suggesting that a longer interval between immunizations

should be used for these SApNPs. The identical trafficking, presentation and retention patterns

for wildtype and glycan-trimmed I3-01v9 SApNPs indicate that the density of mannose glycans

is not as critical for the follicular localization of HIV-1 NP immunogens as previously thought

(*95*). The stronger GC reactions and $CD4^+$ T_{fh} responses observed for the glycan-trimmed I3-

01v9 SApNP were consistent with early studies where mannoses were immunosuppressive

(*152*), and the enzymatic removal of mannose moieties improved the gp120-induced IgG

response when formulated with alum (*153*).

Future research may focus on several fronts. First, antibodies induced by glycan-trimmed

Env need to be mapped onto the wildtype Env surface (*63, 66*) to understand how glycan

trimming alters epitope targeting, as previously reported for influenza virus and SARS-CoV-2

(*154, 155*). Second, the effect of Env glycans on the immune system warrants a more in-depth

investigation, e.g., the involvement of mannose moieties in immunosuppression (*152*) *vs.* vaccine

trafficking and immunogenicity (*94, 95*). Third, the immunization regimen requires extensive

optimization. The 3-week interval used in our screening regimen is unsuitable for SApNPs with

a retention time of 5 weeks or longer. This may explain why the two large SApNPs did not

substantially outperform the trimer in mice and rabbits, as the 3-week interval would interrupt

the generation of memory B cells and long-lived plasma cells and suppress the recall antibody

response. The interval problem was partially addressed in the NHP study, but the regimen still

requires optimization. A systematic comparison would be needed to identify the most effective

adjuvant (*123*). The resulting vaccine formulation, with an optimized regimen, could then be

considered for evaluation in humans.

METHODS

Antibodies

We utilized a panel of bNAbs and non-NAbs to characterize the antigenicity of various native-

like trimers and gp140 nanoparticles. Antibodies were requested from the NIH AIDS Reagent

Program (https://www.aidsreagent.org/) except for 438-B11(*98*), PGT151 (*110*), and M4H2K1

(*71*), which were produced in-house together with the fragment antigen-binding (Fab) regions of

PGT128 and VRC01, using previously described protocols (*71*).

Expression and purification of HIV-1 Env trimers and SApNPs

The BG505 UFO trimer and SApNPs used for the in vitro characterization and in vivo

assessment of immunogens with wildtype glycans were transiently expressed in ExpiCHO cells

(Thermo Fisher) using a previously described protocol (*27*). Briefly, ExpiCHO[TM] cells were

thawed and incubated with ExpiCHO[TM] Expression Medium (Thermo Fisher) in a shaker

incubator at 37°C, 135 rpm, and 8% CO_2. When the cells reached a density of 10×10^6/ml,

ExpiCHO[TM] Expression Medium was added to reduce the cell density to 6×10^6 ml^{-1} for

transfection. ExpiFectamine[TM] CHO/plasmid DNA complexes were prepared for 100-ml

transfection in ExpiCHO cells following the manufacturer's instructions. Kifunensine (10 mg/L,

Tocris Bioscience) and swainsonine (3.4 mg/L, Santa Cruz Biotechnology) were added at the

time of ExpiCHO transfection to inhibit α-mannosidase I and II to enrich oligomannose- and

complex-type glycans, respectively. For Env constructs tested in this study, 100 μg of antigen

plasmid and 320 μl of ExpiFectamine[TM] CHO reagent were mixed in 7.7 ml of cold OptiPRO™

medium (Thermo Fisher). After the first feed on day 1, ExpiCHO cells were cultured in a shaker

incubator at 32°C, 120 rpm, and 8% CO_2 following the Max Titer protocol with an additional

feed on day 5 (Thermo Fisher). Culture supernatants were harvested 13-14 days after

transfection, clarified by centrifugation at 4000 rpm for 20 min, and filtered using a 0.45 μm

filter (Thermo Fisher). For the BG505 UFO trimer, Env protein was extracted from the culture

supernatants using a *Galanthus nivalis* lectin (GNL) column (Vector Labs), whereas for all

SApNPs, BG505 and non-BG505 (CH505 and H078.14) Env-fusion proteins were purified using

PGT145 and 2G12 columns, respectively. Trimer was purified on a Superdex 200 Increase

10/300 GL column or a HiLoad 16/600 Superdex 200 PG column (GE Healthcare), whereas

SApNPs were characterized on a Superose 6 10/300 GL column. Protein concentration was

determined using UV_{280} absorbance with theoretical extinction coefficients.

Generation of stable CHO cell lines expressing two BG505 UFO-Env NP vaccine candidates

Stable CHO cell lines expressing BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P NPs were generated by the cationic lipid mediated transfection of a parental suspension culture adopted CHO K1 host cell line (ATCC no. CCL-61) using bacterial artificial chromosomes (BACs) as gene transfer vehicles (*156*). BAC vector modification and integration of the BG505 Env-NP expression cassettes by recombineering were performed by Gen-H GmbH (Heidelberg, Germany). Parental CHO K1 host cells were cultured in a 125 ml shake flask (Corning) in 20 ml CD CHO medium (Gibco, Thermo Fisher) supplemented with 8 mM L-glutamine, 6.25 mg/ml phenol red and 1:100 Anti-Clumping Agent (Gibco, Thermo Fisher). Shaker flasks were cultivated in a humidified ISF1-X incubator shaker (Kuhner) at 37°C, 125 rpm, and 5% CO_2, and multi-well plates were maintained in a static incubator cabinet at 37°C and 5% CO_2. The passaging of cells for the purpose of culture maintenance was performed every 3-4 days to starting cell densities of 0.3×10^6 cells/ml. Per transfection, 1×10^6 cells were prepared in 1.8 ml CD DG44 medium (Gibco, Thermo Fisher), 5 µg BAC DNA, and 25 µg Lipofectin® (1 mg/ml; Thermo Fisher) diluted in 0.2 ml of CD DG44 medium, and transfection mixes were incubated for up to 48 hours in six-well plates (Greiner). Following incubation, media of transfection pools were replaced with selection medium (CD CHO medium + supplements and 0.5 mg/ml Geneticin sulphate (Sigma, catalog no. G418). Cultures were transferred to 96-well plates (Thermo Fisher) and transfection cell pools were continuously monitored for signs of cell growth and increasing viabilities, as well as BG505 Env-NP expression by ELISA (NP capture: *Galanthus nivalis* lectin (GNL, Sigma) / NP detection: anti-HIV-1 gp120 mAb C2G12 (Polymun Scientific)). High NP expressing cell pools were expanded to 50 ml of TubeSpin® Bioreactor

tubes (TPP) and cell pools were evaluated for specific growth (μ) and specific NP expression

(qp). The top performing cell pools (2-4) were single-cell-cloned following an in-house

established limiting-dilution procedure in 384-well plates (Corning). Clonality was confirmed

using a Cell Metric plate imaging device (Solentim). Clonal cell populations were again screened

in 96-well plates (Thermo Fisher) by ELISA, and the 20 best performing clonal cell lines were

further evaluated in a small-scale screen (5 ml) in 50 ml TubeSpin® Bioreactor tubes (TPP) with

respect to cell growth and NP expression by ELISA in addition to non-reducing SDS-PAGE and

Western blot. Based on the evaluation results for the top 20 clones, the top eight cell clones were

selected for each BG505 Env-NP and challenged in small-scale fed-batch experiments with the

aim to identify clones that may have the potential to serve as prospective production clones

under good manufacturing practice (GMP) conditions. Fed batch experiments were performed in

125 ml shaker flasks with 45 ml culture volume using ActiPro medium (HyClone) and the

corresponding nutrient feeds Cell Boost 7a and 7b (both HyClone) in a shaking incubator. Small-

scale research cell banks (RCBs) were established and cryo-preserved to support future master

cell bank (MCB) establishment. Cell line development was performed at Polymun Scientific

GmbH (Klosterneuburg, Austria) through a contract from Uvax Bio.

SDS-PAGE and BN-PAGE

BG505 Env-NPs were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis

(SDS-PAGE) and blue native-polyacrylamide gel electrophoresis (BN-PAGE). The proteins

were mixed with loading dye and added to either a 10% Tris-Glycine Gel (Bio-Rad) or a 4-12%

Bis-Tris NativePAGE^{TM} gel (Life Technologies). For SDS-PAGE under reducing conditions, the

proteins were first treated with dithiothreitol (DTT, 25 mM) and boiled for 5 min at 100 °C.

SDS-PAGE gels were run for 20 min at 250 V using SDS running buffer (Bio-Rad), and BN-

PAGE gels were run for 2-2.5 hours at 150 V using NativePAGE^{TM} running buffer (Life

Technologies) according to the manufacturer's instructions. The gels were stained using

Coomassie Brilliant Blue R-250 (Bio-Rad) and de-stained using a solution of 6% ethanol and 3%

glacial acetic acid.

Differential scanning calorimetry (DSC)

Thermal melting curves of BG505 Env UFO-FR, UFO-E2p-L4P, and UFO-10GS-I3-01v9-L7P

SApNPs following PGT145 and SEC purification were obtained from a MicroCal PEAQ-DSC

Man instrument (Malvern). Samples of the E2p and I3-01v9 SApNPs after glycan trimming by

endo Hf and SEC purification were also analyzed. Briefly, the purified SApNP protein was

buffer exchanged into 1×PBS buffer and concentrated to 0.8 μM before analysis by the

instrument. Melting was probed at a scan rate of 60 °C·h^{-1} from 20°C to 100°C. Data processing,

including buffer correction, normalization, and baseline subtraction, was conducted using

MicroCal PEAQ-DSC software. Gaussian fitting was performed using Origin 9.0 software.

Dynamic light scattering (DLS)

Particle size distributions of BG505 Env-NPs based on three NP platforms (FR, E2p-L4P, and

I3-01v9-L7P) were obtained from a Zetasizer Ultra instrument (Malvern). PGT145/SEC-purified

NPs from ExpiCHO cells were diluted to 0.2 mg/ml using 1×PBS buffer, and 30 μl of the

prepared NP sample was added to a quartz batch cuvette (Malvern, catalog no. ZEN2112).

Particle size was measured at 25 °C in back scattering mode. Data processing was performed on

the Zetasizer and the particle size distribution was plotted using Origin 9.0 software.

Endo H treatment and removal

Glycan trimming of Env immunogens was performed using Endo-Hf, a fusion of endo H and

MBP (NEB, catalog no. P0703L), by mixing 20 µg of Env protein, 2 µl of 10×GlycoBuffer3, 5

µl of endo H, and H_2O (if necessary) to make a 20 µl reaction. The mixture was kept at room

temperature (25°C) for 5 fours to facilitate enzymatic processing of the Env glycans. The

reaction volume can be scaled up proportionally to generate more glycan-trimmed material for

immunization. After 5 hours of incubation, the mixture was passed through a Superose 6 column

to remove the MBP-tagged endo H. Both NP and endo Hf fractions were collected for

quantitation of the residual endo Hf using an anti-MBP mouse antibody (Sigma, catalog no.

M6295-.2ML) in ELISA. While most endo Hf can be readily removed from NP samples by SEC,

a second purification step must be included for the trimer because trimer and endo Hf have

separate but close SEC peaks. Amylose resin (NEB, catalog no. E8021S) was used to remove

residual endo Hf from the SEC-purified trimer fractions. The flow through was collected to

quantify the residual endo Hf and then adjusted to appropriate concentrations for in vitro and in

vivo studies. To achieve the most complete glycan trimming, 1-20 µg of the Env protein was

mixed with 1 µl of Glycoprotein Denaturing Buffer (NEB, 10×) and H_2O (if necessary) in a

reaction volume of 10 µl. The reaction was then heated to 100°C for 10 min to denature the

protein, prior to endo H treatment at 37°C for 1 hour. This protocol was only used to test the

maximum trimming of Env glycans by endo H in **Fig. 3G**. The denatured, glycan-trimmed

materials were not used in the animal studies.

Enzyme-linked immunosorbent assay (ELISA)

In this study, ELISA was performed to detect residual endo Hf in trimer and SApNP samples

after glycan trimming. Each well of a Costar^{TM} 96-well assay plate (Corning) was first coated

with 50 µl of PBS containing 0.2 µg of appropriate antigens. The plates were incubated

overnight at 4 °C, and then washed five times with wash buffer containing PBS and 0.05% (v/v)

Tween 20. Each well was then coated with 150 µl of a blocking buffer consisting of PBS and 40

mg ml^{-1} blotting-grade blocker (Bio-Rad). The plates were incubated with the blocking buffer for

1 h at room temperature, and then washed five times with wash buffer. The human bNAb

VRC01 and MBP-specific mouse antibody (Sigma, catalog no. M6295-.2ML) were diluted in

blocking buffer to a maximum concentration of 1 µg/ml and to reach ×466 dilution (as

recommended by the manufacturer), respectively, followed by a 10-fold dilution series. For each

antibody dilution, a total 50 µl volume was added to the appropriate wells. Each plate was

incubated for 1 h at room temperature, and then washed five times with PBS containing 0.05%

Tween 20. A 1:5000 dilution of goat anti-human IgG antibody (Jackson ImmunoResearch

Laboratories) or 1:3000 dilution of goat anti-mouse IgG antibody (Jackson ImmunoResearch

Laboratories) was then made in wash buffer (PBS containing 0.05% Tween 20), with 50 µl of

this diluted secondary antibody added to each well. The plates were incubated with the

secondary antibody for 1 h at room temperature, and then washed five times with PBS containing

0.05% Tween 20. Finally, the wells were developed with 50 µl of TMB (Life Sciences) for 3-5

min before stopping the reaction with 50 µl of 2 N sulfuric acid. The resulting plate readouts

were measured at a wavelength of 450 nm. As a rapid screening ELISA assay, only four

antibody concentrations/dilutions were tested.

Negative stain EM analysis

The initial evaluation of various Env-NP samples was performed by the Core Microscopy

Facility at The Scripps Research Institute. All NP samples were prepared at the concentration of

0.01-0.05 mg/ml. Carbon-coated copper grids (400 mesh) were glow-discharged and 8 µl of each

sample was adsorbed for 2 min. Excess sample was wicked away and grids were negatively

stained with 2% uranyl formate for 2 min. Excess stain was wicked away and the grids were

allowed to dry. Samples were analyzed at 80 kV with a Talos L120C transmission electron

microscope (Thermo Fisher) and images were acquired with a CETA 16M CMOS camera. The

structural analysis of glycan trimmed-treated BG505 UFO trimer bound to bNAbs PGT128 and

VRC01 was performed by the Core Microscopy Facility following the protocol described above.

For each complex, a total of 55 images were manually collected at a magnification of 73,000×.

CryoSPARC 2 software (*157*) on the Scripps Garibaldi cluster was used to analyze the EM

images, including imaging processing, particle picking, 2D classification, and 3D reconstruction

(**Fig. S4P**). The validation of BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P NP samples

prior to high-resolution cryo-EM was performed in the Hazen EM facility at The Scripps

Research Institute. The experiments were performed as previously described (*101, 158*). The

concentrated BG505 Env-NP samples were diluted to 50 µg/ml in TBS buffer (Alfa Aesar) and

loaded onto the carbon-coated 400-mesh Cu grid. Prior to sample application the grids were

glow-discharged at 15 mA for 30 s. The samples were blotted off the grids after 10 s and

negatively stained with 2 % (w/v) uranyl-formate for 60 s. A Tecnai Spirit electron microscope

(120 keV) featuring a Tietz 4k × 4k TemCam-F416 CMOS camera was used for data acquisition.

Nominal magnification was 52,000 ×. The resulting pixel size at the specimen plane was 2.05 Å,

and the defocus was set to -1.50 µm. Total electron dose per image was adjusted to 25 e-/Å2.

Images were recorded using the Leginon software suite (*159*). Data were visualized in the

Appion data processing suite (*160*).

Cryo-EM analysis of two BG505 UFO trimer-presenting SApNPs

The cryo-EM grid preparation was performed as the following. BG505 UFO-E2p-L4P and UFO-

10GS-I3-01v9-L7P samples were concentrated to 1.5 and 2.5 mg/ml, respectively. Immediately

prior to grid application, lauryl maltose neopentyl glycol (LMNG) was added to each sample at a

final concentration of 0.005 mM. Quantifoil R 2/1 holey carbon copper grids (Cu 400 mesh)

were the main type of grids used. The grids were pretreated with Ar/O2 plasma (Gatan Solarus

950 plasma system) for 10 s before sample loading. Vitrobot mark IV (Thermo Fisher Scientific)

was used for the sample application, blotting, and vitrification steps. The temperature was set to

10°C, humidity was 100 %, blotting force was set to 0, wait time was 10 s, and blotting time

varied in the range of 3.5-5.5 s. For grid preparation, 3 µl of the NP sample (with LMNG) was

loaded onto plasma-activated grids. Following the blot step the grids were plunge-frozen into

liquid ethane cooled with liquid nitrogen. The cryo-EM data were collected from two

microscopes. Specifically, the I3-01v9-L7P NP data were collected on the Talos Arctica TEM

(Thermo Fisher Scientific) operating at 200 kV. The E2p-L4P NP data were acquired on an FEI

Titan Krios TEM at 300 kV (Thermo Fisher Scientific). Both microscopes were equipped with

the Gatan K2 Summit direct electron detector camera and sample autoloader. The Leginon

software suite (*159*) was used for automated data acquisition. Data collection parameters are

shown in **Table S1**. Cryo-EM data were processed as described previously (*101*). MotionCor2

(*161*) was used for frame alignment and dose-weighting and GCTF was applied for the

estimation of CTF parameters. The early processing steps were performed in CryoSPARC 2

(*157*). Particles were picked using template picker, extracted and subjected to two rounds of 2D

classification to eliminate bad picks and heterogeneously looking particles. After the 2D cleaning

step, 17097 and 4806 particles were retained for BG505 UFO-E2p-L4P and UFO-I3-01v9-L7P

NPs for further processing steps. These particle subsets were transferred to Relion/3.0 (*162*). Ab-

initio reconstruction in cryoSPARC was used to generate the starting reference models for the

3D steps in Relion. Two iterative rounds of 3D classification and refinement with imposed

icosahedral symmetry restraints were applied to produce the final 3D maps of the E2p-L4P and

I3-01v9-L7P NP scaffolds. A soft solvent mask around the corresponding NP scaffold was

introduced for 3D classification, 3D refinement, and postprocessing steps to exclude signal

contributions from flexibly linked BG505 trimers and additional stabilizing domains (e.g., LD4

in E2p-L4P). For the E2p-L4P NP, the final subset had 7,672 particles and yielded a map

resolution of 3.7 Å. For the I3-01v9-L7P NP, the final subset consisted of 4,806 particle

projection images and was reconstructed to 6.0 Å resolution.

For the analysis of NP-attached BG505 UFO trimers, we used the localized

reconstruction method (*163*). Localized reconstruction v1.2.0 was applied to extract subparticles

corresponding to the trimer antigens from pre-aligned E2p-LD4 and I3-01v9 particle datasets.

Each NP presents 20 trimers on the surface. The starting trimer datasets consisted of 153,400 (20

× 7,672) and 96,120 (20 × 4,806) subparticles for BG505 UFO-E2p-L4P and UFO-10GS-I3-

01v9-L7P, respectively. Subparticle datasets were subjected to two rounds of 2D classification

and two rounds of 3D classification in Relion/3.0 (*162*) to eliminate cropped, overlapping, and

heterogeneously looking particles. For the 3D steps, we applied an HIV-1 Env trimer model from

negative stain EM, low-pass filtered to 40 Å resolution. The final subset consisted of 6,726 and

3,723 particles for E2p-L4P and I3-01v9-L7P-displayed BG505 UFO trimers. These subsets

were subjected to 3D refinement. In the case of E2p-L4P-displayed trimers, a soft solvent mask

around the trimer was used for the 3D refinement and postprocessing steps. C3 symmetry

restraints were imposed during refinement. Maps at 7.4 and 10.4 Å resolution were obtained for

E2p-L4P and I3-01v9-L7P-bound BG505 UFO trimers, respectively. The resulting maps after

3D refinement and postprocessing, and corresponding half-maps and solvent masks were

submitted to the Electron Microscopy Data Bank (EMDB). Model refinement was performed

only for the E2p-L4P NP scaffold, using the B-factor-sharpened map after postprocessing. The

structure of dihydrolipoyl transacetylase from PDB entry 1B5S (*104*) was used as a starting

model. Iterative rounds of manual model building in Coot (*164*) and Rosetta relaxed refinement

(*165*) were applied to generate the final structures. Model evaluation was performed using the

EMRinger (*166*) and MolProbity (*167*) packages, with statistics reported in **Table S2**. The model

was submitted to the Protein Data Bank (PDB).

Site-specific glycan analysis of BG505 trimer and SApNP immunogens

Three 50 µg aliquots of each sample were denatured for 1 h in 50 mM Tris/HCl, pH 8.0,

containing 6 M urea and 5 mM DTT. Next, Env proteins were reduced and alkylated by adding

20 mM iodoacetamide (IAA) and incubated for 1 h in the dark, followed by 1 h of incubation

with 20 mM DTT to eliminate residual IAA. The alkylated Env proteins were buffer-exchanged

into 50 mM Tris/HCl, pH 8.0 using Vivaspin columns (3 kDa) and digested separately overnight

using trypsin, chymotrypsin or elastase (Mass Spectrometry Grade, Promega) at a ratio of 1:30

(w/w). The next day, the peptides were dried and extracted using C18 Zip-tip (MerckMilipore).

The peptides were dried again, resuspended in 0.1% formic acid, and analyzed by nanoLC-ESI

MS with an Easy-nLC 1200 (Thermo Fisher Scientific) system coupled to a Fusion mass

spectrometer (Thermo Fisher Scientific) using higher energy collision-induced dissociation

(HCD) fragmentation. Peptides were separated using an EasySpray PepMap RSLC C18 column

(75 µm × 75 cm). A trapping column (PepMap 100 C18 3 µM 75 µM × 2cm) was used in line

with the LC prior to separation with the analytical column. The LC conditions were the

following: 275 min linear gradient consisting of 0-32% acetonitrile in 0.1% formic acid over 240

min followed by 35 min of 80% acetonitrile in 0.1% formic acid. The flow rate was set to 200

nl/min. The spray voltage was set to 2.7 kV and the temperature of the heated capillary was set to

40 °C. The ion transfer tube temperature was set to 275 °C. The scan range was 400-1600 *m/z.*

The HCD collision energy was set to 50%, appropriate for the fragmentation of glycopeptide

ions. Precursor and fragment detection were performed using an Orbitrap at a resolution MS1 =

100,000 MS2 = 30,000. The AGC target for MS1 = 4e5 and MS2 = 5e4 and injection time: MS1

= 50ms MS2 = 54ms.

Glycopeptide fragmentation data were extracted from the raw file using Byonic[TM]

(Version 3.5) and Byologic[TM] software (Version 3.5; Protein Metrics). The glycopeptide

fragmentation data were evaluated manually for each glycopeptide; the peptide was scored as

true-positive when the correct b and y fragment ions were observed along with oxonium ions

corresponding to the glycan identified. The MS data were searched using the Protein Metrics 305

N-glycan library. The relative amounts of each glycan at each site, as well as the unoccupied

proportion, were determined by comparing the extracted chromatographic areas for different

glycotypes with an identical peptide sequence. All charge states for a single glycopeptide were

summed. The precursor mass tolerance was set to 4 part per million (ppm) and 10 ppm for

fragments. A 1% false discovery rate (FDR) was applied. Glycans were categorized according to

the composition detected. HexNAc(2)Hex(9−5) was classified as M9 to M5,

HexNAc(3)Hex(5−6)X as Hybrid with HexNAc(3)Fuc(1)X classified as Fhybrid. Complex-type

glycans were classified according to the number of HexNAc residues, which are attributed to the

number of processed antenna/bisecting GlcNAc (B), and fucosylation (F). For example,

HexNAc(3)Hex(3-4)X was assigned to A1, HexNAc(4)X to A2/A1B, HexNAc(5)X to A3/A2B,

and HexNAc(6)X to A4/A3B. If all of these compositions had a fucose, then they were assigned

to the corresponding FA category. Note that this analytical approach does not distinguish

between isomers, which could influence formal assignment of the number of antennae in some

cases.

1005 Global glycan analysis of SApNP immunogens before and after endo H treatment

SDS-PAGE gel bands corresponding to BG505 UFO-10GS-I3-01v9-L7P SApNP were excised

and washed three times with alternating 1 ml acetonitrile and water, incubating and shaking for 5

minutes following addition of each wash solution. All liquid was removed following the final

wash stages and N-linked glycans were released in-gel using PNGaseF, (2 µg enzyme in 100 µL

H2O) (New England Biolabs) at 37 °C overnight. Following digestion, the liquid was removed

from the gel bands and placed into a separate Eppendorf. The gel bands were then washed twice

with 100 µl MilliQ H2O and this was pooled with the original solution. The extracted glycans

were then dried completely in a speed vac at 30 °C. The released glycans were subsequently

fluorescently labelled with procainamide using 110 mg/ml procainamide and 60 mg/ml sodium

cyanoborohydride in a buffer consisting of 70% DMSO, 30% acetic acid. For each sample, 100

µl of labelling mixture was added. Labelling was performed at 60 °C for 2 hours. Excess label

and PNGaseF were removed using Spe-ed Amide-2 cartridges (Applied Separations). First, the

cartridges were equilibrated sequentially with 1 ml acetonitrile, water and acetonitrile again.

Then 1 ml of 95% acetonitrile was added to the procainamide-released glycan mixture and

applied to the cartridge, allowing the cartridge to drain by gravity flow. After the mixture has

emptied the cartridge, two washes using 97% acetonitrile were performed. To elute the labelled

glycans 1 ml HPLC grade water was added to the cartridges and the elution collected. The

elution was then dried completely using a speed vac, before resuspending in 24 µl of 50 mM

ammonium formate. A 6 µl aliquot of the resuspended glycans were mixed with 24 µl of

acetonitrile and analysed on a Waters Acquity H-Class UPLC instrument with a Glycan BEH

Amide column (2.1 mm × 150 mm, 1.7 μM, Waters), with an injection volume of 10 μl. A

gradient of two buffers; 50 mM ammonium formate (buffer A) and acetonitrile (buffer B) was

used for optimal separation. Gradient conditions were as follows: initial conditions, 0.5 ml/min

22% buffer A, increasing buffer A concentration to 44.1% over 57.75 minutes. Following this

the concentration of buffer A was increase to 100% at 59.25 minutes and held there until 66.75

minutes and the flow rate was dropped to 0.25 ml/min, to fully elute from the column. Finally,

the %A was reduced to 20% to prepare for subsequent runs. Wavelengths used for detection of

the procainamide label were: excitation 310 nm, emission 370 nm. Data were processed using

Empower 3 software (Waters, Manchester, UK). The relative abundance of oligomannose-type

glycans was measured by digestion with Endoglycosidase H (per sample in 20 μl volume) (Endo

H; New England Biolabs). A 6 μl aliquot of labelled glycans was combined with 1 μg endoH to

a final volume of 20 μl. Digestion was performed for 1 hour at 37 °C. Digested glycans were

cleaned using a 96-well PVDF protein-binding membrane (Millipore) attached to a vacuum

manifold. Prior to application to the membrane, 100 μl HPLC-grade H2O was added to each

sample. Following equilibration with 150 μl ethanol, and 2 × 150 μl HPLC-grade H2O, the

sample was added to the 96-well plate and the flow-through was collected in a 96-well collection

plate. Each well was then washed twice with HPLC-grade H2O to a final elution volume of 300

μl. The elution was then dried completely at 30 °C. Prior to analysis the sample was resuspended

in 6 μl ammonium formate and 24 μl acetonitrile.

To generate a global glycan profile, a wildtype or glycan-trimmed I3-01v9 SApNP

sample was subjected to SDS-PAGE. The corresponding NP bands were processed by PNGaseF

to release glycans, which were labelled with procainamide. One aliquot of released glycans was

analyzed by ultra-performance liquid chromatography (UPLC), with the chromatogram

displayed in green. A second aliquot of released glycans was treated with endo H, which cleaved

the first GlcNAc and the procainamide label while depleting oligomannose-type glycans from

the sample. This sample was then analyzed by UPLC. The resulting chromatogram was

normalized to the untreated aliquot and displayed in pink when merged with the first

chromatogram to generate the profile.

Bio-layer interferometry (BLI)

The antigenic profiles of BG505 Env immunogens, the UFO trimers and SApNPs with wildtype

and modified glycan shields, were measured using an Octet RED96 instrument (FortéBio, Pall

Life Sciences) against a large panel of HIV-1 NAbs, bNAbs, and nNAbs in the IgG form. All

assays were performed with agitation set to 1000 rpm in FortéBio 1× kinetic buffer. The final

volume for all solutions was 200 μl per well. Assays were performed at 30 °C in solid black 96-

well plates (Geiger Bio-One). For UFO trimers, 5 μg/ml antibody in 1× kinetic buffer was loaded

onto the surface of anti-human Fc Capture Biosensors (AHC) for 300 s. For UFO trimer-

presenting SApNPs, anti-human Fc Quantitation Biosensors (AHQ) were used because they have

been shown to be more suitable for measuring the avidity effect for particulate immunogens (*27,*

89, 91, 92). Notably, the respective trimer was also measured using AHQ biosensors to facilitate

comparisons of antibody binding with trimer-presenting SApNPs. A 60 s biosensor baseline step

was applied prior to the analysis of the association of the antibody on the biosensor to the antigen

in solution for 200 s. A two-fold concentration gradient of antigen, starting at 266.7 nM for the

UFO trimer, 14.9 nM for the FR SApNP, and 5.5 nM for the E2p and I3-01v9 SApNPs, was

used in a titration series of six. The dissociation of the interaction was followed for 300 s. The

correction of baseline drift was performed by subtracting the mean value of shifts recorded for a

sensor loaded with antibody but not incubated with antigen, and for a sensor without antibody

but incubated with antigen. Octet data were processed by FortéBio's data acquisition software

v.8.1. Peak signals at the highest antigen concentration were summarized in a matrix and color-

coded accordingly to facilitate comparisons between different vaccine platforms and glycan

treatments. A separate Octet experiment was performed for the wildtype E2p SApNP in

duplicate at the highest concentration (5.5 nM) to determine the intra-experiment signal

variation, which is shown in fig. S4P.

Mouse immunization and sample collection

Similar immunization protocols have been used in our previous vaccine studies (*27, 89, 91, 92,*

123). Briefly, Institutional Animal Care and Use Committee (IACUC) guidelines were followed

with animal subjects tested in the immunization study. Eight-week-old BALB/c mice were

purchased from The Jackson Laboratory and housed in ventilated cages in environmentally

controlled rooms at The Scripps Research Institute, in compliance with an approved IACUC

protocol and Association for Assessment and Accreditation of Laboratory Animal Care

(AAALAC) international guidelines. Mice were immunized at weeks 0, 3, 6, and 9 with 200 μl

of antigen/adjuvant mix containing 30 μg of vaccine antigen and 100 μl of adjuvant, AddaVax,

Adju-Phos, or Alhydrogel (InvivoGen), via the intraperitoneal (i.p.) route. Blood was collected 2

weeks after each injection. All bleeds were performed through the retro-orbital sinus using

heparinized capillary tubes into EDTA-coated tubes. Samples were spun at 1200 rpm for 10 min

to separate plasma (top layer) and the rest of the whole blood layer. Upon heat inactivation at 56

°C for 30 min, plasma was spun at 2000 rpm for 10 min to remove precipitates. The rest of the

whole blood layer was diluted with an equal volume of PBS and then overlaid on 4.5 ml of Ficoll

in a 15 ml SepMate^{TM} tube (STEMCELL Technologies) and spun at 1200 rpm for 10 min at 20

°C to separate peripheral blood mononuclear cells (PBMCs). Cells were washed once in PBS and

then resuspended in 1 ml of ACK Red Blood Cell lysis buffer (Lonza). After washing with PBS,

PBMCs were resuspended in 2 ml of Bambanker Freezing Media (Lymphotec). Spleens were

harvested at week 11 and ground against a 70-μm cell strainer (BD Falcon) to release

splenocytes into a cell suspension. Splenocytes were centrifuged, washed in PBS, treated with 5

ml of ACK lysing buffer (Lonza), and frozen with 3 ml of Bambanker freezing media. Total

plasma from week 11 was purified using CaptureSelect™ IgG-Fc (Multispecies) Affinity Matrix

(Thermo Scientific) following the manufacturer's instructions. Purified IgG samples from

individual mice were analyzed in TZM-bl assays to determine the vaccine-induced neutralizing

response.

Rabbit immunization and sample collection

The Institutional Animal Care and Use Committee (IACUC) guidelines were followed for the

animal subjects tested in the immunization studies. For the evaluation of wildtype Env and Env-

NP immunogens using a long regimen, rabbit immunization and blood sampling were performed

under a subcontract at Covance (Denver, PA). Seven groups of female New Zealand White

rabbits, six rabbits per group, were immunized intramuscularly (i.m.) with 100 μg of HIV-1 Env

antigen formulated in 250 μl of adjuvant, AddaVax or Adju-Phos (InvivoGen), with a total

volume of 500 μl, at weeks 0, 8, and 24. Blood samples (15 ml each) were collected at day 0 and

weeks 1, 2, 4, 6, 8, 10, 12, 16, 20, 24, 26, and 28. Plasma was separated from blood and heat

inactivated for the TZM-bl neutralization assays. For the assessment of glycan-trimmed Env-NP

immunogens using a short regimen, rabbit immunization and blood sampling were performed

under a subcontract at ProSci (San Diego, CA). Four groups of female New Zealand White

rabbits, six rabbits per group, were intramuscularly (i.m.) immunized with 30 μg of vaccine

antigen formulated in 250 μl of adjuvant, AddaVax or Adju-Phos (InvivoGen), with a total

volume of 500 µl, at weeks 0, 3, 6, 9, and 12. Blood samples, 20 ml each time, were collected

from the auricular artery at day 0 (Pre) and weeks 2, 5, 8, 11, 14, and 16. More than 100 ml of

blood was taken at week 16, via cardiac puncture, for PBMC isolation. Plasma samples were

heat inactivated for further characterization, and purified rabbit IgGs were assessed in TZM-bl

neutralization assays.

Nonhuman primate immunization and sample collection

Research-naive adult rhesus macaques of Indian origin were genotyped and selected as negative

for the protective major histocompatibility complex (MHC) class I alleles Mamu-A*01, Mamu-

B*08 and Mamu-B*17. Rhesus macaques were sourced from the Southwest National Primate

Research Center (SNPRC) and Tulane National Primate Research Center (TNPRC). Animals

from TNPRC were quarantined at SNPRC for 42 days during which a physical exam, ova and

parasite evaluation, and three TB skin tests were performed. All experimental procedures were

performed at SNPRC in San Antonio, TX, USA according to the guidelines of the AAALAC

standards. These macaque experiments were carried out in compliance with all pertinent US

National Institutes of Health (NIH) regulations and were approved by the IACUC of SNPRC.

Rhesus macaques were moved into research housing and given a physical exam. The animals

were randomly assigned to the study groups, which were balanced for age, weight, and gender.

Animals were allowed to acclimate to research housing for 14 days before initiation of the study

protocol. On the immunization time points, animals received a 250 µl subcutaneous or

intramuscular injections of vaccine into each quadracep. Injection sites were visually observed at

24 and 48 hours post injection. Blood collections were made from the femoral vein under

anesthesia. In the study of wildtype SApNPs, animals were immunized at weeks 0, 8, and 24

with 200 µg antigen formulated with AddaVax (AV, for E2p) or aluminum phosphate (AP, for

I3-01v9) and blood was collected at weeks -2, 1, 2, 4, 8, 9, 10, 12, 16, 24, 25, 26, and 28. In the

study of glycan-trimmed SApNPs, animals were immunized at weeks 0, 4, 12, and 24 with 100

µg antigen formulated with AP (for both E2p and I3-01v9) and blood was collected at weeks 0,

1, 4, 5, 8, 13, 16, 25, and 28. Animals were returned to the colony 28 weeks after the initial

vaccination.

Pseudovirus production and neutralization assays

Pseudoviruses were generated by the transfection of HEK293T cells with an HIV-1 Env

expressing plasmid and an Env-deficient genomic backbone plasmid (pSG3ΔEnv), as previously

described (*168*). HIV-1 Env expressing vectors for BG505 (catalog no. 11518), SF162 (catalog

no. 10463), and the 12-virus global panel (*169*) (catalog no. 12670) were obtained through the

NIH AIDS Reagent Program (https://www.aidsreagent.org/). A T332N mutation was introduced

into BG505 Env to produce the BG505.T332N clone. Other BG505.T332N mutants were created

by introducing mutations as previously described (*56, 60, 69*). Pseudoviruses were harvested 72

h post-transfection for use in the neutralization assays. The neutralizing activity of each mouse,

rabbit or NHP sample was assessed using a single round of replication pseudovirus assay and

TZM-bl target cells, similar to that described previously (*168*). Briefly, pseudovirus was

incubated with serial dilutions of antibodies or heat inactivated rabbit plasma/NHP sera in a 96-

well flat bottom plate for 1 h at 37 °C before TZM-bl cells were seeded in the plate. For purified

mouse and rabbit IgGs, a starting concentration of 300 µg/ml was used, or an initial dilution of

either 40-times or 100-times for rabbit plasma or NHP sera, and then subjected to 3-times serial

dilutions in the TZM-bl assays. Samples were tested in duplicate with a full series of dilutions

unless otherwise specified in the Results, with such exceptions including tests against the 12-

virus global panel performed in singlet to conserve purified IgG samples. Luciferase reporter

gene expression was quantified 60-72 h after infection upon lysis and the addition of Bright-

Glo^{TM} Luciferase substrate (Promega). Data were retrieved from a BioTek microplate reader

with Gen 5 software. Values from experimental wells were compared against a well containing

virus only, with background luminescence from a series of uninfected wells subtracted from

both. Dose-response neutralization curves were then fit by nonlinear regression in GraphPad

Prism 9.3.1, with IC_{30}, IC_{50}, ID_{30} and ID_{50} values calculated with constraints set between 0 to

100. As a negative control, pseudoparticles displaying the envelope glycoproteins of MLV

(MLV-pps) were tested in the TZM-bl assays following the same protocol.

Histology, immunostaining, and imaging

For the mechanistic study of HIV-1 Env vaccine distribution, trafficking, cellular interaction, and

induced GCs in lymph nodes, vaccines were intradermally injected into mouse footpads using a

29-gauge insulin needle under 3% isoflurane anesthesia with oxygen. Similar immunization and

tissue analysis protocols were used in our previous study (*123*). The administered dose was 80 μl

of antigen/adjuvant mix containing 40 μg of vaccine antigen per mouse or 10 μg per footpad.

Mice were euthanized 30 min to 8 weeks after a single-dose injection. The axillary and popliteal

sentinel lymph nodes were collected for histological analysis. Fresh lymph nodes were merged

into frozen section compound (VWR International, catalog no. 95057-838) in a plastic cryomold

(Tissue-Tek at VWR, catalog no. 4565). Sample molds were merged into liquid nitrogen and

then stored at -80°C before shipping to The Centre for Phenogenomics in Canada for sample

processing, immunostaining, and imaging. Tissue sections were sliced 8 μm thick on a cryostat

(Cryostar NX70) and collected on charged slides. Samples were then fixed in 10% neutral

buffered formalin and further permeabilized in PBS that contained 0.5% Triton X-100 before

staining. The slides were blocked with protein Block (Agilent) to prevent nonspecific antibody

binding. Primary antibody was then applied on the sections and incubated overnight at 4°C.

After washing with TBST, secondary antibodies that were conjugated with either biotin or a

fluorophore were used, and the samples were incubated for 1 h at 25 °C. Lymph node sections

were stained with bNAbs VRC01 (*115*), PGT124 (*133*), and PGDM1400 (*134*) (1:50), and

biotinylated goat anti-human secondary antibody (Abcam, catalog no. ab7152, 1:300), followed

by streptavidin-horseradish peroxidase (HRP) reagent (Vectastain Elite ABC-HRP Kit, Vector,

catalog no. PK-6100) and diaminobenzidine (DAB) (ImmPACT DAB, Vector, catalog no. SK-

4105).

To visualize interactions between HIV-1 Env vaccines and cells in mouse lymph nodes,

FDCs were labeled using anti-CD21 primary antibody (Abcam, catalog no. ab75985, 1:1800),

followed by anti-rabbit secondary antibody conjugated with Alexa Fluor 555 (Thermo Fisher,

catalog no. A21428, 1:200). B cells were labeled using anti-B220 antibody (eBioscience, catalog

no. 14-0452-82, 1:100) followed by anti-rat secondary antibody conjugated with Alexa Fluor 647

(Thermo Fisher, catalog no. A21247, 1:200). Subcapsular sinus macrophages were labeled using

anti-sialoadhesin (CD169) antibody (Abcam, catalog no. ab53443, 1:600) followed by anti-rat

secondary antibody conjugated with Alexa Fluor 488 (Abcam, catalog no. ab150165, 1:200).

HIV-1 vaccine-induced GCs were studied using immunostaining, GC B cells stained using rat

anti-GL7 antibody (FITC; BioLegend, catalog no. 144604, 1:250), T_{fh} cells stained using anti-

CD4 antibody (BioLegend, catalog no. 100402, 1:100) followed by anti-rat secondary antibody

conjugated with Alexa Fluor 488 (Abcam, catalog no. ab150165, 1:1000), GC cells stained using

Bcl6 antibody (Abcam, catalog no. ab220092, 1:300) followed by anti-rabbit secondary antibody

conjugated with Alexa Fluor 555 (Thermo Fisher, catalog no. A21428, 1:1000). Nuclei were

labeled using 4′,6-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich, catalog no. D9542, 100

ng/ml).

The stained lymph node sections were scanned using an Olympus VS-120 slide scanner

with a Hamamatsu ORCA-R2 C10600 digital camera for bright-field and fluorescent images.

The transport of HIV-1 BG505 Env immunogens and their induced GCs in mouse lymph nodes

were quantified through bright-field and fluorescent images using ImageJ software (*170*).

Electron microscopy analysis of protein nanoparticles and lymph node tissues

TEM analysis was conducted by the Core Microscopy Facility at The Scripps Research Institute.

To visualize interactions of BG505 UFO Env-SApNPs with FDCs and phagocytic cells in lymph

nodes using TEM, mice were injected with 140 μl of antigen/adjuvant mix containing 100 μg of

vaccine antigen (40 μl of adjuvant) into the two hind footpads or 50 μg per footpad. Popliteal

sentinel lymph nodes were isolated 2, 12, and 48 hours after administration, bisected and

immersed in oxygenated 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1 M sodium

cacodylate buffer (pH 7.4) fixative overnight at 4ºC. The lymph node tissues were washed using

0.1 M sodium cacodylate buffer and post-fixed in buffered 1% osmium tetroxide and 1.5%

potassium ferrocyanide for 1-1.25 hours at 4°C. The samples were rinsed in Corning Cell

Culture Grade Water and stained *en bloc* with 0.5% uranyl acetate overnight at 4°C. The tissues

were rinsed with double-distilled H_2O and dehydrated through a graded series of ethanol

followed by acetone, infiltrated with LX-112 (Ladd) epoxy resin, and polymerized at 60°C.

Ultrathin tissue sections (70 nm) were prepared on copper grids for TEM imaging. Tissue

samples were imaged at 80 kV with a Talos L120C transmission electron microscope (Thermo

Fisher), and images were acquired with a CETA 16M CMOS camera.

Lymph node disaggregation, cell staining, and flow cytometry

GC B cells (GL7$^+$B220$^+$) and T$_{fh}$ cells (CD3$^+$CD4$^+$CXCR5$^+$PD-1$^+$) were characterized using flow cytometry (**Fig. S10A**). Mice were euthanized 2, 5, and 8 weeks after a single-dose injection and 2 and 5 weeks after the boost injection (4×10 μg/footpad each time). Fresh axillary, brachial, and popliteal sentinel lymph nodes were isolated. After mechanically disaggregating the lymph node tissues, samples were merged in enzyme digestion solution in an Eppendorf tube containing 958 μl of Hanks' balanced salt solution (HBSS) buffer (Thermo Fisher Scientific, catalog no. 14185052), 40 μl of 10 mg/ml collagenase IV (Sigma-Aldrich, catalog no. C5138), and 2 μl of 10 mg/ml DNase (Roche, catalog no. 10104159001). Lymph node tissues were incubated at 37°C for 30 min and filtered through a 70 μm cell strainer. Samples were spun down at $400 \times g$ for 10 min to isolate cell pellets, which were resuspended in HBSS blocking buffer with 0.5% (w/v) bovine serum albumin and 2 mM EDTA. Anti-CD16/32 antibody (BioLegend, catalog no. 101302) was added to block the nonspecific binding of Fc receptors, while the sample solution was kept on ice for 30 min. Samples were then transferred to 96-well microplates with pre-prepared cocktail antibodies, which included the Zombie NIR live/dead stain (BioLegend, catalog no. 423106), Brilliant Violet 510 anti-mouse/human CD45R/B220 antibody (BioLegend, catalog no. 103247), FITC anti-mouse CD3 antibody (BioLegend, catalog no. 100204), Alexa Fluor 700 anti-mouse CD4 antibody (BioLegend, catalog no. 100536), PE anti-mouse/human GL7 antibody (BioLegend, catalog no. 144608), Brilliant Violet 605 anti-mouse CD95 (Fas) antibody (BioLegend, catalog no. 152612), Brilliant Violet 421 anti-mouse CD185 (CXCR5) antibody (BioLegend, catalog no. 145511), and PE/Cyanine7 anti-mouse CD279 (PD-1) antibody (BioLegend, catalog no. 135216). The cell samples mixed with antibody cocktail were placed on ice for 30 min and centrifuged to remove excess antibody. After washing with the

HBSS blocking solution, cells were fixed with 1.6% paraformaldehyde (Thermo Fisher

Scientific, catalog no. 28906) in HBSS on ice for 30 min. The samples were then placed in

HBSS blocking solution at 4°C. Sample events were acquired by a 5-laser AZE5 flow cytometer

(Yeti, Bio-Rad) with Everest software at the Core Facility of The Scripps Research Institute. The

data were analyzed using FlowJo 10 software.

Statistical analysis

Data were collected from 8 mice per group in the immunization study and 3-10 mice per group

in the mechanistic analysis. All of the statistical analyses were performed and graphs were

generated using GraphPad Prism 9.3.1 software. For the vaccine transport and GC study in

lymph nodes, different vaccine groups were compared using one-way ANOVA, followed by

Tukey's multiple comparison *post hoc* test. Statistical significance is indicated as the following

in the figures: ns (not significant), $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. To

assess the highest level of improvement in FVR, one comparison was made for each vaccine

platform (trimer or SApNP) between the wildtype group (or a representative wildtype group,

when multiple existed) and the glycan-trimmed group that had the highest FVR (or the highest

average, when multiple repeats existed). This statistical comparison was performed using a two-

tailed Fisher exact test.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at XXX.

Tables S1 and S2.

Figures S1-S9.

ACKNOWLEDGEMENTS

Y.-N.Z. acknowledges support from the Natural Sciences and Engineering Research Council of

Canada (NSERC) for the postdoctoral fellowship. We thank M. Ganguly, K. Duffin, and G.

Ossetchkine at The Centre for Phenogenomics for their technical support in immunohistology.

We thank K. Vanderpool, T. Fassel, and S. Henderson of the Core Microscopy Facility at The

Scripps Research Institute for their expert assistance in the TEM analysis. We thank A. Saluk, B.

Seegers, and B. Monteverde of the Flow Cytometry Core Facility at The Scripps Research

Institute for their technical support in flow cytometry. We thank P. Berman and D. Burton for

helpful discussions and M. Arends and V. Tong for proofreading the manuscript. **Funding:**

This work was supported by the International AIDS Vaccine Initiative (IAVI) through grant

INV-008352/OPP1153692 (M.C.) and the IAVI Neutralizing Antibody Center through the

Collaboration for AIDS Vaccine Discovery grant (INV-034657) (I.A.W., A.B.W.), and INV-

008352/OPP1153692 (M.C.) funded by the Bill and Melinda Gates Foundation; Scripps

Consortium for HIV/AIDS Vaccine Development (CHAVD 1UM1 AI144462) (M.C., A.B.W.

and I.A.W.); HIV Vaccine Research and Design (HIVRAD) program (P01 AI124337) (J.Z.);

NIH grants R01 AI129698 (J.Z.) and R01 AI140844 (J.Z., I.A.W.); Southwest National Primate

Research Center through grants P51 OD011133 and U42OD010442 and Tulane National

Primate Research Center through grant P51 OD011104 from the Office of Research

Infrastructure Programs, NIH (XXX); Ufovax/SFP-2018-0416 (J.Z.) and Ufovax/SFP-2018-1013

(J.Z.). **Author contributions:** Project design by Y.-N.Z., J.P., L.H., I.A.W., and J.Z.;

immunogen expression and purification by M.E. and L.H.; negative stain EM and cryo-EM by

A.A., Y.-Z.L., J.C., and A.B.W.; glycan analysis by J.D.A., M.N., and M.C.; mouse and rabbit

sample preparation, and plasma and IgG neutralization by J.P. and L.H.; NHP immunization and

sample collection by D.C., P.F., A.G., J.D., R.L., and C.C.; NHP serum neutralization and data

analysis by M.E. and L.H.; mouse immunization, lymph node isolation, immunohistology, TEM,

and flow cytometry by Y.-N.Z.; manuscript written by Y.-N.Z., A.A., J.P., J.D.A., I.A.W., M.

C., A.B.W. and J.Z. All authors were asked to comment on the manuscript. The Scripps

Research Institute manuscript number is 30193. **Competing interests:** The authors declare that

they have no competing interests. **Data and material availability:** All data and codes to

understand and assess the conclusions of this research are available in the main text,

Supplementary Materials, PDB (accession code XXX) and EMDB (accession code EMD-YYY).

Additional data related to this paper may be requested from the authors.

References

1308
1.   R. Wyatt, J. Sodroski, The HIV-1 envelope glycoproteins: fusogens, antigens, and
immunogens. *Science* **280**, 1884-1888 (1998).
2.   R. P. Smyth, M. P. Davenport, J. Mak, The origin of genetic diversity in HIV-1. *Virus*
Res. **169**, 415-429 (2012).
3.   P. L. Moore *et al.*, Nature of nonfunctional envelope proteins on the surface of human
immunodeficiency virus type 1. *J. Virol.* **80**, 2515-2528 (2006).
4.   X. P. Wei *et al.*, Antibody neutralization and escape by HIV-1. *Nature* **422**, 307-312
(2003).
5.   D. R. Burton, J. R. Mascola, Antibody responses to envelope glycoproteins in HIV-1
infection. *Nat. Immunol.* **16**, 571-576 (2015).
6.   D. Sok, D. R. Burton, Recent progress in broadly neutralizing antibodies to HIV. *Nat.*
Immunol. **19**, 1179-1188 (2018).
7.   L. E. McCoy, D. R. Burton, Identification and specificity of broadly neutralizing
antibodies against HIV. *Immunol. Rev.* **275**, 11-20 (2017).
8.   D. R. Burton *et al.*, A blueprint for HIV vaccine discovery. *Cell Host Microbe* **12**, 396-
407 (2012).
9.   L. M. Walker, D. R. Burton, Rational antibody-based HIV-1 vaccine design: current
approaches and future directions. *Curr. Opin. Immunol.* **22**, 358-366 (2010).
10.   B. F. Haynes, G. Kelsoe, S. C. Harrison, T. B. Kepler, B-cell-lineage immunogen design
in vaccine development with HIV-1 as a case study. *Nat. Biotechnol.* **30**, 423-433 (2012).
11.   F. Klein *et al.*, Antibodies in HIV-1 vaccine development and therapy. *Science* **341**,
1199-1204 (2013).
12.   R. W. Sanders, J. P. Moore, Native-like Env trimers as a platform for HIV-1 vaccine
design. *Immunol. Rev.* **275**, 161-182 (2017).
13.   A. B. Ward, I. A. Wilson, The HIV-1 envelope glycoprotein structure: nailing down a
moving target. *Immunol. Rev.* **275**, 21-32 (2017).
14.   D. Sok *et al.*, Priming HIV-1 broadly neutralizing antibody precursors in human Ig loci
transgenic mice. *Science* **353**, 1557-1560 (2016).

15. J. G. Jardine *et al.*, Priming a broadly neutralizing antibody response to HIV-1 using a
germline-targeting immunogen. *Science* **349**, 156-161 (2015).
16. J. Jardine *et al.*, Rational HIV immunogen design to target specific germline B cell
receptors. *Science* **340**, 711-716 (2013).
17. L. Scharf *et al.*, Structural basis for germline antibody recognition of HIV-1
immunogens. *eLife* **5**, e13783 (2016).
18. A. J. Borst *et al.*, Germline VRC01 antibody recognition of a modified clade C HIV-1
envelope trimer and a glycosylated HIV-1 gp120 core. *eLife* **7**, e37688 (2018).
19. R. W. Sanders *et al.*, A next-generation cleaved, soluble HIV-1 Env trimer, BG505
SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not non-
neutralizing antibodies. *PLoS Pathog.* **9**, e1003618 (2013).
20. P. Pugach *et al.*, A native-like SOSIP.664 trimer based on an HIV-1 subtype B Env gene.
J. Virol. **89**, 3380-3395 (2015).
21. J.-P. Julien *et al.*, Design and structure of two HIV-1 clade C SOSIP.664 trimers that
increase the arsenal of native-like Env immunogens. *Proc. Natl. Acad. Sci. U. S. A.* **112**,
11947-11952 (2015).
22. K. Sliepen *et al.*, Structure and immunogenicity of a stabilized HIV-1 envelope trimer
based on a group-M consensus sequence. *Nat. Commun.* **10**, 2355 (2019).
23. S. K. Sharma *et al.*, Cleavage-independent HIV-1 Env trimers engineered as soluble
native spike mimetics for vaccine design. *Cell Rep.* **11**, 539-550 (2015).
24. J. Guenaga *et al.*, Glycine substitution at helix-to-coil transitions facilitates the structural
determination of a stabilized subtype C HIV envelope glycoprotein. *Immunity* **46**, 792-
803 (2017).
25. J. Guenaga *et al.*, Structure-guided redesign increases the propensity of HIV Env to
generate highly stable soluble trimers. *J. Virol.* **90**, 2806-2817 (2015).
26. L. Kong *et al.*, Uncleaved prefusion-optimized gp140 trimers derived from analysis of
HIV-1 envelope metastability. *Nat. Commun.* **7**, 12040 (2016).
27. L. He *et al.*, HIV-1 vaccine design through minimizing envelope metastability. *Sci. Adv.*
4, eaau6769 (2018).
28. Q. Wang *et al.*, Stabilized diverse HIV-1 envelope trimers for vaccine design. *Emerg.*
Microbes & Infect. **9**, 775-786 (2020).
29. R. Derking, R. W. Sanders, Structure-guided envelope trimer design in HIV-1 vaccine
development: a narrative review. *J. Int. AIDS Soc.* **24**, e25797 (2021).
30. I. del Moral-Sanchez *et al.*, High thermostability improves neutralizing antibody
responses induced by native-like HIV-1 envelope trimers. *Npj Vaccines* **7**, 27 (2022).
31. M. Crispin, A. B. Ward, I. A. Wilson, Structure and immune recognition of the HIV
glycan shield. *Annu. Rev. Biophys.* **47**, 499-523 (2018).
32. K. J. Doores, The HIV glycan shield as a target for broadly neutralizing antibodies. *FEBS*
J. **282**, 4679-4691 (2015).
33. K. Wagh, B. H. Hahn, B. Korber, Hitting the sweet spot: exploiting HIV-1 glycan shield
for induction of broadly neutralizing antibodies. *Curr. Opin. HIV AIDS* **15**, 267-274
(2020).
34. V. Hariharan, R. S. Kane, Glycosylation as a tool for rational vaccine design. *Biotechnol.*
Bioeng. **117**, 2556-2570 (2020).
35. C. N. Scanlan, J. Offer, N. Zitzmann, R. A. Dwek, Exploiting the defensive sugars of
HIV-1 for drug and vaccine design. *Nature* **446**, 1038-1045 (2007).

36.   G. E. Seabright, K. J. Doores, D. R. Burton, M. Crispin, Protein and glycan mimicry in
HIV vaccine design. *J. Mol. Biol.* **431**, 2223-2247 (2019).
37.   A. J. Behrens *et al.*, Molecular architecture of the cleavage-dependent mannose patch on
a soluble HIV-1 envelope glycoprotein trimer. *J. Virol.* **91**, e01894-01816 (2017).
38.   A. J. Behrens *et al.*, Composition and antigenic effects of individual glycan sites of a
trimeric HIV-1 envelope glycoprotein. *Cell Rep.* **14**, 2695-2706 (2016).
39.   L. W. Cao *et al.*, Global site-specific N-glycosylation analysis of HIV envelope
glycoprotein. *Nat. Commun.* **8**, 13 (2017).
40.   F. Garces *et al.*, Affinity maturation of a potent family of HIV antibodies is primarily
focused on accommodating or avoiding glycans. *Immunity* **43**, 1053-1063 (2015).
41.   M. Pancera *et al.*, Structural basis for diverse N-glycan recognition by HIV-1-
neutralizing V1-V2-directed antibody PG16. *Nat. Struct. Mol. Biol.* **20**, 804-813 (2013).
42.   J. S. McLellan *et al.*, Structure of HIV-1 gp120 V1/V2 domain with broadly neutralizing
antibody PG9. *Nature* **480**, 336-343 (2011).
43.   H. B. Gristick *et al.*, Natively glycosylated HIV-1 Env structure reveals new mode for
antibody recognition of the CD4-binding site. *Nat. Struct. Mol. Biol.* **23**, 906-915 (2016).
44.   R. Pejchal *et al.*, A potent and broad neutralizing antibody recognizes and penetrates the
HIV glycan shield. *Science* **334**, 1097-1103 (2011).
45.   E. M. Cale *et al.*, Virus-like particles identify an HIV V1V2 apex-binding neutralizing
antibody that lacks a protruding loop. *Immunity* **46**, 777-791.e710 (2017).
46.   M. Pancera *et al.*, Structure and immune recognition of trimeric pre-fusion HIV-1 Env.
Nature **514**, 455-461 (2014).
47.   J. H. Lee, G. Ozorowski, A. B. Ward, Cryo-EM structure of a native, fully glycosylated,
cleaved HIV-1 envelope trimer. *Science* **351**, 1043-1048 (2016).
48.   J. H. Lee, N. de Val, D. Lyumkis, A. B. Ward, Model building and refinement of a
natively glycosylated HIV-1 Env protein by high-resolution cryoelectron microscopy.
Structure **23**, 1943-1951 (2015).
49.   D. Lyumkis *et al.*, Cryo-EM structure of a fully glycosylated soluble cleaved HIV-1
envelope trimer. *Science* **342**, 1484-1490 (2013).
50.   R. Derking *et al.*, Enhancing glycan occupancy of soluble HIV-1 envelope trimers to
mimic the native viral spike. *Cell Rep.* **35**, 108933 (2021).
51.   R. Andrabi, J. N. Bhiman, D. R. Burton, Strategies for a multi-stage neutralizing
antibody-based HIV vaccine. *Curr. Opin. Immunol.* **53**, 143-151 (2018).
52.   V. Dubrovskaya *et al.*, Vaccination with glycan-modified HIV NFL envelope trimer-
liposomes elicits broadly neutralizing antibodies to multiple sites of vulnerability.
Immunity **51**, 915-929.e917 (2019).
53.   V. Dubrovskaya *et al.*, Targeted N-glycan deletion at the receptor-binding site retains
HIV Env NFL trimer integrity and accelerates the elicited antibody response. *PLoS*
Pathog. **13**, e1006614 (2017).
54.   B. Briney *et al.*, Tailored immunogens direct affinity maturation toward HIV neutralizing
antibodies. *Cell* **166**, 1459-1470.e1411 (2016).
55.   J. K. Hu *et al.*, Murine antibody responses to cleaved soluble HIV-1 envelope trimers are
highly restricted in specificity. *J. Virol.* **89**, 10383-10398 (2015).
56.   L. E. McCoy *et al.*, Holes in the glycan shield of the native HIV envelope are a target of
trimer-elicited neutralizing antibodies. *Cell Rep.* **16**, 2327-2338 (2016).

57.   E. T. Crooks *et al.*, Vaccine-elicited tier 2 HIV-1 neutralizing antibodies bind to
quaternary epitopes involving glycan-deficient patches proximal to the CD4 binding site.
PLoS Pathog. **11**, e1004932 (2015).
58.   J. E. Voss *et al.*, Elicitation of neutralizing antibodies targeting the V2 apex of the HIV
envelope trimer in a wild-type animal model. *Cell Rep.* **21**, 222-235 (2017).
59.   R. P. Ringe *et al.*, Closing and opening holes in the glycan shield of HIV-1 envelope
glycoprotein SOSIP trimers can redirect the neutralizing antibody response to the newly
unmasked epitopes. *J. Virol.* **93**, e01656-01618 (2019).
60.   P. J. Klasse *et al.*, Epitopes for neutralizing antibodies induced by HIV-1 envelope
glycoprotein BG505 SOSIP trimers in rabbits and macaques. *PLoS Pathog.* **14**, e1006913
(2018).
61.   P. J. Klasse *et al.*, Sequential and simultaneous immunization of rabbits with HIV-1
envelope glycoprotein SOSIP.664 trimers from clades A, B and C. *PLoS Pathog.* **12**,
e1005864 (2016).
62.   S. Bale *et al.*, Cleavage-independent HIV-1 trimers from CHO cell lines elicit robust
autologous tier 2 neutralizing antibodies. *Front. Immunol.* **9**, 1116 (2018).
63.   M. Bianchi *et al.*, Electron-microscopy-based epitope mapping defines specificities of
polyclonal antibodies elicited during HIV-1 BG505 envelope trimer immunization.
Immunity **49**, 288-300.e288 (2018).
64.   Y. H. R. Yang *et al.*, Autologous antibody responses to an HIV envelope glycan hole are
not easily broadened in rabbits. *J. Virol.* **94**, e01861-01819 (2020).
65.   A. Schorcht *et al.*, The glycan hole area of HIV-1 envelope trimers contributes
prominently to the induction of autologous neutralization. *J. Virol.* **96**, e0155221 (2022).
66.   A. Antanasijevic *et al.*, Polyclonal antibody responses to HIV Env immunogens resolved
using cryoEM. *Nat. Commun.* **12**, 4817 (2021).
67.   F. Z. Zhao *et al.*, Mapping neutralizing antibody epitope specificities to an HIV Env
trimer in immunized and in infected rhesus macaques. *Cell Rep.* **32**, 108122 (2020).
68.   H. L. Turner *et al.*, Disassembly of HIV envelope glycoprotein trimer immunogens is
driven by antibodies elicited via immunization. *Sci. Adv.* **7**, eabh2791 (2021).
69.   L. Lei *et al.*, The HIV-1 envelope glycoprotein C3/V4 region defines a prevalent
neutralization epitope following immunization. *Cell Rep.* **27**, 586-598.e586 (2019).
70.   Z. Y. Xu *et al.*, Induction of tier-2 neutralizing antibodies in mice with a DNA-encoded
HIV envelope native like trimer. *Nat. Commun.* **13**, 695 (2022).
71.   S. Kumar *et al.*, Neutralizing antibodies induced by first-generation gp41-stabilized HIV-
1 envelope trimers and nanoparticles. *mBio* **12**, e00429-00421 (2021).
72.   T. P. Charles *et al.*, The C3/465 glycan hole cluster in BG505 HIV-1 envelope is the
major neutralizing target involved in preventing mucosal SHIV infection. *PLoS Pathog.*
17, e1009257 (2021).
73.   D. W. Kulp *et al.*, Structure-based design of native-like HIV-1 envelope trimers to
silence non-neutralizing epitopes and eliminate CD4 binding. *Nat. Commun.* **8**, 1655
(2017).
74.   P. J. M. Brouwer, R. W. Sanders, Presentation of HIV-1 envelope glycoprotein trimers on
diverse nanoparticle platforms. *Curr. Opin. HIV AIDS* **14**, 302-308 (2019).
75.   B. Asbach, R. Wagner, Particle-based delivery of the HIV envelope protein. *Curr. Opin.*
HIV AIDS **12**, 265-271 (2017).

76.   M. Brinkkemper, K. Sliepen, Nanoparticle vaccines for inducing HIV-1 neutralizing
1474   antibodies. *Vaccines* **7**, 76 (2019).
1475   77.   S. Thalhauser, D. Peterhoff, R. Wagner, M. Breunig, Critical design criteria for
1476   engineering a nanoparticulate HIV-1 vaccine. *J. Control. Release* **317**, 322-335 (2020).
1477   78.   Z. Y. Xu, D. W. Kulp, Protein engineering and particulate display of B-cell epitopes to
1478   facilitate development of novel vaccines. *Curr. Opin. Immunol.* **59**, 49-56 (2019).
1479   79.   D. J. Irvine, B. J. Read, Shaping humoral immunity to vaccines through antigen-
1480   displaying nanoparticles. *Curr. Opin. Immunol.* **65**, 1-6 (2020).
1481   80.   B. Nguyen, N. H. Tolia, Protein-based antigen presentation platforms for nanoparticle
1482   vaccines. *Npj Vaccines* **6**, (2021).
1483   81.   Q. Zhao, S. Li, H. Yu, N. Xia, Y. Modis, Virus-like particle-based human vaccines:
1484   quality assessment based on structural and functional properties. *Trends Biotechnol.* **31**,
1485   654-663 (2013).
1486   82.   W. A. Rodriguez-Limas, K. Sekar, K. E. J. Tyo, Virus-like particles: the future of
1487   microbial factories and cell-free systems as platforms for vaccine development. *Curr.*
1488   *Opin. Biotechnol.* **24**, 1089-1093 (2013).
1489   83.   G. T. Jennings, M. F. Bachmann, Coming of age of virus-like particle vaccines. *Biol.*
1490   *Chem.* **389**, 521-536 (2008).
1491   84.   C. Ludwig, R. Wagner, Virus-like particles - universal molecular toolboxes. *Curr. Opin.*
1492   *Biotechnol.* **18**, 537-545 (2007).
1493   85.   E. V. L. Grgacic, D. A. Anderson, Virus-like particles: Passport to immune recognition.
1494   *Methods* **40**, 60-65 (2006).
1495   86.   B. Zakeri *et al.*, Peptide tag forming a rapid covalent bond to a protein, through
1496   engineering a bacterial adhesin. *Proc. Natl. Acad. Sci. U. S. A.* **109**, E690-E697 (2012).
1497   87.   J. B. Bale *et al.*, Accurate design of megadalton-scale two-component icosahedral protein
1498   complexes. *Science* **353**, 389-394 (2016).
1499   88.   L. He *et al.*, Presenting native-like trimeric HIV-1 antigens with self-assembling
1500   nanoparticles. *Nat. Commun.* **7**, 12041 (2016).
1501   89.   L. He *et al.*, Proof of concept for rational design of hepatitis C virus E2 core nanoparticle
1502   vaccines. *Sci. Adv.* **6**, eaaz6225 (2020).
1503   90.   T. U. J. Bruun, A. M. C. Andersson, S. J. Draper, M. Howarth, Engineering a rugged
1504   nanoscaffold to enhance plug-and-display vaccination. *ACS Nano* **12**, 8855-8866 (2018).
1505   91.   L. He *et al.*, Single-component, self-assembling, protein nanoparticles presenting the
1506   receptor binding domain and stabilized spike as SARS-CoV-2 vaccine candidates. *Sci.*
1507   *Adv.* **7**, eabf1591 (2021).
1508   92.   L. He *et al.*, Single-component multilayered self-assembling nanoparticles presenting
1509   rationally designed glycoprotein trimers as Ebola virus vaccines. *Nat. Commun.* **12**, 2633
1510   (2021).
1511   93.   M. F. Bachmann, G. T. Jennings, Vaccine delivery: a matter of size, geometry, kinetics
1512   and molecular patterns. *Nat. Rev. Immunol.* **10**, 787-796 (2010).
1513   94.   T. Tokatlian *et al.*, Innate immune recognition of glycans targets HIV nanoparticle
1514   immunogens to germinal centers. *Science* **363**, 649-654 (2019).
1515   95.   B. J. Read *et al.*, Mannose-binding lectin and complement mediate follicular localization
1516   and enhanced immunogenicity of diverse protein nanoparticle immunogens. *Cell Rep.* **38**,
1517   110217 (2022).

96. R. P. Ringe *et al.*, Neutralizing antibody induction by HIV-1 envelope glycoprotein
SOSIP trimers on iron oxide nanoparticles may be impaired by mannose binding lectin. *J.*
Virol. **94**, e01883-01819 (2020).
97. T. J. Moyer *et al.*, Engineered immunogen binding to alum adjuvant enhances humoral
immunity. *Nat. Med.* **26**, 430–440 (2020).
98. S. Kumar *et al.*, A VH1-69 antibody lineage from an infected Chinese donor potently
neutralizes HIV-1 by targeting the V3 glycan supersite. *Sci. Adv.* **6**, eabb1328 (2020).
99. D. S. Rosa, F. Tzelepis, M. G. Cunha, I. S. Soares, M. M. Rodrigues, The pan HLA DR-
binding epitope improves adjuvant-assisted immunization with a recombinant protein
containing a malaria vaccine candidate. *Immunol. Lett.* **92**, 259-268 (2004).
100. P. J. M. Brouwer *et al.*, Immunofocusing and enhancing autologous Tier-2 HIV-1
neutralization by displaying Env trimers on two-component protein nanoparticles. *Npj*
Vaccines **6**, 24 (2021).
101. A. Antanasijevic *et al.*, Structural and functional evaluation of de novo-designed, two-
component nanoparticle carriers for HIV Env trimer immunogens. *PLoS Pathog.* **16**,
e1008665 (2020).
102. P. J. M. Brouwer *et al.*, Enhancing and shaping the immunogenicity of native-like HIV-1
envelope trimers with a two-component protein nanoparticle. *Nat. Commun.* **10**, 4272
(2019).
103. G. Ueda *et al.*, Tailored design of protein nanoparticle scaffolds for multivalent
presentation of viral glycoprotein antigens. *eLife* **9**, e57659 (2020).
104. T. Izard *et al.*, Principles of quasi-equivalence and Euclidean geometry govern the
assembly of cubic and dodecahedral cores of pyruvate dehydrogenase complexes. *Proc.*
Natl. Acad. Sci. U. S. A. **96**, 1240-1245 (1999).
105. Y. Hsia *et al.*, Design of a hyperstable 60-subunit protein icosahedron. *Nature* **535**, 136-
139 (2016).
106. A. J. Behrens, M. Crispin, Structural principles controlling HIV envelope glycosylation.
Curr. Opin. Struct. Biol. **44**, 125-133 (2017).
107. R. Kong *et al.*, Fusion peptide of HIV-1 as a site of vulnerability to neutralizing antibody.
Science **352**, 828-833 (2016).
108. T. Schoofs *et al.*, Broad and Potent Neutralizing Antibodies Recognize the Silent Face of
the HIV Envelope. *Immunity* **50**, 1513-+ (2019).
109. Z. T. Berndsen *et al.*, Visualization of the HIV-1 Env glycan shield across scales. *Proc.*
Natl. Acad. Sci. U. S. A. **117**, 28014-28025 (2020).
110. E. Falkowska *et al.*, Broadly neutralizing HIV antibodies define a glycan-dependent
epitope on the prefusion conformation of gp41 on cleaved envelope trimers. *Immunity* **40**,
657-668 (2014).
111. C. Blattner *et al.*, Structural delineation of a quaternary, cleavage-dependent epitope at
the gp41-gp120 interface on intact HIV-1 Env trimers. *Immunity* **40**, 669-680 (2014).
112. L. W. Cao *et al.*, Differential processing of HIV envelope glycans on the virus and
soluble recombinant trimer. *Nat. Commun.* **9**, 3693 (2018).
113. L. Kong *et al.*, Complete epitopes for vaccine design derived from a crystal structure of
the broadly neutralizing antibodies PGT128 and 8ANC195 in complex with an HIV-1
Env trimer. *Acta Crystallogr. D Biol. Crystallogr.* **71**, 2099-2108 (2015).
114. T. Zhou *et al.*, Structural basis for broad and potent neutralization of HIV-1 by antibody
VRC01. *Science* **329**, 811-817 (2010).

115. X. Wu *et al.*, Rational design of envelope identifies broadly neutralizing human
monoclonal antibodies to HIV-1. *Science* **329**, 856-861 (2010).
116. L. Kong *et al.*, Supersite of immune vulnerability on the glycosylated face of HIV-1
envelope glycoprotein gp120. *Nat. Struct. Mol. Biol.* **20**, 796-803 (2013).
117. J. Huang *et al.*, Broad and potent HIV-1 neutralization by a human antibody that binds
the gp41–gp120 interface. *Nature* **515**, 138-142 (2014).
118. D. Sok *et al.*, Promiscuous glycan site recognition by antibodies to the high-mannose
patch of gp120 broadens neutralization of HIV. *Sci. Transl. Med.* **6**, 236ra263 (2014).
119. L. Kong *et al.*, Key gp120 glycans pose roadblocks to the rapid development of VRC01-
class antibodies in an HIV-1-infected Chinese donor. *Immunity* **44**, 939-950 (2016).
120. T. Zhou *et al.*, Multidonor analysis reveals structural elements, genetic determinants, and
maturation pathway for HIV-1 neutralization by VRC01-class antibodies. *Immunity* **39**,
245-258 (2013).
121. R. Derking *et al.*, Comprehensive antigenic map of a cleaved soluble HIV-1 envelope
trimer. *PLoS Pathog.* **11**, e1004767 (2015).
122. R. Henderson *et al.*, Disruption of the HIV-1 envelope allosteric network blocks CD4-
induced rearrangements. *Nat. Commun.* **11**, 520 (2020).
123. Y. N. Zhang *et al.*, Mechanism of a COVID-19 nanoparticle vaccine candidate that elicits
a broadly neutralizing antibody response to SARS-CoV-2 variants. *Sci. Adv.* **7**, eabj3107
(2021).
124. R. W. Sanders *et al.*, HIV-1 neutralizing antibodies induced by native-like envelope
trimers. *Science* **349**, aac4223 (2015).
125. M. G. Pauthner *et al.*, Vaccine-induced protection from homologous tier 2 SHIV
challenge in nonhuman primates depends on serum-neutralizing antibody titers. *Immunity*
50, 241-252.e246 (2019).
126. M. Pauthner *et al.*, Elicitation of robust tier 2 neutralizing antibody responses in
nonhuman primates by HIV envelope trimer immunization using optimized approaches.
Immunity **46**, 1073-1088 (2017).
127. P. Martinez-Murillo *et al.*, Particulate array of well-ordered HIV clade C Env trimers
elicits neutralizing antibodies that display a unique V2 cap approach. *Immunity* **46**, 804-
817 (2017).
128. K. M. Cirelli *et al.*, Slow Delivery Immunization Enhances HIV Neutralizing Antibody
and Germinal Center Responses via Modulation of Immunodominance. *Cell* **177**, 1153-+
(2019).
129. K. V. Houser *et al.*, Safety and immunogenicity of an HIV-1 prefusion-stabilized
envelope trimer (Trimer 4571) vaccine in healthy adults: A first-in-human open-label,
randomized, dose-escalation, phase 1 clinical trial. *eClinicalMedicine* **Published Online**,
(2022).
130. R. Rappuoli, Glycoconjugate vaccines: principles and mechanisms. *Sci. Transl. Med.* **10**,
eaat4615 (2018).
131. J. G. Cyster, B cell follicles and antigen encounters of the third kind. *Nat. Immunol.* **11**,
989-996 (2010).
132. B. A. Heesters, R. C. Myers, M. C. Carroll, Follicular dendritic cells: dynamic antigen
libraries. *Nat. Rev. Immunol.* **14**, 495-504 (2014).

133. L. M. Walker *et al.*, A limited number of antibody specificities mediate broad and potent
serum neutralization in selected HIV-1 infected individuals. *PLoS Pathog.* **6**, e1001028
(2010).
134. D. Sok *et al.*, Recombinant HIV envelope trimer selects for quaternary-dependent
antibodies targeting the trimer apex. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 17624-17629
(2014).
135. Y. N. Zhang *et al.*, Nanoparticle size influences antigen retention and presentation in
lymph node follicles for humoral immunity. *Nano Lett* **19**, 7226-7235 (2019).
136. A. Singh, Eliciting B cell immunity against infectious diseases using nanovaccines. *Nat.*
Nanotechnol. **16**, 16-24 (2021).
137. R. Rappuoli, D. Serruto, Self-assembling nanoparticles usher in a new era of vaccine
design. *Cell* **176**, 1245-1247 (2019).
138. G. D. Victora, M. C. Nussenzweig, Germinal centers. *Annu. Rev. Immunol.* **30**, 429-457
(2012).
139. J. López-Sagaseta, E. Malito, R. Rappuoli, M. J. Bottomley, Self-assembling protein
nanoparticles in the design of vaccines. *Comput Struct Biotechnol J* **14**, 58-68 (2016).
140. Y. Li *et al.*, Mechanism of neutralization by the broadly neutralizing HIV-1 monoclonal
antibody VRC01. *J. Virol.* **85**, 8954-8967 (2011).
141. T. G. Phan, I. Grigorova, T. Okada, J. G. Cyster, Subcapsular encounter and
complement-dependent transport of immune complexes by lymph node B cells. *Nat.*
Immunol. **8**, 992-1000 (2007).
142. T. G. Phan, J. A. Green, E. E. Gray, Y. Xu, J. G. Cyster, Immune complex relay by
subcapsular sinus macrophages and noncognate B cells drives antibody affinity
maturation. *Nat. Immunol.* **10**, 786-793 (2009).
143. S. F. Gonzalez *et al.*, Capture of influenza by medullary dendritic cells via SIGN-R1 is
essential for humoral immunity in draining lymph nodes. *Nat. Immunol.* **11**, 427-434
(2010).
144. C. D. Allen, T. Okada, J. G. Cyster, Germinal-center organization and cellular dynamics.
Immunity **27**, 190-202 (2007).
145. L. Mesin, J. Ersching, G. D. Victora, Germinal center B cell dynamics. *Immunity* **45**,
471-482 (2016).
146. S. Crotty, T follicular helper cell differentiation, function, and roles in disease. *Immunity*
41, 529-542 (2014).
147. C. Viant *et al.*, Antibody affinity shapes the choice between memory and germinal center
B cell fates. *Cell* **183**, 1298-1311.e1211 (2020).
148. E. T. Crooks *et al.*, Engineering well-expressed, V2-immunofocusing HIV-1 envelope
glycoprotein membrane trimers for use in heterologous prime-boost vaccine regimens.
PLoS Pathog. **17**, e1009807 (2021).
149. K. Xu *et al.*, Epitope-based vaccine design yields fusion peptide-directed antibodies that
neutralize diverse strains of HIV-1. *Nat. Med.* **24**, 857–867 (2018).
150. P. J. Klasse, G. Ozorowski, R. W. Sanders, J. P. Moore, Env exceptionalism: why are
HIV-1 Env glycoproteins atypical immunogens? *Cell Host Microbe* **27**, 507-518 (2020).
151. T. Q. Zhou *et al.*, Quantification of the impact of the HIV-1-glycan shield on antibody
elicitation. *Cell Rep.* **19**, 719-732 (2017).
152. M. M. Shan *et al.*, HIV-1 gp120 mannoses induce immunosuppressive responses from
dendritic cells. *PLoS Pathog.* **3**, 1637-1650 (2007).

153. K. Banerjee *et al.*, Enzymatic removal of mannose moieties can increase the immune
response to HIV-1 gp120 in vivo. *Virology* **389**, 108-121 (2009).
154. H.-Y. Huang *et al.*, Vaccination with SARS-CoV-2 spike protein lacking glycan shields
elicits enhanced protective responses in animal models. *Sci. Transl. Med.* **14**, eabm0899
(2022).
155. J. R. Chen *et al.*, Vaccination of monoglycosylated hemagglutinin induces cross-strain
protection against influenza virus infections. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 2476-
2481 (2014).
156. K. Zboray *et al.*, Heterologous protein production using euchromatin-containing
expression vectors in mammalian cells. *Nucleic Acids Res.* **43**, e102 (2015).
157. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).
158. G. Ozorowski *et al.*, Effects of adjuvants on HIV-1 envelope glycoprotein SOSIP trimers
in vitro. *J. Virol.* **92**, e00381-00318 (2018).
159. C. Suloway *et al.*, Automated molecular microscopy: The new Leginon system. *J. Struct.*
Biol. **151**, 41-60 (2005).
160. G. C. Lander *et al.*, Appion: An integrated, database-driven pipeline to facilitate EM
image processing. *J. Struct. Biol.* **166**, 95-102 (2009).
161. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat. Methods* **14**, 331-332 (2017).
162. J. Zivanov *et al.*, New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *eLife* **7**, e42166 (2018).
163. S. L. Ilca *et al.*, Localized reconstruction of subunits from electron cryomicroscopy
images of macromolecular complexes. *Nat. Commun.* **6**, 8843 (2015).
164. P. Emsley, M. Crispin, Structural analysis of glycoproteins: building N-linked glycans
with Coot. *Acta Crystallogr. D Biol. Crystallogr.* **74**, 256-263 (2018).
165. R. Y. R. Wang *et al.*, Automated structure refinement of macromolecular assemblies
from cryo-EM maps using Rosetta. *Elife* **5**, e17219 (2016).
166. B. A. Barad *et al.*, EMRinger: side chain directed model and map validation for 3D cryo-
electron microscopy. *Nat. Methods* **12**, 943-946 (2015).
167. C. J. Williams *et al.*, MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci.* **27**, 293-315 (2018).
168. M. Sarzotti-Kelsoe *et al.*, Optimization and validation of the TZM-bl assay for
standardized assessments of neutralizing antibodies against HIV-1. *J. Immunol. Methods*
409, 131-146 (2014).
169. A. deCamp *et al.*, Global panel of HIV-1 Env reference strains for standardized
assessments of vaccine-elicited neutralizing antibodies. *J. Virol.* **88**, 2489-2507 (2014).
170. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**, 671-675 (2012).

1693
1694

Figure Legends

Fig. 1. Rational design and characterization of BG505 UFO trimer-presenting

nanoparticles. (**A**) BG505 UFO Env (middle) and three UFO trimer-presenting SApNPs (FR:

left; E2p and I3-01: right). In the inset (center), ribbon representation of BG505 UFO gp140

protomer (PDB ID: 6UTK) is shown in transparent molecular surface with gp120 in cornflower

blue, gp41 in magenta, and the redesigned $HR1_N$ loop in yellow (as both ribbon and molecular

surface models). Surface representations are shown for FR 24-mer and E2p/I3-01 60-mers

presenting 8 and 20 BG505 UFO trimers, respectively. The NP carrier is colored in gray and the

three protomers of each BG505 UFO trimer are colored differently. The NP size is indicated by

diameter (in nm). (**B**) Schematic representation of a multilayered NP (left), which contains an

inner layer of locking domains (LD) and a cluster of T-help epitopes (PADRE), and molecular

surface model of E2p (brown, PDB ID: 1B5S) with stepwise incorporation of LD4 (cyan, PDB

ID: 2MG4) and PADRE (purple). (**C**) SEC profiles of BG505 UFO trimer-presenting FR (panel

1), E2p (panel 2), E2p-LD1-7 (panel 3), E2p-LD4-PADRE (or E2p-L4P, panel 4), I3-01 (panel

5), I3-01-LD4-5/7-9 (panel 6), and I3-01/v9-LD7-PADRE (or I3-01/v9-L7P, panel 7) after

PGT145 purification. (**D**) and (**E**) Reducing SDS-PAGE and BN-PAGE of BG505 UFO-FR,

E2p, and I3-01/v9 SApNPs after PGT145 purification, with LD and PADRE variants included

for E2p and I3-01. (**F**) Negative stain EM images of PGT145-purified BG505 UFO-FR, E2p-

L4P, and I3-01v9-L7P SApNPs. (**G**) Thermostability of BG505 UFO-FR, E2p-L4P, and I3-

01v9-L7P SApNPs, with T_m, $\Delta T_{1/2}$, and T_{on} measured by DSC. (**H**) Particle size distribution of

BG505 UFO-FR, E2p-L4P, and I3-01v9-L7P SApNPs. The hydrodynamic diameter (D_h) was

measured by DLS. Multiple measurements were obtained for each SApNP, with the average

particle size labeled on the distribution plot.

Fig. 2. Cryo-EM analysis of the multilayered BG505 UFO trimer-presenting nanoparticles.

Two PGT145/SEC-purified SApNPs, BG505 UFO-E2p-L4P (**A**) and UFO-10GS-I3-01v9-L7P

(**B**), were analyzed. Representative raw cryo-EM micrographs, 2D classes and 3D maps

featuring full NPs (with trimer antigens) are displayed in the top left part of the corresponding

panels. Focused classification and refinement were applied to reconstruct a 3D map of the

SApNP core (top and bottom right), while the localized reconstruction approach was used for

extraction and analysis of trimer subparticles (bottom left). The maps are represented as

transparent gray mesh and the corresponding models are shown as ribbons (NPs - orange; trimer

antigens - blue). The refined model is presented for the E2p NP backbone, while the I3-01v9 and

trimer maps are fitted with previously reported models (PDB ID: 5KP9 and 6UTK).

Fig. 3. Glycan analysis of BG505 Env immunogens with wildtype and modified glycans. (**A**)

Comparison of site-specific occupancy and oligomannose-type glycan content for the BG505

UFO trimer (wildtype, swainsonine, and kifunensine) and three SApNPs (wildtype). The plotted

data represent the mean of three analytical repeats, with trimers shown as circles and NPs as

squares. The oligomannose content was determined by adding all glycan compositions

containing HexNAc(2). (**B**) SEC profiles of BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P

SApNPs upon endo H treatment. (**C**) Reducing SDS-PAGE of BG505 UFO-E2p-L4P and UFO-

10GS-I3-01v9-L7P SApNPs before and after glycan trimming by endo H. (**D**) EM images of

glycan-trimmed BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P SApNPs. (**E**)

Thermostability of glycan-trimmed BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P

SApNPs, with T_m, $\Delta T_{1/2}$, and T_{on} measured by DSC. (**F**) Reducing SDS-PAGE of BG505 UFO-

E2p-L4P and UFO-10GS-I3-01v9-L7P SApNPs before and after endo H treatment under two

different conditions: 25°C for 5 hours (condition 1) and protein denaturation at 100°C followed

by treatment at 37°C for 1 hour (condition 2). (**G**) UPLC analysis of released glycans from

BG505 UFO-10GS-I3-01v9 SApNP. Top: natively glycosylated; Middle: endo H-treated;

Bottom: overlay of two UPLC traces with the treated trace (from middle panel) recolored in

white. The endo H-treated trace has been normalized to the untreated trace using the complex-

type glycan peak eluting at ~8 min. In top and middle panels: oligomannose-type and complex-

type glycans are colored in green and pink, respectively. In the top panel, oligomannose-type

glycan peaks are labelled with their representative structures. For both UPLC traces, the

percentage of each oligomannose type is listed on the right.

Fig. 4. Antigenicity of BG505 UFO Env immunogens with wildtype and modified glycans.

(**A**) Antigenic profiles of BG505 UFO trimer (top) and its three SApNPs based on FR, E2p-L4D,

and I3-01v9-L7P (bottom) with different glycan treatments (wildtype, expressed in the presence

of kifunensine, and endo H treatment). A total of 13 NAbs/bNAbs and eight non-NAbs were

tested by BLI. Sensorgrams were obtained from an Octet RED96 using an antigen titration series

of six concentrations (starting at 266.7 nM for the UFO trimer, 14.9 nM for the FR SApNP, 5.5

nM for the E2p and I3-01v9 SApNPs followed by two-fold dilution) and are shown in Fig. S4.

The peak values at the highest concentration are summarized in the matrix, in which cells are

colored in red and green for (b)NAbs and non-NAbs, respectively. Higher color intensity

indicates greater binding signal measured by Octet. (**B**) Negative-stain EM analysis of the

glycan-trimmed BG505 UFO trimer in complex with bNAb PGT128. Examples of 2D class

images (top) and comparison with a cryo-EM model of BG505 SOSIP/PGT128 complex

(bottom, EMDB-3121, cornflower blue). (**C**) Negative-stain EM analysis of the glycan-trimmed

BG505 UFO trimer in complex with bNAb VRC01. Examples of 2D class images (top) and

superposition of the 3D reconstruction onto two previously reported BG505 SOSIP/VRC01

models derived from negative-stain EM (EMDB-6252, orange) and cryo-EM (PDB ID: 6V8X,

cyan and green).

Fig. 5. Immunogenicity of HIV-1 UFO Env immunogens with wildtype glycans in mice and

rabbits. (**A**) Schematic representation of the mouse immunization regimen for two BG505 UFO

trimer-presenting E2p and I3-01v9 SApNPs. (**B**) Neutralization of tier 2 clade A BG505.T332N

and tier 1 clade B SF162 by purified mouse IgG from week 11. (**C**) Percent neutralization of

week 11 mouse IgG from four vaccine responders against BG505.T332N and its I396R mutant.

Vaccine responders are those subjects with $IC_{50} < 300$ μg/ml in IgG neutralization against

BG505.T332N. Color coding indicates the level of reduction in % neutralization relative to

BG505.T332N. In both (B) and (C), the TZM-bl assay was performed in duplicate, starting at an

IgG concentration of 300 μg/ml (unless otherwise indicated by *) and a 3-fold dilution series.

(**D**) Schematic representation of the rabbit immunization protocol. Seven groups of rabbits, 6 per

group, were immunized to test the BG505 UFO trimer and its FR, E2p, and I3-01v9 SApNPs,

with a cocktail group added for each SApNP by mixing equal amounts of three SApNPs that

display BG505 UFO, CH505 UFO-BG, or H078.14 UFO-BG trimers (termed Env mix). (**E**)

Longitudinal analysis of rabbit plasma neutralization between weeks 2 and 28. For each group,

average ID_{50} values against BG505.T332N are shown with the SEM. Groups that were

immunized with the same vaccine platform are plotted on the same graph: trimer (red), FR

(dark/light purple), E2p (dark/light green) and I3-01v9 (dark/light blue). Week 0 data are omitted

due to nonspecific background signals. (**F**) Percentage neutralization of select week 26 rabbit

plasma (≥ 35% neutralization against BG505.T332N at the first dilution) against BG505.T332N

and four glycan hole mutants. For each mutant virus, rabbit plasma was tested in duplicate,

starting at 100-fold dilution followed by a 3-fold dilution series. Results are shown for

BG505.T332N from (E) and the mutant virus at the 100-fold dilution. Color coding indicates the

level of reduction in % neutralization relative to BG505.T332N.

Fig 6. Immunogenicity of BG505 UFO Env immunogens with trimmed glycans in mice and rabbits. (**A**) Schematic representation of mouse and rabbit immunization regimen 1, where glycan-trimmed immunogens were used throughout, and regimen 2, where glycan-trimmed and wildtype immunogens were used as prime (doses 1 and 2) and boost (doses 3 and 4), respectively. A three-week interval was used to facilitate the rapid evaluation of various vaccine formulations. The number of animals in each group is the same as in **Fig. 5**. (**B**) Frequency of vaccine responders (FVR) calculated for groups in which mice were immunized with the BG505 UFO trimer (left), E2p SApNP (middle), and I3-01v9 SApNP (right). Purified mouse IgGs from week 11 were tested against tier 2 clade A BG505.T332N. The TZM-bl neutralization assay was done in duplicate at a starting concentration of 300 μg/ml and followed by a 3-fold dilution series. A vaccine responder is defined as a subject with $IC_{50} \leq 300$ μg/ml. No IgG was available for one mouse in the UFO trimer/AP regimen 1 group and two mice in the I3-01v9/AP regimen 1 (1[st] repeat) group. In the comparator groups, no IgG was available for one mouse each in the wildtype E2p/AP and E2p/AV groups. (**C**) Percent neutralization values of week 11 mouse IgG from select mouse samples ($\geq$ 30% autologous neutralization at the highest IgG concentration) against BG505.T332N and its I396R mutant. Color coding indicates the level of reduction in % neutralization relative to BG505.T332N. (**D**) Neutralization of tier 2 clade A BG505.T332N by purified rabbit IgG from week 11. (**E**) FVR of four rabbit groups based on the week 11 autologous neutralization data, with vaccine responder defined as in (B). (**F**) Percentage neutralization of select week 11 rabbit IgG samples ($\geq$ 30% autologous neutralization at the highest IgG concentration) against BG505.T332N and its four glycan hole mutants. The TZM-bl assay was done in duplicate at a starting concentration of 300 ug/ml and followed by a 3-fold

dilution series. Color coding from red to green indicates greater to lesser reduction in %

neutralization relative to BG505.T332N.

Fig. 7. Immunogenicity of BG505 UFO Env immunogens with wildtype and trimmed

glycans in NHPs. (**A**) Schematic representation of the immunization regimen for wildtype

BG505 UFO trimer-presenting E2p and I3-01v9 SApNPs. E2p was formulated with AddaVax

and I3-01v9 with aluminum phosphate (N = 6 animals/group). (**B**) Neutralization of tier 2 clade

A BG505.T332N by rhesus macaque (RM) sera from week 28. Neutralization curves and ID_{50}

titers are shown. (**C**) Schematic representation of the immunization regimen 1, where glycan-

trimmed immunogens were used throughout, and regimen 2, where glycan-trimmed and wildtype

immunogens were used as prime (doses 1 and 2) and boost (doses 3 and 4), respectively.

SApNPs were formulated with aluminum phosphate for all groups in both regimens (N = 4

animals/group). (**D**) Neutralization of tier 2 clade A BG505.T332N by NHP sera from week 28.

Neutralization curves and ID_{50} titers are shown. (**E**) FVR comparison for E2p and I3-01v9

SApNPs using the week-28 NHP serum neutralization data against BG505.T332N. Left:

wildtype; middle and right: glycan-trimmed (regimens 1 and 2). (**F**) Select NHP sera from week

28 (≥ 40% autologous neutralization) against BG505.T332N and its variant with the T465N

mutation in the C3/465 epitope. The TZM-bl neutralization assay was performed in duplicate at a

starting dilution of 40 and followed by a 3-fold dilution series. A vaccine responder is defined as

a subject with $ID_{50} \geq 40$.

Fig. 8. BG505 UFO Env-SApNP vaccines induce long-term retention in lymph node

follicles. (**A**) BG505 UFO trimer-presenting I3-01v9 SApNP distribution in a lymph node 12

hours after a single-dose injection (10 μg/injection, 40 μg/mouse). Anti-Env VRC01, PGT124,

and PGDM1400 bNAbs were used to stain the lymph node tissue sections. (**B**) Schematic

illustration of I3-01v9 SApNP accumulation in a lymph node. (**C**) Histological images of the

distribution of the BG505 UFO trimer and trimer-presenting E2p and I3-01v9 SApNP trafficking

and retention in lymph node follicles 30 minutes to 8 weeks after a single-dose injection. Scale

bar = 50 μm for each image. All immunogens had wildtype glycans except that a glycan-

trimmed (GT) I3-01v9 SApNP was included for comparison. (**D**) Time-dependent curve and (**E**)

Area under the curve (AUC) of the VRC01-stained area in histological images of the vaccine

retention in lymph node follicles over 8 weeks. (**F**) Quantification of vaccine accumulation in

lymph node follicles 48 hours after a single-dose injection. Data were collected from more than

10 lymph node follicles (n = 3-5 mice/group). (**G**) BG505 UFO trimer and trimer-presenting

SApNP vaccine interaction with FDC networks in lymph node follicles 48 hours after a single-

dose injection. All BG505 Env-SApNP immunogens, regardless of the NP platform and glycan

treatment, were colocalized with FDC networks. Immunofluorescent images are pseudo-color-

coded (CD21$^+$, green; CD169$^+$, red; VRC01, white). Scale bars = 500 and 100 μm for complete

lymph node and enlarged image of a follicle, respectively. The data points are expressed as mean

± SEM for (**D**) and SD for (**F**). The data were analyzed using one-way ANOVA followed by

Tukey's multiple comparison post hoc test. ****$p < 0.0001$.

Fig. 9. Representative TEM images of BG505 UFO Env-SApNPs interacting with FDCs

and phagocytic cells in lymph nodes. (**A**) TEM images of BG505 UFO-E2p-L4P SApNPs

(yellow arrows) and AddaVax adjuvant (green arrows) presented on the FDC dendrites in a

lymph node 12 hours after a single-dose injection (50 μg/injection). (**B**) and (**C**) TEM images of

BG505 UFO-10GS-I3-01v9-L7P SApNPs (yellow arrows), wildtype and glycan-trimmed (GT),

aligned on FDC dendrites without observable aluminum phosphate (AP) adjuvant 12 hours after

a single-dose injection. (**D**) TEM images of BG505 UFO-E2p-L4P SApNPs (yellow arrows) and

AddaVax adjuvant (green arrows) colocalized on the surface or inside endolysosomes of a

macrophage 12 hours after injection. (**E**) and (**F**) TEM images of BG505 UFO-10GS-I3-01v9-

L7P SApNPs (yellow arrows), wildtype and glycan-trimmed (GT), and aluminum phosphate

adjuvant (green arrows) internalized inside endolysosomes of a macrophage, with only SApNPs

(yellow arrows) on the macrophage surface (E) 48 hours, or (F) 12 hours after the injection.

Fig. 10. H**IV-1 SApNP vaccines induce robust long-lived germinal centers in lymph nodes.**

(A) Top: Representative immunohistological images of BG505 UFO-10GS-I3-01v9 SApNP

vaccine induced germinal centers (GCs) at two weeks after a single-dose injection (10

μg/injection, 40 μg/mouse). Bottom left: GC B cells (GL7+, red) adjacent to FDCs (CD21+,

green) in lymph node follicles. Bottom Right: Tfh cells in the light zone (LZ) of GCs. Scale bars

= 500 and 100 μm for complete lymph node and enlarged image of a follicle, respectively. (**B**)

and (**C**) Quantification of GCs using immunofluorescent images at weeks 2, 5, and 8 after (B) a

single-dose or (C) prime-boost injections (n = 5-10 mice/group). The GC/FDC ratio and GC size

were determined and plotted here. (**D**) and (**E**) Representative immunohistological images of

GCs included by the BG505 UFO trimer, E2p, I3-01v9, and GT I3-01v9 SApNP vaccines at

week 8 after (D) a single-dose or (E) prime-boost injections. Scale bar = 50 μm for each image.

(**F**) and (**G**) Quantification of GCs by flow cytometry after (F) a single-dose or (G) prime-boost

injections. Percentage and number of GC B cells and T_{fh} cells were determined and plotted. The

data points are shown as mean ± SD. The data were analyzed using one-way ANOVA followed

by Tukey's multiple comparison post hoc test for each timepoint. $*p < 0.05$, $**p < 0.01$, $***p <$

0.001, $****p < 0.0001$.

 SUPPLEMENTAL LEGENDS

Table S1. Cryo-EM data collection information.

Table S2. Model building and refinement statistics for the BG505 UFO E2p-L4P NP core.

Figure S1. Design and in vitro characterization of BG505 UFO trimer-presenting SApNPs.

Figure S2. Cryo-EM analysis of BG505 UFO-E2p-L4P and UFO-10GS-I3-01v9-L7P SApNPs.

Figure S3. Site-specific glycan analysis of BG505 UFO trimer and SApNPs.

Figure S4. Antigenic characterization of BG505 UFO trimer and trimer-presenting SApNPs with

wildtype and modified glycans.

Figure S5. Neutralization data from the evaluation of Env immunogens with wildtype glycans in

mice and rabbits.

Figure S6. Neutralization data from the evaluation of BG505 immunogens with trimmed glycans

in mice and rabbits.

Figure S7. Neutralization data from the evaluation of BG505 SApNP immunogens with

wildtype and trimmed glycans in NHPs.

Figure S8. Immunohistological images of BG505 UFO trimer and SApNPs in lymph nodes.

Figure S9. TEM images of BG505 UFO trimer and SApNPs interacting with FDCs and

phagocytic cells in a lymph node.

Figure S10. Immunohistological analysis of BG505 UFO and SApNP vaccine-induced GCs.

Figure S11. Flow cytometry analysis of BG505 UFO trimer and SApNP vaccine-induced GCs.

Figure 2

Figure 3

A

Trimer comparison

	PGDM1400	PGT145	PG16	PGT121	PGT128	PGT135	438-B11	2G12	VRC01	VRC34	PGT151	35O22	SF12	M4H2K1	F105	b6	19b	446-52D	F240	7B2	17b	A32	RM20A3
BG505 UFO trimer (WT)	1.0	0.5	0.5	0.7	0.8	0.4	0.7	0.6	0.9	0.8	0.6	0.6	0.6	0.7	0.1	0.3	0.7	0.5	0.0	0.0	0.0	0.1	0.5
BG505 UFO trimer (KIF)	0.2	0.2	0.1	0.3	0.6	0.2	0.4	0.4	0.5	0.5	0.0	0.5	0.5	0.4	0.1	0.2	0.4	0.2	0.0	0.0	0.1	0.0	0.5
BG505 UFO trimer (endo H)	0.2	0.2	0.1	0.3	0.4	0.4	0.3	0.4	0.8	0.4	0.1	0.3	0.5	0.4	0.1	0.3	0.3	0.2	0.1	0.0	0.1	0.1	0.5

Trimer-SApNP comparison

	PGDM1400	PGT145	PG16	PGT121	PGT128	PGT135	438-B11	2G12	VRC01	VRC34	PGT151	35O22	SF12	M4H2K1	F105	b6	19b	446-52D	F240	7B2	17b	A32	RM20A3
BG505 UFO trimer (WT)	1.1	0.6	0.5	0.7	0.7	0.4	0.8	0.6	0.8	0.9	0.6	0.6	0.9	0.6	0.2	0.3	0.7	0.5	0.0	0.0	0.0	0.1	0.8
BG505 UFO-FR (WT)	2.7	2.1	1.7	1.5	2.1	0.8	1.5	1.9	1.1	0.4	0.2	0.3	0.6	1.3	0.2	0.3	1.8	1.4	0.0	0.0	0.0	0.1	0.0
BG505 UFO-E2p-L4P (WT)	3.3	2.8	2.3	1.7	2.7	1.0	1.9	2.6	1.3	0.5	0.1	0.4	0.5	1.6	0.2	0.3	1.9	2.0	0.1	0.1	0.1	0.1	0.2
BG505 UFO-10GS-I3-01v9-L7P (WT)	2.9	2.5	2.0	1.5	2.3	1.0	1.7	2.3	1.3	0.5	0.4	0.4	0.5	1.5	0.2	0.3	1.8	1.7	0.1	0.0	0.0	0.1	0.1
BG505 UFO trimer (KIF)	0.4	0.5	0.2	0.6	0.7	0.3	0.7	0.6	0.8	0.8	0.0	0.7	0.8	0.6	0.2	0.4	0.7	0.4	0.0	0.0	0.1	0.1	0.8
BG505 UFO-FR (KIF)	2.3	2.2	1.6	1.2	2.0	0.7	1.7	1.7	1.0	0.2	0.0	0.2	1.2	1.2	0.2	0.3	1.5	1.3	0.0	0.0	0.0	0.1	0.0
BG505 UFO-E2p-L4P (KIF)	2.6	2.7	1.7	1.3	2.6	1.1	2.2	2.4	0.9	0.3	0.0	0.5	1.5	1.3	0.2	0.3	1.8	1.9	0.0	0.0	0.1	0.1	0.0
BG505 UFO-10GS-I3-01v9-L7P (KIF)	2.3	2.4	1.7	1.3	2.3	1.2	1.9	2.1	1.1	0.3	0.0	0.5	1.3	1.4	0.3	0.4	1.9	1.7	0.1	0.0	0.0	0.1	0.0
BG505 UFO trimer (endo H)	0.5	0.4	0.1	0.6	0.6	0.6	0.6	0.6	1.1	0.6	0.4	0.5	0.7	0.6	0.3	0.5	0.6	0.4	0.0	0.0	0.0	0.1	0.8
BG505 UFO-FR (endo H)	2.6	2.1	1.9	1.5	1.9	1.4	1.6	1.9	2.2	0.7	0.1	0.7	0.6	1.7	0.4	0.5	1.9	1.6	0.1	0.0	0.1	0.1	0.0
BG505 UFO-E2p-L4P (endo H)	2.8	2.2	2.5	1.7	2.4	1.7	1.7	2.2	2.5	0.7	0.1	0.7	0.7	2.1	0.2	0.3	1.9	1.8	0.1	0.0	0.1	0.1	0.1
BG505 UFO-10GS-I3-01v9-L7P (endo H)	2.3	2.2	2.1	1.6	2.2	1.6	1.6	2.2	2.4	0.7	0.0	0.8	0.6	2.0	0.3	0.5	1.8	1.8	0.0	0.0	0.0	0.1	0.1

B — Glycan-trimmed (GT) BG505 UFO trimer with PGT128

2D class comparison

BG505 SOSIP/PGT128 (EMDB-3121)

C — Glycan-trimmed (GT) BG505 UFO trimer with VRC01

3D model fitting

90°

BG505 SOSIP/VRC01 (EMDB-6252)

BG505 SOSIP F14/VRC01 (PDB ID: 6V8X)

A

B

C

% neutralization of mouse IgG against BG505.T332N.I396R mutant [a.]				
Vaccine:	BG505 E2p SApNP		BG505 I3-01v9 SApNP	
Animal ID:	G2-M4*	G3-M2	G3-M1	G4-M6
BG505.T332N	33	84	94	93
BG505.T332N.I396R	6	17	52	26

a. Color coding indicates percentage reduction with respect to the % neutralization value obtained for BG505.T332N, <25% green, 25-50% yellow, 50-75% orange, >75% red.
*results shown for this IgG are at ~100 µg/ml for both BG505.T332N & mutant viruses

D

E

F

% neutralization of W26 rabbit sera against BG505.T332N glycan hole mutants [b.]																						
Vaccine	BG505 UFO trimer				BG505 FR SApNP				Env Mix FR SApNP		BG505 E2p SApNP					Env Mix E2p SApNP		BG505 I3-01v9 SApNP		Env Mix I3-01v9 SApNP			
Animal ID	121	122	123	124	126	127	128	130	132	143	137	156	158	159	160	141[c.]	143[c.]	146	147	149	150	151	152	154
BG505.T332N	72	83	64	82	95	95	49	95	37	73	93	45	47	55	96	78		94	84	98	88	86	99	67	42
BG505.T332N.**Q130N**	86	86	75	89	98	90	69	98	61	90	98	45	59	72	96	60		85	84	99	93	67	98	92	24
BG505.T332N.**S241N**	6	69	54	-19	82	85	63	51	37	72	86	48	56	12	88	73		87	74	82	83	77	82	69	51
BG505.T332N.**P291T**	-12	47	31	-14	92	80	27	94	24	63	73	43	37	25	91	68		94	67	97	83	65	96	31	17
BG505.T332N.**T465N**	79	85	42	79	50	43	23	67	24	38	83	24	23	64	36	33		72	52	54	33	62	38	58	21

b. Color coding indicates percentage reduction with respect to the % neutralization value obtained for BG505.T332N, <25% green, 25-50% yellow, 50-75% orange, >75% red.
c. Against BG505.T332N.Q130N: W28 plasma used for rabbits 141 and 143. The % neutralization change is <25% in comparison to BG505.T332N at either W28 (data not shown) or W26 (above).

Figure 6

A

| GT: glycan trimmed | WT: wildtype |

Regimen 1: GT (W0), GT (W3), GT (W6), GT (W9), blood draws W2, W5, W8, W11

Regimen 2: W0, GT (W3), WT (W6), WT (W9), blood draws W2, W5, W8, W11

Immunization
Immunogen: Protein (30 μg)
Adjuvant: Aluminum phosphate, AddaVax
Total volume: 200 μl
Route: Intraperitoneal

B Neutralization against tier 2 BG505.T332N by purified mouse IgG from week 11

BG505 Trimmer — ** (p=0.001)
BG505 E2p SApNP — * (p=0.041)
BG505 I3-01v9 SApNP — ns (p=0.282)

Legend: Responder (orange), Non-responder (grey)

C

% neutralization of mouse IgG against BG505.T332N.I396R mutant [a.]																			
Vaccine	**BG505 UFO trimer (Aluminum phosphate)**															**BG505 E2p SApNP (AddaVax)**			
Glycan trimming regimen	**Regimen 1**							**Regimen 2**								**Regimen 1**			**Regimen 2**
Animal ID	G3-M1	G3-M2	G3-M4	G3-M5	G3-M6	G3-M7	G3-M8	G4-M1	G4-M2	G4-M3	G4-M4	G4-M5	G4-M6	G4-M7	G4-M8	G1-M2	G1-M3	G1-M5	G2-M4	G2-M8
BG505.T332N	95	51	76	98	97	56	37	70	86	97	68	95	33	38	84	51	32	90	32	86
BG505.T332N. I396R	37	28	63	28	29	46	29	32	24	30	34	33	31	42	42	89	36	27	13	95

Vaccine	**BG505 E2p SApNP (Aluminum phosphate)**														**BG505 I3-01v9 SApNP (Aluminum phosphate)**			
Glycan trimming regimen	**Regimen 1**							**Regimen 2**						**Regimen 1**			**Regimen 2**	
Animal ID	G1-M1	G1-M2	G1-M3	G1-M4	G1-M5	G1-M6	G1-M7	G2-M2	G2-M3	G2-M4	G2-M6	G2-M7	G2-M8	G3-M1	G3-M3	G3-M8	G4-M2	G4-M3	G4-M4	G4-M5
BG505.T332N	49	71	73	71	93	39	89	92	98	60	88	91	38	70	49	91	93	82	91	87
BG505.T332N. I396R	40	71	87	80	86	39	30	89	58	62	92	41	37	66	43	43	98	94	96	46

a. Color coding indicates percentage reduction with respect to the % neutralization value obtained for BG505.T332N, <25% green, 25-50% yellow, 50-75% orange, >75% red.

D Neutralization against tier 2 BG505.T332N by purified rabbit IgG from week 11

BG505 E2p SApNP (AddaVax) — Regimen 1, Regimen 2
BG505 I3-01v9 SApNP (Aluminum phosphate) — Regimen 1, Regimen 2

Legend: RB1, RB2, RB3, RB4, RB5, RB6, RB7, RB8

E

% rabbits with detectable autologous neutralization; E2p (AV) regimen 1, E2p (AV) regimen 2, I3-01v9 (AP) regimen 1, I3-01v9 (AP) regimen 2; Responder / Non-responder

F

% neutralization of rabbit IgG against BG505.T332N glycan hole mutants [b.]										
Vaccine	**BG505 E2p SApNP**				**BG505 I3-01v9 SApNP**					
Glycan trimming regimen	**Regimen 1**		**Regimen 2**		**Regimen 1**			**Regimen 2**		
Animal ID	G1-2	G1-6	G2-1	G2-3	G3-2	G3-3	G3-4	G4-2	G4-3	G4-4	G4-5	G4-6
BG505.T332N	97	54	59	34	96	61	84	82	32	87	64	94
BG505.T332N.**Q130N**	97	53	69	38	97	58	90	87	5	91	77	96
BG505.T332N.**S241N**	99	2	89	26	100	32	96	90	71	91	62	98
BG505.T332N.**P291T**	96	2	74	7	97	11	88	86	53	88	59	95
BG505.T332N.**T465N**	16	59	16	39	36	59	26	65	54	39	49	95

b. Color coding indicates percentage reduction with respect to the % neutralization value obtained for BG505.T332N, <25% green, 25-50% yellow, 50-75% orange, >75% red.

A

B **Neutralization against tier 2 BG505.T332N by RM sera from week 28. Immunization with wildtype SApNPs**

Vaccine	Adjuvant	W28 ID$_{50}$ titers					
		RM1	RM2	RM3	RM4	RM5	RM6
BG505 E2p SApNP	AddaVax	<40	49	<40	<40	<40	<40
BG505 I3-01v9 SApNP	Aluminum phosphate	<40	<40	<40	<40	<40	<40

C

D **Neutralization against tier 2 BG505.T332N by RM sera from week 28. Immunization with glycan trimmed SApNPs**

Vaccine	Adjuvant	Glycan trimming regimen	W28 ID$_{50}$ titers			
			RM1	RM2	RM3	RM4
BG505 E2p SApNP	Aluminum phosphate	Regimen 1	78	<40	<40	404
		Regimen 2	<40	121	<40	80
BG505 I3-01v9 SApNP	Aluminum phosphate	Regimen 1	83	<40	<40	97
		Regimen 2	<40	44	<40	<40

E

F

% neutralization of W28 RM sera against BG505.T332N with the V5 mutation (T465N) [a.]									
Vaccine	BG505 E2p SApNP (AddaVax)	BG505 E2p SApNP (Aluminum phosphate)					BG505 I3-01v9 SApNP (Aluminum phosphate)		
Glycan trimming regimen	WT	Regimen 1			Regimen 2		Regimen 1		Regimen 2
Animal ID	G2-RM2	G1-RM1	G1-RM3	G1-RM4	G2-RM2	G2-RM4	G2-RM1	G3-RM2	G3-RM4	G4-RM2
BG505.T332N	62	69	43	96	80	65	69	47	80	56
BG505.T332N.**T465N**	16	28	31	17	7	29	8	-9	12	66

a. Color coding indicates percentage reduction with respect to the % neutralization value obtained for BG505.T332N, <25% green, 25-50% yellow, 50-75% orange, >75% red.

Figure 8

Figure 9

Figure 10