When should bees be flower constant? An agent-based model

highlights the importance of social information and foraging

conditions

Lucy Hayes & Christoph Grüter

School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, BS8 1TQ Bristol,

UK

Correspondence: c.grueter@bristol.ac.uk

Abstract

1. Many bee species show flower constancy, *i.e.* a tendency to visit flowers of one type

during a foraging trip. Flower constancy is important for plant reproduction, but whether

bees also benefit from flower constancy remains unclear. Social bees, which often use

communication about food sources, show particularly strong flower constancy.

2. We hypothesised that the sharing of social information increases the benefits of flower

constancy because foragers share information selectively about high-quality food

sources, thereby reducing the need to sample alternatives. We also asked if foraging

landscapes affect flower constancy. We developed an agent-based model that allowed

us to simulate bee colonies with and without communication and flower constancy in

different foraging environments.

3. Flower constancy alone performed poorly in all environments, while indiscriminate

flower choice was often the most successful strategy. However, communication

improved the performance of flower constant colonies in nearly all tested environments.

This combination was particularly successful when high-quality food sources were

abundant and competition was weak.

4. Our findings help explain why social bees tend to be more flower constant than solitary

bees and suggest that flower constancy can be an adaptive strategy in social bees.

Simulations suggest that anthropogenic changes of foraging landscapes will have

different effects on the foraging performance of bees that vary in flower constancy.

Introduction

Most flowering plant species are animal pollinated and bees, in particular, are important

pollinators of wild and agricultural plants (Bawa, 1990; Klein et al., 2007; Ollerton et al., 2011).

Several biological features explain why bees are helpful agents of reproduction for plants,

including their abundance and their often broad (*i.e.* polylectic) diet in combination with a

tendency to specialise on a particular flower type during an individual foraging bout. The latter

behaviour, called flower constancy (Bateman, 1951; Chittka et al., 1999; Darwin, 1876; Grüter

& Ratnieks, 2011; Waser, 1986), reduces conspecific pollen loss and heterospecific pollen

deposition, both of which can reduce plant fitness (Ashman & Arceo-Gómez, 2013; Campbell

& Motten, 1985; Chittka et al., 1999; Morales & Traveset, 2008; Waser, 1986). Flower

constancy is also thought to enhance the coexistence of different plant species and, thus, shapes

plant community structure (Morales & Traveset, 2008; Song & Feldman, 2014).

From a pollinator perspective, however, the benefits of flower constancy are less

obvious. Ignoring potentially superior flower species appears to contradict optimal foraging

theory (King & Marshall, 2022; Latty & Trueblood, 2020; Waser, 1986; Wells & Wells, 1983,

1986). Why then are pollinators flower constant? The most widely accepted view is that flower

constancy is driven by cognitive limitations, which can include (*i*) slow learning to forage

efficiently on a new flower species, (*ii*) an inability to memorise more than one or a few flower

types, (*iii*) unstable short-term memories which are prone to being erased by competing

information or (*iv*) an inability to retrieve long-term memory about different flower species fast

enough to be an efficient generalist (Darwin, 1876; Heinrich, 1979; Lewis, 1986; Menzel, 1999;

Raine & Chittka, 2007; Waser, 1986; for reviews see Chittka et al., 1999; Grüter & Ratnieks,

2011). These cognitive limitations are likely to cause time delays as a bee tries to extract nectar

from a flower after switching from a different species and they may increase switching times

(Chittka et al., 1999; Goulson et al., 1997; Lewis, 1986; Raine & Chittka, 2007).

The "cognitive limitations hypothesis" as an explanation for flower constancy is not

without challenges. Given that efficient foraging is likely to be under strong natural selection

due to its effects on reproductive success (Heinrich, 1979), why does natural selection not lead

to the evolution of lower flower constancy in all bees? How can we explain the finding that

individual bees often show plasticity in their flower constancy, e.g. by being more flower

constant after finding good rewards (Chittka et al., 1997; Grüter et al., 2011; Wells & Rathore,

1994; but see Hill et al., 1997) or the distances between food sources are shorter (Gegear &

Thomson, 2004; Kunin, 1993; Mardern & Waddington, 1981)? Why do bee species vary in

their degree of flower constancy? Social bees, in particular, are often highly flower constant

(Cholis et al., 2020; Free, 1963; Heinrich, 1976, 1979; Hill et al., 1997; Kozuharova, 2018;

Pangestika et al., 2017; Rossi et al., 2015; Slaa et al., 2003; White et al., 2001; but see Martínez-

Bauer et al., 2021), while flower constancy seems to be less pronounced in solitary bees

(Bateman, 1951; Campbell & Motten, 1985; Eckhardt et al., 2014; Jakobsson et al., 2008; Pohl

et al., 2011; Smith et al., 2019; Waser, 1986; Williams & Tepedino, 2003). Smith et al. (2019),

for example, studied pollen composition of 56 bee species and found that individual social bees

showed a higher degree of specialisation during a foraging bout than solitary species. Different

ecological needs could explain this difference. Solitary bees need to collect all required

nutrients by themselves, potentially favouring a strategy of mixing resources during a foraging

trip even if this has energetic costs (e.g. Eckhardt et al., 2014; Williams & Tepedino, 2003). In

social species, on the other hand, different bees from the same colony can specialise on different

types of resources and flower species to cover their nutritional needs.

Foragers of many social bees share information about profitable food sources, and this could affect the value of flower constancy. Honeybees use the waggle dance to indicate the odour (type) and location of profitable food sources (von Frisch, 1967) and some stingless bees lay pheromone trails (Grüter, 2020; Jarau, 2009; Lindauer & Kerr, 1960; Nieh, 2004). Stingless bees and bumblebees inform nestmates about the availability and odour of a good food source by means of excitatory or jostling runs inside their nest (Dornhaus & Chittka, 2004; Hrncir, 2009). Trophallaxis – food transfer between bees – is performed by honey bees and stingless bees (Farina et al., 2012; Farina & Grüter, 2009; Hart & Ratnieks, 2002; Hrncir et al., 2006; Krausa et al., 2017; von Frisch, 1967) and is another behaviour that allows nestmates to learn the odour of available food sources (Aguilar et al., 2005; Farina et al., 2005; Lindauer & Kerr, 1960; von Frisch, 1967). These different communication behaviours share two common features. First, they depend on food quality. Dances, pheromone trails, jostling runs and trophallaxes occur at higher frequencies if the exploited food sources are of higher quality (Farina et al., 2012; Hrncir, 2009; Krausa et al., 2017; Lindauer & Kerr, 1960; von Frisch, 1967). Second, during these social interactions, nestmates can learn the odour of the exploited flower species and acquire a preference for this flower species in the field.

Heinrich (1976) was probably the first to propose a link between recruitment communication and flower constancy. Since recruiting bees share information selectively about high-quality food sources, recruits can discover profitable flower types without the costs of sampling different, lower-quality plant species. This does not require that foragers are able to direct nestmates to a specific location, as in honeybees and some stingless bees, but depends more generally on foragers biasing the food search towards flower types that are more profitable. Among the social bees, bumble bees seem to be less flower constant that honey bees (Bateman, 1951; Grant, 1950; Martínez-Bauer et al., 2021; Smith et al., 2019), possibly because their communication system is less sophisticated than that of honey bees (Heinrich, 1976).

Experimental studies of the benefits of flower constancy and how they depend on social

and ecological traits are logistically challenging for several reasons. For example, it is often not

possible to manipulate the degree of flower constancy while keeping other factors constant.

Agent-based simulation models can be a useful complementary tool to evaluate how biological

and ecological factors affect the benefits of a behavioural strategy. We developed an agent-

based simulation model to test the hypothesis that flower constancy is more beneficial in bees

that communicate about profitable food sources than in bees without communication. Colonies

consisting of virtual bees (agents) were either flower constant or they chose food sources

randomly (indiscriminately) in environments that varied in the number and quality of food

sources. Some studies have found that bees adjust the degree of flower constancy depending on

the foraging conditions, being more flower constant if the rewards on offer are better (Chittka

et al., 1997; Grüter et al., 2011; Wells & Rathore, 1994; but see Hill et al., 1997) and the

distances between food sources (or density) are shorter (Gegear & Thomson, 2004; Kunin,

1993; Marden & Waddington, 1981). We, therefore, expected flower constancy to be more

beneficial in environments with more flowers and larger reward sizes. Exploitation competition,

on the other hand, is expected to favour a indiscriminate choice because greater competition

increases the costs of rejecting a reward (Pulliam, 1974).

The model

We built an agent-based model (ABM) using the programming software NetLogo 6.1

(Wilensky, 1999) (see supplementary material for NetLogo file). It is an extension and further

analysis of the model developed in Grüter & Hayes (in preparation), which analysed foraging

distances. The model simulates an environment with a colony surrounded by food sources. The

agents ("bees") operate on a two-dimensional square grid with 400 x 400 patches. A single

patch length corresponds to 5 meters and 1 tick corresponds to 1 second. Thus, the size of the

virtual world corresponds to 2 x 2 km. The nest of the colony is positioned in the centre of the

grid (x=0, y=0). In the default situation, environments contained two different flower types that

differed in the rewards they offered.

The model allows simulating a wide range of parameter values, but for the purpose of

this study we based our default parameters, such as the nest stay time (t_{nest}), flight speed (v_{flight}),

metabolic costs of flying (M_{cost}), and crop capacity (*Crop*) on the Western honeybee *Apis*

mellifera because we have accurate information about these relevant biological parameters in

Apis mellifera. Other values were tested (see Table 1 and section *Sensitivity analysis and model*

exploration).

Foragers in social bees use different behavioural mechanisms to transmit social

information and, thereby, influence the food source preferences of their nestmates (see

introduction). The model does not simulate a particular behaviour, but a generic process that

biases the food preferences of nestmates, which could correspond to jostling runs, trophallaxis

or the waggle dance.

Entities and state variables

<u>Bees</u>

The default colony size was 100 agents (forager bees), which corresponds to the size of the

forager pool in many species of bumble bees (Westphal et al., 2006) and stingless bees (Grüter,

2020). Agents could assume any of the following states: (1) *generalists*, (2) *feeding forager*,

(3) *searching forager*, (4) *returning forager*, (5) *inside-nest-worker* and (6) *influencer* (see Fig.

1).

Fig. 1. State diagram showing the different states of the agents and the possible transitions between states. Here, yellow flowers were arbitrarily chosen to represent a lower-quality food source, therefore, the default probability that foragers visiting yellow flowers would become influencers after their return to the nest was 0.

Agents begin the simulation in the centre of the nest with energy = 0 as *generalists*. They then move at a flying speed of 1.4 patch/tick (v_{flight}), corresponding to a flight speed of *Apis mellifera* (7 m/sec, von Frisch 1967). Their random search behaviour follows a Lévy-flight pattern (with $\mu = 2.4$ as default) (Reynolds, 2009; Reynolds et al., 2007). A Lévy-flight consists of a random sequence of flight segments whose lengths, l, come from a probability distribution function having a power-law tail, $P(l) \sim l^{-\mu}$, with $1 < \mu < 3$ (Reynolds et al., 2007). The speed of agents moving inside the nest (v_{nest}) was arbitrarily chosen to be 0.1 (patch/tick). Flying has a

metabolic cost (M_{cost}) of 0.032 Joule (J) per tick in the default condition (Heinrich, 1975;

Willmer, 2011). Once an agent encounters a food source, they remain on the food source for 60

ticks ($t_{flower-stay}$) under default conditions (*feeding foragers*), irrespective of whether they were

choosing indiscriminately or are flower constant. Thus, we assume that the time spent handling

a flower or flowers in a patch and extracting the reward is the same for flower constant and

indiscriminate foragers. This was chosen as the default condition to explore whether flower

constancy can be an adaptive strategy in the absence of cognitive constraints.

The agent then continues to forage (*searching foragers*) until its crop is full, after which

it returns to the nest (*returning foragers*) to unload its energy and stay in the nest for 300 ticks

($t_{nest-stay}$) (Farina, 2000; Seeley, 1986; von Frisch, 1967). In the default condition, only foragers

visiting the high-quality food source could become *influencers* (*i.e.* bees that bias the food

choice of other bees) upon return to the nest. *Influencers* target *inside-nest-workers* that are not

yet flower-constant to the high-quality food type by changing the latter's preference if they

encountered each other on same patch inside the nest. Following such an encounter, *inside-*

nest-workers become flower constancy for the high-quality type.

Since recruitment behaviours often depend on the food source distance (with greater

foraging distances lowering the probability of recruitment), we simulated recruitment curves

where the probability of becoming an *influencer* decreased with increasing distance of the last

visited food patch (Fig. S1).

Food sources

In the default condition, two different types of food sources can be found in the environment,

mimicking the typical situation in experimental flower constancy studies (e.g. Chittka et al.,

1997; Goulson & Wright, 1998; Grüter et al., 2011; Ishii & Masuda, 2014; Wells & Wells,

1983). The food source types differ in the rewards they offer per visit. Natural bee-visited

flowers offer between 0.1 and 10 µL of nectar per flower (Willmer, 2011, p. 203). For the

default condition, we chose 5μL (29.07 J) for the high-quality type and 2.5μL (14.535 J) for the

low-quality type. This reward could represent an individual flower that offers a large reward or

a small patch of several flowers, each offering smaller quantities, or it could represent a larger

patch of flowers that is shared by several bees.

We tested different refill times (t_{refill}) for food sources: 0, 1200 and 3600 ticks (Stout &

Goulson, 2002). When $t_{refill}= 0$, food sources became rewarding again immediately after the

visit of a bee. This simulates conditions under which bees have a high probability of finding a

reward after landing on food source, which might occasionally occur at food patches. With

$t_{refill}= 3600$, a food source (flower or patch) was empty for the equivalent of an hour after it had

been visited by a bee, leading to intense exploitation competition among bees. The number of

food sources per type in the simulated environment varied between 1500 (low abundance) and

4500 (high abundance). Default conditions simulated even numbers of food sources for both

food source types, but we also explored uneven food source abundances (Table 1). We

measured the average foraging distance of bees during a simulation run to confirm that the

simulated conditions led to naturally realistic average foraging distances for many social bees

(271 ± 130 m; range 63-581 m; N = 1800 simulations in default conditions) (Kohl et al., 2020;

Van Nieuwstadt & Iraheta, 1996; Walther-Hellwig & Frankl, 2000).

The energy collected by agents with a full crop was estimated in the following way:

Apis mellifera can carry up to ~70 μL of nectar in their crop, but they usually carry less (I'Anson

Price et al., 2019; Núñez, 1966). The crop load has been shown to depend on the quality of the

visited food source, with lower quality food sources leading to smaller crop loads (Núñez, 1966,

1970). Agents visiting the low-quality flower type foraged until their crop contained 25μL,

whereas agents visiting the high-quality food type collected 50μL per foraging trip. Generalist

bees that choose indiscriminately have an intermediate crop load, reflecting the relative number

of high- and low-quality food sources in the environment. For example, in an environment with

an even number of high- and low-quality food sources, they collect 37.5μL per foraging trip.

Sugar concentration of collected nectar varies considerably from c. 10-70% (I'Anson Price,

2018; Seeley, 1986; Willmer, 2011). We chose an average sugar concentration of 35%,

providing 5.814 J/μL.

Each simulation lasted 36,000 ticks (i.e. 10 hours), simulating a day with good foraging

conditions. We measured the total energy collected by a colony during this period divided by

the number of agents (Energy/bee). Our main questions were if the energy/bee depended on

flower constancy (vs. indiscriminate choice), communication (vs. no communication), refill

time, the number of food sources and reward size. We also tested situations when flower

constancy was lower after visiting a low-quality food source ($Constancy_{LQ}$) (Grüter et al.,

2011), when there were 4 food source types and when indiscriminate flower choice increased

the time to extract a reward from a food source (*i.e.* to simulate cognitive constraints) (Chittka

et al., 1999).

Sensitivity analysis and model exploration

We varied a range of other factors to explore how they affected our results. These included

colony size, crop load size, flower stay time, metabolic costs, nest stay time, Lévy flight μ,

selectivity of communication (*i.e.* bees foraging on low-quality food source become *influencers*

with the same probability as those foraging on the high-quality type) and the shape of the

recruitment curve (see Fig. S1).

We performed 30 runs per parameter combination. We do not provide *p*-values due to

the arbitrariness of the simulation number but indicate 95%-confidence intervals to facilitate

interpretation of effect sizes.

Table 1: Overview of the model variables and the used values.

Variables	Description	Default values	Other values tested	Information source

Colony size	Number of foragers in a colony	100	5-300	Westphal et al. 2006; Grüter 2020
FS_{number}	Number of food sources per type	3000	1500, 4500	arbitrary
FS_{size}	Size of food sources	1 patch		arbitrary
FS types	Number of food source types	2	4	arbitrary
$Reward_{HQ}$	Nectar amount per food source	5μl	2.5μl, 10μl	Willmer 2011
$Reward_{LQ}$	Nectar amount per food source	2.5μl	1.25μl, 5μl	Willmer 2011
v_{flight}	Flight speed	1.4 patch/tick		von Frisch 1967
M_{cost}	Metabolic costs of flight, J/tick	0.032	0.016, 0.064	Heinrich 1975; Willmer 2011
$t_{flower-stay}$	Time spent at food source	60 ticks	20, 180	arbitrary
v_{nest}	Movement speed inside nest	0.1 patch/tick		arbitrary
$t_{nest-stay}$	Time in nest between trips	300 ticks	150, 450	Farina 2000
$Crop_{HQ}$	Crop load when flower constant to HQ food sources	50μl	25 μl, 100 μl	Núñez 1966
$Crop_{LQ}$	Crop load when flower constant to LQ food sources	25μl	12.5 μl, 50μl	Núñez 1966
$Crop_{Random}$	Crop load without flower constancy	37.5μl	50μl	Núñez 1966
t_{refill}	Time until food sources offer food again	0 ticks	1200, 3600	Stout & Goulson, 2002
Lévy μ	Lévy flight parameter	2.4	1.8, 3.0	Reynolds et al. 2009
$Constancy_{HQ}$	Probability to remain constant to a type after leaving a high-quality food source during flower constant simulations	100%		Grüter et al. 2011
$Constancy_{LQ}$	Probability to remain constant to a type after leaving a low-quality food source during flower constant simulations	100%	90%, 95%	Grüter et al. 2011

Results

Food source abundance and refill speed

We found that communication about the high-quality flower type did not affect the collected

energy if bees chose food sources indiscriminately (Fig. 2) (the raw data from which the figures

were built can be found in the supplementary material). However, if colonies were flower

constant, communication increased the energy collected by bees in all situations when the two

246   flower types were equally abundant (Fig. 2, see also Figs. 4-8), showing an interaction between

247   flower constancy and communication. The combination of communication and flower

248   constancy was relatively more beneficial when high-quality food sources were easy to find,

249   either because they were highly abundant (Fig. 2c,f,i) or because visited food sources

250   replenished quickly (Fig. 2a,b,c).

Fig. 2. Energy collected per bee (Joule) under varying food abundances (1500, 3000 and 4500 food
sources per type) and refill times (0, 1200 and 3600 ticks). Colonies either showed flower constancy
(constant) or they chose food source indiscriminately (random). Plots show the mean and the 95%-
confidence based on 30 simulations (grey dots). Numbers show % of change compared to random
choice without communication.

In the most favourable conditions, flower constancy in combination with communication was

the most successful combination (Fig. 2b, c). In all other conditions, indiscriminate choice was

the most successful strategy.

The relative abundance of the two food source types also played an important role.

Flower constancy combined with communication was relatively more successful when high-

quality food sources were more common than the low-quality flower type compared to when

they were rarer than the lower-quality flower type (Fig. 3). When high-quality food sources

represented the common flower type, colonies with flower constancy and communication were

either more successful (Fig. 3b) or not much less successful than colonies with indiscriminate

choice (Fig. 3d).

Fig. 3. Energy collected per bee (Joule) when high-quality food sources were either rare (1500) or

common (4500) compared to the low-quality food sources (3000). Default values were used for the

other parameters (Table 1).

However, indiscriminate choice was considerably more successful when high-quality food

sources were in the minority (Fig. 3a,c). When high-quality food sources were particularly

difficult to find, communication lowered the foraging success of flower constant colonies (Fig.

3c). Under these circumstances, communication directs the foragers of a colony towards a rare

food source, leading to long search times.

Reward sizes

Reward quantities are known to affect flower constancy, with bees becoming more flower

constant with increasing reward quantities (Chittka et al., 1997; Grüter et al., 2011; Wells &

Rathore, 1994). In accordance with this observation, we found that flower constancy became

relatively more successful (energy/bee) as reward sizes of both high- and low-quality food

sources increased (Fig. 4). However, indiscriminate choice was the most successful strategy in

many tested environments.

Fig. 4. Energy collected per bee (Joule) when rewards were (a) smaller (2.5μl and 1.25μl) or (c) larger
(10μl and 5μl) than in the (b) default situation (5μl and 2.5μl). Medium food source abundance was
simulated; default values were used for the other parameters (Table 1). Blue shows means and 95%-
CI for colonies using communication, red shows data for colonies choosing indiscriminately.

Time needed to collect a reward

The time bees need to extract a reward from a flower will affect the time costs of foraging

decisions and, if the refill time is >0, it will affect the number of depleted food sources in the

environment. Under default conditions, bees needed 60 ticks (1 minute) to obtain the reward

from a flower/food patch. We explored how different values for $t_{Flower-stay}$ affected the benefits

of flower constancy and communication. Increasing the time needed to obtain a reward

increased the relative benefits of combined flower constancy and communication compared to

short reward collection times (Fig. 5).

Fig. 5. Energy collected per bee (Joule) when the time required to obtain a reward from a food source
was (a,d) shorter (20 ticks) or (c,f) longer (180 ticks) than in the (b,e) default situation (60 ticks).
Default values and a medium food source abundance were simulated (Table 1).

Quality dependent flower constancy

Under default conditions, flower constancy did not depend on the quality of the food source

("spontaneous flower constancy", Hill et al., 1997). We simulated situations when bees visiting

a low-quality food source were slightly less flower constant (they had a 90% or a 95% chance

to remain flower constant on the subsequent visit, as in Grüter et al., 2011). Our results show

that this quality-dependent flower constancy considerably improves the energy collected by

colonies following this strategy of quality-dependent flower constancy (Fig. 6).

Fig. 6. Energy collected per bee (Joule) when bees foraged indiscriminately, when they showed
reduced flower constancy after visiting a low-quality food source (90% LQ constant or 95% LQ
constant) and when they were strictly flower constant. Refill time was 0, default values were used for
the other parameters (Table 1).

Exploring environments with 4 food types

When environments provide four different types of food sources rather than two, flower
constancy is less favourable overall (Fig. 7). In other words, indiscriminate flower choice is
highly beneficial in an environment where flower constancy would limit the options a forager
has to a small subset (25% of all food sources) of all available food sources than with two food
source types (Fig. 7).

We tested situations where one, two or three of the four plant types were of high quality,
while the remaining food sources were of low quality. While it was always beneficial to use
communication when colonies were also flower constant, the relative benefits of
communication diminished as the number of high-quality food types and the refilling time
increased. Unsurprisingly, therefore, foraging in an environment that consists mainly of high-
quality food sources belonging to different plant species somewhat diminishes the value of
using communication to direct foragers towards higher-quality food sources.

Fig. 7. Energy collected per bee (Joule) when bees foraged in an environment of four flower species. In
(a) & (d), one of the four plant types was of high quality, while the remaining three types were of low
quality. In (b) & (e), two of four types were of high quality and in (c) & (f), three of flower types were of
high quality. Food sources were refilling either at a fast or a medium rate.

Time penalty for non-specialists

So far, we have assumed that there are no additional time costs (e.g. as a result of cognitive

limitations) for bees that do not specialise on a particular type of food source. To explore the

consequences of cognitive limitations, we simulated situations when indiscriminate bees

require more time to extract a reward from a food source compared to flower constant bees. A

time penalty for indiscriminate bees favours flower constant colonies, especially those that also

communicate the high-quality flower type to nestmates (Fig. 8).

Fig. 8. Energy collected per bee (Joule) when non-specialists needed 50% (b, e) or 200% (c, f) more time than flower constant bees to obtain a reward from a food source. Food sources were refilling either at a fast (t_{refill} =0) or a medium rate (t_{refill} = 1200). Default values were used for the other parameters (Table 1).

Sensitivity analysis and model exploration

Varying colony size from 5 to 300 (Fig. S2) does not greatly affect the general pattern observed for the default colony size of 100 (see Fig. 2). When food sources refill at a fast rate (Fig. S2a), bees do not experience exploitation competition and colony size does not affect the energy collected by individual bees. Increasing the refill time while also increasing the number of agents searching for food, on the other hand, increases exploitation competition and, therefore, lowers the energy collected by individual bees (Fig. S2b,c).

Using different recruitment curves (Fig. S1) had no noticeable effect on the energy collected by bees, but non-selective recruitment (recruitment to both high- and low-quality food sources) lowers the collected energy to levels similar to those of flower constant colonies without communication (Fig. S3). Changing the metabolic costs of flying or the Lévy-flight μ has little effect on the overall pattern (Fig. S4, S5), whereas increasing the time spent inside the nest in-between foraging trips reduces the energy collected by bees, but less so in colonies with

flower constancy (Fig. S6). Thus, longer nest stay times favour flower constancy. Flower

constancy was also favoured when bees had smaller crop sizes (Fig. S7).

Discussion

Results from our simulations suggest that flower constancy without communication is less

successful than indiscriminate choice under all tested conditions. Flower constancy imposes

significant costs because it (*i*) limits the available options to a subset of all available flowers,

thereby increasing time and energy costs during foraging, and (*ii*) causes many foragers to

specialise on a sub-optimal flower type. Communication about the high-quality flower type

positively interacted with flower constancy (Fig. 2) and considerably improved the foraging

success of flower constant colonies by. Communication allows a colony to focus on high-

quality flowers, thereby reducing the second type of cost (*ii*). Many species of social bees have

evolved mechanisms of reward-quality dependent recruitment communication, which allow

influencers to affect the foraging decisions of their nestmates towards a particular flower type,

mainly via olfactory learning (Dornhaus & Chittka, 1999; Farina et al., 2012; Jarau & Hrncir,

2009; Lindauer & Kerr, 1960; von Frisch, 1967). This, in turn, lowers the benefits of sampling

alternative flower species and highlights the importance of social information use as a process

of information-filtering (Grüter et al., 2010; Rendell et al., 2010). Our findings can help explain

why social bees tend to be more flower constant than solitary bees (e.g. Smith et al., 2019;

Waser, 1986).

The general foraging conditions had a strong effect on the value of flower constancy

and the strength of its interaction with communication. Flower constancy in combination with

communication was the most successful strategy when foraging conditions were very

favourable, while indiscriminate choice was the better strategy when foraging options were

more limited. For instance, flower constancy in combination with communication was

beneficial when foragers did not encounter empty food sources (refill time of 0) and food

sources were abundant (Fig. 2b,c), when most food sources were of high-quality (Fig. 3b) and

when rewards were large (Fig. 4c). These findings are consistent with empirical studies showing

that bees are more flower constant when flower density is higher (Chittka et al., 1997; Kunin,

1993; Marden & Waddington, 1981) and rewards are larger (Chittka et al., 1997; Greggers &

Menzel, 1993; Grüter et al., 2011). Similarly, predator-prey models show that the abundance of

a prey item has a positive effect on diet specialisation of the predator (Pulliam, 1974). If food

sources took time to replenish, resulting in many empty food sources due to exploitation

competition, indiscriminate choice was more successful (Fig. 2d-i), suggesting that rejecting

flowers due to flower constancy is more costly in environments that offer fewer options.

Changes in the temporal dynamics of foraging trips affected the performance of the

different strategy by changing the relative costs of ignoring flowers (*i*) and choosing suboptimal

food sources (*ii*). Flower constancy in combination with communication performed relatively

better if bees required more time to extract a reward from a food source (Fig. 5). One

explanation for this is that visiting low-quality food sources, which is common with

indiscriminate choice, becomes relatively more costly as the time costs of a flower visit

increase. Thus, longer flower handling times, e.g. due to a complex flower morphology, favour

flower constancy from both an adaptive and a constraints-based perspective (see Chittka et al.,

1999 for arguments based on cognitive constraints). Flower constancy in combination with

communication performed relatively better when bees stayed in their nest longer (Fig. S6) and

had smaller crops (Fig. S7). These findings are somewhat puzzling, but one explanation could

be that longer nest stay times provide *influencers* with more opportunities to communicate their

findings to other bees. When food sources need time to replenish, longer nest stay times will

reduce the number of depleted food sources a bee encounters, which favours flower constancy

in combination with communication (Fig. 2b,c). Similarly, when bees have smaller crop loads,

they visit fewer food sources per trip and spend a larger proportion of their time in the nest,

reducing exploitation competition and the number of depleted food sources. Crop size will

depend on body size and one might, therefore, predict that smaller bees are more flower

constant, which is consistent with comparative data (Smith et al. 2019). However, it is unlikely

that there is straight forward relationship between crop size, body size and flower constancy in

nature because body size covaries with numerous other extrinsic and intrinsic factors, including

foraging conditions, metabolic costs, flying speed or sensory acuity (Gervais et al., 2020; Grab

et al., 2019; Spaethe et al., 2007), all of which might affect flower constancy.

The Western honey bee *Apis mellifera* is strongly flower constant, but there is

disagreement about whether and when flower constancy depends on the profitability of visited

flowers. Some studies have suggested that flower constancy is often "spontaneous", *i.e.*

unrelated to reward size (Wells & Wells 1983; Hill et al. 1996, 2001; Sanderson et al. 2006),

whereas others have found that honey bees adjust flower constancy according to the

profitability of rewards (Greggers & Menzel 1993; Chittka et al. 1997; bumble bees: Heinrich

1976, 1979b; reviewed in Grüter & Ratnieks 2011). Our simulations show that context-

dependent flower constancy is more successful than strict ("spontaneous") flower constancy

(Fig. 6). When bees visiting the less profitable food type were only 90-95% flower constant,

colonies collected about 25% more energy than colonies with strict flower constancy. As is the

case with communication, context-dependent flower constancy allows bees to switch from the

low-quality to the high-quality flower species over time (type (*ii*) costs).

Human impacts have significantly affected the diversity of plant species found in some

environments, especially in intensively farmed habitats (e.g. Potts et al., 2010; Tew et al., 2021),

which is likely to affect the costs and benefits of flower constancy. In our simulations, flower

constancy performed considerably worse when there were four rather than two flower types

(Fig. 7). With more plant species present, flower constant bees will ignore most of the available

options and focus on a small subset of all food sources, thereby dramatically increasing

opportunity costs (type (*i*) costs). Thus, bees should be less flower constant in more diverse

foraging environments. Flower constant bees, in turn, might suffer a reduction in foraging

success in more biodiverse habitats. These findings challenge the reasoning behind the "costly

information hypothesis", which argues that flower constancy is an adaptive foraging strategy

because acquiring information about suitable alternatives would cost too much time and energy

if there are several plant species available (Chittka et al., 1999; Grüter & Ratnieks, 2011). In

flower diverse environments, bees should accept even low-quality food source if it means they

can cut time and energy costs imposed by flower constancy. Empirical studies on the links

between floral diversity and flower constancy provide contrasting results. While Gervais et al.

(2020) and Martínez-Bauer et al. (2021) found that increasing plant diversity was associated

with lower flower constancy in *Bombus impatiens* and *B. terrestris*, Austin et al. (2019) found

that bumble bees became more flower constant when there are more options available. The

latter finding is more consistent with a "cognitive limitations" perspective, since deciding

among more options would be cognitively more challenging and flower constancy, therefore, a

possible solution to avoid switching costs (see also Chittka et al. 1997; Gegear & Thomson

2004). Decision making is often impaired as the number of choices increases (Latty &

Trueblood, 2020). The different studies differ in that the first two were performed under natural

conditions, whereas Austin et al. (2019) was experimental. Non-experimental surveys can be

confounded by numerous factors, such as differences in rewards, clustering of flowers or

management, whereas experimental studies might fail to capture crucial features of natural

environments that affect decision-making (Fawcett et al., 2014).

Agent-based models have important limitations. Simulation outcomes depend on the

underlying assumptions and the parameters chosen when building the model, some of which

are arbitrary or simplistic. As a result, ABMs potentially miss important natural features that

shape decision-making (Fawcett et al., 2014). For example, we assumed that food sources are

randomly distributed, whereas natural foraging environments are often spatially heterogenous

and patchy, which is likely to affect the value of flower constancy. Patchiness can lead to flower

constancy "by accident" if bees forage in large patches, even if they choose flowers

indiscriminately. We might, therefore, expect increasing patchiness to lead to more similar

outcomes for flower constant and indiscriminate foragers. Pulliam's (1974) predator-prey

model found that an increasingly clumped prey distribution favours a more specialised diet in

predators and we might expect a similar finding in plant pollinator interactions. Agent-based

models also have important strengths because they allow us to systematically vary factors that

cannot be manipulated experimentally, such as tuning flower constancy or recruitment

communication while keeping all other factors constant. ABMs should be seen as a useful tool

to complement empirical studies.

One of the aims of our model was to test whether flower constancy could be an adaptive

strategy *per se* under some foraging conditions, *i.e.* in the absence of cognitive constraints. If,

however, switching between flower species leads to increased time costs or reduced reward

sizes (Chittka et al., 1999; Darwin, 1876; Grüter & Ratnieks, 2011; Lewis, 1986; Raine &

Chittka, 2007), flower constancy, with or without communication, becomes a much more

beneficial strategy under a wide range of conditions (Fig. 8c,f). The reasons for flower

constancy in pollinators are likely to be complex and depend on both constraints and adaptive

processes, to varying degrees in different species. However, our results suggest that a more

pronounced flower constancy in social bees is more likely due to increased performance of

flower constancy in social species due to social traits, rather than the result of poorer cognitive

abilities in social bees compared to solitary bees (Amaya-Márquez & Wells, 2008; Dukas &

Real, 1991).

Acknowledgements

We thank the University of Bristol Ecology group for comments and feedback on the results of

this study.

References

References

Aguilar, I., Fonseca, A., & Biesmeijer, J. C. (2005). Recruitment and communication of food source

location in three species of stingless bees (Hymenoptera, Apidae, Meliponini). *Apidologie*, *36*,

313–324.

Amaya-Márquez, M., & Wells, H. (2008). Social complexity and learning foraging tasks in bees.

Caldasia, *30*(2), 469–477.

Ashman, T.-L., & Arceo-Gómez, G. (2013). Toward a predictive understanding of the fitness costs of

heterospecific pollen receipt and its importance in co-flowering communities. *American*

Journal of Botany, *100*(6), 1061–1070.

Austin, M. W., Horack, P., & Dunlap, A. S. (2019). Choice in a floral marketplace: The role of

complexity in bumble bee decision-making. *Behavioral Ecology*, *30*(2), 500–508.

Bateman, A. J. (1951). The taxonomic discrimination of bees. *Heredity*, *5*(2), 271–278.

Bawa, K. S. (1990). Plant-pollinator interactions in Tropical rain forests. *Annual Review of Ecology*

and Systematics, *21*, 399–422.

Campbell, D. R., & Motten, A. F. (1985). The Mechanism of Competition for Pollination between

Two Forest Herbs. *Ecology*, *66*(2), 554–563.

Chittka, L., Gumbert, A., & Kunze, J. (1997). Foraging dynamics of bumble bees: Correlates of

movements within and between plant species. *Behavioral Ecology*, *8*, 239–249.

Chittka, L., Thomson, J. D., & Waser, N. M. (1999). Flower constancy, insect psychology, and plant

evolution. *Naturwissenschaften*, *86*, 361–377.

Cholis, M. N., Alpionita, R., Prawasti, T. S., & Atmowidi, T. (2020). *Pollen Load and Flower*

Constancy of Stingless Bees Tetragonula laeviceps (Smith) and Heterotrigona itama

(Cockerell) (Apidae: Meliponinae). *8*, 285–289.

Darwin, C. (1876). *Cross and self fertilization in the vegetable kingdom*. Murray.

Dornhaus, A., & Chittka, L. (1999). Evolutionary origins of bee dances. *Nature*, *401*(6748), 38–38.

Dornhaus, A., & Chittka, L. (2004). Information flow and regulation of foraging activity in bumble

bees (Bombus spp.). *Apidologie*, *35*(2), 183–192.

Dukas, R., & Real, L. A. (1991). Learning foraging tasks by bees: A comparison between social and

solitary species. *Animal Behaviour*, *42*(2), 269–276.

Eckhardt, M., Haider, M., Dorn, S., & Müller, A. (2014). Pollen mixing in pollen generalist solitary

bees: A possible strategy to complement or mitigate unfavourable pollen properties? *Journal*

of Animal Ecology, *83*(3), 588–597.

Farina, W. M. (2000). The interplay between dancing and trophallactic behavior in the honey bee *Apis*

mellifera. Journal of Comparative Physiology A-Neuroethology Sensory Neural and

Behavioral Physiology, *186*(3), 239–245.

Farina, W. M., & Grüter, C. (2009). Trophallaxis—A mechanism of information transfer. In S. Jarau

& M. Hrncir (Eds.), *Food exploitation by social insects: Ecological, behavioral, and*

theoretical approaches (pp. 173–187). CRC Press.

Farina, W. M., Grüter, C., & Arenas, A. (2012). Olfactory information transfer during recruitment in

honey bees. In C. G. Galizia, D. Eisenhardt, & M. Giurfa (Eds.), *Honeybee Neurobiology and*

Behavior—A Tribute to Randolf Menzel (pp. 89–101). Springer.

Farina, W. M., Grüter, C., & Diaz, P. C. (2005). Social learning of floral odours within the honeybee

hive. *Proceedings of the Royal Society of London Series B-Biological Sciences*, *272*, 1923–

1928.

Fawcett, T. W., Fallenstein, B., Higginson, A. D., Houston, A. I., Mallpress, D. E. W., Trimmer, P. C.,

& McNamara, J. M. (2014). The evolution of decision rules in complex environments. *Trends*

in Cognitive Sciences, *18*(3), 153–161.

Free, J. B. (1963). The flower constancy of honeybees. *Journal of Animal Ecology*, *32*, 119–131.

Gegear, R. J., & Thomson, J. D. (2004). Does the Flower Constancy of Bumble Bees Reflect Foraging

Economics? *Ethology*, *110*(10), 793–805.

Gervais, A., Courtois, È., Fournier, V., & Bélisle, M. (2020). Landscape composition and local floral

resources influence foraging behavior but not the size of Bombus impatiens Cresson

(Hymenoptera: Apidae) workers. *PLOS ONE*, *15*(6), e0234498.

Goulson, D., Stout, J., & Hawson, S. A. (1997). Can flower constancy in nectar butterflies be

explained by Darwin's interference hypothesis? *Oecologia*, *112*, 225–231.

Goulson, D., & Wright, N. P. (1998). Flower constancy in the hoverfies *Episyrphus balteatus* (Degeer)

and *Syrphus ribesii* (L.) (Syrphidae). *Behavioral Ecology*, *9*, 213–219.

Grab, H., Brokaw, J., Anderson, E., Gedlinske, L., Gibbs, J., Wilson, J., Loeb, G., Isaacs, R., &

Poveda, K. (2019). Habitat enhancements rescue bee body size from the negative effects of

landscape simplification. *Journal of Applied Ecology*, *56*, 2144–2154.

Grant, V. (1950). The flower constancy of bees. *The Botanical Review*, *16*, 379–398.

Greggers, U., & Menzel, R. (1993). Memory dynamics and foraging strategies of honeybees.

Behavioral Ecology and Sociobiology, *32*(1), 17–29.

Grüter, C. (2020). *Stingless Bees: Their Behaviour, Ecology and Evolution*. Springer International

Publishing.

Grüter, C., Leadbeater, E., & Ratnieks, F. L. W. (2010). Social learning: The importance of copying

others. *Current Biology*, *20*, R683–R685.

Grüter, C., Moore, H., Firmin, N., Helanterä, H., & Ratnieks, F. L. W. (2011). Flower constancy in

honey bee foragers (*Apis mellifera*) depends on ecologically realistic rewards. *Journal of*

Experimental Biology, *214*, 1397–1402.

Grüter, C., & Ratnieks, F. L. W. (2011). Flower constancy in insect pollinators: Adaptive foraging

behaviour or cognitive limitation? *Communicative and Integrative Biology*, *4*, 633–636.

Hart, A. G., & Ratnieks, F. L. W. (2002). Task-partitioned nectar transfer in stingless bees: Work

organisation in a phylogenetic context. *Ecological Entomology*, *27*, 163–168.

Heinrich, B. (1975). Energetics of pollination. *Annual Review of Ecology and Systematics*, *6*, 139–170.

Heinrich, B. (1976). The Foraging Specializations of Individual Bumblebees. *Ecological Monographs*,

46(2), 105–128.

Heinrich, B. (1979). "Majoring" and "Minoring" by Foraging Bumblebees, Bombus Vagans: An

Experimental Analysis. *Ecology*, *60*(2), 246–255.

Hill, P. S. M., Wells, P. H., & Wells, H. (1997). Spontaneous flower constancy and learning in honey

bees as a function of colour. *Animal Behaviour*, *54*, 615–627.

Hrncir, M. (2009). Mobilizing the Foraging Force Mechanical Signals in Stingless Bee Recruitment.

In S. Jarau & M. Hrncir (Eds.), *Food Exploitation by Social Insects: Ecological, Behavioral,*

and Theoretical Approaches (Vol. 11). CRC Press, Taylor & Francis Group.

Hrncir, M., Schmidt, V. M., Schorkopf, D. L. P., Jarau, S., Zucchi, R., & Barth, F. G. (2006).

Vibrating the food receivers: A direct way of signal transmission in stingless bees (*Melipona*

seminigra). *Journal of Comparative Physiology A*, *192*(8), 879–887.

I'Anson Price, R. (2018). *The adaptive significance of communication and learning in bees* [PhD

Thesis]. University of Lausanne, Switzerland.

I'Anson Price, R., Dulex, N., Vial, N., Vincent, C., & Grüter, C. (2019). Honeybees forage more

successfully without the "dance language" in challenging environments. *Science Advances*, *5*,

eaat0450.

Ishii, H. S., & Masuda, H. (2014). Effect of flower visual angle on flower constancy: A test of the

search image hypothesis. *Behavioral Ecology*, *25*, 933–944.

Jakobsson, A., Padrón, B., & Traveset, A. (2008). Pollen transfer from invasive Carpobrotus spp. To

natives – A study of pollinator behaviour and reproduction success. *Biological Conservation*,

141(1), 136–145.

Jarau, S. (2009). Chemical Communication during Food Exploitation in Stingless Bees. In S. Jarau &

M. Hrncir (Eds.), *Food Exploitation by Social Insects: Ecological, Behavioral, and*

Theoretical Approaches (Vol. 12). CRC University Press.

Jarau, S., & Hrncir, M. (2009). *Food Exploitation by Social Insects: Ecological, Behavioral, and*

Theoretical Approaches (S. Jarau & M. Hrncir, Eds.). CRC Press, Taylor & Francis Group.

King, A. J., & Marshall, H. H. (2022). Optimal foraging. *Current Biology*, *32*(12), R680–R683.

Klein, A.-M., Vaissiere, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., &

Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops.

Proceedings of the Royal Society B: Biological Sciences, *274*(1608), 303–313.

Kohl, P. L., Thulasi, N., Rutschmann, B., George, E. A., Steffan-Dewenter, I., & Brockmann, A.

(2020). Adaptive evolution of honeybee dance dialects. *Proceedings of the Royal Society B:*

Biological Sciences, *287*(1922), 20200190.

Kozuharova, E. (2018). Flower Constancy of Bumblebees – The Case of (Fabaceae) Pollinators.

Journal of Apicultural Science, *62*(1), 135–140.

Krausa, K., Hager, F. A., & Kirchner, W. H. (2017). The effect of food profitability on foraging

behaviors and vibrational signals in the African stingless bee *Plebeina hildebrandti*. *Insectes*

Sociaux, *64*(4), 567–578.

Kunin, W. E. (1993). Sex and the single mustard: Population density and pollinator behavior effects

on seed-set. *Ecology*, *74*, 2145–2160.

Latty, T., & Trueblood, J. S. (2020). How do insects choose flowers? A review of multi-attribute

flower choice and decoy effects in flower-visiting insects. *Journal of Animal Ecology*, *89*(n/a),

2750–2762.

Lewis, A. C. (1986). Memory Constraints and Flower Choice in Pieris rapae. *Science*.

Lindauer, M., & Kerr, W. E. (1960). Communication between the workers of stingless bees. *Bee*

World, *41*, 29–71.

Marden, J. H., & Waddington, K. D. (1981). Floral choices by honey bees in relation to the relative

distances to flowers. *Physiological Entomology*, *6*, 431–435.

Martínez-Bauer, A. E., Chadwick, F. J., Westmoreland, A. J., & Lander, T. A. (2021). Novel pollen

analogue technique shows bumblebees display low floral constancy and prefer sites with high

floral diversity. *Landscape Ecology*.

Menzel, R. (1999). Memory dynamics in the honeybee. *Journal of Comparative Physiology A-*

Neuroethology Sensory Neural and Behavioral Physiology, *185*(4), 323–340.

Morales, C. L., & Traveset, A. (2008). Interspecific Pollen Transfer: Magnitude, Prevalence and

Consequences for Plant Fitness. *Critical Reviews in Plant Sciences*, *27*(4), 221–238.

Nieh, J. C. (2004). Recruitment communication in stingless bees (Hymenoptera, Apidae, Meliponini).

Apidologie, *35*(2), 159–182.

Núñez, J. A. (1966). Quantitative Beziehungen zwischen den Eigenschaften von Futterquellen und

dem Verhalten von Sammelbienen. *Zeitschrift Für Vergleichende Physiologie*, *53*, 142–164.

Núñez, J. A. (1970). The relationship between sugar flow and foraging and recruiting behaviour of

honey bees (*Apis mellifera* L.). *Animal Behaviour*, *18*, 527–538.

Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals?
Oikos, *120*(3), 321–326.

Pangestika, N. W., Atmowidi, T., & Kahono, S. (2017). Pollen Load and Flower Constancy of Three
Species of Stingless Bees (Hymenoptera, Apidae, Meliponinae). *Tropical Life Sciences*
Research, *28*(2), 179–187.

Pohl, N. B., Wyk, J. V., & Campbell, D. R. (2011). Butterflies show flower colour preferences but not
constancy in foraging at four plant species. *Ecological Entomology*, *36*(3), 290–300.

Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010).
Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology & Evolution*, *25*(6),
345–353.

Pulliam, H. R. (1974). On the Theory of Optimal Diets. *The American Naturalist*, *108*(959), 59–74.

Raine, N. E., & Chittka, L. (2007). Pollen foraging: Learning a complex motor skill by bumblebees
(Bombus terrestris). *Naturwissenschaften*, *94*, 459–464.

Rendell, L., Boyd, R., Cownden, D., Enquist, M., Eriksson, K., Feldman, W. M., Fogarty, L.,
Ghirlanda, S., Lillicrap, T., & Laland, K. N. (2010). Why copy others? Insights from the social
learning strategies tournament. *Science*, *328*, 208–213.

Reynolds, A. M. (2009). Lévy flight patterns are predicted to be an emergent property of a
bumblebees' foraging strategy. *Behavioral Ecology and Sociobiology*, *64*(1), 19.

Reynolds, A. M., Smith, A. D., Reynolds, D. R., Carreck, N. L., & Osborne, J. L. (2007). Honeybees
perform optimal scale-free searching flights when attempting to locate a food source. *Journal*
of Experimental Biology, *210*(21), 3763–3770.

Rossi, N., Santos, E., Salvarrey, S., Arbulo, N., & Invernizzi, C. (2015). Determination of Flower
Constancy in *Bombus atratus* Franklin and *Bombus bellicosus* Smith (Hymenoptera: Apidae)
through Palynological Analysis of Nectar and Corbicular Pollen Loads. *Neotropical*
Entomology, *44*(6), 546–552.

Seeley, T. D. (1986). Social foraging by honeybees—How colonies allocate foragers among patches
of flowers. *Behavioral Ecology and Sociobiology*, *19*(5), 343–354.

Slaa, E. J., Tack, A. J. M., & Sommeijer, M. J. (2003). The effect of intrinsic and extrinsic factors on

flower constancy in stingless bees. *Apidologie*, *34*, 457–468.

Smith, C., Weinman, L., Gibbs, J., & Winfree, R. (2019). Specialist foragers in forest bee

communities are small, social or emerge early. *Journal of Animal Ecology*, *88*(8), 1158–1167.

Song, Z., & Feldman, M. W. (2014). Adaptive foraging behaviour of individual pollinators and the

coexistence of co-flowering plants. *Proceedings of the Royal Society B: Biological Sciences*,

281(1776), 20132437.

Spaethe, J., Brockmann, A., Halbig, C., & Tautz, J. (2007). Size determines antennal sensitivity and

behavioral threshold to odors in bumblebee workers. *Naturwissenschaften*, *94*, 733–739.

Stout, J. C., & Goulson, D. (2002). The Influence of Nectar Secretion Rates on the Responses of

Bumblebees (Bombus spp.) to Previously Visited Flowers. *Behavioral Ecology and*

Sociobiology, *52*(3), 239–246.

Tew, N. E., Memmott, J., Vaughan, I. P., Bird, S., Stone, G. N., Potts, S. G., & Baldock, K. C. R.

(2021). Quantifying nectar production by flowering plants in urban and rural landscapes.

Journal of Ecology, *109*, 1747–1757.

Van Nieuwstadt, M. G. L., & Iraheta, C. R. (1996). Relation between size and foraging range in

stingless bees (Apidae, Meliponinae). *Apidologie*, *27*, 219–228.

von Frisch, K. (1967). *The dance language and orientation of bees*. Harvard University Press.

Walther-Hellwig, K., & Frankl, R. (2000). Foraging Distances of Bombus muscorum, Bombus

lapidarius, and Bombus terrestris (Hymenoptera, Apidae). *Journal of Insect Behavior*, *13*(2),

239–246.

Waser, N. M. (1986). Flower constancy: Definition, cause, and measurement. *American Naturalist*,

127, 593–603.

Wells, H., & Rathore, R. R. S. (1994). Foraging ecology of the Asian hive bee, *Apis cerana indica*,

within artificial flower patches. *Journal of Apicultural Research*, *33*, 219–230.

Wells, H., & Wells, P. H. (1983). Honey bee foraging ecology: Optimal diet, minimal uncertainty or

individual constancy? *Journal of Animal Ecology*, *52*, 829–836.

Wells, H., & Wells, P. H. (1986). Optimal diet, minimal uncertainty and individual constancy in the

foraging of honey bees, *Apis mellifera*. *Journal of Animal Ecology*, *55*, 881-891.

Westphal, C., Steffan-Dewenter, I., & Tscharntke, T. (2006). Bumblebees experience landscapes at

different spatial scales: Possible implications for coexistence. *Oecologia*, *149*(2), 289–300.

White, D., Cribb, B. W., & Heard, T. A. (2001). Flower constancy of the stingless bee *Trigona*

carbonaria Smith (Hymenoptera: Apidae: Meliponini). *Australian Journal of Entomology*,

40(1), 61–64.

Wilensky, U. (1999). *NetLogo. Http://ccl.northwestern.edu/netlogo/*. Northwestern University,

Evanston, IL.

Williams, N. M., & Tepedino, V. J. (2003). Consistent mixing of near and distant resources in

foraging bouts by the solitary mason bee Osmia lignaria. *Behavioral Ecology*, *14*(1), 141–149.

Willmer, P. (2011). *Pollination and Floral Ecology*. Princeton University Press.

715   **Fig. S1**

717   **Fig. S1**. Probability that a returning bee that visited the high-quality food type (red) becomes an

718   influencer inside the nest. Bees visiting the low-quality food type did not become influencers under

719   default conditions. Thick red line shows the default probability. The other two lines show other tested

720   probability curves. 1 patch ~ 5 m.

736 **Fig. S2**

738 **Fig. S2**. Energy collected per bee for different colony sizes and different refill speeds (fast refill, $t_{refill} = 0$;

739 medium refill, $t_{refill} = 1200$; slow refill, $t_{refill} = 3600$). Lines in (a) show best fit lines based on linear

740 regression. In (b) and (c), generalised additive models (GAM) were used to fit curves. Grey bands show

741 95% confidence intervals. Default values were used for other parameters.

Fig. S3

Fig. S3. Energy collected per bee for different types of communication, with or without flower constancy.

The different recruitment strengths (weak, default and strong) correspond to the three different curves

shown in S1. Unselective refers to a situation where bees communicate equally about high- and low-

quality food sources (assuming the default curve shown in Fig. S1). Means and 95%-confidence

intervals are shown.

775   **Fig. S4**

777   **Fig. S4**. Energy per bee (Joule) collected using different Lévy-flight μ values, which determines the

778   distribution of the segment lengths l that constitute a Lévy-flight, with $P(l) \sim l^{\mu}$. Food sources that refill

779   immediately were simulated. Default values were used for the other parameters (Table 1).

801 **Fig. S5**

803 **Fig. S5**. Energy per bee (Joule) collected using different flight energy costs (low = 0.016 J/tick; default

804 = 0.032 J/tick; high = 0.064 J/tick). Food sources refilled immediately. Default values were used for the

805 other parameters (Table 1).

825 **Fig. S6**

827 **Fig. S6**. Energy per bee (Joule) collected using different nest stay times. Food sources refilled

828 immediately. Default values were used for the other parameters (Table 1).

Fig. S7

Fig. S7. Energy per bee (Joule) collected with different crop loads. (a) Small crop loads mean that bees constant to high-quality food sources filled their crop to 25µl, whereas bees constant to low-quality food sources returned to their nest if their crop load was 12.5µl. Bees choosing food sources randomly had intermediate crop loads (18.75µl). (c) Bees with large crop loads foraged until they had 100µl (constant to high-quality food sources), 50µl (constant to low-quality food sources or 75µl (randomly choosing bees). Food sources refilled immediately. Default values were used for the other parameters (Table 1).