1

Emotional Vocalizations Alter Behaviors and

Neurochemical Release into the Amygdala

Zahra Ghasemahmad[1,2], Aaron Mrvelj[1], Rishitha Panditi[1], Bhavya Sharma[1], Karthic
Drishna Perumal[1], and Jeffrey J. Wenstrup[1,2,3]•

[1]Department of Anatomy and Neurobiology and Hearing Research Group, Northeast
Ohio Medical University, Rootstown, Ohio, USA

[2]School of Biomedical Sciences, Kent State University, Kent, Ohio, USA

[3]Brain Health Research Institute, Kent State University, Kent, Ohio, USA

* Correspondence:

Jeffrey J. Wenstrup

jjw@neomed.edu

Keywords: acetylcholine, basolateral amygdala, dopamine, estrus, mating, microdialysis,
mouse, restraint, vocalization playback

ABSTRACT

The basolateral amygdala (BLA), a brain center of emotional expression, contributes to

acoustic communication by first interpreting the meaning of social sounds in the context

of the listener's internal state, then organizing the appropriate behavioral responses.

We propose that modulatory neurochemicals such as acetylcholine (ACh) and dopamine

(DA) provide internal-state signals to the BLA while an animal listens to social

vocalizations. We tested this in a vocal playback experiment wherein we sampled fluids

within the BLA and observed behavioral responses of male and female mice while

presenting highly affective vocal sequences associated with either mating or restraint

behaviors. In male mice, playback of restraint vocalizations increased ACh release and

decreased DA release, while playback of mating sequences evoked the opposite

neurochemical release patterns. In non-estrus female mice, the ACh and DA release

patterns to mating playback were similar to males. Estrous females, however, showed

increased ACh, associated with vigilance, as well as increased DA, associated with

reward-seeking. Across these groups, increased ACh concentration was correlated with

an increase in defensive behavior. These neurochemical release patterns and several

behavioral responses depended on prior experience with the mating and restraint

behaviors. Our results support a model in which ACh and DA provide contextual

information to sound-analyzing BLA neurons that modulates their output to

downstream brain regions responsible for appropriate behavioral outcomes.

INTRODUCTION

In social interactions involving vocal communication signals, an animal receives and

analyzes acoustic information, compares it with previous experiences, identifies the

salience and valence of such information, and responds with appropriate behaviors.

These integrated functions depend on brain circuits that include the amygdala, a region

located within the temporal lobe that is recognized to play a role in orchestrating

responses to salient sensory stimuli (LeDoux, 2000, 2003; McGaugh, 2002; Sah et al.,

2003; Wenstrup et al., 2020). The amygdalar target of auditory inputs from the

thalamus and cortex is the basolateral amygdala (BLA) (J. E. LeDoux et al., 1991; X. F. Li

et al., 1996; Sacchetti et al., 1999; Keifer et al., 2015). Through reciprocal connections

with areas such as the hippocampus, the BLA compares received sensory information

with previous experiences (Pitkänen et al., 2000). By integrating this information with

other sensory inputs and inputs from other limbic areas, BLA neurons shape appropriate

behavioral responses (Namburi et al., 2015, 2016; Beyeler et al., 2016; Gründemann et

al., 2019) via projections to downstream targets such as the nucleus accumbens

(Ambroggi et al., 2008; Stuber et al., 2011) and central nucleus of the amygdala (Ciocchi

et al., 2010).

The BLA processes vocal and other acoustic information in a context-dependent manner

(Goosens et al., 2000; Sander et al., 2005; Wiethoff et al., 2009; Leitman et al., 2010;

Parsana et al., 2012; Grimsley et al., 2013; Gadziola et al., 2016; Li et al., 2017;

Gründemann et al., 2019). Contextual information may arise from inputs associated

with other sensory modalities (e.g., somatic sensation or olfaction) (Mcdonald, 1998;

Lanuza et al., 2004; Grimsley et al., 2013b), but in other cases the contextual

information is associated with an animal's internal state. Sources of internal state cues

to BLA include brain circuits involving modulatory neurochemicals (i.e.,

neuromodulators), known to affect the processing of sensory signals, thus shaping

attention, emotion, and goal-directed behaviors (Smythies, 2005; Krichmar, 2008;

Bargmann, 2012; Picciotto et al., 2012; Schofield & Hurley, 2018; Likhtik & Johansen,

2019).

This study investigates contextual information provided by neuromodulatory inputs to

the BLA in response to vocal communication signals. Our hypothesis is that these salient

vocalizations elicit distinct patterns of neuromodulator release into the BLA, by which

they shape the processing of subsequent meaningful sensory information. We further

hypothesize that these neuromodulatory patterns may depend on longer term

processes that are critical to vocal communication: to experience with these behaviors

and the accompanying vocalizations, to sex, and to estrous stage in females. To test

these hypotheses, we conducted playback experiments in a mouse model to understand

the behavioral and neuromodulator responses to salient vocalizations associated with

very different behavioral states.

RESULTS

To study how vocalizations affect behaviors and release of neurochemicals within BLA,

we first developed highly salient vocal stimuli associated with appetitive (mating) and

aversive (restraint) behaviors of CBA/CaJ mice (Figs. 1, S1). During interactions between

adult male and female mice, we analyzed mating-related behaviors and vocalizations to

assess the intensity of the interactions (see Materials and Methods). Comparing lower

and higher intensity interactions, we found significant differences in acoustic features

representing emotional intensity of the vocalizations produced (Fig. S2) (Lahvis et al.,

2011; Altenmüller et al., 2013; Cowen et al., 2019; Huang et al., 2021). From higher

intensity mating interactions, we selected several sequences of vocalizations to form a

20-minute mating vocal stimulus. These sequences included ultrasonic vocalizations

(USVs) with harmonics, steps, and complex structure, mostly emitted by males, and low

frequency harmonic calls (LFHs) emitted by females (Figs. 1A, S1A,C, S2B)

(Ghasemahmad, 2020). During short periods of restraint, mice produce distinctive

vocalizations that are associated with anxiety-related behaviors and increased release of

the stress hormone corticosterone (Grimsley et al., 2016). From vocal sequences

obtained in restrained mice, we created a 20-minute vocal stimulus, primarily containing

mid-frequency vocalizations (MFVs) and fewer USV and LFH syllables (Figs. 1B, S1B,C).

We next asked whether these salient vocal stimuli, associated with very different

behavioral states, could elicit distinct behaviors and patterns of neuromodulator release

into the BLA, where neurons respond to vocalizations in context-dependent ways

(Grimsley et al., 2013; Gadziola et al., 2016; Matsumoto and Okanoya, 2016). Because

previous studies suggest that interaction of cholinergic and dopaminergic systems

shapes the emission of positive and negative vocalizations (Silkstone & Brudzynski,

2020; Inagaki et al., 2020; Rojas-Carvajal et al., 2022), we specifically hypothesized that

patterns of acetylcholine (ACh) and dopamine (DA) release show differences in response

to playback of mating and restraint vocalizations. We further hypothesized that sex,

female estrous stage, and prior behavioral experience shape the behavioral and

neurochemical responses to vocal playback. We tested these hypotheses through

experiments combining playback of the vocal stimuli, behavioral tracking and

observations, and microdialysis of BLA extracellular fluid in freely moving mice (Figs. 1C,

S1D). Fluids were analyzed using a liquid chromatography-mass spectrometry (LC-MS)

technique that allowed simultaneous measurement of several neurochemicals and their

metabolites in the same dialysate samples, including ACh, DA and the serotonin

metabolite 5-HIAA (see Materials and Methods).

Prior to the study, male and female mouse subjects had no experience with sexual or

restraint behaviors. On the first two days of the experiment, mice in the experienced

group (EXP, n=31) were each exposed to 90-min sessions with mating and restraint

behaviors in a counterbalanced design (Fig. 1C). Mice in the inexperienced group (INEXP,

n=21) received no mating or restraint experience when placed in the behavioral arena

on those days. All mice were then implanted with a guide cannula for microdialysis (Fig.

1C). On the playback/sample collection day (Day 6), a microdialysis probe was inserted

into the guide cannula. After a 4-hour period of mouse habitation and probe

equilibration, we recorded behavioral reactions and sampled extracellular fluid from the

BLA before (Pre-Stim) and during a 20 min playback period, divided into two 10-min

collection/observation periods (Stim 1 and Stim 2) (Fig. 1C, S1D). Each mouse received

playback of either the mating or restraint stimuli, but not both. Data are reported only

from mice with more than 75% of the microdialysis probe implanted within the BLA

(Figs. 1D, S3).

We first tested whether playback of mating and restraint vocalizations results in

different behavioral responses in male mice. In both playback groups (n_{mating}=7,

$n_{restraint}$=6), increased attending (Fig 2A) indicated a generalized response to the vocal

stimuli. Other behaviors differed between groups, however. For instance, still-and-alert

and flinching behaviors displayed pronounced increases with restraint playback that

were not observed in the mating playback group (Fig. 2B,C).

For these groups of male mice, analysis of microdialysis samples revealed distinct,

vocalization-dependent modulation of cholinergic and dopaminergic release in the BLA.

That is, in response to mating vocalizations, ACh release decreased below pre-stimulus

levels during both playback windows, while restraint vocalizations resulted in an

increase compared to pre-stimulus ACh concentration (Fig. 2D). DA release, however,

displayed opposite patterns to ACh, increasing during playback of mating vocalizations

but decreasing during playback of restraint vocal sequences (Fig. 2E). In contrast to

these distinct patterns, the serotonin metabolite 5-HIAA showed no significant change

over time as a result of playback of either vocal stimulus (Fig. S5A). These findings

suggest that both behavioral responses and ACh and DA release are modulated in

listening male mice by the affective content of social vocalizations. Further, some

behavioral and neurochemical responses were significantly correlated, especially the

percentage change of ACh concentration (re Pre-Stimulus) with the number of flinching

behaviors (Fig. 2F).

As male and female mice emit different vocalizations (Neunuebel et al., 2015; Sangiamo

et al., 2020; Warren et al., 2020) during mating, we sought to understand whether

playback of vocalizations associated with intense mating interactions results in different

behavioral and neurochemical responses in listening male and female mice. Since our

testing included females in both estrus and non-estrus stages, we further examined the

estrous effect on neurochemical release and behavioral reactions. We used a

generalized linear model with repeated measure comparisons to examine how estrous

stage of females on the day of microdialysis affected neurochemical levels and

behaviors during playback of mating vocalizations.

Playback of mating vocalizations resulted in some general and sex-based differences in

behavioral responses. For instance, all groups displayed increased attending behavior

(Fig. S4A). In females, regardless of estrous stage, exploratory behaviors such as rearing

were significantly reduced during playback (Fig. 3A), while still-and-alert behavior

increased (Fig. S4B), compared to male mice. Other behaviors differed by estrous stage

during mating playback: females in estrus displayed a strikingly higher number of

flinching behaviors and reduced self-grooming, compared to males, while females in

non-estrus stages did not (Fig. 3B,C).

We then examined neuromodulator responses to mating vocalization playback for sex-

or estrous-stage-dependent patterns of release. This analysis revealed an estrous-

dependent modulation of ACh levels during playback: ACh concentration in estrus

females increased significantly compared to males during the first and second playback

windows (Fig. 3D), but decreased in both non-estrus females and males. DA release,

however, showed a consistent increase with mating playback across all three

experimental groups (Fig. 3E). Similar to male groups in restraint and mating playback,

the 5-HIAA release patterns in females showed no modulation during mating vocal

playback (Fig. S5B).

Like male mice exposed to restraint vocalizations, estrus females showed increases in

flinching behavior in response to mating vocalizations. Both of these groups displayed

increased ACh release. Among all mice exposed to mating playback, we observed a

positive correlation between ACh concentration and the number of flinching behaviors

(Fig. 3F). This supports the possible involvement of ACh in shaping such behavior in both

males listening to restraint calls and in estrus females listening to mating vocalizations.

All EXP mice used in the above experiments had undergone a single session each to

experience mating and restraint conditions prior to the playback session on Day 6 (Fig.

1C). Does such experience shape the release patterns of these neuromodulators in

response to vocal playback? We tested male and female mice under identical vocal

playback conditions as previous groups, except that they did not receive the restraint

and mating experiences (INEXP groups). Since only one INEXP female was in a non-

estrus stage during the playback session, our analysis of the effect of experience

included only estrus females and males.

Several behavioral responses to vocalization playback differed between EXP and INEXP

mice in a sex- or context-dependent manner. For example, only estrus females showed

experience-dependent increases in flinching behaviors (Fig. 4A) and experience-

dependent decreases in attending behaviors (not shown) in response to mating vocal

sequences. These experience effects were not observed in males in response to mating

or restraint vocal playback. Males, however, showed striking experience-dependent

increases in rearing behaviors (Fig. 4B) and locomotion in response to mating

vocalizations (not shown), but this pattern was not observed during restraint playback

for males or mating playback in estrus females. These findings indicate that behavioral

responses to salient vocalizations result from interactions between sex of the listener or

context of vocal stimuli with the previous behavioral experience associated with these

vocalizations.

A major finding is that prior experience with mating and restraint behaviors also shaped

patterns of ACh and DA release in response to vocal playback (Fig. 4C,D, S6). Our results

show that the significant effects of vocal stimulus type and of sex on ACh and DA

release, observed in our EXP groups, were absent in the INEXP groups (Figs. 4C,D,

S6A,B). For example, recall that in EXP males, restraint vocalizations increased ACh and

decreased DA, while mating vocalizations decreased ACh and increased DA. When male

mice lacked the previous mating and restraint experiences, the concentrations of ACh

and DA showed similar, non-significant changes in response to both vocal playback

types (Figs. 4C,D, S6A,B).

Also recall that in EXP animals exposed to mating vocalization playback, ACh release

increased in estrus females while decreasing in males and non-estrus females. In INEXP

estrus females, however, the ACh pattern was similar to INEXP male mice. Although we

did not include INEXP non-estrus females, the current data suggests that the effect of

estrus in modulating ACh release is experience dependent (Figs. 4C, S6A). DA release

during mating playback also appears to be experience-dependent: in both male and

female INEXP groups, DA release during mating playback remained similar to baseline

levels, unlike the comparable EXP groups (Fig. 4D). Further, 5-HIAA concentrations,

which were unaffected by sex, estrous stage, or playback type, were also unaffected by

experience (Fig. S6C). Finally, INEXP groups showed no significant correlations between

concentrations of ACh with behavior in response to vocalization playback.

Collectively, these data suggest that the playback vocalization type and estrous effects

observed in neuromodulator release patterns and behavioral reactions are mediated by

previous experience with the corresponding behaviors.

DISCUSSION

In a mouse model of acoustic communication, we showed that the motivational state of

a "sender" is reflected in the acoustics of social vocalizations. We then showed that

these vocalizations related to intense experiences of restraint and mating affect

behavioral responses in listening conspecifics. Further, behavioral responses to mating

vocalizations are shaped by sex and estrous stage. The link between hearing and

responding to vocal communication signals is formed by brain circuits that integrate

acoustic information in vocal signals with other sensory inputs, with internal state

signals, and with previous experiences of the listener (Gadziola et al., 2016; Rouby et al.,

2016; Wenstrup et al., 2020; Frühholz and Schweinberger, 2021). With access to centers

shaping behavioral responses to these sensory cues, including those involved in social

communication (Bickart) (Sah et al., 2003; Bickart et al., 2014; Hsu et al., 2014;

Wenstrup et al., 2020), the amygdala participates substantially in this process. Finally,

we showed that a single 90-min experience with intense behaviors is sufficient to

establish strong, consistent patterns of neuromodulatory release into the amygdala in

estrous- and context-dependent manners. These release patterns are expected to alter

the functional properties of the vocal communication circuits that evoke behavioral

responses.

Neuromodulatory Contributions to Behavioral Output

Functional imaging studies in humans and mechanistic studies in other species provide

substantial evidence of amygdalar involvement in processing vocalizations (Sander et

al., 2003; Andics et al., 2010; Frühholz et al., 2012, 2014, 2016; Viinikainen et al., 2012;

Abrams et al., 2016; Liebenthal et al., 2016; Pannese et al., 2016), in valence coding of

appetitive and aversive cues (Kyriazi et al., 2018; O'Neill et al., 2018; Pignatelli &

Beyeler, 2019; W. C. Huang et al., 2020; Kong & Zweifel, 2021; Smith & Torregrossa,

2021), and in shaping appropriate behavioral responses to these cues (Lim et al., 2009;

Senn et al., 2014; Zhang & Li, 2018; Gründemann et al., 2019). Nonetheless, how

contextual information is delivered to the amygdala and contributes to vocal processing

is not well understood. Since the amygdala receives strong projections from

neuromodulatory brain centers (Carlsen et al., 1985; Asan, 1997, 1998; Bigos et al.,

2008; Bocchio et al., 2016; Aitta-aho et al., 2018), and since the role of these

neurochemicals in providing internal state and contextual information is well-proven

(Bocchio et al., 2016; L. Jiang et al., 2016; Likhtik & Johansen, 2019), we hypothesized

that release patterns of neuromodulators into the BLA provide contextual information

during processing of affective vocalizations. Our results show that these emotionally

charged vocalizations result in distinct release patterns of acetylcholine (ACh) and

dopamine (DA) into the BLA of male and female mice. Further, female hormonal state

appears to influence ACh but not DA release into the BLA when processing mating

vocalizations. Such context- or state-dependent changes were not observed in release

patterns of other neurochemicals (e.g., 5-HIAA). These data indicate that during analysis

of affective vocalizations in the BLA, ACh and DA provide state- and context-related

information that can, potentially, modulate sensory processing within the BLA and thus

shape an individual's response to these vocalizations.

The BLA receives strong cholinergic projections from the basal forebrain (Carlsen et al.,

1985; Zaborszky et al., 1999; Aitta-aho et al., 2018) that contribute to ACh-dependent

processing of aversive cues and fear learning in the amygdala (Mascagni et al., 2009;

Baysinger et al., 2012; Pidoplichko et al., 2013; Tingley et al., 2014; Gorka et al., 2015; L.

Jiang et al., 2016; Minces et al., 2017). Our findings support these studies by

demonstrating increased ACh release in BLA in response to playback of aversive

vocalizations. Although the exact mechanism by which ACh affects vocal information

processing in BLA is not clear yet, the result of ACh release onto BLA neurons seems to

enhance arousal during emotional processing (Likhtik & Johansen, 2019). Several studies

suggest mechanisms by which vocalizations affect ACh release and in turn drive

behavioral responses (Fig. 5A). Cholinergic modulation in the BLA is mediated via

muscarinic and nicotinic ACh receptors on BLA pyramidal neurons and inhibitory

interneurons (Mesulam et al., 1983; Pidoplichko et al., 2013; Unal et al., 2015; Aitta-aho

et al., 2018). During the processing of sensory information in the BLA, partially non-

overlapping populations of neurons respond to cues related to positive or negative

experiences (Paton et al., 2006; Shabel & Janak, 2009; Namburi et al., 2015; Beyeler et

al., 2018). These neurons then project to different target areas involved in appetitive or

aversive behaviors—the nucleus accumbens or central nucleus of the amygdala,

respectively (Namburi et al., 2015).

In response to an aversive cue or experience (Fig. 5A), released ACh affects neurons

according to their activity. If projection neurons are at rest, ACh may exert an inhibitory

effect by activating nicotinic ACh receptors on local GABAergic interneurons, which in

turn synapse onto the quiescent pyramidal neurons. This results in GABA-A mediated

inhibitory postsynaptic potentials (IPSPs) in the pyramidal neurons. Alternately, direct

activation of M1 ACh receptors on pyramidal neurons, activating inward rectifying K+

currents, may result in additional ACh-mediated inhibition (Fig. 5A.) (Pidoplichko et al.,

2013; Unal et al., 2015; Aitta-aho et al., 2018). When BLA pyramidal neurons are already

active due to strong excitatory input associated with aversive cues, M1 receptor

activation can result in long afterdepolarizations that produce persistent firing lasting as

long as ACh is present (Unal et al., 2015; Jiang et al., 2017). Such a process may explain

persistent firing observed in single neuron responses to aversive social vocalizations in

bats (Gadziola et al., 2012; Peterson & Wenstrup, 2012). Through this process of

inhibiting quiescent neurons and enhancing activation and persistent firing in active

neurons, ACh sharpens the population signal-to-noise ratio (SNR) during the processing

of salient, aversive signals in the BLA. These neurons, processing negative cues, likely

project to the central nucleus of the amygdala (CeA) to regulate defensive behaviors

such as escape and avoidance (Fig. 5A) (Namburi et al., 2015; Beyeler et al., 2016, 2018).

In agreement, our behavioral findings show increased behaviors such as flinching

correlated with increased release of ACh during processing of aversive vocalizations.

Such prolonged afterdepolarizations provide the appropriate condition for associative

synaptic plasticity (Likhtik & Johansen, 2019) that underlies an increase in AMPA/NMDA

currents in CeA-projecting neurons during processing of aversive cues (Namburi et al.,

2015; van Vugt et al., 2020).

Dopaminergic innervation from the ventral tegmental area (VTA) (Asan, 1997, 1998)

acts on BLA neurons via D1 and D2 receptors, both G-protein coupled receptors.

Dopamine is important in reward processing, fear extinction, decision making, and

motor control (Di Ciano & Everitt, 2004; Ambroggi et al., 2008; Lutas et al., 2019). We

observed increased DA release in the BLA in response to mating vocalizations both for

males and for females across estrous stages. Electrophysiological studies show that DA

enhances sensory processing in BLA neurons by increasing the population response SNR

in a process like ACh (See et al., 2001; Kröner et al., 2005; Vander Weele et al., 2018).

Thus, during processing of mating vocalizations or those related to other rewarding

experiences, DA release in the vicinity of BLA pyramidal neurons and interneurons is

enhanced (Fig. 5B). For neurons with elevated spiking activity during processing of

appetitive vocalizations or other sensory stimuli, DA acts on D2 receptors of pyramidal

cells to further enhance neuronal firing and result in persistent firing of these projection

neurons. Conversely, in BLA projection neurons that do not respond to such positive

cues, such as CeA-projecting neurons, DA exerts a suppressive effect directly via D1

receptors and indirectly by activating inhibitory interneuron feedforward inhibition

(Kröner et al., 2005). The net result of this process in response to appetitive

vocalizations is an enhancement of activity in the reward-responding neurons and

suppression of activity in aversive-responding neurons. This process likely depends on

the increase in synaptic plasticity via enhanced AMPA/NMDA current during processing

such cues in NAc-projecting neurons in the BLA (Otani et al., 2003; Namburi et al., 2015;

van Vugt et al., 2020). Our findings suggest that this may occur in the BLA in response to

appetitive vocalizations.

Influence of sex, hormonal state, and experience

As the results with males listening to restraint vocalizations demonstrate, increased ACh

release is associated with processing aversive cues. How, then, should the increased

ACh release in estrus females during mating vocal playback be interpreted? Many

studies have suggested changes in forebrain activity and neuromodulation due to

hormonal changes in males and females (Egozi et al., 1986; van Huizen et al., 1994;

Matsuda et al., 2002; Sosa et al., 2004; Kirry et al., 2019; Mizuno et al., 2022; Krentzel et

al., 2022). For instance, the cholinergic neurons that project to the BLA, originating in

the basal forebrain, exhibit high expression of estrogen receptors that is influenced by a

female's hormonal state (Shughrue et al., 2000). During estrus, the enhanced circulating

estrogen affects release of ACh and may influence neuronal networks and behavioral

phenotypes in a distinct manner (Gibbs, 1996; McEwen, 1998). Thus, increased ACh

release in estrus females may underlie increased attentional and risk assessment

behaviors in response to vocalization playback. Combined with DA increase, it may

trigger both NAc and CeA circuit activation, resulting in both reward-seeking and

cautionary behaviors in estrus females.

Our results demonstrate the strong impact of even limited experience in shaping

behavioral and neuromodulatory responses associated with salient social vocalizations.

In the adult mice, a single 90-min session of mating and of restraint, occurring 4-5 days

prior to the playback experiment, resulted in consistent behavioral responses and

consistent and enhanced ACh and DA release into BLA for both vocalization types. As

previous work shows (Pawlak et al., 2010; S. Y. Huang et al., 2012; Nadim & Bucher,

2014) neuromodulatory inputs play crucial roles in regulating experience-dependent

changes in the brain. However, it remains unclear whether the experience shapes

neuromodulator release, or whether neuromodulators deliver the experience-related

effect into the BLA.

The interaction between ACh and DA is thought to shape motor responses to external

stimuli (Lester et al., 2010). Our results support the view that a balance of DA and ACh

may regulate the proper behavioral response to appetitive and aversive auditory cues.

For instance, increased reward-seeking behavior (rearing and locomotion) in EXP males

after mating vocalizations playback may result from the differential release of the two

neuromodulators—decreased ACh and increased DA. Further, the lack of this

differential release may be the underlying cause for the lack of such responses in INEXP

male mice. This supports the role of experience in tuning interactions of these two

neuromodulators throughout the BLA, for shaping appropriate behaviors. Overall, the

behavioral changes orchestrated by the BLA in response to emotionally salient stimuli

are most likely the result of the interaction between previous emotional experiences,

hormonal state, content of sensory stimuli, and sex of the listening animals.

1 ACKNOWLEDGMENTS

We are indebted to Sharad Shanbhag and Daniel Gavazzi for software used to condition

vocal sequences and analyze audio and behavioral data, respectively, Kristin Yeager for

statistical consulting. We thank Anthony Zampino, Austin Poth, Debin Lei, and Krish Nair

for technical assistance. We further thank Vanderbilt University Neurochemistry Core,

supported by Vanderbilt Brain Institute and the Vanderbilt Kennedy Center, that

performed the neurotransmitter sample analyses. We are grateful to Drs. Alexander

Galazyuk, Brett Schofield, Merri Rosen, Mahtab Tehrani, and Sharad Shanbhag for their

comments on this manuscript.

Funding: This work was supported by research grant R01 DC00937 to J.J.W and

Alexander Galazyuk from the National Institute on Deafness and Other Communication

Disorders of the U.S. Public Health Service. Additional support was through awards to

Z.G from the Kent State University Graduate Student Senate and Northeast Ohio

Medical University Biomedical Sciences Graduate Program Committee.

Author Contributions (CreDit). https://credit.niso.org: Conceptualization: ZG, JJW; Data

curation: ZG; Formal analysis: ZG, AM, RP, BS, KDP; Funding acquisition: ZG, JJW;

Investigation: ZG; Methodology: ZG, JJW; Project Administration: ZG, JJW; Resources ZG,

JJW; Supervision: JJW; Validation: ZG; Visualization: ZG, JJW, AM; Writing—original

draft: ZG, JJW; Writing—review and editing: ZG, JJW.

Main Figures

Figure 1. Behavioral / microdialysis experiments test how playback of affective vocal signals alters behaviors and neuromodulator release into the BLA. A. Short sample of mating vocal sequence used in playback experiments. Recording was obtained during high-intensity mating interactions and included USVs, likely emitted by males, as well as LFH calls likely emitted by females. **B.** Short sample of restraint vocal sequence used in playback experiments. Recording was obtained from an isolated, restrained mouse (see Material and Methods) and consisted mostly of MFV syllables. **C**. Experimental design in playback experiment. Days 1 and 2: each animal experienced restraint and mating behaviors (counterbalanced order across subjects). Day 2: a microdialysis guide tube was implanted in the brain above the BLA. Day 6: the microdialysis probe was inserted through guide tube into the BLA. Playback experiments began after several hours of habituation/equilibration. Behavioral observations and microdialysis sampling were obtained before, during, and after playback of one vocalization type. Microdialysis samples were analyzed using LC/MS method described in Materials and Methods. **D.** Location of microdialysis probe in BLA was confirmed by perfusion of dextran-fluorescein through probe at the end of experiment. Abbreviations: *B,* basal nucleus of the amygdala; *La*, lateral nucleus of the amygdala.

Figure 2. Behavioral and neuromodulator responses to vocal playback in male mice differ by behavioral context of vocalizations. A-C. Boxplots show number of occurrences of specified behavior in 10-min observation periods before (Pre-Stim) and during (Stim 1, Stim 2) playback of mating or restraint vocal sequences (n_{Mating} = 7, $n_{Restraint}$ = 6). Note that playback sequences during Stim 1 and Stim 2 periods were identical within each group. **A.** Both mating and restraint vocal playback similarly increased attending behaviors (time: F (2,22) =19.3, p=0.0001, partial η^2=0.64; time*context: F (2,22) =0.53, p=0.6, η^2=0.05). **B, C.** During playback, the restraint group increased still-and-alert (time: F(2,22) =6.8, p=0.005, η^2=0.4; context: F (1,11) = 9.6, p=0.01, η^2=0.5) and flinching (time: F (1.1,12.2) =14.2, p=0.002, η^2=0.6; time*context: F (1.1,12.2) =6.3, p=0.02, η^2=0.4) behaviors compared to the mating group. **D, E.** Boxplots show differential release of acetylcholine (ACh) or dopamine (DA), relative to the Pre-Stim period, during mating and restraint vocal playback (n_{Mating} = 9, $n_{Restraint}$ = 7). **D.** Significant differences in ACh for mating (decrease) vs restraint (increase) playback in stim 1 and stim2 (time: F(2,28)=0.3, p=0.7, η^2=0.02; time*context: F(2,28)=6.6,p=0.004, η^2=0.32) **E.** Significant differences for DA for mating (increase) vs restraint (decrease) playback (time: F(2,28)=0.25,p=0.8, η^2=0.02; time*context: F(2,28)=6.9,p=0.004, η^2=0.33). **F.** Across EXP male subjects in vocal playback, the number of flinching behaviors was positively correlated with change in ACh concentration relative to the Pre-Stim period. (n=15, Pearson r=0.54, p=0.03); **A-E.** GLM repeated measures with Bonferroni post hoc test, *p<0.05, **p<0.01, ***p<0.001.

Figure 3. Behavioral and neuromodulator responses to playback of mating vocal sequences differ by sex and female estrous stage. A-C. Occurrences of specified behaviors in 10-min periods before (Pre-Stim) and during (Stim 1, Stim 2) mating vocal playback (n_{Male} = 7, $n_{Estrous\ Fem}$ = 6, $n_{Non-estrous\ Fem}$ = 5). **A.** Females reared less than males during playback (time: (1.4,20.8) =1.8, p=0.2, η^2=0.1; sex: F (1,15) =10.22, p=0.006; η^2=0.4; estrus: F (1,15) =0.2, p=0.7, η^2=0.01). **B, C**. In comparison to males, estrus females but not non-estrus females showed a significant increase in flinching behavior (time: F(2,30) =20.5, p=0.000, η^2= 0.6; time* sex: F(2,30) =3.1, p=0.06, η^2= 0.2; time*estrus: F (2,30) =9.0, p=0.001, η^2=0.4) and reduced self-grooming (time: F (2,30) =2.4, p=0.1, η^2= 0.14 ; sex: F(1,15) =0.34, p=0.6, η^2= 0.02; estrus for stim1: estrous female vs male: t =2.5, p=0.02, η^2=0.3; non-estrous female vs male: t =0.7, p=0.5, η^2=0.03). **D, E.** Changes in concentration of acetylcholine (ACh) or dopamine (DA) relative to the Pre-Stim period, evoked during Stim 1 and Stim 2 periods of vocal playback (n_{Male} = 9, $n_{Estrous\ Fem}$ = 8, $n_{Non-estrous\ Fem}$ = 7). **D.** Release of ACh during mating playback showed a significant estrous effect (F (2,42) =10.0, p=0.000, η^2=0.32), increasing above Pre-Stim period in estrous females (time: F (2,42) =1.9 p=0.2, η^2=0.08). **E.** DA release during mating playback increased in all groups relative to Pre-

Stim period (time: $F_{(2,42)} = 12.4$, p=0.000, η^2=0.4), with no significant sex ($F_{(1,21)}$ =0.2, p=0.6, η^2=0.01) or estrous effect ($F_{(1,21)}$ =0.8, p=0.4, η^2=0.04). **F.** Across EXP subjects responding to mating playback, number of flinching behaviors was positively correlated with change in ACh concentration, relative to Pre-Stim period (n=18, Pearson r=0.54, p=0.02). A-E: GLM repeated measures with Bonferroni post hoc test, *p<0.05, **p<0.01, ***p<0.001.

Figure 4. A single bout each of mating and restraint experience altered behaviors and enhanced neuromodulator release in response to vocal playback. In all graphs, dots represent measures from individual animals obtained during the first 10-min playback period; connected thick lines represent mean values across subjects. **A.** Experience increased flinching responses in estrus female mice, but not in males in mating or restraint vocal playback (Females: $n_{EXP} = 6$, $n_{INEXP} = 8$, t=5.0, p=0.000; Male-mating: $n_{EXP} = 8$, $n_{INEXP} = 7$, t=0.4, p=0.7; Male-restraint: $n_{EXP} = 7$, $n_{INEXP} = 7$, t=0.6, p=0.6). **B.** Experience increased rearing responses in males exposed to mating playback (Male-mating: $n_{EXP} = 8$, $n_{INEXP} = 7$, t=3.2, p=0.007), but not in females (t=1.4, p=0.2) or males in restraint playback (t=0.4, p=0.7). **C.** EXP mice displayed consistent estrus-(estrous female vs male: t=4.5, p=0.000) and context-dependent (mating vs restraint: t=3.2, p=0.006) changes in Ach. Such changes were not observed in INEXP groups during playback period (estrous female vs male: t= 0.1 , p=0.9; mating vs restraint: t=1.6, p=0.1). **D**. Context dependent increase of DA in EXP mice during mating playback (estrous female vs male: t=0.5, p=0.6, mating vs restraint: t=2.5, p=0.02) was absent in INEXP animals (estrous female vs male: t=0.7, p=0.5, mating vs restraint: t=0.5, p=0.6). A-D. GLM repeated measures with Bonferroni post hoc test, *p<0.05, **p<0.01, ***p<0.001.

Figure 5. Proposed model for neuromodulation of salient vocalization processing via acetylcholine (ACh) and dopamine (DA) in the basolateral amygdala (BLA). A. Cholinergic modulation of CeA-projecting neurons during aversive vocalization cue processing in the BLA. In the presence of aversive cues, ACh released from the basal forebrain acts on m1 ACh receptors (m1AChRs) to enhance the cue-induced excitatory responses of CeA -projecting neurons. In contrast, NAc-projecting neurons are quiescent, because they do not respond to aversive cues and are inhibited by interneurons that are activated through nicotinic ACh receptors (nAChRs). **B.** Dopaminergic modulation and enhancement of signal-to-noise ratio in response to reward-associated cues (appetitive vocalizations). When vocalizations or other rewarding cues are present, release of DA from VTA enhances D2R-mediated excitation in NAc-projecting neurons that are responsive to positive cues. In contrast, CeA -projecting neurons are not responsive to rewarding vocalizations and are inhibited by local interneurons. DA is thought to act on D1 DA receptors (D1Rs) in these local interneurons to shape a direct inhibition onto CeA-projecting neurons. Abbreviations: ACx: auditory cortex; BF, basal forebrain; CeA, central nucleus of amygdala; MGB, medial geniculate body; NAc, nucleus accumbens; VTA, ventral tegmental area.

Supplemental Figure Captions

Supplemental Figure 1. Features of vocal stimuli in playback experiments. A. Two-second segments from three different vocal sequence exemplars in mating illustrate common syllable types and sequencing. **B.** Two-second segments from three different vocal sequence exemplars in restraint illustrate common syllable types and sequencing. **C.** Syllable types in mating playback sequences differ substantially from those in restraint playback sequences. Percentages indicate frequency-of-occurrence of syllable types across all examplars used in mating or restraint vocal stimuli (n_{Mating} = 545, $n_{Restraint}$ = 622 syllables). **D.** Detailed sequencing of vocal stimuli, shown here for mating playback. A 20-min period of vocal playback was formed by seven repeated stimulus blocks of 170 s. The stimulus blocks were composed of five vocal exemplars of variable length, with each exemplar followed by an equal duration of silence. Stimuli during the Stim1 and Stim 2 playback windows thus included identical blocks but in slightly altered patterns.

Supplemental Figure 2. Features of mating vocalizations differ based on interaction intensity.
A. Behaviors associated with lower and higher intensity of mating, respectively. **B.** Percentage of different syllable types associated with lower and higher intensity mating interactions (n $_{pairs}$ =5; recording duration, 30 min each). **C.** The mean duration of syllables is significantly higher when emitted during higher intensity mating interactions. Each line represents one mating pair (n$_{LI}$ = 3107, n$_{HI}$ = 3093 syllables, F (12, 6168.1) = 289.7, p=0.000, linear mixed model). **D.** The mean inter-syllable interval is significantly lower when emitted during higher intensity mating interactions. Each line represents one mating pair (n$_{LI}$ = 2944, n$_{HI}$ = 2920 syllables, F (12,5826.3) =3.31, p<0.001, linear mixed model). *p<0.05, **p<0.01, ***p<0.001.

EXP Groups

INEXP Groups

Supplemental Figure 3. Microdialysis probe locations for EXP and INEXP Groups. Colored lines indicate recovered probe tracks that resulted from infusion of fluorescent tracers. Black solid lines indicate external capsule; black dashed lines indicate major amygdalar subdivisions. Abbreviations: *B,* basal nucleus of amygdala; *CeA*, central nucleus of amygdala; *La*, lateral nucleus of amygdala.

Supplemental Figure 4. Behavioral changes during mating vocal playback in EXP male and female mice. Boxplots show number of occurrences of the specified behavior in 10-min periods before (Pre-Stim) and during (Stim 1, Stim 2) vocal sequence playback in EXP mice. **A.** Attending behavior increased in response to mating vocal playback, regardless of sex or estrus stage (time: $F_{(2,30)} = 32.6$; $p<0.001$, $\eta 2=0.7$; time*sex: $F_{(2,30)} = 0.12$; $p=0.9$, $\eta 2=0.008$ time*estrus: $F_{(2,30)} = 1.1$; $p=0.4$, $\eta 2=0.07$). **B.** Still-and-Alert behavior increased in response to mating vocalizations for female but not male mice (time: $F_{(2,30)} = 4.8$; $p=0.01$, $\eta 2=0.24$; Sex: $F_{(1,15)} =5.17$, $p=0.04$, partial $\eta 2= 0.3$, estrus: $(1,15) =0.07$, $p=0.8$, partial $\eta 2= 0.005$). GLM repeated measures with Bonferroni post hoc test, *$p<0.05$, ***$p<0.001$.

Supplemental Figure 5. Concentration of serotonin metabolite 5-HIAA in EXP animals showed no significant context-, sex- or estrous-dependent changes with vocal stimulus playback. Boxplots show change in concentration of 5-HIAA in 10-min periods from the baseline level (Pre-Stim) during (Stim 1, Stim 2) vocal sequence playback. **A.** Comparison of mating and restraint groups of male mice (n Mating=9, n Restraint=7; time: $F_{(2,28)}$ =0.02, p=0.9, η^2 =0.001; time*context: $F_{(2,28)}$ = 0.97, p=0.4, η^2 =0.06). **B.** Comparison of male and female mice during mating sequence playback (n_{Male} = 9, $n_{Estrous\ Fem}$ = 8, $n_{Non-estrous\ Fem}$ = 7, time: $F_{(2,42)}$ =1.7, p=0.2, η^2 =0.07; time*sex: $F_{(2,42)}$ = 0.2, p=0.8, η^2 =0.01; time*estrus: $F_{(2,42)}$ =1.2, p=0.3, η^2 =0.05). GLM repeated measures with Bonferroni post hoc test.

Supplemental Figure 6. In INEXP mice, vocal playback failed to evoke significant differences in neuromodulator release between experimental groups. Boxplots show change in concentration of the specified neuromodulator in 10-min periods compared to the baseline level (Pre-Stim) during (Stim 1, Stim 2) vocal sequence playback. **A.** ACh release shown for both playback periods (mating vs restraint: time: $F_{(2,24)} = 0.14$; p=0.9; time*context: $F_{(2,24)} = 0.56$; p=0.6; male vs female: time: $F_{(2,24)} = 0.9$; p=0.4; time*sex**:** $F_{(2,24)} = 0.01$; p=0.9). **B.** DA release shown for both playback periods (mating vs restraint: time: $F_{(2,24)} = 1.2$; p=0.3; time*context**:** $F_{(2,24)} = 0.2$; p=0.8; male vs female: time: $F_{(2,24)} = 0.4$; p=0.7; time*sex: $F_{(2,24)} = 0.4$; p=0.7). **C.** 5-HIAA concentration shown for both playback periods (mating vs restraint: time: $F_{(1.3,15.6)} = 0.005$, p=0.9; time*context: $F_{(1.3,15.6)} = 2.1$; p=0.17; male vs female: time: $F_{(1.3,15.7)} = 0.9$; p=0.4; time* sex: $F_{(1.3,15.7)} = 0.3$; p=0.7). GLM repeated measure with Bonferroni post hoc test, $n_{INEXP} = 21$.