Emotional Vocalizations Alter Behaviors and

Neurochemical Release into the Amygdala

Authors:

Zahra Ghasemahmad[1,2,3], Aaron Mrvelj[1], Rishitha Panditi[1], Bhavya Sharma[1], Karthic Drishna
Perumal[1], Jeffrey J. Wenstrup[1,2,3]

Affiliations:

[1]Department of Anatomy and Neurobiology and Hearing Research Group, Northeast Ohio
Medical University; Rootstown, Ohio, 44272.

[2]School of Biomedical Sciences, Kent State University; Kent, Ohio, 44240.

[3]Brain Health Research Institute, Kent State University; Kent, Ohio, 44240.

Corresponding author:

Jeffrey J. Wenstrup, Ph.D.

Northeast Ohio Medical University, 4209 State Route 44, Rootstown, OH 44272

Voice: 330.325.6630, Email: jjw@neomed.edu

Keywords:

acetylcholine, acoustic communication, basolateral amygdala, dopamine, estrus, mating,
microdialysis, mouse, restraint, serotonin, vocalization playback

ABSTRACT

The basolateral amygdala (BLA), a brain center of emotional expression, contributes to acoustic

communication by first interpreting the meaning of social sounds in the context of the listener's

internal state, then organizing the appropriate behavioral responses. We propose that modulatory

neurochemicals such as acetylcholine (ACh) and dopamine (DA) provide internal-state signals to

the BLA while an animal listens to social vocalizations. We tested this in a vocal playback

experiment utilizing highly affective vocal sequences associated with either mating or restraint,

then sampled and analyzed fluids within the BLA for a broad range of neurochemicals and

observed behavioral responses of adult male and female mice. In male mice, playback of

restraint vocalizations increased ACh release and usually decreased DA release, while playback

of mating sequences evoked the opposite neurochemical release patterns. In non-estrus female

mice, patterns of ACh and DA release with mating playback were similar to males. Estrus

females, however, showed increased ACh, associated with vigilance, as well as increased DA,

associated with reward-seeking. Experimental groups that showed increased ACh release also

showed the largest increases in an aversive behavior. These neurochemical release patterns and

several behavioral responses depended on a single prior experience with the mating and restraint

behaviors. Our results support a model in which ACh and DA provide contextual information to

sound analyzing BLA neurons that modulate their output to downstream brain regions

controlling behavioral responses to social vocalizations.

SIGNIFICANCE STATEMENT

In social communication by sound, an animal interprets the meaning of vocalizations based on its

prior experience, other sensory stimuli, and its internal state. The basolateral amygdala (BLA), a

brain center of emotional expression, contributes to this analysis. We found that the modulatory

neurochemicals acetylcholine and dopamine were released differentially into the BLA depending

on the emotional content of the vocalizations, the sex and hormonal state of the animal, as well

as its prior experience. Our results suggest that acetylcholine and dopamine provide context- and

hormonal state-dependent information to sound-analyzing BLA neurons that modulates their

output to downstream brain centers controlling behavioral responses to social vocalizations.

INTRODUCTION

In social interactions utilizing vocal communication signals, the acoustic features of the signals

carry emotional information (Altenmüller et al., 2013; Darwin, 1872; Seyfarth & Cheney, 2003).

Listeners receive and analyze the acoustic information, compare it with previous experiences,

identify the salience and valence of such information, and respond with appropriate behaviors.

These integrated functions depend on brain circuits that include the amygdala, a region located

within the temporal lobe that is recognized to play a role in orchestrating emotional responses to

salient sensory stimuli (J. E. LeDoux, 2000; McGaugh, 2002; Sah et al., 2003). The amygdalar

target of auditory inputs from the thalamus and cortex is the basolateral amygdala (BLA) (J.

LeDoux et al., 1984; Romanski & LeDoux, 1993; Shi & Cassell, 1997; Tsukano et al., 2019). By

integrating this auditory input with other sensory inputs and inputs from other limbic areas, BLA

neurons shape appropriate behavioral responses (Beyeler et al., 2016; Gründemann et al., 2019;

Namburi et al., 2015, 2016) via projections to downstream targets such as the nucleus accumbens

(Ambroggi et al., 2008; Stuber et al., 2011) and central nucleus of the amygdala (Ciocchi et al.,

2010).

The BLA processes vocal and other acoustic information in a context-dependent manner

(Gadziola et al., 2016; Grimsley et al., 2013; Matsumoto et al., 2016; Wenstrup et al., 2020;

Wiethoff et al., 2009). Contextual information may arise from inputs associated with other

sensory modalities (e.g., somatic sensation or olfaction) (Grimsley et al., 2013; Lanuza et al.,

2004; McDonald, 1998), but in other cases the contextual information is associated with an

animal's internal state. Sources of internal state cues to BLA include brain circuits involving

modulatory neurochemicals (i.e., neuromodulators), known to affect the processing of sensory

signals, thus shaping attention, emotion, and goal-directed behaviors (Bargmann, 2012; Likhtik

& Johansen, 2019; Schofield & Hurley, 2018). While previous work has demonstrated a role for some neuromodulators—dopamine (DA) and acetylcholine (ACh)—in the production of social vocalizations (Rojas-Carvajal et al., 2022; Silkstone & Brudzynski, 2020), it remains unclear how these and other neuromodulators contribute to vocal processing in social interactions.

This study investigates contextual information provided by neuromodulatory inputs to the BLA in response to vocal communication signals. Our hypothesis is that these salient vocalizations elicit distinct patterns of neuromodulator release into the BLA, by which they shape the processing of subsequent meaningful sensory information. We further hypothesize that these neuromodulatory patterns may depend on longer term processes that are critical to vocal communication: to experience with these behaviors and the accompanying vocalizations, to sex, and to estrous stage in females. To test these hypotheses, we conducted playback experiments in a mouse model to understand the behavioral and neuromodulator responses to salient vocalizations associated with very different behavioral states.

RESULTS

To study how vocalizations affect behaviors and release of neurochemicals within BLA, we first developed highly salient vocal stimuli associated with appetitive (mating) and aversive (restraint) behaviors of CBA/CaJ mice. From more intense, mating-related interactions between adult male and female mice that included female head-sniffing and attempted or actual mounting (Gaub et al., 2016; Ghasemahmad, 2020, see Materials and Methods), we selected several sequences of vocalizations to form a 20-minute mating vocal stimulus. These sequences included ultrasonic vocalizations (USVs) with harmonics, steps, and complex structure, mostly emitted by males, and low frequency harmonic calls (LFHs) emitted by females (Fig. 1A,C) (Finton et al., 2017;

Gaub et al., 2016; Ghasemahmad, 2020; Hanson & Hurley, 2012). During short periods of restraint, mice produce distinctive mid-frequency vocalizations (MFVs) that are associated with anxiety-related behaviors and increased release of the stress hormone corticosterone (Dornellas et al., 2021; Grimsley et al., 2016; Niemczura et al., 2020). From vocal sequences produced by restrained mice, we created a 20-minute vocal stimulus, primarily containing MFVs and fewer USV and LFH syllables (Fig. 1B,C).

We next asked whether these salient vocal stimuli, associated with very different behavioral states, could elicit distinct behaviors and patterns of neuromodulator release into the BLA. We focused on the neuromodulators acetylcholine (ACh) and dopamine (DA), since previous work suggests that these neuromodulatory systems interact in the emission of positive and negative vocalizations (Rojas-Carvajal et al., 2022; Silkstone & Brudzynski, 2020). Our experiments combined playback of the vocal stimuli, behavioral tracking and observations (see Table 1 for descriptions), and microdialysis of BLA extracellular fluid in freely moving mice (Fig. 1D).

Prior to the study, male and female mouse subjects had no experience with sexual or restraint behaviors. On the first two days of the experiment, mice in an experienced group (EXP, n=31) were each exposed to 90-min sessions with mating and restraint behaviors in a counterbalanced design (Fig. 1D). Mice were then implanted with a guide cannula for microdialysis. On the playback/sample collection day (Day 6), a microdialysis probe was inserted into the guide cannula. After a 4-hour period of mouse habituation and probe equilibration, we recorded behavioral reactions and sampled extracellular fluid from the BLA before (Pre-Stim) and during a 20 min playback period, divided into two 10-min stimulation/collection/observation periods that are designated Stim 1 and Stim 2 (Fig. 1D,E; see also Materials and Methods). Each mouse received playback of either the mating or restraint stimuli, but not both: same-day presentation of

both stimuli would require excessively long playback sessions, the condition of the same probe

would likely change on subsequent days, and quality of a second implanted probe on a

subsequent day was uncertain. Fluids were analyzed using a liquid chromatography/mass

spectrometry (LC/MS) technique that allowed simultaneous measurement of several

neurochemicals and their metabolites in the same dialysate samples, including ACh, DA, and the

serotonin metabolite 5-HIAA (see Materials and Methods). All neurochemical results during

Stim 1 and Stim 2 periods are expressed as a percentage relative to the Pre-Stim control period.

However, raw values of ACh, DA, and 5-HIAA are reported in supplementary tables 1, 2, and 3,

respectively. Data are reported only from mice with more than 75% of the microdialysis probe

implanted within the BLA (Fig. 2).

We first describe tests to examine whether playback of mating and restraint vocalizations results

in different behavioral and neurochemical responses in male mice. We observed that two

behaviors, still-and-alert and flinching, showed differential effects of playback type, increasing

during restraint playback relative to mating playback during the Stim 1 period (Fig. 3A, B).

There were also distinct patterns of ACh and DA release into the BLA depending on the type of

vocalization playback (Fig 3C, D). Thus, in response to restraint vocalizations, we observed an

increase in ACh concentration during both Stim 1 ($M \pm SD$: +24 ± 31% re Pre-Stim baseline) and

Stim 2 (+25 ± 16%) playback windows. Mating vocalizations, however, resulted in a decrease in

ACh release during both playback periods (Stim 1: -19 ± 21.0%; Stim 2: -16.4 ± 24.0%). (Fig.

3C). DA release displayed opposite patterns to ACh, increasing during playback of mating

vocalizations during Stim 1 (+12.0 ± 27.4%) and Stim 2 (+25 ± 39% re baseline), but decreasing

during playback of restraint vocal sequences (Stim 1: -11.0± 11.2%; Stim 2: -22 ± 25.6%) (Fig.

3D). For ACh, levels differed significantly between the mating and restraint groups during both

Stim 1 and Stim 2, while DA levels differed significantly only during Stim 2 (Fig. 3C, D). In

contrast, the serotonin metabolite 5-HIAA showed no distinct pattern over time following

playback of either vocal stimulus, nor significant differences between groups (Fig. 3E). These

findings suggest that both behavioral responses and ACh and DA release are modulated in

listening male mice by the affective content of social vocalizations.

As male and female mice emit different vocalizations during courtship and mating (Finton et al.,

2017; Grimsley et al., 2013; Neunuebel et al., 2015; Sales (née Sewell), 1972), we tested whether

playback of vocal interactions associated with mating (Fig. 1A) results in different behavioral

and neurochemical responses in listening male and female mice. Since our testing included both

estrus and non-estrus females, we further examined the estrous effect on neurochemical release

and behavioral reactions.

Playback of mating vocalizations resulted in some general and some sex-based differences in

behavioral responses. For instance, all groups displayed increased attending behavior (Fig. 4A).

In females, regardless of estrous stage, rearing decreased and still-and-alert behavior increased

significantly relative to male mice (Fig. 4B, C). This change in motor activity was further

supported by video tracking results showing a significant decrease in distance traveled by female

mice in response to mating vocal playback (significant sex effect ($F_{(1,15)}$ =9.3, p=0.008,

$\eta2$=0.4). However, one behavioral change was estrous dependent: females in estrus displayed a

strikingly higher number of flinching behaviors compared to males and non-estrus females

during both Stim 1 and Stim 2 periods (Fig. 4D). Our analysis of neuromodulator responses to

mating vocalization playback revealed an estrous-dependent modulation of ACh levels during

playback. ACh concentration in estrus females increased during both Stim 1 and Stim 2 periods,

whereas ACh decreased in both males and non-estrus females (Fig. 4E). Moreover, during

playback, post hoc comparison with Bonferroni correction showed that the ACh in estrus females

was significantly higher (Stim 1, +29.3 ± 27.0 %; Stim 2, +25.5 ± 22%) than both males (Stim 1,

-18 ± 21.0%; Stim 2, -16.0 ± 24.0%) and non-estrus females (Stim 1, -6.0 ± 14 %; Stim 2, -35.0

± 29.6%). DA release showed increases during mating playback within all three experimental

groups (Males: Stim 1, +12.0 ± 27.0%, Stim 2, +26.0 ± 39.1%;Estrus females: Stim 1, +24.1 ±

37.0%, Stim 2, +23.0 ± 27.0%; Non-estrus females: Stim 1, +27.0 ± 17.0%, Stim 2, +49.0 ±

48.0%), but no significant differences among groups (Fig. 4F). Similar to male groups in

restraint and mating playback, the 5-HIAA release patterns in females showed no clear

modulation pattern during mating vocal playback or differences among groups (Fig. 4G).

Like male mice exposed to restraint vocalizations, estrus females showed robust and significant

increases in flinching behavior in response to mating vocalizations. Both groups displayed

increased ACh release. This supports the possible involvement of ACh in shaping such behavior

in both males listening to restraint calls and in estrus females listening to mating vocalizations.

Note that neuromodulator release, including ACh, has been previously linked to motor behaviors

(Wall & Woolley, 2020), but the changes in ACh that we observed showed no relationship with

behaviors involving motor activity such as rearing (restraint males: Stim 1: n = 6, r = 0.1, p =

0.8; Est females: n = 6, r = 0.7, p=0.08), locomotion (Stim 1: restraint males, n = 6, r = -0.2, p =

0.7; Est females: n = 6, r = 0.3, p = 0.6), or distance traveled (Stim 1: restraint males, n = 6 , r =

0.16, p = 0.8; Est females: n = 6, r = -0.6, p = 0.2). This suggests that the observed changes in

ACh reflect the valence of these vocalizations.

All EXP mice used in the above experiments had undergone a single session each to experience

mating and restraint conditions prior to the playback session on Day 6 (Fig. 1D). Does such

experience shape the release patterns of these neuromodulators in response to vocal playback?

We tested male and female mice under identical vocal playback conditions as previous groups,

except that they did not receive the restraint and mating experiences (INEXP groups). Since only

one INEXP female was in a non-estrus stage during the playback session, our analysis of the

effect of experience included only estrus females and males.

Several behavioral responses to vocalization playback differed between EXP and INEXP mice in

a sex- or context-dependent manner. For example, EXP estrus females showed significantly

increased sound-evoked flinching behaviors compared to INEXP estrus females (Fig. 5A) in

response to mating vocal sequences. These experience effects were not observed in males in

response to mating or restraint vocal playback. Further, the differences in flinching behavior

among EXP groups (male-mating vs male-restraint or estrus female-mating) were not evident

among INEXP groups. For rearing behavior, EXP males responding to mating vocalizations

showed an increase compared to INEXP males (Fig. 5B). This pattern was not observed during

restraint playback for males or mating playback in estrus females. These findings indicate that

behavioral responses to salient vocalizations result from interactions between sex of the listener

or context of vocal stimuli with the previous behavioral experience associated with these

vocalizations.

A major finding is that prior experience with mating and restraint behaviors shaped patterns of

ACh release in response to vocal playback; there was a significant interaction of context and

experience for ACh ($F(1, 39) = 12.7$, $P<0.001$; Figs. 5C, 6A). Thus, ACh release was

significantly different between INEXP and EXP males in restraint playback groups. A similar

pattern was observed in INEXP estrus females in response to mating vocal playback, i.e., the

concentration of ACh re baseline failed to show such pronounced increases as observed in EXP

females and showed a significant experience effect (Sex*experience interaction: $F (1, 39) = 7.8$,

P= 0.008; Figs. 5C, 6A). Further, INEXP males in mating playback group failed to show the

reduced ACh concentration observed in EXP males (INEXP Stim 1: +9.6 ± 15.0%; INEXP Stim

2: +6.1 ± 30.0; EXP Stim 1: -19.0 ± 21.0%; EXP Stim 2: -16.0 ± 24.0 %), even though these

differences were not significant (Figs. 5C, 6A).

DA release patterns, which were weaker in EXP groups compared to ACh patterns (Figs. 3C vs

3D; 4E vs 4F) showed no significant experience-dependent changes in any group (Fig. 5D).

There was no evidence that vocal playback altered DA release in any of the three groups of

INEXP mice (Fig 6B), unlike what was observed in EXP groups (Figs. 3D, 4F). 5-HIAA

concentrations, which were unaffected by sex, estrous stage, or playback type (Figs. 3E, 4G),

were also unaffected by experience (Fig. 6C).

Collectively, these data suggest that the playback vocalization type and estrous effects observed

in ACh release patterns and behavioral reactions depend on previous experience with the

corresponding behaviors.

DISCUSSION

Functional imaging studies in humans and mechanistic studies in other species provide substantial evidence that the amygdala participates in circuits that process vocalizations (Frühholz et al., 2016; Liebenthal et al., 2016; Sander et al., 2003; Wenstrup et al., 2020; Voytenko et al., 2023), assess the valence of appetitive and aversive cues (Kyriazi et al., 2018; O'Neill et al., 2018; Pignatelli & Beyeler, 2019; Smith & Torregrossa, 2021), and shape appropriate behavioral responses to these cues (Gründemann et al., 2019; Lim et al., 2009; Schönfeld et al., 2020; Zhang & Li, 2018). Nonetheless, how contextual information is delivered to the amygdala and contributes to vocal processing is not well understood. Since the amygdala receives strong projections from neuromodulatory brain centers (Aitta-aho et al., 2018; Asan, 1997, 1998; Bigos et al., 2008; Bocchio et al., 2016; Carlsen et al., 1985), and since the role of these neurochemicals in providing internal state and contextual information is well documented (Bocchio et al., 2016; Jiang et al., 2016; Likhtik & Johansen, 2019), we hypothesized that release patterns of neuromodulators into the BLA provide contextual information during processing of affective vocalizations. Our results show that these emotionally charged vocalizations result in distinct release patterns of acetylcholine (ACh) and dopamine (DA) into the BLA of male and female mice. Further, female hormonal state appears to influence ACh but not DA release into the BLA when processing mating vocalizations. Such context- or state-dependent changes were not observed in patterns of other neurochemicals (e.g., 5-HIAA), nor in the absence of experience with behaviors associated with the vocalizations. In particular, we showed that a single 90-min experience with intense behaviors is sufficient to establish strong, consistent patterns of ACh release into the amygdala. These data indicate that during analysis of affective vocalizations in the BLA, ACh and DA provide experience-dependent, state- and context-related

information that can potentially modulate sensory processing within the BLA and thus shape an individual's response to these vocalizations.

The linkage between neuromodulator release and behavioral responses to vocalizations varies. The strongest case is for a link between increased ACh release and flinching behavior. Males in the restraint playback group and estrus females in the mating playback group both displayed significantly increased ACh release and flinching behavior, occurring in both playback periods. This suggests a mechanistic relationship. Other significant behavioral responses to vocal stimuli did not closely match the timing of neurochemical changes, suggesting a weak mechanistic link. Overall, it is not surprising that most behaviors and neurochemicals do not match well, given the poor temporal resolution of the neurochemical sampling. We believe that our results show the need for a finer analysis of DA and ACh release based on techniques for high temporal resolution of neurochemical measurements (e.g., genetically encoded ACh indicator (Jing et al., 2018)), coupled with short interval behavioral observations. Only then can the role of the BLA and neuromodulator inputs in behavioral vocal responses be established more strongly. Further, we note that no conclusion can be made regarding relationships between noradrenalin and serotonin concentrations and vocal playback, since these were not detected by the neurochemical analysis.

The BLA receives strong cholinergic projections from the basal forebrain (Aitta-aho et al., 2018; Carlsen et al., 1985) that contribute to ACh-dependent processing of aversive cues and fear learning in the amygdala (Baysinger et al., 2012; Gorka et al., 2015; Jiang et al., 2016; Tingley et al., 2014). Our findings support these studies by demonstrating increased ACh release in BLA in EXP animals in response to playback of aversive vocalizations. Although the exact mechanism by which ACh affects vocal information processing in BLA is not clear yet, the result of ACh

release onto BLA neurons seems to enhance arousal during emotional processing (Likhtik &

Johansen, 2019). Our results, in conjunction with previous work, suggest mechanisms by which

vocalizations affect ACh release and in turn drive behavioral responses (Fig. 7A). Cholinergic

modulation in the BLA is mediated via muscarinic and nicotinic ACh receptors on BLA

pyramidal neurons and inhibitory interneurons (Aitta-aho et al., 2018; Mesulam et al., 1983;

Pidoplichko et al., 2013; Unal et al., 2015). During the processing of sensory information in the

BLA, partially non-overlapping populations of neurons respond to cues related to positive or

negative experiences (Namburi et al., 2015; Paton et al., 2006; Smith & Torregrossa, 2021).

These neurons then project to different target areas involved in appetitive or aversive

behaviors—the nucleus accumbens or central nucleus of the amygdala, respectively (Namburi et

al., 2015).

In response to an aversive cue or experience (Fig. 7A), released ACh affects BLA neurons

according to their activity. If projection neurons are at rest, ACh may exert an inhibitory effect

by activating nicotinic ACh receptors on local GABAergic interneurons, which in turn synapse

onto the quiescent pyramidal neurons. This results in GABA-A mediated inhibitory postsynaptic

potentials (IPSPs) in the pyramidal neurons. Alternately, direct activation of M1 ACh receptors

on pyramidal neurons, activating inward rectifying K+ currents, may result in additional ACh-

mediated inhibition (Fig. 7A) (Aitta-aho et al., 2018; Pidoplichko et al., 2013; Unal et al., 2015).

When BLA pyramidal neurons are already active due to strong excitatory input associated with

aversive cues, M1 receptor activation can result in long afterdepolarizations that produce

persistent firing lasting as long as ACh is present (Jiang et al., 2016; Unal et al., 2015). Such a

process may explain persistent firing observed in single neuron responses to aversive social

vocalizations in bats (Gadziola et al., 2012; Peterson & Wenstrup, 2012). Through this process of

inhibiting quiescent neurons and enhancing activation and persistent firing in active neurons,

ACh sharpens the population signal-to-noise ratio (SNR) during the processing of salient,

aversive signals in the BLA. These neurons, processing negative cues, likely project to the

central nucleus of the amygdala (CeA) to regulate defensive behaviors such as escape and

avoidance (Fig. 7A) (Beyeler et al., 2016; Davis et al., 2010; Namburi et al., 2015). In

agreement, our behavioral findings show increased behaviors such as flinching along with

increased release of ACh during processing of aversive vocalizations. Such prolonged

afterdepolarizations provide the appropriate condition for associative synaptic plasticity (Likhtik

& Johansen, 2019) that underlies an increase in the AMPA/NMDA current ratio in CeA-

projecting neurons during processing of aversive cues.

Dopaminergic innervation from the ventral tegmental area (VTA) (Asan, 1998) acts on BLA

neurons via D1 and D2 receptors, both G-protein coupled receptors. Dopamine is important in

reward processing, fear extinction, decision making, and motor control (Ambroggi et al., 2008;

Di Ciano & Everitt, 2004; Lutas et al., 2019). We observed increased DA release in the BLA in

EXP groups in response to mating vocalizations both for males and for females across estrous

stages. Electrophysiological studies show that DA enhances sensory processing in BLA neurons

by increasing the population response SNR in a process like ACh (Kröner et al., 2005; Vander

Weele et al., 2018). Thus, during processing of mating vocalizations or those related to other

rewarding experiences, DA presence in the vicinity of BLA pyramidal neurons and interneurons

is enhanced (Fig. 7B). For neurons with elevated spiking activity during processing of appetitive

vocalizations or other sensory stimuli, DA acts on D2 receptors of pyramidal cells to further

enhance neuronal firing and result in persistent firing of these projection neurons. Conversely, in

BLA projection neurons that do not respond to such positive cues, such as CeA-projecting

neurons, DA exerts a suppressive effect directly via D1 receptors and indirectly by activating

inhibitory interneuron feedforward inhibition (Kröner et al., 2005). The net result of this process

in response to appetitive vocalizations is an enhancement of activity in the reward-responding

neurons and suppression of activity in aversive-responding neurons. This process likely depends

on the increase in synaptic plasticity via an enhanced AMPA/NMDA current ratio during

processing such cues in NAc-projecting neurons in the BLA (Namburi et al., 2015; Otani et al.,

2003; van Vugt et al., 2020). Our findings suggest that this may occur in the BLA in response to

appetitive vocalizations.

As the results with males listening to restraint vocalizations demonstrate, increased ACh release

in BLA is associated with processing aversive cues. How, then, should the increased ACh release

in estrus females during mating vocal playback be interpreted? Previous work shows that

neuromodulation of amygdalar and other forebrain activity is altered by sex hormone/receptor

changes in males and females (Egozi et al., 1986; Kalinowski et al., 2023; Kirry et al., 2019;

Matsuda et al., 2002; Mizuno et al., 2022; van Huizen et al., 1994). For instance, the cholinergic

neurons that project to the BLA, originating in the basal forebrain, exhibit high expression of

estrogen receptors that is influenced by a female's hormonal state (Shughrue et al., 2000).

During estrus, the enhanced circulating estrogen affects release of ACh and may influence

neuronal networks and behavioral phenotypes in a distinct manner (Gibbs, 1996; McEwen,

1998). Thus, increased ACh release in estrus females may underlie increased attentional and risk

assessment behaviors in response to vocalization playback, as we observed with flinching

behavior in this experimental group. Combined with DA increase, it may trigger both NAc and

CeA circuit activation, resulting in both reward-seeking and cautionary behaviors in estrus

females.

Our results demonstrate the strong impact of even limited experience in shaping behavioral and neuromodulatory responses associated with salient social vocalizations. In the adult mice, a single 90-min session of mating and of restraint, occurring 4-5 days prior to the playback experiment, resulted in consistent behavioral responses and consistent and enhanced ACh release into BLA for both vocalization types. As previous work shows (Huang et al., 2012; Nadim & Bucher, 2014; Pawlak et al., 2010), neuromodulatory inputs play crucial roles in regulating experience-dependent changes in the brain. However, it remains unclear whether the experience shapes neuromodulator release, or whether neuromodulators deliver the experience-related effect into the BLA. It is also unknown whether these experiences were specific to the type of vocalization playback associated with the experience, or whether either experience had a more generalized effect on responses to all vocalization playback, as suggested by the broad physiological effects of mating or restraint (Leuner et al 2010; Arnold et al., 2019).

The interaction between ACh and DA is thought to shape motor responses to external stimuli (Lester et al., 2010). Our results support the view that a balance of DA and ACh may regulate the proper behavioral response to appetitive and aversive auditory cues. For instance, increased reward-seeking behavior (rearing and locomotion) in EXP males during mating vocalizations playback may result from the differential release of the two neuromodulators—decreased ACh and increased DA. Further, the lack of this differential release may be the underlying cause for the lack of such responses in INEXP male mice. This supports the role of experience in tuning interactions of these two neuromodulators throughout the BLA, for shaping appropriate behaviors. Overall, the behavioral changes orchestrated by the BLA in response to emotionally salient stimuli are most likely the result of the interaction between previous emotional experiences, hormonal state, content of sensory stimuli, and sex of the listening animals.

MATERIALS AND METHODS

Animals

Experimental procedures were approved by the Institutional Animal Care and Use Committee at

Northeast Ohio Medical University. A total of 83 adult CBA/CaJ mice (Jackson Labs, p90-

p180), male and female, were used for this study. Animals were maintained on a reversed

dark/light cycle and experiments were performed during the dark cycle. The mice were housed in

same-sex groups until the week of the experiments, during which they were singly housed. Food

and water were provided *ad libitum* except during the experiment.

The estrous stage of female mice was evaluated based on vaginal smear samples obtained by

sterile vaginal lavage. Samples were collected using glass pipettes filled with double distilled

water, placed on a slide, stained using crystal violet, and coverslipped for microscopic

examination. Estrous stage was determined by the predominant cell type: squamous epithelial

cells (estrus), nucleated cornified cells (proestrus), or leukocytes (diestrus) (McLean et al.,

2012). To confirm that the stage of estrous did not change during the experiment day, samples

obtained prior to and after data collection on the experimental day were compared.

Experimental overview

The basic experimental structure that occurred over six days is illustrated in Figure 1D. Briefly,

each mouse was placed in an arena on Days 1 and 2 to provide one-time experiences of mating

and sustained restraint. After the experience on Day 2, the subject was anesthetized for

implantation of a guide tube for the microdialysis probe. On Day 6, the vocalization playback

session occurred. The mouse was briefly anesthetized for insertion of the microdialysis probe

into the amygdala, followed by several hours recovery. Before, during, and after vocal playback,

we sampled extracellular fluid from the amygdala through the microdialysis probe and recorded

video to analyze the subject's behavior. A liquid chromatographic/mass spectrometric method

was used to measure concentrations of several neurochemicals.

The microdialysis sampling interval of 10 mins was used to establish the temporal framework for

vocal playback and analysis of neurochemicals and behaviors (Fig. 1E). The analysis compared

neurochemicals and behaviors in a period immediately preceding the vocal stimuli (Pre-Stim)

and in two 10-min periods featuring playback of the same sequence of vocal stimuli (Stim 1 and

Stim 2). Each playback session featured only one category of vocalizations (associated with

mating or restraint) and each animal participated in a single playback session. Additional details

of this experimental structure are described in the sections below.

Acoustic methods

Vocalization recording and analysis

To record vocalizations for use in playback experiments, mice were placed in an open-topped

plexiglass chamber (width, 28 cm; length, 28 cm; height, 20 cm), housed within a darkened,

single-walled acoustic chamber (Industrial Acoustics, New York, NY) lined with anechoic foam

(Grimsley et al., 2016; Niemczura et al., 2020). Acoustic signals were recorded using ultrasonic

condenser microphones (CM16/CMPA, Avisoft Bioacoustics, Berlin, Germany) connected to a

multichannel amplifier and A/D converter (UltraSoundGate 416H, Avisoft Bioacoustics). The

gain of each microphone was independently adjusted once per recording session to optimize the

signal-to-noise ratio (SNR) while avoiding signal clipping. Acoustic signals were digitized at

500 kHz and 16-bit depth, monitored in real time with RECORDER software (Version 5.1,

Avisoft Bioacoustics), and Fast Fourier Transformed (FFT) at a resolution of 512 Hz. A night

vision camera (VideoSecu Infrared CCTV), centered 50 cm above the floor of the test box,

recorded the behaviors synchronized with the vocal recordings (VideoBench software,

DataWave Technologies, version 7).

To record mating vocalizations, ten animals (5 male-female pairs) were used in sessions that

lasted for 30 minutes. A male mouse was introduced first into the test box, followed by a female

mouse 5 min later. Vocalizations were recorded using two ultrasonic microphones placed 30 cm

above the floor of the recording box and 13 cm apart. See below for analysis of behaviors during

vocal recordings.

To record vocalizations during restraint, six mice (4 male, 2 female) were briefly anesthetized

with isoflurane and then placed in a restraint jacket as described previously (Grimsley et al.,

2016). Vocalizations were recorded for 30 minutes while the animal was suspended in the

recording box. Since these vocalizations are usually emitted at lower intensity compared to

mating vocalizations, the recording microphone was positioned 2-3 cm from the snout to obtain

the best SNR.

Vocal recordings were analyzed offline using Avisoft-SASLab Pro (version 5.2.12, Avisoft

Bioacoustics) with a hamming window, 1024 Hz FFT size, and an overlap percentage of 98.43.

For every syllable the channel with the higher amplitude signal was extracted using a custom-

written Python code and analyzed. Since automatic syllable tagging did not allow distinguishing

some syllable types such as noisy calls and mid-frequency vocalizations (MFVs) from

background noise, we manually tagged the start and end of each syllable, then examined

spectrograms to measure several acoustic features and classify syllable types based on Grimsley

et al. (Grimsley et al., 2011, 2016).

Vocalization playback

Vocalization playback lasting 20 min was constructed from a sequence of seven repeating stimulus blocks lasting 2:50 min each (Fig. 1E). Each block was composed of a set of vocal sequence exemplars that alternated with an equal duration of background sound ("Silence") associated with the preceding exemplar. The exemplars were recorded during mating interactions and restraint, and selected based on high SNR, correspondence with behavioral category by video analysis, and representation of the spectrotemporal features of vocalizations emitted during mating and restraint (Ghasemahmad, 2020; Grimsley et al., 2016). Mating stimulus blocks contained five exemplars of vocal sequences emitted during mating interactions. These exemplars ranged in duration from 15.0 – 43.6 s. Restraint stimulus blocks included seven vocal sequences, emitted by restrained male or female mice, with durations ranging from 5.7 – 42.3 s. Across exemplars, each stimulus block associated with both mating and restraint included different sets of vocal categories (Fig. 1A-C).

Playback sequences, i.e., exemplars, were conditioned in Adobe Audition CC (2018), adjusted to a 65 dB SNR level, then normalized to 1V peak-to-peak for the highest amplitude syllable in the sequence. This maintained relative syllable emission amplitude in the sequence. For each sequence, an equal duration of background noise (i.e., no vocal or other detected sounds) from the same recording was added at the end of that sequence (Fig. 1E). A 5-ms ramp was added at the beginning and the end of the entire sequence to avoid acoustic artifacts. A MATLAB app (EqualizIR, Sharad Shanbhag; https://github.com/TytoLogy/EqualizIR) compensated and calibrated each vocal sequence for the frequency response of the speaker system. Vocal sequences were converted to analog signals at 500 KHz and 16-bit resolution using DataWave (DataWave SciWorks, Loveland, CO), anti-alias filtered (TDT FT6-2, fc=125KHz), amplified (HCA-800II, Parasound, San Francisco, CA), and sent to the speaker (LCY, K100, Ying Tai

Audio Company, Hong Kong). Each sequence was presented at peak level equivalent to 85 dB

SPL.

Behavioral methods

Behaviors during both vocalization recording and playback sessions were recorded using a night

vision camera (480TVL 3.6mm, VideoSecu), centered 50 cm above the floor of the test box, and

SciWorks (DataWave, VideoBench version 7) for video acquisition and analysis.

Analysis of mating behaviors during vocal recordings

Interactions between male and female mice were video-recorded, then analyzed second-by-

second. Among more general courtship interactions, we identified a set of behaviors that are

mating-related, as described previously (Gaub et al., 2016; Heckman et al., 2016). All vocal

sequences selected as exemplars for playback of "mating" vocalizations were recorded in

association with these male mating behaviors: head-sniffing, attempted mounting, or mounting.

Vocalizations during these behaviors included chevron, stepped, and complex USVs emitted

with longer durations and higher repetition rates, and more LFH calls (Gaub et al., 2016;

Ghasemahmad, 2020; Hanson & Hurley, 2012).

Experience and playback sessions

Prior to playback experiments, each animal underwent 90-min sessions on two consecutive days

(Days 1 and 2) that provided both mating and restraint experiences to the EXP group (n = 31

animals) or no experiences of these to the INEXP group (n = 22 animals). EXP sessions were

presented in a counterbalanced pattern across subjects (Fig. 1D). For the mating experience,

mounting or attempted mounting was required for the animal to be included in the remainder of

the experiment. However, we did not record detailed behaviors or track estrous stage during the

mating experience session. After the Day 2 session, mice underwent surgery for implantation of

a microdialysis guide canula (see below), then recovered for four days.

On Day 6, the day of the playback experiment, male mice were randomly assigned to either

restraint or mating vocal playback groups. Females were only tested with mating playback, since

a preliminary behavioral study suggested no difference in male and female responses to restraint.

Video recording was performed simultaneously with microdialysis, beginning 10 min before

vocal playback (pre-stimulus) and continuing for 20 min during playback (Fig. 1D).

Analysis of behaviors during vocal playback

Behavioral analysis was based on previous descriptions (Bakshi & Kelley, 1993, 1994;

Blanchard et al., 2003; Füzesi et al., 2016; Grimsley et al., 2015; Lezak et al., 2017; Saibaba et

al., 1996) (Mouse Ethogram: https://mousebehavior.org). The list of behaviors assessed was

determined through pilot analysis and previous studies characterizing rodent defense and fear

behaviors. They are defined as the following: abrupt attending, flinching, locomotion, rearing,

grooming, still-and-alert, and stretch-attend (description in Table 1; (Blanchard et al., 2003)). A

behavior was only counted when its occurrence lasted two or more seconds, except for flinching,

which could take place more quickly.

For analysis of behaviors during vocal playback, all videos were examined blind to the sex of the

animal and the context of vocalizations. Video recordings were analyzed manually (n=48 mice)

in 10-second intervals for the seven behaviors identified above. The number of occurrences of

every behavior per 10-second block was marked. The numbers were added for every block of

2:50 min of vocal playback (see Fig. 1E), then averaged for the Pre-Stim, Stim 1, and Stim 2

periods separately.

Videos were further analyzed automatically using the video tracker within VideoBench

(DataWave Technologies, version 7) for total distance traveled and time spent in the periphery

and center of the test arena. For video-tracking analysis of time spent, the floor was divided into

16 equal squares with 4 central squares identified as "center" and the 4 corner squares identified

as periphery. The two squares in the middle of each wall were excluded from the analysis to

avoid attraction to the holes affect the animal's preference for the periphery vs center.

Procedures related to microdialysis

Surgery

Since both left and right amygdala are responsive to vocal stimuli in human and experimental

animal studies (Wenstrup et al., 2020), we implanted microdialysis probes into the left amygdala

to maintain consistency with other studies in our laboratory. Mice were anaesthetized with

Isoflurane (2-4%, Abbott Laboratories, North Chicago, IL) and hair overlying the skull was

removed using depilatory lotion. A midline incision was made, then the skin was moved laterally

to expose the skull. A craniotomy ($\simeq 1$ mm^2) was made above the basolateral amygdala (BLA)

on the left side (stereotaxic coordinates from bregma: -1.65 mm rostrocaudal, +3.43 mm

mediolateral). A guide cannula (CMA-7, CMA Microdialysis, Sweden) was implanted to a depth

of 2.6 mm below the cortical surface and above the left BLA (Fig. 1D, Day 2), then secured

using dental cement. After surgery, the animal received a subcutaneous injection of carprofen (4

mg/kg, s.c.) and topical anesthetic (lidocaine) and antibiotic cream (Neosporin). It was returned

to its cage and placed on a heating pad until fully recovered from anesthesia. Recovery from the

surgical procedures occurred in the animal's home cage and was assessed by resumption of

normal eating, drinking, exploratory, and grooming behaviors and postures. Playback

experiments occurred in all recovered animals four days after surgery.

Microdialysis

On the day before microdialysis, the probe was conditioned in 70% methanol and artificial

cerebrospinal fluid (aCSF) (CMA Microdialysis, Sweden). On playback day (Day 6), the animal

was briefly anesthetized and the probe with 1 mm membrane length and 0.24 mm outer diameter

(MWCO 6 kDa) was inserted into the guide cannula (Fig. 1D).

Using a spiral tubing connector (0.1 mm ID x 50 cm length) (CT-20, AMUZA Microdialysis,

Japan), the inlet and outlet tubing of the probe was connected to the inlet /outlet Teflon tubing of

the microdialysis lines. A swivel device for fluids (TCS-2-23, AMUZA Microdialysis, Japan),

secured to a balance arm, held the tubing, and facilitated the animal's free movement during the

experiment.

After probe insertion, a four-hour period allowed animals to habituate and the neurochemicals to

equilibrate between aCSF fluid in the probe and brain extracellular fluid (ECF). Sample

collection then began, in 10-min intervals, beginning with four background samples, then two

samples during playback of restraint or mating vocal sequences (total 20 minutes), then one or

more samples after playback ended. To account for the dead volume of the outlet tubing, a flow

rate of 1.069 µl/min was established at the syringe pump to obtain a 1 µl sample per minute.

Samples collected during this time were measured for volume to assure a consistent flow rate. To

prevent degradation of collected neurochemicals, the outlet tubing was passed through ice to a

site outside the sound-proof booth where samples were collected on ice, then stored in a −80°C

freezer. The choice of 10-min sampling intervals was based on the minimum collection time

needed to provide 5µl samples both for immediate testing and for a backup sample in the event

of catastrophic failure of the processes associated with the neurochemical analysis.

Neurochemical analysis

Samples were analyzed using a liquid chromatography (LC) / mass spectrometry (MS) technique

(Fig. 1D) at the Vanderbilt University Neurochemistry Core. This method allows simultaneous

measurement of several neurochemicals in the same dialysate sample: acetylcholine (ACh),

dopamine (DA) and its metabolites (3,4-Dihydroxyphenylacetic acid (DOPAC), and

homovanillic acid (HVA)), serotonin (5-HT) and its metabolite (5-hydroxyindoleacetic acid (5-

HIAA)), norepinephrine (NE), gamma aminobutyric acid (GABA), and glutamate. Due to low

recovery of NE and 5-HT from the mouse brain, we were unable to track these two

neuromodulators in this experiment.

Before each LC/MS analysis, 5 μl of the sample was derivatized using sodium carbonate,

benzoyl chloride in acetonitrile, and internal standard (Wong et al., 2016). LC was performed on

a 2.0 × 50 mm, 1.7 μM particle Acquity BEH C18 column (Waters Corporation, Milford, MA,

USA) with 1.5% aqueous formic acid as mobile phase A and acetonitrile as mobile phase B.

Using a Waters Acquity Classic UPLC, samples were separated by a gradient of 98–5 % of

mobile phase A over 11 min at a flow rate of 0.6 ml/min with delivery to a SCIEX 6500+ Qtrap

mass spectrometer (AB Sciex LLC, Framingham, MA, USA). The mass spectrometer was

operated using electrospray ionization in the positive ion mode. The capillary voltage and

temperature were 4KV and 350°C, respectively (Wong et al., 2016; Yohn et al., 2020).

Chromatograms were analyzed using MultiQuant 3.0.2 Software (AB SCIEX, Concord, Ontario,

Canada).

Verification of probe location

We verified placement of microdialysis probes to minimize variability that could arise due to

placement in regions surrounding BLA that receive different sources of neurochemical inputs

(e.g., cholinergic inputs to putamen and central amygdala). To verify probe placement within the

BLA, each probe was perfused with 2% dextran-fluorescein (MW 4Kda) (Sigma Aldrich Inc,

Atlanta, GA) at the end of the experiment. The location of the probe was then visualized in

adjacent cleared and Nissl-stained sections (e.g., Fig. 2, insets). Sections were photographed

using a SPOT RT3 camera and SPOT Advanced Plus imaging software (version 4.7) mounted on

a Zeiss Axio Imager M2 fluorescence microscope. Adobe Photoshop CS3 was used to adjust

brightness and contrast globally. Animals were included in statistical analyses only if $\geq$75% of

the probe membrane was located within the BLA (Fig. 2).

Data analysis

For neurochemical analyses, the total numbers of animals used were N_{EXP}=31 and N_{INEXP}=22.

Only mice with usable neurochemical data were further evaluated for behaviors. This included

46 mice: (N_{EXP}=25 and N_{INEXP}=21). Note that some animals used in neurochemical analyses

were not included in the behavioral analyses because their behavioral data were unavailable.

Since only one INEXP female was in a non-estrus stage during the playback session, our analysis

of the effect of experience included only estrus females and males. Further, one INEXP female

without detectable DA levels was removed from the INEXP group neurochemical analysis.

Only the following 10-min sample collection windows were used for statistical analysis: Pre-

Stim, Stim 1, and Stim 2 (Fig. 1E). The purpose of the two collection windows during vocal

playback was to allow detection of different epochs of neurochemical release, as may occur in

ACh release into the amygdala (Aitta-aho et al., 2018). All neurochemical data were normalized

to the background level, obtained from a single pre-stimulus sample immediately preceding

playback. This provided clarity in representations and did not result in different outcomes of

statistical tests compared to use of three pre-stimulus samples. The fluctuation in all background

samples was $\leq$20%. The percentage change from background level was calculated based on the

formula: *% change from background = (100 x stimulus sample concentration in pg)/ background*

sample concentration in pg. Values are represented as mean ± one standard error unless stated

otherwise. Raw neurochemical values are reported in supplementary tables 1 (ACh), 2 (DA), and

3 (5-HIAA).

All statistical analyses were performed using SPSS (IBM, V. 26 and 27). To examine

vocalizations in different stages of mating, we used a linear mixed model to analyze the changes

of interval and duration (dependent variables) of vocalizations based on mating interaction

intensity (fixed effect). For behavioral and neurochemical analyses in playback experiments, we

initially compared the output using a linear mixed model and a generalized linear model (GLM)

with a repeated measure for neurochemical data. Since both statistical methods resulted in

similar findings, we chose to use the GLM for statistical comparisons of the neurochemicals and

behaviors. Where Mauchly's test indicated that the assumption of sphericity had been violated,

degrees of freedom were corrected using Greenhouse Geisser estimates of sphericity.

To further clarify the differences observed between groups for every comparison in the GLM

repeated measure, multivariate contrast was performed. All multiple comparisons were corrected

using Bonferroni post hoc testing. 95% confidence intervals were used to compare values

between timepoints (Julious, 2004).

For both microdialysis and behavioral data, we tested the hypotheses that the release of

neurochemicals into the BLA and the number of behaviors is differently modulated by the vocal

playback type (mating and restraint) in male mice, by estrous stage of females or sex of the

animals during mating vocal playback, and by previous experience in any of these groups. In the

GLM model for Figure 3, context of vocalizations was defined as the fixed factor and time (Stim

1 and Stim 2) as the within-subject factor. The model in Figure 4 tested the effect of sex and

estrous state as a nested variable within sex as fixed factors, which were considered in the model

design (design = intercept + Sex + Estrous(sex)). The GLM model in Figure 5 tested sex,

context, and experience as fixed factors with the interaction terms of sex*experience and

context*experience. Here we avoided using a full factorial model due to the absence of females

in the restraint group, as that would have made the model unbalanced (design = intercept + Sex +

Context + Experience + Sex*Experience + Context*Experience). Further, estrous state was not

tested in examining the experience effect due to lack of INEXP non-estrus females. Dependent

(response) variables included the normalized concentration of ACh, DA, 5-HIAA in Stim 1 and

Stim 2 or the numbers of behaviors in Pre-Stim, Stim 1, and Stim 2 as previously described in

the behavioral analysis section.

Throughout the figures, the distributions of data points are visualized by box plots, computed

using the "inclusive median" formula in Excel. The box indicates first (Q1) and third (Q3)

quartiles with the median line between the two. The whiskers normally stretch between

minimum and maximum values, but this computation sometimes shows data points outside of the

whiskers. Excel identifies these values as outliers in the boxplot function. However, our GLM

statistical analysis includes these values because there was no compelling reason to exclude

them. The mean value is marked with an "x".

ACKNOWLEDGMENTS:

We are indebted to Sharad Shanbhag and Daniel Gavazzi for software used to condition vocal sequences and analyze audio and behavioral data, respectively, to Kristin Yeager for statistical consulting, and to Sheila Fleming for advice on behavioral assessments. We thank Anthony Zampino, Austin Poth, Debin Lei, and Krish Nair for technical assistance. We further thank Vanderbilt University Neurochemistry Core, supported by Vanderbilt Brain Institute and the Vanderbilt Kennedy Center, that performed the neurotransmitter sample analyses. We are grateful to Drs. Alexander Galazyuk, Brett Schofield, Merri Rosen, Mahtab Tehrani, and Sharad Shanbhag for their comments on previous versions of this manuscript.

FUNDING:

National Institutes of Health grant R01DC00937 (JJW, Alexander Galazyuk)

Kent State University Graduate Student Senate (ZG)

Northeast Ohio Medical University Biomedical Sciences Program Committee (ZG)

REFERENCES

Aitta-aho, T., Hay, Y. A., Phillips, B. U., Saksida, L. M., Bussey, T. J., Paulsen, O., & Apergis-Schoute, J. (2018a). Basal Forebrain and Brainstem Cholinergic Neurons Differentially Impact Amygdala Circuits and Learning-Related Behavior. *Current Biology*, *28*(16), 2557–2569. https://doi.org/10.1016/j.cub.2018.06.064

Aitta-aho, T., Hay, Y. A., Phillips, B. U., Saksida, L. M., Bussey, T. J., Paulsen, O., & Apergis-Schoute, J. (2018b). Basal Forebrain and Brainstem Cholinergic Neurons Differentially Impact Amygdala Circuits and Learning-Related Behavior. *Current Biology*, *28*(16), 2557-2569.e4. https://doi.org/10.1016/j.cub.2018.06.064

Altenmüller, E., Schmidt, S., & Zimmermann, E. (2013). *Evolution of Emotional Communication* (E. Altenmüller, S. Schmidt, & E. Zimmermann, Eds.; Edition One). Oxford University Press.

Ambroggi, F., Ishikawa, A., Fields, H. L., & Nicola, S. M. (2008). Basolateral Amygdala Neurons Facilitate Reward-Seeking Behavior by Exciting Nucleus Accumbens Neurons. *Neuron*, *59*(4), 648–661. https://doi.org/10.1016/j.neuron.2008.07.004

Arnold, M.R., Thallon, C.L., Pitkofsky, J.A., & Meerts, S.H. (2019) Sexual experience confers resilience to restraint stress in female rats. *Hormones and Behavior,* 107:61-66. doi: 10.1016/j.yhbeh.2018.12.003.

Asan, E. (1997). Ultrastructural features of tyrosine-hydroxylase-immunoreactive afferents and their targets in the rat amygdala. *Cell and Tissue Research*, *288*(3), 449–469. https://doi.org/10.1007/s004410050832

Asan, E. (1998). The catecholaminergic innervation of the rat amygdala. *Advances in Anatomy, Embryology, and Cell Biology*, *142*, 1–118. https://doi.org/10.1007/978-3-642-72085-7

Bakshi, V. P., & Kelley, A. E. (1993). Feeding induced by opioid stimulation of the ventral striatum: role of opiate receptor subtypes. *Journal of Pharmacology and Experimental Therapeutics*, *265*(3), 1253–1260.

Bakshi, V. P., & Kelley, A. E. (1994). Sensitization and conditioning of feeding following multiple morphine microinjections into the nucleus accumbens. *Brain Research*, *648*(2), 342–346. https://doi.org/10.1016/0006-8993(94)91139-8

Bargmann, C. I. (2012). Beyond the connectome: How neuromodulators shape neural circuits. *BioEssays*, *34*(6), 458–465. https://doi.org/10.1002/bies.201100185

Baysinger, A. N., Kent, B. A., & Brown, T. H. (2012). Muscarinic Receptors in Amygdala Control Trace Fear Conditioning. *PLoS ONE*, *7*(9). https://doi.org/10.1371/journal.pone.0045720

Beyeler, A., Namburi, P., Glober, G. F., Simonnet, C., Calhoon, G. G., Conyers, G. F., Luck, R., Wildes, C. P., & Tye, K. M. (2016). Divergent Routing of Positive and Negative Information from the Amygdala during Memory Retrieval. *Neuron*, *90*(2), 348–361. https://doi.org/10.1016/j.neuron.2016.03.004

Bigos, K. L., Pollock, B. G., Aizenstein, H. J., Fisher, P. M., Bies, R. R., & Hariri, A. R. (2008). Acute 5-HT reuptake blockade potentiates human amygdala reactivity. *Neuropsychopharmacology*, *33*(13), 3221–3225. https://doi.org/10.1038/npp.2008.52

Blanchard, D. C., Griebel, G., & Blanchard, R. J. (2003). The Mouse Defense Test Battery: pharmacological and behavioral assays for anxiety and panic. *European Journal of Pharmacology*, *463*(1–3), 97–116. https://doi.org/10.1016/S0014-2999(03)01276-7

Bocchio, M., McHugh, S. B., Bannerman, D. M., Sharp, T., & Capogna, M. (2016). Serotonin, amygdala and fear: Assembling the puzzle. *Frontiers in Neural Circuits*, *10*(APR), 1–15. https://doi.org/10.3389/fncir.2016.00024

Carlsen, J., Záborszky, L., & Heimer, L. (1985). Cholinergic projections from the basal forebrain to the basolateral amygdaloid complex: A combined retrograde fluorescent and immunohistochemical study. *Journal of Comparative Neurology*, *234*(2), 155–167. https://doi.org/10.1002/cne.902340203

Ciocchi, S., Herry, C., Grenier, F., Wolff, S. B. E., Letzkus, J. J., Vlachos, I., Ehrlich, I., Sprengel, R., Deisseroth, K., Stadler, M. B., Müller, C., & Lüthi, A. (2010). Encoding of conditioned fear in central amygdala inhibitory circuits. *Nature*, *468*(7321), 277–282. https://doi.org/10.1038/nature09559

Darwin, C. (1872). *The Expression of the Emotions in Man and Animals*. John Murray.

Davis, M., Walker, D. L., Miles, L., & Grillon, C. (2010). Phasic vs sustained fear in rats and humans: Role of the extended amygdala in fear vs anxiety. In *Neuropsychopharmacology* (Vol. 35, Issue 1, pp. 105–135). https://doi.org/10.1038/npp.2009.109

Di Ciano, P., & Everitt, B. J. (2004). Direct interactions between the basolateral amygdala and nucleus accumbens core underlie cocaine-seeking behavior by rats. *Journal of Neuroscience*, *24*(32), 7167–7173. https://doi.org/10.1523/JNEUROSCI.1581-04.2004

Dornellas, A. P. S., Burnham, N. W., Luhn, K. L., Petruzzi, M. V., Thiele, T. E., & Navarro, M. (2021). Activation of locus coeruleus to rostromedial tegmental nucleus (RMTg) noradrenergic pathway blunts binge-like ethanol drinking and induces aversive responses in mice. *Neuropharmacology*, *199*. https://doi.org/10.1016/j.neuropharm.2021.108797

Egozi, Y., Kloog, Y., & Sokolovsky, M. (1986). Acetylcholine rhythm in the preoptic area of the rat hypothalamus is synchronized with the estrous cycle. *Brain Research*, *383*(1–2), 310–313. https://doi.org/10.1016/0006-8993(86)90030-2

Finton, C. J., Keesom, S. M., Hood, K. E., & Hurley, L. M. (2017). What's in a squeak? Female vocal signals predict the sexual behaviour of male house mice during courtship. *Animal Behaviour*, *126*, 163–175. https://doi.org/10.1016/j.anbehav.2017.01.021

Frühholz, S., Trost, W., & Kotz, S. A. (2016). The sound of emotions-Towards a unifying neural network perspective of affective sound processing. In *Neuroscience and Biobehavioral Reviews* (Vol. 68, pp. 96–110). Neurosci Biobehav Rev. https://doi.org/10.1016/j.neubiorev.2016.05.002

Füzesi, T., Daviu, N., Wamsteeker Cusulin, J. I., Bonin, R. P., & Bains, J. S. (2016). Hypothalamic CRH neurons orchestrate complex behaviours after stress. *Nature Communications*, *7*(1), 11937. https://doi.org/10.1038/ncomms11937

Gadziola, M. A., Grimsley, J. M. S., Shanbhag, S. J., & Wenstrup, J. J. (2012). A novel coding mechanism for social vocalizations in the lateral amygdala. *Journal of Neurophysiology*, *107*(4), 1047–1057. https://doi.org/10.1152/jn.00422.2011

Gadziola, M. A., Shanbhag, S. J., & Wenstrup, J. J. (2016). Two distinct representations of social vocalizations in the basolateral amygdala. *Journal of Neurophysiology*, *115*(2), 868–886. https://doi.org/10.1152/jn.00953.2015

Gaub, S., Fisher, S. E., & Ehret, G. (2016). Ultrasonic vocalizations of adult male Foxp2-mutant mice: behavioral contexts of arousal and emotion. *Genes, Brain and Behavior*, *15*(2), 243–259. https://doi.org/10.1111/gbb.12274

Ghasemahmad, Z. (2020). *Behavioral and neuromodulatory responses to emotional vocalizations in mice* [Kent State University]. https://www.proquest.com/dissertations-

theses/behavioral-neuromodulatory-responses-emotional/docview/2572567360/se-2?accountid=36418

Gibbs, R. B. (1996). Fluctuations in relative levels of choline acetyltransferase mRNA in different regions of the rat basal forebrain across the estrous cycle: Effects of estrogen and progesterone. *Journal of Neuroscience*, *16*(3), 1049–1055. https://doi.org/10.1523/jneurosci.16-03-01049.1996

Gorka, A. X., Knodt, A. R., & Hariri, A. R. (2015). Basal forebrain moderates the magnitude of task-dependent amygdala functional connectivity. *Social Cognitive and Affective Neuroscience*, *10*(4), 501–507. https://doi.org/10.1093/scan/nsu080

Grimsley, C. A., Longenecker, R. J., Rosen, M. J., Young, J. W., Grimsley, J. M., & Galazyuk, A. V. (2015). An improved approach to separating startle data from noise. *Journal of Neuroscience Methods*, *253*, 206–217. https://doi.org/10.1016/j.jneumeth.2015.07.001

Grimsley, J. M. S., Hazlett, E. G., & Wenstrup, J. J. (2013). Coding the meaning of sounds: Contextual modulation of auditory responses in the basolateral amygdala. *Journal of Neuroscience*, *33*(44), 17538–17548. https://doi.org/10.1523/JNEUROSCI.2205-13.2013

Grimsley, J. M. S., Monaghan, J. J. M., & Wenstrup, J. J. (2011). Development of Social Vocalizations in Mice. *PLoS ONE*, *6*(3), e17460. https://doi.org/10.1371/journal.pone.0017460

Grimsley, J. M. S., Sheth, S., Vallabh, N., Grimsley, C. A., Bhattal, J., Latsko, M., Jasnow, A., & Wenstrup, J. J. (2016a). Contextual modulation of vocal behavior in mouse: Newly identified 12 kHz "Mid-frequency" vocalization emitted during restraint. *Frontiers in Behavioral Neuroscience*, *10*(MAR). https://doi.org/10.3389/fnbeh.2016.00038

Grimsley, J. M. S., Sheth, S., Vallabh, N., Grimsley, C. A., Bhattal, J., Latsko, M., Jasnow, A., & Wenstrup, J. J. (2016b). Contextual modulation of vocal behavior in mouse: Newly identified 12 kHz "Mid-frequency" vocalization emitted during restraint. *Frontiers in Behavioral Neuroscience*, *10*(MAR). https://doi.org/10.3389/fnbeh.2016.00038

Gründemann, J., Bitterman, Y., Lu, T., Krabbe, S., Grewe, B. F., Schnitzer, M. J., & Lüthi, A. (2019a). Amygdala ensembles encode behavioral states. *Science*, *364*(6437). https://doi.org/10.1126/science.aav8736

Gründemann, J., Bitterman, Y., Lu, T., Krabbe, S., Grewe, B. F., Schnitzer, M. J., & Lüthi, A. (2019b). Amygdala ensembles encode behavioral states. *Science*, *364*(6437). https://doi.org/10.1126/science.aav8736

Hanson, J. L., & Hurley, L. M. (2012). Female presence and estrous state influence mouse ultrasonic courtship vocalizations. *PLoS ONE*, *7*(7). https://doi.org/10.1371/journal.pone.0040782

Heckman, J., McGuinness, B., Celikel, T., & Englitz, B. (2016). Determinants of the mouse ultrasonic vocal structure and repertoire. In *Neuroscience and Biobehavioral Reviews* (Vol. 65, pp. 313–325). Neurosci Biobehav Rev. https://doi.org/10.1016/j.neubiorev.2016.03.029

Huang, S. Y., Treviño, M., He, K., Ardiles, A., DePasquale, R., Guo, Y., Palacios, A., Huganir, R., & Kirkwood, A. (2012). Pull-Push neuromodulation of LTP and LTD enables bidirectional experience-induced synaptic scaling in visual cortex. *Neuron*, *73*(3), 497–510. https://doi.org/10.1016/j.neuron.2011.11.023

Inagaki, R. T., Raghuraman, S., Chase, K., Steele, T., Zornik, E., Olivera, B., & Yamaguchi, A. (2020). Molecular characterization of frog vocal neurons using constellation pharmacology. *Journal of Neurophysiology*, *123*(6), 2297–2310. https://doi.org/10.1152/jn.00105.2020

Jiang, L., Kundu, S., Lederman, J. D. D., López-Hernández, G. Y. Y., Ballinger, E. C. C., Wang, S., Talmage, D. A. A., & Role, L. W. W. (2016). Cholinergic Signaling Controls Conditioned Fear Behaviors and Enhances Plasticity of Cortical-Amygdala Circuits. *Neuron*, *90*(5), 1057–1070. https://doi.org/10.1016/j.neuron.2016.04.028

Jing, M., Zhang, P., Wang, G., Feng, J., Mesik, L., Zeng, J., Jiang, H., Wang, S., Looby, J. C., Guagliardo, N. A., Langma, L. W., Lu, J., Zuo, Y., Talmage, D. A., Role, L. W., Barrett, P. Q., Zhang, L. I., Luo, M., Song, Y., … Li, Y. (2018). A genetically encoded fluorescent acetylcholine indicator for in vitro and in vivo studies. *Nature Biotechnology*, *36*(8), 726–737. https://doi.org/10.1038/nbt.4184

Julious, S. A. (2004). Using confidence intervals around individual means to assess statistical significance between two means. *Pharmaceutical Statistics*, *3*(3), 217–222. https://doi.org/10.1002/pst.126

Kalinowski, D., Bogus-Nowakowska, K., Kozłowska, A., & Równiak, M. (2023). Dopaminergic and cholinergic modulation of the amygdala is altered in female mice with oestrogen receptor β deprivation. *Scientific Reports*, *13*(1), 897. https://doi.org/10.1038/s41598-023-28069-2

Kirry, A. J., Durigan, D. J., Twining, R. C., & Gilmartin, M. R. (2019). Estrous cycle stage gates sex differences in prefrontal muscarinic control of fear memory formation. *Neurobiology of Learning and Memory*, *161*, 26–36. https://doi.org/10.1016/j.nlm.2019.03.001

Kröner, S., Rosenkranz, J. A., Grace, A. A., & Barrionuevo, G. (2005). Dopamine modulates excitability of basolateral amygdala neurons in vitro. *Journal of Neurophysiology*, *93*(3), 1598–1610. https://doi.org/10.1152/JN.00843.2004

Kyriazi, P., Headley, D. B., & Pare, D. (2018). Multi-dimensional Coding by Basolateral Amygdala Neurons. *Neuron*, *99*(6), 1315-1328.e5. https://doi.org/10.1016/j.neuron.2018.07.036

Lanuza, E., Nader, K., & Ledoux, J. E. (2004). Unconditioned stimulus pathways to the amygdala: effects of posterior thalamic and cortical lesions on fear conditioning. *Neuroscience*, *125*(2), 305–315. https://doi.org/10.1016/j.neuroscience.2003.12.034

Leuner, B., Glasper, E.R., & Gould, E. (2010) Sexual experience promotes adult neurogenesis in the hippocampus despite an initial elevation in stress hormones. *PLoS One*, 5(7):e11597. doi: 10.1371/journal.pone.0011597.

LeDoux, J. (2003). The emotional brain, fear, and the amygdala. *Cellular and Molecular Neurobiology*, *23*(4/5), 727–738. https://doi.org/10.1023/A:1025048802629

LeDoux, J. E. (2000). Emotion Circuits in the Brain. *Annual Review of Neuroscience*, *23*(1), 155–184. https://doi.org/10.1146/annurev.neuro.23.1.155

LeDoux, J., Sakaguchi, A., & Reis, D. (1984). Subcortical efferent projections of the medial geniculate nucleus mediate emotional responses conditioned to acoustic stimuli. *The Journal of Neuroscience*, *4*(3), 683–698. https://doi.org/10.1523/JNEUROSCI.04-03-00683.1984

Lester, D. B., Rogers, T. D., & Blaha, C. D. (2010). Acetylcholine-dopamine interactions in the pathophysiology and treatment of CNS disorders. *CNS Neuroscience and Therapeutics*, *16*(3), 137–162. https://doi.org/10.1111/j.1755-5949.2010.00142.x

Lezak, K. R., Missig, G., & Carlezon Jr, W. A. (2017). Behavioral methods to study anxiety in rodents. *Dialogues in Clinical Neuroscience*, *19*(2), 181–191. https://doi.org/10.31887/DCNS.2017.19.2/wcarlezon

Liebenthal, E., Silbersweig, D. A., & Stern, E. (2016). The language, tone and prosody of emotions: Neural substrates and dynamics of spoken-word emotion perception. In *Frontiers in Neuroscience* (Vol. 10, Issue NOV, p. 506). Frontiers Media S.A. https://doi.org/10.3389/fnins.2016.00506

Likhtik, E., & Johansen, J. P. (2019a). Neuromodulation in circuits of aversive emotional learning. In *Nature Neuroscience* (Vol. 22, Issue 10, pp. 1586–1597). Nature Publishing Group. https://doi.org/10.1038/s41593-019-0503-3

Likhtik, E., & Johansen, J. P. (2019b). Neuromodulation in circuits of aversive emotional learning. *Nature Neuroscience*, *22*(10), 1586–1597. https://doi.org/10.1038/s41593-019-0503-3

Lim, S. L., Padmala, S., & Pessoa, L. (2009). Segregating the significant from the mundane on a moment-to-moment basis via direct and indirect amygdala contributions. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(39), 16841–16846. https://doi.org/10.1073/pnas.0904551106

Lutas, A., Kucukdereli, H., Alturkistani, O., Carty, C., Sugden, A. U., Fernando, K., Diaz, V., Flores-Maldonado, V., & Andermann, M. L. (2019). State-specific gating of salient cues by midbrain dopaminergic input to basal amygdala. *Nature Neuroscience*, *22*(11), 1820–1833. https://doi.org/10.1038/s41593-019-0506-0

Matsuda, Y., Hirano, H., & Watanabe, Y. (2002). Effects of estrogen on acetylcholine release in frontal cortex of female rats: Involvement of serotonergic neuronal systems. *Brain Research*, *937*(1–2), 58–65. https://doi.org/10.1016/S0006-8993(02)02465-4

Matsumoto, J., Nishimaru, H., Takamura, Y., Urakawa, S., Ono, T., & Nishijo, H. (2016). Amygdalar auditory neurons contribute to self-other distinction during ultrasonic social vocalization in rats. *Frontiers in Neuroscience*, *10*(SEP). https://doi.org/10.3389/fnins.2016.00399

McDonald, A. J. (1998). Cortical pathways to the mammalian amygdala. *Progress in Neurobiology*, *55*(3), 257–332. https://doi.org/10.1016/S0301-0082(98)00003-3

McEwen, B. S. (1998). Multiple ovarian hormone effects on brain structure and function. *The Journal of Gender-Specific Medicine: JGSM: The Official Journal of the Partnership for Women's Health at Columbia*, *1*(1), 33–41.

McLean, A. C., Valenzuela, N., Fai, S., & Bennett, S. A. L. (2012). Performing vaginal lavage, crystal violet staining, and vaginal cytological evaluation for mouse estrous cycle staging identification. *Journal of Visualized Experiments : JoVE*, *67*. https://doi.org/10.3791/4389

Mesulam, M. M., Mufson, E. J., Wainer, B. H., & Levey, A. I. (1983). Central cholinergic pathways in the rat: An overview based on an alternative nomenclature (Ch1-Ch6). *Neuroscience*, *10*(4), 1185–1201. https://doi.org/10.1016/0306-4522(83)90108-2

Mizuno, I., Matsuda, S., Tohyama, S., & Mizutani, A. (2022). The role of the cannabinoid system in fear memory and extinction in male and female mice. *Psychoneuroendocrinology*, *138*. https://doi.org/10.1016/j.psyneuen.2022.105688

Nadim, F., & Bucher, D. (2014). Neuromodulation of neurons and synapses. *Current Opinion in Neurobiology*, *29*, 48–56. https://doi.org/10.1016/J.CONB.2014.05.003

Namburi, P., Al-Hasani, R., Calhoon, G. G., Bruchas, M. R., & Tye, K. M. (2016). Architectural Representation of Valence in the Limbic System. *Neuropsychopharmacology*, *41*(7), 1697–1715. https://doi.org/10.1038/npp.2015.358

Namburi, P., Beyeler, A., Yorozu, S., Calhoon, G. G., Halbert, S. A., Wichmann, R., Holden, S. S., Mertens, K. L., Anahtar, M., Felix-Ortiz, A. C., Wickersham, I. R., Gray, J. M., & Tye,

K. M. (2015a). A circuit mechanism for differentiating positive and negative associations. *Nature*, *520*(7549), 675–678. https://doi.org/10.1038/nature14366

Namburi, P., Beyeler, A., Yorozu, S., Calhoon, G. G., Halbert, S. A., Wichmann, R., Holden, S. S., Mertens, K. L., Anahtar, M., Felix-Ortiz, A. C., Wickersham, I. R., Gray, J. M., & Tye, K. M. (2015b). A circuit mechanism for differentiating positive and negative associations. *Nature*, *520*(7549), 675–678. https://doi.org/10.1038/nature14366

Neunuebel, J. P., Taylor, A. L., Arthur, B. J., & Egnor, S. R. (2015). Female mice ultrasonically interact with males during courtship displays. *ELife*, *4*. https://doi.org/10.7554/eLife.06203

Niemczura, A. C., Grimsley, J. M., Kim, C., Alkhawaga, A., Poth, A., Carvalho, A., & Wenstrup, J. J. (2020). Physiological and Behavioral Responses to Vocalization Playback in Mice. *Frontiers in Behavioral Neuroscience*, *14*, 155. https://doi.org/10.3389/fnbeh.2020.00155

O'Neill, P. K., Gore, F., & Salzman, C. D. (2018). Basolateral amygdala circuitry in positive and negative valence. *Current Opinion in Neurobiology*, *49*, 175–183. https://doi.org/10.1016/j.conb.2018.02.012

Otani, S., Daniel, H., Roisin, M. P., & Crepel, F. (2003). Dopaminergic Modulation of Long-term Synaptic Plasticity in Rat Prefrontal Neurons. *Cerebral Cortex*, *13*(11), 1251–1256. https://doi.org/10.1093/cercor/bhg092

Paton, J. J., Belova, M. A., Morrison, S. E., & Salzman, C. D. (2006). The primate amygdala represents the positive and negative value of visual stimuli during learning. *Nature*, *439*(7078), 865–870. https://doi.org/10.1038/nature04490

Pawlak, V., Wickens, J. R., Kirkwood, A., & Kerr, J. N. D. (2010). Timing is not everything: Neuromodulation opens the STDP gate. *Frontiers in Synaptic Neuroscience*, *2*(OCT). https://doi.org/10.3389/fnsyn.2010.00146

Peterson, D. C., & Wenstrup, J. J. (2012). Selectivity and persistent firing responses to social vocalizations in the basolateral amygdala. *Neuroscience*, *217*, 154–171. https://doi.org/10.1016/j.neuroscience.2012.04.069

Pidoplichko, V. I., Prager, E. M., Aroniadou-Anderjaska, V., & Braga, M. F. M. (2013). α7-Containing nicotinic acetylcholine receptors on interneurons of the basolateral amygdala and their role in the regulation of the network excitability. *Journal of Neurophysiology*, *110*(10), 2358–2369. https://doi.org/10.1152/jn.01030.2012

Pignatelli, M., & Beyeler, A. (2019). Valence coding in amygdala circuits. *Current Opinion in Behavioral Sciences*, *26*, 97–106. https://doi.org/10.1016/j.cobeha.2018.10.010

Rojas-Carvajal, M., Chinchilla-Alvarado, J., & Brenes, J. C. (2022). Muscarinic regulation of self-grooming behavior and ultrasonic vocalizations in the context of open-field habituation in rats. *Behavioural Brain Research*, *418*, 113641. https://doi.org/10.1016/j.bbr.2021.113641

Romanski, L. M., & LeDoux, J. E. (1993). Information Cascade from Primary Auditory Cortex to the Amygdala: Corticocortical and Corticoamygdaloid Projections of Temporal Cortex in the Rat. *Cerebral Cortex*, *3*(6), 515–532. https://doi.org/10.1093/cercor/3.6.515

Sah, P., Faber, E. S. L., Lopez De Armentia, M., & Power, J. (2003). *The Amygdaloid Complex: Anatomy and Physiology*. https://doi.org/10.1152/physrev.00002.2003.-A

Saibaba, P., Sales, G. D., Stodulski, G., & Hau, J. (1996). Behaviour of rats in their home cages: daytime variations and effects of routine husbandry procedures analysed by time sampling techniques. *Laboratory Animals*, *30*(1), 13–21. https://doi.org/10.1258/002367796780744875

Sales (née Sewell), G. D. (1972). Ultrasound and mating behaviour in rodents with some observations on other behavioural situations. *Journal of Zoology*, *168*(2), 149–164. https://doi.org/10.1111/j.1469-7998.1972.tb01345.x

Sander, D., Grafman, J., & Zalla, T. (2003). The Human Amygdala: An Evolved System for Relevance Detection. *Reviews in the Neurosciences*, *14*(4), 303–316. https://doi.org/10.1515/REVNEURO.2003.14.4.303

Schofield, B. R., & Hurley, L. (2018). *Circuits for Modulation of Auditory Function* (pp. 235–267). https://doi.org/10.1007/978-3-319-71798-2_9

Schönfeld, L.-M., Zech, M.-P., Schäble, S., Wöhr, M., & Kalenscher, T. (2020). Lesions of the rat basolateral amygdala reduce the behavioral response to ultrasonic vocalizations. *Behavioural Brain Research*, *378*, 112274. https://doi.org/10.1016/j.bbr.2019.112274

Seyfarth, R. M., & Cheney, D. L. (2003). Meaning and Emotion in Animal Vocalizations. *Annals of the New York Academy of Sciences*, *1000*, 32–55. https://doi.org/10.1196/annals.1280.004

Shi, C. J., & Cassell, M. D. (1997). Cortical, thalamic, and amygdaloid projections of rat temporal cortex. *Journal of Comparative Neurology*, *382*(2), 153–175. https://doi.org/10.1002/(SICI)1096-9861(19970602)382:2<153::AID-CNE2>3.0.CO;2-2

Shughrue, P. J., Scrimo, P. J., & Merchenthaler, I. (2000). Estrogen binding and estrogen receptor characterization (ERα and ERβ) in the cholinergic neurons of the rat basal forebrain. *Neuroscience*, *96*(1), 41–49. https://doi.org/10.1016/S0306-4522(99)00520-5

Silkstone, M., & Brudzynski, S. M. (2020). Dissimilar interaction between dopaminergic and cholinergic systems in the initiation of emission of 50-kHz and 22-kHz vocalizations. *Pharmacology Biochemistry and Behavior*, *188*. https://doi.org/10.1016/j.pbb.2019.172815

Smith, D. M., & Torregrossa, M. M. (2021a). Valence encoding in the amygdala influences motivated behavior. *Behavioural Brain Research*, *411*. https://doi.org/10.1016/j.bbr.2021.113370

Smith, D. M., & Torregrossa, M. M. (2021b). Valence encoding in the amygdala influences motivated behavior. *Behavioural Brain Research*, *411*, 113370. https://doi.org/10.1016/j.bbr.2021.113370

Stuber, G. D., Sparta, D. R., Stamatakis, A. M., Van Leeuwen, W. A., Hardjoprajitno, J. E., Cho, S., Tye, K. M., Kempadoo, K. A., Zhang, F., Deisseroth, K., & Bonci, A. (2011). Excitatory transmission from the amygdala to nucleus accumbens facilitates reward seeking. *Nature*, *475*(7356), 377–382. https://doi.org/10.1038/nature10194

Tingley, D., Alexander, A. S., Kolbu, S., de Sa, V. R., Chiba, A. A., & Nitz, D. A. (2014). Task-phase-specific dynamics of basal forebrain neuronal ensembles. *Frontiers in Systems Neuroscience*, *8*(SEP). https://doi.org/10.3389/fnsys.2014.00174

Tsukano, H., Hou, X., Horie, M., Kitaura, H., Nishio, N., Hishida, R., Takahashi, K., Kakita, A., Takebayashi, H., Sugiyama, S., & Shibuki, K. (2019). Reciprocal connectivity between secondary auditory cortical field and amygdala in mice. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-56092-9

Unal, C. T., Pare, D., & Zaborszky, L. (2015). Impact of basal forebrain cholinergic inputs on basolateral amygdala neurons. *Journal of Neuroscience*, *35*(2), 853–863. https://doi.org/10.1523/JNEUROSCI.2706-14.2015

van Huizen, F., March, D., Cynader, M. S., & Shaw, C. (1994). Muscarinic Receptor Characteristics and Regulation in Rat Cerebral Cortex: Changes during Development,

Aging and the Oestrous Cycle. *European Journal of Neuroscience*, *6*(2), 237–243. https://doi.org/10.1111/j.1460-9568.1994.tb00266.x

van Vugt, B., van Kerkoerle, T., Vartak, D., & Roelfsema, P. R. (2020). The contribution of AMPA and NMDA receptors to persistent firing in the dorsolateral prefrontal cortex in working memory. *Journal of Neuroscience*, *40*(12), 2458–2470. https://doi.org/10.1523/JNEUROSCI.2121-19.2020

Vander Weele, C. M., Siciliano, C. A., Matthews, G. A., Namburi, P., Izadmehr, E. M., Espinel, I. C., Nieh, E. H., Schut, E. H. S., Padilla-Coreano, N., Burgos-Robles, A., Chang, C. J., Kimchi, E. Y., Beyeler, A., Wichmann, R., Wildes, C. P., & Tye, K. M. (2018). Dopamine enhances signal-to-noise ratio in cortical-brainstem encoding of aversive stimuli. *Nature*, *563*(7731), 397–401. https://doi.org/10.1038/s41586-018-0682-1

Wall, E. M., & Woolley, S. C. (2020). Acetylcholine in action. *ELife*, *9*. https://doi.org/10.7554/eLife.57515

Wenstrup, J. J., Ghasemahmad, Z., Hazlett, E., & Shanbhag, S. J. (2020). The Amygdala – A Hub of the Social Auditory Brain. In *The Senses: A Comprehensive Reference* (pp. 812–837). Elsevier. https://doi.org/10.1016/b978-0-12-809324-5.24194-1

Wiethoff, S., Wildgruber, D., Grodd, W., & Ethofer, T. (2009). Response and habituation of the amygdala during processing of emotional prosody. *NeuroReport*, *20*(15), 1356–1360. https://doi.org/10.1097/WNR.0b013e328330eb83

Wong, J. M. T., Malec, P. A., Mabrouk, O. S., Ro, J., Dus, M., & Kennedy, R. T. (2016). Benzoyl chloride derivatization with liquid chromatography-mass spectrometry for targeted metabolomics of neurochemicals in biological samples. *Journal of Chromatography A*, *1446*, 78–90. https://doi.org/10.1016/j.chroma.2016.04.006

Yohn, S. E., Foster, D. J., Covey, D. P., Moehle, M. S., Galbraith, J., Garcia-Barrantes, P. M., Cho, H. P., Bubser, M., Blobaum, A. L., Joffe, M. E., Cheer, J. F., Jones, C. K., Lindsley, C. W., & Conn, P. J. (2020). Activation of the mGlu1 metabotropic glutamate receptor has antipsychotic-like effects and is required for efficacy of M4 muscarinic receptor allosteric modulators. *Molecular Psychiatry*, *25*(11), 2786–2799. https://doi.org/10.1038/s41380-018-0206-2

Zhang, X., & Li, B. (2018). Population coding of valence in the basolateral amygdala. *Nature Communications*, *9*(1). https://doi.org/10.1038/s41467-018-07679-9

Figure 1

Figure 1. Behavioral / microdialysis experiments test how playback of affective vocal
signals alters behaviors and neuromodulator release into the BLA. A. Short sample of
mating vocal sequence used in playback experiments. Recording was obtained during high-
intensity mating interactions and included ultrasonic vocalizations (USVs), likely emitted by the
male, as well as low frequency harmonic (LFH) calls likely emitted by the female. **B.** Short
sample of restraint vocal sequence used in playback experiments. Recording was obtained from
an isolated, restrained mouse (see Material and Methods) and consisted mostly of mid-frequency
vocalization (MFV) syllables. **C.** Syllable types in mating playback sequences differ
substantially from those in restraint playback sequences. Percentages indicate frequency-of-
occurrence of syllable types across all examplars used in mating or restraint vocal stimuli (n_{Mating}

= 545, $n_{Restraint}$ = 622 syllables). **D**. Experimental design in playback experiment. Days 1 and 2: each animal experienced restraint and mating behaviors (counterbalanced order across subjects). Day 2: a microdialysis guide tube was implanted in the brain above the BLA. Day 6: the microdialysis probe was inserted through guide tube into the BLA. Playback experiments began after several hours of habituation/equilibration. Behavioral observations and microdialysis sampling were obtained before, during, and after playback of one vocalization type. Microdialysis samples were analyzed using LC/MS method described in Materials and Methods. **E.** Schematic illustration of detailed sequencing of vocal stimuli, shown here for mating playback. A 20-min period of vocal playback was formed by seven repeated stimulus blocks of 170 s. The stimulus blocks were composed of five vocal exemplars (each represented by a different color) of variable length, with each exemplar followed by an equal duration of silence. Stimuli during the Stim1 and Stim 2 playback windows thus included identical blocks but in slightly altered patterns. See Material and Methods for in-depth description of vocalization playback.

1 **Figure 2**
2

EXP Groups

INEXP Groups

3
4
Figure 2. Microdialysis probe locations for EXP and INEXP Groups. Labels above BLA
outlines indicate groups based on playback type, sex, and hormonal state. Colored lines indicate
recovered probe tracks that resulted from infusion of fluorescent tracers. Black solid lines
indicate external capsule; black dashed lines indicate major amygdalar subdivisions. Arrows
indicate tracks related to insets. Insets: photomicrographs shows dextran-fluorescein labeling that
marks the placement of the microdialysis probes for two cases, 1340 and 1404. Abbreviations: *B,*
basal nucleus of amygdala; *CeA*, central nucleus of amygdala; *La*, lateral nucleus of amygdala.
12
13

1   **Figure 3**
2

3
4
Figure 3. Behavioral and neuromodulator responses to vocal playback in male mice differ
by behavioral context of vocalizations. A, B. Boxplots show number of occurrences of
specified behavior in 10-min observation periods before (Pre-Stim) and during (Stim 1, Stim 2)
playback of mating or restraint vocal sequences (n_{Mating} = 7, $n_{Restraint}$ = 6). Note that playback
sequences during Stim 1 and Stim 2 periods were identical within each group. During playback,
the restraint group increased still-and-alert behavior compared to the mating group **(A)** (context:
F $(1,11)$ = 9.6, p=0.01, η^2=0.5), and increased flinching behavior **(B)** (time*context: F $(1.1,12.2)$
=6.3, p=0.02, η^2=0.4) compared to the Pre-Stim baseline and to the mating group. **C-E.** Boxplots
show differential release of acetylcholine (ACh), dopamine (DA), and 5-HIAA relative to the
Pre-Stim period, during mating and restraint vocal playback (n_{Mating} = 9, $n_{Restraint}$ = 7). **C.**
Significant differences in ACh for restraint (increase) playback in Stim 1 and Stim 2 vs males in
mating playback (Main effect of context: F $(1,14)$ = 22.6,p<0.001, η^2=0.62). **D.** Significant
differences for DA for mating (increase) vs restraint playback (Main effect of context: F $(1,14)$ =

7.4, p=0.02, η^2=0.35). **E**. No significant changes in 5-HIAA during vocal playback in male mice (context: F (1,14) = 1.36, p=0.3, η^2 = 0.09). **A-E.** Statistical testing examined time windows within groups for behavioral observations and all intergroup comparisons within time windows for both behavior and neuromodulator data. Only normalized neuromodulators in Stim 1 and Stim 2 were used for statistical comparisons. Only significant tests are shown. Repeated measures GLM: *p<0.05, **p<0.01, ***p<0.001 (Bonferroni post hoc test). Time windows comparison: 95% confidence intervals. See Data Analysis section in *Materials and Methods* for description of box plots.

1 **Figure 4**

Figure 4

Figure 4. Behavioral and neuromodulator responses to playback of mating vocal sequences
differ by sex and female estrous stage. A-D. Occurrences of specified behaviors in 10-min
periods before (Pre-Stim) and during (Stim 1, Stim 2) mating vocal playback (n_{Male} = 7, $n_{Estrus\ Fem}$
= 6, $n_{Non-estrus\ Fem}$ = 5). **A.** Attending behavior increased during Stim 1 in response to mating vocal
playback, regardless of sex or estrous stage (time: $F_{(2,30)}$ = 32.6, $p<0.001$, $\eta2=0.7$; time*sex: F
$(2,30)$ = 0.12; $p=0.9$, $\eta2=0.008$; time*estrous: $F_{(2,30)}$ = 1.1; $p=0.4$, $\eta2=0.07$). **B-C.** Females
regardless of estrous stage reared less (sex: $F_{(1,15)}$ =10.22, $p=0.006$; $\eta^2=0.4$; estrous: F (1,15)
=0.2, $p=0.7$, $\eta^2=0.01$) and-displayed more Still-and-Alert behaviors (Sex: $F_{(1,15)}$ =5.2, $p=0.04$,
partial $\eta2$= 0.3, estrous: $(1,15)$ =0.07, $p=0.8$, partial $\eta2$= 0.005) than males during mating vocal

playback. **D**. Estrus females, but not non-estrus females or males, showed a significant increase in flinching behavior during Stim 1 and Stim 2 periods (time*estrous: $F_{(2,30)} = 9.0$, $p<0.001$, $\eta^2 = 0.4$). **E-G.** Changes in concentration of acetylcholine (ACh), dopamine (DA), and 5-HIAA relative to the Pre-Stim period, evoked during Stim 1 and Stim 2 periods of vocal playback (n_{Male} = 9, $n_{Estrus\ Fem}$ = 8, $n_{Non-estrus\ Fem}$ = 7). **E.** Release of ACh during mating playback increased in estrus females (Stim1, Stim 2) but decreased in males and non-estrus females (Stim 2). Among groups, there was a significant estrous effect (estrous: $F_{(1,21)} = 29.0$, $p<0.001$, $\eta^2 = 0.0.6$. **F.** DA release during mating playback increased in all groups relative to Pre-Stim period , with no significant sex ($F_{(1,21)} = 0.8$, $p=0.4$, $\eta^2 = 0.04$) or estrous effect ($F_{(1,21)} = 0.9$, $p=0.4$, $\eta^2 = 0.04$). **G.** No significant changes in 5-HIAA during mating sequence playback (sex: $F_{(1,21)} = 0.07$, $p=0.8$, $\eta^2 = 0.004$; estrus: $F_{(1,21)} = 1.6$, $p=0.22$, $\eta^2 = 0.07$). **A-G**: Repeated measures GLM: *$p<0.05$, **$p<0.01$, ***$p<0.001$ (Bonferroni post hoc test). Time windows comparison: 95% confidence intervals.

Figure 5

4
5

Figure 5. A single bout each of mating and restraint experience altered behaviors and ACh release in response to vocal playback. In all graphs, dots represent measures from individual animals obtained during the Stim 1 playback period; thick horizontal lines represent mean values across subjects. There were no differences in behavioral counts for Pre-Stim values between INEXP and EXP mice for any group (e.g., males-restraint, males-mating, estrus females-mating. Neuromodulator values are normalized to the baseline level. **A.** Experience increased flinching responses in estrus female mice but not in males in mating or restraint vocal playback (time*sex*experience: $F(1.6, 56) = 4.1$, $p=0.03$, $\eta^2 = 0.11$)). **B.** Experience increased rearing responses in males exposed to mating playback (sex*experience: *F(1,35)=5.3, P=0.03,* η^2 *=0.13*). **C.** Estrus female mice -(sex*experience: $F(1, 39) = 8.0$, $p=0.008$, , $\eta^2 = 0.2$,) and restraint male mice (context*experience $= F(1, 39) = 13.0$, $p<0.001$, $\eta^2 = 0.2$)displayed consistent experience effect for changes in ACh. **D**. DA did not show the EXP effect observed in ACh during vocal playback (sex*experience: $F(1, 39) =0.12$, $p=0.7$, $\eta^2 = 0.003$; context*experience $= F(1, 39) = 4.0$, $p=0.052$, $\eta^2 =0.09$. GLM repeated measures with Bonferroni post hoc test: *$p<0.05$, **$p<0.01$, ***$p<0.001$.

21
22

1 **Figure 6**

2
3
Figure 6. In INEXP mice, vocal playback failed to evoke distinct patterns in
neuromodulator release. Boxplots show change in concentration of the specified
neuromodulator in 10-min playback periods (Stim 1, Stim 2) compared to the baseline level (Pre-
Stim). Male-restraint, $n_{INEXP} = 7$; Male-mating, $n_{INEXP} = 7$; Estrus Female-mating, $n_{INEXP} = 7$). **A.**
ACh release shown for both playback periods. **B.** DA release shown for both playback periods.
C. 5-HIAA concentration shown for both playback periods.
10

Figure 7

Figure 7. Proposed model for neuromodulation of salient vocalization processing via
acetylcholine (ACh) and dopamine (DA) in the basolateral amygdala (BLA). A. Cholinergic
modulation of CeA-projecting neurons during aversive vocalization cue processing in the BLA.
In the presence of aversive cues, ACh released from the basal forebrain acts on M1 ACh
receptors (M1 mAChRs) to enhance the cue-induced excitatory responses of CeA -projecting
neurons. In contrast, NAc-projecting neurons are quiescent, because they do not respond to
aversive cues and are inhibited by interneurons that are activated through nicotinic ACh
receptors (nAChRs). B. Dopaminergic modulation and enhancement of signal-to-noise ratio in
response to reward-associated cues (appetitive vocalizations). When vocalizations or other
rewarding cues are present, release of DA from VTA enhances D2R-mediated excitation in NAc-
projecting neurons that are responsive to positive cues. In contrast, CeA -projecting neurons are
not responsive to rewarding vocalizations and are inhibited by local interneurons. DA is thought
to act on D1 DA receptors (D1Rs) in these local interneurons to shape a direct inhibition onto
CeA-projecting neurons. Abbreviations: ACx: auditory cortex; BF, basal forebrain; CeA, central
nucleus of amygdala; MGB, medial geniculate body; NAc, nucleus accumbens; VTA, ventral
tegmental area.

Table 1: Classification of manually evaluated behaviors during playback of vocalizations.

Behavior	Definition
Abrupt attending	Sudden and quick pause in locomotion followed by abrupt change in head and body position lasting 2+ seconds. This behavior is accompanied by fixation of the eyes and ears during attending. Appears similar to "freezing" behavior described in fear conditioning but occurs in the context of natural response to vocalizations.
Flinch	A short-duration twitch-like movement in response to vocal sequences, occurring at any location within arena. Unlike acoustic startle (Grimsley et al., 2015), this behavior occurs in free-moving animals in response to non-repetitive, variable-level vocal stimuli. Flinching movements are of smaller magnitude than those observed in acoustic startle.
Locomotion	Movement of all four limbs from one quadrant of the arena to another.
Rearing	A search behavior during which the body is upright and the head is elevated to investigate more distant locations.
Self-grooming	Licking fur and using forepaws to scratch and clean fur on head. The behavior can extend to other parts of the body.
Still and alert	Gradual reduction or lack of movement for 2+ seconds, during which animal appears to be listening or responding to external stimulus. Distinguished from abrupt attending by gradual onset.
Stretch-attend posture	A risk-assessment behavior during which hind limbs are fixed while the head, forelimbs, and the body are stretched sequentially in different directions.