Vertebrates but not ants protect rainforest trees from herbivorous insects along an elevational

gradient in Papua New Guinea

Running title: Exclosures Mt. Wilhelm

Katerina Sam[*,a,b], Leonardo Re Jorge[b], Bonny Koane[c], Pita Amick[c], Elise Sivault[a,b]

[a] Biology Centre of Czech Academy of Sciences, Institute of Entomology, Ceske Budejovice, Czech

Republic

[b] University of South Bohemia, Faculty of Science, Ceske Budejovice, Czech Republic

[c] The New Guinea Binatang Research Center, Madang, Papua New Guinea

* corresponding author: Katerina Sam, katerina.sam.cz@gmail.com, Biology Centre CAS, Institute of

Entomology, Ceske Budejovice, Branisovska 31, Czech Republic

Draft: 2022-07-22

Acknowledgements: We are thankful to villagers from Kausi, Numba, Degenumbu, Sinopass, Bruno

Sawmill and Kegesugl for allowing work on their land and assistance in the field. The work wouldn't

be possible without the logistical support and help of The New Guinea Binatang Research Centre.

We are thankful for various support of Prof. Vojtech Novotny, Prof. Nigel Stork and Prof. Roger

Kitching for support during the field and laboratory work, which was carried during the maternal

leave of KS. The work was supported by Czech Science Foundation Grant 18-23794Y and European

Research Council Starting Grant BABE 805189 (PI: K. Sam).

Aim: The theory on trophic interactions between plants, insect herbivores, and their predators

predicts that predators increase plant biomass by feeding on herbivores. However, it remains

unclear whether different types of predators regulate herbivores to the same degree, and how the

trophic interactions affect lower trophic levels along elevational gradients where predator

communities differ significantly. Therefore, we aimed to investigate the impact of vertebrate

predators and ants (individually and in combination) on arthropod communities and leaf herbivory

along a complete tropical forest gradient.

Location: Papua New Guinea

Taxon: Multi-taxon

Methods: We excluded predators from 560 saplings in two six-month long predator exclusion

experiments spanning wet and dry seasons. Saplings were spread across 8 study sites which were

evenly spaced at 500 m elevational increments from 200 to 3700m a.s.l..

Results: On average the density of arthropods increased significantly by 37% and 33% respectively

when vertebrate predators, and both ants and vertebrates predators, were removed. Both season

and elevation mediated this effect significantly. At lower trophic levels, both the exclusion of both

vertebrates alone, and exclusion of vertebrates plus ants, led to a significant increase in leaf damage

by 50% and 36% respectively. In contrast, the exclusion of ants alone had no significant effect on

arthropod density or leaf damage, which increased by 12% and 9% respectively.

Main conclusions: Our results indicate that the relative contribution of birds and bats changes at

different elevational sites, while the overall effect of vertebrate predators remains consistent

along the whole elevational gradient. This contrasts with ant driven trophic cascades which

brought about increased herbivory only at the most productive sites of the elevational gradient,

where ant abundance is highest. We conclude that disappearance of insectivorous vertebrate

predators can lead to substantial negative consequences for plants.

Keywords: Altitudinal gradient, herbivory, predation, predator exclosures, top-down control, trophic

cascades

Introduction

Insect herbivores are the primary consumers of plant leaf tissue in tropical forests (Coley and Aide 1991, Coley and Barone 1996, but Suzuki, et al. 2013), and their impact is notably increased in the absence of top-down control (Mooney, et al. 2010, Sam, et al. 2022). This top-down control is therefore a high value ecological service across a range of tropical ecosystems (Schmitz 2006, Sekercioglu 2006). It is usually assumed that disparate insectivorous predator groups have similar effects and are thus treated as a single functional unit, leaving their individual or joint effects largely unknown (Mooney 2007, Perfecto and Vandermeer 1996, Richards and Coley 2007, Sih, et al. 1998). Resolving the often-complex trophic dynamics of multiple types of predators in the same community and their cascading effect on plant performance remains an important challenge in ecology (Singer, et al. 2017). This challenge is primarily rooted in the variable strength of trophic cascades between predators, prey and plants both within (Moon and Stiling 2004, Mooney and Linhart 2006) and between communities (Shurin, et al. 2002).

Bats and birds, and to a lesser extent insectivorous vertebrate, consume large quantities of insects and other arthropods (Kalka and Kalko 2006, Mooney, et al. 2010, Nyffeler, et al. 2018). They are likely to act as intraguild predators, feeding both on predatory as well as herbivorous arthropods (Ingala, et al. 2021, Karp and Daily 2014, Milne, et al. 2016, Sam, et al. 2017). The impact of ubiquitous insectivorous birds on lower trophic levels was shown in some but not all studies (Mooney, et al. 2010, Van Bael, et al. 2008). The impact of bats on arthropods and herbivory might be comparable to the effect of birds according to some studies (Kalka and Kalko 2006, Williams-Guillén, et al. 2008). However, bats can be erroneously overlooked in exclusion studies (Mooney, et al. 2010, Sam, et al. 2022), as the typical approach of covering foliage with a net effectively quantifies the combined effect of excluding both birds and bats (Greenberg, et al. 2000, Holmes, et al. 1979, Van Bael, et al. 2003, but see Kalka, et al. 2008).

Ants are recognised as other important predators in many habitats (Hölldobler and Wilson 1990, Rosumek, et al. 2009). Despite their study as natural enemies and biological control agents (Mestre, et al. 2012, Philpott and Armbrecht 2006, Rosumek, et al. 2009, Way and Khoo 1992), their importance as predators remains equivocal. Previous efforts failed to detect their effect on lower trophic levels or showed temporal inconsistency (Tanhuanpää, et al. 2001, Sam, et al. 2022). Plant damage in the absence of ants was reported to increase between 0 and 250% on average, with the duration of the ant exclosure (Sam, et al. 2022), and productivity of the study site (Rosumek, et al. 2009, Sam, et al. 2022) having signficant impacts. Ant exclosures are difficult to conduct (Hood, et al. 2022, Rosumek, et al. 2009) thus bait removal is often considered as a proxy of predation (e.g., Liu,

et al. 2020, Roslin, et al. 2017, Burger, et al. 2021). Unfortunately, this method may not accurately

measure the realized top-down control and trophic cascade to plants from ants.

Changes in the diversity and abundance of animal taxa along elevational gradients is one of

the most striking biogeographic patterns on Earth (Colwell, et al. 2016, McCain 2005, Rahbek 1995,

Sam, et al. 2019). These distributional changes lead to changes in the relative importance of

predator groups at different sites on the same gradient. As such, elevational gradients provide an

excellent experimental environment for studies focussing on the individual and combined effects of

insectivorous predators on prey and plants (Schemske, et al. 2009). As climate change is expected to

lead to further changes in predator distribution and thus ecosystem function (Chen, et al. 2011), it is

increasingly important to understand individual and combined effects of various groups of predators

along large environmental gradients.

Changes in diversity along elevational gradients are also accompanied by changes in body

size where lowland animals are generally larger than animals at higher elevational sites. As predator

size also determines preferred prey sizes, we can also expect the effect of predators to be detectable

in the mean body size of insect communities. For example, birds were shown to reduce the mean

size of caterpillars by 12% through preferentially feeding on large individuals (Singer, et al. 2017).

Similarly, birds and bats reduced mean body size of Araneae, Formicidae and arthropod larvae (Karp

and Daily 2014). In contrast to vertebrate predators, ants are expected to hunt smaller prey, thus

their absence from an ecosystem might lead to an increase in mean arthropod body size.

Here we investigated the impact of vertebrate predators and ants (individually and in

combinations) on arthropod communities and leaf herbivory along a complete tropical forest

gradient in Papua New Guinea. We aimed to answer the following questions: (1) Do predators

significantly affect the density of arthropods in various tropical habitats along the elevational

gradient? We hypothesise that the removal of vertebrate predators, but not ants, will result in an

increased density of arthropods on plants. Moreover, we expect that the effect of predators will

decrease with increasing elevation (1A) and will be higher in the wet season than in the dry season

(1B) due to the differences in productivity and the assumption that tropic interactions are stronger

in more productive environments. (2) Are the effect of ants and vertebrate predators detectable on

the mean body size of arthropod communities? We propose that arthropods will be relatively larger

in the absence of vertebrate predators and relatively smaller in the absence of ants. (3) Do any of

the predator groups display disproportionate impacts on predatory arthropod abundance compared

with herbivore abundance? We expect that both focal predator groups (i.e., vertebrate predators

and ants) act as intraguild predators whereby they feed on both herbivorous and predatory

arthropods similarly, thus diminishing the effect of trophic cascades but not so much as it wouldn't

be detectable. (4) Are the effects of ants and vertebrate predators independent of one another or

do they interact? We hypothesise that the effect will be mostly independent and additive, as ants

and vertebrate predators feed on different prey items and birds (and presumably also bats) consume

few ants (Sam, et al. 2017).(5) Does the potential effect of predators cascade down to the trophic

level of plants and is it detectable on plant herbivory? Considering our first hypothesis, we suggest

that the removal of vertebrate predators, but not ants, will have a measurable effect on plants thus

resulting in increased herbivory.

Materials and methods

Study sites and experimental trees

Our study was performed along the slopes of Mt. Wilhelm (4,509 m a.s.l.) in the Central Range of

Papua New Guinea (Figure S1). The rainforest gradient spans from the lowland floodplains of the

Ramu river (200 m a.s.l., 5° 44' S 145° 20' E) to the timberline (3,700 m a.s.l., 5° 47' S 145° 03' E). The

gradient is 30 km long and consists of eight sites evenly spaced at 500 m elevational increments.

Average annual precipitation is 3,288 mm (local meteorological station) in the lowlands, rising to

4,400 mm at 3,700 m a.s.l., with a distinct condensation zone around 2500 – 2700 m a.s.l. Mean

annual temperature (measured by our data loggers – Sam, et al. 2019) decreases from 27.4°C at the

lowland site to 8.4°C at the tree line at a constant rate of 0.54°C per 100 elevational metres. The

habitats and zonation of forest types are described elsewhere (McAlpine, et al. 1983, Paijmans 1976,

Sam, et al. 2019, Tvardikova 2013).

We selected and tagged saplings belonging to 3 – 7 tree species at each elevational study

site prior to the experiment (Table S1). Unfortunately, no single tree species or genus is distributed

along the complete elevational gradient. Therefore, we aimed to work with several *Ficus* species

which represent the most dominant and ecologically important genus from 200 up to 2700 m a.s.l.

We had to work with other locally dominant species at the two highest elevations (Table S1). In

total, we selected 80 (or 40 at 3,200 and 3,700m asl) individual saplings per elevational site, i.e., 560

saplings along the entirety of the elevational gradient (Table S2). For statistical independence, we

allowed at least 50 m between any pair of tree individuals. We visually assessed saplings of the focal

species and selected individuals which had approximately 500 leaves growing within a well-

developed crown 2.5 – 4 m above the ground, did not have any ant nests, and did not have

abnormally high herbivory or fungal damage. The selected tree species did not produce any

exudates or sugar droplets for attracting ants. Average leaf-sizes of the selected species ranged from

16.31 to 154.10 cm^2 (mean ± S.E. = 52.16 ± 6.12). Mean sapling DBH at the beginning of the

experiment was 1.16 cm for exclosures and 1.17 cm for controls.

Experimental design

We prevented vertebrate predators (i.e., birds and bats together, VER) from accessing 20 out of the

80 preselected saplings at each elevational site between 200 and 2,700 m a.s.l. We also prevented

ants accessing another 20 saplings (ANT) and both vertebrate predators and ants accessing another

20 saplings (ALL) at each elevation. We left 20 saplings without predator protection as controls

(CON). At the elevational sites 3,200 and 3,700 m, we protected 20 saplings against vertebrates

(VER) and kept 20 as control saplings (CON). There were no ant exclosures at these two sites as they

are above the natural elevation range of ants (Moses 2015, Colwell et al. 2016). Thus, at each

elevation, 3-7 tree species were selected for conducting our experiments, with a total of 18 tree

species used across the 6 elevation bands between 200 m and 3700 m . Most tree species occurred

in multiple elevation bands (see Table S1 for details).

Vertebrate (VER) exclosures were constructed from PVC tubes (1 cm in diameter) joined

together by PVC corners and covered with agricultural nylon netting (mesh opening 3 x 3 cm,

transparent green). Each exclosure had dimensions 2 x 2 x 2.5 m and comprised a volume of 10 m^3,

which enclosed an average of 1.63 (± S.D. 1.02) m^2 of leaf area. The mesh size was comparable to

other exclosure studies [e.g., 29×29 mm in Greenberg, et al. (2000); 25×25 mm in Mols and Visser

(2002); 20×20 mm in Van Bael, et al. (2003)]. The exclosure materials did not attract arthropods, did

not damage leaves or branches, and did not significantly reduce light. We attached a system of ropes

to neighbouring vegetation (i.e., strong branches) in order to secure the exclosures in place (thus

preventing the exclosure from moving and disturbing the foliage of the target saplings).

Observations confirmed that netting did not exclude small lizards as they were seen crawling under

the cage. However, one ca. 50 cm long lizard was found tangled in the netting and was subsequently

released, suggesting the exclosures partially prevented access of non-flying insectivores which were

not quantified in the study.

To exclude ants (ANT), we used the adhesive Tanglefoot® pest barrier (Philpott, et al. 2008,

Philpott, et al. 2004) which we applied in a 30cm wide strip around the trunk of the sapling at breast

height. We removed all lianas and branches touching the sapling and we carefully checked for

potential ant nests in plant hollows. The ants foraging on foliage were removed by both beating and

manual removal. We placed tuna baits (one teaspoon under a gauze tied to the trunk of the tree) 20

cm above the Tanglefoot layer and checked for the presence of ants two hours later. All individuals

were removed. Tanglefoot was reapplied every 3 – 4 weeks and new tuna bait was placed at the

same place to confirm that ants were not present. Some ants crossed the Tanglefoot barriers so our

analysis excluded all ant exclosure saplings where ant activity exceeded 3 ants on a single checking

date (3 saplings in total).

To exclude all (ALL) predators (ants and vertebrate predators) we used a combination of the

above methods, i.e., by applying Tanglefoot to the trunk, installing a cage around the sapling, and

preventing the foliage from touching the cage. Tanglefoot was also applied to all of the support

ropes used to lower the cage and hold it in place so as to prevent ants from accessing the sapling via

neighbouring vegetation.

Measured variables

We sampled the effect of predator exclusion after 6 and 12 months, corresponding to the dry and

wet season respectively (Table S2). We first carefully opened the protective nettings where needed.

Next the trunk of the sapling was slowly lowered above a 2x2m mosquito net, wrapped into the net,

and sprayed by fast knock-down insecticide (Mortein®). Shortly after we shook the foliage firmly,

opened the net, collected all arthropods (>1mm) and preserved them in vials filled with DNA grade

ethanol. Arthropods were later identified into orders or families, feeding specialization, life stage

and measured to the nearest 0.1 mm (Table S3).

To assess herbivory leaf damage, we randomly selected two branches (ca. 50 leaves in total)

and collected all their leaves. We estimated how many other leaves in total were present on the

sapling and used this number to calculate total leaf area. We took photographs of the flattened

leaves of each sapling against a 50 x 50 cm^2 white background. Using Adobe Photoshop CS6 (Adobe

Systems Inc., USA), we outlined the missing edges of the leaves based on their expected shape. We

then used ImageJ version 1.47 (National Institute of Health, USA) to calculate the remaining leaf area

(a, in cm^2), the extrapolated leaf area without any herbivore damage (b), and the area lost to

herbivory (c = b - a). We then estimated the percentage of leaf–area loss per m^2 of foliage and total

standing leaf area per each whole focal sapling using the count of the number of leaves (i.e., number

of leaves * mean leaf area a) which we later used to calculate densities of arthropods per m^2.

We completed the second survey on the same saplings approximately six months after the

first survey (Table S2). After the first survey, all exclosures were renewed and only some (n= 16)

saplings (including their treatments) were replaced completely due to their death or damage (e.g.,

vandalism of villagers, fallen branches, landslides). The second survey was conducted exactly the

same way as the first, with the only difference being that *all* foliage was now collected from each

sapling. We placed all leaves collected from individual saplings into a bag, randomly selected ca. 50

leaves for photographing and further analyses, and weighed all leftover foliage as well as the leaves

used for photography. This allowed us to calculate total leaf area (from known weight and leaf area

of leaves used to measure herbivorous damage) of the saplings more accurately than during the first

survey. We found that leaf herbivory within a branch (i.e., herbivory on leaves from first survey) was

correlated similarly (P < 0.001, R = 0.854) to herbivory on leaves randomly collected within the

whole plant (i.e., herbivory on leaves from first survey; P < 0.001, R = 0.628). Further, estimated leaf

area (from the first survey) was correlated (P < 0.001, R = 0.358) to the real leaf area based on leaf-

weight (from the second survey), despite some saplings growing significantly more than others

during the project.

Predator survey

Bird communities here have been repeatedly surveyed by point counts as part of long-term

monitoring efforts, with this methodology and data published in several studies. In short, point

counts at each elevational site were carried out at 16 points regularly spaced along a 2,350 m

transect (successive points were 150 ± 5 m apart to avoid overlap). All birds seen or heard within a

fixed radial distance of 0 - 50 m were recorded. Each count lasted 15 min. Bat communities were

surveyed in the understory during two expeditions conducted in wet (February – March 2015) and

dry seasons (June – July 2015). This method and data were published in Sivault, et al. (2022). In

short, we used an ultrasonic bat call detector coupled with a recorder to detect echolocating bat

species. We recorded bats at five points (i.e., 15 minutes per point) separated by 200 meters at each

elevation, in line with the bird transect methods described above. Surveys were conducted for four

days per site after sunset (6 pm). The ant communities at each of the study sites were sampled by

hand collection. The trunk of each sapling was inspected for ants at breast-height for 10 minutes,

and also by the tuna bait method in May-June 2014. More details on the respective survey are

provided in Supplementary material.

Data analysis

Arthropod densities: We used a linear mixed model and model selection to determine the effect of

our experimental treatments on arthropod densities. We used arthropod density (per m^2 of foliage)

as our response variable and treatment, season and elevation and possible two-way interactions

between them as our predictor variables. Elevation was modelled as a second-order polynomial to

allow for non-linear elevational trends. We used tree species as a random effect in our model to

account for differences in baseline arthropod densities between species, an expectation arising from

variation in defensive traits. Individual saplings were used as a second random factor as each of the

560 focal saplings was surveyed twice i.e., in both wet and dry seasons.

Herbivory: The effect of our experimental treatments on herbivory damage (as the

proportion of the leaf area lost per m^2 of foliage) was analysed using a generalised linear mixed

model with a beta error structure and model selection. As models with a beta error structure only

allow values between 0 and 1, the 0 values in our study (n = 2) were converted into 0.0001 prior to

analysis. We used herbivory as the response variable, while experimental treatment, elevation band

and their interactions were predictor variables with tree species as a random effect. Again, elevation

was modelled as a second-order polynomial. We used the glmmTMB package (Brooks, et al. 2017)

within R 4.0.2 (Team 2020) to build our generalised linear mixed models.

Body sizes: We modelled the mean body size of arthropods from each of the saplings using

a similar model to that described above for arthropod density.

Effect on groups of arthropods: We based this analysis on the unweighted natural log

response ratios LRR (Curtis and Wang 1998, Hedges, et al. 1999) calculated from mean responses of

individual groups of arthropods in the presence and absence of predators. LRR is an effect size

measure that quantifies the results of experiments as the log-proportional change between the

means of the treatment (in the absence of insectivorous predators, μ_{I-}) and control group (in the

presence of insectivorous predators, μ_{I+}) and was thus calculated as ($\ln[\mu_{I-}/\mu_{I+}]$) for each of the three

treatments at each elevational study site and season. We then constructed linear models, selected

the best one, and used the function *get_model_data* from package *metan* to obtain effect estimates

(https://github.com/KatkaSam/GRAD_Exclosures).

Abundance, species richness, and biomass of insectivorous birds and bats were correlated

(Pearson correlation) with the log response ratio (LRR, calculated from the raw data) of the

vertebrate exclosure treatment at each elevation. The number of saplings with ants on them and

abundance of ants on baits were correlated with log response ratio LRR of the ant exclosure

treatment at each elevation. All data were normalized on the scale of 0-1 and averages across both

seasons were taken as the data from the predator surveys did not exactly match the timing of the

experimental predator exclusion.

Results

Change in arthropod densities: In general, the density of arthropods was higher and increased more

in the wet season than in the dry season (Table S6). Based on the model predictions, the density of

arthropods increased on average by 37% when the vertebrate predators were removed (Table S6).

While the effect of vertebrates on arthropods was significant even when combined with the effect of

ants (33% increase, Table S6), the exclusion of ants alone had no significant effect on arthropod

density, which only increased by 12% based on the model predictions (Table S6). Ant removal was

successful. There was a total of 538 and 884 ants found on control saplings (CON) and on saplings

protected against vertebrates (VER) respectively, but only 13 and 30 on saplings protected against

both types of predators (ALL) and ants only (ANT) respectively, across the whole experiment. Thus,

ants represented 2.3% and 1.9% of all arthropods in CON and VER saplings respectively but less than

0.05% of arthropods on ANT and ALL saplings. Elevation and the interaction between elevation and

treatment, and elevation and season, also had significant effects on the resulting density of

arthropods (Figure 1, Table 1).

Change in herbivory: Predator exclusion, elevation and season had a significant additive

effect on leaf area loss (Figure 2, Table 1). The natural mean standing herbivorous damage on

control saplings at 200 m a.s.l. was 2.24 ± 1.7% in the wet season and 2.49 ± 1.7% in the dry season

and decreased with increasing elevation [see Sam, et al. (2020) for more results]. Overall, exclusion

of insectivorous vertebrates as well as the exclusion of both types of predators (ALL) led to a

significant increase in leaf damage by 50 and 36% respectively (Table S7). However, the increase in

leaf damage by ca. 9% in the ant exclosures was not significant (Table S7). Insects consumed

significantly more leaf area in the wet than in the dry season (Figure 2). The increase in leaf damage

seemed to be highest in the lowest and upper most elevations (Figure 2, Table S7).

Change in mean body size of arthropods: Treatment, elevation, season, and interaction

between elevation and treatment, and elevation and season, had an impact on the mean body sizes

of arthropods (Figure 3, Table 1). While model selection identified season as a significant factor due

to its importance in model interactions, the effect of season alone was not significant (Table 1). This

means that seasonal effect cannot be described properly because of its variation across elevations.

The arthropods in caged treatments were significantly bigger than arthropods collected from control

saplings and saplings from which ants were excluded (Figure 3, Table 1). The majority of arthropods

collected during the experiment were very small, with larger arthropods (i.e,. greater than ca. 2.2 cm

in length) found exclusively in the caged treatments (VER and ALL) to which vertebrate predators

had no access.

Changes in community composition: Overall, 12,177 arthropod individuals were identified

as predatory arthropods, while only 4,085 were herbivores potentially responsible for leaf damage

and 10,065 were arthropods which had no relationship to chewing herbivory (Table S3). The

majority of predatory arthropods (8,221) were spiders. Absolute abundance as well as relative

abundances of spiders were higher in treatments where ants were excluded (both ALL and ANT) but

the change was not detectable in vertebrate exclosures (VER) (Figure 4). Across the whole study,

Araneae, Diptera, Hemiptera, and Lepidoptera larvae increased their abundances significantly after

the removal of ants. Araneae, Coleoptera, Lepidoptera larvae, Orthoptera and all other arthropods

increased their densities (per m^2 of foliage) on saplings from which the insectivorous predators were

excluded. Densities of Hymenoptera (other than ants) and Hemiptera tended to decrease after

vertebrate predators were excluded (Figure 4).

338    ***Predator communities and their relation to the observed cascading effect***: Overall, we

found only weak relationships between the communities of predators and the cascades they cause.

In general, the abundances of predators correlated with the effect measured on abundances of prey

(Figure 5), but not with the effect measured on herbivory damage (Figure S2). This may simply be

the result of having just eight points in the correlation thus limiting the robustness of the analyses.

Abundances of insectivorous birds, and thus those potentially responsible for attacks, peaked at mid

elevations (1700 - 2700 m) and correlated positively and significantly (given the low power of the

analysis and strong effect size) with the effect of vertebrate exclosure treatment on arthropod

abundances (R = 0.71, P = 0.049, Figure 5a). Abundances of insectivorous bats decreased steeply

from 200 to 1200 m a.s.l. and then peaked again at 2700m a.s.l. This pattern was again positively,

but not significantly, correlated with the effect of vertebrate exclosure (R = 0.56, P = 0.15, Figure 5b).

We detected more bats, and especially gleaning bats, in lowlands than at higher elevations [see also

Sivault, et al. (2022) for more detailed results]. It seems that our results indicate that bats are at

least partly responsible for predation at low elevations, while insectivorous birds are more important

predators at the mid-elevations. Neither richness nor biomass of insectivorous birds or insectivorous

bats correlated well with the effect of the exclosures measured on abundance of prey, and

abundance of these insectivorous predators thus seemed to be a better correlate for predation

pressure (R < 0.21, P > 0.78 in all cases). The number of trees on which ants occurred as well as ant

abundances decreased with elevation. Despite the positive relationship, neither the number of trees

with ants on them (R = 0.25, P = 0.56) nor abundance of the ants (R = 0.14, P = 0.74, Figure 5c)

correlated significantly with the effect of the ant exclosures on abundances of arthropods.

Discussion

Predator removal increases arthropod richness on plants, which is in line with our first hypothesis

(*H1*). Overall, arthropod richness increased by ca. 37% after the removal of insectivorous

vertebrates, which is in line with the estimates from two global reviews reporting an increase by ca.

39% (Mooney, et al. 2010) and 69 % (Sam, et al., 2022). The effect of ant removal was overall rather

low, with arthropod abundance increasing by 12%, while the global effect of ant removal is

estimated to be ca. 12% (Sam, et al., 2022) but also up to more than 100% (Rosumek, et al. 2009).

However, it is important to note that Rosumek's, et al. (2009) study analyses datasets comprising

70% myrmecophytic and only 30% non-myrmecophytic plants, while Sam's, et al. (2022) analyses

mostly data from non-myrmecophytic plants. Similarly, the plant species used in the current study

have no or very weak associations with ants. It is important to note that the treatment itself almost

completely removed ants from our experimental saplings, and ants represented ca. 2% of all

arthropods in samples from our controls, thus impacting the effect of the treatment on the total

density of the arthropods.

The overall effect of predators on lower trophic levels was however affected by season

(*H1A*). Predator exclosures had a smaller effect on arthropod richness in the dry season than in the

wet season. The seasonal differences are likely to be much higher in more seasonal habitats than

those of the northern side of Papua New Guinea, which is known to be only weakly seasonal

(Novotny and Basset 1998, Wright, et al. 1997). Birds also appear to follow the patterns observed

here (Bell 1982). Observations made during our long-term surveys indicate that most breeding of

insectivores occurs at the onset of rains, when the arthropods become more abundant in response

to leaf flushing (Sam, et al. 2015, Sam, et al. 2017), which can lead to strong effects of predator

exclosures.

Contrary to our expectations (*H1B*) the effect of insectivorous vertebrates on arthropod

richness increased both towards the lowest (200 and 700m) and the highest elevations (above

2700m). Based on the productivity of the environment and previous observations on dummy prey

predation rate by birds along the same gradient (Sam, et al. 2015), we expected that predator

exclusion would have the highest effect between 700 and 1700 m a.s.l. where the abundances of

insectivorous birds peaked. Our current results imply that the predation rate observed on dummy

prey may not translate directly into the actual effect of predator exclusion, as there is an interplay

between the natural abundances of arthropods and their predators. It is important to again note,

that both groups of predators (birds and bats) were excluded in the current study. While the

abundance of insectivorous bats was high at low and high elevations but low at mid-elevations, the

abundance of insectivorous birds peaked at the mid-elevations. Moreover, bat communities

between lowlands and highlands are different, with most of the gleaners found in lowlands. We can

only speculate that bats are responsible, at least partly, for the predation at low elevations as our

data are not suited to detect subtle elevational trends. We showed that predation risk on

arthropods was correlated with the abundance of predatory birds (and less so for bats), supporting

thus the hypothesis that predator density drives the effect of predation on lower trophic levels.

These results are in line with some but not all earlier studies (Nordberg and Schwarzkopf 2019). Our

data further indicates that the abundances of birds are a better correlate for the effect of predators

than their richness or biomass.

Predator exclusion was affected not only arthropod densities but also their mean body sizes
(*H2*). In general, the absence of insectivorous predators allowed larger arthropods to survive. This
effect was detectable despite the fact that the mesh size (3x3 cm) of the cages was likely also
preventing access of large arthropods. Arthropod mean body size increased primarily at low
elevations where many large arthropods occur (Sam, et al. 2017, Horne, et al. 2018), but also at the
highest elevations. In contrast to vertebrate insectivores, the exclusion of ants did not have any
effect on the mean body size of arthropod communities. As such this contradicts results of an earlier
study where ants preyed selectively upon small-bodied caterpillars, increasing thus mean caterpillar
length by 6% (Singer, et al. 2017). However, we believe that this is because ants hunt not only
individually for tiny prey, but also hunt collectively for large arthropods (Schmidt and Dejean 2018).
This might lead to a balanced consumption of arthropods of all sizes.
Overall, the absence of predators allowed various groups of arthropods to increase their
abundances (*H3*). First of all, insectivorous vertebrates did not affect ant density significantly,
implying limited intraguild predation between these taxa in our study systems. The absolute
numbers of ants increased on saplings protected against vertebrates, and ants represented very
similar proportions of arthropods in samples from the control and vertebrate-protected saplings.
Diet analyses of birds from the same gradient also showed that ants are not important prey for birds
(Sam, et al. 2017). Other studies concur with this, where Coleoptera, Aranea, Lepidoptera larvae
(Poulin, et al. 1994, Sam, et al. 2017) and many other smaller groups of arthropods (Bodawatta, et
al. 2022, Poulin, et al. 1994) were found to be preferred prey of birds, but not ants (*H4*). This seems
to be reflected in our current results, as these groups of arthropods increased their abundances
significantly after the insectivore predators were removed, while other groups (e.g., Hymenoptera,
Diptera, Hemiptera) did not respond strongly to the absence of predators. In the ant exclosures,
Hemiptera, Lepidoptera, and Aranea increased their abundances significantly, but spiders seemed to
be the most affected. This is in line with results of earlier studies which showed that spiders
increased their abundances by more than 100% (Rosumek, et al. 2009) and by 84% (Sam, et al. 2022)
in the absence of ants. Even though our results can't resolve whether the spiders are typical prey of
ants or their competitors, we believe that competition might be at least partially responsible for the
patterns observed as spider abundance was higher in the absence of ants than in the absence of
vertebrates. Results of existing studies are vague, and despite most of the studies typically show ants
to feed on eggs and insect larvae primarily (Gathalkar and Sen 2018), there is some evidence of them
hunting spiders also (Risch and Carroll 1982).
In contrast to previous reports, the standing herbivory on control saplings along Mt Wilhelm
was 5 to 10 times lower than the global estimates and we discuss this matter and potential reasons

in an earlier study (Sam, et al. 2020). The change in herbivory following the removal of predators

was also surprising (*H5*). Herbivory damage increased significantly both after removal of ants and

vertebrate predators but varied in time and space. Herbivory across the whole gradient in the wet

season increased by 112, 78 and 78% after the removal of all predators, vertebrate predators, and

ants respectively. In contrast, the dry season saw increases of 58, 52 and 42% respectively. All of

these increases are higher than the previous global estimates which predicted increases in leaf

herbivory of ca. 40 - 47% in the absence of vertebrate predators (Mooney, et al. 2010, Sam, et al.,

2022). The effect of ants on herbivory at low elevations was actually stronger than their effect on

the density of herbivore arthropods, where ants increased herbivory by ca. 130% and herbivorous

arthropods only by 3%. Similarly, Rosumek et al. (2009) showed that ant removal increased

herbivory by more than 95%, whereas the abundances of herbivorous arthropods increased by ca.

60% in the tropics.

Season had an effect on both the densities of arthropods and herbivory. The removal of

vertebrate predators in the wet season led to increases in herbivory of ca. 100-230%, where peaks

towards the lowest and higher elevations created a distinct U-shaped pattern. We saw the highest

increase in herbivory during the dry season following the removal of vertebrates above 2200m

which ranged between 160% and 555%. This pattern suggests that insectivorous predators provide

more effective services to plants at the highest and lowest than at middle elevations. However, this

is likely heavily dependent on plant defences and bottom-up control of the trophic cascades.

According to one hypothesis, plants are expected to invest progressively less into

chemical defences at higher altitudes as arthropod richness is generally lower there (Garibaldi,

et al. 2011, Pellissier, et al. 2014). Our results did not support this however, as we did not

observe a steep decrease in arthropod densities along our elevational gradient. A second

hypothesis states that replacing biomass at higher elevations comes at a higher cost due to the

harsh, low nutrients environments of higher elevations . These conditions favour higher

investment into defences at the expense of growth (Defossez, et al. 2018, Salgado, et al. 2016).

Plants may also defend their biomass directly and indirectly through insectivorous predators

(Marquis and Whelan 1996). We might speculate that plants do not invest much into direct

defences at higher elevations as arthropods are relatively rare there, and insectivorous birds

more abundant, but instead rely on indirect defences which work in response to herbivory

attacks . While this would explain the pattern we detected, it appears unlikely given our

understanding of plant defences. In an earlier study from the same elevational gradient using

several of the plant species studied here (Sam, et al. 2020), showed that there is no universal

elevational increase in plant defences (i.e., levels of alkaloids and flavonoids), and several *Ficus*

species from mid elevations were also well defended (i.e., high protein precipitation capacity). If

we wish to better understand elevational trends in plant defences then it is necessary to

perform analyses based on multiple traits of direct and indirect defences and to link these to

datasets on arthropods herbivores.

In conclusion, we show that vertebrate insectivores suppress arthropods effectively, yet

their impact differs among various groups of arthropods. The pressure of vertebrate insectivores

is robust across the whole elevational gradient and translates to a significantly higher plant

herbivory in the absence of predators. Our results imply that the relative contribution of birds

and bats differs along the elevational gradient. In contrast, ants do not suppress arthropods

significantly, yet the trophic cascades caused by them do translate into increased herbivory at

the most productive sites of the elevational gradient, where the abundances of ants are also the

highest. Insectivorous vertebrates, but not ants, had a stronger negative effect on large

arthropods which can further negatively impact herbivory rates. While generalities emerge, we

suggest that future experiments need to consider intraguild predation and mesopredators and

plant defences, and that these should be accompanied by more detailed predator surveys as all

these factors may significantly affect the strength of trophic cascades.

Data availability statement: All data and scripts are available at GitHub

https://github.com/KatkaSam/GRAD_Exclosures

Bioscatch: Katerina Sam is a head of Laboratory of Multitrophic interactions and the Department of

Ecology at the Entomology Institute, where she studies relationships between insectivorous

predators, insect and plants. This work represents her earlier postdoctoral experiment, which

further grew into a European Research Council funded project. She and her colleagues study various

aspects of ecology and biodiversity of Mt. Wilhelm from 2008.

Author contributions: KS conceived the ideas, secured funding and conducted the experiment with

help of BK. PA and ES conducted the survey of insectivorous predators, LRJ assisted with statistical

analyses to KS. All authors contributed to writing and edited the first draft written by KS.

Figure 1. Effect of the predator exclusion (treatment) on the densities of all arthropods on saplings

along the elevational gradient of Mt Wilhelm in the wet (a) and dry season (b). Each individual data

point shows the resulting density of arthropods (per m^2 of foliage) on a focal sapling. The curves

show predicted values from the linear model which includes season, elevation and treatment, using

the interaction between elevation and treatment as a fixed factor and plant species and identity of

the sapling as a random effect. Treatments: CON = control saplings with free access of predators,

ANT = saplings from which ants are excluded by tanglefoot glue, VER = saplings from which

insectivorous vertebrates were excluded by cages, ALL = saplings from which both ants and

insectivorous vertebrates were excluded.

Figure 2. Effect of the predator exclusion (treatment) on leaf damage (m^2 per sapling) on saplings

along the elevational gradient of Mt Wilhelm in the wet (a) and dry season (b). Each individual data

point shows the resulting herbivory damage of a focal sapling. The curves show predicted values

from the glmmTMB models which include season, elevation and treatment as fixed factors and plant

species and identity of the sapling as a random effect. Treatments: CON = control saplings with free

access of predators, ANT = saplings from which ants are excluded by tanglefoot glue, VER = saplings

from which insectivorous vertebrates were excluded by cages, ALL = saplings from which both ants

and insectivorous vertebrates were excluded.

Figure 3. Effect of the predator exclusion on mean body size of the arthropods collected from

saplings along the elevational gradient of Mt Wilhelm in the wet (a) and dry season (b). Each

individual data point shows the resulting mean body size of the arthropod community on a given

sapling. The curves show predicted values from the linear model which included season, elevation,

and treatment as a fixed factor. See Table S6 for model results. The occurrence (log of the total

abundance) of arthropods of a given body size in each of the four treatments is shown in (c).

Treatments: CON = control saplings with free access of predators, ANT = saplings from which ants

are excluded by tanglefoot glue, VER = saplings from which insectivorous vertebrates were excluded

by cages, ALL = saplings from which both ants and insectivorous vertebrates were excluded.

Figure 4. Mean proportion of spiders in control treatments (grey area), insectivorous vertebrate

exclosures (VER - magenta line), ant exclosures (ANT - blue line) and both insectivorous vertebrates

and ants exclosures (ALL- yellow line) across the elevational gradient (a). Caterpillar graph showing

how individual orders of arthropods responded to the predator exclosure treatments. The X-axis

shows mean effect sizes of natural log ratios (LRR = ln(exclosure/control)) with 95% CI of individual

response variables. Effect size values above zero indicated that the absence of predators (treatment)

was more harmful to plants than the control (presence of predators), as the abundances of

arthropods in exclosures increased and caused potentially higher herbivory damage to plants (b).

Treatments: ANT = saplings from which ants are excluded by tanglefoot glue, VER = saplings from

which insectivorous vertebrates were excluded by cages, ALL = saplings from which both ants and

insectivorous vertebrates were excluded.

Figure 5. Correlations between the effect of vertebrate exclosures (a, b) and ant (c) exclosures (LRR

calculated from raw data and normalized data) on abundances of arthropods and abundances

(normalized) of insectivorous birds (a), insectivorous bats (b) and ants (c) potentially feeding on the

experimental saplings.

Figure 1.

Figure 1. Effect of the predator exclusion (treatment) on the densities of all arthropods on saplings

along the elevational gradient of Mt Wilhelm in the wet (a) and dry season (b). Each individual data

point shows the resulting density of arthropods (per m^2 of foliage) on a focal sapling. The curves

show predicted values from the linear model which includes season, elevation and treatment, using

the interaction between elevation and treatment as a fixed factor and plant species and identity of

the sapling as a random effect. Treatments: CON = control saplings with free access of predators,

ANT = saplings from which ants are excluded by tanglefoot glue, VER = saplings from which

insectivorous vertebrates were excluded by cages, ALL = saplings from which both ants and

insectivorous vertebrates were excluded.

583
Figure 2.

Figure 2. Effect of the predator exclusion (treatment) on leaf damage (m^2 per sapling) on saplings

along the elevational gradient of Mt Wilhelm in the wet (a) and dry season (b). Each individual data

point shows the resulting herbivory damage of a focal sapling. The curves show predicted values

from the glmmTMB models which include season, elevation and treatment as fixed factors and plant

species and identity of the sapling as a random effect. Treatments: CON = control saplings with free

access of predators, ANT = saplings from which ants are excluded by tanglefoot glue, VER = saplings

from which insectivorous vertebrates were excluded by cages, ALL = saplings from which both ants

and insectivorous vertebrates were excluded.

Figure 3.

Figure 3. Effect of the predator exclusion on mean body size of the arthropods collected from

saplings along the elevational gradient of Mt Wilhelm in the wet (a) and dry season (b). Each

individual data point shows the resulting mean body size of the arthropod community on a given

sapling. The curves show predicted values from the linear model which included season, elevation,

and treatment as a fixed factor. See Table S6 for model results. The occurrence (log of the total

abundance) of arthropods of a given body size in each of the four treatments is shown in (c).

Treatments: CON = control saplings with free access of predators, ANT = saplings from which ants

are excluded by tanglefoot glue, VER = saplings from which insectivorous vertebrates were excluded

by cages, ALL = saplings from which both ants and insectivorous vertebrates were excluded.

Figure 4.

Figure 4. Mean proportion of spiders in control treatments (grey area), insectivorous vertebrate

exclosures (VER - magenta line), ant exclosures (ANT - blue line) and both insectivorous vertebrates

and ants exclosures (ALL- yellow line) across the elevational gradient (a). Caterpillar graph showing

how individual orders of arthropods responded to the predator exclosure treatments. The X-axis

shows mean effect sizes of natural log ratios (LRR = ln(exclosure/control)) with 95% CI of individual

response variables. Effect size values above zero indicated that the absence of predators (treatment)

was more harmful to plants than the control (presence of predators), as the abundances of

arthropods in exclosures increased and caused potentially higher herbivory damage to plants (b).

Treatments: ANT = saplings from which ants are excluded by tanglefoot glue, VER = saplings from

which insectivorous vertebrates were excluded by cages, ALL = saplings from which both ants and

insectivorous vertebrates were excluded.

645 Figure 5.

646 **Figure 5.** Correlations between the effect of vertebrate exclosures (a, b) and ant (c) exclosures (LRR

647 calculated from raw data and normalized data) on abundances of arthropods and abundances

648 (normalized) of insectivorous birds (a), insectivorous bats (b) and ants (c) potentially feeding on the

649 experimental saplings.

Table 1. Corrected Akaike Information Criterion (AICc) of regression models examining abundances of all arthropods, herbivory damage, and mean body size of arthropods observed on saplings in the wet and dry season (season) and excluding various predators (treatment) along the elevational gradient of Mt. Wilhelm (elevation) in Papua New Guinea (a). Results of the analysis of variance of the most parsimonious models selected based on the AIC (b).

	Arthropod abundances		Herbivory		Body sizes	
	dAICc	df	dAICc	df	dAICc	df
Null	216.1	4	102.5	4	162.2	4
Treatment	174.5	7	80.3	7	86	7
Elevation	186.2	6	50.2	6	129.8	6
Season	140	5	75.6	5	167.1	5
Treatment + Season	97.9	8	52.3	8	90.9	8
Treatment + Elevation	145.3	9	25.8	9	48.1	9
Season + Elevation	110.6	7	25.7	7	90.9	8
Treatment : Season	87	11	60.2	11	93.9	11
Treatment : Elevation	133	12	47.7	12	89.6	12
Season : Elevation	165.6	8	52.1	8	106.7	8
Treatment : Season + Elevation	58.5	13	7.4	13	56	13
Treatment : Elevation + Season	57.4	13	22.9	13	94.6	13
Season : Elevation + Treatment	124.7	11	27.9	11	24.9	11
Treatment + Elevation + Season	69.2	10	0.3	10	53	10
Treatment + Elevation + Season + Treatment : Elevation	21.5	16	0	16	23.3	16
Treatment + Elevation + Season + Treatment : Season	61.3	13	5.4	13	56	13
Treatment + Elevation + Season + Elevation : Season	47.7	12	0.8	12	29.8	12
Treatment + Elevation + Season + Elevation : Season + Elevation : Treatment	0	18	0.5	18	0	18
Treatment + Elevation + Season + Elevation : Treatment + Season : Treatment	13.6	19	4.8	19	26.3	19
Treatment + Elevation + Season + Elevation : Treatment + Season : Treatment + Season : Elevation	7.8	21	5.3	21	8	21
Results of the best model	Chisq (df)	P	Chisq (df)	P	Chisq (df)	P
Elevation	24.41 (2)	<0.001	73.18 (2)	<0.001	50.13 (2)	<0.001
Treatment	43.58 (3)	<0.001	30.87 (3)	<0.001	104.21 (3)	<0.001
Season	83.22 (1)	<0.001	28.21 (1)	<0.001	0.23 (1)	0.632
Elevation : Treatment	5.09 (6)	0.532	12.57 (6)	0.051	11.87 (6)	0.065
Elevation : Season	9.78 (2)	0.011			19.95 (2)	<0.001

693   References

694   Bell, H. 1982. Abundance and seasonality of the savanna avifauna at Port Moresby, Papua New

695   Guinea. Ibis **124**:252-274.

696   Bodawatta, K. H., I. Klečková, J. Klečka, K. Pužejová, B. Koane, M. Poulsen, K. A. Jønsson, and K. Sam.

697   2022. Specific gut bacterial responses to natural diets of tropical birds. Scientific Reports

698   **12**:1-15.

699   Bonaccorso, F. J. 1998. Bats of Papua New Guinea. Conservation International.

700   Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J. Skaug, M.

701   Machler, and B. M. Bolker. 2017. glmmTMB balances speed and flexibility among packages

702   for zero-inflated generalized linear mixed modeling. The R journal **9**:378-400.

703   Burger, H., K. Hylander, B. Ayalew, N. van Dam, E. Mendesil, A. Schedl, T. Shimales, B. Zewdie, and A.

704   Tack. 2021. Bottom-up and top-down drivers of herbivory on Arabica coffee along an

705   environmental and management gradient. Basic and Applied Ecology.

706   Chen, I.-C., J. K. Hill, R. Ohlemüller, D. B. Roy, and C. D. Thomas. 2011. Rapid range shifts of species

707   associated with high levels of climate warming. Science **333**:1024-1026.

708   Coley, P. D., and T. M. Aide. 1991. Comparison of herbivory and plant defences in temperate and

709   tropical broad- leaved forests, in plant-animal interactions: Evolutionary ecology in the

710   tropical and temperate regions. John Wiley & Sons Ltd., , Brisbane.

711   Coley, P. D., and J. Barone. 1996. Herbivory and plant defenses in tropical forests. Annual Review of

712   Ecology and Systematics:305-335.

713   Colwell, R. K., N. J. Gotelli, L. A. Ashton, J. Beck, G. Brehm, T. M. Fayle, K. Fiedler, M. L. Forister, M.

714   Kessler, and R. L. Kitching. 2016. Midpoint attractors and species richness: Modelling the

interaction between environmental drivers and geometric constraints. Ecology Letters

19:1009-1022.

Curtis, P. S., and X. Wang. 1998. A meta-analysis of elevated CO_2 effects on woody plant mass, form,

and physiology. Oecologia **113**:299-313.

Defossez, E., L. Pellissier, and S. Rasmann. 2018. The unfolding of plant growth form-defence

syndromes along elevation gradients. Ecology Letters **21**:609-618.

Flannery, T. F. 1995. Mammals of new guinea. Reed.

Garibaldi, L. A., T. Kitzberger, and E. J. Chaneton. 2011. Environmental and genetic control of insect

abundance and herbivory along a forest elevational gradient. Oecologia **167**:117-129.

Gathalkar, G., and A. Sen. 2018. Foraging and predatory activities of ants. The complex world of

ants:51-70.

Greenberg, R., P. Bichier, A. C. Angon, C. MacVean, R. Perez, and E. Cano. 2000. The impact of avian

insectivory on arthropods and leaf damage in some Guatemalan coffee plantations. Ecology

81:1750-1755.

Hedges, L. V., J. Gurevitch, and P. S. Curtis. 1999. The meta-analysis of response ratios in

experimental ecology. Ecology **80**:1150-1156.

Hölldobler, B., and E. O. Wilson. 1990. The Ants Springer Berlin.

Holmes, R. T., R. E. J. Bonney, and S. W. Pacala. 1979. Guid structure of the Hubbard Brook bird

community: a multivariate approach. Ecology **60**:512-520.

Hood, A. S., A. A. K. Aryawan, A. D. Advento, W. R. Suberkah, A. Ashton-Butt, S. Ps, J. P. Caliman, M.

Naim, W. A. Foster, and E. C. Turner. 2022. A whole-ecosystem method for experimentally

suppressing ants on a small scale. Methods in Ecology and Evolution **13**:852-865.

Hoyo, d. J., J. Elliott, J. Sargatal, and D. A. Christie, editors. 1992-2011. Handbook of the Birds of the

World (vol. 1-16). Lynx Editions, Barcelona, Spain.

Ingala, M. R., N. B. Simmons, C. Wultsch, K. Krampis, K. L. Provost, and S. L. Perkins. 2021. Molecular

diet analysis of neotropical bats based on fecal DNA metabarcoding. Ecology and evolution

11:7474-7491.

Kalka, M., and E. K. Kalko. 2006. Gleaning bats as underestimated predators of herbivorous insects:

diet of Micronycteris microtis (Phyllostomidae) in Panama. Journal of Tropical Ecology **22**:1-

10.

Kalka, M. B., A. R. Smith, and E. K. V. Kalko. 2008. Bats Limit Arthropods and Herbivory in a Tropical

Forest. Science **320**:71.

Karp, D. S., and G. C. Daily. 2014. Cascading effects of insectivorous birds and bats in tropical coffee

plantations. Ecology **95**:1065-1074.

Kerbiriou, C., Y. Bas, I. Le Viol, R. Lorrilliere, J. Mougnot, and J. F. Julien. 2019. Potential of bat pass

duration measures for studies of bat activity. Bioacoustics **28**:177-192.

Liu, X., Z. Wang, C. Huang, M. Li, F. Bibi, S. Zhou, and A. Nakamura. 2020. Ant assemblage

composition explains high predation pressure on artificial caterpillars during early night.

Ecological Entomology **45**:547-554.

Marki, P. Z., K. Sam, B. Koane, J. B. Kristensen, J. D. Kennedy, and K. A. Jønsson. 2016. New and

noteworthy bird records from the Mt. Wilhelm elevational gradient, Papua New Guinea.

Bull. Brit. Ornitholog. Club **137**:263-271.

Marquis, R. J., and C. Whelan. 1996. Plant morphology and recruitment of the third trophic level:

subtle and little-recognized defenses? Oikos:330-334.

McAlpine, J. R., R. Keig, and R. Falls. 1983. Climate of Papua New Guinea. CSIRO and Australian

National University Press, Canberra.

McCain, C. M. 2005. Elevational gradients in diversity of small mammals. Ecology **86**:366–372.

Mestre, L., J. Piñol, J. Barrientos, A. Cama, and X. Espadaler. 2012. Effects of ant competition and

bird predation on the spider assemblage of a citrus grove. Basic and Applied Ecology **13**:355-

362.

Milne, D. J., C. J. Burwell, and C. R. Pavey. 2016. Dietary composition of insectivorous bats of the Top

End of Australia. Australian Mammalogy **38**:213-220.

Mols, C. M., and M. E. Visser. 2002. Great tits can reduce caterpillar damage in apple orchards.

Journal of Applied Ecology **39**:888-899.

Moon, D. C., and P. Stiling. 2004. The influence of a salinity and nutrient gradient on coastal vs.

upland tritrophic complexes. Ecology **85**:2709-2716.

Mooney, K. A. 2007. Tritrophic effects of birds and ants on a canopy food web, tree growth, and

phytochemistry. Ecology **88**:2005-2014.

Mooney, K. A., D. S. Gruner, N. A. Barber, S. A. Van Bael, S. M. Philpott, and R. Greenberg. 2010.

Interactions among predators and the cascading effects of vertebrate insectivores on

arthropod communities and plants. Proceedings of the National Academy of Sciences.

Mooney, K. A., and Y. B. Linhart. 2006. Contrasting cascades: insectivorous birds increase pine but

not parasitic mistletoe growth. Journal of Animal Ecology **75**:350–357.

Moses, J., T. M. Fayle, V. Novotny, and P. Klimes. 2021. Elevation and leaf litter interact in

determining the structure of ant communities on a tropical mountain. BIOTROPICA **53**:906-

919.

Nordberg, E. J., and L. Schwarzkopf. 2019. Predation risk is a function of alternative prey availability

rather than predator abundance in a tropical savanna woodland ecosystem. Scientific

Reports **9**:1-11.

Novotny, V., and Y. Basset. 1998. Seasonality of sap-sucking insects (Auchenorrhyncha, Hemiptera)

feeding on Ficus (Moraceae) in a lowland rain forest in New Guinea. Oecologia **115**:514-522.

Nyffeler, M., Ç. H. Şekercioğlu, and C. J. Whelan. 2018. Insectivorous birds consume an estimated

400–500 million tons of prey annually. The Science of Nature **105**:47.

Paijmans, K., editor. 1976. New Guinea Vegetation. National University Press, Canberra.

Pellissier, L., A. Roger, J. Bilat, and S. Rasmann. 2014. High elevation Plantago lanceolata plants are

less resistant to herbivory than their low elevation conspecifics: is it just temperature?

Ecography **37**:950-959.

Perfecto, I., and J. Vandermeer. 1996. Microclimatic changes and the indirect loss of ant diversity in

a tropical agroecosystem. Oecologia **108**:577-582.

Philpott, S., I. Perfecto, and J. Vandermeer. 2008. Effects of predatory ants on lower trophic levels

across a gradient of coffee management complexity. Journal of Animal Ecology **77**:505-511.

Philpott, S. M., and I. Armbrecht. 2006. Biodiversity in tropical agroforests and the ecological role of

ants and ant diversity in predatory function. Ecological Entomology **31**:369-377.

Philpott, S. M., R. Greenberg, and I. P. Bichier. 2004. Impacts of major predators on tropical

agroforest arthropods: comparisons within and across taxa. Oecologia:140-149.

Poulin, B., G. Lefebvre, and R. McNeil. 1994. Diets of Land Birds from Northeastern Venezuela. The

Condor **96**:354-367.

Pratt, T. K., and B. M. Beehler. 2015. Birds of New Guinea. Princeton University Press.

Rahbek, C. 1995. The elevational gradient of species richness: a uniform pattern? Ecography **18**:200-

205.

Richards, L. A., and P. D. Coley. 2007. Seasonal and habitat differences affect the impact of food and

predation on herbivores: a comparison between gaps and understory of a tropical forest.

Oikos **116**:31-40.

Risch, S. J., and C. R. Carroll. 1982. Effect of a keystone predaceous ant, Solenopsis geminata, on

arthropods in a tropical agroecosystem. Ecology:1979-1983.

Roslin, T., B. Hardwick, V. Novotny, W. K. Petry, N. R. Andrew, A. Asmus, I. C. Barrio, Y. Basset, A. L.

Boesing, and T. C. Bonebrake. 2017. Higher predation risk for insect prey at low latitudes and

elevations. Science **356**:742-744.

Salgado, A. L., T. Suchan, L. Pellissier, S. Rasmann, A.-L. Ducrest, and N. Alvarez. 2016. Differential

phenotypic and genetic expression of defence compounds in a plant–herbivore interaction

along elevation. Royal Society Open Science **3**:160226.

Sam, K., and B. Koane. 2014. New avian records along the elevational gradient of Mt. Wilhelm,

Papua New Guinea. Bulletin of the British Ornithologists' Club **134**:116-133.

Sam, K., B. Koane, D. C. Bardos, S. Jeppy, and V. Novotny. 2019. Species richness of birds along a

complete rain forest elevational gradient in the tropics: Habitat complexity and food

resources matter. Journal of Biogeography **46**:279-290.

Sam, K., B. Koane, S. Jeppy, J. Sykorova, and V. Novotny. 2017. Diet of land birds along an elevational

gradient in Papua New Guinea. Scientific Reports **7**: **44018**.

Sam, K., B. Koane, and V. Novotny. 2015. Herbivore damage increases avian and ant predation of

caterpillars on trees along a complete elevational forest gradient in Papua New Guinea.

Ecography **38**:293-300.

Sam, K., B. Koane, L. Sam, A. Mrazova, S. Segar, M. Volf, M. Moos, P. Simek, M. Sisol, and V. Novotny.

2020. Insect herbivory and herbivores of Ficus species along a rain forest elevational

gradient in Papua New Guinea. BIOTROPICA **52**:263-276.

Sam, K., Houska Tahadlova, M., Mrazova, A., Freiberga, I., Toszogyova, A., Sreekar, R. 2022. The

impact of ants and vertebrate predators on arthropods and plants: a meta-analysis. -

BIORXIV/2022/498005

Schemske, D. W., G. G. Mittelbach, H. V. Cornell, J. M. Sobel, and K. Roy. 2009. Is there a latitudinal

gradient in the importance of biotic interactions? Annual Review of Ecology, Evolution, and

Systematics **40**:245-269.

Schmidt, M., and A. Dejean. 2018. A dolichoderine ant that constructs traps to ambush prey

collectively: convergent evolution with a myrmicine genus. Biological Journal of the Linnean

Society **124**:41-46.

Schmitz, O. J. 2006. Predators have large effects on ecosystem properties by changing plant

diversity, not plant biomass. Ecology **87**:1432-1437.

Sekercioglu, C. H. 2006. Increasing awareness of avian ecological function. Trends in Ecology &

Evolution **21**:464-471.

Shurin, J. B., E. T. Borer, E. W. Seabloom, K. Anderson, C. A. Blanchette, B. Broitman, S. D. Cooper,

and B. S. Halpern. 2002. A cross-ecosystem comparison of the strength of trophic cascades. .

Ecology Letters **5**:785–791.

Sih, A., G. Englund, and D. Wooster. 1998. Emergent impacts of multiple predators on prey. Trends in

Ecology & Evolution **13**:350-355.

Singer, M., R. Clark, I. Lichter-Marck, E. Johnson, and K. Mooney. 2017. Predatory birds and ants

partition caterpillar prey by body size and diet breadth. Journal of Animal Ecology.

Sivault, E., P. K. Amick, K. N. Armstrong, V. Novotny, and K. Sam. 2022. Species richness and

community structure of bats along a forest elevational transect in Papua New Guinea.

bioRxiv.

Suzuki, S., K. Kitayama, S.-i. Aiba, M. Takyu, and K. Kikuzawa. 2013. Annual leaf loss caused by

folivorous insects in tropical rain forests on Mt. Kinabalu, Borneo. Journal of Forest Research

18:353-360.

Tanhuanpää, M., K. Ruohomäki, and E. Uusipaikka. 2001. High larval predation rate in

non-outbreaking populations of a geometrid moth. Ecology **82**:281-289.

Team, R. C. 2020. R: a language and environment for statistical computing. Version 4.0. 2. Vienna,

Austria.

Tvardikova, K. 2013. Trophic relationships between insectivorous birds and insect in Papua New

Guinea. University of South Bohemia, Ceske Budejovice.

Van Bael, S. A., J. D. Brawn, and S. K. Robinson. 2003. Birds defend trees from herbivores in a

Neotropical forest canopy. Proceedings of the National Academy of Sciences **100**:8304-8307.

Van Bael, S. A., S. M. Philpott, Greenberg. R., P. Bichier, N. A. Barber, K. A. Mooney, and D. S. Gruner.

2008. Birds as predators in tropical agroforestry systems. Ecology **89**:928-934.

Véle, A., J. Holuša, and J. Frouz. 2009. Sampling for ants in different-aged spruce forests: a

comparison of methods. European journal of soil biology **45**:301-305.

Way, M., and K. Khoo. 1992. Role of ants in pest management. Annual Review of Entomology

37:479-503.

Williams-Guillén, K., I. Perfecto, and J. Vandermeer. 2008. Bats limit insects in a neotropical

agroforestry system. Science **320**:70-70.

Wright, D. D., J. H. Jessen, P. Burke, and H. G. de Silva Garza. 1997. Tree and Liana Enumeration and

Diversity on a One-Hectare Plot in Papua New Guinea 1. BIOTROPICA **29**:250-260.

873