Title: Selection for greater dispersal in early life leads to faster age-dependent decline in
locomotor activity and shorter lifespan

Authors: B G Ruchitha[1,2], Nishant Kumar[1], Chand Sura[1], Sudipta Tung[1]*

Affiliations:

1. Integrated Genetics and Evolution Laboratory (IGEL), Department of Biology,
Ashoka University, Sonipat, Haryana, India, 131029.

2. Indian Institute of Science Education and Research (IISER) Pune, Pune, Maharashtra,
India, 411008.
9
1. ruchitha.bg@students.iiserpune.ac.in
2. nishant.kumar@ashoka.edu.in
3. chand@ashoka.edu.in
4. sudipta.tung@ashoka.edu.in [ORCID 0000-0002-3201-3303]

* **Name and address of the corresponding author:**

Sudipta Tung
DBT/Wellcome Trust India Alliance Early Career Fellow
Department of Biology, Ashoka University
Plot No. 2, Rajiv Gandhi Education City,
Post Office Rai, Near Rai Police Station,
Sonipat, Rai, Haryana, India 131029
21
22 Tel: +91-130 2300669
23 Email: sudipta.tung@ashoka.edu.in
24

Acknowledgements:
The authors thank Prof. Sutirth Dey from Indian Institute of Science Education and Research
(IISER)-Pune for kindly providing us the experimental populations. The authors also thank
Prof. Dey and Akhila Mudunuri for their valuable comments on an earlier version of the
manuscript. RBG acknowledges the support from Kishore Vaigyanik Protsahan Yojana
(KVPY) fellowship, NK and CS acknowledge the support of the Research and Development
Office, Ashoka University. ST acknowledges the support of DBT/Wellcome Trust India
Alliance Early Career Fellowship (#IA/E/18/1/504347) and Ashoka University.

Conflict of interest:
The authors declare no conflict of interest.

Abstract

Locomotor activity is one of the major traits that is affected by age. Greater locomotor activity is also known to evolve in the course of dispersal evolution. However, the impact of dispersal evolution on the functional senescence of locomotor activity is largely unknown. We addressed this knowledge gap using large outbred populations of *Drosophila melanogaster* selected for increased dispersal. We tracked locomotor activity of these flies at regular intervals until a late age. Longevity of these flies was also recorded. We found that locomotor activity declines with age in general. However interestingly, activity level of dispersal selected populations never drops below the ancestry-matched-controls, despite the rate of age-dependent decline in activity of the dispersal selected populations being greater than their respective controls. Dispersal selected population was also found to have shorter lifespan as compared to its control, a potential cost of elevated level of activity throughout their life. These results are crucial in the context of invasion biology as contemporary climate change, habitat degradation, and destruction provide congenial conditions for dispersal evolution. Such controlled and tractable studies investigating the ageing pattern of important functional traits are important in the field of biogerontology as well.

Introduction

Dispersal is a central life-history trait (Bonte and Dahirel 2017). At an individual level, it can confer survival advantage against proximate environmental stresses (Studds et al. 2008), as well as reproductive advantage through a greater prospect of finding mates and suitable breeding grounds in the case of sexually reproducing organisms (Greenwood and Harvey 1982, Studds et al. 2008). At the population level, it modulates the rate and extent of adaptive dynamics by affecting the degree of gene flow between populations (Lenormand 2002, Garant et al. 2007). Dispersal also shapes local ecology by influencing processes such as territoriality, range expansion, species invasion, etc. (Hargreaves and Eckert 2014, Andrade-Restrepo et al. 2019).

It has been shown through both theoretical and empirical evidence that the dispersal abilities of individuals can evolve rather quickly (Perkins et al. 2013, Huang et al. 2015, Ochocki and Miller 2017, Tung et al. 2018a) in the presence of environmental conditions like climate change (Travis et al. 2013), habitat fragmentation (McPeek and Holt 1992, Berg et al. 2010), and habitat degradation and destruction (Fronhofer et al. 2014). These conditions are prevalent in the contemporary times due to, *inter alia*, climate change, human-induced landscape changes, and habitat destruction. Notably, dispersal is known to have strong associations with a number of traits of an individual, such as locomotor activity, aggression, body size, boldness, etc. (Wahlström 1994a, Michelangeli et al. 2017, Searcy et al. 2018, McGhee et al. 2021, Wu and Seebacher 2022), which are collectively known as dispersal syndrome (Clobert et al. 2009). Consequently, these traits also often evolve as a result of dispersal evolution (Fjerdingstad et al. 2007, Tung et al. 2018b). Such changes in organismal trait distributions can lead to the evolution of organisms with combinations of dispersal and dispersal syndromic traits with significantly higher potential for range expansion and

biological invasion, and thereby threaten local ecological stability (Lambrinos 2004, Perkins

et al. 2013, Renault et al. 2018). Thus, not surprisingly, investigating different aspects of

dispersal evolution is a prime focus among contemporary ecologists, evolutionary biologists,

and conservation biologists alike.

Despite the evident advantages that greater dispersal ability offers, such as avoiding local

stressors, enhancing foraging opportunities, and facilitating mate finding, the diversity in

dispersal properties among organisms in nature remains puzzling. This diversity can be

understood by considering the costs associated with dispersal (reviewed in (Bonte et al.

2012)). These costs primarily arise from the energy-intensive nature of the dispersal process.

Previous research involving pre- and post-dispersal analyses, has demonstrated that the

dispersal process exacts a significant toll on an organism's physiology (Mishra et al. 2022)

and increases stress levels (Maag et al. 2019), although their long-term impacts across

generations remain unclear. It follows logically that these physiological impacts would

influence organismal performance later in life by accelerating the age-dependent decline in

physiological functionality, also known as functional senescence of organismal traits

(Grotewiel et al. 2005). Accelerated functional aging could diminish motor ability of an

organism, thereby reducing its capacity to disperse and colonize new areas in later life, and

may also alter age-dependent reproductive efforts and survival. For instance, functional

senescence might result in decreased capacity for long-distance travel or increased

vulnerability to predation during dispersal (Greenwood and Harvey 1982, Harvey et al. 1984,

Newton 1993, Badyaev and Faust 1996, Forero et al. 1999, Pyle et al. 2001, Andreu and

Barba 2006). These changes can significantly impact subsequent evolution of dispersal traits,

thus providing a potential explanation for why we see diversity in dispersal properties among

organisms in nature. Hence, it's crucial to incorporate an understanding of senescence into

studies on the evolution of dispersal, particularly concerning the functional senescence of

traits associated with dispersal, or dispersal syndrome. Failure to do so could result in

incomplete or inaccurate conclusions about the evolution of dispersal and its associated traits.

However, the costs of dispersal evolution and its trade-offs with the functional senescence of

dispersal syndromic traits are rarely investigated.

Locomotor activity is a prominent dispersal syndromic trait with strong association with

greater dispersal ability across multiple species (Hanski et al. 2006; Cote et al. 2010;

Fronhofer et al. 2015; Kosmala et al. 2017; Mishra et al. 2018). It is known to evolve as a

correlated response to dispersal evolution (Tung et al. 2018b). However, the impact of this

evolutionary adaptation on the functional senescence of locomotor activity remains unclear.

Moreover, the relationship between age and locomotor activity varies among species. For

example, some animals, such as senescence-accelerated mice (Miyamoto et al. 1986) and

rabbits in open-field tests (Deyo et al. 1989) show an increase in activity with age; humans,

mice, rhesus monkeys and some rats show a decrease in activity with age (Goodrick 1971,

Dean et al. 1981, Emborg et al. 1998, McGibbon and Krebs 2004). In contrast, no significant

change in activity with age has been seen in cats and some rats (Levine et al. 1987, Nagahara

and Handa 1997). In the case of the insect model *Drosophila melanogaster*, (Le Bourg 1987)

has shown an age-dependent decrease in activity in females, and an increase in activity in

males till 5 weeks of age, after which activity declines. However, some studies have shown

that such sexual dimorphism in the age-dependent activity of *Drosophila* is strongly

influenced by the genotype of the flies used (van Dijken et al. 1979, Fernández et al. 1999).

Therefore, it is not yet clear how mobility changes with age in sexually dimorphic organisms,

such as fruit flies *Drosophila melanogaster*. Given that males and females of sexually

dimorphic organisms tend to defer with respect to their resource allocation strategies in

reproduction and somatic maintenance as well as differ in metabolic rates, it is likely that

selection for dispersal ability can shape functional senescence in males and females

differently (Garratt 2020). However, there is limited understanding of how evolution for

increased dispersal ability affects different sexes at various ages.

In this study, we aimed to fill this knowledge gap by investigating locomotor activity and

lifespan of male and female individuals from four large, outbred populations selected for

greater dispersal ability and propensity (Tung et al. 2018a). We examined the locomotor

activity of the populations selected for greater dispersal and their ancestry-matched controls

after maintaining both types of populations under identical conditions for nine generations, or

approximately five months, without any further selection for dispersal. The age-dependent

locomotor activity patterns of the evolved and control populations are compared for males

and females at multiple time points across the lifespan to understand the impact of dispersal

evolution on the functional senescence of locomotor activity in these flies. We began by

analysing the aging profile in a cross-sectional study to understand how the trait has changed

in these populations. Next, considering variation in lifespan among individuals within a

population and to eliminate potential survivorship bias because of it, we conducted a

longitudinal study by following flies individually and measuring their locomotor activity

across age. Additionally, in this longitudinal study, we compared the longevity of the evolved

and control populations to understand the impact of functional aging of locomotor activity on

demographic aging by measuring lifespan.

Through these experiments, we provide insights on functional senescence of locomotor

activity and demographic senescence in the two sexes and how the evolution of dispersal

affects them. These aspects are particularly relevant as locomotor activity plays a critical role

in reproduction, encompassing activities such as mate searching and successful courtship for

males, as well as locating suitable egg-laying surfaces for females, ultimately influencing

their post-dispersal colonization success.

Methods

Experimental Populations

We used four large (n = ~2400), outbred, laboratory-maintained populations of *Drosophila*

melanogaster (VB_{1-4}) that were selected for greater dispersal propensity and ability for 153

generations (>6 years), and their respective ancestry-matched control (VBC_{1-4}) populations.

For clarity and ease of reading, henceforth, we will refer to the VB populations as 'dispersal-

selected', 'selected', or 'D' populations interchangeably. Similarly, the VBC populations will

be referred to as 'control' or 'C' populations. These populations were derived from a common

ancestry, therefore had comparable genetic architecture, and were maintained identically

under standard laboratory conditions. Details of the selection protocol and maintenance

regime of these populations were reported previously (Tung et al. 2018a). Briefly, the D_{1-4}

and C_{1-4} populations were maintained in 15-day discrete generation cycles on a standard

banana-jaggery medium (following the recipe of Sheeba and Joshi 1998) at 25°C, with

constant light. On the 12^{th} day from the day of egg collection, only the most dispersive half of

the population were artificially selected. This was done by subjecting the D_{1-4} populations to

a three-compartment source-path-destination dispersal selection setup, where the flies from

D_{1-4} populations were placed in the source compartment, allowed to disperse via the path to

reach the destination compartment. Among these flies, only the first ~50% of the population

estimated visually to reach the destination compartment were allowed to lay eggs for the next

generation. To maintain the consistent breeding population size of the D_{1-4} populations across

generations while also not biasing the density of the population in the source compartment,

two such source-path-destination setups would be created for each population of the D_{1-4}

populations. Thus, collecting approximately the first 50% of the population from each of the

two source-path-destination setups ensured a consistent breeding population size in the

selected populations comparable to that of the corresponding control populations. These

populations were maintained at a high population size of approximately 2400 flies per

population to minimize the impact of inbreeding. C_{1-4} populations were not subjected to

selection for dispersal, but the rest of the rearing conditions were identical to that of the D_{1-4}

populations. To obtain the next generation of flies, in both D and C populations, eggs were

collected on the 15th day post egg collection, ensuring that no differential age-dependent

selection is acting on the D and C populations.

Experiments

Experiment 1: For investigating population level effect

Prior to this experiment, selection pressure for dispersal was relaxed for eight generations i.e.

D_{1-4} and C_{1-4} populations were maintained identically without selection pressure to minimize

non-genetic parental effects. ~300 eggs were collected in drosophila-bottles (Laxbro®

#FLBT20) containing 50mL banana-jaggery food. Eight such replicates were made for each

population. On 11th day from egg collection, all flies from eight bottles of same population

were pooled into a single cage (25 cm × 20 cm × 15 cm) and maintained with fresh food

being provided every alternate day. This was done for each of the eight populations (D_{1-4} and

C_{1-4}). From each of these population cages, 32 flies of each sex were randomly sampled at

predetermined age-points (see "Assay Details" below). Overall, locomotor activity of 2048

flies (32 flies × 2 sexes × 8 populations × 4 age-points) was measured for this experiment.

Figure S1 shows a schematic diagram of the experimental design for investigating population

level effect.

Experiment 2: For tracking individual flies

In experiment 2, we collected longitudinal data as opposed to the cross-sectional data

gathered in experiment 1. Since the lifespans of individuals can vary within a population,

cross-sectional activity data can be affected by survivorship bias, which can influence the

interpretation of results. Therefore, experiment 2 was designed as a proof-of-principle

validation of the findings from experiment 1, and due to logistical constraints was conducted

using one set of ancestry-matched selected and control populations (D_2 and C_2), chosen at

random.

This experiment was performed on the dispersal selected population, after relaxation of

selection pressure for dispersal over nine generations, and its ancestry matched control. In

order to generate experimental flies for each population, ~300 eggs were collected in

drosophila-bottles (Laxbro® FLBT20) containing 50mL banana-jaggery food. Three such

replicates were made for each population. From the three bottles, 30 individuals of each sex

were chosen at random during the P13-P15 stage of pupa and hosted individually in glass

vials containing ~4mL of food and incubated at 25°C. On eclosion, these 120 virgin adult

flies (*i.e.* 30 individuals per sex per population) were checked daily for survival until death,

and their age-dependent locomotor activity was tracked by measuring the trait at

predetermined age-points (see "Assay Details" below). The flies were provided fresh food

every alternate day, while strictly maintaining their vial identity. Figure S2 shows a

schematic diagram of the experimental design for tracking individual flies.

Assay Details

Locomotor activity assay (used for experiments 1 and 2)

Locomotor activity was measured as the number of times per hour that each fly present in a

glass tube (diameter: 5 mm, length: 8 cm) crossed the middle of the tube. This was done

using Drosophila Activity Monitoring (DAM2) data collection system (Trikinetics Inc,

Waltham, MA) which recorded the number of times each fly would cross the infrared beams

at the centre of the glass tube every 30 seconds using a standard protocol (Chiu et al. 2010).

To minimize any potential bias from variation in movement patterns of the flies, all

treatments in our experiment were handled consistently.

For experiment 1, we tracked locomotor activity of four independent but identically

maintained, large, outbred, dispersal-selected populations D_{1-4} across age and contrasted the

same against corresponding four ancestry-matched control populations C_{1-4}. To do this, we

randomly sampled 32 flies of each sex from each of these eight populations (~2400 adults on

the 12^{th} day in each population) on the 12^{th}, 26^{th}, 40^{th}, and 47^{th} day from the day of egg

collection and measured their locomotor activity. On the specified days of activity

measurement, flies were collected from population cages and introduced into clean glass

tubes (diameter: 5 mm, length: 8 cm), without CO_2 anaesthesia. Each glass tube was secured

with cotton plugs on both sides. Activity was recorded for 2h and 5mins, and the initial 5 min

was excluded to minimize the effect of disturbance caused at the time of setup. At the end of

the recording, mortality was recorded and all the flies in the glass tubes were discarded.

Overall, for experiment 1, locomotor activity of 256 flies (32 flies $\times$ 2 sexes $\times$ 4 age-points)

of each population was measured.

For experiment 2, we repeatedly measured locomotor activity of same set of 30 male and 30

female flies each of dispersal-selected and control populations on the 12^{th}, 26^{th}, 33^{rd}, 40^{th}, and

47^{th} day from the day of egg collection. On the specified days of activity measurement, flies

were introduced into individually labelled (to maintain identity of each fly), clean glass tubes

(diameter: 5 mm, length: 8 cm), without CO_2 anaesthesia. Each glass tube contained 1.5%

w/vol agar, and was sealed with parafilm on one end and secured with cotton plug on the

other. Activity was recorded for 4h and 5mins, and the initial 5 min was excluded to

minimize the effect of disturbance caused at the time of setup. At the end of the recording,

the flies were transferred back to the corresponding individually labelled glass vials (to

maintain identity of each fly) with food. Overall, for experiment 2, 60 flies (30 flies × 2

sexes) of each population were individually tracked by maintaining them in labelled

containers and their locomotor activity was measured at five pre-determined age-points.

Longevity assay (used for experiment 2)

Longevity of each fly was assessed as the number of days it was alive. Each day, we

manually looked at each of the flies that were being individually tracked (from experiment 2)

until all flies were dead. Complete absence of movement of limbs, abdomen and proboscis

was recorded as death.

Statistical Analysis

Locomotor activity assay of experiment 1

We disregarded the data of the flies which were found dead at the end of the recording

(details of sample size is present in Table S1). For all the remaining flies, out of the 2 hours

of recording, we only considered the first 1 hour of data. We did this to minimize the effect of

potential artifacts generated by inactivity due to death and hyperactivity due to starvation (Yu

et al. 2016). From the data obtained after these corrections, we calculated the average of one

hour locomotor activity of 32 flies for each of 16 treatment groups (2 sexes × 8 populations)

across 4 age-points.

To test how evolution of greater dispersal affects age-dependent locomotor activity of

dispersal selected population, we analysed the locomotor activity data using a linear mixed

effects model with Selection (dispersal-selected vs. control), Sex (male vs. female), Age (12,

26, 40, and 47) as fixed factors, and population identity (1, 2, 3 and 4) as random factor. We

used 'lmer' function from 'lme4' package in R v4.3.2 with a model *Activity ~ Selection+*

*Sex+ Age + (1|Block) + Selection*Sex + Selection*Age + Sex*Age*. When the main or

interaction effect was significant, post-hoc was performed using Tukey's HSD test using

'emmeans' function from 'lsmeans' package in R v4.3.2.

Locomotor activity assay of experiment 2

The 4h locomotor activity data was used to calculate per hour activity. In order to distinguish

between within-individual variation in locomotor activity across age and between-individual

variation in locomotor activity resulting from differences in longevity, we divided age into

two components: Delta age and Average age, respectively (van de Pol and Wright 2009).

Here, Average age represents the mean of all timepoints for an individual, while Delta age

reflects timepoint specific deviations from the average. Now, to test how evolution of greater

dispersal affects locomotor activity of individual flies (*i.e.* here, we measure longitudinal data

as opposed to the cross-sectional data obtained from experiment 1) with age, the per hour

activity data was subjected to a linear mixed effects model with Selection (dispersal-selected

vs. control), Sex (male vs. female), Average age and Delta age as fixed factors, and identity

of individual flies as random factor. If a significant Selection $\times$ Delta age interaction is

observed, it would imply that the selection for increased dispersal is associated with the

selection for accelerated locomotor senescence.

Longevity assay of experiment 2

To check whether there is change in lifespan due to evolution of greater dispersal, we

performed a full-factorial 2-way fixed factor ANOVA on longevity data (measured as

mention in "Assay Details" section above) with Selection (dispersal-selected vs. control) and

Sex (male vs. female) as fixed factors using STATISTICA® v5 (StatSoft. Inc., Tulsa,

Oklahoma) software.

Additionally, we conducted a comparison of survivorship curves employing a Cox

proportional-hazards model. The 'Surv' function from the 'survival' package in R v4.3.2 was

used to construct the survival object, incorporating both time and event status. A full factorial

model, incorporating Selection (dispersal-selected vs. control) and Sex (male vs. female) as

fixed factors, along with their interaction was fitted using the 'coxph' function from the same

R package, and summary statistics were extracted. The proportional hazards assumption in

the Cox proportional hazards model was assessed graphically by checking the parallelism of

the log-minus-log survival plots created using the 'survfit' and 'plot' functions in R. The

relatively parallel lines suggest that the proportional hazards assumption is reasonably met

(Figure S3).

To discern whether dispersal-selection impacts longevity through baseline mortality or age-

dependent mortality rate, we fitted a two-parameter Gompertz function using the 'flexsurvreg'

function from the 'flexsurv' package in R v4.3.2. We conducted separate analyses for males

and females in both the selected and control populations. In this analysis, the shape and rate

parameters correspondingly signify the baseline mortality early in life and the age-dependent

mortality rate for each group of individuals.

To determine if there is a correlation between locomotor activity and lifespan within

treatment groups, fitted a linear model for lifespan with Average locomotor activity,

Selection, Sex and their interaction as fixed effects (*Longevity ~ Selection*Sex*AvgActivity*)

using 'lm' function from R v4.3.2. The same analysis was also repeated with locomotor

activity measured at the first time point as a fixed factor instead of Average locomotor

activity of the individuals.

All the graphs were generated using *matplotlib* (Version: 3.5.1) module in Python 3.8.3

(https://www.python.org/).

 Dispersal evolution leads to greater rate of functional senescence

Figure 1. Age-and sex-dependent locomotor activity profile from cross-sectional data. (A) Mean (± 95% CI) locomotor activity of populations from dispersal-selected D (circle) and its ancestry-matched control C (triangle) populations are plotted against age, measured in days from the time of egg collection. Activity level of each population at the tested age-point of D and C populations are represented as blue and red scatter points respectively. Some of the error bars are not visible due to their small size. *p* value for Selection × Age interaction is mentioned in red font. (B) Box-plot representing locomotor activity of dispersal-selected D and its ancestry-matched control populations across males and females. The edges of the box denote 25th and 75th percentiles, while the black solid line represents the median. The blue and red scatter points represent the data for all the replicates of D and C populations respectively. *p* value for Selection × Sex interaction is mentioned in red font. Different lower-case alphabets denote statistically significant differences (p<0.05).

In the cross-sectional study (experiment 1) of locomotor activity across age, although activity

declines in both dispersal selected populations and their ancestry-matched controls, we found

a significant interaction between Age and Selection (Figure 1A; Age × Selection $p = 2 \times 10^{-22}$)

with faster decline in locomotor activity in the dispersal-selected populations than in the

ancestral control populations. This observation is further validated, when we tracked

locomotor activity of individual flies from both dispersal-selected and control populations

across age in an independent longitudinal study i.e., experiment 2 (Figure 2A). For the

analysis of this data, we split age into a Delta age and an Average age component to delineate

the effect of within-individual variation in locomotor activity across age and between-

individual variation in locomotor activity due to difference in longevity, respectively. We

found a significant Selection × Delta age interaction (Figure S5, $p = 2\times10^{-11}$) demonstrating

that selection for greater dispersal is accompanied with selection for accelerated locomotor

senescence. Further, we found that the main effect of Average age ($p = 0.4$) and Selection ×

Average age interaction (Figure S6, $p = 0.68$) are not significant, indicating that the above

observation is not due to any variation in longevity between the individuals or survivorship

bias in the experiment. This is further validated as we found that there is no significant

correlation between average locomotor activity and longevity of the individuals ($F_{1, 103}$ =

0.07, $p = 0.79$) highlighting that increased locomotor activity is not linked with reduced

lifespan within treatment groups. The result remained same when we evaluated correlation

between early-life activity with longevity of the individuals ($F_{1, 103} = 0.01, p = 0.91$).

Interestingly, our cross-sectional analysis showed a significant Age × Sex interaction (Figure

S4, p = 5×10^{-8}). Additionally, our longitudinal investigation revealed a significant Sex ×

Average age interaction ($p = 0.02$), whereas the Sex × Delta age interaction ($p = 0.33$) was

not significant. This suggests that the observed sex-specific variation in locomotor activity

across age points predominantly stem from variations in lifespan between sexes rather than

differences in the rate of senescence of locomotor activity.

Despite elevated functional senescence, dispersal selected flies have high locomotor

activity across age

Figure 2. Age-and sex-dependent locomotor activity profile from longitudinal data.
(A) Mean (± 95% CI) locomotor activity of individuals from dispersal selected population D
(circle) and its ancestry-matched control population C (triangle) are plotted against age, measured
in days from the time of egg collection. Activity level of each individual at the tested age-point of
dispersal selected and control populations are represented as red and blue scatter points
respectively. Scatter points for same individual are connected by solid line. (B) Box-plot
representing locomotor activity of individuals from dispersal-selected D and its ancestry-matched
control C populations across males and females. The blue and red scatter points represent the data
for all the replicates of D and C populations respectively. The edges of the box denote 25^{th} and
75^{th} percentiles, while the black solid line represents the median. p value for Selection × Sex
interaction is mentioned in red font. Different lower-case alphabets denote statistically significant
differences (p<0.05). Some of the error bars are not visible due to their small size.

We see that despite dispersal evolution causing greater rate of decline in locomotor activity

with age (as mentioned above), the locomotor activity of the dispersal selected flies does not

drop below the activity level of their ancestry-matched control flies even at late age in both

cross-sectional (experiment 1, Figure 1A and Table S1) and longitudinal (experiment 2,

Figure 2A and Table S2) studies. This observation is consistent across both males and

females. However, interestingly, the increment in locomotor activity in selected males was

greater than that in females, leading to a significant Selection × Sex interaction in both the

experiment 1 (Figure 1B, $p = 1 \times 10^{-4}$) and experiment 2 (Figure 2B, $p = 3 \times 10^{-6}$). A

comprehensive table containing mean activity levels and p values for experiment 1 and

experiment 2 can be found in Table S1 and Table S2.

Selection for greater dispersal properties trades-off with lifespan

Figure 3. Survivorship curve and lifespan of dispersal-selected and control populations. (A) Survivorship probability (i.e. proportion of viable individuals) of dispersal-selected D (blue solid line) and control C (red dashed line) populations across time in days. *p* value for the main effect of Selection as per the Cox proportional hazard model is mentioned in red font. (B) In the box plot, the middle line, lower and upper edge of the box, lower and upper end of the whiskers represent mean, 25[th], 75[th], 5[th], 95[th] percentile of the longevity data respectively. The scatter points denote longevity of each individual of the dispersal-selected D (blue circles) and control C (red triangles) populations. ANOVA *p* value for the main effect of Selection is mentioned in red font.

In experiment 2, as we tracked each fly individually, we were able to measure the lifespan of

these flies. From this data, we see that the dispersal selected flies live significantly shorter

(Figure 3 and Table S3; $F_{1,107} = 4.52$, p = 0.036) than their ancestry-matched control. We also

see that longevity between the dispersal-selected and ancestral population does not vary

differentially due to sex i.e. the difference in longevity between the dispersal selected and

their ancestry-matched control flies is comparable in both males and females (Selection × Sex

$F_{1,107} = 0.76$, p = 0.4). We arrived at the same conclusion when we compared the survivorship

curves through a cox proportional hazards model, yielding a significant main effect of

Selection (Figure 3A, $p = 5 \times 10^{-4}$) and a non-significant effect of Selection × Sex interaction

($p = 0.23$). Interestingly, fitting the lifespan data of the selected (D) and control (C)

populations with a two-parameter Gompertz function revealed that the reduced longevity in

the selected populations primarily results from an accelerated age-dependent mortality rate.

This result is consistent in both males (shape parameter: 0.14 for selected vs. 0.07 for control)

and females (shape parameter: 0.18 for selected vs. 0.097 for control). In contrast, the

baseline mortality in early life was actually lower in the selected population, as indicated by

the rate parameter values: -9.322 for selected males compared to -6.853 for control males,

and -10.037 for selected females compared to -7.515 for control females.

Discussion

In this study, we have used four large (n ~2400) outbred populations of *Drosophila*

melanogaster, which were selected for greater dispersal at their early life. In these dispersal-

evolved populations, we quantified the status of functional senescence in locomotor activity

of both male and female flies, and contrasted the data against their ancestry matched control

populations, which haven't undergone dispersal evolution. Interestingly, we found that

dispersal evolution leads to faster decline in locomotor activity with age (Figure 1) indicating

that evolving greater dispersal at early life, indeed, has adverse effect in the late life stages

resulting in accelerated functional senescence. Further, in order to demonstrate that the above

result is not an artefact of cross-sectional experiment involving population level averages, as

a proof-of-principle, we tracked individual flies throughout their lifespan and measured

locomotor activity at predetermined age-points. This experiment supported the above insight

by yielding the same results (Figure 2).

Moreover, we also found that despite faster age-dependent decline in locomotor activity in

dispersal evolved flies, their locomotor activity does not drop below the activity level of their

ancestry-matched control flies at any given age (Figure 1A and Figure 2A). It is further

noteworthy that this elevated locomotor activity in the dispersal selected flies persisted even

in the absence dispersal selection for over nine generations. Together, these imply that the

dispersal evolved individuals continue to carry and exhibit correlated traits of dispersal

evolution like increased locomotor activity even after they reach their destination, and it can

last over multiple generations long after the selection pressure for dispersal ceases. At the

mechanistic level, these observations indicate that the metabolic remodelling as a correlated

response to dispersal evolution proposed in previous study (Tung et al. 2018b) is likely to

have become constitutive in the dispersal selected flies leading to persistent elevated

expression of locomotor activity in the selected flies throughout their life. Elevated locomotor

activity in the more dispersive individuals throughout their life have important ecological

implications as well. Previous studies have indicated that increased locomotor activity

enhances mating success by elevating courtship behaviour (Kyriacou 1981, Cobb et al. 1987)

and assists females in locating suitable oviposition sites (Ferguson et al. 2015). Consequently,

under similar circumstances, the heightened locomotor activity observed in dispersal-selected

populations would theoretically yield greater reproductive success in a natural environment,

thus contributing to greater evolutionary success. However, we also note a shorter lifespan in

dispersal-selected populations, which could potentially counterbalance the reproductive

advantages conferred by increased locomotor activity. Consequently, determining the net

impact of dispersal evolution on overall reproductive output across the lifespan is not

straightforward. Additionally, it is important to keep in mind that the sustained elevated

locomotor activity in the dispersal-selected populations we see in our study are from

laboratory conditions of *ad libitum* food and absence of predation. The environmental

conditions in the wild would be much harsh and thus sustained elevated locomotor activity

may not be selected for in wild populations. Therefore, further studies are required to

understand the nuanced effect of dispersal selection on reproductive and evolutionary success

of populations selected for greater dispersal.

Furthermore, the observed faster functional senescence and shorter lifespan in dispersal-

selected flies could potentially be a negative consequence of constitutively elevated

expression of locomotor activity in the selected flies throughout their life (Figure 1A and

Figure 2A). Although we do not have direct evidence for a mechanistic explanation behind

these observations, the following mechanisms might be responsible for this. Firstly, the

dispersal-selected flies were previously reported to have evolved an elevated level of

metabolite 3-HK (Tung et al. 2018b), which is associated with age-dependent neural

degeneration in *Drosophila* (Savvateeva et al. 2000) and hence the greater rate of

neuromuscular impairment, leading to deterioration of locomotor activity and overall reduced

lifespan. Secondly, studies have shown that elevated physical activity can lead to the

activation of stress-responsive pathways such as the MAPK and NF-κB pathways (Kramer

and Goodyear 2007, Egan and Zierath 2013), which can lead to increased oxidative stress and

inflammation. Similarly, the selected flies were found to have a greater level of cellular

respiration resulting in an elevated level of ATP in these flies (Tung et al. 2018b). It is also

known that free radicals are produced during ATP production via oxidative phosphorylation

in mitochondria (Cadenas and Davies 2000). These responses can ultimately lead to the

activation of pro-aging pathways and the observed acceleration of aging-associated declines

in physiological function in the selected populations. Thirdly, *Drosophila* insulin-TOR

signalling pathways are known to affect cardiac functional aging, and thereby shorten

lifespan (Wessells et al. 2009). Systemic glucose level is reported to be higher in dispersal-

selected flies (Tung et al. 2018b), which can lead to relatively greater activation of insulin-

TOR signalling pathways in these flies resulting in faster senescence.

While further targeted experiments are necessary to fully understand the exact underlying

mechanisms, this study presents the first empirical evidence that aging evolves as a direct

consequence of dispersal evolution. Moreover, the metabolic changes mentioned above

emerged in response to elevated cellular respiration to meet the increased energy demands for

dispersal in early life (Tung et al., 2018b). The evolutionary theories of aging suggest that

aging evolves as a result of weakened selection pressure later in life (Medawar 1946;

Williams 1957). Given that, for the artificial selection procedure, eggs from these flies are

collected within three days after dispersal, the later part of life remain under selection

shadow. Thus, it is conceivable that over the course of evolution, the evolved populations

have accumulated alleles that facilitate greater dispersal in early life, albeit at the expense of

negative pleiotropic effects in late life, and increasing the likelihood of death. Thus, this

study serves as an empirical demonstration of antagonistic pleiotropy, a population-genetic

concept to explain the evolution of aging based on the broader idea of evolutionary

constraints and life history tradeoffs (Medawar 1946; Williams 1957). Even in natural

population of neotropical butterflies, higher dispersal rates led to decreased longevity (Tufto

et al. 2012). On the other hand, from the existing literature, we also know that often dispersal,

particularly in severely patchy habitats, results in populations with small effective population

size, which leads to strong random genetic drift in these population (Polechová 2022), which

can result in reduced lifespan and accelerated aging, as seen in a study with water fleas,

Daphnia magna (Lohr et al. 2014). However, in a natural setting, it is difficult to dissect out

whether reduced lifespan and accelerated senescence is due to dispersal or due to genetic drift

caused by habitat fragmentation and/or isolation. Whereas in our study, using large outbred

populations we could show that, independent of genetic drift route, dispersal evolution can

have direct negative impact on the physiological function of organisms at late life and their

lifespan.

Moreover, it is worth noting that in a previous study on the same populations (Tung et al.

2018a), the difference between the selected and control populations in terms of dispersal

ability and propensity was apparent as early as after 10 generations of selection. However, the

lifespan of the dispersal-selected and control populations was not significantly different,

although there was a trend (Figure 3C in Tung et al. 2018b). Whereas, the same trade-off

became apparent after 153 generations of dispersal selection. This demonstrates the fact that

the rate of evolution of the correlated traits can differ drastically compared to the traits which

are under direct selection. Further, it also highlights that the trait correlations observed in the

short-term can change over a longer evolutionary timescale.

Finally, our findings demonstrate that although the trade-off between dispersal and longevity

logically indicates a rapid decline of dispersal and associated traits such as locomotor activity

upon relaxation of selection for greater dispersal, it may not occur as per the expectations in

all circumstances. Specifically, we observed elevated levels of locomotor activity in our

dispersal-evolved flies even after nine generations of relaxation of selection pressure. This

apparent discrepancy can be explained by the fact that these flies are maintained in a 15-day

discrete-generation lifecycle, and therefore, they do not experience their full lifespan in these

laboratory conditions. Thus, one may speculate that the trade-off may be avoided in natural

settings where organisms reproduce rapidly at an early stage of life. This hypothesis can be

tested in future research using natural populations. Furthermore, the effect of faster aging and

shortened lifespan on other life-history and physiological traits is an important avenue for

future research to explore.

In summary, evolutionary history has an impact on age-dependent locomotor activity in fruit

flies. The flies that evolved greater dispersal at their early life suffer from steeper age-

dependent decline in locomotor activity, and have shorter lifespan. This pattern of trade-off

between dispersal and senescence is observed in both male and females. The exact molecular

mechanism behind this antagonistic effect (shorter lifespan as a result of selection for

dispersal in early life) remains to be discovered. Additionally, we found that a correlated

increase in locomotor activity due to dispersal selection does not disappear even after nine

generations of removal of selection pressure and the increased locomotor activity remains

higher than that of the control populations until later ages. Please note that locomotor activity

is not the same as dispersal. In our experiment here, locomotor activity change is a correlated

response to dispersal selection. In the current global scenario, where habitat degradation and

destruction, and rapid climate change are rampant, species that are better at dispersing have a

distinct evolutionary advantage. This highlights the importance of gaining a complete

understanding of how dispersal evolution and senescence influence each other. As senescence

greatly affects the performance of an organism, and thereby impacts its long-term persistence

and adaptation, such knowledge can inform us about the distribution and interactions of

species with their environment, and have far-reaching implications for various eco-

evolutionary processes. For example, it can help us predict the success of species in

colonizing new areas, invading new territories, expanding their range, and spreading diseases,

as well as the stability of populations with strong dispersal abilities. Thus, this understanding

is pertinent and relevant to the research community, especially those working in the fields of

evolution, ecology, and conservation.

References

Abrams, P. A. 1991. The fitness costs of senescence: The evolutionary importance of events in early adult life. - Evol. Ecol. 5: 343–360.

Alex Perkins, T. et al. 2013. Evolution of dispersal and life history interact to drive accelerating spread of an invasive species. - Ecol. Lett. 16: 1079–1087.

Andrade-Restrepo, M. et al. 2019. Local adaptation, dispersal evolution, and the spatial eco-evolutionary dynamics of invasion. - Ecol. Lett. 22: 767–777.

Andreu, J. and Barba, E. 2006. Breeding dispersal of Great Tits Parus major in a homogeneous habitat: effects of sex, age, and mating status. in press.

Badyaev, A. V. and Faust, J. D. 1996. Nest Site Fidelity in Female Wild Turkey: Potential Causes and Reproductive Consequences. - The Condor 98: 589–594.

Berg, M. P. et al. 2010. Adapt or disperse: understanding species persistence in a changing world. - Glob. Change Biol. 16: 587–598.

Bonte, D. and Dahirel, M. 2017. Dispersal: a central and independent trait in life history. - Oikos 126: 472–479.

Bonte, D. et al. 2012. Costs of dispersal. - Biol. Rev. 87: 290–312.

Cadenas, E. and Davies, K. J. 2000. Mitochondrial free radical generation, oxidative stress, and aging. - Free Radic. Biol. Med. 29: 222–230.

Chiu, J. C. et al. 2010. Assaying Locomotor Activity to Study Circadian Rhythms and Sleep Parameters in Drosophila. - JoVE J. Vis. Exp.: e2157.

Cobb, M. et al. 1987. The relationship between locomotor activity and courtship in the melanogaster species sub-group of Drosophila. - Anim. Behav. 35: 705–713.

Cote, J. et al. 2010. Personality-dependent dispersal: characterization, ontogeny and consequences for spatially structured populations. - Philos. Trans. R. Soc. B Biol. Sci. 365: 4065–4076.

Clobert J, Le Galliard J-F, Cote J, et al. 2009. Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. Ecol Lett 12:197–209.

Daunt, F. et al. 1999. Experimental evidence that age-specific reproductive success is independent of environmental effects. - Proc. R. Soc. Lond. B Biol. Sci. 266: 1489–1493.

Dean, R. L. et al. 1981. Age-related differences in behavior across the life span of the C57BL/6J mouse. - Exp. Aging Res. 7: 427–451.

Desrochers, A. and Magrath, R. D. 1993. Age-Specific Fecundity In European Blackbirds (Turdus Merula): Individual And Population Trends. - The Auk 110: 255–263.

Deyo, R. A. et al. 1989. Nimodipine ameliorates aging-related changes in open-field behaviors of the rabbit. - Exp. Aging Res. 15: 169–175.

Egan, B. and Zierath, J. R. 2013. Exercise Metabolism and the Molecular Regulation of Skeletal Muscle Adaptation. - Cell Metab. 17: 162–184.

Emborg, M. E. et al. 1998. Age-related declines in nigral neuronal function correlate with motor impairments in rhesus monkeys. - J. Comp. Neurol. 401: 253–265.

Espie, R. H. M. et al. 2000. Age-Dependent Breeding Performance in Merlins(falco Columbarius). - Ecology
81: 3404–3415.

Ferguson, C. T. J. et al. 2015. The sexually dimorphic behaviour of adult Drosophila suzukii: elevated female
locomotor activity and loss of siesta is a post-mating response. - J. Exp. Biol. 218: 3855–3861.

Fernández, J. R. et al. 1999. Differences in locomotor activity across the lifespan of Drosophila melanogaster☆.
- Exp. Gerontol. 34: 621–631.

Fjerdingstad, E. J. et al. 2007. Evolution of dispersal and life history strategies – Tetrahymena ciliates. - BMC
Evol. Biol. 7: 133.

Forero, M. G. et al. 1999. Causes and Consequences of Territory Change and Breeding Dispersal Distance in the
Black Kite. - Ecology 80: 1298–1310.

Forslund, P. and Pärt, T. 1995. Age and reproduction in birds — hypotheses and tests. - Trends Ecol. Evol. 10:
374–378.

Fronhofer, E. A. et al. 2014. Spatially correlated extinctions select for less emigration but larger dispersal
distances in the spider mite Tetranychus Urticae. - Evolution 68: 1838–1844.

Fronhofer E. A. et al. 2015. Condition-dependent movement and dispersal in experimental metacommunities.
Ecol Lett 18:954–963.

Garant, D. et al. 2007. The Multifarious Effects of Dispersal and Gene Flow on Contemporary Adaptation. -
Funct. Ecol. 21: 434–443.

Garratt M. 2020. Why do sexes differ in lifespan extension? Sex-specific pathways of aging and underlying
mechanisms for dimorphic responses. Nutr Healthy Aging 5:247–259.

Goodrick, C. L. 1971. Free Exploration and Adaptation Within an Open Field as a Function of Trials and
Between-Trial-Interval for Mature-Young, Mature-Old, and Senescent Wistar Rats. - J. Gerontol. 26:
58–62.

Green, D. J. 2001. The influence of age on reproductive performance in the Brown Thornbill. - J. Avian Biol.
32: 6–14.

Greenwood, P. J. and Harvey, P. H. 1982. The Natal and Breeding Dispersal of Birds. - Annu. Rev. Ecol. Syst.
13: 1–21.

Grotewiel, M. S. et al. 2005. Functional senescence in Drosophila melanogaster. - Ageing Res. Rev. 4: 372–397.

Hanski, I. et al. 2006. Dispersal-related life-history trade-offs in a butterfly metapopulation. - J. Anim. Ecol. 75:
91–100.

Hargreaves, A. L. and Eckert, C. G. 2014. Evolution of dispersal and mating systems along geographic
gradients: implications for shifting ranges. - Funct. Ecol. 28: 5–21.

Harvey, P. H. et al. 1984. Breeding Dispersal of the Pied Flycatcher (Ficedula hypoleuca). - J. Anim. Ecol. 53:
727–736.

Huang, F. et al. 2015. Rapid evolution of dispersal-related traits during range expansion of an invasive vine
Mikania micrantha. - Oikos 124: 1023–1030.

Kim, S.-Y. et al. 2007. Effects of breeding success, mate fidelity and senescence on breeding dispersal of male
and female blue-footed boobies. - J. Anim. Ecol. 76: 471–479.

Kosmala G. et al. 2017. Locomotor performance of cane toads differs between native-range and invasive
populations. R Soc Open Sci 4:170517.

Kramer, H. F. and Goodyear, L. J. 2007. Exercise, MAPK, and NF-κB signaling in skeletal muscle. - J. Appl.
Physiol. 103: 388–395.

Kyriacou, C. P. 1981. The relationship between locomotor activity and sexual behaviour in ebony strains of
Drosophila melanogaster. - Anim. Behav. 29: 462–471.

Lambrinos, J. G. 2004. How Interactions Between Ecology and Evolution Influence Contemporary Invasion
Dynamics. - Ecology 85: 2061–2070.

Le Bourg, E. 1987. The rate of living theory. Spontaneous locomotor activity, aging and longevity in Drosophila
melanogaster. - Exp. Gerontol. 22: 359–369.

Lenormand, T. 2002. Gene flow and the limits to natural selection. - Trends Ecol. Evol. 17: 183–189.

Levine, M. S. et al. 1987. Sensory, motor and cognitive alterations in aged cats. - Neurobiol. Aging 8: 253–263.

Lohr, J. N. et al. 2014. Reduced lifespan and increased ageing driven by genetic drift in small populations: low
genetic diversity reduces lifespan. - Evolution 68: 2494–2508.

Maag N, Cozzi G, Bateman A, et al. 2019. Cost of dispersal in a social mammal: body mass loss and increased
stress. Proc R Soc B Biol Sci 286:20190033.

Mashau, A. C. et al. 2021. Plant height and lifespan predict range size in southern African grasses. - J. Biogeogr.
48: 3047–3059.

McGhee, K. E. et al. 2021. Maternal stress during pregnancy affects activity, exploration and potential dispersal
of daughters in an invasive fish. - Anim. Behav. 171: 41–50.

McGibbon, C. A. and Krebs, D. E. 2004. Discriminating age and disability effects in locomotion:
neuromuscular adaptations in musculoskeletal pathology. - J. Appl. Physiol. 96: 149–160.

McPeek, M. A. and Holt, R. D. 1992. The Evolution of Dispersal in Spatially and Temporally Varying
Environments. - Am. Nat. 140: 1010–1027.

Medawar PB. 1946. Old age and natural death. Mod Quart 2:30–49

Michelangeli, M. et al. 2017. Aggression mediates dispersal tendency in an invasive lizard. - Anim. Behav. 133:
29–34.

Mishra, A. et al. 2018. Sex differences in dispersal syndrome are modulated by environment and evolution. -
Philos. Trans. R. Soc. B Biol. Sci. 373: 20170428.

Mishra A, et al. 2022. Desiccation Stress Acts as Cause as well as Cost of Dispersal in Drosophila melanogaster.
Am Nat 199:E111–E123.

Miyamoto, M. et al. 1986. Age-related changes in learning and memory in the senescence-accelerated mouse
(SAM). - Physiol. Behav. 38: 399–406.

Nagahara, A. H. and Handa, R. J. 1997. Age-related changes in c-fos mRNA induction after open-field exposure
in the rat brain. - Neurobiol. Aging 18: 45–55.

Narimanov, N. et al. 2022. Accelerated invasion through the evolution of dispersal behaviour. - Glob. Ecol.
Biogeogr. 31: 2423–2436.

Newton, I. 1993. Age and site fidelity in female sparrowhawks, Accipiter nisus. - Anim. Behav. 46: 161–168.

Ochocki, B. M. and Miller, T. E. X. 2017. Rapid evolution of dispersal ability makes biological invasions faster
and more variable. - Nat. Commun. 8: 14315.

Phillips, B. L. et al. 2008. Reid's Paradox Revisited: The Evolution of Dispersal Kernels during Range
Expansion. - Am. Nat. 172: S34–S48.

Polechová, J. 2022. The costs and benefits of dispersal in small populations. - Philos. Trans. R. Soc. B Biol. Sci.
377: 20210011.

Pyle, P. et al. 2001. Effects of age, breeding experience, mate fidelity and site fidelity on breeding performance
in a declining population of Cassin's auklets. - J. Anim. Ecol. 70: 1088–1097.

Reid, J. M. et al. 2003. Age-specific reproductive performance in red-billed choughs Pyrrhocorax pyrrhocorax:
patterns and processes in a natural population. - J. Anim. Ecol. 72: 765–776.

Renault, D. et al. 2018. Environmental Adaptations, Ecological Filtering, and Dispersal Central to Insect
Invasions. - Annu. Rev. Entomol. 63: 345–368.

Savvateeva, E. et al. 2000. Age-dependent memory loss, synaptic pathology and altered brain plasticity in the
Drosophila mutant cardinal accumulating 3-hydroxykynurenine. - J. Neural Transm. 107: 581–601.

Searcy, C. A. et al. 2018. Positive correlation between dispersal and body size in Green Frogs (Rana clamitans)
naturally colonizing an experimental landscape. - Can. J. Zool. 96: 1378–1384.

Sheeba, V. and Joshi, A. 1998. A test of simple models of population growth using data from very small
populations of Drosophila melanogaster. - Curr. Sci. 75: 1406–1410.

Shine, R. et al. 2021. Increased rates of dispersal of free-ranging cane toads (Rhinella marina) during their
global invasion. - Sci. Rep. 11: 23574.

Studds, C. E. et al. 2008. Natal dispersal driven by environmental conditions interacting across the annual cycle
of a migratory songbird. - Proc. Natl. Acad. Sci. 105: 2929–2933.

Travis, J. M. J. et al. 2013. Dispersal and species' responses to climate change. - Oikos 122: 1532–1540.

Tufto, J. et al. 2012. Estimating Brownian motion dispersal rate, longevity and population density from spatially
explicit mark–recapture data on tropical butterflies. - J. Anim. Ecol. 81: 756–769.

Tung, S. et al. 2018a. Simultaneous evolution of multiple dispersal components and kernel. - Oikos 127: 34–44.

Tung, S. et al. 2018b. Evolution of dispersal syndrome and its corresponding metabolomic changes. - Evolution
72: 1890–1903.

van de Pol M, Wright J. 2009. A simple method for distinguishing within- versus between-subject effects using
mixed models. Anim Behav 77:753–758.

van Dijken, F. R. et al. 1979. Divergent selection on locomotor activity in Drosophila melanogaster. III. Genetic
analysis. - Behav. Genet. 9: 563–570.

Wahlström, L. K. 1994a. The significance of male-male aggression for yearling dispersal in roe deer (Capreolus
capreolus). - Behav. Ecol. Sociobiol. 35: 409–412.

Wahlström, L. K. 1994b. The significance of male-male aggression for yearling dispersal in roe deer (Capreolus
capreolus). - Behav. Ecol. Sociobiol. 35: 409–412.

Wessells, R. et al. 2009. d4eBP acts downstream of both dTOR and dFoxo to modulate cardiac functional aging
in Drosophila. - Aging Cell 8: 542–552.

Williams GC. 1957. Pleiotropy, Natural Selection, and the Evolution of Senescence. Evolution 11:398–411.

Wu, N. C. and Seebacher, F. 2022. Physiology can predict animal activity, exploration, and dispersal. -
Commun. Biol. 5: 1–11.

Yu, Y. et al. 2016. Regulation of starvation-induced hyperactivity by insulin and glucagon signaling in adult
661   Drosophila. - eLife 5: e15693.